The *Drosophila* dystonia gene homolog *Neurocalcin* facilitates sleep

by inhibiting a movement-promoting neural network

Ko-Fan Chen, Angélique Lamaze, Patrick Krätschmer and James E.C. Jepson*

Department of Experimental and Clinical Epilepsy, UCL Institute of Neurology, UK

*Correspondence to: j.jepson@ucl.ac.uk

Abstract

Primary dystonia is a hyperkinetic movement disorder linked to altered dopaminergic

signaling and synaptic plasticity in regions of the brain involved in motor control.

Mutations in *HPCA*, encoding the neuronal calcium sensor Hippocalcin, are

associated with primary dystonia, suggesting a function for Hippocalcin in regulating

the initiation and/or maintenance of activity. However, such a role for Hippocalcin or

Hippocalcin homologs has yet to be demonstrated in vivo. Here we investigate the

cellular and organismal functions of the *Drosophila* Hippocalcin homolog

Neurocalcin (NCA), and define a role for NCA in promoting sleep by suppressing

nighttime hyperactivity. We show that NCA acts in a common pathway with the D1-

type Dop1R1 dopamine receptor and facilitates sleep by inhibiting neurotransmitter

release from a multi-component activity-promoting circuit. Our results suggest

conserved roles for Hippocalcin homologs in modulating motor control through

dopaminergic pathways, suppressing aberrant movements in humans and

inappropriate nighttime locomotion in *Drosophila*.

Introduction

Primary dystonia is characterized by repetitive involuntary movements or sustained abnormal postures, and represents the third most common movement disorder after benign tremor and Parkinson's disease (Fahn, 1988; Wenning et al., 2005). Dystonia has been linked to altered neurotransmission and/or plasticity of circuits within the basal ganglia, a group of nuclei involved in action initiation and maintenance, as well as associated inputs/outputs (Calabresi et al., 2016; Karimi and Perlmutter, 2015; Pappas et al., 2015; Weisheit and Dauer, 2015). However, the molecular pathways underlying this disorder remain unclear.

Identifying genetic loci linked to hereditary forms of primary dystonia represents a useful strategy to uncover such pathways. Mutations in *GNAL*, encoding $G\alpha_{olf}$, have been linked to primary dystonia (Fuchs et al., 2013), and since the $G\alpha_{olf}$ G-protein α-subunit couples to D1-type dopamine receptors, this finding supports a link between dystonia and altered dopaminergic signaling (Karimi and Perlmutter, 2015). Indeed, current models suggest that the basal ganglia regulate motor control through two distinct circuits, the direct and indirect pathways, that are defined by expression of D1- and D2-type type dopamine receptors respectively (Gerfen and Surmeier, 2011; Tecuapetla et al., 2016). These pathways antagonistically influence activity of large-scale brain networks (Lee et al., 2016), and an imbalance of direct versus indirect pathway signaling has been proposed to cause dystonia (Breakefield et al., 2008; Peterson et al., 2010; Tanabe et al., 2009; Yokoi et al., 2015). Mutations in *dTorsin*, the *Drosophila* homolog of the dystonia gene *TOR1A*, reduce dopamine levels by suppressing expression of the dopamine synthesis factor GTP cyclohydrolase (Wakabayashi-Ito et al., 2015; Wakabayashi-Ito et al., 2011), further linking dystonia to altered dopamine signaling.

Recent studies have identified an array of further loci linked to primary

dystonia, including *THAP1*, *ANO3*, *KCTD17* and *HPCA* (Charlesworth et al., 2015;

Charlesworth et al., 2012; Fuchs et al., 2009; Mencacci et al., 2015). However, in

vivo data linking these genes to motor control and/or dopaminergic pathways is

limited. Interestingly, mutations in the *Drosophila KCTD17* homologue *insomniac*

result in reduced sleep (i.e increased movement), a phenotype that can be rescued by

inhibiting dopamine synthesis (Pfeiffenberger and Allada, 2012; Stavropoulos and

Young, 2011). Thus, we sought to examine whether other dystonia gene homologs in

Drosophila also impacted sleep, beginning with the *Drosophila HPCA* homolog

Neurocalcin (*Nca*).

NCA and Hippocalcin (the *HPCA* gene product) act as neuronal calcium

sensors, cytoplasmic proteins that bind to calcium via EF hand domains and

translocate to lipid membranes through a calcium-dependent myristoylation switch

(Burgoyne and Haynes, 2012). This switch modulates interactions with membrane-

bound ion channels and receptors, altering target function or localization (Braunewell

and Klein-Szanto, 2009; Burgoyne and Haynes, 2012). Missense mutations in *HPCA*

have been linked to DYT2 primary isolated dystonia, which predominantly affects the

upper limbs, cervical and cranial regions (Charlesworth et al., 2015). The index N75K

mutation in *HPCA* alters a neutral asparagine residue in the second EF hand to a

positively charged lysine, likely interfering with calcium binding in a loss-of-function

manner.

Hippocalcin undertakes pleiotropic roles in mammalian neurons (Braunewell

and Klein-Szanto, 2009), including acting as a calcium sensor to gate the slow

afterhyperpolarisation (sAHP), a calcium-dependent potassium current, and

facilitating NMDA receptor endocytosis during LTD (Jo et al., 2010; Tzingounis et

al., 2007). In *Drosophila*, NCA has previously been shown to be broadly expressed

throughout the adult nervous system (Teng et al., 1994), yet the neuronal and

organismal roles of NCA are unknown. Here we show that NCA is required to

promote sleep by suppressing locomotor activity during the night. We demonstrate

that NCA acts in a common pathway with the D1-type Dop1R1 dopamine receptor

and identify a multi-component wake-promoting neuronal network in which NCA

facilitates sleep by suppressing synaptic release. Thus, we propose that Hippocalcin

homologs play conserved roles in regulating aspects of motor control.

Results

Generation of *Nca* knockout flies

The *Drosophila* genome contains a single *HPCA* homolog, *Neurocalcin* (*Nca*).

Hippocalcin and NCA share > 90% amino-acid identity (Figure 1 – figure supplement

1A), suggesting conservation of function. To investigate the neuronal and behavioral

roles of NCA we generated a *Nca* null allele by replacing the entire *Nca* locus

(including 5' and 3' UTRs) with a mini-*white*$^+$ marker sequence using ends-out

homologous recombination (Baena-Lopez et al., 2013) (Figure 1A, Figure 1 – figure

supplement 1B-C). The mini-*white*$^+$ sequence is flanked by loxP sites, allowing

removal by Cre recombinase and leaving single attP and loxP sites in place of the *Nca*

locus (Figure 1A). As expected, no *Nca* mRNA expression was detected in

homozygotes for the deleted *Nca* locus (Figure 1 – figure supplement 1D, E). Thus,

we term this allele *Nca*^{KO} (*Nca* knockout).

***Nca* knockout flies exhibit reduced night sleep**

Following outcrossing into an isogenic *iso31* control background, *Nca*^{KO}

homozygotes were viable to the adult stage, allowing us to test whether NCA impacts

locomotor control. To do so, we first used a high-throughout yet low resolution

system, the *Drosophila* Activity Monitor (DAM) (Pfeiffenberger et al., 2010a), which

counts locomotor activity via perturbations of an infrared beam intersecting a glass

tube housing individual flies. Using this system, under 12 h light: 12 h dark

conditions (12L: 12D, 25°C) we found that *Nca*^{KO} males exhibited two distinct

locomotor phenotypes. Firstly, reduced maximal locomotor activity in response to

lights-off (also termed the startle response or masking) (Figure 1 – figure supplement

2A, B). Secondly, increased locomotor activity during the night but not the day

(Figure 1 – figure supplement 2A, C, D). This later phenotype was suggestive of

reduced night sleep. We therefore quantified sleep levels in Nca^{KO} males and controls

using a 5 min period of inactivity to define a sleep bout – the standard definition in

the field (Pfeiffenberger et al., 2010b). Indeed, Nca^{KO} males exhibited reduced night

sleep but not day sleep relative to controls (Figure 1 – figure supplement 3).

Interestingly, the impact of removing NCA on night sleep appeared further enhanced

by shortening photoperiod, such that night (but not day) sleep was substantially

reduced in Nca^{KO} males under 8L: 16D (Figure 1B-D). Similarly to 12L: 12D, peak

locomotor activity following lights-off was reduced in Nca^{KO} males under 8L: 16D,

and the number of beam breaks during the normally quiescent period of the night was

increased (Figure 1 – figure supplement 4).

To obtain a higher resolution analysis of locomotor patterns in Nca^{KO} flies we

next utilized a video-tracking method - the DART (*Drosophila* ARousal Tracking)

system (Faville et al., 2015). Continual video-monitoring of Nca^{KO} males and *iso31*

controls confirmed robust sleep loss in Nca^{KO} males specifically during the night

(Figure 1E-G) accompanied by significantly shortened sleep bouts (Figure 1 – figure

supplement 5). By examining locomotor patterns in individual flies, we found that

Nca^{KO} males consistently displayed prolonged activity relative to controls following

lights-off and frequent bouts of movement even in the middle of the night, a period of

quiescence in *iso31* controls (Figure 1H, I). The velocity of locomotion in Nca^{KO}

males was also significantly increased relative to moving controls in the normally

quiescent period of the night (4-12 h after lights-off), whereas peak velocity during

the startle response to lights-off was reduced (Figure 1 – figure supplement 6), in

agreement with data collected using the DAM system. Collectively, the above data

suggest that complete loss of NCA results in both locomotor deficits and profound

nighttime hyperactivity, resulting in reduced peak levels of activity and night sleep.

NCA is required in neurons to promote night sleep

To identify the cellular substrates in which NCA acts to regulate locomotion and sleep

we performed cell-specific knockdown of *Nca* expression using transgenic RNA

interference (RNAi). NCA has been shown to be widely expressed in neuropil regions

throughout the *Drosophila* brain (Teng et al., 1994). Thus, we initially examined

neurons as a potential cellular candidate. We used three independent RNAi lines

(*kk108825, hmj21533* and *jf03398,* termed *kk, hmj* and *jf* respectively) to reduce *Nca*

expression in adult male neurons using the pan-neuronal *elav*-Gal4 driver. The *kk* and

jf dsRNAs target a partially overlapping sequence of *Nca*, whereas *hmj* targets a

distinct, non-overlapping upstream sequence (Figure 2 – figure supplement 1A). For

each RNAi line, we confirmed reduced *Nca* expression using qPCR (Figure 2 – figure

supplement 1B). Transcription of *Nca* occurs from promoter regions shared with the

downstream locus *cg7646*, yet *cg7646* transcription was not affected by *Nca* RNAi

(Figure 2 – figure supplement 1C), nor were any common off-target mRNAs

predicted for the *kk*, *hmj* or *jf* dsRNA hairpins (data not shown). Importantly, under

8L: 16D conditions, pan-neuronal expression of all three *Nca* RNAi lines specifically

reduced night sleep in males as measured by the DAM system (Figure 2A-C, Figure 2

– figure supplement 2). However, *Nca* knockdown did not result in a reduction in

peak locomotor activity as observed in *Nca*^{KO} males (Figure 2 – figure supplement 3).

Thus, neuronal NCA predominantly impacts sleep/activity during the night, with

NCA potentially acting in other cell types to regulate peak locomotor levels. We

therefore focused on elucidating the genetic pathways and neuronal circuits in which

NCA acts to promote night sleep.

To test whether sleep loss caused by neuronal *Nca* knockdown flies was due to

an indirect effect on the circadian clock, we examined whether *Nca* knockdown

altered circadian patterns of locomotor activity under constant dark conditions (Figure

2 – figure supplement 4A, B). Importantly, knockdown of *Nca* in neurons did not alter

circadian rhythmicity (Figure 2 – figure supplement 4A, B), nor did *Nca* expression

cycle in whole fly heads (Figure 2 – figure supplement C). Thus, it is unlikely that

sleep loss caused by neuronal *Nca* knockdown is due to circadian clock dysfunction.

We extended our initial findings in *Nca* knockdown males harbouring the pan-

neuronal *elav*-Gal4 driver and found that night sleep loss due to neuronal *Nca*

knockdown was also observed in adult virgin females (Figure 2 – figure supplement

5), and in male flies expressing the *kk Nca* RNAi using distinct pan-neuronal (*nsyb*-

Gal4) or broadly expressed (*insomniac*-Gal4) drivers (Figure 3A). In contrast,

expression of the *kk Nca* RNAi in muscle cells did not alter night sleep in adult males

(Figure 2 – figure supplement 6), supporting the premise that NCA acts in the nervous

system to regulate sleep. Video tracking further confirmed that neuronal expression of

kk Nca RNAi reduced night sleep (Figure 2D-F). Collectively, the above data

demonstrate that NCA acts in neurons to promote night sleep in *Drosophila*. For

simplicity, we use the *kk Nca* RNAi for all subsequent experiments, and refer to flies

expressing *kk Nca* RNAi under *elav*-Gal4 as Nca^{KD} (*Nca* knockdown). Given the

similar results obtained using the DAM and DART systems for both Nca^{KO} and

Nca^{KD} flies, we use DAM as a high-throughput method for all sleep measurements

detailed below, which are performed in 8L: 16D conditions at 25°C.

NCA acts in a common pathway with the Dop1R1 dopamine receptor

In *Drosophila*, dopamine is a pro-arousal factor, with elevated dopaminergic neurotransmission strongly reducing sleep (Kume et al., 2005). Recent studies have shown that the pro-arousal effect of elevated dopamine is mediated by the D1-type Dop1R1 dopamine receptor (Liu et al., 2012; Ueno et al., 2012). Furthermore, both hypo- and hyper-dopaminergic signalling within the human basal ganglia has been proposed to underlie forms of primary dystonia (Breakefield et al., 2008). We therefore tested whether *Nca* promotes sleep in a common pathway with genes involved in dopaminergic signalling. Indeed, we found that heterozygosity for a null or strongly hypomorphic allele of the *Dop1R1* dopamine receptor (*Dop1R1*$^{MI03085-GFST.2}$, a homozygous lethal MiMIC insertion) rescued night sleep loss in *Nca*^{KD} flies (Figure 2G, H). Importantly, in both *elav*-Gal4/+ and *kk*/+ control backgrounds, heterozygosity for *Dop1R1*$^{MI03085-GFST.2}$ did not alter sleep levels (Figure 2G, H; p > 0.05, Kruskal-Wallis test with Dunn's post-hoc test). A similar epistatic interaction between *Nca* and *Dop1R1* was observed using a second, weaker *Dop1R1* allele (*Dop1R1*^{MI004437}) (Figure 2 – figure supplement 7).

Mammalian Hippocalcin regulates cell-surface levels of NMDA receptors during LTD (Jo et al., 2010), and *Drosophila nmda receptor 1* (*dNR1*) mutants exhibit reduced sleep during the night (Tomita et al., 2015), similarly to *Nca* knockout and knockdown flies. However, in contrast to *Dop1R1*, we found no signatures of genetic interaction between *Nca* and *dNR1*, suggesting that these loci act in distinct pathways to promote sleep (Figure 2 – figure supplement 8).

NCA acts in two distinct circuits to promote night sleep

We next sought to delineate the neural circuits in which NCA functions to promote

night sleep. Using transgenic RNAi, we performed an extensive screen of sleep

relevant circuits defined by numerous *promoter*-Gal4 driver lines (Figure 3A, Figure

3 – figure supplement 1). These include clock neurons, dopaminergic and other

neurotransmitter-specific subtypes, fan-shaped body neurons, mushroom body (MB),

and sensory neurons (Figure 3A) (Donlea et al., 2011; Joiner et al., 2006; Lamaze et

al., 2017; Liu et al., 2014; Pitman et al., 2006; Seidner et al., 2015; Sitaraman et al.,

2015). Given the genetic interaction between *Nca* and *Dop1R1*, we also utilised

genomic enhancer elements in the *Dop1R1* locus to drive *Nca* knockdown in subsets

of potential Dop1R1-expressing neurons (Figure 3A, Figure 3 – figure supplement 1)

(Jenett et al., 2012; Jiang et al., 2016). However, in contrast to broadly expressed

drivers (*elav*-, *nsyb*- and *inc*-Gal4), *Nca* knockdown in restricted neural subsets was

insufficient to significantly reduce night sleep (Figure 3A, Figure 3 – figure

supplement 1).

These results suggested a complex sleep-relevant circuit requirement for

NCA. We therefore reduced NCA levels in multiple sub-circuits to test for a

simultaneous role of NCA in distinct anatomical regions. Through this approach, we

found that *Nca* knockdown using two *enhancer*-Gal4 lines (*R14A05* – an enhancer in

the *single-minded* locus, and *R72C01* – an enhancer in the *Dop1R1* locus; see Figure

3B for expression patterns) was sufficient to strongly phenocopy the effect of pan-

neuronal *Nca* knockdown on night sleep (Figure 3C, D; compare Figure 3C with

Figure 2A). For simplicity we refer to these drivers as *A05* and *C01* respectively.

The *A05* enhancer drives expression in approximately 70 neurons, as

quantified using a fluorescent nuclear marker (Figure 3 – figure supplement 2A, B)

that include a subset of MB neurons, a cluster of cell bodies adjacent to the anterior

ventrolateral protocerebrum (AVP), and two visual sub-circuits: optic lobe (OL) and

anterior optic tubercle (AOTU) neurons (Figure 3B). *C01* drives expression in

approximately 250 neurons (Figure 3 – figure supplement 2C, D) that include the

MBs, neurons projecting to the MB γ-lobes, the antennal mechanosensory and motor

center (AMMC) (Figure 3B) and the superior medial protocerebrum (SMP). Both

drivers label additional cell bodies of unknown identity. The potential overlap of *A05*

and *C01* in the MBs raised the possibility that sleep loss in *A05/C01 > Nca* RNAi

flies was due to strong NCA knockdown in neurons common to both the *A05* and *C01*

enhancers. If so, driving *Nca* RNAi with two copies of either *A05* or *C01* should

mimic sleep loss in *A05/C01 > Nca* RNAi flies. However, this was not the case

(Figure 4 – figure supplement 3). Thus, NCA is simultaneously required in two non-

overlapping sub-circuits defined by the *A05* and *C01* enhancers.

Given that *C01* is a *Dop1R1* enhancer element, that *Nca* and *Dop1R1*

genetically interact to regulate sleep (Figure 2G, H), and that Dop1R1 is highly

expressed in the MBs (Lebestky et al., 2009), we tested whether the MBs were a

constituent of the *C01* expression domain by swapping *C01* for the MB-specific

driver *ok107* and measuring sleep in flies expressing *Nca* RNAi in both *A05* and MB

neurons. Indeed, knockdown of *Nca* in both *A05* and MB neurons also specifically

reduced night sleep (Figure 3 – figure supplement 4), albeit to a weaker degree

compared to knockdown in *A05* and *C01* neurons (compare with Figure 3C, D). Thus,

we conclude that the MBs are a component of a complex network defined by *C01*-

Gal4 with additional, as yet undefined, neurons acting within both the *C01* and *A05*

domains to regulate night sleep.

NCA promotes sleep by suppressing synaptic output from a wake-promoting circuit

We next assessed how NCA impacts excitability of *C01* and *A05* neurons. To do so, we expressed a genetically-encoded fluorescent indicator of neurotransmitter release, UAS-*synaptopHluorin* (*spH*) (Miesenbock, 2012), in *C01* and *A05* neurons of either wild type or *Nca*^{KD} males. spH is localised to presynaptic neurotransmitter-containing vesicles and increases in fluorescence in a pH-dependent manner upon fusion of synaptic vesicles with the presynaptic membrane, providing an optical read-out of neurotransmitter release (Miesenbock, 2012). Intriguingly, we found that *Nca* knockdown significantly enhanced spH fluorescence in the MB α/β-lobes and the AMMC but not in the MB γ-lobe region or the SMP (Figure 4).

Since neurotransmitter release is enhanced in subsets of *C01* and *A05* neurons following *Nca* knockdown, this suggested that NCA normally acts to inhibit synaptic output in these circuits. If sleep loss in *Nca* knockdown flies directly results from a loss of such inhibition in *C01* and *A05* neurons (thus enhancing neurotransmitter release), we predicted the following: firstly, that artificial activation of *C01* and *A05* neurons should be sufficient to promote locomotor activity (and thus sleep loss), and secondly, that silencing *C01* and *A05* neurons should suppress sleep loss in *Nca* knockdown flies. To test our first prediction, we enhanced the excitability of *C01* and *A05* neurons by expressing the temperature-sensitive channel TrpA1 in either neuronal subset or both and shifting flies from a non-activating temperature (22°C) to an activating temperature (27°C) sufficient to cause neural excitation through TrpA1-mediated cation influx (Hamada et al., 2008) (Figure 5A). At the non-activating temperature, over-expression of TrpA1 in either circuit or both did not alter sleep levels (Figure 5B). At the activating temperature, excitation of *A05* neurons did not

alter night sleep levels relative to controls (Figure 5C, D). In contrast, excitation of

C01 neurons profoundly reduced night sleep (Figure 5C, D) as well as day sleep

(Figure 5C). Interestingly, simultaneous activation of *C01* and *A05* neurons further

reduced night sleep but not day sleep relative to activation of *C01* neurons alone

despite the lack of effect of *A05* neuron activation on sleep (Figure 5C, D).

To test our second prediction, we over-expressed a non-inactivating outward

rectifying potassium channel (dORKΔC2) in *C01* and *A05* neurons with and without

Nca knockdown via RNAi. Here, expression of dORKΔC2 is predicted to suppress

neuronal firing by hyperpolarizing the resting membrane potential (Nitabach et al.,

2002; Park and Griffith, 2006). Silencing *C01* and *A05* neurons with dORKΔC2 in an

otherwise wild type background did not alter day or night sleep levels (Figure 5E, F).

However, consistent with our above prediction, dORKΔC2 expression significantly

suppressed night sleep loss due to *Nca* knockdown in *C01* and *A05* neurons (Figure

5E, F). Thus, we propose that NCA promotes night sleep by limiting synaptic output

from a multi-component activity-promoting circuit coordinately defined by subsets of

the *C01*- and *A05*-Gal4 expression domains. Our results further suggest that the *C01*

and *A05* circuits interact to suppress sleep, with *C01* neurons acting as a predominant

pro-arousal circuit and *A05* neurons acting in a modulatory manner to enhance the

impact of *C01* activation on night sleep.

Discussion

Previous work has demonstrated that *Drosophila* Neurocalcin (NCA), a member of

the neuronal calcium sensor family, is widely expressed throughout the fly brain and

localizes to synaptic regions (Teng et al., 1994). However, the neurobiological

functions of NCA have remained unclear. Here we define a role for NCA in

promoting night sleep and show that NCA acts via inhibiting synaptic output from a

complex movement-promoting circuit.

Previous genetic screens have identified an array of sleep-promoting factors in

Drosophila (Tomita et al., 2017). However, despite extensive circuit analyses, the

activity of such proteins does not fully map onto known sleep-regulatory neurons, and

the complete neural substrates in which these factors act have yet to be determined

(Afonso et al., 2015; Rogulja and Young, 2012; Shi et al., 2014; Stavropoulos and

Young, 2011; Tomita et al., 2015; Wu et al., 2014). Our results are consistent with

these findings and offer a tentative explanation for the difficulties in defining circuit

requirements for sleep-relevant proteins. We show that NCA activity is not required

within a single cell-type or neuropil region to promote sleep. Instead, sleep-relevant

NCA activity can largely be localised to *two* distinct domains of the *Drosophila*

nervous system defined by the *A05*- and *C01*-Gal4 drivers. One relevant cell-type

within these domains is the mushroom bodies (MBs), a known sleep-regulatory center

(Joiner et al., 2006; Pitman et al., 2006; Sitaraman et al., 2015). However, the MBs

only partially contribute to one of the two sleep-relevant domains, indicating that

NCA acts in a dispersed network that likely incorporates several non-overlapping

wake-promoting circuits.

Interestingly, whereas activation of *C01* neurons is sufficient to suppress

sleep, activation of *A05* neurons reduces night but not day sleep only in the context of

C01 neuron activation. We note that activation of *C01* neurons profoundly reduces

sleep during both day and night periods, whereas only night sleep is impaired in flies

with reduced NCA expression in *C01* and *A05* neurons. Based on these findings, we

hypothesize that reduction of NCA in *C01* and *A05* neurons causes mild

hyperexcitation, which in *C01* neurons alone is insufficient to modulate sleep but in

both *C01* and *A05* neurons simultaneously causes an increase in network excitability

sufficient to reduce night sleep. A selective inhibition of neurotransmitter release by

NCA in subsets of *C01* and *A05* neurons is supported by our in vivo imaging data

showing that *Nca* knockdown increases synaptic output in the MB α/β-lobes and

AMMC but not the MB γ-lobe region or the SMP. Since NCA is only partly required

in the MBs, we consider it unlikely that increased synaptic output within the MBs and

AMMC alone is sufficient to drive sleep loss in *Nca* knockdown flies. Rather, we

propose that coincident hyperactivity of several sub-circuits within the *C01* and *A05*

domains collectively drives night sleep loss. Temporal information encoded by light-

sensing or circadian pathways may further demarcate the period in which NCA

promotes sleep (i.e the night), with such inputs likely to be bypassed by ectopic

activation of *C01* neurons, resulting in sleep loss across 24 h.

Since silencing of *C01* and *A05* neurons does not alter basal sleep, we suggest

that these circuits are normally activated by an ethological stimulus absent during our

experimental conditions. We further postulate that modulatory dopamine signalling

through Dop1R1 promotes neural activity in wake-promoting *C01* neurons. These

neurons are defined by a *Dop1R1* enhancer element, and thus it is likely that at least a

subset of *C01* neurons express Dop1R1. Indeed, dopamine signalling via Dop1R1 has

been shown to enhance synaptic transmission between antennal lobe projection

neurons and MB neurons (Ueno et al., 2013). This model is attractive since reducing

such input could balance the increased synaptic release caused by *Nca* knockdown,

providing an explanatory basis for the genetic interaction between *Nca* and *Dop1R1*.

How might NCA inhibit synaptic output from *C01* and *A05* neurons? The

mammalian NCA homolog Hippocalcin acts pleiotropically in several pathways that

control neuronal excitability and plasticity, facilitating NMDA receptor endocytosis

during LTD and gating the slow afterhyperpolarisation, a calcium-activated potassium

current controlling spike frequency adaptation that is thought to be mediated by a

complex array of potassium channels (Andrade et al., 2012; Jo et al., 2010;

Tzingounis et al., 2007). Recent data suggest that Hippocalcin also negatively

regulates calcium influx through N-type voltage-gated calcium channels (Helassa et

al., 2017). Given the strong homology between Hippocalcin and NCA, it is possible

that NCA suppresses neurotransmitter release through similar pathways in *Drosophila*

neurons, although the lack of genetic interaction between *Nca* and the *dNR1* NMDA

receptor suggests that an enhancement of NMDA receptor levels is unlikely to

significantly contribute to sleep loss in *Nca* knockout/knockdown flies.

Mutations in Hippocalcin also cause the movement disorder DYT2 primary

isolated dystonia (Charlesworth et al., 2015). What is the common neurobiological

link between dystonic movements in humans and sleep loss in *Drosophila*? We

suggest that both phenotypes fundamentally reflect a dysregulation of motor control

i.e when activity is initiated, and for how long such activity is maintained.

Hippocalcin/Neurocalcin homologs have the potential to act as molecular toggle

switches, inhibiting sustained rapid-firing and neurotransmitter release in activity-

promoting neurons via simultaneous modulation of potassium and calcium channel

function (Helassa et al., 2017; Tzingounis et al., 2007). Mutations in these calcium

sensors may thus result in prolonged bouts of synaptic output and corresponding

motor activity. We note that *insomniac*, another sleep-promoting gene in *Drosophila*,

is homologous to the myoclonus dystonia-gene *KCTD17* (as well the paralogs *KCTD2*

and *KCTD5*) (Li et al., 2017; Mencacci et al., 2015; Stavropoulos and Young, 2011).

It will thus be intriguing to test if other dystonia gene homologs also promote sleep in

Drosophila, and whether human dystonia and *Drosophila* sleep loss can thus be

considered homologous phenotypes (or phenologs) linked to mutations in a conserved

genetic network that functions to suppress inappropriate activity in both humans and

flies (Lehner, 2013; McGary et al., 2010).

Materials and Methods

Fly husbandry

Flies were maintained on standard fly food at constant temperature 25^{o}C under 12 h:

12 h light-dark cycles (12L: 12D). The following strains were obtained from the

Bloomington and VDRC stock centers: kk108825 (v100625), hmj21533 (54814),

jf03398 (29461), *Dop1R1*$^{MI03085-GFSTF.2}$ (59802), *Dop1R1*^{MI04437}(43773), *ple*-Gal4

(8848), *Chat*-Gal4 (6798), *vGlut*-Gal4 (26160), *GAD*-Gal4 (51630), *Ddc*-Gal4(7010),

GMR-Gal4 (1104), *Trh.1*-Gal4 (38388), *Tdc2*-Gal4(9313), *C5*-Gal4 (30839), *ok107*-

Gal4 (854) and *A502* (16130). The remaining lines obtained from the Bloomington

stock center are part of the Janelia Flylight collection with identifiable prefixes:

R23E10-Gal4, R55B01-Gal4, R52H12-Gal4, Hdc-Gal4 (R17F12-Gal4), R14A05-

Gal4, R72B05-Gal4, R72B07-Gal4, R72B08-Gal4, R72B11-Gal4, R72C01-Gal4, and

R72C02-Gal4. The following lines were gifts from laboratories of Kyunghee Koh:

elav-Gal4, *nsyb*-Gal4, *tim*-Gal4 and *TUG*-Gal4; Joerg Albert: *nompC*-Gal4

(Kamikouchi et al., 2009) and Nicolas Stavropouplos: *inc*-Gal4:2 (Stavropoulos and

Young, 2011). *ppk*-Gal4 and *TrpA1*-CD-Gal4 were described previously (Zhong et

al., 2012). *GMR-hid*, *tim*^{KO} and *cry*^{02} were previously described in (Lamaze et al.,

2017). Except for *Ddc*-Gal4, *Trh.1*-Gal4, *Tdc2*-Gal4, *nompC*-Gal4 and *Hdc*-Gal4, all

Drosophila strains above were either outcrossed five times into an isogenic control

background (*iso31*) or insertion-free chromosomes were exchanged with the *iso31*

line (*hmj21533, jf03398, Dop1R1*$^{MI03085-G}$, *Dop1R1*^{MI04437} and *NMDAR1*^{MI11796}) before

testing for sleep-wake activity behaviour. Note: R14A05-Gal4 was initially

mislabelled as R21G01-Gal4 in Bloomington shipment. The clear mismatch between

the image of R21G01>GFP in FlyLight database and our immunostaining data (A05,

Fig3B) led us to clarify the identity of the line, as R14A05-Gal4, by sequencing

genomic PCR product using primers pair: pBPGw_ampF: agggttattgtctcatgagcgg and

pBPGw_Gal4R: ggcgcacttcggttttttctt.

Generation of the *Nca^{KO}* line

Null alleles of *Nca* were generated using homologous recombination as described

previously (Baena-Lopez et al., 2013). Briefly, genomic DNA was extracted from 20

wild type flies (Canton S) using the BDGP buffer A-LiCl/KAc precipitation protocol

(http://www.fruitfly.org/about/methods/inverse.pcr.html). The 5' (Arm 1) and 3'

(Arm 2) genomic regions flanking the *Nca* coding sequence were PCR amplified via

high fidelity DNA polymerase (Q5 high-fidelity 2X master mix, M0492S, NEB) with

the following primers: NotI_Arm1F1:gcggccgctaatttgcagctctgcatcg,

NotI_Arm1R1:gcggccgcatggtaagaagcacgcaacc,

AscI_Arm2F1:ggcgcgccttatgaccgttccaaaacacc,

AvrII_Arm2R1:cctagggggctaaatacgttgaccaagc. The corresponding Arm1 and Arm2

fragments (~2.5kb) were gel purified (Wizard® SV Gel and PCR Clean-Up System,

A9281, Promega) and cloned into pCR-Blunt II-TOPO vector (Zero Blunt® TOPO®

PCR Cloning Kit, 450245, ThermoFisher Scientific), and subsequently sub-cloned via

NotI (R3189S, NEB) and AscI/AvrII digestion (R0558S and R0174S, NEB) and T4

ligation (M0202S, NEB) into pTV vector, a P-element construct containing the mini-

white^+ marker and UAS-*reaper* flanked by FRT and I-SceI sites (Baena-Lopez et al.,

2013). The sequence identifies of Arm 1 and Arm 2 fragments within the pTV vector

were verified via Sanger sequencing using the following primers: nca1_f:

cagctctgcatcgctttttgt, nca1_3_f: ccctcgcgcatggtacttta, nca1_r: agcgtcacataagttctccca,

nca1_4_f: tggacgaaaataacgatggtca, nca1_5_f: agactacttagccatgttttcatact, nca1_2_f:

tgacgaagccacaattaaagagtg, nca1_1_f: gcaaccctgttcccctttca,

nca2_f: gaccgttccaaaacacccca, nca2_3_f: ttgttgtgcgccacgttttc, nca2_r:

acgtatgctccatgattcctct

nca2_4_f: tgcaggtcggttaatcaatgc, nca2_5_f: tcaatcgatttggggccagg, nca2_2_f:

ccttctccaggctcagcaaa, nca2_1_f: actctgcatttcgataagattagcc. Donor lines containing

pTV vector with Arm1 and Arm2 homologous fragments (pTV_nca1+2) were then

generated via embryonic injection and random P-element mediated genomic

insertions (Bestgene, inc.). To initiate homologous recombination between

pTV_nca1+2 and the endogenous *Nca* locus, donor lines were crossed to *yw*; hs-*flp*,

hs-I-*SceI*/CyO and the resulting larvae were heat shocked at 48 h and 72 h after egg

laying for 1 h at 37°C. Around 200 female offspring with mottled/mosaic red eyes

were crossed in pools of three to *ubiquitin*-Gal4[3xP3-GFP] males to remove

nonspecific recombination events (via UAS-*reaper*-mediated apoptotic activity). The

crossings were flipped once over and the progeny (~12000 adults) was screened for

the presence of red-eyed and GFP-positive flies. Three independent GFP[+] red-eyed

lines (*ko1*, *ko2*, and *ko3*) were identified. The exchange of endogenous *Nca* locus with

pTV_nca1+2 fragments was confirmed by detecting a 2.6 kb PCR product (Figure 1

figure supplement 1C) in the genomic DNA samples of the above three lines (pre-

digested by EcoRI/NotI) using the following primer pairs: ncaKO-F2:

tgggaattgactgatacagcct; ncaKO-R2: ggcactacggtacctgcat. ncaKO-F2 matches to the

region between 24 bp and 2 bp upstream of Arm1 and ncaKO-R2 overlaps with attP

site (Figure 1A). The absence of endogenous *Nca* mRNA in *ko1* flies was confirmed

by standard and quantitative RT-PCR (Figure 1 figure supplement 1D; see also the

below RNA section). The min-*white*[+] cassette and majority of pTV vector sequences

were further removed from the *ko1* genome via Cre-loxP recombination (Figure 1A).

This "Cre-out" strain was then backcrossed five times to a *Nca*[A502] line (where *A502*

is a P-element insertion 2 kbp upstream of the *Nca* CDS) that was outcrossed

previously into the *iso31* background (see Fly husbandry section). Before testing for

changes in sleep/wake behaviour, the resulted line, termed *Nca* knockout (*Nca*[KO]),

was lastly verified by sequencing a 576 bp genomic PCR product (using primer pair:

nca1_5_f and nca2_r), confirming the absence of *Nca* CDS sequence and the insertion

an attP site in the *Nca* locus.

RNA extraction and Quantitative PCR

For RNA extractions, 10-20 fly heads per genotype were collected with liquid

nitrogen and dry ice. Total RNA was extracted using TRIzol™ reagent following

manufacturer's manual (Thermo Fisher Scientific). cDNA was reverse transcribed

from 250 or 500 ng of DNase I (M0303S, NEB) treated RNA via MMLV RT

(M170A, Promega). A set of five or six standards across 3125-fold dilution was

prepared from the equally pooled cDNA of all genotypes in each experiment.

Triplicated PCR reactions were prepared in 96-well or 384-well plates for standards

and the cDNA sample of each genotype (20 to 40 fold dilution) by mixing in Power

SYBR Green Master Mix (Thermo Fisher Scientific) and the following primer sets:

ncaqF2: acagagttcacagacgctgag, ncaqR2: ttgctagcgtcaccatatggg; cg7646F:

gcctttcgaatgtacgatgtcg, cg7646R: cctagcatgtcataaattgcctgaac or

rp49F:cgatatgctaagctgtcgcaca, rp49R: cgcttgttcgatccgtaacc. PCR reactions were

performed in Applied Biosystems StepOne (96-wells module) or QuantStudio 6Flex

instruments (384 wells module) using standard thermocycle protocols. Melting curve

analysis was also performed to evaluate the quality of the PCR product and avoid

contamination. The Ct values were exported as csv files and a standard curve between

Ct values and logarithm of dilution was calculated using the liner regression function

in Graphpad. The relative expression level for *Nca*, *cg7646* and *rp49* of each sample

were estimated by interpolation and anti-logarithm. The expression levels of *Nca* and

cg7646 for each genotype were further normalized to their respective average *rp49*

expression level. Statistical differences between the normalized expressions levels of

each genotype were determined by Mann-Whitney test or Kruskal-Wallis test with

Dunn's post-hoc test using Graphpad software.

Sleep-wake behavioral analysis

Three to five days old male or virgin female flies were collected and loaded into glass

tubes containing 4% sucrose and 2% agar (w/v). Sleep-wake behavior was recorded

using the *Drosophila* Activity Monitor (DAM, TriKinetics, inc.) system or

Drosophila ARousal Tracking (DART, BFKlab) system for 3 days in the designated

LD regime (L12: D12 or L8: D16) at 25$^{\text{o}}$C. Behavioral recordings from the third day

of the given LD regime were then analyzed. All flies were entrained to 12L: 12D

prior to entering designated LD regimes. For ectopic activation experiments involving

UAS-*TrpA1*, flies were cultured in 18$^{\text{o}}$C during development and then entrained to

L8: D16 at 22$^{\text{o}}$C before entering L8: D16 condition at 27$^{\text{o}}$C. *Drosophila* activity (or

wake) is measured by infra-red beam crosses in DAM or by direct movement tracking

in DART. Sleep is defined by 5 minutes of inactivity (where inactivity is defined as

no beam crosses during 1 min in the DAM or less than 3 mm movement in 5 s in

DART). The csv output files with beam crosses (DAM) or velocity data (DART)

were processed by a customized Excel calculators (Supplementary file 1) and R-

scripts (https://github.com/PatrickKratsch/DAM_analysR) to calculate the following

parameters for individual flies: *Onset and offset of each sleep bout*, *sleep bout length*,

day and night sleep minutes, *daily total sleep minutes*, and *daily sleep profile* (30

minutes interval). An established MATLAB® based tool, Flytoolbox, was used for

circadian rhythmicity analysis (Levine et al., 2002a, b). Briefly, the strength of

rhythmicity (RI) was estimated using the height of the third peak coefficient in the

auto-correlogram calculated for the activity time series of each fly. Rhythmic

Statistics values were then obtained from the ratio of the RI value to the 95%

confidence interval for the correlogram ($2/\sqrt{N}$, where N is the number of

observations, which correlatively increase with the sampling frequency), in order to

determine statistical significance of any identified period (RS is ≥ 1)

Immunohistochemistry and confocal microscopy

Adult male R72C01 > CD8::*GFP* and R21G01 > CD4::*tdTomato* flies were

anesthetized in 70% ethanol before brains were dissected in PBT (0.1M phosphate

buffer with 0.3% TritonX100) and collected in 4% paraformaldehyde/PBT on ice.

The fixation was then performed at room temperature for 15 min before washing 3

times with PBT. The brain samples were blocked using 5% goat serum/PBT for 1 h at

room temperature before incubation with primary antibodies. The samples were

washed 6 times with PBT before incubated with Alexa Fluor secondary antibodies in

5% goat serum/PBT at 4°C over 24 h. After washing 6 times with PBT, the samples

were mounted in SlowFade Gold antifade reagent (S36936, Thermo Fisher Scientific)

on microscope slides and stored at 4°C until imaged using an inverted confocal

microscope Zeiss LSM 710. Primary antibody concentrations were as follows: mouse

anti-nc82 (Developmental Studies Hybridoma Bank) - 1:200; rabbit anti-GFP

(Invitrogen) - 1:1000; rabbit anti-dsRED (Clontech) - 1:2000. Alexa Fluor

secondaries (Invitrogen) were used as follows: Alexa Fluor 647 goat anti-mouse IgG -

1:500, Alexa Fluor 488 goat anti-rabbit IgG - 1:2000, Alexa Fluor 555 goat anti-rabbit

IgG - 1:2000. For quantification of nuclei number in *C01 > red-stinger* and *A05 >*

red-stinger brains, unstained Red-Stinger fluorescence was captured via confocal

microscopy. DAPI (Sigma Aldrich) was used to counterstain nuclei (at a dilution of

1:5000). The number of Red-Stinger-positive nuclei in each brain was subsequently

quantified using the ImageJ 3D Objects Counter tool, with a variable threshold used

to incorporate all of the visible Red-Stinger-positive nuclei.

Synapto-pHluorin imaging

Synaptic activity of C01/A05 neurons was monitored in *ex vivo* fly brains using UAS-

super-eclipse-synaptopHluorin construct (UAS-*spH*) (Miesenbock, 2012). Adult male

C01/A05 > UAS-*spH* or *C01/A05* > UAS-*spH, kk* flies were housed in normal

behaviour tubes (see behaviour analysis section) and entrained for 3 days in L8: D16

condition at 25°C. Individual flies of either genotype were carefully captured between

ZT9 and ZT11 and fly brains were immediately dissected in HL3 *Drosophila* saline

(70 mM NaCl, 5 mM KCl, 1.5 mM $CaCl_2$, 20 mM $MgCl_2$, 10 mM $NaHCO_3$, 5 mM

Trehalose, 115 mM Sucrose and 5 mM HEPES, pH 7.2) at room temperature. Fly

brains were transferred into 200 µl HL3 in a poly-lysine treated glass bottom dish (35

mm, 627860, Greiner Bio-One) before imaging using an inverted confocal Zeiss LSM

710 microscope (20x objective with maximum pinhole). Three to five image stacks

(16 bits) were taken within two minutes to minimise tissue degradation and to cover

the depth of all spH-positive anatomical regions. Z-projections of the image stacks of

each brain were generated by ImageJ software before the fluorescent intensity of the

indicated neuropil centres was quantified using free drawn ROIs. Background

fluorescence measured by the same ROIs from areas with no brain tissue was then

subtracted to obtain the final fluorescent value. Mean fluorescent values of the

indicated neuropil regions in each hemisphere were calculated and compared between

genotypes. The statistical difference was determined by Mann-Whitney U-test using

Graphpad software.

Bioinformatics

Conservation of amino acid residues between *Drosophila* Neurocalcin and human

Hippocalcin was determined using ClustalW2 software for multiple sequence

alignment. Amino-acid identity and similarity was visualised using BOXSHADE.

Acknowledgements

We thank Jack Humphrey for performing initial work on *Neurocalcin* knockdown

flies, and Kyunghee Koh and Simon Lowe for helpful comments on the manuscript.

This study was supported by the Wellcome Trust (Synaptopathies strategic award

[104033]), and by the MRC [New Investigator Grant MR/P012256/1]. P.K is

supported by a Wellcome Trust Neuroscience PhD studentship.

Competing interests

The authors have no financial or non-financial competing interests.

573   **Figures and Figure legends**

Figure 1

Figure 1. *Neurocalcin* (*Nca*) knockout flies exhibit enhanced night activity and

reduced sleep. (A) Schematic illustration of the procedure used to generate a *Nca*

knockout allele. Homologous arms upstream (Arm 1) and downstream (Arm 2) of the

Nca locus are indicated. Following homologous recombination, the endogenous *Nca*

locus is replaced by a cassette containing the mini-*white* selection marker (red bar),

and attP (blue bar) and loxP sites (yellow bars). The mini-*white* cassette was

subsequently removed via Cre-loxP recombination. (B) Mean sleep levels in 8L: 16D

conditions for Nca^{KO} adult males and *iso31* controls measured using the *Drosophila*

Activity Monitor (DAM). (C-D) Median day (C) and night (D) sleep levels in the

above genotypes. (E) Mean sleep levels in 8L: 16D conditions for Nca^{KO} adult males

and *iso31* controls measured by the video-based *Drosophila* ARousal Tracking

system (DART). (F-G) Median day (F) and night (G) sleep levels in the above

genotypes. Data are presented as Tukey box plots. The 25^{th}, Median, and 75^{th}

percentiles are shown. Whiskers represent 1.5 x the interquartile range. Identical

representations are used in all subsequent box plots. B-D: n = 32 per genotype; E-G: n

= 16. (H-I) The longitudinal movement for individual *iso31* (H) and Nca^{KO} (I) flies

are shown as rows of traces plotting vertical position (Y-axis) over 24 h (X-axis)

under 8L: 16D condition. ***$p < 0.001$, ns - $p > 0.05$, Mann-Whitney U-test.

Figure 1 figure supplement 1

A

```
Hs HPCA    1 MGKQNSKLRPEMLQDLRENTEFSELELQEWYKGFLKDCPTGILNVDEFKKIYANFFPYGD
Dm NCA     1 MGKQNSKLRPEVLEDLKQNTEFTDAEIQEWYKGFLKDCPSGHLSVEEFKKIYGNFFPYGD

Hs HPCA   61 ASKFAEHVFRTFDTNSDGTIDFREFIIALSVTSRGRLEQKLMWAFSMYDLDGNGYISREE
Dm NCA    61 ASKFAEHVFRTFDANGDGTIDFREFLCALSVTSRGKLEQKLKWAFSMYDLDGNGYISRQE

Hs HPCA  121 MLEIVQAIYKMVSSVMKMPEDESTPEKRTEKIFRQMDTNNDGKLSLEEFIRGAKSDPSIV
Dm NCA   121 MLEIVTAIYKMVGSVMKMPEDESTPEKRTDKIFRQMDRNKDGKLSLEEFIEGAKSDPSIV

Hs HPCA  181 RLLQCDPSSASQF
Dm NCA   181 RLLQCDPQSH---
```

B

C

D

E

Figure 1 supplemental figure 1. Generation of the *Nca* knockout allele. (A)

Hippocalcin and Neurocalcin are highly conserved neuronal calcium sensors. Amino-

acid alignment of human (Hs) Hippocalcin (HPCA) and *Drosophila* (Dm)

Neurocalcin (NCA). Identical amino-acids are shaded in black, with functionally

similar amino-acids shaded in grey. Total homology between Hippocalcin and

Neurocalcin is > 90%. (B-C) PCR validation of homologous recombination events.

Correct recombination was verified using primers designed to the attP site and

upstream of the neighbouring locus *cg7646* (B), which will only generate a ~ 3 kb

product following homologous recombination between the targeting vector and the

Nca locus (C). Three independent targeting events (ko1-3) are shown in (C), one of

which was selected for mini-*white* removal as described in Figure 1. WT: wild-type

genome lacking an attP site neighbouring the *cg7646* locus. (D-E) No *Nca* mRNA

was detected in *Nca*[KO] using either standard RT-PCR (D) or quantitative RT-PCR (E;

n = 3 qPCR reactions for *iso31* control and *Nca*[KO] flies).

Figure 1 figure supplement 2

Figure 1 supplemental figure 2. Reduced startle response and elevated night time locomotor activity in *Nca*^{KO} flies. (A) Mean level of locomotor activity (beam crosses) for *iso31* control and *Nca*^{KO} males under 12L: 12D conditions. (B) Median beam breaks during the hour immediately following lights-off (ZT12-13). The rapid light-dark transition initiates a startle response and thus elevated locomotor activity, the degree of which is reduced in *Nca*^{KO} males. (C-D) Median total beam crosses during the day (C) and night (D) periods. Locomotor activity is significantly increased in *Nca*^{KO} males during the night but not the day. n = 32 per genotype. **$p < 0.01$, ***$p < 0.001$, ns – $p > 0.05$, Mann-Whitney U-test.

Figure 1 figure supplement 3

626 **Figure 1 supplemental figure 3.** Night time sleep loss in *Nca*^{KO} flies. (A) Mean sleep

627 level of *iso31* control and *Nca*^{KO} flies under 12L: 12D conditions. (B-C) Median total

628 day (B) and night (C) sleep in *iso31* control and *Nca*^{KO} flies. Night sleep is

629 significantly reduced in *Nca*^{KO} flies compared to controls. n = 32 per genotype.

630 ***p < 0.001, ns – p > 0.05, Mann-Whitney U-test.

Figure 1 figure supplement 4

Figure 1 supplemental figure 4. Reduced startle response and elevated night time

locomotor activity in *Nca*^KO flies under 8L: 16D conditions. (A) Mean level of

locomotor activity (beam crosses) for *iso31* control and *Nca*^KO males. (B) Median

beam breaks during the hour immediately following lights-off (ZT8-9). (C-D) Median

total beam crosses during the normally quiescent period of the night (ZT12-20). n =

16 per genotype. ***p < 0.001, Mann-Whitney U-test.

Figure 1 figure supplement 5

Figure 1 supplemental figure 5. Reduced long sleep bouts in Nca^{KO} flies.

(A) Following detailed fly movement detection using the DART system, individual

sleep bout durations were further estimated using a custom-made R program and

visualised by plotting sleep bout offset against onset for night sleep bouts in control

iso31 and Nca^{KO} adult males under 8L: 16D conditions. In control flies, longer sleep

bouts initiated early during the night (note the black dots deviating from the

diagonal), which are largely absent in Nca^{KO} adult males (red dots). n = 16 for each

genotype. (B) Distribution of sleep bout lengths in Nca^{KO} and control adult males.

Note the significant shift towards shorter sleep bout lengths in Nca^{KO} flies (Nca^{KO} vs.

iso31 control: χ^2, *df*: 85.59, 4, p < 0.001).

Figure 1 figure supplement 6

Figure 1 supplemental figure 6. Detailed tracking reveals increased locomotor

velocity during the night and reduced velocity during the startle response to lights-off

in *Nca*[KO] flies. (A) Mean waking locomotor velocity in *iso31* control and *Nca*[KO] adult

males under L8: 16D condition. (B) Locomotor velocity is significantly reduced in

Nca[KO] flies following the startle response to lights-off (ZT8-ZT9). (C) *Nca*[KO] flies

exhibit a significant increase in locomotor velocity between ZT12-ZT20 compared to

controls – a normally quiescent period. n = 16 per genotype. **p < 0.01, ***p <

0.001, Mann-Whitney U-test.

Figure 2

Figure 2. *Nca* and *Dop1R1* act in a common pathway to regulate night sleep. (A)

Mean sleep levels measured using the DAM system under 8L: 16D conditions for

pan-neuronal *Nca* knockdown (*Nca*[KD]) male adult flies (*elav > kk*) and associated

controls (*elav*-Gal4 driver or *kk* RNAi transgene heterozygotes). (B-C) Median day

and night sleep levels in the above genotypes. n = 54-55. (D) Mean sleep levels

measured using DART system in 8L: 16D conditions for above genotypes. (E-F)

Median day and night sleep levels in the above genotypes. n = 20 per genotype. (G-H)

heterozygosity for the null or strongly hypomorphic Dop1R1 allele *Dop1R1*^{MI03085-}

^{GFST.2} (*D1*/+) suppressed sleep loss in *Nca*^{KD} adult males, but did not alter sleep in

control males ($p > 0.05$, between +/+ and *D1*/+ backgrounds for *elav > +* or *+ > kk*

flies). Mean sleep patterns in 8L: 16D conditions are shown in (G). Median total night

sleep levels are shown in (H). *elav > kk, D1/+*: n = 32; *elav > +, D1/+*: n = 15; *+ >*

kk, D1/+: n = 32; *elav > kk*: n = 48*; elav > +*: n = 47; *+ > kk*: n = 48. **$p < 0.01$,

***$p < 0.001$, ns – $p < 0.05$, Kruskal-Wallis test with Dunn's post-hoc test.

Figure 2 figure supplement 1

Figure 2 supplemental figure 1. Robust knockdown of *Nca* using independent *elav*-

Gal4-driven *Nca* RNAi lines. (A) Schematic showing the *Nca* locus alongside the

neighbouring *cg7646* locus, which shares a common promoter region with *Nca*.

Regions of the *Nca* transcript targeted by the *kk108825, hmj21533* and *jf03398* RNAi

lines (termed *kk, hmj* and *jf* respectively) are depicted by red bars. (B) qPCR

verification of *Nca* knockdown by *kk, hmj* and *jf*. RNAi. Transgene insertions lacking

the *elav*-Gal4 driver were used as controls. (C) Knockdown of *Nca* had no effect on

transcription of the neighbouring *cg7646* locus. Expression levels of *Nca* or *cg7646*

were normalised to the *rp49* control transcript and are displayed as the ratio to the

mean level of the respective RNAi alone controls (+ > *kk*, + > *hmj* or + > *jf*). n = 6 for

all qPCRs (two independent biological repetitions of RNA extraction with triplicated

qPCR reactions for each genotype). **p < 0.01, ns – p > 0.05, Mann-Whitney U-test.

Figure 2 figure supplement 2

Figure 2 supplemental figure 2. Robust night sleep loss in adult male flies

expressing two independent *Nca* RNAi lines *driven by elav*-Gal4. (A-B) Mean sleep

profiles under 8L: 16D conditions for *elav*-gal4 drive*, hmj* (A) or *jf* (B) *Nca* RNAi.

(C) Median night sleep amounts for genotypes shown in (A-B). *elav > +:* n = 32; *+ >*

hmj: n = 26; *elav > hmj*: n = 17; *+ > jf*: n = 32; *elav > jf*: n = 32. **p < 0.01, ***p <

0.001 as compared to driver and RNAi alone controls, Kruskal-Wallis test with

Dunn's post-hoc test.

Figure 2 figure supplement 3

Figure 2 supplemental figure 3. Elevated night time locomotor activity in Nca^{KD}

flies under 8L: 16D conditions, but no alteration in peak locomotor activity. (A) Mean

level of locomotor activity (beam crosses) for Nca^{KD} males (*elav > kk*) and associated

controls. (B) Median beam breaks during the hour immediately following lights-off

(ZT8-9) in the above genotypes. (C) Median total beam crosses during the normally

quiescent period of the night (ZT12-20) in the above genotypes. n = 54-55 per

genotype. ***p < 0.001, ns – p > 0.05, Mann-Whitney U-test.

Figure 2 figure supplement 4

Figure 2 supplemental figure 4. *Nca* knockdown does not alter circadian

rhythmicity. (A) Actograms showing representative individual patterns of locomotor

activity across two weeks under constant dark conditions. (B) Quantification of

locomotor rhythm strength. Robust circadian patterns of locomotor activity were still

observed following in adult males expressing *Nca* RNAi (*kk*) under *elav*-Gal4 relative

to controls. $15 > n > 14$. (C) qPCR analysis of wild type (*Canton-S*) head *Nca*

expression across a 24 h period. No evidence of circadian oscillations in *Nca*

expression was observed. $n = 3$ (triplicated qPCR reactions for each time point).

Expression levels were normalised to the *rp49* control transcript. ns – $p > 0.05$,

Kruskal-Wallis test with Dunn's post-hoc test.

Figure 2 figure supplement 5

Figure 2 supplemental figure 5. Neuronal *Nca* knockdown results in reduced night

sleep in adult virgin female *Drosophila*. (A) Mean sleep patterns of neuronal *Nca*

knockdown females (*elav > kk*) and associated controls under 8L:16D conditions. (B-

C) Median day sleep is unaffected relative to controls (B), whereas median night

sleep is significantly reduced (C). n = 16 per genotype. *p < 0.05, **p < 0.01,

Kruskal-Wallis test with Dunn's post-hoc test.

Figure 2 figure supplement 6

750   **Figure 2 supplemental figure 6.** *Nca* knockdown in muscle cells does not affect

751   sleep in *Drosophila*. (A) Mean sleep patterns of adult male flies with muscle-specific

752   *Nca* knockdown via *mef2*-Gal4 (*mef2 > kk*) and associated controls under 8L: 16D.

753   (B-C) Median day (B) and night (C) sleep levels are unaffected relative to controls. n

754   = 16 per genotype. ns – p > 0.05, Kruskal-Wallis test with Dunn's post-hoc test.

Figure 2 figure supplement 7

Figure 2 supplemental figure 7. Genetic interaction between *Nca* and a second

independent *Dop1R1 allele*. (A) Mean sleep patterns of Nca^{KD} males (*elav > kk*) and a

heterozygous *kk* alone control with and without one copy of the $Dop1R1^{MI04437}$ allele

under 8L: 16D conditions. $Dop1R1^{MI04437}$ is a hypomorphic allele of Dop1R1 that is

homozygous viable, and is therefore weaker compared to the homozygous lethal

$Dop1R1^{MI03085-GFST.2}$ insertion. (B) Median night sleep levels during ZT8-16 are

shown. Heterozygosity for $Dop1R1^{MI04437}$ had no impact on sleep in *kk* heterozygote

controls but partially rescued night sleep during ZT8-16 in Nca^{KD} males. *elav > kk*: n

= 64; *+ > kk*: n = 64; *elav > kk*; $Dop1R1^{MI04437}$/+: n = 39; *+ > kk*; $Dop1R1^{MI04437}$/+: n =

45. *p < 0.05, ***p < 0.001, Kruskal-Wallis test with Dunn's post-hoc test.

Figure 2 figure supplement 8

Figure 2 supplemental figure 8. NCA regulate sleeps in a distinct pathway to the

dNR1 NMDA receptor. (A) Mean sleep patterns of Nca^{KD} males ($elav > kk$) and

heterozygous controls with and without one copy of the $dNR1^{MI11796}$ ($dNR1/+$) allele

in 8L: 16D conditions are shown. (B) Median night sleep levels in the above

genotypes. Heterozygosity for $dNR1^{MI11796}$ resulted in sleep loss in $elav$-Gal4 driver

controls, and reduced sleep further in an additive manner in $elav > kk$ Nca^{KD} males,

suggesting that NCA and dNR1 act in separate pathways to promote night sleep. *Elav*

> kk; *dNR1/+*: n = 32; *elav > +*; *dNR1/+*: n = 26; *+ > kk*; *dNR1/+*: n = 38; *elav > kk*:

n = 32; *elav > +*: n = 30; *+ > kk*: n = 34. ns – p > 0.05, *p < 0.05, **p < 0.01, ***p <

0.001, Kruskal-Wallis test with Dunn's post-hoc test.

Figure 3

Figure 3. NCA acts in a two distinct neural networks to regulate night sleep. (A)

Transgenic RNAi-based mini-screen to identify key NCA-expressing neurons. NCA

knockdown with broadly expressed drivers resulted in reduced night sleep in adult

males under 8L: 16D conditions. In contrast, NCA knockdown in previously defined

sleep-regulatory centers, clock neurons, the visual system or subsets of Dop1R1-

expressing neurons did not impact night sleep. FSB: fan-shaped body. MB:

mushroom body. Grey and blue box plots: control lines. Red box plots: experimental

lines showing reduced night sleep relative to controls. Green box plots: experimental

lines failing to show reduced night sleep relative to one or both controls. See Figure 3

figure supplement 1 for n-values and statistical comparisons. (B) Confocal z-stacks of

adult male brains expressing genetically-encoded fluorophores under the *A05* or *C01*-

Gal4 drivers. Neuropil regions are labelled with anti-Bruchpilot (BRP). Scale bar =

100 μm. Arrows point to neuropil centers. AOTU: anterior optic tubercle. MBNs:

mushroom body neurons. OL: optic lobe. AMMC: antennal mechanosensory and

motor center. AVP: anterior ventrallateral protocerebrum. SMP: supra-medial

protocerebrum. (C-D) *Nca* knockdown in both *A05* and *C01*-neurons recapitulates the

effect of pan-neuronal *Nca* knockdown, whereas *Nca* knockdown in either neuronal

subpopulation alone did not reduce sleep relative to controls. Mean sleep patterns in

8L: 16D conditions are shown in (C). Median night sleep levels are shown in (D). + >

kk: n = 80; *C01* > +: n = 64, *C01* > *kk*: n = 80; *A05* > +: n = 31; *A05* > *kk*: n = 31;

C01/A05 > +: n = 42; *C01/A05* > *kk*: n = 71. *p < 0.05, **p < 0.01, ***p < 0.001, ns

– p > 0.05 compared to driver and RNAi alone controls, Kruskal-Wallis test with

Dunn's post-hoc test.

Figure 3 figure supplement 1. Night sleep levels in flies with *kk Nca RNAi* driven by neuron sub-type-specific Gal4 drivers.

Genotype	n-values	25% Percentile (min)	Median (min)	75% Percentile (min)	Mean (min)	Std. Deviation (min)	Dunn's test vs +>kk	vs Gal4 >+
+>kk	275	602	671	742	668	110	-	-
elav>+	86	518	633	698	604	133	-	-
elav>kk	99	306	428	530	413	168	***	***
nsyb>+	40	491	576	654	565	121	-	-
nsyb>kk	39	248	409	490	388	196	***	*
inc>+	14	655	684	705	688	56	-	-
inc>kk	30	460	521	637	534	128	***	**
tim>+	16	577	607	707	636	91	-	-
tim>kk	16	668	709	767	707	80	ns	ns
tug>+	16	590	658	688	650	117	-	-
tug>kk	15	580	701	805	670	134	ns	ns
ple>+	32	556	663	718	634	132	-	-
ple>kk	30	545	670	718	642	101	ns	ns
Ddc>+	16	616	658	721	662	63	-	-
Ddc>kk	16	538	608	667	610	84	ns	ns
Chat>+	16	601	656	758	677	111	-	-
Chat>kk	14	607	704	766	672	140	ns	ns
vGlut>+	16	536	632	660	607	91	-	-
vGlut>kk	24	495	594	679	586	111	ns	ns
GAD>+	16	662	703	752	689	82	-	-
GAD>kk	16	705	758	817	739	100	ns	ns
Trh.1>+	16	598	635	696	640	83	-	-
Trh.1>kk	16	617	649	720	668	71	ns	ns
Tdc2>+	16	659	715	742	682	105	-	-
Tdc2>kk	15	657	733	794	695	128	ns	ns
C5>+	40	509	568	648	583	103	-	-
C5>kk	49	513	567	665	581	107	***	ns
R23E10>+	16	651	670	705	680	48	-	-
R23E10>kk	16	535	632	712	621	122	ns	ns
R55B01>+	16	621	653	714	652	107	-	-
R55B01>kk	16	618	670	721	680	80	ns	ns
OK107>+	23	644	702	777	695	94	-	-

OK107>kk	32	601	668	751	664	106	ns	ns
nompC>+	16	708	769	842	768	76	-	-
nompC>kk	16	638	753	804	725	92	ns	ns
ppk>+	16	661	701	742	706	62	-	-
ppk>kk	16	712	766	798	760	78	ns	ns
trpACD>+	16	586	639	677	630	62	-	-
trpACD>kk	15	702	764	825	758	75	ns	*
R52H12>+	16	536	654	689	629	92	-	-
R52H12>kk	32	525	603	684	581	155	ns	ns
GMR>+	16	582	649	685	640	73	-	-
GMR>kk	14	592	656	714	652	79	ns	ns
Hdc>+	16	527	599	694	575	145	-	-
Hdc>kk	16	566	641	686	624	86	ns	ns
R21G01>+	15	584	646	694	631	75	-	-
R21G01>kk	16	440	489	598	519	109	**	ns
R72B05>+	15	341	388	502	397	146	-	-
R72B05>kk	16	611	706	795	710	107	ns	***
R72B07>+	13	570	668	697	645	66	-	-
R72B07>kk	14	492	541	585	545	69	**	ns
R72B08>+	16	452	595	643	561	129	-	-
R72B08>kk	32	348	436	591	474	159	***	ns
R72B11>+	16	403	515	651	513	123	-	-
R72B11>kk	15	512	580	645	583	88	ns	ns
R72C01>+	48	528	590	662	591	84	-	-
R72C01>kk	64	496	582	639	569	104	***	ns
R72C02>+	16	500	551	595	544	62	-	-
R72C02>kk	16	591	627	694	643	60	ns	ns

- Not significant, ns-p>0.05, *p < 0.05, **p < 0.01, ***p < 0.001, Kruskal-Wallis test with Dunn's

post-hoc test.

Figure 3 figure supplement 2

Figure 3 supplemental figure 2. *C01*-Gal4 and *A05*-Gal4 label small subsets of

neurons in the adult fly brain. (A, C) Representative confocal z-stacks of adult male

brains expressing nuclear RFP marker (Red-stinger) under *A05*-Gal4 (A) or *C01*-Gal4

driver (C). DAPI-staining labels nuclei within the *Drosophila* brain. (B, D) Number

of cells expressing Red-stinger driven by *A05*- (B) or *C01*-Gal4 (D). n = 3 for each

genotype. Scale bar =100 μm.

Figure 3 figure supplement 3

Figure 3 supplemental figure 3. (A) Mean sleep patterns of adult males homozygous

for the *C01*-Gal4 driver with and without the *kk Nca* RNAi insertion. (B) Mean sleep

patterns of adult males homozygous for the *A05*-Gal4 driver with and without the *kk*

Nca RNAi insertion. (C) Median night sleep levels for heterozygous RNAi transgene

and homozygous driver controls, and males expressing *Nca* RNAi with two Gal4

driver copies. No night sleep loss was observed using two copies of either driver

relative to controls. *+ > kk*: n = 80 (the same population was used in Figure 3C, as the

experiments were performed side by side), *C01/C01 > +*: n = 24, *C01/C01 > kk*: n =

39; *A05/A05 >* : n = 22, *A05/A05 > kk*: n = 23. ns – p > 0.05, Kruskal-Wallis test with

Dunn's post-hoc test.

Figure 3 figure supplement 4

Figure 3 supplemental figure 4. Knockdown of *Nca* in mushroom body and *A05*-

neurons results in partial night sleep loss. (A-B) *Nca* knockdown in both *A05* and

MB-neurons (defined by *ok107*-Gal4; *ok107*) results in reduced night sleep (also see

Figure 3A showing *ok107 > kk* alone does not cause night sleep loss). Mean sleep

patterns in 8L: 16D conditions are shown in (A). Median night sleep levels are shown

in (B). *+ > kk*: n = 31; *A05/ok107 > +*: n = 33; *A05/ok107 > kk*: n = 42. *p < 0.05, **p

< 0.01, Kruskal-Wallis test with Dunn's post-hoc test.

Figure 4

Figure 4. NCA inhibits synaptic release in subsets of *C01/A05* neurons. Expression of

a fluorescent marker for synaptic release (synapto-pHluorin, spH) in *C01*- and *A05*-

positive neurons in control (*C01/A05 > spH*) or *Nca* knockdown (*C01/A05 > spH, kk*)

backgrounds. Adult brains were dissected and imaged between ZT9-ZT11, when

robust sleep loss occurs in *C01/A05 > kk* flies (see Figure 3C). (A, C, E, G)

Representative pseudo-coloured images for α/β mushroom body lobe Kenyon cells

(α/β MB, A; α/β lobes are outlined in white), the MB γ-lobe region (C), AMMC (E)

and SMP (G) regions in each genotype. Fluorescent intensity range is shown in the

horizontal bar, illustrating minimum to maximum spectrum. Scale bars = 10 μm. (B,

D, F, H) Mean fluorescent intensity for individual hemispheres in the above regions. n

= 22-24. *$p < 0.05$, ***$p < 0.001$, Mann-Whitney U-test.

Figure 5

Figure 5. Sleep loss in *Nca* knockdown flies is caused by enhanced excitability of

C01/A05 neurons. (A) Experimental paradigm for acute activation of *A05* or *C01*-

neurons. 22°C: non-activating temperature for TrpA1. 27°C: activating temperature.

Sleep measurements were measured over two days in 8L: 16D conditions. (B-C)

Mean sleep levels across 8L: 16D following expression of TrpA1 in *A05*-, *C01*- or

A05- and *C01*-neurons (and associated controls) at 22°C (B) or 27°C (C). (D) Median

change in night sleep levels (Δ night sleep) following the shift from 22°C on day 1 to

27°C on day 2. + > *TrpA1*: n = 53; *A05* > +: n = 23; *A05* > *TrpA1*: n = 68; *C01* > +: n

= 24; *C01* > *TrpA1*: n = 40; *C01/A05* > +: n = 33; *C01/A05* > *TrpA1*: n = 40. ***p <

0.001, ns – p > 0.05, as compared to *TrpA1* or driver alone controls by Kruskal-Wallis

test with Dunn's post-hoc test (for *C01, A05* or *C01/ A05* > *TrpA1* compared to

controls) or Mann-Whitney U-test (for *C01/A05* > *TrpA1* compared to *C01* > *TrpA1*).

(E-F) Inhibition of *C01/A05* neurons by expressing dORKΔC2 rescues sleep loss due

to *Nca* knockdown in *C01/A05* neurons, while expression of dORKΔC2 does not

change base line sleep. Mean sleep patterns in 8L: 16D conditions are shown in (E).

Median night sleep levels are shown in (F). + > *kk*: n = 47, *C01/A05* > +: n = 47,

C01/A05 > *kk*: n = 62, *C01/A05* > *dORKΔC2, kk*: n = 50, *C01/A05* > *dORKΔC2*: n =

39, + > *dORKΔC2*: n = 30. *p < 0.05, ***p < 0.001, ns – p > 0.05, Kruskal-Wallis

test with Dunn's post-hoc test.

References

Afonso, D.J., Liu, D., Machado, D.R., Pan, H., Jepson, J.E., Rogulja, D., and Koh, K.

(2015). TARANIS Functions with Cyclin A and Cdk1 in a Novel Arousal Center to

Control Sleep in *Drosophila*. Curr Biol *25*, 1717-1726.

Andrade, R., Foehring, R.C., and Tzingounis, A.V. (2012). The calcium-activated

slow AHP: cutting through the Gordian knot. Front Cell Neurosci *6*, 47.

Baena-Lopez, L.A., Alexandre, C., Mitchell, A., Pasakarnis, L., and Vincent, J.P.

(2013). Accelerated homologous recombination and subsequent genome modification

in *Drosophila*. Development *140*, 4818-4825.

Braunewell, K.H., and Klein-Szanto, A.J. (2009). Visinin-like proteins (VSNLs):

interaction partners and emerging functions in signal transduction of a subfamily of

neuronal Ca2+ -sensor proteins. Cell Tissue Res *335*, 301-316.

Breakefield, X.O., Blood, A.J., Li, Y., Hallett, M., Hanson, P.I., and Standaert, D.G.

(2008). The pathophysiological basis of dystonias. Nat Rev Neurosci *9*, 222-234.

Burgoyne, R.D., and Haynes, L.P. (2012). Understanding the physiological roles of

the neuronal calcium sensor proteins. Mol Brain *5*, 2.

Calabresi, P., Pisani, A., Rothwell, J., Ghiglieri, V., Obeso, J.A., and Picconi, B.

(2016). Hyperkinetic disorders and loss of synaptic downscaling. Nat Neurosci *19*,

868-875.

Charlesworth, G., Angelova, P.R., Bartolome-Robledo, F., Ryten, M., Trabzuni, D.,

Stamelou, M., Abramov, A.Y., Bhatia, K.P., and Wood, N.W. (2015). Mutations in

HPCA cause autosomal-recessive primary isolated dystonia. Am J Hum Genet *96*,

657-665.

Charlesworth, G., Plagnol, V., Holmstrom, K.M., Bras, J., Sheerin, U.M., Preza, E.,

Rubio-Agusti, I., Ryten, M., Schneider, S.A., Stamelou, M.*, et al.* (2012). Mutations

in ANO3 cause dominant craniocervical dystonia: ion channel implicated in

pathogenesis. Am J Hum Genet *91*, 1041-1050.

Donlea, J.M., Thimgan, M.S., Suzuki, Y., Gottschalk, L., and Shaw, P.J. (2011).

Inducing sleep by remote control facilitates memory consolidation in *Drosophila*.

Science *332*, 1571-1576.

Fahn, S. (1988). Concept and classification of dystonia. Adv Neurol *50*, 1-8.

Faville, R., Kottler, B., Goodhill, G.J., Shaw, P.J., and van Swinderen, B. (2015).

How deeply does your mutant sleep? Probing arousal to better understand sleep

defects in *Drosophila*. Sci Rep *5*, 8454.

Fuchs, T., Gavarini, S., Saunders-Pullman, R., Raymond, D., Ehrlich, M.E.,

Bressman, S.B., and Ozelius, L.J. (2009). Mutations in the THAP1 gene are

responsible for DYT6 primary torsion dystonia. Nat Genet *41*, 286-288.

Fuchs, T., Saunders-Pullman, R., Masuho, I., Luciano, M.S., Raymond, D., Factor, S.,

Lang, A.E., Liang, T.W., Trosch, R.M., White, S.*, et al.* (2013). Mutations in GNAL

cause primary torsion dystonia. Nat Genet *45*, 88-92.

Gerfen, C.R., and Surmeier, D.J. (2011). Modulation of striatal projection systems by

dopamine. Annu Rev Neurosci *34*, 441-466.

Hamada, F.N., Rosenzweig, M., Kang, K., Pulver, S.R., Ghezzi, A., Jegla, T.J., and

Garrity, P.A. (2008). An internal thermal sensor controlling temperature preference in

Drosophila. Nature *454*, 217-220.

Helassa, N., Antonyuk, S.V., Lian, L.Y., Haynes, L.P., and Burgoyne, R.D. (2017).

Biophysical and functional characterization of hippocalcin mutants responsible for

human dystonia. Hum Mol Genet *26*: 2426-2435.

Jenett, A., Rubin, G.M., Ngo, T.T., Shepherd, D., Murphy, C., Dionne, H., Pfeiffer,

B.D., Cavallaro, A., Hall, D., Jeter, J.*, et al.* (2012). A GAL4-driver line resource for

Drosophila neurobiology. Cell Rep *2*, 991-1001.

Jiang, Y., Pitmon, E., Berry, J., Wolf, F.W., McKenzie, Z., and Lebestky, T.J. (2016).

A Genetic Screen To Assess Dopamine Receptor (DopR1) Dependent Sleep

Regulation in *Drosophila*. G3 (Bethesda) *6*, 4217-4226.

Jo, J., Son, G.H., Winters, B.L., Kim, M.J., Whitcomb, D.J., Dickinson, B.A., Lee,

Y.B., Futai, K., Amici, M., Sheng, M.*, et al.* (2010). Muscarinic receptors induce

LTD of NMDAR EPSCs via a mechanism involving hippocalcin, AP2 and PSD-95.

Nat Neurosci *13*, 1216-1224.

Joiner, W.J., Crocker, A., White, B.H., and Sehgal, A. (2006). Sleep in *Drosophila* is

regulated by adult mushroom bodies. Nature *441*, 757-760.

Kamikouchi, A., Inagaki, H.K., Effertz, T., Hendrich, O., Fiala, A., Göpfert, M.C.,

and Ito, K. (2009). The neural basis of *Drosophila* gravity-sensing and hearing.

Nature *458*, 165-171.

Karimi, M., and Perlmutter, J.S. (2015). The role of dopamine and dopaminergic

pathways in dystonia: insights from neuroimaging. Tremor Other Hyperkinet Mov (N

Y) *5*, 280.

Kume, K., Kume, S., Park, S.K., Hirsh, J., and Jackson, F.R. (2005). Dopamine is a

regulator of arousal in the fruit fly. J Neurosci *25*, 7377-7384.

Lamaze, A., Ozturk-Colak, A., Fischer, R., Peschel, N., Koh, K., and Jepson, J.E.

(2017). Regulation of sleep plasticity by a thermo-sensitive circuit in *Drosophila*. Sci

Rep *7*, 40304.

Lebestky, T., Chang, J.S., Dankert, H., Zelnik, L., Kim, Y.C., Han, K.A., Wolf, F.W.,

Perona, P., and Anderson, D.J. (2009). Two different forms of arousal in *Drosophila*

are oppositely regulated by the dopamine D1 receptor ortholog DopR via distinct

neural circuits. Neuron *64*, 522-536.

Lee, H.J., Weitz, A.J., Bernal-Casas, D., Duffy, B.A., Choy, M., Kravitz, A.V.,

Kreitzer, A.C., and Lee, J.H. (2016). Activation of Direct and Indirect Pathway

Medium Spiny Neurons Drives Distinct Brain-wide Responses. Neuron *91*, 412-424.

Lehner, B. (2013). Genotype to phenotype: lessons from model organisms for human

genetics. Nat Rev Genet *14*, 168-178.

Levine, J.D., Funes, P., Dowse, H.B., and Hall, J.C. (2002a). Advanced analysis of a

cryptochrome mutation's effects on the robustness and phase of molecular cycles in

isolated peripheral tissues of *Drosophila*. BMC neuroscience *3*, 5.

Levine, J.D., Funes, P., Dowse, H.B., and Hall, J.C. (2002b). Signal analysis of

behavioral and molecular cycles. BMC neuroscience *3*, 1.

Li, Q., Kellner, D.A., Hatch, H.A.M., Yumita, T., Sanchez, S., Machold, R.P., Frank,

C.A., and Stavropoulos, N. (2017). Conserved properties of *Drosophila insomniac*

link sleep regulation and synaptic function. PLoS Genet *13*, e1006815.

Liu, Q., Liu, S., Kodama, L., Driscoll, M.R., and Wu, M.N. (2012). Two

dopaminergic neurons signal to the dorsal fan-shaped body to promote wakefulness in

Drosophila. Curr Biol *22*, 2114-2123.

Liu, S., Lamaze, A., Liu, Q., Tabuchi, M., Yang, Y., Fowler, M., Bharadwaj, R.,

Zhang, J., Bedont, J., Blackshaw, S*., et al.* (2014). WIDE AWAKE mediates the

circadian timing of sleep onset. Neuron *82*, 151-166.

McGary, K.L., Park, T.J., Woods, J.O., Cha, H.J., Wallingford, J.B., and Marcotte,

E.M. (2010). Systematic discovery of nonobvious human disease models through

orthologous phenotypes. Proc Natl Acad Sci U S A *107*, 6544-6549.

Mencacci, N.E., Rubio-Agusti, I., Zdebik, A., Asmus, F., Ludtmann, M.H., Ryten,

M., Plagnol, V., Hauser, A.K., Bandres-Ciga, S., Bettencourt, C.*, et al.* (2015). A

missense mutation in KCTD17 causes autosomal dominant myoclonus-dystonia. Am

J Hum Genet *96*, 938-947.

Miesenbock, G. (2012). Synapto-pHluorins: genetically encoded reporters of synaptic

transmission. Cold Spring Harb Protoc *2012*, 213-217.

Nitabach, M.N., Blau, J., and Holmes, T.C. (2002). Electrical silencing of *Drosophila*

pacemaker neurons stops the free-running circadian clock. Cell *109*, 485-495.

Pappas, S.S., Darr, K., Holley, S.M., Cepeda, C., Mabrouk, O.S., Wong, J.M., LeWitt,

T.M., Paudel, R., Houlden, H., Kennedy, R.T.*, et al.* (2015). Forebrain deletion of the

dystonia protein torsinA causes dystonic-like movements and loss of striatal

cholinergic neurons. Elife *4*, e08352.

Park, D., and Griffith, L.C. (2006). Electrophysiological and anatomical

characterization of PDF-positive clock neurons in the intact adult *Drosophila* brain. J

Neurophysiol *95*, 3955-3960.

Peterson, D.A., Sejnowski, T.J., and Poizner, H. (2010). Convergent evidence for

abnormal striatal synaptic plasticity in dystonia. Neurobiol Dis *37*, 558-573.

Pfeiffenberger, C., and Allada, R. (2012). Cul3 and the BTB adaptor insomniac are

key regulators of sleep homeostasis and a dopamine arousal pathway in *Drosophila*.

PLoS Genet *8*, e1003003.

Pfeiffenberger, C., Lear, B.C., Keegan, K.P., and Allada, R. (2010a). Locomotor

activity level monitoring using the *Drosophila* Activity Monitoring (DAM) System.

Cold Spring Harb Protoc *2010*, pdb prot5518.

Pfeiffenberger, C., Lear, B.C., Keegan, K.P., and Allada, R. (2010b). Processing sleep

data created with the *Drosophila* Activity Monitoring (DAM) System. Cold Spring

Harb Protoc *2010*, pdb prot5520.

Pitman, J.L., McGill, J.J., Keegan, K.P., and Allada, R. (2006). A dynamic role for

the mushroom bodies in promoting sleep in *Drosophila*. Nature *441*, 753-756.

Rogulja, D., and Young, M.W. (2012). Control of sleep by cyclin A and its regulator.

Science *335*, 1617-1621.

Seidner, G., Robinson, J.E., Wu, M., Worden, K., Masek, P., Roberts, S.W., Keene,

A.C., and Joiner, W.J. (2015). Identification of Neurons with a Privileged Role in

Sleep Homeostasis in *Drosophila melanogaster*. Curr Biol *25*, 2928-2938.

Shi, M., Yue, Z., Kuryatov, A., Lindstrom, J.M., and Sehgal, A. (2014). Identification

of Redeye, a new sleep-regulating protein whose expression is modulated by sleep

amount. Elife *3*, e01473.

Sitaraman, D., Aso, Y., Jin, X., Chen, N., Felix, M., Rubin, G.M., and Nitabach, M.N.

(2015). Propagation of Homeostatic Sleep Signals by Segregated Synaptic

Microcircuits of the *Drosophila* Mushroom Body. Curr Biol *25*, 2915-2927.

Stavropoulos, N., and Young, M.W. (2011). insomniac and Cullin-3 regulate sleep

and wakefulness in Drosophila. Neuron *72*, 964-976.

Tanabe, L.M., Kim, C.E., Alagem, N., and Dauer, W.T. (2009). Primary dystonia:

molecules and mechanisms. Nat Rev Neurol *5*, 598-609.

Tecuapetla, F., Jin, X., Lima, S.Q., and Costa, R.M. (2016). Complementary

Contributions of Striatal Projection Pathways to Action Initiation and Execution. Cell

166, 703-715.

Teng, D.H., Chen, C.K., and Hurley, J.B. (1994). A highly conserved homologue of

bovine neurocalcin in *Drosophila melanogaster* is a Ca(2+)-binding protein expressed

in neuronal tissues. J Biol Chem *269*, 31900-31907.

Tomita, J., Ban, G., and Kume, K. (2017). Genes and neural circuits for sleep of the

fruit fly. Neurosci Res *118*, 82-91.

Tomita, J., Ueno, T., Mitsuyoshi, M., Kume, S., and Kume, K. (2015). The NMDA

Receptor Promotes Sleep in the Fruit Fly, *Drosophila melanogaster*. PLoS One *10*,

e0128101.

Tzingounis, A.V., Kobayashi, M., Takamatsu, K., and Nicoll, R.A. (2007).

Hippocalcin gates the calcium activation of the slow afterhyperpolarization in

hippocampal pyramidal cells. Neuron *53*, 487-493.

Ueno, K., Naganos, S., Hirano, Y., Horiuchi, J., and Saitoe, M. (2013). Long-term

enhancement of synaptic transmission between antennal lobe and mushroom body in

cultured *Drosophila* brain. J Physiol *591*, 287-302.

Ueno, T., Tomita, J., Tanimoto, H., Endo, K., Ito, K., Kume, S., and Kume, K. (2012).

Identification of a dopamine pathway that regulates sleep and arousal in *Drosophila*.

Nat Neurosci *15*, 1516-1523.

Wakabayashi-Ito, N., Ajjuri, R.R., Henderson, B.W., Doherty, O.M., Breakefield,

X.O., O'Donnell, J.M., and Ito, N. (2015). Mutant human torsinA, responsible for

early-onset dystonia, dominantly suppresses GTPCH expression, dopamine levels and

locomotion in *Drosophila melanogaster*. Biol Open *4*, 585-595.

Wakabayashi-Ito, N., Doherty, O.M., Moriyama, H., Breakefield, X.O., Gusella, J.F.,

O'Donnell, J.M., and Ito, N. (2011). Dtorsin, the *Drosophila* ortholog of the early-

onset dystonia TOR1A (DYT1), plays a novel role in dopamine metabolism. PLoS

One *6*, e26183.

Weisheit, C.E., and Dauer, W.T. (2015). A novel conditional knock-in approach

defines molecular and circuit effects of the DYT1 dystonia mutation. Hum Mol Genet

24, 6459-6472.

Wenning, G.K., Kiechl, S., Seppi, K., Muller, J., Hogl, B., Saletu, M., Rungger, G.,

Gasperi, A., Willeit, J., and Poewe, W. (2005). Prevalence of movement disorders in

men and women aged 50-89 years (Bruneck Study cohort): a population-based study.

Lancet Neurol *4*, 815-820.

Wu, M., Robinson, J.E., and Joiner, W.J. (2014). SLEEPLESS is a bifunctional

regulator of excitability and cholinergic synaptic transmission. Curr Biol *24*, 621-629.

Yokoi, F., Dang, M.T., Liu, J., Gandre, J.R., Kwon, K., Yuen, R., and Li, Y. (2015).

Decreased dopamine receptor 1 activity and impaired motor-skill transfer in Dyt1

DeltaGAG heterozygous knock-in mice. Behav Brain Res *279*, 202-210.

Zhong, L., Bellemer, A., Yan, H., Ken, H., Jessica, R., Hwang, R.Y., Pitt, G.S., and

Tracey, W.D. (2012). Thermosensory and nonthermosensory isoforms of *Drosophila*

melanogaster TRPA1 reveal heat-sensor domains of a thermoTRP Channel. Cell Rep

1, 43-55.

1080

1081