Iron-dependent essential genes in *Salmonella* Typhimurium

Sardar Karash[1,2] and Young Min Kwon[1,3*]

[1] Cell and Molecular Biology Program, University of Arkansas, Fayetteville, Arkansas

[2] Department of Biology, College of Education, Salahaddin University, Erbil, Kurdistan, Iraq

[3] Department of Poultry Science, University of Arkansas, Fayetteville, AR 72701

*Corresponding author:

Email : ykwon@uark.edu (YK)

Abstract

Background: The molecular mechanisms underlying bacterial cell death due to stresses or bactericidal antibiotics are complex and remain puzzling. Previously, it was shown that iron is required for effective killing of bacterial cells by numerous bactericidal antibiotics. Here, our high-resolution Tn-seq analysis demonstrated that transposon mutants of *S*. Typhimurium with insertions in essential genes escaped immediate killing or growth inhibition under iron-restricted conditions for approximately one-third of all essential genes. **Results:** We grouped essential genes into two categories, iron-dependent and iron-independent essential genes. The iron-dependency of the iron-dependent essential genes was further validated by the fact that the relative abundance of these essential gene mutants increased further with more severe iron restrictions. Our unexpected observation can be explained well by the recently proposed common killing mechanisms of bactericidal antibiotics via production of reactive oxygen species (ROS). In this model iron restriction would inhibit production of ROS, leading to reduced killing activity following blocking of an essential function. Interestingly, the targets of most antibiotics currently in use clinically, whether bacteriostatic or bactericidal, are iron-dependent essential genes. **Conclusions:** Our result suggests that targeting iron-independent essential genes may be a better strategy for future antibiotic development, because blocking these genes would lead to immediate cell death regardless of iron concentration. On the contrary, blocking iron-dependent pathways under iron limited *in vivo* environment could lead to reduced killing action, which might increase drug-resistance by mutagenic action of sublethal concentrations of ROS. This work expands our knowledge on the role of iron to a broader range of essential pathways, and provides novel insights for development of more effective antibiotics.

Keywords: Essential genes, Iron-dependent essential genes, ROS, Tn-seq, Antibiotic targets, *Salmonella* Typhimurium

Background

Essential genes encode the proteins that are essentially required for cell viability or growth. These genes have been exploited as pivotal targets for antibacterial drugs, because blocking their proteins cause cell impairment and ultimately growth inhibition or death of bacterial cells. Thus, nearly all antibiotics in clinical use target these essential pathways. However, for many natural antibiotics, the molecular targets remain unknown [1] and even if the target is known, in case of bactericidal antibiotics, the cellular events that follow in response to disruption of essential pathways leading to bacterial cell death remain puzzling.

Numerous studies have shown the role of reactive oxygen species (ROS) in cell death for eukaryotes as well as prokaryotes. In eukaryotes, apoptosis and necroptosis are associated with ROS [2, 3]. Ferroptosis is an iron-dependent nonapoptotic form of oxidative cell death in mammalian cancer cells. These cells die as a result of ROS accumulation and the death can be prevented via iron chelators [4]. In bacteria, contribution of ROS to cell death due to bactericidal antibiotics is supported by numerous studies. Kohanski et al. [5] proposed that bactericidal antibiotics, regardless of their molecular targets, induce production of ROS, which consequently damages biomolecules contributing to cell death, and demonstrated the death process can be mitigated via iron chelators. This model asserts that upon antibiotic-target interactions, consecutive specific intracellular events induce ROS formation, specifically hydroxyl radical, via Fenton reaction through the process that involves TCA cycle-NADH depletion and destabilization of Fe-S clusters [5, 6]. Furthermore, it was also shown that ROS generation elevates in bacterial cells by the attack of competitor bacteria or P1vir phage via type VI secretion system [7]. In addition, mammalian peptidoglycan recognition protein-induced bacterial killing requires ROS and the lethality of this protein can be inhibited via an iron chelator [8]. Immune cells also produce ROS to kill bacterial pathogens [9]. However, despite these numerous evidences on the role of ROS in bacterial

cell death, it is unknown if this role of ROS can be generalized to all death process in

bacterial cells, and if not, what the scope of the ROS-mediated cellular death process is.

A pathogenic bacterium can possess a few hundred essential genes that are indispensable for

maintaining cell viability. Empirically, essential genes are defined by the genes that when

inactivated lead to loss of cell viability. In *E. coli* Keio collection, single-gene deletions were

made for all known open reading frames, excluding 302 genes which could not tolerate

disruptions and these 302 genes were considered essential [10, 11]. On the other hand,

random genome-wide transposon mutagenesis coupled with next generation sequencing (Tn-

seq) is a powerful method to identify essential genes [12]. Tn-seq experiments have shown

that there are 353 essential genes in *Salmonella* Typhimurium SL326 [13]; 461 in

Mycobacterium tuberculosis H37Rv [14]; and 227 in *Streptococcus pyogenes* [15]. Recently,

a team chemically synthesized *Mycoplasma mycoid* JCVI-syn3.0 based on 473 essential

genes [16]. Clustered Regularly Interspaced Short Palindromic Repeats Interference

(CRISPRi) was employed for phenotypic analysis of 289 essential genes in *Bacillus subtilis*

that were identified by Tn-seq, and confirmed that approximately 94% of the putative

essential genes were genuine essential genes [17].

Nearly all studies on defining essential genomes in bacteria have been conducted using

stress-free nutrient-rich media for the given bacterial species under the assumption that a

minimum set of the core essential genes would be best revealed under such "optimal" growth

conditions. In this study, on the contrary, we analyzed our Tn-seq data to determine essential

genes in *S.* Typhimurium under the stress conditions created by restricting iron

concentrations using iron chelator 2,2`-Dipyridyl (Dip) ranging from 0 to 400 μM. Our initial

focus was to identify conditionally essential genes required for fitness under iron-restriction

conditions. However, we unexpectedly found that a considerable portion of the genes that are

categorized as essential genes in LB media (no Dip) were rendered non-essential under iron-

restriction conditions. Furthermore, the relative abundance of the transposon mutants in those

essential genes increased with the increasing severity of iron restrictions. We reason that this

finding has significant implications for the current efforts to overcome the crisis in public

health due to increasing antibiotic resistance, and may provide valuable insights for future

direction for the development of new antibiotics with inherently reduced chance of

developing drug resistance. Therefore, in this study we focus on the analysis of the essential

genes under iron-restricted conditions, and discuss the implications of our discovery.

Results and Discussion

Library selection and initial Tn-seq analysis

We constructed two genome-saturating Tn5 transposon libraries (Library–A and –AB) in

which 92.6% of all ORFs had insertions (Table 1). To track the relative abundance of mutants

in the libraries in response to iron restriction, appropriate library was inoculated into LB

media supplemented with iron chelator 2,2`-Dipyridyl (Dip) at final concentrations of 0

(controls: LB-I, LB-II and LB-III), 100 (Dip100), 150 (Dip150), 250 (Dip250-I and Dip250-

II), or 400 µM (Dip400) as shown in Fig. 1. The cultures were grown until the bacteria

reached mid-log phase. Since the growth rate of *S.* Typhimurium was reduced by Dip

treatment in a concentration-dependent manner (Fig. 2), the cultures were incubated for

different lengths of time to reach the mid-log phase (Table 1). We also included Library-A

itself (LB-I) for Tn-seq analysis as an additional control group (Fig. 1). From Library-AB, we

obtained 273 million (M) sequence reads from Tn5 genomic junctions in the chromosome of

S. Typhimurium for all conditions combined, and 185 M sequence reads were mapped to the

completed genome of *S.* Typhimurium 14028 (Table 2).

The accurate genome-mapping based on long Tn5 genomic junctions and high number of

read counts allowed us to define essentiality and conditional essentiality of the genes with a

high precision. Our initial goal in this study was to elucidate the conditionally essential genes

that are required for fitness under different levels of iron restriction using Tn-seq. During the

data analysis, however, we found the read counts corresponding to the mutants in numerous

known essential genes increased significantly and consistently with iron restriction. This

observation was in contrary to the currently accepted working definition of essential genes

that cannot tolerate disruptions. It required further detailed analysis before we could accept

this interesting, yet unexpected finding. Therefore, we have conducted a systematic analysis

for essential genes and comparatively analyzed the results between iron-replete and iron-

restricted conditions.

Essential genome of *S*. Typhimurium in iron-replete and iron-restricted conditions
116
We used a rigorous analytical pipeline for essential gene identification as outlined in

additional file 1. As a result, we identified 336 essential genes that are required for aerobic

growth of *S*. Typhimurium 14028 in LB broths and on LB agar plates (Additional file 2). We

compared the essential genes in *S*. Typhimurium 14028 to those in *S*. Typhimurium SL3261

identified by TraDIS [13]. Interestingly, out of 336 genes in our essential gene list, 265

(80%) orthologous genes in *S*. Typhimurium 14028 were also essential in *S*. Typhimurium

SL3261 (Additional file 3). This is a significant overlap considering variations in the genetic

backgrounds of the two strains. Further, KEGG pathway analysis recognized 306 out of the

336 genes, which were categorized into 23 essential pathways (Additional file 4).

We also used the same method to identify essential genes from the Tn-seq data obtained from

iron-restricted conditions. Surprisingly, the number of essential genes under iron-restricted

conditions decreased to 215 genes, which indicated that 121 genes (36%) of the 336 essential

genes were considered non-essential under iron-restriction conditions (Additional file 5). The

number of reads corresponding to the insertions in these 121 genes significantly increased

under iron-restricted conditions: the average read counts in the 121 genes were 4.3 in LB-III

whereas this elevated to 68 in Dip400 (Additional file 6). This is a clear evidence that the

mutants of the 121 genes escaped immediate killing or growth arrest, and were able to

multiply slowly under iron-restricted conditions. In another word, chelation of iron in the

media allowed the mutants of these 121 genes to become no longer "essential" according to

the working definition of an essential gene (Fig. 3, Additional file 7).

Fitness of the mutants of iron-dependent essential genes
138
We next measured the changes of mutant fitness for the 121 genes in the presence of high

concentrations of Dip using Dip250-I, Dip250-II and Dip400 Tn-seq data as outputs in

comparison to LB-III as the input. Strikingly, the mutant fitness of 97 out of 121 genes (78%)

increased numerically (fitness expressed in Log_2 fold change in read counts) in Dip400, and 8

other genes showed increased mutant fitness in Dip250-I, Dip250-II or both. Further

comparative analysis revealed that 33 essential genes showed significant increase in fitness in

the presence of Dip, including *gyrA*, *gyrB*, and *ileS* (p value < 0.05; Fig. 4). For these 33

genes, the mutants showed at least Log_2 fold change in read counts $\geq$ 3.64 in Dip400 in

comparison to LB-III. Interestingly, *gyrA*, *gyrB* and *ileS* were not in the list of 121 iron-

dependent essential genes (e.g. considered as iron-independent essential genes), whereas

fitness of these mutants increased significantly by iron restriction. In the case of *murA*, the

read counts increased from 5 (LB-III) to 14,981 (Dip400), exhibiting highest increase in

abundance (Fig. 4). These are additional strong evidences supporting that the genes with

positive fitness in iron-restricted conditions are rendered nonessential and multiplied in iron-

restriction conditions.

Iron-dependent essential genes are not condition-specific essential genes
155

It is well known that the essential genes are operationally defined depending on the variations

in the specific optimal growth conditions used for the experiment. We then asked if our

discovery could be considered as a general extension of the concept to iron-restricted

conditions. To answer this question, we closely examined our Tn-seq data that were

generated for other stress conditions such as H_2O_2 [18] and H_2O_2 coupled with Dip

(unpublished). However, we could not find any similar patterns that a significant portion of

the essential genes in LB medium became nonessential under stress conditions. We also

examined other studies on identification of essential genes in other bacteria, including

Salmonella Typhimurium SL326 [13], *Mycobacterium tuberculosis* H37Rv [14] and

Mycoplasma mycoid JCVI-syn3.0 [16] as well as *E. coli* [10]. Nearly all orthologous

essential genes in these bacteria, specifically the 33 genes, barely tolerated insertions and thus

considered essential in their stress-free conditions (Additional file 8). Further, Lee et al [19]

used Tn-seq to identify essential genes in *Pseudomonas aeruginosa* under 6 different

conditions, and found that the 352 genes are general essential genes important in all 6

different media, while 199 essential genes were condition-specific. The condition-specific

essential genes constitute 11-23% essential genes depending on the growth medium. It is

important to note that all these 6 media support the growth of *P. aeruginosa* well, and thus

are not considered as stress conditions. In contrast, in our study 36% of the essential genes in

LB medium became nonessential under iron-restricted stress conditions, and there were no

condition-specific essential genes for iron-restriction condition (Fig. 3). Therefore, we reason

that these 121 genes should be still considered as general essential genes, instead of LB-

specific essential genes. In other words, these 121 genes are general essential genes, but

became non-essential under the iron-restricted condition due to the critical role iron plays in

broad range of death processes in *S.* Typhimurium following inactivation of the essential

functions.

Validation of the Tn-seq results

We next asked whether the increase of insertions and read counts for the 121 essential genes under iron-restricted conditions were due to a bias in Tn-seq data analysis. We conducted the analysis for identification of essential genes in this study without data normalization. Typically normalization of read counts is critical for reliable identification of conditionally essential genes for the mutant libraries before and after the selection. On the contrary, essential gene analysis is usually conducted without normalization. It is because the process is based on one Tn-seq profile from the condition of interest under which the essential genes are studied, and the information on insertion sites is critical for discovery of essential genes, while relative abundance of mutants is not. In this study, there were differences in read counts in Tn-seq conditions (Table 2). Furthermore, when focused on the reads in ORFs only, the total read counts of LB-III was ~30 millions (M) versus ~16 M in Dip400, excluding intergenic regions (Additional file 9). Interestingly, the insertions in the two genes, *STM14_2422* (*umuC*) and *STM14_2428*, consumed 8.7% of all reads in LB-III and 27% in Dip400. Consequently, excluding the reads from these two genes, an insertion in LB-III is expected had on average 227 reads, while 100 reads for an insertion in Dip400 (Additional file 9). This indicates that the bias in read counts was in favor to not see insertions and read counts in essential genes in iron-restricted conditions (Dip400). Despite this bias in read counts, which was further intensified by reads from these two genes, the read counts in the 121 iron-dependent essential genes were higher in Dip400 compared to LB-III. This is a strong evidence that the mutants in these 121 genes did indeed multiply slowly in iron-restricted conditions.

Iron-dependent and -independent essential genes

We came to the conclusion that iron-restriction alleviated the immediate killing or growth

arrest of the mutants in the 121 iron-dependent essential genes, allowing growth of the

mutants, although the underlying mechanism(s) is unknown. Here, we propose to divide the

essential genes in *S.* Typhimurium for convenience into two categories according to the

dependency of their gene essentiality on iron: iron-dependent and iron-independent essential

genes (Fig. 3). The 121 essential genes that allowed growth of the mutants under iron-

restricted conditions are iron-dependent essential genes (Fig. 3, Fig. 5, and Additional file 6).

On the contrary, the 215 essential genes that did not tolerate growth of the mutants regardless

of iron concentrations are iron-independent essential genes (Fig. 3, Fig. 5, and Additional file

5).

One note for caution for this classification is that for any iron-independent essential genes

reported in this study the classification can be temporary. It is because the decision on gene

essentiality can be dependent on the sequencing depth, and a substantial increase in the

sequencing depth may reclassify some of the iron-independent genes as iron-dependent

essential genes.

Then, we compared the average read counts between iron-independent and iron-dependent

essential genes under iron-replete or iron-restricted condition. For the 215 iron-independent

essential genes, their average read counts were 9.6 in Dip400 (Additional file 5). On the

contrary, for the 121 iron-dependent essential genes, their average read counts were 67.9 in

Dip400 (Additional file 6). Whereas in iron-replete condition, LB-III, the average read counts

were only 2.2 for the 215 iron-independent essential genes and 4.3 for 121 iron-dependent

essential genes. In addition, we asked if the 121 iron-dependent essential genes represent

unique pathways in comparison to iron-independent essential genes. The result of KEGG

pathway analysis showed that iron-independent and dependent genes are represented in 15

and 10 pathways, respectively, with 7 overlapping ones (Fig. 6).

It is also important to note that all known targets of antibiotics, with only one exception of

colistin, are encoded by iron-dependent essential genes (Fig. 3). It is uncertain if this strong

bias toward iron-dependent genes reflects the preference of the antibiotic-producing

microorganisms in nature to target these essential genes. If that's the case, it would be very

intriguing to explore the possible mechanistic reasons for the bias in relation to the roles of

antibiotic production in the natural environments.

Fitting iron-dependent essential genes in ROS-mediated antibiotic killing model
237
It has been well established that iron contributes to cell death due to various causes. 2,2`-

Dipyridyl iron chelator inhibits the Fenton reaction, which is required for ROS generation [5,

20]. We speculate that following disruption of the essential genes with transposon insertions,

ROS might contribute to the death or growth arrest depending on the target genes, in addition

to the disruption of the essential protein functions. We reason that our results have significant

implications in understanding and expanding the current model of ROS-mediated common

killing mechanisms of bactericidal antibiotics.

Since its first proposal by Kohanski et al. [5], this hypothesis has been substantiated by

numerous studies using different bacterial species and bactericidal antibiotics [21-24].

Traditionally, the mechanisms of antibiotic action have been studied largely in terms of

antibiotic-target interactions. However, numerous researches supporting the ROS-mediated

killing mechanism have shown that the interaction of antibiotic-target leads to production of

ROS, contributing to the killing activity mediated by direct blocking of the essential

pathways for living cells. In our study, we did not use bactericidal antibiotics to disrupt an

essential pathway. However, inactivation of the essential gene functions was achieved in a

permanent and irreversible manner through Tn5 insertions in the essential genes, instead of

reversible antibiotics treatment. In most studies focused on understanding the killing

mechanisms of bactericidal antibiotics, the experiments were conducted using optimized sub-lethal concentrations of relevant antibiotics to allow reliable measurements of quantitative changes in killing effects (e.g. by addition of iron chelator) [5]. However, inactivation of the essential protein functions by Tn5 insertions in our study is expected to exhibit the lethal effect equivalent to or even stronger than that caused by high lethal concentrations of bactericidal antibiotics. This situation may limit the effect of iron chelation to very small, making it difficult to detect and measure reduced killing by iron chelator. However, the Tn-seq procedure conducted with deep sequencing in combination with the high concentrations of Dip (up to 400 μM, which is much higher than typically used) employed in this study allowed to detect the small reductions in killing or growth arrest under the iron-restricted conditions.

Until now the ROS-mediated killing mechanism has been studied and discussed in the context of a few bactericidal antibiotics and their molecular targets. Our Tn-seq data show that this ROS-mediated killing mechanism might be linked at least to one-third of the essential genes in *S.* Typhimurium, thereby potentially expanding the ROS-mediated lethal pathway as a more general mechanism connected to a broad range of basic essential pathways in *S.* Typhimurium.

Iron-independent essential genes may be better targets for antibiotic development

We believe that our findings have profound implications for understanding current crisis in public health due to the rapid increase in antibiotic resistance to most antibiotics in clinical use. Our deeper understanding of the underlying mechanisms would help in development of novel antibiotics with inherently reduced chance of developing drug resistance. Here, we propose that iron-independent essential genes may serve as better targets for antibiotic development for the following reasons: it has been shown that there are two opposing aspects

of ROS-mediated killing mechanism. When ROS production is high, it would lead to

facilitated killing of bacterial cells. On the contrary, when ROS production is low, it would

lead to production of resistant mutants through mutagenic action of ROS on DNA [25]. When

Salmonella infects the host, iron-restricted host niches would suppress ROS production from

bactericidal antibiotics, which would in turn reduce ROS-mediated killing process and thus

overall killing effect by the antibiotics. Furthermore, depending on the iron restriction levels,

and reduced local concentrations of antibiotics, it might allow production of low levels of

ROS, which might facilitate production of antibiotic resistant mutants through its mutagenic

action on DNA. In contrast, if a certain antibiotic targets an iron-independent essential

pathway (such as those encoded by the 215 iron-independent essential genes discovered in

this study), we speculate that since ROS production is not a part of their lethal processes,

blocking the pathways will lead to iron-independent killing, without increasing the chance of

developing antibiotic resistant population via mutagenic action of ROS [25].

Our Tn-seq results show clearly that mutants of iron-dependent essential genes can grow

slowly in iron-restricted conditions, and the same phenomenon may happen in the host

tissues, because iron-restriction by host is a vital mechanism to combat the pathogen. As a

result, it may be hard to completely kill, and eliminate *Salmonella* by targeting iron-

dependent essential genes. Conversely, blocking iron-independent essential pathways would

allow immediate killing of *Salmonella* regardless of iron concentration. Thus, the possibility

will be higher to eradicate this pathogenic bacterium by targeting the iron-independent

essential pathways in comparison to iron-dependent essential pathways as is the case for all

antibiotics in clinical use except colistin.

A mechanism that bacteria exploit for antibiotic resistance is alteration of drug interaction

site. Our results emphasize that the majority of genes encoding currently known drug targets

are iron-dependent essential genes (Fig. 3). Prevalence of antibiotic resistance in clinical

isolates due to mutations in drug targets has been rising. Mutations in a peptidoglycan

synthesis gene *fts* which is target of β-lactams in *Haemophilus influenzae* cause resistance to

antibiotics [26, 27]. *E. coli* strains harboring mutations in *murA* are resistant to fosfomycin

[28]. UDP-N-acetylglucosamine enolpyruvyl transferase (MurA) catalyzes the reaction in the

first step of biosynthesis of peptidoglycan in bacterial cell wall, and the protein is the target

of fosfomycin [29]. Our Tn-seq results show that *murA* mutants did grow very well in iron-

restricted conditions and the mutant had 14,981 read counts in Dip400 but there were only 5

reads of this mutant in LB-III (Fig. 4). It has been reported that *Pseudomonas putida*

develops intrinsic fosfomycin resistance due to the presence of a salvage pathway that

bypasses *de novo* biosynthesis of MurA [30]. Since *murA* is an iron-dependent essential gene

in our study, we reason that almost all *murA* mutants died effectively in LB-III because of the

contribution of ROS in the death process. However, in Dip400, reduced ROS formation and

the salvage pathway biosynthesis of MurA might have caused *S*. Typhimurium to grow well.

Further, Fluoroquinolone-resistant bacteria are also present in clinical isolates due to

mutations in drug targets, *gyrA*, *gyrB*, *parC*, and *parE*, in pathogens such as *Shigella flexneri*

[31], *Salmonella* Typhi [32], and group B *Streptococcus* [33]. Rifampin-resistant

Mycobacterium tuberculosis isolates are associated with mutations in their targets, *rpoB* and

rpoC [34, 35]. Mutations in *rplC* contributed to *Staphylococcus aureus* resistance to linezolid

in a clinical isolate [36]. Finally, Telithromycin resistant-isolates of *S. aureus* due mutations

in a ribosomal gene, *rplB*, were detected *in vitro* [37]. All together, these antibiotic targets,

which are iron-dependent essential genes, can mutate and alter the structure of corresponding

proteins in order to evade lethal interactions with the antibiotics.

One example of an antibiotic targeting a protein encoded by iron-independent gene is

colistin. Colistin (polymyxin E) is a last resort antibiotic for treatment of infections caused by

multidrug resistant Gram-negative bacteria [38]. This bactericidal drug interacts with the

lipid A moiety of lipopolysaccharide (LPS) and ultimately causes membrane lysis [39]. We

showed that the genes encoding the molecular targets of Colistin, *lpxABCDHK* are iron-

independent essential genes. Over the last 60 years, Colistin has been used for fighting

infectious diseases, but it has been used with caution and limitation due to its known toxicity.

This resulted in less frequent use of Colistin, which is considered as the main reason why

bacterial resistance is low for Colistin. Our Tn-seq results indicate that disruption of LPS is

lethal in *S.* Typhimurium and there is no contribution of ROS to death process caused by

Colistin. However, one study demonstrated that Colistin induced *Acinetobacter baumannii*

killing through ROS production [40]. The contradicted findings are not surprising,

considering that there has been continued debates on the common antibiotic killing

mechanism via ROS. Although this model is widely accepted, a few studies challenged it [41,

42].

Conclusion

In this work we employed Tn-seq to elucidate the essential genes exhibiting iron-dependent

and iron-independent mutant phenotypes in *S.* Typhimurium. Our Tn-seq data indicated that

when transposon mutant library was treated with an iron chelator, the mutants of

approximately one-third of essential genes escaped immediate growth inhibition or killing,

and multiplied slowly, increasing their relative abundance. Based on this observation, we

speculate that the iron chelator reduces ROS formation via inhibition of the Fenton reaction,

thereby lessening the lethal outcomes following inactivation of the essential functions.

Accordingly, we divided 336 essential genes in *S.* Typhimurium 14028 into iron-dependent

vs. iron-independent essential genes depending on their dependency of the essentiality on

iron concentration. We propose that iron-independent essential genes may serve as better

targets to develop new antibiotics, because targeting these pathways would lead to immediate

killing or growth inhibition regardless of local iron concentrations in the host tissues. The

proposed model illustrating our concept of iron-dependent and iron-independent essential

genes is summarized in Fig 5. Obviously, further studies providing experimental evidences

on the direct involvement of ROS production in the process would be necessary to support

our model. Due to the difficulty associated with the phenotype of essentiality, the use of

inducible knockouts such as CRISPRi [17] would be helpful in elucidating the role of ROS in

the process following inactivation of the essential pathways. It would be also interesting to

see if we could come to a similar conclusions using the same Tn-seq approach, when

thiourea, hydroxyl radical scavenger, is used in place of Dip to suppress ROS production or

the mutants are cultured anaerobically.

This study provides new insights on the previously unknown aspect of essential pathways in

Salmonella. Although further studies are needed to gain better understanding on the scope of

this observation (e.g. other bacteria), and to establish mechanistic basis of the iron-dependent

essential genes, this study points us to a new direction of research that would be important to

understand and overcome current crisis of antibiotic drug resistance.

Methods

Bacterial strains and Tn5 mutant library construction

Salmonella enterica subsp. enterica serovar Typhimurium ATCC 14028 with spontaneous

mutation conferring resistance to nalidixic acid (NA) was used in this study. All procedures

involving this pathogen (Biosafety level 2) were conducted according to the protocol

approved by Institutional Biosafety Committee (IBC) at the University of Arkansas.

Transposon mutant libraries were prepared as previously described by Karash et al., [18].

Briefly, *S.* Typhimurium ATCC 14028 was subjected to transposon mutagenesis by

biparental mating using *Escherichia coli* SM10 λ*pir* carrying a pBAM1 transposon-delivery

plasmid vector [43] as the donor. An equal volume of overnight growth cultures of the donor

and recipient bacteria (*S.* Typhimurium) were washed with 10 mM $MgSO_4$ and concentrated

on the nitrocellulose filter, which was then incubated for 5 h at 37°C on a surface of LB agar

plate. After the incubation, the cells were washed with 10 mM $MgSO_4$ and plated on LB agar

plates containing 50 µg/ml NA and 50 µg/ml Kanamycin (Km). The plates were incubated at

37°C for 24 h. Then, the colonies were scrapped off, added into LB broth supplemented with

7% DMSO, and stored at −80°C in aliquots. We constructed two mutant libraries, A and B.

Each library contains approximately 325,000 mutants.

Growth rate measurement for *S.* Typhimurium
391
A single colony of *S.* Typhimurium ATCC 14028 was inoculated into 2 ml LB broth medium

in a 5 ml tube and incubated overnight (~16 h). Freshly prepared LB broth media

supplemented with different concentrations of 2,2`-Dipyridyl (Dip) were inoculated with the

overnight culture at a 1:200 dilution. The cultures were immediately added into a 96-well

microplate (200 µl/well) and incubated in a TECAN Infinite 200 microplate reader at 37°C

(shaking amplitude of 1.5 mm, and shaking duration of 5 s), and OD_{600} was measured every

10 min for 24 h. The data were used to determine lag time phase, growth rate, and maximum

OD_{600} for each concentration using GrowthRates script [44].

Selection of mutant libraries for Tn-seq analysis
401
An aliquot of transposon libraries in stock was thawed at room temperature and diluted 1:10

in LB broth. The library was incubated at 37°C with shaking at 225 rpm for an hour and then

washed twice in PBS. The library-A was inoculated to 20 ml LB broth (LB-II) and LB

supplemented with either 100 (Dip100) or 150 µM Dip (Dip150) in a 300 ml flask, with

seeding CFUs of 3.5 x 10^6 per ml. We also included LB-I for which Library-A was directly

subjected to Tn-seq after activation and washing. In addition, to accomplish a higher

saturation level, Library-A was combined with Library-B (termed Library-AB; Fig. 1).

Library-AB was inoculated to 20 ml LB broth in a 300 ml flask (LB-III) and LB

supplemented with either 250 (Dip250-I and Dip250-II) or 400 µM Dip (Dip400), at seeding

CFUs of 8 x 10^6 per ml as described above. Dip100, Dip150, Dip250-I, Dip250-II, and

Dip400 along with LB-II and LB-III were incubated at 37°C with shaking at 225 rpm until

the cultures reached mid-log phase (OD_{600} of ~2.7). Then, the cultures were immediately

centrifuged to get the cell pellets, which were stored at -20°C for downstream analysis.

Preparation of Tn-seq libraries for HiSeq sequencing
416
The preparation of Tn-seq libraries was performed as previously described by Karash et al.,

[18]. Briefly, genomic DNA was extracted for each sample using DNeasy Blood & Tissue kit

(Qiagen), and quantified using Qubit dsDNA RB Assay kit (Invitrogen). To remove the

pseudo Tn5 mutants generated by chromosomal integration of pBAM1, genomic DNA was

digested with PvuII-HF (New England Biolabs), and purified with DNA Clean &

Concentrator-5 kit (Zymo Research). Then, a linear PCR extension was performed using

Tn5-DPO (5'-AAGCTTGCATGCCTGCAGGTIIIIICTAGAGGATC-3'). The PCR reaction

was performed in a 50 µl reaction containing GoTaq Colorless Master Mix (Promega), 20

µM Tn5-DPO primer, 100 ng gDNA, and H_2O. The PCR cycles consisted of 95°C for 2 min,

followed by 50 cycles at 95°C for 30 sec, 62°C for 45 sec, and 72°C for 10 sec. The PCR

product was purified with the DNA Clean & Concentrator-5 kit. The C-tailing reaction was

conducted with terminal transferase (TdT; New England Biolabs), $CoCl_2$, dCTP, ddCTP, TdT

buffer, and the purified linear PCR product. dCTP and ddCTP were included at the molar

ratio of 20:1. The mixture was incubated at 37°C for 1 h, which was followed by 10 min

incubation at 70°C. The C-tailed product was purified. Next, the exponential PCR was

performed   with   P5-BRX-TN5-MEO   primer,

AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGCTCTTCCGA

TCTNNNNAG-BC-CCTAGGCGGCCTTAATTAAAGATGTGTATAAGAG (where "BC'

denotes different sample index barcodes of 8nt long) and P7-16G primer,

CAAGCAGAAGACGGCATACGAGCTCTTCCGATCTGGGGGGGGGGGGGGGGG. The

PCR reaction was performed in a 50 µl reaction containing GoTaq Green Master Mix, P5-

BRX-TN5-MEO primer, P7-16G primer, purified C-tailed Tn5-junction fragments, and H_2O;

the PCR cycles consisted of 95°C for 2 min, followed by 30 cycles of 95°C for 30 sec, 60°C

for 30 sec, and 72°C for 20 sec, with the final extension at 72°C for 5 min. Then, the 50 µl

PCR products were run on an agarose gel and the DNA fragments of size 325–625 bp were

extracted using Zymoclean Gel DNA Recovery kit (Zymo Research). The DNA libraries

were quantified using Qubit dsDNA RB Assay kit. The DNA libraries were combined and

sequenced on a flow cell of HiSeq 3000 using single-end read and 151 cycles (Illumina) at

the Center for Genome Research & Biocomputing in Oregon State University.

Analysis of Tn-seq data
448
The HiSeq sequence results were downloaded onto High Performance Computing Center

(AHPCC) at the University of Arkansas. The libraries were de-multiplexed using a custom

Python script. The script searched for the unique 8-nucleotide barcode for each library for

perfect matches. The transposon genomic junctions were extracted using Tn-Seq Pre-

Processor (TPP) tool [45]. The TPP searched for the 19 nucleotide inverted repeat (IR) in a

fixed sequence window and identified five nucleotides (GACAG) at the end of the IR

sequence, allowing one nucleotide mismatch. The genomic junctions that start immediately

after the GACAG were extracted and the C-tails were removed. The junction sequences of

less than 20 nucleotides were removed and remaining junction sequences were mapped to the

Salmonella enterica serovar Typhimurium 14028s genome and plasmid using BWA-0.7.12

[46]. The TPP counted and reported the number of total sequence reads after filtering, total

mapped read, and total unique insertions in each library.

Identification of essential genes

LB-I, LB-II, and LB-III were analyzed to identify the essential genes in *S*. Typhimurium

14028. We used two different tools for Tn-seq essential gene analysis. First, TRANSIT [45]

analysis of essentiality on gaps in entire genome was conducted using tn5gaps algorithm. The

5% of N-terminal and 10% of C-terminal of open reading frames (ORF) were removed and

even insertions with only one reads were considered for the analysis. The gene was

considered essential if its p value $\leq$ 0.05. Secondly, Tn-Seq Explorer [47], was used for

essential gene analysis by applying a 550 window size. The 5% of N-terminal and 10% of C-

terminal ORFs were removed and insertions with only one reads were included in the

analysis. The gene was considered essential if its Essentiality Index was $\leq$ 2. Then, the

essentiality analysis results by both methods were combined. Finally, we demanded that for a

gene to considered essential for growth on LB agar or LB broth the following three criteria

should be met: (i) the gene is essential in LB-III by Tn-Seq Explorer analysis (ii) the gene is

essential in LB-III by TRANSIT analysis (iii) the gene is essential in at least 5 out of the 6

analysis that was performed for the LB-I, LB-II, and LB-III by the two analysis tools

(Additional file 1). We made an exception for 17 genes to be considered essential:

requirements for essential genes in at least 5 out of the 6 analysis were reduced to 4. This

exception was based on the other analysis for the same libraries selected and analyzed under

different growth conditions. The same strategy was used to identify essential genes under

Dip250-I, Dip250-II, and Dip400.

Calculation of fitness for selected essential genes

Fitness of essential genes for all iron-restricted conditions were analyzed by using TRANSIT,

with resampling option. LB-I was used as the input for Dip100 and Dip150, while LB-II was

used as the input for Dip250-I, Dip250-II, and Dip400. For data normalization for fitness

calculation, Trimmed Total Reads (TTR) option in TRANSIT was used and 10,000 samples

were used for the analysis. The 5% of N-terminal and 10% of C-terminal of ORFs were

removed and the gene fitness was considered if p value was ≤ 0.05.

Acknowledgments
492
493 This research was supported by the Arkansas High Performance Computing Center (AHPCC)

494 which is funded through multiple National Science Foundation grants and the Arkansas

495 Economic Development Commission.

496 **Funding**
497
498 This study was conducted with the financial support from Arkansas Biosciences Institute

499 (ABI). The first author was supported by his parents, Cell and Molecular Biology (CEMB)

500 program at the University of Arkansas, and Human Capacity Development Program-

501 Kurdistan Regional Government (HCDP-KRG).

502
503 **Availability of data and materials**
504
505 All Tn-seq sequencing data are available on NCBI Sequence Read Archive under BioProject

506 number PRJNA397775.

507 **Authors' contributions**
508
509 Conceived and designed the experiments: YK SK. Performed the experiments, analyzed the

510 data, wrote the manuscript: SK. Revised the manuscript: YK SK.

511 **Competing interests**
512
513 The authors declare that they have no competing interests

515 **References**
516
1. Lewis K. Platforms for antibiotic discovery. Nat Rev Drug Discov. 2013;12: 371-387.

2. Perrone GG, Tan S, Dawes IW. Reactive oxygen species and yeast apoptosis. Biochim

Biophys Acta. 2008;1783: 1354-1368.

3. Zhang D, Shao J, Lin J, Zhang N, Lu B, Lin S, et al. RIP3, an Energy Metabolism Regulator
That Switches TNF-Induced Cell Death from Apoptosis to Necrosis. Science. 2009;325: 332-
336.

4. Dixon SJ, Lemberg KM, Lamprecht MR, Skouta R, Zaitsev EM, Gleason CE, et
al. Ferroptosis: an iron-dependent form of nonapoptotic cell death. Cell. 2012;149: 1060-1072.

5. Kohanski MA, Dwyer DJ, Hayete B, Lawrence CA, Collins JJ. A Common Mechanism of
Cellular Death Induced by Bactericidal Antibiotics. Cell. 2007;130: 797-810.

6. Dwyer DJ, Belenky PA, Yang JH, MacDonald IC, Martell JD, Takahashi N, et al. Antibiotics
induce redox-related physiological alterations as part of their lethality. Proc Natl Acad Sci U
S A. 2014;111: E2100-E2109.

7. Dong TG, Dong S, Catalano C, Moore R, Liang X, Mekalanos JJ. Generation of reactive
oxygen species by lethal attacks from competing microbes. Proc Natl Acad Sci U S
A. 2015;112: 2181-2186.

8. Kashyap DR, Wang M, Liu L, Boons G, Gupta D, Dziarski R. Peptidoglycan recognition
proteins kill bacteria by activating protein-sensing two-component systems. Nat Med.
2011;17: 676-683.

9. Kovacs I, Horvath M, Lanyi A, Petheo G, Geiszt M. Reactive oxygen species-mediated
bacterial killing by B lymphocytes. J Leukoc Biol. 2015;97: 1133-1137.

10. Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, et al. Construction of
Escherichia coli K-12 in-frame, single-gene knockout mutants: the Keio collection. Mol Syst
Biol. 2006;2: 2006.0008.

11. Yamamoto N, Nakahigashi K, Nakamichi T, Yoshino M, Takai Y, Touda Y, et al. Update on
the Keio collection of *Escherichia coli* single-gene deletion mutants. Mol Syst Biol. 2009;5:
335.

12. Opijnen T, Bodi KL, Camilli A. Tn-seq: high-throughput parallel sequencing for fitness and
genetic interaction studies in microorganisms. Nat Methods. 2009;6: 767-772.

13. Barquist L, Langridge G, Turner D, Phan M, Turner A, Bateman A, et al. A comparison of
dense transposon insertion libraries in the *Salmonella* serovars Typhi and Typhimurium.
Nucleic Acids Res. 2013;41: 4549-4564.

14. DeJesus MA, Gerrick ER, Xu W, Park SW, Long JE, Boutte CC, et al. Comprehensive
Essentiality Analysis of the *Mycobacterium tuberculosis* Genome via Saturating Transposon
Mutagenesis. Mbio. 2017;8: e02133-16.

15. Breton Y, Belew AT, Valdes KM, Islam E, Curry P, Tettelin H, et al. Essential Genes in the
Core Genome of the Human Pathogen *Streptococcus pyogenes*. Sci Rep. 2015;5: 9838.

16. Hutchison C, Chuang R, Noskov V, Assad-Garcia N, Deerinck T, Ellisman M, et al. Design
and synthesis of a minimal bacterial genome. Science. 2016;351: 1414-U73.

17. Peters J, Colavin A, Shi H, Czarny T, Larson M, Wong S, et al. A Comprehensive, CRISPR-
based Functional Analysis of Essential Genes in Bacteria. Cell. 2016;165: 1493-1506.

18. Karash S, Liyanage R, Qassab A, Lay J, Kwon Y. A Comprehensive Assessment of the
Genetic Determinants in Salmonella Typhimurium for Resistance to Hydrogen Peroxide

Using Proteogenomics. Sci Rep. 2017;7: 17073.

19. Lee SA, Gallagher LA, Thongdee M, Staudinger BJ, Lippman S, Singh PK, et al. General and
condition-specific essential functions of *Pseudomonas aeruginosa*. Proc Natl Acad Sci U S A.
2015;112: 5189.

20. Ezraty B, Barras F. The 'liaisons dangereuses' between iron and antibiotics. FEMS Microbiol.
Rev. 2016;40: 418–435.

21. Davies BW, Kohanski MA, Simmons LA, Winkler JA, Collins JJ, Walker GC. Hydroxyurea
induces hydroxyl radical-mediated cell death in Escherichia coli. Molecular cell. 2009;36(5):
845-860.

22. Ling J, Cho C, Guo L-T, Aerni H, Rinehart J, Söll D. Protein aggregation caused by
aminoglycoside action is prevented by a hydrogen peroxide scavenger. Molecular cell.
2012;48(5): 713-722.

23. Van Acker H, Coenye T. The role of reactive oxygen species in antibiotic-mediated killing of
bacteria. Trends Microbiol. 2017;25: 456–466

24. Liou J, Hung Y, Yang C, Chen Y. The Antimicrobial Activity of Gramicidin A Is Associated
with Hydroxyl Radical Formation. Plos One. 2015;10: e0117065.

25. Kohanski MA, DePristo MA, Collins JJ. Sublethal antibiotic treatment leads to multidrug
resistance via radical-induced mutagenesis. Mol Cell. 2010;37: 311-20.

26. Kim I, Ki C, Kim S, Oh WS, Peck KR, Song J, et al. Diversity of Ampicillin Resistance
Genes and Antimicrobial Susceptibility Patterns in *Haemophilus influenzae* Strains Isolated
in Korea. Antimicrob Agents Chemother. 2007;51: 453-460.

27. Kishii K, Chiba N, Morozumi M, Hamano-Hasegawa K, Kurokawa I, Masaki J, et al. Diverse
mutations in the *ftsI* gene in ampicillin-resistant *Haemophilus influenzae* isolates from
pediatric patients with acute otitis media. J Infect Chemother. 2010;16: 87-93.

28. Takahata S, Ida T, Hiraishi T, Sakakibara S, Maebashi K, Terada S, et al. Molecular
mechanisms of fosfomycin resistance in clinical isolates of *Escherichia coli*. Int J Antimicrob
Agents. 2010;35: 333-337.

29. Eschenburg S, Priestman M, Schonbrunn E. Evidence That the Fosfomycin Target Cys115 in
UDP-N-acetylglucosamine Enolpyruvyl Transferase (MurA) Is Essential for Product Release.
J Biol Chem. 2005;280: 3757-3763.

30. Gisin J, Schneider A, Nagele B, Borisova M, Mayer C. A cell wall recycling shortcut that
bypasses peptidoglycan de novo biosynthesis. Nat Chem Biol. 2013;9: 491-U86.

31. Qin T, Bi R, Fan W, Kang H, Ma P, Gu B. Novel mutations in quinolone resistance-
determining regions of *gyrA*, *gyrB*, *parC* and *parE* in *Shigella flexneri* clinical isolates from
eastern Chinese populations between 2001 and 2011. Eur J Clin Microbiol Infect Dis. 2016;
35: 2037-2045.

32. Misra R, Thakare R, Amrin N, Prasad K, Chopra S, Dhole T. Antimicrobial susceptibility
pattern and sequence analysis of DNA gyrase and DNA topoisomerase IV in *Salmonella*
enterica serovars Typhi and Paratyphi A isolates with decreased susceptibility to
ciprofloxacin. Trans R Soc Trop Med Hyg. 2016;110: 472-479.

33. Neemuchwala A, Teatero S, Patel S, Fittipaldi N. Fluoroquinolone Resistance among Clonal

Complex 1 Group B *Streptococcus* Strains. Can J Infect Dis Med Microbiol. 2016;2016:
6403928.

34. Ma X, Wang H, Deng Y, Liu Z, Xu Y, Pan X, et al. rpoB Gene Mutations and Molecular
Characterization of Rifampin-Resistant *Mycobacterium tuberculosis* Isolates from Shandong
Province, China. J Clin Microbiol. 2006;44: 3409-3412.

35. Vos Md, Müller B, Borrell S, Black PA, P. D. van Helden, Warren RM, et al. Putative
Compensatory Mutations in the *rpoC* Gene of Rifampin-Resistant *Mycobacterium*
tuberculosis Are Associated with Ongoing Transmission. Antimicrob Agents Chemother.
2013;57: 827-832.

36. Locke JB, Morales G, Hilgers M, C KG, Rahawi S, Picazo JJ, et al. Elevated Linezolid
Resistance in Clinical cfr-Positive *Staphylococcus aureus* Isolates Is Associated with Co-
Occurring Mutations in Ribosomal Protein L3. Antimicrob Agents Chemother. 2010;54:
5352-5355.

37. Gentry DR, Holmes DJ. Selection for High-Level Telithromycin Resistance in
Staphylococcus aureus Yields Mutants Resulting from an *rplB*-to-*rplV* Gene Conversion-
Like Event. Antimicrob Agents Chemother. 2008;52: 1156-1158.

38. Nation R, Li J. Colistin in the 21st century. Curr Opin Infect Dis. 2009;22: 535-543.

39. Hancock RE. Peptide antibiotics. The Lancet. 1997;349: 418-422.

40. Sampson TR, Liu X, Schroeder MR, Kraft CS, Burd EM, Weiss DS. Rapid Killing of
Acinetobacter baumannii by Polymyxins Is Mediated by a Hydroxyl Radical Death Pathway.
Antimicrob Agents Chemother. 2012;56: 5642-5649.

41. Keren I, Wu Y, Inocencio J, Mulcahy L, Lewis K. Killing by Bactericidal Antibiotics Does
Not Depend on Reactive Oxygen Species. Science. 2013;339: 1213-1216.

42. Liu Y, Imlay J. Cell Death from Antibiotics Without the Involvement of Reactive Oxygen
Species. Science. 2013;339: 1210-1213.

43. Martínez-García E, Calles B, Arévalo-Rodríguez M, de Lorenzo V. pBAM1: an all-synthetic
genetic tool for analysis and construction of complex bacterial phenotypes. BMC Microbiol.
2011;11: 1-38.

44. Hall BG, Acar H, Nandipati A, Barlow M. Growth rates made easy. Mol Biol Evol. 2014;31:
232-238.

45. DeJesus M, Ambadipudi C, Baker R, Sassetti C, Ioerger T. TRANSIT - A Software Tool for
Himar1 TnSeq Analysis. PLoS Comput Biol. 2015;11: e1004401.

46. Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform.
Bioinformatics. 2009;25: 1754.

47. Solaimanpour S, Sarmiento F, Mrazek J. Tn-Seq Explorer: A Tool for Analysis of High-
Throughput Sequencing Data of Transposon Mutant Libraries. Plos One. 2015;10: e0126070.

637
638

Figure Legends

Fig.1 Schematic representation of study design. Transposon library-A inoculated to LB broth (LB-II) or the LB contained either 100 μM iron chelator Dip (Dip100) or 150 μM Dip (Dip150). LB-I was library-A that subjected to Tn-seq without growth. Transposon library-B inoculated to LB broth (LB-III) or the LB contained either 250 μM Dip (Dip250) or 400 μM Dip (Dip400). The cultures were grown till mid-log phase and then subjected to Tn-seq.

Fig. 2 Effect of 2,2`-Dipyridyl (Dip) on *S*. Typhimurium growth. An overnight culture diluted 1:200 in LB broth supplemented with 100 μM Dip (Dip150), 150 μM Dip (Dip150), 250 μM Dip (Dip250), 400 μM Dip (Dip400), or Dip free (LB). The cultures were added to a 96-well plate and directly incubated at 37∘C in a plate reader, reading OD_{600} every 10 minuets. The maximum OD_{600} reduction is shown as a percentage.

Fig. 3 Iron-independent essential genes and iron-dependent essential genes of *Salmonella* Typhimurium. Essential genes of *S*. Typhimurium were identified using Tn-seq method in LB media (iron-replete) and LB media supplemented with 2,2`-Dipyridyl (Dip;

iron-restricted). Among all 336 essential genes identified, 215 genes were essential under

both conditions (iron-independent essential genes), while 121 genes were essential only in

iron-replete condition (iron-dependent essential genes). The iron-dependent essential genes

included 11 genes encoding well known antibiotic targets as shown in the table below.

Fig. 4 Iron-dependent essential genes have increased insertions and read counts in iron-

restricted conditions. *Salmonella* Typhimurium transposon mutants were grown in LB

media (iron-replete) and LB media supplemented with Dip (iron-restricted). Tn-seq analysis

identified 121 genes including these genes as essential in iron-replete, but not in iron-

restricted conditions. The fitness of these 33 genes increased in Dip400. Gene fitness is Log_2

fold change of sequence reads in Dip400 in comparison to LB-III.

Fig. 5 Identification of iron-dependent essential genes by Tn-seq. *Salmonella*

Typhimurium Tn5 mutant library was inoculated to iron-replete media (LB media) and iron-

restricted media (LB supplemented with an iron chelator 2,2`-Dipyridyl). The cultures were

grown till mid-log phase and then Tn-seq was used to identify essential under each condition.

The essential genes that were considered essential only in iron-replete condition are iron-

dependent essential genes.

Fig 6. Fe-dependent essential pathways in *Salmonella* Typhimurium identified by Tn-

seq. Transposon libraries were grown in iron-replete and iron restricted conditions. Essential

genes identified for both conditions. The genes that were essential in iron-replete condition

but not in iron-restricted conditions were considered Fe-dependent. KEGG pathway analysis

was used for pathway description.

Fig. 7 Iron-independent essential genes may serve as better targets for development of

new antibiotics. The proposed model indicates that targeting iron-independent essential

pathways would lead to effective killing or growth inhibition of *Salmonella* Typhimurium

even in iron-restricted host niches.

Additional files

Additional files 1: Algorithms utilized for essential gene identification. Two tools were

used for essential gene analysis, TRANSIT (Gumbel) and Tn-Seq Explorer. LB-I, LB-II, and

Lb-III were analyzed separately by both tools for identification of essential genes. The gene

was considered essential if 5 out of 6 analyses was essential (E) or essentiality index (EI) < 3.

LB-I was transposon library inoculum subjected to Tn-seq without growth. LB-II and LB-III

were grown in LB broth till mid-log phase. Dip is abbreviation of iron chelator 2,2`-

Dipyridyl in μM. The number with Dip is concentration of Dip. (PDF)

Additional files 2: Full list of *Salmonella* Typhimurium essential genes in iron-replete

(LB) conditions identified by Tn-seq. (XLSL)

Additional files 3: The overlapped essential genes that identified by our Tn-seq method

of *Salmonella* Typhimurium 14028S in iron-replete (LB) conditions and essential genes

that identified by TraDIS method in *S*. Typhimurium SL3261. (XLSL)

Additional files 4: Required essential pathways of *Salmonella* Typhimurium in rich

media. Tn-seq libraries were grown in LB broth media till mid-log phase and on LB agar.

The total number of essential genes were 336 and KEGG pathway analysis categorized them

to 23 essential pathways. (PDF)

Additional files 5: Full list of *Salmonella* Typhimurium essential genes in iron-restricted

conditions identified by Tn-seq (Fe-independent essential genes). Iron chelator 250 and

400 μM 2,2`-Dipyridyl were used. (XLSL)

Additional files 6: Full list of *Salmonella* Typhimurium essential genes in iron-restricted

conditions identified by Tn-seq (Fe-dependent essential genes). Iron chelator 250 and 400

μM 2,2`-Dipyridyl were used. (XLSL)

Additional files 7: Average sequencing read counts in essential genes of *Salmonella* Typhimurium in iron-replete (LB) and iron-restricted (Dip400) conditions identified by Tn-seq. (XLSL)

Additional files 8: Iron-dependent essential genes are essential genes in other bacteria.
Almost all orthologous iron-dependent essential genes are genuine essential genes in (Keio) *E. coli*, *Salmonella Typhimurium* SL326, *Bacillus subtilis*, *Mycoplasma mycoid* JCVI-syn3.0, *Mycobacterium tuberculosis* H37Rv, and (LB-III) *Salmonella Typhimurium* 14028S in stress free conditions. (XLSL)

Additional files 9: Bias in read sequencing is in favor of LB (iron-replete) conditions.
Total sequencing reads and insertions in ORFs are shown. Sequencing reads consumed by two mutants, *umuC* and *STM14_2428* are also shown. An insertion in iron-replete (LB) has a chance to get 227 sequence reads but this reduced to 100 reads in iron-restricted condition (Dip400). (XLSL)

	Seeding CFUs/ml	CFUs/ml at mid-log phase	Time to reach mid-log phase h:min	OD600
colspan header	colspan	colspan	colspan	colspan

	Seeding CFUs/ml	CFUs/ml at mid-log phase	Time to reach mid-log phase h:min	OD600
Library-A	325,000 mutant colonies recovered on 50 LB agar plates			
LB-II	3,500,000	117,000,000	5	2.63
Dip100	3,500,000	190,000,000	5:35	2.61
Dip150	3,500,000	164,000,000	6:05	2.565
Library-AB	325,000 mutant colonies recovered on 50 LB agar plates + Library-A (total: 650,000 mutants)			
LB-III	8,000,000	700,000,000	5:30	2.567
Dip250-I	8,000,000	2,000,000,000	10	2.446
Dip250-II	8,000,000	2,010,000,000	10	2.462
Dip400	8,000,000	550,000,000	24	1.843

Table 1. Transposon inoculum densities and CFUs at mid-log phase.

731 **Table 1. Transposon inoculum densities and CFUs at mid-log phase.** The seeding CFUs

732 of all cultures counted following inoculation at time zero and at the mid-log phase when the

733 growth stopped. OD_{600} measurements were used to monitor the growth. LB is broth free of

734 Dip. Dip is abbreviation of iron chelator 2,2`-Dipyridyl in μM. The number with Dip is

735 concentration of Dip.

Table 2. Sequence read counts used in this study.				
Total reads with Tn5	**Extracted reads >20bp**	**Mapped Reads**	**Unique Insertions**	**Mean genomic length bp**
Library-A	**325,000 mutant colonies recovered on 50 LB agar plates**			
LB-I	18,225,644	14,437,819	12,289,451	115,784	93.8
LB-II	38,808,640	31,728,005	25,223,444	125,449	92.9
Dip100	25,788,698	21,034,947	16,991,894	117,474	93
Dip150	36,677,408	29,905,496	24,364,738	121,132	93.1
Library-AB	**325,000 mutant colonies recovered on 50 LB agar plates + Library-A (total: 650,000)**			
LB-III	57,779,778	47,575,248	39,248,662	193,728	90.6
Dip250-I	29,832,849	25,082,465	21,096,630	179,562	90.1
Dip250-II	35,439,669	28,119,351	23,104,233	181,534	89.1
Dip400	30,028,187	26,382,625	23,135,546	169,666	91
Total	272,580,873	224,265,956	185,454,598	1,204,329	91.7

Table 2. Sequence read counts used in this study. Total reads with Tn5 represent the

sequence reads that passed the quality control and had sequence of Tn5. Extracted reads > 20

bp represent sequence reads that had trimmed C-tail (if present) and their length were above

20 nucleotides. Number of mapped reads, unique insertions in chromosome with mean length

of mapped reads are shown. LB is broth free of Dip. Dip is abbreviation of iron chelator 2,2`-

Dipyridyl in μM. The number with Dip is concentration of Dip.

Salmonella Typhimurium Tn5 libraries grown till mid-log phase

Library-A
325,000 mutants

Library-AB
650,000 mutants

2,2`-Dipyridyl µM	0	100	150	0	250	250	400

Tn-seq	LB-I	LB-II	Dip100	Dip150	LB-III	Dip250-I	Dip250-II	Dip400

Essential genome of *S.* Typhimurium in iron-replete (LB) and iron-restricted (Dip)

121 Iron-dependant essential genes

215 Iron-independant essential genes

Of the 121 Iron-dependant essential genes, fitness of 33 mutants increased significantly in iron-restricted conditions (Dip400), $p < 0.05$, and proteins of 11 genes are well known antibiotic targets

Gene	Read counts		Gene Fitness (Log2FC)	Antibiotic	Target
	LB-III	**Dip400**			
ftsI	1	210	7.71	β-lactams	Cell wall synthesis
gyrA	0	106	4.78	Fluoroquinolones	DNA synthesis
gyrB	0	152	5.29	Fluoroquinolones	DNA synthesis
murA	5	14981	11.55	Fosfomycin	Cell wall synthesis
parC	1	98	6.61	Fluoroquinolones	DNA synthesis
parE	0	125	5.01	Fluoroquinolones	DNA synthesis
rplB	0	74	3.97	Aminoglycoside	Protein synthesis
rplC	1	38	5.25	Aminoglycoside	Protein synthesis
rpoB	0	392	4.96	Rifampin	RNA synthesis
rpoC	1	150	7.23	Rifampin	RNA synthesis
rpsD	0	242	5.94	Aminoglycoside	Protein synthesis

Essential genes have increased insertions and read counts in iron-restricted conditions
(LB: iron-replete, Dip: iron-restricted, I: insertion, R: read count, fitness: Log2 fold change)

Biological Process	Gene	LB-III		Dip250		Dip400			
		I	R	I	R	I	R	Fitness	p value
Cell division	*mukB*	3	9	6	184	5	120	3.73	0.025
Cell membrane	*yfiO*	1	3	4	129	6	180	5.9	0.0015
Cell wall biosynthesis	*ftsI*	1	1	1	0	5	210	7.71	0.0137
Cell wall biosynthesis	*mrdB*	0	0	5	24	5	98	4.36	0.0087
Cell wall biosynthesis	*murA*	1	5	3	45	4	14981	11.55	0.0008
Coenzyme A biosynthetic process	*dfp*	0	0	1	0	6	152	4.33	0.0001
Coenzyme A biosynthetic process	*coaA*	0	0	5	185	5	125	4.45	0.0021
DNA replication	*dnaE*	0	0	3	9	8	155	4.35	0.0004
DNA replication	*dnaX*	0	0	2	2	5	65	3.81	0.0085
DNA replication	*nrdA*	0	0	4	141	6	286	5.39	0.0006
DNA replication	*parC*	1	1	1	15	4	98	6.61	0.0458
DNA replication	*parE*	0	0	4	22	4	125	5.01	0.0287
Fatty acid metabolic process	*fabH*	0	0	3	42	5	74	3.97	0.006
Glutamine metabolic process	*glmS*	0	0	4	126	4	98	4.36	0.0058
Glutamine metabolic process	*pyrG*	1	7	4	5744	4	185	4.72	0.0462
Phospholipid biosynthetic process	*pssA*	1	7	4	88	5	139	4.31	0.016
Protein transport	*secY*	1	1	5	49	5	19	4.25	0.0139
Protein transport	*yidC*	1	1	6	165	4	144	7.17	0.0241
Transcription	*gyrA*	0	0	2	2	3	106	4.78	0.0253
Transcription	*gyrB*	0	0	2	0	3	152	5.29	0.0308
Transcription	*rpoB*	0	0	12	164	13	392	4.96	0
Transcription	*rpoC*	1	1	9	186	10	150	7.23	0
Translation	*glyS*	0	0	2	31	7	250	5.2	0.0007
Translation	*ileS*	0	0	1	2	4	84	4.47	0.0284
Translation	*infB*	0	0	2	12	5	57	3.64	0.0057
Translation	*proS*	0	0	2	0	5	71	3.92	0.0088
Translation	*rplB*	0	0	3	134	6	74	3.97	0.0074
Translation	*rplC*	1	1	1	6	4	38	5.25	0.046
Translation	*rpsD*	0	0	1	13	3	242	5.94	0.0284
Translation	*thrS*	2	4	6	5855	10	207	5.69	0.0001
Translation	*valS*	2	3	4	2446	7	147	5.61	0.0031
Translation	*glyQ*	0	0	2	55	3	133	4.79	0.0071
Unknown	*yfgM*	1	2	3	7	5	98	5.61	0.0077