The most developmentally truncated fishes show extensive

Hox gene loss and miniaturized genomes

Martin Malmstrøm[1,2,*], Ralf Britz[3], Michael Matschiner[1,2], Ole K. Tørresen[1], Renny

K. Hadiaty[4], Norsham Yaakob[5], Heok H. Tan[6], Kjetill S. Jakobsen[1], Walter

Salzburger[1,2] & Lukas Rüber[7,8,*]

[1]Centre for Ecological and Evolutionary Synthesis (CEES), Department of Biosciences, University

of Oslo, Oslo, Norway. [2]Zoological Institute, University of Basel, Basel, Switzerland. [3]Department

of Life Sciences, Natural History Museum, London, UK.[4]Museum Zoologicum Bogoriense,

Research Center for Biology, Indonesian Institute of Sciences, Cibinong, Indonesia. [5]Forest

Research Institute Malaysia, Kuala Lumpur, Malaysia. [6]Lee Kong Chian Natural History Museum,

National University of Singapore, Singapore. [7]Naturhistorisches Museum der Burgergemeinde

Bern, Bern, Switzerland. [8]Institute of Ecology and Evolution, University of Bern, Bern,

Switzerland. *Corresponding authors Ma.M. and L.R.

Abstract

Hox genes play a fundamental role in regulating the embryonic development of

all animals. Manipulation of these transcription factors in model organisms has

unraveled key aspects of evolution, like the transition from fin to limb. However,

by virtue of their fundamental role and pleiotropic effects, simultaneous

knockouts of several of these genes pose significant challenges. Here, we report

on evolutionary simplification in two species of the dwarf minnow genus

Paedocypris using whole genome sequencing. The two species feature

unprecedented *Hox* gene loss and genome reduction in association with their

massive developmental truncation. We also show how other genes involved in

the development of musculature, nervous system, and skeleton have been lost in

Paedocypris, mirroring its highly progenetic phenotype. Further, we identify two

mechanisms responsible for genome streamlining: severe intron shortening and

reduced repeat content. As a naturally simplified system closely related to

zebrafish, *Paedocypris* provides novel insights into vertebrate development.

Introduction

The developmental mechanisms that determine how genotypes translate into phenotypes and how selection acts on genotypes to shape morphological differences are central to our understanding of the diversity of living organisms[1,2]. Model organisms have been instrumental in our quest to decipher how genetic differences are associated with morphological and physiological disparity[3], and improved technologies for genetic modifications will aid in further elucidating genotype-phenotype interrelations[4]. One interesting example is the recent work by Nakamura *et al.* (2016), which utilized CRISPR-Cas9 to knock out three *Hox13* copies (*Aa, Ab,* and *D*) in zebrafish (*Danio rerio*) to investigate the role of these genes in the transition from fins to limbs. Although experiments like these are now feasible, the pleiotropic effects of genes involved in fundamental developmental processes, such as those of the large *Hox* gene family, present challenges and limitations to the phenotypic variation that can easily be induced in model organisms through genetic engineering. Examining naturally occurring extreme phenotypes in close relatives of model organisms thus provides a novel source of phenotypic and genotypic variation that is becoming increasingly important in improving our understanding of the molecular basis of evolutionary changes[5].

Discovered around 200 years ago[6], the zebrafish has been used as a molecular model organism since the 1980s and is currently one of the most important model systems for studying vertebrate development, genome evolution, toxicology, physiology, behavior, and disease[7,8]. Comparative efforts have so far focused on closely related *Danio* species[9,10], but recent studies have revealed that the diversity of zebrafish mutants is surpassed by the range of phenotypic variation among several of its related species in the wild[3,11,12]. Some of these other members of Cyprinidae (e.g *Paedocypris*, *Sundadanio*, and *Danionella*) are characterized by developmental truncation[13-16] and morphological novelties[16-18] and thus offer additional, underappreciated potential for comparative studies and promise fundamental and transformative advances in our understanding of the molecular underpinnings of evolutionary change leading to novelty and adaptation at a genetic and phenotypic level[3].

The Series Otophysi, which includes Cyprinidae, represents one of the

earliest diverging teleostean lineages, and the phylogeny of this lineage remains

controversial[13,19-23]. One particularly difficult taxon to place confidently is the

recently discovered miniaturized dwarf minnow genus *Paedocypris*[13,15,21,22,24]

found in the highly acidic blackwater of endangered peat swamp forests in

Southeast Asia. This genus of tiny vertebrates includes the world's smallest fish,

maturing at ~8 mm[25]. *Paedocypris* exhibits an extreme case of organism-wide

progenesis or developmental truncation, resulting in an anatomical adult

condition closely resembling that of a 7.5 mm zebrafish larva, with over 40 bones

not developed[15]. To investigate the genomic signatures of developmental

truncation, we sequenced and compared the genomes of two representatives of

the genus *Paedocypris*; *P. carbunculus* and *P. micromegethes*. The genome

signatures of these two species allow the distinction of genus-specific from

species-specific genomic changes and thus enable identification of features

associated with the extreme phenotype of *Paedocypris*. By comparing their

genomes with those of other teleosts, including the closely related zebrafish, we

identify *Paedocypris*-specific genomic signatures of developmental truncation in

the form of loss of various key developmental genes, mirroring their progenetic

phenotype. We further demonstrate how the *Paedocypris* genome size has been

reduced through shorter genes due to significantly shorter introns compared to

zebrafish, while exon lengths and gene numbers have remained relatively

unchanged. We find that the *Paedocypris* genomes are comparable in size and

gene structure (i.e short introns and compact genomes) to those of the two

pufferfish model organisms; *Takifugu rubripes* and *Dichotomyctere nigroviridis*,

which have the smallest vertebrate genomes known[26]. Additionally, we show

that the accumulation of transposable elements (TEs), especially DNA

transposons, is very low, following the diversification of the genus *Paedocypris*,

and propose a potential mechanism for enhanced transposon-silencing activity

through duplication of the PIWI-like (*PIWIl1*) gene in this lineage. Highly

progenetic fish species like *Paedocypris* will be important resources for future

studies on vertebrate development, presenting a novel opportunity to

investigate phenotype–genotype relationships in early vertebrate development

and the genetic mechanisms of developmental truncation.

Results

Loss of *Hox*- and other developmental genes

Hox genes encode transcription factors essential in body patterning along the
anterior-posterior body axis during early development of all animals[27]. Although
a large number of genes is involved in developmental processes, *Hox* genes are
especially interesting as they remain organized in conserved clusters and are
ordered along the chromosomes according to where and when they are
activated[28].

As all known *Paedocypris* species have a progenetic phenotype, we
investigated whether this condition is reflected in their *Hox* gene repertoire. We
compared 70 *Hox* gene transcripts, encoded by the 49 *Hox* genes in zebrafish, to
detect syntenic *Hox* cluster regions in the two *Paedocypris* genomes (Methods
Supplementary Note and Supplementary Table 1). For *Hox* clusters *Aa*, *Ab*, and
Bb, we recover all genes on one to three contiguous sequences in synteny with
Danio rerio (Figure 1a). Two scaffolds in both *Paedocypris* species cover the
HoxBa cluster; however, functional copies of *hoxB10a* and *hoxB7a* cannot be
identified in this region. In total, 10 of the zebrafish *Hox* genes appear to be
absent in *Paedocypris*, and their predicted positions are illustrated as red lines in
Figure 1a*.* While the remaining *Hox* clusters are recovered as more fragmented,
most of the genetic regions are covered by contiguous sequences containing
intact copies of other *Hox* genes in at least one of the two *Paedocypris* species.
None of the 10 *Hox* genes could be detected in other parts of the genomes by
sequence similarity searches, strengthening the hypothesis that these genes are
indeed lost in this lineage. Although extensive *Hox* gene loss has been reported
for other cyprinids, these are secondary losses, following genome duplication
events[29]. Figure 1b shows the positions of both the missing and the present *Hox*
genes in *Paedocypris* compared to currently available otophysan genomes,
illustrating the most plausible reconstruction of *Hox*-cluster evolution in this fish
lineage.

Figure 1 | Sequence similarity plots and reconstructed evolutionary history of teleost *Hox* genes. a) Pairwise sequence similarity between zebrafish (*D. rerio*) and *P. carbunculus* (top) and *P. micromegethes* (bottom) for syntenic *Hox* regions. Sequence similarity from 50–100% is shown with green bars representing the location of complete copies of recovered *Hox* genes, while red bars represent the expected location of the unrecovered genes. Grey shading represent contiguous sequences in either of the *Paedocypris* species with sufficient similarity to be mapped to the zebrafish chromosomes. b) Cladogram of selected otophysan species including hypothetical ancestors and ancestral losses (topology according to Stout *et al.* 2017). Genus-specific *Hox* gene losses are shown in red. Only intact gene copies are shown. *Hox* cluster information from other species, and the ancestral states, are based on Henkel *et al.* (2012)[29], Pascual-Anaya *et al.* (2013)[30], Yang *et al.* (2016)[31], and Ensembl[32].

Given its progenetic phenotype and the fact that ~20% of the zebrafish *Hox* gene repertoire is apparently absent in *Paedocypris*, we also investigated the presence

of other key genes involved in various developmental pathways. We first

determined the overlap in gene space between the two *Paedocypris* species, the

zebrafish, and the two tetraodontid models *Di. nigroviridis* and *T. rubripes,* as

these pufferfishes have similarly small genomes as the *Paedocypris* species.

Figure 2a shows the number of shared orthogroups for all species, identified

using OrthoFinder[33], and highlights the number of orthogroups without

orthologs in *Paedocypris* (numbers in bold). We used the set of genes from these

1,160 orthogroups to identify a comprehensive list of 1,581 genes that had gene

ontologies associated with different system development pathways and pattern

specific processes in *D. rerio* (Methods and Supplementary Table 2). These 1,581

genes were then used as queries to screen the genomic sequences and annotated

protein sets of *Paedocypris*. Fourteen of these genes, primarily involved in

skeletal- (*ucmab*, *disc1* and *lbh*), muscle- (*ccdc78* and *marcksa*), and nervous

system (*nkx2.9*, *pho*, *cntf*, *thfrsf1b*, *tnfrsf19* and *lepa*) development, were not

found in either of the *Paedocypris* species (Figure 2b). Adjacent flanking genes in

zebrafish could, however, be identified in *Paedocypris* for all 14 genes

(Supplementary Table 3).

163
164
Figure 2 | Gene loss and duplication of developmental process genes. a) Five-species

comparison of shared orthogroups for the identification of genes lost in *Paedocypris*. Bold

numbers represent orthogroups without any *Paedocypris* orthologs. Orthogroups were identified

using OrthoFinder[33] on the basis of full protein datasets for all included species. b) Lost

developmental pathway genes and schematic representation of phenotypically affected body

segments in *Paedocypris*. c) Gene duplicates retained in *Paedocypris*. d) Genus specific gene

duplications. Genes in bold are associated with a truncated phenotype in *D. rerio*.

Because phenotypic changes can also result from gene duplication[34,35] , we

identified genes with two copies in *Paedocypris*, but only a single copy in

zebrafish and pufferfishes. Based on the topology of gene trees generated for

these genes, we differentiated between genes originating from duplication

events predating the divergence of *Paedocypris* and *Danio*, where only one copy

is retained in zebrafish (Figure 2c), and *Paedocypris*-specific gene duplication

events (Figure 2d). Interestingly, although only four genes could be identified as

recent duplicates in *Paedocypris*, two of these (*ttc21b* and *ruvbl1*) are associated

with deformed phenotypes in zebrafish, including shortening of the anterior-

posterior axis and decreased head size[32].

Genome miniaturization through intron shortening

In parallel to their miniaturized body size, the two *Paedocypris* species show a

surprising evolutionary trajectory in terms of genome miniaturization.

Compared to the genome sizes of zebrafish (~1,5 Gb)[7,32] and other cyprinid

fishes (0.81–3.5 Gb)[36], we find that the two *Paedocypris* species have

substantially smaller genomes (0.45–0.52 Gb), yet similar numbers of genes

(Table 1). Comparative analyses of vertebrate genomes have shown that genome

size reduction is typically characterized by shorter introns and reduced repeat

content[37,38]. However, substantial loss of protein-coding genes[38,39], large

segmental deletions facilitated by fission of macrochromosomes[38], and a reduced

rate of large insertions have also been demonstrated to play a role in reducing or

constricting genome size[37,40].

As the *Paedocypris* genome sizes are comparable to those of tetraodontid

pufferfishes, we compared the gene repertoire of *Di. nigroviridis* and *T. rubripes*

to that of *Paedocypris* and *D. rerio* to determine what genomic features are

shared among these lineages. In order to obtain a consistent gene set for

comparative analysis of gene-, exon-, and intron lengths, we categorized the full

gene set of all five species into orthologous groups using the software

OrthoFinder[33]. We identified 10,368 orthogroups containing genes from all five

species (Figure 2a), which included 16,142 genes in zebrafish, 15,287 in *P.*

carbunculus, 14,181 in *P. micromegethes*, 14,529 in *Di. nigroviridis*, and 14,393 in

T. rubripes. The total gene length and the proportions of exonic and intronic

regions for this gene set are shown in Figure 3a. Although the accumulative

length of this gene set is significantly smaller in both *Paedocypris* species and the

two tetraodontids than in the zebrafish, the proportional change is more subtle,

with the gene set constituting on average 29–30% of the total genome length in

Paedocypris and the tetraodontids and 40% in zebrafish. Further, even though

we detected minor differences in this gene set regarding the average number of

exons per gene in *Paedocypris* (10.04) compared to *D. rerio* (11.21), the observed

47% reduction in total exon length of *Paedocypris* in relation to zebrafish cannot

be attributed to exon loss alone, as the two tetraodontids show very similar

results to *Paedocypris* with even higher average exon count (11.89). However,

the majority of the total gene length reduction observed in *Paedocypris* and

tetraodontids is due to an 80–84% reduction in overall intron size (Figure 2a).

To rule out that the overall reduction in intron size is driven by a highly deviant

fraction of the *Paedocypris* and tetraodontid gene sets, we also investigated the

distribution of gene-, exon-, and intron-lengths of this common gene set (Fig 2b).

We observe that zebrafish has substantially fewer short genes compared to all

other species, with an average gene length 4.3–5.5 longer than that of

Paedocypris and tetraodontids. This is not unexpected, as previous studies have

reported a lineage-specific expansion of intron size[41], resulting in an additional

peak of intron lengths between 1,000–2,000 bp in zebrafish, shown in Figure 3b.

Based on our results, it is apparent that the reduced average gene length in

Paedocypris is driven by a substantial shift towards consistently shorter introns,

similar to, but not as extreme as, in the two tetraodontids. This is further

illustrated by the calculated median sizes depicted in Fig. 3b.

a

D. rerio
91.8%   559 kb

P. carbunculus
80.4%   125 kb

P. micromegethes
80.7%   127 kb

Di. nigroviridis
74.7%   100 kb

T. rubripes
78.0%   120 kb

P. carbunculus female   5 mm

P. micromegethes female

Total exons   Total introns

b

Genes
18,190 bp
5,773 bp
5,019 bp
4,643 bp
3,683 bp

Exons
134 bp
124 bp
123 bp
122 bp
120 bp

Introns
1,169 bp
279 bp
283 bp
144 bp
120 bp

Count (thousands)

Length (bp)   0   50,000
Length (bp)   0   500
Length (bp)   0   6,000

D. rerio   *P. micromegethes*   *P. carbunculus*   *T. rubripes*   *Di. rubripes*   *Di. nigroviridis*

Figure 3 | Comparative analyses of repeat content and gene space in zebrafish and *Paedocypris*. a) Total gene length of 14–16,000 genes belonging to common orthogroups, and the proportional contribution of exonic and intronic regions, in zebrafish, *Paedocypris*, and tetraodontids. b) Frequency plot and median values of gene-, exon-, and intron lengths for the common gene set. In all distributions, the values for both *Paedocypris* species are significantly lower than in zebrafish, but significantly greater than those of both tetraodontids. (Wilcoxon Rank Sum test, $p < 10^{-15}$). Significant results were not detected for intron- and exon lengths between the two *Paedocypris* species; however, in terms of gene lengths, *P. micromegethes* was reported to be significantly longer.

Repeat content reduction and genome evolution

The genome size of an organism evolves through the relative rate of insertions and deletions and through the effect of natural selection that either favors or eliminates these changes. Genome shrinkage has thus been postulated to evolve through a bias in the rate of insertions relative to deletions[42], but the impact of this mechanism, and its importance in genome size evolution is still debated[43]. Importantly, even though deletions indeed appear to be more frequent than insertions, the latter tend to include significantly more base pairs, resulting in the gradual increase in genome size in eukaryotes[42]. Although several other types of mutational activity can promote genome-size expansion, self-replicating mobile elements (*i.e.* transposons) have been identified as the most prominent

contributor in this regard[44,45], and recent studies have confirmed a strong

correlation between genome size and the amount of transposable elements (TEs)

in teleost fishes and other vertebrates[46,47]. Because the more compact genomes

have previously been shown to harbor fewer repeats[38], we investigated the

degree to which this was the case in *Paedocypris*. Using a *Danio*-specific repeat

library, we determined the total amount of TEs; DNA-transposons, long- and

short interspersed repeats (LINEs and SINEs), and long terminal repeats (LTRs)

in the genomes of *Paedocypris*, zebrafish, and four other cyprinid species

(*Cyprinus carpio*, *Pimephales promelas*, *Sinocyclocheilus grahami*, and *Leuciscus*

waleckii) in addition to the characiform cave tetra (*Astyanax mexicanus*) using

Repeatmasker[48]. The repeat landscape graphs, illustrating the relative amount of

each TE class and the Kimura distance[49] for each of these are shown in Figure 4a

along with a time-calibrated phylogeny of these eight species, as inferred using

BEAST[50] (see Methods). Figure 4a also shows the percentage of each repeat type

of various age categories, calculated following Kapusta, *et al.* 2017)[47]. The total

number of repetitive elements of each class is shown in Figure 4b, clearly

illustrating that even though the proportional differences in repeat content is

similar between *Paedocypris* and the other cyprinids, the total number of

elements is much lower.

Figure 4 | Repeat landscape graphs showing the prevalence and relative age of the four

main repeat classes in *Paedocypris* compared to six other teleost genome assemblies. a)

Phylogeny with divergence times for the internal nodes for the eight species analyzed. Colored

numbers on the branch leading to *Paedocypris* show the percentages of each of the

corresponding repeat classes that originate from the time interval corresponding to this branch

(117.7–79.9, 79.9–12.3, and more recent than 12.3 Ma). Repeat landscapes represent

transposable elements of the four main classes as well as the unclassified ones. The x-axis

indicates the Kimura distance[49] as a proxy for time while the y-axis gives the relative coverage of
each repeat class based on the genome size. b) Total number of repetitive elements in each of the
cyprinid genomes (excluding zebrafish); colored by class.

Consistent with previous reports[46], we find the that the zebrafish genome is

dominated by repetitive elements, constituting 58.1% of its genome. In contrast,

only 15.3 –15.5% of the *Paedocypris* genomes consist of repetitive elements

(transposable elements, satellites, simple repeats, and low complexity regions), a

percentage that is comparable to that of the other cyprinids investigated (12.3–

19.01%). However, in terms of TE content, the *Paedocypris* species have

considerably fewer and shorter elements that comprise in total 7.24 and 7.30%

of their genomes, compared to the four other cyprinids in which we find on

average a genomic proportion of 13.81% (not including zebrafish in which these

constitute 53.28% of the genome)(Supplementary Table 4). Based on the timing

of the divergence events leading to crown *Paedocypris* and the calculated

substitution rate (see Methods), we investigated the proportions of "ancient" and

"lineage specific" TEs. Interestingly, only 0.05 – 0.25% of the *Paedocypris*

transposons appear to have been incorporated into their genomes after these

two species diverged (Figure 4).

As *Paedocypris* thus appears to have reversed the general trend of DNA

gain[43,43] towards gradual DNA loss, as indicated by their low TE content, we

investigated whether this could have been achieved through silencing of

transposon activity[40]. We thus explored the genomic content of the PIWI-like

genes (*PIWIl1* and *PIWIl2*), which are known for silencing transposons in

zebrafish and other vertebrates[51,52]. Interestingly, we identified additional copies

of *PIWIl1* in both *Paedocypris* genomes, located in a conserved region ~4 Mb

upstream of *PIWIl1* on zebrafish chromosome 8, as shown in Figure 5. One of

these gene duplications is shared between the two *Paedocypris* species and

therefore appears to have evolved prior to their divergence, while an additional,

third copy is found exclusively in *P. carbunculus* (Fig. 5a). These lineage-specific

duplications suggest that the gradual genome miniaturization has been, at least

in part, attained through increased transposon silencing, leading to a bias of DNA

loss over DNA gain. This needs to be tested further, utilizing genomic sequences

from other closely related cyprinids like *Danionella* and *Sundadanio*. Probably

resulting from the lineage-specific whole genome duplication, a single duplicate

of the *PIWIl1* gene was also found in *C. carpio,* but no extra copies could be

detected in the other cyprinids; *P. promelas*, *L. waleckii*, or *S. graham*i. No

additional copies of *PIWIl2* were detected in either *Paedocypris* or the other

cyprinids.

Figure 5 | Synteny of the *PIWIl1* loci and surrounding regions in the zebrafish and

***Paedocypris* genomes.** a) Synteny between zebrafish (top) and *P. carbunculus*, showing the

original *PIWIl1* copy (right red rectangle) and the two additional copies (left and middle red

rectangles) detected in this genome assembly. b) Same as a) for *P. micromegethes*, showing the

single, presumably ancestral, duplicated copy of *PIWIl1* in this genome. Transparent red boxes

depict the exon-intron structure of all *PIWIl1* genes in *Paedocypris* with black lines representing

the scaffolds covering each region. Not all exons could be recovered for all gene copies.

Chromosome fusion

Another interesting feature of the miniaturized genome of *Paedocypris* is the

reduced number of chromosomes compared to that of zebrafish. While most

Cypriniformes have ≥ 24 chromosomes[53] in the haploid metaphase and zebrafish

has 25, *P. carbunculus* only features 15 chromosomes[54]. Since we did not find any

indication of large-scale chromosome loss, as would be indicated by substantial

gene loss, this discrepancy implies that genome miniaturization in *Paedocypris*

spp. has been accompanied by chromosomal fusion since they diverged from

their last common ancestor with zebrafish. In order to identify regions on

different chromosomes in the zebrafish genome that are now tightly linked in the

Paedocypris genome, we looked for syntenic regions in *Paedocypris* that contain

at least six genes from two different chromosomes in zebrafish. Figure 6 shows

the three scaffolds identified in each of the *Paedocypris* species fulfilling these

criteria. In both species, nine genes from chromosomes 9 and 21 in zebrafish are

co-localized on a single scaffold. For chromosome 11 in zebrafish, we find good

evidence in different regions for a fusion with chromosome 1 in *P. carbunculus*

but also chromosome 2 in *P. micromegethes*, suggesting that either all three

chromosomes may have fused or that species-specific fusion events may have

occurred. Although the remaining two putative fusion events can only be

detected in one of the species, there are no contradicting results between the

two *Paedocypris* species, indicating that fusion of the chromosome pairs 6 and 19

and 22 and 25 is likely to have occurred in the last common ancestor of both

species.

Figure 6 | Syntenic regions between zebrafish chromosomes and *Paedocypris* scaffolds.

352 Syntenic regions between zebrafish (*D. rerio*) and *P. carbunculus* (a) or *P. micromegethes* (b).

353 Links between *Paedocypris* scaffolds and *D. rerio* chromosomes indicate orthologous genes. Only

354 the relevant parts of *D. rerio* chromosomes are shown and the locations of all genes in these

355 regions are illustrated in the gray shaded areas. Numbers on scaffolds and chromosomes are

356 given as kbp.

Discussion

Miniature body size has evolved several times within Cyprinidae[13]. However,

little is known about the molecular underpinnings of these miniaturization

events, whether these are similar in several instances of miniaturization, and to

what extent the parallel evolution of genomic and phenotypic traits is coupled.

Given the identification of several classical signatures of genome reduction,

including reduced TE content and smaller introns, we conclude that the small

Paedocypris genomes represent a derived state, having evolved from an ancestor

with a substantially larger genome and a larger body size. Reduced genome size

has also been reported in other vertebrates[37,38], yet these reductions include

large segmental deletions, chromosome fissions, and massive losses of protein

coding genes, which we do not observe in the miniaturized genomes of

Paedocypris. Moreover, while reduced genome sizes in birds and bats can

probably be explained by the adaptation to flight through the need for higher

metabolic rates[55], the causes of genome size reduction in fishes seem in general

less apparent. However, the reduced genome sizes in *Paedocypris* may also have

evolved as a response to a selection for higher metabolic rate, mediated in this

case by its extreme habitat (*i.e.* low oxygen concentrations and pH < 4.0) rather

than rapid locomotion.

Small body size has arisen among miniature cyprinids in two different

ways, either as proportionate dwarfism, resulting in miniature but otherwise

identical copies of their larger ancestors, or through developmental truncation

or progenesis leading to larval appearance of the tiny mature fish[13,14].

Evolutionary novelties such as highly modified pelvic and pectoral girdles[14], an

anterior shift in genital pore and anus[13], and fang-like structures forming from

the jaw bones[16], however, appear to be restricted to developmentally truncated

miniatures (*e.g. Paedocypris*, *Danionella,* and *Sundadanio*). The absence of

novelties in proportioned dwarfs would suggest that developmental truncation

plays an important role in escaping evolutionary and developmental constraints

imposed by the species' *Bauplan* and in opening up new evolutionary avenues

for drastic morphological change[14,16,18].

This study represents the first report on genomes of highly

developmentally truncated fish species. It illustrates how the unique features of

Paedocypris at the phenotypic level (*i.e.* extreme miniaturization and dramatic

developmental truncation) are paralleled at the genomic level, in the form of

substantially smaller genomes via reduced intron lengths and lower repeat

content, and also through the loss of numerous genes related to development,

including highly important *Hox* genes. Although a causal link between

miniaturized body size and reduced genome size remains to be established, our

study highlights the potential to investigate such connections in a comparative

framework, including the sequencing of additional species representing other

miniature genera. Nevertheless, these naturally simplified species, having

escaped the developmental, and, as we report here, genetic constraints of the

cyprinid *Bauplan*, present a novel opportunity to investigate phenotype–

genotype relationships in vertebrate development. These naturally occurring

extreme phenotypes will greatly aid future research on model organisms and our

quest to understand how phenotypic diversity is generated during vertebrate

evolution and development.

406   Recent investigation of the Gulf pipefish genome[56] has revealed a loss of the

last *Hox7* paralog in this species. This has revived the disputed hypothesis that

the lack of ribs in both pufferfish and the pipefishes can be attributed to the fact

that none of these species have any remaining *Hox7* paralogs[57]. Interestingly, the

Paedocypris species show a similar loss of the last *Hox7* paralog (*hoxb7a*), yet the

two species investigated here do feature ribs. The ribs are, however, reduced and

remain poorly ossified[14], suggesting that although the *Hoxa7/Hoxb7* genes are

not the only essential genes for rib development, they do appear to have an

influence on their development, as suggested by earlier experiments in mice[58].

415   Finally, the observed loss of *Hox13* genes (*Cb* and *Da*) in the two

Paedocypris species is of special interest as these genes represent the

termination of the patterning system. *Hoxd13* was also one of the genes targeted

in the successful CRISPR-Cas9 knockout experiment in the zebrafish[59], that

showed that the *Hox* genes play a very similar role in patterning during

development of both fins bones, and thus elucidated the evolutionary transition

from fins to limbs. We propose that future studies of similar nature should

consider using *Paedocypris* species as a complementary system to the zebrafish

model, due to its already limited *Hox* gene repertoire, yet relatively close

phylogenetic relationship with zebrafish.

Materials and methods

Specimens used

P. carbunculus was caught at the type locality in Pangkalanbun, Kalimantan

Tengah, on the island of Borneo and *P. micromegethes* was caught in Sibu,

Sarawak, Malaysia, on the island of Borneo. Both species were caught using dip

nets. Immediately upon capture, specimens were killed by an overdose of

anesthesia (MS222) following guidelines by the American Society of

Ichthyologists and Herpetologists (ASIH) (http://www.asih.org/pubs/; issued

2013). Individuals were preserved in 96% ethanol for subsequent DNA analyses.

DNA isolation

A single whole specimen of each *Paedocypris* species, stored on 96% ethanol,

was used for isolation of high molecular genomic DNA with the EZNA Tissue

DNA Kit (Omega Bio-Tek, Norcross, Gerogia, USA), following manufacturer`s

instructions. Sample identifiers were LR12004 (*Paedocypris carbunculus*) and

LR7898 (*Paedocypris micromegethes*), supplied by Lukas Rüber.

Sequencing library preparation

Genomic DNA samples were fragmented to lengths of ~550 bp by sonication on a

Covaris E220 (Life Technologies, Carlsbad, California, USA) with the following

settings: 200 cycles for 45 seconds with Peak Incident Power of 175 W and

frequency sweeping mode. All sequencing libraries were constructed following

Illumina's TrueSeq PCR-free library preparation protocol for 550 bp fragments.

Whole genome sequencing

Based on expected genome sizes of 315-350 Mb[54], both *Paedocypris* genomes

were sequenced to ~90✕ coverage on the Illumina HiSeq 2500 platform, with

the Illumina 500 cycles kit (Rapid mode) with on-board clustering. Per species,

this produced 151-155 M paired reads of 250 bp each.

The Kapa Library quantification kit for Illumina (Kapa Biosciences, Wilmington,

Massachusetts, USA) was used to find the correct molarity (nM) before

sequencing.

Genome assembly

Both *Paedocypris* genomes were initially assembled using two different assembly

programs, the "Overlap-Layout-Consensus" based Celera Assembler[60] and the

"de Bruijn graph" based DISCOVAR *de novo*[61]. We then used the Metassembler

software[62] to merge and optimize these two assemblies, producing a

reconsolidated single assembly with superior quality for each of the two

Paedocypris species (Table 1, Supplementary Note, and Supplementary Table 5).

Assembly quality assessment

The quality of the different genome assemblies was assessed by comparing the

proportion of conserved genes detected, as a measure of gene-space

completeness. We used the program BUSCO v2.0 (Benchmarking Universal

Single-Copy Orthologs)[63], which searches for 4,584 highly conserved single-copy

actinopterygian genes. Results are listed in Table 1.

We also assessed the assembly quality of the three different assembly

versions with the software FRC*bam* [64], which identifies "features" (incorrectly

mapped reads, incorrect insert size, coverage issues, etc.) in each of the

assemblies, and ranks the different assembly versions according to the number

of detected features. Additional information on the execution of these programs

is available in the Supplementary Note and the resulting graphs are presented in

Supplementary Figs. 1 and 2.

480
Annotation

Structural and functional annotation of both reconsolidated *Paedocypris*

genomes was performed with two iterative rounds of the MAKER2 (v 2.31.8)[65-67]

pipeline, following the instructions by Sujai Kumar (available at

https://github.com/sujaikumar/assemblage/blob/master/README-

annotation.md). Numbers of genes annotated are listed in Table 1. See

Supplementary Note for in-depth descriptions of additional software and

commands used.

489
490   **Table 1**
491   Assembly statistics, gene-space completeness-, and annotation metrics.

	P. carbunculus	*P. micromegethes*
Total assembly length (bp)	430,790,821	414,707,736
Estimated assembly coverage (CA / DDN)	67× / 73×	76× / 91×
Number of scaffolds	18,953	15,158
N50 scaffold length	59,252	61,901
Longest scaffold (bp)	653,275	678,207
BUSCOs found[2]	4,113	4,073
BUSCOs complete[2]	3,719	3,564

Total gene count[3]	25,567	25,453

Hox gene search

In addition to the two *Paedocypris* genomes, *Hox* gene content was also investigated in the genomes of *Pimephales promelas* (GCA_000700825.1_FHM_SOAPdenovo_genomic.fna) and *Leuciscus waleckii* (GCA_900092035.1_Amur_ide_genome_genomic.fa) based on BLAST[68] searches, using 70 zebrafish *Hox* transcripts from Ensembl as queries (including some truncated variants and isoforms), representing the 49 unique *Hox* genes (Supplementary Table 1). The similarity stringency threshold used in these searches was $1e^{-20}$. We also conducted additional searches using Exonerate v2.2.0[69] for genes whose presence could not be established based on the BLAST search. For those genes that could not be detected using either method, additional searches were conducted using the orthologous protein sequence from the cave tetra (*Astyanax mexicanus*). Contiguous sequences from both *Paedocypris* genomes were extracted from the scaffolds assembled with Metassembler, spanning all hit regions plus 10 kb upstream and downstream of each hit. These scaffold sequences were aligned to the orthologous sequences extracted from the zebrafish genome (GRCz10) using mVista[70].

Identification of lost and expanded developmental genes

In order to obtain a complete list of genes from *D. rerio* that were associated with various developmental pathways that could be compromised in *Paedocypris*, we started out with the orthogroups found not to contain orthologs from either *Paedocypris* species, as identified using OrthoFinder[33]. We further utilized the gene ontology information associated with all genes belonging to these orthogroups, and identified key gene ontology terms: GO:0009948 (anterior–posterior axis formation), GO:0009950 (dorsal–ventral axis formation), GO:0040007 (growth), GO:0007517 (muscle organ development), GO:0007399 (nervous system development), GO:0001501 (skeletal system development), and GO:0007379 (segment specification). This data set contained 1,581 unique genes from *D. rerio*, and 64 of these genes could not be detected in the *Paedocypris*

genome sequences using TBLASTN[68] with a similarity cutoff of $1e^{-20}$, and their

presence was further examined by running Exonerate[69] (v2.2.0). Based on the

Exonerate results, reciprocal BLAST searches, annotation, and identification of

flanking genes, we could confidently determine that 14 of these genes were

indeed not present in either of the *Paedocypris* genomes. Using expression data

from Ensembl[32], we reconstructed the hypothetically affected body segments in

Paedocypris, as illustrated in Figure 2b.

We further investigated whether we could confidently identify

developmental genes that are now duplicated in *Paedocypris* but not in *D. rerio*.

From the full list of orthogroups, we identified 138 groups that were represented

by a single copy in the two tetraodontid pufferfishes and in *D. rerio*, but by two

apparent copies in both *Paedocypris* species. The alignments of these genes were

then screened for missing data, and based on manual inspection of these

alignments, only alignments with < 35% missing data were included in further

analysis. Out of the 35 genes that met this criterion, 19 could be confirmed

through annotation, and had a gene tree consistent with the two hypotheses

outlined in Figure 2c and 2d.

Gene space evolution

In order to assess the changes of gene-, exon-, and intron sizes in *Paedocypris* we

first identified the proteomic overlap of *Paedocypris*, *D. rerio*, *Di. nigroviridis*, and

T. rubripes by running the software OrthoFinder[33] on the complete protein sets

of these five species. We used the full protein sets from Ensembl[32] (v. 80): *D.*

rerio (GRCz10) *, Di. nigroviridis* (TETRAODON8), and *T. rubripes*(FUGU4).

However, as some of the *D. rerio* genes have more than one protein or transcript

in the Ensembl database, the output from BioMart (31,953 genes and 57,349

proteins) was filtered so that only the longest protein sequence from each gene

was used in the analysis, and genes without protein sequences were removed.

This resulted in a set of 25,460 genes with a single protein prediction. For the

Paedocypris species, the "standard" gene sets resulting from the annotation were

used as input. These sets were filtered to include only genes with AED

(Annotation Edit Distance) scores < 1 or those with a Pfam domain. By using only

genes belonging to the 10,368 orthogroups found to contain orthologs from all

these species (Fig. 2a), we obtained a comprehensive but conservative dataset as

the basis for these analyses. Information about each of the corresponding genes

in *D. rerio*, *Di. nigroviridis*, and *T. rubripes* was obtained from BioMart, and

included the Ensembl gene and protein ID, and the chromosome name in

addition to the start and stop position for each gene, transcript, and exon. Intron

sizes were then calculated on the basis of exon positions, using a custom script

("gene_stats_from_BioMart.rb"). In some cases, the sum of exons and introns did

not equal the total length of a gene, which appears to be caused by inconsistency

in the registration of UTR regions in the Ensembl database for individual genes.

In these cases, to be conservative with regard to intron length estimates, the

gene length was shortened to correspond to the sum of the exons and the

corresponding introns between these.

Intron and exon lengths for the two *Paedocypris* species were calculated

in a similar manner, but on the basis of the "standard" filtered annotation file in

"gff" format produced as part of the annotation pipeline. Also for these species,

the intron lengths were determined on the basis of identified exons, with another

custom script ("gene_stats_from_gff.rb"). Gene-, exon-, and intron length

histograms were plotted with the R package ggplot2. Custom Ruby scripts are

available for download at https://github.com/uio-cees/Paedocypris_gene_stats

Repeat content analysis

The repeat contents of the two *Paedocypris* genomes and the model organism

genomes were assessed using RepeatMasker (v4.0.6)[48] with the *Danio*-specific

repeat library included in the program, and with the "-s" setting to increase

sensitivity. The following model organism genome assemblies were used in this

comparison: *D. rerio* (Danio_rerio.GRCz10.dna.toplevel.fa), *P. promelas*

(GCA_000700825.1_FHM_SOAPdenovo_genomic.fna), *L. waleckii*

(GCA_900092035.1_Amur_ide_genome_genomic.fa), *Sinocyclocheilus grahami*

(GCA_001515645.1_SAMN03320097.WGS_v1.1_genomic.fna), *Cyprinus carpio*

(GCA_001270105.1_ASM127010v1_genomic.fna), and *A. mexicanus*

(Astyanax_mexicanus.AstMex102.dna.toplevel.fa). Repeat landscape graphs (Fig.

4a) were plotted with the R package ggplot2 based on the ".aligned" output file

from RepeatMasker. The proportion of repeats originating from specific time

intervals was calculated using the parseRM.pl script[47] (available at

https://github.com/4ureliek/Parsing-RepeatMasker-Outputs), using the

Paedocypris-specific substitution rate and the "--age" setting set to "12.31,79.98",

according to divergence times estimated with BEAST2. All four repeat classes

were analyzed individually using the "--contain" setting.

Calculation of substitution rate for *Paedocypris*

The *Paedocypris*-specific substitution rate was calculated on the basis of a whole

genome alignment of the two *Paedocypris* species. These alignments were

created by mapping the sequencing reads of *P. micromegethes* to the *P.*

carbunculus assembly using BWA (Burrows-Wheeler Aligner) (v. 0.7.12)[71] and

SAMTOOLS (v. 1.3.1)[72,73]. The number of nucleotide differences in each of the

18,953 alignments (one per scaffold) was identified using a custom script

("find_variable_sites.rb", available for download at https://github.com/uio-

cees/Paedocypris_gene_stats). The substitution rate per million years

(0.001288) was then calculated as the sum of total differences (13,415,043)

divided by the total number of aligned sites (422,829,036) and two times the

estimated crown age of *Paedocypris* (12.31), as inferred by the BEAST v.2.4.5[50]

analysis.

Phylogenetic inference of selected otophysan species

To allow the estimation of divergence times between the two *Paedocypris*

species and six other otophysan taxa for which genomic resources were available

(*D. rerio*, *A. mexicanus, C. carpio*, *P. promelas*, *S. grahami*, and *L. waleckii*), we

followed the pipeline for phylogenetic marker selection presented in Malmstrøm

et al. (2016 and 2017)[74,75] with few modifications. These modifications included

the following changes: Marker selection began with a set of 3,238 cave fish (*A.*

mexicanus) exons, for which at least five orthologs were known among the seven

species *D. rerio*, *Gadus morhua, Gasterosteus aculeatus*, *Oreochromis niloticus*,

Oryzias latipes, *Poecilia formosa*, and *T. rubripes*, according to version 87 of the

Ensembl database. This marker set was then used to identify potential orthologs

from the genomes of the eight selected otophysan taxa based on TBLASTN[68]

searches followed by a strict filtering procedure. Compared to Malmstrøm et al.

(2016), we applied a lower dN/dS threshold of 0.25 to exclude markers

potentially affected by positive selection, and we removed all markers for which

no homologs could be detected in one or more of the eight otophysan genomes.

We also applied stricter thresholds on clock-like evolution of candidate markers,

so that all genes with an estimated coefficient of rate variation above 0.8 as well

as those with a mean mutation rate above 0.0004 per site per million years were

excluded. We identified 138 genes with a total alignment length of 135,286 bp

which were subsequently used for analysis with BEAST v.2.4.5[50]. Since the

topology of otophysan taxa has previously been resolved with a more

comprehensive phylogenetic dataset[21], we here focused on the inference of

divergence times only, by using the topology inferred by Stout et al. (2016) as a

starting tree and excluding all of BEAST2's operators on the tree topology.

Divergence times were estimated by calibrating the most recent common

ancestor of Cypriniformes and Characiformes with a lognormal distribution

centered at 121 Ma (standard deviation on log scale: 0.1) according to the results

of Malmstrøm et al. (2016). We performed two replicate BEAST2 analyses with

800 million MCMC iterations, of which the first 100 million were discarded as

burnin. Convergence was assessed based on similarity of parameter traces

between run replicates and effective sample sizes (ESS) greater than 200. A

maximum clade credibility (MCC) summary tree with node heights according to

mean age estimates was produced with TreeAnnotator v.2.1.2[50].

Identification of PIWI-like genes

We investigated the presence of PIWI-like genes in the genomes of the two

Paedocypris species and the other cyprinids using Exonerate[69] (v2.2.0) with the

longest transcripts available for the two PIWI-like homologs from zebrafish;

PIWIl1 (ENSDARG00000041699) and *PIWIl2* (ENSDARG00000062601). Regions

containing sequences spanning more than three introns were aligned to the

zebrafish exons using mafft[76] as implemented in AliView[77] (v1.17.1) . Intron

sequences were aligned manually based on the established exon structure, using

the full-length scaffold sequences. Local gene synteny to zebrafish chromosome

8, surrounding the putative *PIWIl1* copies, was confirmed through reciprocal

BLAST searches using both the MAKER2 annotated proteins and proteins

predicted by GeneScan (online version)[78] as queries.

Identification of chromosomal rearrangements (fusions)

As *P. carbunculus* has been shown to have a haploid chromosome count of 15

[54], potential chromosomal fusions were investigated on the basis of disrupted

synteny of zebrafish chromosomes in relation to *Paedocypris*.

We identified putative homologous regions between the zebrafish genome

assembly and each of the *Paedocypris* species' genome assemblies by using

MCScanX[79]. In short, the predicted proteins for each *Paedocypris* species were

merged with predicted proteins from zebrafish into a single file, and BLASTP[68]

was executed with this file as both query and target, thus identifying putative

homologs both within each species and between. The genomics positions of the

proteins were extracted from the annotation files, and the BLASTP results and

the genomic positions were provided to MCScanX for identifying the putative

homologous regions.

Acknowledgments

Fieldwork in the peat swamp forests in Malaysia and Indonesia was funded by

the Natural Environmental Research Council (NERC; NE/F003749/1, to L.R. and

R.B.), National Geographic (8509-08, to L.R.), and the North of England

Zoological Society (to L.R.). Fieldwork in Sarawak was conducted under permits

issued by the Economic Planning Unit, Prime Minister's Department, Malaysia

UPE 40/200/19/2534) and the Forest Department Sarawak (NCCD.970.4.4[V]-

43) and fieldwork in Indonesia was conducted under permits issued by the

Indonesian Institute of Sciences (LIPI) and the Kementerian Negara Riset dan

Teknology (RISTEK; 3/TKPIPA/FRP/SM/III/2012). We thank E. Adamson H.

Budianto H. Ganatpathy, S. Lavoué, M. Lo, H. Michael, and S. Sauri for their help in

the field. All computational work was performed on the Abel Supercomputing

Cluster (Norwegian metacenter for High Performance Computing (NOTUR) and

the University of Oslo) operated by the Research Computing Services group at

USIT, the University of Oslo IT-department. Sequencing library creation and

high-throughput sequencing were carried out at the Norwegian Sequencing

Centre (NSC), University of Oslo, Norway. This work was funded by grants from

the Naturhistorisches Museum der Burgergemeinde Bern to L.R. and the

Research Council of Norway (RCN grants 199806 and 222378) to K.S.J. H.H.T

acknowledges funding from the National University of Singapore (NUS, R-154-

000-318-112) and Lee Kong Chian Natural History Museum. W.S. acknowledges

funding from the European Research Council (ERC) and the Swiss National

Science Foundation (SNF).

Competing financial interests

699 The authors declare no competing financial interests.

References

1. Carroll, S. P., Hendry, A. P., Reznick, D. N. & Fox, C. W. Evolution on
ecological time-scales. *Func. Ecol.* **21,** 387–393 (2007).
2. Moczek, A. P., Sultan, S., Foster, S., Ledón-Rettig, C., Dworkin, I., Nijhout, H.
F., Abouheif, E. & Pfennig, D. W. The role of developmental plasticity in
evolutionary innovation. *Proc. R. Soc. B.* **278,** 2705–2713 (2011).
3. Braasch, I., Peterson, S. M., Desvignes, T., McCluskey, B. M., Batzel, P. &
Postlethwait, J. H. A new model army: emerging fish models to study the
genomics of vertebrate Evo-Devo. *J. Exp. Zool.* **324,** 316–341 (2014).
4. Shalem, O., Sanjana, N. E. & Zhang, F. High-throughput functional genomics
using CRISPR-Cas9. *Nat. Rev. Genet.* **16,** 299–311 (2015).
5. Streelman, J. T., Peichel, C. L. & Parichy, D. M. Developmental genetics of
adaptation in Fishes: the case for novelty. *Annu. Rev. Ecol. Evol. Syst.* **38,**
655–681 (2007).
6. Hamilton, F. *An Account of the Fishes Found in the River Ganges and Its*
Branches. Hurst, Robinson, and Co, Edinburgh (1822).
7. Howe, K., Clark, D. M., Torroja, C. F., Torrance, J., Berthelot, C., Muffato, M.,
Collins, J. E., Humphray, S., McLaren, K., Matthews, L., McLaren, S., Sealy, I.,
Caccamo, M., Churcher, C., Scott, C., Barrett, J. C., Koch, R., Rauch, G-J.,
White, S., Chow, W. *et al.* The zebrafish reference genome sequence and its
relationship to the human genome. *Nature* **496,** 498–503 (2013).
8. Kettleborough, R. N. W. Busch-Nentwich, E. M., Harvey, S. A., Dooley, C. M.,
de Bruijn, E., van Eeden, F., Sealy, I., White, R. J., Herd, C., Nijman, I. J.,
Fényes, F., Mehroke, S., Scahill, C., Gibbons, R.,n Wali, N., Carruthers, S.,
Hall, A., Yen, J., Cuppen, E. & Stemple, D. L. A systematic genome-wide
analysis of zebrafish protein-coding gene function. *Nature* **496,** 494–497
(2013).
9. McMenamin, S. K., Bain, E. J., McCann, A. E., Patterson, L. B., Eom, D. S.,
Zachary P. Waller, Z. P., Hamill, J. C., Kuhlman, J. A., Eisen, J. S. & Parichy, D.
M. Thyroid hormone-dependent adult pigment cell lineage and pattern in
zebrafish. *Science* **345,** 1358–1361 (2014).
10. McCluskey, B. M. & Postlethwait, J. H. Phylogeny of zebrafish, a "model

species," within *Danio,* a 'model genus'. *Mol. Biol. Evol.* **32,** 635–652
(2015).

11.   Patterson, L. B., Bain, E. J. & Parichy, D. M. Pigment cell interactions and
differential xanthophore recruitment underlying zebrafish stripe
reiteration and *Danio* pattern evolution. *Nat. Commun.* **5,** 5299 (2014).

12.   Parichy, D. M. The natural history of model organisms: advancing biology
through a deeper understanding of zebrafish ecology and evolution. *eLife*
4, e05635 (2015).

13.   Rüber, L., Kottelat, M., Tan, H. H., Ng, P. K. L. & Britz, R. Evolution of
miniaturization and the phylogenetic position of *Paedocypris*, comprising
the world's smallest vertebrate. *BMC Evol. Biol.* **7,** 38 (2007).

14.   Britz, R. & Conway, K. W. Osteology of *Paedocypris*, a miniature and highly
developmentally truncated fish (Teleostei: Ostariophysi: Cyprinidae). *J.*
Morphol. **270,** 389–412 (2009).

15.   Britz, R., Conway, K. W. & Rüber, L. Miniatures, morphology and molecules:
Paedocypris and its phylogenetic position (Teleostei, Cypriniformes). *Zool.*
J. Linnean Soc. **172,** 556–615 (2014).

16.   Britz, R. & Conway, K. W. *Danionella dracula*, an escape from the
cypriniform Bauplan via developmental truncation? *J. Morphol.* **277,** 147–
166 (2015).

17.   Conway, K. W. & Britz, R. Sexual dimorphism of the Weberian apparatus
and pectoral girdle in *Sundadanio axelrodi* (Ostariophysi: Cyprinidae). *J.*
Fish Biol. **71,** 1562–1570 (2007).

18.   Conway, K. W., Kubicek, K. M. & Britz, R. Morphological novelty and modest
developmental truncation in *Barboides*, Africa's smallest vertebrates
(Teleostei: Cyprinidae). *J. Morphol.* **14,** 33–18 (2017).

19.   Fang, F., Norï n, M., Liao, T. Y., Källersjö, M. & Kullander, S. O. Molecular
phylogenetic interrelationships of the south Asian cyprinid genera *Danio*,
Devario and *Microrasbora* (Teleostei, Cyprinidae, Danioninae). *Zool.*
Scripta **38,** 237–256 (2009).

20.   Tang, K. L., Agnew, M. K., Vincent Hirt, M., Sado, T., Schneider, L. M.,
Freyhof, J., Sulaiman, Z., Swartz, E., Vidthayanon, C., Miya, M., Saitoh, K.,
Simons, A. M., Wood, R. M. & Mayden, R. L. Systematics of the subfamily
Danioninae (Teleostei: Cypriniformes: Cyprinidae). *Mol. Phylogenet. Evol.*
57, 189–214 (2010).

21.   Stout, C. C., Tan, M., Lemmon, A. R., Lemmon, E. M. & Armbruster, J. W.
Resolving Cypriniformes relationships using an anchored enrichment
approach. *BMC Evol. Biol.* **16,** 244 (2016).

22.   Arcila, D., Ortí, G., Vari, R., Armbruster, J. W., Stiassny, M. L. J., Ko, K. D.,
Sabaj, M. H., Lundberg, J., Revell, L. J. & Betancur-R, R. Genome-wide
interrogation advances resolution of recalcitrant groups in the tree of life.
Nat. Ecol. Evol. **1,** 0020 (2017).

23.   Hirt, M. V., Arratia, G., Chen, W. & Mayden, R. L. Effects of gene choice, base
composition and rate heterogeneity on inference and estimates of
divergence times in cypriniform fishes. *Biol. J. Linnean Soc.* (2017).
doi.org/10.1093/biolinnean/blw045

24.   Mayden, R. L. & Chen, W.-J. The world's smallest vertebrate species of the
genus *Paedocypris*: a new family of freshwater fishes and the sister group
to the world's most diverse clade of freshwater fishes (Teleostei:

782   Cypriniformes). *Mol. Phylogenet. Evol.* **57,** 152–175 (2010).

783   25.   Kottelat, M., Britz, R., Hui, T. H. & Witte, K.-E. *Paedocypris*, a new genus of
784   Southeast Asian cyprinid fish with a remarkable sexual dimorphism,
785   comprises the world's smallest vertebrate. *Proc. Biol. Sci.* **273,** 895–899
786   (2006).

787   26.   Jaillon, O., Aury, J.-M., Brunet, F., Petit, J.-L., Stange-Thomann, N., Mauceli,
788   E., Bouneau, L., Fischer, C., Ozouf-Costaz, C., Bernot, A., Nicaud, S., Jaffe, D.,
789   Fisher, S., Lutfalla, G., Dossat, C., Segurens, B., Dasilva, C., Salanoubat, M.,
790   Levy, M., Boudet, N. *et al.* Genome duplication in the teleost fish *Tetraodon*
791   *nigroviridis* reveals the early vertebrate proto-karyotype. *Nature* **431,**
792   946–957 (2004).

793   27.   Mallo, M. & Alonso, C. R. The regulation of *Hox* gene expression during
794   animal development. *Development* **140,** 3951–3963 (2013).

795   28.   Gaunt, S. J. The significance of *Hox* gene collinearity. *Int. J. Dev. Biol.* **59,**
796   159–170 (2015).

797   29.   Henkel, C. V., Burgerhout, E., de Wijze, D. L., Dirks, R. P., Minegishi, Y.,
798   Jansen, H. J., Spaink, H. P., Dufour, S., Weltzien, F.-A., Tsukamoto, K. & van
799   den Thillart G. E. E. J. M. Primitive duplicate *Hox* clusters in the European
800   Eel's genome. *PLoS ONE* **7,** e32231–9 (2012).

801   30.   Pascual-Anaya, J., D'Aniello, S., Kuratani, S. & Garcia-Fernàndez, J.
802   Evolution of *Hox* gene clusters in deuterostomes. *BMC Dev. Biol.* **13,** 26
803   (2013).

804   31.   Yang, J., Chen, X., Bai, J., Fang, D., Qiu, Y., Jiang, W., Yuan, H., Bian, C., Lu, J.,
805   He, S., Pan, X., Zhang, Y., Wang, X., You, X., Wang, Y., Sun, Y., Mao, D., Liu, Y.,
806   Fan, G., Zhang, H. *et al.* The *Sinocyclocheilus* cavefish genome provides
807   insights into cave adaptation. *BMC Biol.* **14,** 1–13 (2016).

808   32.   Cunningham, F., Amode, M. R., Barrell, D., Beal, K., Billis, K., Brent, S.,
809   Carvalho-Silva, D., Clapham, P., Coates, G., Fitzgerald, S., Gil, L., Girón, C. G.,
810   Gordon, L., Hourlier, T., Hunt, S. S., Janacek, S. H., Johnson, N., Juettemann,
811   T., Kähäri, A. K., Keenan, S. *et al.* Ensembl 2015. *Nucleic Acids Res.* **43,**
812   D662–669 (2015).

813   33.   Emms, D. M. & Kelly, S. OrthoFinder: solving fundamental biases in whole
814   genome comparisons dramatically improves orthogroup inference
815   accuracy. *Genome Biol.* **16,** 157 (2015).

816   34.   Kaessmann, H. Origins, evolution, and phenotypic impact of new genes.
817   *Genome Res.* **20,** 1313–1326 (2010).

818   35.   Peterson, T. & Müller, G. B. Phenotypic novelty in EvoDevo: the distinction
819   between continuous and discontinuous variation and its importance in
820   evolutionary theory. *Evol. Biol.* **43,** 314–335 (2016).

821   36.   Gregory, T. R. Animal genome size database (Available at
822   http://genomesize.com).

823   37.   Neafsey, D. E. & Palumbi, S. R. Genome size evolution in pufferfish: a
824   comparative analysis of diodontid and tetraodontid pufferfish genomes.
825   *Genome Res.* **13,** 821–830 (2003).

826   38.   Zhang, G., Li, C., Li, Q., Li, B., Larkin, D. M., Lee, C., Storz, J. F., Antunes,A.,
827   Greenwold, M. J., Meredith, R. W., Ödeen, A., Cui, J., Zhou, Q., Xu, L., Pan, H.,
828   Wang, Z., Jin, L., Zhang, P., Hu, H., Yang, W. *et al.* Comparative genomics
829   reveals insights into avian genome evolution and adaptation. *Science* **346,**
830   1311–1320 (2014).

831 39. Hughes, A. L. & Friedman, R. Genome size reduction in the chicken has
832   involved massive loss of ancestral protein-coding genes. *Mol. Biol. Evol.* **25,**
833   2681–2688 (2008).

834 40. Vinogradov, A. E. Evolution of genome size: multilevel selection, mutation
835   bias or dynamical chaos? *Curr. Opin. Genet. Dev.* **14,** 620–626 (2004).

836 41. Moss, S. P., Joyce, D. A., Humphries, S., Tindall, K. J. & Lunt, D. H.
837   Comparative analysis of teleost genome sequences reveals an ancient
838   intron size expansion in the zebrafish lineage. *Genome Biol. Evol.* **3,** 1187–
839   1196 (2011).

840 42. Petrov, D. A. Mutational equilibrium model of genome size evolution.
841   *Theor. Pop. Biol.* **61,** 531–544 (2002).

842 43. Gregory, T. R. Insertion–deletion biases and the evolution of genome size.
843   *Gene* **324,** 15–34 (2004).

844 44. Sun, C., Shepard, D. B. & Chong, R. A. LTR retrotransposons contribute to
845   genomic gigantism in plethodontid salamanders. *Genome Biol.* **4,** 168-183
846   (2012).

847 45. Chénais, B., Caruso, A., Hiard, S. & Casse, N. The impact of transposable
848   elements on eukaryotic genomes: from genome size increase to genetic
849   adaptation to stressful environments. *Gene* **509,** 7–15 (2012).

850 46. Chalopin, D., Naville, M., Plard, F., Galiana, D. & Volff, J. N. Comparative
851   analysis of transposable elements highlights mobilome diversity and
852   evolution in vertebrates. *Genome Biol. Evol.* **7,** 567–580 (2015).

853 47. Kapusta, A., Suh, A. & Feschotte, C. Dynamics of genome size evolution in
854   birds and mammals. *Proc. Natl. Acad. Sci. USA* **114,** E1460–E1469 (2017).

855 48. Smith, A.F.A., Hubley R., & Green, P. RepeatModeler Open-1.0. (2015)
856   (Available at http://www.repeatmasker.org)

857 49. Kimura, M. A simple method for estimating evolutionary rates of base
858   substitutions through comparative studies of nucleotide sequences. *J. Mol.*
859   *Evol.* **16,** 111–120 (1980).

860 50. Bouckaert, R., Heled, J., Kühnert, D., Vaughan, T., Wu, C.-H., Xie, D., Suchard,
861   M. A., Rambaut, A. & Drummond, A. J. BEAST 2: A software platform for
862   Bayesian evolutionary analysis. *PLoS Comput. Biol.* **10,** e1003537 (2014).

863 51. Aravin, A. A., Hannon, G. J. & Brennecke, J. The Piwi-piRNA pathway
864   provides an adaptive defense in the transposon arms race. *Science* **318,**
865   761–764 (2007).

866 52. Levin, H. L. & Moran, J. V. Dynamic interactions between transposable
867   elements and their hosts. *Nat. Rev. Genet.* **12,** 615–627 (2011).

868 53. Yu, X., Zhou, T., Li, K., Li, Y. & Zhou, M. On the karyosystematics of cyprinid
869   fishes and a summary of fish chromosome studies in China. *Genetica* **72,**
870   225–235 (1987).

871 54. Liu, S., Hui, T. H., Tan, S. L. & Hong, Y. Chromosome evolution and genome
872   miniaturization in minifish. *PLoS ONE* **7,** e37305 (2012).

873 55. Zhang, Q. & Edwards, S. V. The Evolution of intron size in amniotes: a role
874   for powered flight? *Genome Biol. Evol.* **4,** 1033–1043 (2012).

875 56. Small, C. M., Bassham, S., Catchen, J., Amores, A., Fuiten, A. M., Brown, R. S.,
876   Jones, A. G. & Cresko, W. A. The genome of the Gulf pipefish enables
877   understanding of evolutionary innovations. *Genome Biol.* **17,** 258 (2016).

878 57. Amores, A., Suzuki, T., Yan, Y.-L., Pomeroy, J., Singer, A., Amemiya, C. &
879   Postlethwait, J. H. Developmental roles of pufferfish *Hox* clusters and

genome evolution in ray-fin fish. *Genome Res.* **14,** 1–10 (2004).

58.   Chen, F., Greer, J. & Capecchi, M. R. Analysis of *Hoxa7/Hoxb7* mutants
suggests periodicity in the generation of the different sets of vertebrae.
Mech. Dev. **77,** 49–57 (1998).

59.   Nakamura, T., Gehrke, A. R., Lemberg, J., Szymaszek, J. & Shubin, N. H.
Digits and fin rays share common developmental histories. *Nature* **537,**
225–228 (2016).

60.   Miller, J. R., Delcher, A. L., Koren, S., Venter, E., Walenz, B. P., Brownley, A.,
Johnson, J., Li, K., Mobarry, C. & Sutton G. Aggressive assembly of
pyrosequencing reads with mates. *Bioinformatics* **24,** 2818–2824 (2008).

61.   Weisenfeld, N. I., Yin, S., Sharpe, T., Lau, B., Hegarty, R., Holmes, L., Sogoloff,
B., Tabbaa, D., Williams, L., Russ, C., Nusbaum, C., Lander, E. S., MacCallum I.
& Jaffe, D. B. Comprehensive variation discovery in single human genomes.
Nat. Genet. **46,** 1350–1355 (2014).

62.   Wences, A. H. & Schatz, M. C. Metassembler: merging and optimizing de
novo genome assemblies. *Genome Biol.* **16,** 207 (2015).

63.   Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V. & Zdobnov,
E. M. BUSCO: assessing genome assembly and annotation completeness
with single-copy orthologs. *Bioinformatics* **31,** 3210–3212 (2015).

64.   Vezzi, F., Narzisi, G. & Mishra, B. Reevaluating assembly evaluations with
feature response curves: GAGE and assemblathons. *PLoS ONE* **7,** e52210
(2012).

65.   Cantarel, B. L., Korf, I., Robb, S. M. C., Parra, G., Ross, E., Moore, B., Holt, C.,
Alvarado, A. S. & Yandell, M. MAKER: An easy-to-use annotation pipeline
designed for emerging model organism genomes. *Genome Res.* **18,** 188–
196 (2007).

66.   Holt, C. & Yandell, M. MAKER2: An annotation pipeline and genome-
database management tool for second-generation genome projects. *BMC*
Bioinform. **12,** 491 (2011).

67.   Campbell, M. S., Law, M. Y, Holt, C., Stein, J. C., Moghe, G. D., Hufnagel, D. E.,
Lei, J., Achawanantakun, R., Jiao, D., Lawrence, C. J., Ware, D., Shiu, S.-H.,
Childs, K. L., Sun, Y., Jiang, N. & Yandell, M. MAKER-P: A tool kit for the
rapid creation, management, and quality control of plant genome
annotations. *Plant Physiol.* **164,** 513–524 (2014).

68.   Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. Basic local
alignment search tool. *J. Mol. Biol.* **215,** 403–410 (1990).

69.   Slater, G. & Birney, E. Automated generation of heuristics for biological
sequence comparison. *BMC Bioinform.* **6,** 31 (2005).

70.   Frazer, K. A., Pachter, L., Poliakov, A., Rubin, E. M. & Dubchak, I. VISTA:
computational tools for comparative genomics. *Nucleic Acids Res.* **32,**
W273–W279 (2004).

71.   Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-
Wheeler transformation. *Bioinformatics* **25,** 1754–1760 (2009).

72.   Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G.,
Abecasis, G., Durbin, R. & 1000 Genome Project Data Processing Subgroup.
The sequence alignment/map format and SAMtools. *Bioinformatics* **25,**
2078–2079 (2009).

73.   Li, H. A statistical framework for SNP calling, mutation discovery,
association mapping and population genetical parameter estimation from

sequencing data. *Bioinformatics* **27,** 2987–2993 (2011).

74.   Malmstrøm, M., Matschiner, M., Tørresen, O. K., Star, B., Snipen, L. G.,
Hansen, T. F., Baalsrud, H. T., Nederbragt, A. J., Hanel, R., Salzburger, W.,
Stenseth, N. C., Jakobsen, K. S. & Jentoft, S. Evolution of the immune system
influences speciation rates in teleost fishes. *Nat. Genet.* **48**, 1204–1210
(2016).

75.   Malmstrøm, M., Matschiner, M., Tørresen, O. K., Jakobsen, K. S. & Jentoft, S.
Whole genome sequencing data and de novo draft assemblies for 66
teleost species. *Sci. Data* **4,** 160132–13 (2017).

76.   Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software
version 7: improvements in performance and usability. *Mol. Biol. Evol.* **30,**
772–780 (2013).

77.   Larsson, A. AliView: a fast and lightweight alignment viewer and editor for
large datasets. *Bioinformatics* **30,** 3276–3278 (2014).

78.   Burge, C. & Karlin, S. Prediction of complete gene structures in human
genomic DNA. *J. Mol. Biol.* **268,** 78–94 (1997).

79.   Wang, Y., Tang, H., DeBarry, J. D., Tan, X., Li, J., Wang, X., Lee, T.-H., Jin, H.,
Marler, B., Guo, H., Kissinger, J. C. & Paterson, A. H. MCScanX: a toolkit for
detection and evolutionary analysis of gene synteny and collinearity.
Nucleic Acids Res. **40,** e49 (2012).