1

Cyanobacterial Growth on Municipal Wastewater Requires

Low Temperatures

Travis C. Korosh[a,b], Andrew Dutcher[c], Brian F. Pfleger [a,d,], and Katherine D. McMahon [c,e] *

[a] Department of Chemical and Biological Engineering, University of Wisconsin-Madison, Madison, WI 53706, United States
[b] Environmental Chemistry and Technology Program, University of Wisconsin-Madison, Madison, WI 53706, United States
[c] Department of Civil and Environmental Engineering, University of Wisconsin-Madison, Madison, WI 53706, United States
[d] Microbiology Doctoral Training Program, University of Wisconsin-Madison, Madison, WI 53706, United States
[e] Department of Bacteriology, University of Wisconsin-Madison, Madison, WI 53706, United States

* Corresponding author. 5552 Microbial Science Building, 1550 Linden Drive, Madison, WI 53706, United States. Phone: +1 608 890 2836. Fax: +1 608 262-9865. E-mail address: trina.mcmahon@wisc.edu.

Keywords: Nutrient removal, cyanobacteria, Synechococcus strain PCC 7002

ABSTRACT

Side-streams in wastewater treatment plants can serve as concentrated sources of nutrients (i.e. nitrogen and phosphorus) to support the growth of photosynthetic organisms that ultimately serve as feedstock for production of fuels and chemicals. However, other chemical characteristics of these streams may inhibit growth in unanticipated ways. Here, we evaluated the use of liquid recovered from municipal anaerobic digesters via gravity belt filtration as a nutrient source for growing the cyanobacterium *Synechococcus* sp. strain PCC 7002. The gravity belt filtrate (GBF) contained high levels of complex dissolved organic matter (DOM), which seemed to negatively influence cells. We investigated the impact of GBF on physiological parameters such as growth rate, membrane integrity, membrane composition, photosystem composition, and oxygen evolution from photosystem II. At 37°C, we observed an inverse correlation between GBF concentration and membrane integrity. Radical production was also detected upon exposure to GBF at 37°C. However, at 27°C the dose dependent relationship between GBF concentration and lack of membrane integrity was abolished. Immediate resuspension of strains in high doses of GBF showed markedly reduced oxygen evolution rates relative to the control. Together, this suggests that one mechanism responsible for GBF toxicity to *Synechococcus* is the interruption of photosynthetic electron flow and subsequent phenomena. We hypothesize this is likely due to the presence of phenolic compounds within the DOM.

IMPORTANCE

Cyanobacteria are viewed as promising platforms to produce fuels and/or high-value chemicals as part of so-called "bio-refineries". Their integration into wastewater treatment systems is particularly interesting because removal of the nitrogen and phosphorus in many wastewater

streams is an expensive but necessary part of wastewater treatment. In this study, we evaluated

strategies for cultivating *Synechococcus* strain PCC 7002 on media comprised of two wastewater

streams; treated secondary effluent supplemented with the liquid fraction extracted from sludge

following anaerobic digestion. This strain is commonly used for metabolic engineering to produce

a variety of valuable chemical products and product precursors (e.g. lactate). However, initial

attempts to grow PCC 7002 under otherwise standard conditions of light and temperature failed.

We thus systematically evaluated alternative cultivation conditions and then used multiple

methods to dissect the apparent toxicity of the media under standard cultivation conditions.

INTRODUCTION

The need to develop non-petroleum based platforms for fuel and chemical production is driving researchers to explore alternatives that harness renewable energy sources while minimizing other environmental impacts such as freshwater depletion, eutrophication, and the use of arable land for non-food production. Cyanobacteria are particularly attractive such platforms due to their genetic tractability, rapid growth rates, halotolerance, and ability to be grown on non-productive land with simple nutrient requirements (1, 2). According to published life cycle assessments, a large portion of the associated costs of culturing photoautotrophs are tied to upstream costs, such as CO_2 delivery and fertilizer application (3). High phosphorus/nitrogen removal rates and energy efficiencies have been reported for photobioreactor and open pond cultivation systems using wastewater streams rich in nitrogen and phosphorus (4, 5). Therefore, it may be possible to offset the requirement for fertilizer by reclaiming nutrients from wastewater. This approach could yield the sought-after non-petroleum based alternative while also providing a more effective means of nutrient and metal removal than conventional wastewater treatment (6–8). Side-streams from common wastewater treatment facilities such as supernatants or filtrates from solids-separation processes are particularly promising nutrient sources, assuming that cyanobacterial strains can efficiently use them.

Of the many streams available in common wastewater facilities, the liquid fraction of anaerobic digestate is thought to be the most attractive nutrient source (8–11). Although digestate is rich in the inorganic constituents necessary for growth, it also contains dissolved organic matter (DOM) that has been shown to limit photosynthetic activity (12). DOM is a heterogenous mixture of aliphatic and aromatic compounds derived from the decomposition of living organisms (13). The chemical nature of wastewater-derived DOM is largely governed by the type of treated waste

and the treatment process, but it is largely comprised of hydrophilic, fulvic, and humic substances

(14, 15). Humics can induce damaging permeability in model and bacterial membranes (16, 17).

Various studies have also demonstrated that fulvic and humic acids can enhance the solubility of

many organic compounds (18), which in turn would augment their bioavailability and potential

membrane permeability. Many of these compounds are also photo-reactive, producing toxic

hydrogen peroxide and hydroxyl radicals (19, 20), which is of significant concern for phototroph

cultivation.

The mode of DOM toxicity is thought to involve interactions with the protein-pigment

complex of photosystem II (PSII) in photosynthetic organisms, although the exact molecular

mechanism remains unclear (21). When the rate of light induced damage to PSII exceeds its rate

of repair, growth suppression and chlorosis result from the phenomena known as photoinhibition

(22). When damage by photoinhibition is sufficient to hamper the natural ability to consume

electrons generated by photosynthesis, reactive oxygen species (ROS) are concomitantly produced

as an undesired by-product. Prolonged oxidative stress halts protein translation through oxidation

of specific cysteine residues in the ribosomal elongation factors (23). Given these findings, it is

increasingly evident that under conditions of sustained stress, regulation of electron flow is critical

to maintain homeostasis in photosynthetic organisms (24, 25). Thus, it is important to understand

the mechanisms by which DOM may be interrupting electron flow in order to capitalize on the

potential of cyanobacteria to remediate wastewater and generate high-value chemicals.

In this study, we tested the practicality of using combined streams from a municipal

wastewater plant as a nutrient source for cyanobacterial cultivation. We used the euryhaline

cyanobacterium *Synechococcus* sp. strain PCC 7002 (PCC 7002) due to its exceptional tolerance

to high light intensity, salt, and other environmental stresses (26, 27). Initial attempts to grow PCC

7002 in this nutrient source under standard environmental conditions for this strain (1% (v/v) CO_2,

a temperature of 37°C, and illumination of 200 μmol photons $m^{-2}s^{-1}$) resulted in photobleached

(white-yellow) cultures. In an effort to explain this observation while developing more feasible

cultivation conditions, we assessed the effects of many environmental parameters during PCC

7002 cultivation in wastewater-based media by monitoring changes in growth rate, photopigment

abundance, oxygen evolution rates, membrane integrity, and membrane composition. High GBF

concentrations were associated with elevated DOM levels. Decreased photosynthetic oxygen

production rates were also noted upon exposure to high levels of GBF. We observed marked

membrane permeability, photosystem degradation, low growth rates, and ROS production in

cultures exposed to GBF at 37°C. In comparison, at 27°C the cultures grown on GBF had reduced

membrane permeability, robust growth rates, as well as high levels of total fatty acids and an

elevated unsaturated fatty acid content relative to control. This suggests bioavailability of the

photoinhibitory compound in GBF is governed by changes in membrane content and composition

that occur during growth.

MATERIALS AND METHODS

Media and Growth Conditions

Synechococcus sp. strain PCC 7002 was obtained from the Pasteur Culture Collection of

Cyanobacteria. Experiments were performed with a strain of PCC 7002 harboring the *aaC1* gene

encoding a gentamicin resistance marker in the A2842 locus (*glpK*) to maintain axenic cultures.

Strains were grown and maintained on solid media composed of Medium A^+ (28) supplemented

with 5 μM $NiSO_4$ (29) with 1.5% Bacto-Agar. Strains were cultured from a single colony in 250

ml baffled flasks with 50 mL media with 1% CO_2-enriched air at 150 rpm in a Kuhner ISF1-X

orbital shaker. Temperature was maintained at 37°C or 27°C and light intensity was fixed at

approximately 200 μmol photons $m^{-2}s^{-1}$ or 100 μmol photons $m^{-2}s^{-1}$ via a custom LED panel.

Strains were pre-acclimated to the culture conditions overnight before inoculating in fresh media.

Optical density at 730 nm was measured in a Tecan M1000 plate reader.

Wastewater-derived media was obtained from the Nine Springs Wastewater Treatment

Plant (Dane County, Wisconsin, USA). The plant is configured for biological nutrient removal via

a modified University of Cape Town process with no internal nitrate recycling and stable

performance yielding high secondary effluent quality (total phosphorus < 1 mg P/L, ammonia < 1

mg N/L, nitrate ~ 15 mg N/L) (30). Anaerobic digesters are used for solids stabilization and the

resulting digested material is passed over a gravity belt filter for dewatering. The system includes

an Ostara WASSTRIP process to recover phosphorus. This filtrate (GBF) served as the primary

source of phosphorus and reduced nitrogen for our cultures, and effluent from the post-mainstream

secondary treatment clarifier (secondary effluent) served as a diluent. GBF was gravity filtered

through a paper coffee filter to remove any exceptionally large flocs, then stored at -80°C until

use. Secondary effluent was collected one to four days before each experiment and held under

refrigeration at approximately 2°C. Experimental media was comprised primarily of secondary

clarifier effluent and GBF, combined in different proportions. Unless otherwise noted, GBF was

used at a concentration of 12.5% (v/v) in secondary effluent, supplemented with trace metals and

vitamin B12 at the concentrations found in Medium A^+, as well as KH_2PO_4 at a molar ratio of 1:32

soluble reactive phosphorus (SRP) to bioavailable nitrogen (the sum of NH_4^+ and NO_3^-). All media

was buffered with Tris-HCl and adjusted to pH 8.0 with potassium hydroxide or hydrochloric acid

before sterilization by autoclaving, and gentamycin was added at working concentrations (30

μg/mL) after cooling.

Staining, Flow Cytometry, and Fluorescence Measurements

The membrane permeability of cyanobacteria cells was recorded by a flow cytometer

(FACSCalibur, BD Biosciences, San Jose, CA, USA). After growth in the respective media, cells

were centrifuged (2 minutes, 5000 RCF), decanted, and resuspended in 1 mL of Tris-Buffered

Saline (TBS) solution (pH 8.0). To identify membrane-compromised cells, SYTOX Green (Life

Technologies) was also added to each sample (1 μM). SYTO 59 (Life Technologies) was added to

each sample (1 μM) as a nucleic acid counterstain. As a control for a permeabilized membrane,

cells were resuspended in 190 proof ethanol. SYTOX green fluorescence was visualized using 488

nm laser excitation and emission area was read using a 530/30 nm bandpass filter. The 633 nm

laser coupled with a 661/16 bandpass filter was used for SYTO 59 visualization. Analysis of the

cytometric data was carried out with CellQuest Pro (BDBiosciences, San Jose,CA, USA) software.

To assess reactive oxygen species production, cells (OD_{730} = 1) were incubated overnight

in Medium A$^+$, 12.5 % (v/v) GBF, 12.5 % (v/v) GBF + 1 mM Dithiothreitol (DTT), 12.5 % (v/v)

GBF + 1 mM N-acetylcysteine (NAC), or Medium A$^+$ + 100 μM methyl viologen as a positive

control at 37°C with 5% CO_2 at a light intensity of 200 μmol photons $m^{-2}s^{-1}$. Cells were washed in

TBS, and either Sytox Green (1 μM) or CellROX Orange reagent (Life Technologies) (5 μM) was

added. After 30 min incubation in darkness at 37°C, fluorescence was measured (Ex/Em 545/565

nm for Cell ROX Orange) and (Ex/Em 504/523 nm for Sytox Green) in a Tecan M1000 plate

reader.

GBF Characterization

SRP, ammonia, nitrate, and nitrite concentrations were determined for all secondary

clarifier effluent and GBF samples used in these experiments. In addition, the GBF was tested for

total suspended solids (TSS), volatile suspended solids (VSS) total solids (TS), and chemical

oxygen demand (COD). Ammonia, SRP, and COD concentrations were determined by

colorimetric tests using reagents from Hach. Nitrate and nitrite were determined using high

performance liquid chromatography (Shimadzu) with a C18 column and photodiode array detector

(31). TSS, VSS, and TS were measured according to Standard Method 2540 D, 2540 E, and 2540

B, respectively, with 47mm diameter glass fiber filters (Whatman) used for TSS and VSS (32).

Fluorescence EEM measurements and UV-Vis absorbance scans were conducted using a M1000

Tecan plate reader using a UV-transparent plate (Costar 3635). Fluorescence intensity was

normalized to quinine sulfate units (QSU), where 1 QSU is the maximum fluorescence intensity

of 1 ppm of quinine sulfate in 0.1 N H_2SO_4 at Ex/Em = 350/450. Rayleigh scatter effects were

removed from the data set.

Biochemical analyses

Batch cultures were further assayed for dry cell weight (DCW), fatty acid content, oxygen

evolution rates, and chlorophyll a content. Cells were concentrated by centrifugation, washed in

TBS, and lyophilized overnight to obtain dry cell weights (DCW). Fatty acids from approximately

10 mg of DCW with 10 mg ml^{-1} pentadecanoic acid as an internal standard were converted to

methyl-esters, extracted with n-hexane and analyzed by GC-FID on a Restek Stabilwax column

(60m, 0.53 mm ID, 0.50 μm) (33). Photosynthetic oxygen evolution from whole cells was

measured with a Unisense MicroOptode oxygen electrode with 10 mM $NaHCO_3$ illuminated with

a slide projector at photosynthetic photon flux densities ranging from 76-2700 μmol

photons photons m^{-2} s^{-1} for 30 min at room temperature (34). Cells were collected by

centrifugation and resuspended in the appropriate media to give an OD_{730} = 1.0. Chlorophyll a

measurements were done via a 100% chilled methanol extraction procedure (35). Chlorophyll a

was calculated via the following equation: $Chl_a = 16.29 * A^{665} - 8.54 * A^{652}$ (36).

RESULTS

GBF and secondary effluent characteristics

We measured nutrient concentrations from the batches of GBF collected over the 6 month

experimental period (Table 1 and Table 2), to ascertain if it was a stable and reliable nutrient source

for cultivating the cyanobacteria. Sampling points from the Nine Springs Wastewater Treatment

Plant (Dane County, Wisconsin, USA) are shown in Figure 1. In order to calculate nutrient

stoichiometries, we took the sum of NH_3-N and NO_3-N to be the bioavailable N. Nutrient levels

in the GBF were markedly more variable than in the secondary effluent. The average molar ratio

of bioavailable N to SRP was 35 ± 7 in GBF (12.5% v/v) diluted with secondary effluent, as

compared to 32 in Medium A^+.

We also measured DOM quality in the secondary effluent and GBF using

excitation−emission matrix (EEM) fluorescence spectroscopy (37) because we hypothesized that

DOM was linked to toxicity during cultivation, as has been shown in prior studies (16, 17, 38).

EEM fluorescence spectroscopy revealed that secondary effluent contained diffuse constituents,

including humic [excitation wavelengths (>280 nm) and emission wavelengths (>380 nm)] and

fulvic acid-like [excitation wavelengths (<250 nm) and emission wavelengths (>350 nm)] spectra

relative to Medium A^+ (Figure 2). The distinction between the two substances is historically based

on solubility (39), but compositionally, fulvic acids contain more acidic functional groups than

humic acids (40). The fulvic acid content in media preparations rose with increasing GBF

concentrations. Additionally, at high concentrations of GBF, we detected absorbance in a distinct

region [excitation wavelengths (270 -290 nm) and emission wavelengths (340-400 nm)] that points

to the presence of soluble microbial products, which include aromatic amino acids, carbohydrates,

or phenols (37, 41). Prior work has shown that the GBF stream of the Nine Springs Treatment

Plant contained high molecular weight DOM, which was dissimilar in composition to other

internal streams (42)

Dose-dependent tolerance to GBF is a function of growth temperature

To evaluate the effects of GBF dosage on PCC 7002 physiology, we measured biomass

accumulation (Figure 3) and growth rates (Table 3) in media with GBF concentrations ranging

from 6.25%-12.5% (v/v), as a function of temperature (27°C vs 37°C) and light intensity (100

μmol photons m^{-2} s^{-1} vs 200 μmol photons m^{-2} s^{-1}). Medium A^{+} served as a control. Higher

temperatures depressed growth rates in GBF-based media. At 37°C, growth rates with 6.25% GBF

at both 100 μmol photons m^{-2} s^{-1} and 200 μmol photons m^{-2} s^{-1} were most comparable to Medium

A^{+}. Higher GBF concentrations had a more extreme effect on growth rates. However, this dose-

dependent effect of GBF on growth rate was abolished when the cultivation temperature was

lowered to 27°C. At 200 μmol photons m^{-2} s^{-1} and 27°C, GBF cultures grew twice as slowly as

the control and there was no significant difference between the tested GBF concentrations. Under

100 μmol photons m^{-2} s^{-1} and 27°C, growth rates were comparable across media conditions. Thus,

successful cultivation using the more concentrated GBF media required adjusting both the light

and temperature regimes.

Dose-dependent membrane permeability of GBF is a function of growth temperature

We wondered whether the decreased growth rates in GBF at 37°C was a result of

membrane permeability, given the known effect of humic acids on membrane integrity (16, 17).

To track the dynamics of GBF induced membrane permeability, we employed forward scatter flow

cytometry using SYTO 59 as a counterstain to identify cells, which were subsequently visualized

for membrane permeability using Sytox Green. As Sytox Green is a membrane impermeable dye

that fluoresces when binding to nucleic acids (43), fluorescence would indicate compromised outer

membrane structures. Two distinct phases were identified upon exposure to GBF, which we

interpreted as initial and chronic membrane permeability (Figure 4). Initial membrane permeability

was defined as the Sytox Green positive events for samples analyzed within the first 10 hours of

growth, while chronic membrane permeability accounted for the Sytox Green positive events

during subsequent time points. As was the case for growth rate, a relationship between both initial

and chronic membrane permeability and increasing GBF concentrations was found at 37°C (Figure

4a). While we still detected considerable initial membrane permeability with GBF exposure at

27°C, chronic membrane permeability decreased over time, mostly likely to the increase in

biomass (Figure 3). Altogether, this suggested that there was a temperature dependent adaptation

that ameliorated the susceptibility of cultures to GBF induced membrane permeability.

Exposure to GBF at high temperatures generates radicals and destroys the photosynthetic

pigments

257   We directly measured the ROS content and membrane permeability in response to

overnight GBF exposure at 37°C using the fluorophores Sytox Green and CellROX Orange (Figure

5). We examined the capacity of reducing agents Dithiothreitol (DTT) or N-acetylcysteine (NAC)

to quench the media toxicity, since they have anti-oxidant properties due to the direct reduction of

disulfide bonds or as precursors for the anti-oxidant glutathione (44). To measure their effect on

the ROS production and culture viability after GBF exposure, we assessed ROS content and

membrane integrity with either concurrent addition or preincubation of these quenching

compounds in the diluted (12.5% v/v) GBF media. Addition of 100 μM methyl viologen to

Medium A[+] served as a positive control. Exposure of cells to 12.5%-GBF media resulted in marked

ROS production and membrane permeability, which no thiol treatment alleviated.

Photobleaching of the photosynthetic pigments is also a common symptom of oxidative

stress in photosynthetic organisms and is caused by the accumulation of ROS (45). To investigate

the effects of GBF media on the photosynthetic pigmentation, we performed whole cell absorbance

scans of cultures cultivated at two different temperatures (27°C vs 37°C) and the same light

intensity (200 μmol photons $m^{-2} s^{-1}$). High GBF concentrations yielded enhanced chlorophyll,

phycobilisome, and carotenoid degradation at 37°C (Figure 7a). At 27°C, photosynthetic pigments

maintained intact relative to the control, regardless of GBF concentration, implying less ROS

production at this temperature (Figure 7b).

Exposure to GBF retards oxygen evolution

To better delineate the cause of initial toxicity associated with high GBF concentrations,

we measured maximal oxygen evolution rates for strains briefly exposed to media with 12.5% v/v

GBF while increasing the light intensity. Measurements of photosynthetic oxygen evolution would

allow for an indirect assessment of PSII activity and electron transfer (46). Cultures were grown

to early linear phase in Medium A[+] or GBF media, washed, and resuspended in the appropriate

media. Resuspension media was saturated with HCO_3^- (10 mM) in order to prevent inorganic

carbon limitation. As expected, cells grown and assayed in Medium A[+] at 37°C showed a clear

increase in O_2 evolution rate as the light intensity approached saturation at 2700 μm photons

$m^{-2} s^{-1}$, reaching a maximal rate of 240 ± 8 μmol O_2 (mg Chl a)$^{-1}$ h^{-1} (Figure 7a). Cells grown in

Medium A[+] at 37°C but assayed in 12.5% GBF had diminished O_2 evolution rates at all light

intensities, plateauing with a rate of 79 ± 6 μmol O_2 (mg Chl a)$^{-1}$ h^{-1} at an intensity of 270 μmol

photons $m^{-2} s^{-1}$ (Figure 7a). Thus, exposure to GBF under these conditions caused an immediate

decrease in O_2 production.

Next, we examined the effect of temperature in a similar experiment. Assays carried out in

Medium A^+ after growth in Medium A^+ at 27°C showed much higher maximal O_2 evolution rates

at all tested light intensities than with cells grown at 37°C (Figure 7b), peaking at a rate 556 ± 102

µmol O_2 (mg Chl a)$^{-1}$ h^{-1}). This was expected because elevated O_2 evolution rates in low

temperature grown cells have been previously reported and were attributed to a substantial change

in photosystem stoichiometry (47). The O_2 evolution rates of cells grown in Medium A^+ at 27°C

and then resuspended in 12.5% GBF stayed relatively constant at all of the tested intensities and

were roughly 5-fold lower than in controls, with a maximal rate of 109 ± 22 µmol O_2 (mg Chl a)$^{-}$

^{1} h^{-1} at an intensity of 75 µmol photons $m^{-2} s^{-1}$. We compared the above rates to those from

cultures grown in 12.5% GBF at 27°C to test if adaptation to GBF was met with changes in

photosynthetic activity. At the examined light intensities, O_2 evolution rates with 27°C GBF-

adapted cultures were not statistically different from those measured in cultures grown at the same

temperature in Medium A^+. Finally, we conducted the inverse experiment, using cultures grown

in GBF media at 27°C but assayed in Medium A^+ under saturating light. Interestingly, they

displayed the highest evolution rate of any tested condition, at 944 ± 96 µmol O_2 (mg Chl a)$^{-1}$ h^{-1}

(Figure 7b). This suggested that there is a period of dynamic photosynthetic adaptation to

overcome the stress of GBF, and that when the stress is removed, the cells have an enhanced

capacity for photosynthetic activity.

Additional assays were performed at light saturation (2700 µmol $m^{-2} s^{-1}$) with cultures

adapted to 37°C (Figure 8) to further examine the effects of media conditions on oxygen evolution

rates. Again, oxygen evolution rates in GBF (73 ± 8 µmol O_2 (mg Chl a)$^{-1}$ h^{-1}) were much lower

than Medium A$^+$ (248 $\pm$ 21 μmol O$_2$ (mg Chl a)$^{-1}$ h^{-1}). In order to rule out the effect of differences

in overall ionic strength, we increased the osmolarity of the GBF medium by adding NaCl to match

the levels found in Medium A$^+$, and found no statistical difference in oxygen evolution as

compared to experiments without added NaCl. Attempts to eliminate the inhibitory effect of GBF

on oxygen evolution by gravity filtration through powdered activated carbon were unsuccessful.

We also measured oxygen evolution rates in control experiments designed to inhibit

photosynthetic electron transfer through 10 μM DCMU addition to Medium A$^+$ (48), and to prevent

de-novo protein synthesis by pre-treating cultures with 800 μg ml^{-1} Lincomycin in Medium A$^+$

(49), prior to incubation at light saturation for 1 hour. However, evolution rates with these controls

were higher than those treated with GBF. Therefore, no definitive conclusion about the molecular

mechanism responsible for reduced oxygen evolution rates in the presence of GBF could be made.

Acclimation to GBF changes lipid content and composition

Based on the results described above, we hypothesized that the temperature dependent

adaptation may be related to changes in membrane content and composition. We extracted total

fatty acids of cultures grown at 27°C for 72 hours in 12.5% GBF or Medium A$^+$, and analyzed the

content after derivatization. We could detect and resolve all major saturated and unsaturated fatty

acid species (Table 4). Cultures grown in GBF had greater totals of assayed fatty acid species (27

$\pm$ 7 mg FAME gDCW^{-1}) when compared to cells grown in Medium A$^+$ (15 $\pm$ 1 mg FAME gDCW$^-$

^{1}) (Table 4). The most abundant fatty acid in all samples was 16:0 (42-45% of the total fatty acids).

C18:2 Δ9,12 fatty acids comprised a significant fraction of Medium A$^+$ grown cultures with 22%

of the total fatty acid species. However, in GBF-grown cells C18:2 Δ9,12 fatty acids were only

15% of the total fatty acid pool, while C18:3 Δ9,12,15 fatty acids were twice as high in GBF grown

cells (16%) than in Medium A$^+$ grown cells (9%) (Table 4). These results suggest that cells were

altering their membrane homeostasis when grown in GBF, as compared to standard growth in

Medium A$^+$. While we did not differentiate the fatty acid content and composition between the

outer, plasma, or thylakoid membranes in our study, prior work done with the cyanobacterium

Synechocystis PCC 6803 found similar fatty acid composition between thylakoid and plasma

membranes (50).

DISCUSSION

The amount and distribution of arable land and potable water are projected to change over

the next several decades due to climate change, while the rise in global population and standard of

living are expected to increase demand for these resources (51). Microalgal cultivation integrated

with industrial and municipal wastewater treatment circumvents many of the resource concerns

raised over biofuel production (2), while simultaneously removing additional nutrients and

pollutants present in the wastewater (52). *Synechococcus* PCC 7002 is an intriguing possible

platform for this purpose, since it can be readily engineered to produce high-value chemicals (53).

However, under standard environmental conditions we were unable to obtain robust growth of

PCC 7002 using a diluted municipal side stream (GBF) as a nutrient source. We hypothesized that

this effect may be due to the presence of DOM, which has been demonstrated to cause a decrease

in photosynthetic performance in various cyanobacteria strains (54–56). We investigated the

effects of light intensity and temperature on the physiology of cultures grown in GBF, in an effort

to find conditions conducive to high growth rates and biomass generation, and to better understand

the mechanisms of GBF toxicity.

We propose that the herbicidal effect of GBF is primarily due to PSII inhibition as shown

in Figures 7 and 8, brought about by quinone or phenolic compounds within the "soluble microbial

products" found in our EEM scans (Figure 2). Humic material enriched in quinone and phenolic

compounds can inhibit cyanobacterial growth (57, 58). Phenolic photosynthetic electron transfer

inhibitors, such as 2,5-Dibromo-3-methyl-6-isopropyl-p-benzoquinone (DBMIB), alter the redox

potential of the PQ pool of PSII by blocking forward electron transfer to the cytochrome b_6/f

complex (59). The immediate decrease in oxygen evolution (Figure 7) and high rates of initial

membrane permeability (Figure 4) upon GBF exposure suggests that the toxic compound(s)

rapidly cross the outer membrane, where it subsequently interrupts photosynthetic electron flow.

Chronic exposure to phenolic herbicides eventually leads to radical-catalyzed back reactions that

trigger ROS formation (60) (Figure 5) that facilitates complex destruction (Figure 6) and cell death

(45, 61–64). Some phenolic herbicides may also act as arylating agents, causing covalent binding

to macromolecules via Michael addition and thiol pool depletion (65). The inability of the reducing

agents we tested to maintain membrane integrity suggests that GBF-induced cytomembrane

permeability is likely caused by redox cycling, and not arylation.

We found that cultivation temperature was an important factor determined whether PCC

7002 could grow under high GBF concentrations. Numerous physiological processes are altered

at low temperatures (66). Notably, the fatty acyl chains in membranes undergo a transition from a

fluid to nonfluid state (67). Cyanobacterial membrane organization is complicated by the

simultaneous existence of the outer, plasma, and thylakoid membranes, each with a designated

physiological function (68). As part of the homeoviscous response upon a shift to a lower

temperature, cyanobacteria alter the expression of desaturases in their plasma and thylakoid

membranes (69) to increase unsaturated fatty acid content and maintain optimal membrane

function (70). Optimal thylakoid membrane fluidity is a critical factor in photosynthetic electron

transport, because co-utilized redox components must be able to move quickly between

photosynthetic and respiratory complexes to ensure ideal electron flow (71, 72). Temperature is

known to influence the kinetics governing the redox state of plastiquinone (PQ) through alterations

in thylakoid membrane composition and fluidity (73). We hypothesize that alteration of the

thylakoid membrane to circumvent GBF induced changes in the overall redox state is also an

important component of the adaptive response to lower temperatures, given the increase in total

unsaturated fatty acids observed during cultivation in GBF-based media (Table 4). Upon shifts to

a lower temperature (74), PCC 7002 has also been shown to translocate the phycobilisomes from

PSII to PSI, thereby decreasing the ratio of reducing power to proton-motive force (75). Mutant

cells of the cyanobacterium *Synechocystis* PCC 6803 that lacked polyunsaturated fatty acids were

unable to perform these state transitions at low temperatures (76). This change in photosystem

arrangement at 27°C may also contribute to acclimation to the photoinhibitory compound. Future

efforts to increase tolerance to these toxicants might include disruption of the inhibitor binding

sites (77, 78) or optimizing thylakoid membrane fluidity (79).

CONCLUSIONS

Under standard cultivation conditions of 37°C with the cyanobacteria *Synechococcus* sp.

strain PCC 7002, there was a dose-dependent relationship between liquid anaerobic digestate

(GBF) concentration and membrane permeability. This dose-dependent relationship mirrored ROS

production and photopigment degradation. Digestate contained dissolved organic matter

constituents that were likely affecting photosynthetic electron transport. Decreasing the cultivation

temperature to 27°C enabled robust cultivation at high digestate concentrations, resulting in high

biomass productivities. This temperature dependent tolerance may be due to changes in membrane

properties. Our study highlights an unanticipated influence of dissolved organic matter on

photosynthetic growth and physiology in wastewater based media, as well as a potential

mechanism for tolerance in low temperatures.

ACKNOWLEDGEMENTS

This work was funded by the US National Science Foundation (EFRI-1240268). TCK is

the recipient of a National Institutes of Health (NIH) Biotechnology Training Fellowship (NIGMS-

5 T32 GM08349) and a fellowship from the UW-Madison College of Engineering's Graduate

Engineering Research Scholars (GERS) program. The authors are grateful to Richard Mikel,

Matthew Dysthe, and Derek Jacobs for help with routine sampling, and Andrew Maizel and

Christina Remucal for advice on the excitation−emission matrix (EEM) fluorescence

spectroscopy.

REFERENCES

1. Oliver JWK, Atsumi S. 2014. Metabolic design for cyanobacterial chemical synthesis. Photosynth Res 120:249–61.

2. Pate R, Klise G, Wu B. 2011. Resource demand implications for US algae biofuels production scale-up. Appl Energy 88:3377–3388.

3. Clarens AF, Resurreccion EP, White M a, Colosi LM. 2010. Environmental life cycle comparison of algae to other bioenergy feedstocks. Environ Sci Technol 44:1813–9.

4. Posadas E, García-Encina PA, Domínguez A, Díaz I, Becares E, Blanco S, Muñoz R. 2014. Enclosed tubular and open algal-bacterial biofilm photobioreactors for carbon and nutrient removal from domestic wastewater. Ecol Eng 67:156–164.

5. Sturm BSM, Lamer SL. 2011. An energy evaluation of coupling nutrient removal from wastewater with algal biomass production. Appl Energy 88:3499–3506.

6. Hoffmann JP. 1998. Wastewater Treatment with Suspended and Nonsuspended Algae. J Phycol 34:757–763.

7. de la Noüe J, Laliberté G, Proulx D. 1992. Algae and waste water. J Appl Phycol 4:247–254.

8. Olguín EJ. 2012. Dual purpose microalgae–bacteria-based systems that treat wastewater and produce biodiesel and chemical products within a Biorefinery. Biotechnol Adv 30:1031–1046.

9. Wang L, Min M, Li Y, Chen P, Chen Y, Liu Y, Wang Y, Ruan R. 2010. Cultivation of green algae Chlorella sp. in different wastewaters from municipal wastewater treatment plant. Appl Biochem Biotechnol 162:1174–86.

10. Peccia J, Haznedaroglu B, Gutierrez J, Zimmerman JB. 2013. Nitrogen supply is an

important driver of sustainable microalgae biofuel production. Trends Biotechnol 31:134–8.

11.   Rawat I, Ranjith Kumar R, Mutanda T, Bux F. 2011. Dual role of microalgae: Phycoremediation of domestic wastewater and biomass production for sustainable biofuels production. Appl Energy 88:3411–3424.

12.   Pflugmacher S, Spangenberg M, Steinberg CEW. 1999. Dissolved organic matter (DOM) and effects on the aquatic macrophyte Ceratophyllum demersum in relation to photosynthesis, pigmentpattern and activity of detoxication enzymes. J Appl Bot 73:184–190.

13.   Zsolnay Á. 2003. Dissolved organic matter: Artefacts, definitions, and functions, p. 187–209. *In* Geoderma.

14.   Imai A, Fukushima T, Matsushige K, Kim Y-H, Choi K. 2002. Characterization of dissolved organic matter in effluents from wastewater treatment plants. Water Res 36:859–870.

15.   Akhiar A, Torrijos M, Battimelli A, Carrère H. 2016. Comprehensive characterization of the liquid fraction of digestates from full-scale anaerobic co-digestion. Waste Manag.

16.   Vigneault B, Percot A, Lafleur M, Campbell PGC. 2000. Permeability changes in model and phytoplankton membranes in the presence of aquatic humic substances. Environ Sci Technol 34:3907–3913.

17.   Ojwang' LM, Cook RL. 2013. Environmental Conditions That Influence the Ability of Humic Acids to Induce Permeability in Model Biomembranes. Environ Sci Technol 47:8280–8287.

18.   Chiou CT, Malcolm RL, Brinton TI, Kile DE. 1986. Water solubility enhancement of some organic pollutants and pesticides by dissolved humic and fulvic acids. Environ Sci

Technol;(United States) 20.

19.   Zhang D, Yan S, Song W. 2014. Photochemically Induced Formation of Reactive Oxygen Species (ROS) from Effluents Organic Matter. Environ Sci Technol.

20.   Lee E, Glover CM, Rosario-Ortiz FL. 2013. Photochemical formation of hydroxyl radical from effluent organic matter: Role of composition. Environ Sci Technol 47:12073–12080.

21.   Laue P, Bährs H, Chakrabarti S, Steinberg CEW. 2014. Natural xenobiotics to prevent cyanobacterial and algal growth in freshwater: Contrasting efficacy of tannic acid, gallic acid, and gramine. Chemosphere 104:212–220.

22.   Keren N, Krieger-Liszkay A. 2011. Photoinhibition: molecular mechanisms and physiological significance. Physiol Plant 142:1–5.

23.   Nagano T, Kojima K, Hisabori T, Hayashi H, Morita EH, Kanamori T, Miyagi T, Ueda T, Nishiyama Y. 2012. Elongation factor G is a critical target during oxidative damage to the translation system of Escherichia coli. J Biol Chem 287:28697–704.

24.   Foyer CH, Neukermans J, Queval G, Noctor G, Harbinson J. 2012. Photosynthetic control of electron transport and the regulation of gene expression. J Exp Bot 63:1637–61.

25.   Voss I, Sunil B, Scheibe R, Raghavendra AS. 2013. Emerging concept for the role of photorespiration as an important part of abiotic stress response. Plant Biol (Stuttg) 15:713–22.

26.   Ruffing AM. 2011. Engineered cyanobacteria: Teaching an old bug new tricks. bioe 2:136–149.

27.   Xu Y, Alvey RM, Byrne PO, Graham JE, Shen G, Bryant DA. 2011. Expression of genes in cyanobacteria: adaptation of endogenous plasmids as platforms for high-level gene expression in Synechococcus sp. PCC 7002. Methods Mol Biol 684:273–93.

28. Stevens SE, Patterson COP, Myers J. 1973. The Production of Hydrogen Peroxide by Blue-Green Algae: A Survey. J Phycol 9:427–430.

29. Sakamoto T, Bryant DA. 2001. Requirement of Nickel as an Essential Micronutrient for the Utilization of Urea in the Marine Cyanobacterium Synechococcus sp. PCC 7002. Microbes Environ 16:177–184.

30. Flowers JJ, Cadkin TA, McMahon KD. 2013. Seasonal bacterial community dynamics in a full-scale enhanced biological phosphorus removal plant. Water Res 47:7019–7031.

31. He S, McMahon KD. 2011. "Candidatus Accumulibacter" gene expression in response to dynamic EBPR conditions. ISME J 5:329–40.

32. Eaton AD, Franson MAH. 2005. Standard Methods for the Examination of Water & Wastewater, 21sted. American Public Health Association.

33. Lennen RM, Braden DJ, West RA, Dumesic JA, Pfleger BF. 2010. A process for microbial hydrocarbon synthesis: Overproduction of fatty acids in Escherichia coli and catalytic conversion to alkanes. Biotechnol Bioeng 106:193–202.

34. Sakamoto T, Bryant D a. 2002. Synergistic effect of high-light and low temperature on cell growth of the Delta12 fatty acid desaturase mutant in Synechococcus sp. PCC 7002. Photosynth Res 72:231–42.

35. Miyashita H, Adachi K, Kurano N, Ikemot H, Chihara M, Miyach S. 1997. Pigment composition of a novel oxygenic photosynthetic prokaryote containing chlorophyll d as the major chlorophyll. Plant cell Physiol 38:274–281.

36. Porra RJ, Thompson WA, Kriedemann PE. 1989. Determination of accurate extinction coefficients and simultaneous equations for assaying chlorophylls a and b extracted with four different solvents: verification of the concentration of chlorophyll standards by atomic

absorption spectroscopy. Biochim Biophys Acta (BBA)-Bioenergetics 975:384–394.

37. Chen W, Westerhoff P, Leenheer JA, Booksh K. 2003. Fluorescence Excitation-Emission Matrix Regional Integration to Quantify Spectra for Dissolved Organic Matter. Environ Sci Technol 37:5701–5710.

38. Neilen A, W. Hawker D, R. O'Brien K, Burford M. 2017. Phytotoxic effects of terrestrial dissolved organic matter on a freshwater cyanobacteria and green algae species is affected by plant source and DOM chemical compositionChemosphere.

39. Lehmann J, Kleber M. 2015. The contentious nature of soil organic matter. Nature 1–9.

40. Peña-méndez ME, Havel J, Patočka J. 2005. Humic substances – compounds of still unknown structure : applications in agriculture , industry , environment , and biomedicine. J Appl Biomed 3:13–24.

41. Cowgill RW. 1963. Fluorescence and the structure of proteins. I. Effects of substituents on the fluorescence of indole and phenol compounds. Arch Biochem Biophys 100:36–44.

42. Maizel AC, Remucal CK. 2017. The effect of advanced secondary municipal wastewater treatment on the molecular composition of dissolved organic matter. Water Res 122:42–52.

43. Roth BL, Poot M, Yue ST, Millard PJ. 1997. Bacterial viability and antibiotic susceptibility testing with SYTOX green nucleic acid stain. Appl Environ Microbiol 63:2421–31.

44. Samuni Y, Goldstein S, Dean OM, Berk M. 2013. The chemistry and biological activities of N-acetylcysteine. Biochem Biophys 1830:4117–4129.

45. Latifi A, Ruiz M, Zhang C-C. 2009. Oxidative stress in cyanobacteria. FEMS Microbiol Rev 33:258–78.

46. van Gorkom HJ, Gast P. 1996. Measurement of Photosynthetic Oxygen Evolution, p. 391–405. *In* Amesz, J, Hoff, AJ (eds.), Biophysical Techniques in Photosynthesis. Springer

Netherlands, Dordrecht.

47.   Sakamoto T, Bryant D a. 1998. Growth at low temperature causes nitrogen limitation in the cyanobacterium Synechococcus sp. PCC 7002. Arch Microbiol 169:10–9.

48.   Huber SC, Edwards GE. 1975. Effect of DBMIB, DCMU and antimycin A on cyclic and noncyclic electron flow in C4 mesophyll chloroplasts. FEBS Lett 58:211–214.

49.   Tyystjärvi E, Aro EM. 1996. The rate constant of photoinhibition, measured in lincomycin-treated leaves, is directly proportional to light intensity. Proc Natl Acad Sci U S A 93:2213–2218.

50.   Los DA, Murata N. 1996. Characterization of the Fad12 mutant of Synechocystis that is defective in Δ12 acyl-lipid desaturase activity. Biochim Biophys Acta - Lipids Lipid Metab 1299:117–123.

51.   Parry ML, Rosenzweig C, Iglesias A, Livermore M, Fischer G. 2004. Effects of climate change on global food production under SRES emissions and socio-economic scenarios. Glob Environ Chang 14:53–67.

52.   Wang Y, Ho SH, Cheng CL, Guo WQ, Nagarajan D, Ren NQ, Lee DJ, Chang JS. 2016. Perspectives on the feasibility of using microalgae for industrial wastewater treatment. Bioresour Technol 222:485–497.

53.   Davies FK, Work VH, Beliaev AS, Posewitz MC. 2014. Engineering Limonene and Bisabolene Production in Wild Type and a Glycogen-Deficient Mutant of Synechococcus sp. PCC 7002. Front Bioeng Biotechnol 2:1–11.

54.   Sun B, Tanji Y, Unno H. 2006. Extinction of cells of cyanobacterium Anabaena circinalis in the presence of humic acid under illumination. Appl Microbiol Biotechnol 72:823–828.

55.   Shao J, Wu Z, Yu G, Peng X, Li R. 2009. Allelopathic mechanism of pyrogallol to

Microcystis aeruginosa PCC7806 (Cyanobacteria): From views of gene expression and antioxidant system. Chemosphere 75:924–928.

56. Gjessing ET, Alberts J., Bruchet A, Egeberg PK, Lydersen E, McGown LB, Mobed JJ, Münster U, Pempkowiak J, Perdue M, Ratnawerra H, Rybacki D, Takacs M, Abbt-Braun G. 1998. Multi-method characterisation of natural organic matter isolated from water: characterisation of reverse osmosis-isolates from water of two semi-identical dystrophic lakes basins in Norway. Water Res 32:3108–3124.

57. Bährs H, Putschew A, Steinberg CEW. 2013. Toxicity of hydroquinone to different freshwater phototrophs is influenced by time of exposure and pH. Environ Sci Pollut Res 20:146–154.

58. Bährs H, Menzel R, Kubsch G, Stößer R, Putschew A, Heinze T, Steinberg CEW. 2012. Does quinone or phenol enrichment of humic substances alter the primary compound from a non-algicidal to an algicidal preparation? Chemosphere 87:1193–1200.

59. Schoepp B, Brugna M, Riedel A, Nitschke W, Kramer DM. 1999. The Qo-site inhibitor DBMIB favours the proximal position of the chloroplast Rieske protein and induces a pK-shift of the redox-linked proton. FEBS Lett 450:245–250.

60. Pallett KE, Dodge AD. 1980. Studies into the Action of Some Photosynthetic Inhibitor Herbicides. J Exp Bot 31:1051–1066.

61. Krieger-Liszkay A, Fufezan C, Trebst A. 2008. Singlet oxygen production in photosystem II and related protection mechanism. Photosynth Res 98:551–564.

62. Idedan I, Tomo T, Noguchi T. 2011. Herbicide effect on the photodamage process of photosystem II: Fourier transform infrared study. Biochim Biophys Acta - Bioenerg 1807:1214–1220.

63. Nishiyama Y, Allakhverdiev SI, Murata N. 2011. Protein synthesis is the primary target of reactive oxygen species in the photoinhibition of photosystem II. Physiol Plant 142:35–46.

64. Dominy PJ, Williams WP. 1987. Chlorophyll Photobleaching is Dependent on Photosystem II Inhibition, p. 35–38. *In* Biggins, J (ed.), Progress in Photosynthesis Research: Volume 4 Proceedings of the VIIth International Congress on Photosynthesis Providence, Rhode Island, USA, August 10--15, 1986. Springer Netherlands, Dordrecht.

65. Wang X, Thomas B, Sachdeva R, Arterburn L, Frye L, Hatcher PG, Cornwell DG, Ma J. 2006. Mechanism of arylating quinone toxicity involving Michael adduct formation and induction of endoplasmic reticulum stress. Proc Natl Acad Sci U S A 103:3604–9.

66. Morgan-Kiss RM, Priscu JC, Pocock T, Gudynaite-Savitch L, Huner NPA. 2006. Adaptation and Acclimation of Photosynthetic Microorganisms to Permanently Cold Environments. Microbiol Mol Biol Rev 70:222–252.

67. De Mendoza D. 2014. Temperature Sensing by Membranes. Annu Rev Microbiol 68:101–16.

68. Pisareva T, Kwon J, Oh J, Kim S, Ge C, Wieslander A, Choi J-S, Norling B. 2011. Model for membrane organization and protein sorting in the cyanobacterium Synechocystis sp. PCC 6803 inferred from proteomics and multivariate sequence analyses. J Proteome Res 10:3617–31.

69. Mustardy L, Los DA, Gombos Z, Murata N. 1996. Immunocytochemical localization of acyl-lipid desaturases in cyanobacterial cells: evidence that both thylakoid membranes and cytoplasmic membranes are sites of lipid desaturation. Proc Natl Acad Sci U S A 93:10524–7.

70. Singh SC, Sinha RP, Häder D. 2002. Role of Lipids and Fatty Acids in Stress Tolerance in

Cyanobacteria. Acta Protozool 41:297–308.

71.   Mullineaux CW. 2014. Electron transport and light-harvesting switches in cyanobacteria. Front Plant Sci 5:1–6.

72.   Liu L-N. 2015. Distribution and dynamics of electron transport complexes in cyanobacterial thylakoid membranes. Biochim Biophys Acta - Bioenerg.

73.   Klementiev KE, Tsoraev G V, Tyutyaev E V, Zorina AA, Feduraev P V, Allakhverdiev SI, Paschenko VZ. 2017. Membrane fluidity controls redox-regulated cold stress responses in cyanobacteria. Photosynth Res 0:0.

74.   Mackey KRM, Paytan A, Caldeira K, Grossman AR, Moran D, McIlvin M, Saito MA. 2013. Effect of temperature on photosynthesis and growth in marine Synechococcus spp. Plant Physiol 163:815–29.

75.   Mao H Bin, Li GF, Ruan X, Wu QY, Gong YD, Zhang XF, Zhao NM. 2002. The redox state of plastoquinone pool regulates state transitions via cytochrome b6f complex in Synechocystis sp. PCC 6803. FEBS Lett 519:82–86.

76.   El Bissati K, Delphin E, Murata N, Etienne a, Kirilovsky D. 2000. Photosystem II fluorescence quenching in the cyanobacterium Synechocystis PCC 6803: involvement of two different mechanisms. Biochim Biophys Acta 1457:229–242.

77.   Ajlani G, Kirilovsky D, Picaud M, Astier C. 1989. Molecular analysis of psbA mutations responsible for various herbicide resistance phenotypes in Synechocystis 6714. Plant Mol Biol 13:469–79.

78.   Lee TX, Metzger SU, Cho YS, Whitmarsh J, Kallas T. 2001. Modification of inhibitor binding sites in the cytochrome bf complex by directed mutagenesis of cytochrome b6 in Synechococcus sp. PCC 7002. Biochim Biophys Acta - Bioenerg 1504:235–247.

79. Gombos Z, Wada H, Murata N. 1994. The recovery of photosynthesis from low-temperature photoinhibition is accelerated by the unsaturation of membrane lipids: a mechanism of chilling tolerance. Proc Natl Acad Sci U S A 91:8787–91.

LEGENDS

Figure 1. Flow diagram and nutrient streams obtained from the Nine Springs Wastewater Treatment Plant (Dane County, Wisconsin, USA). Stars indicate sampling points.

Figure 2. Excitation Emission Matrix of (A) Medium A+, (B) Secondary Effluent, (C) 6.25% GBF, and (D) 12.5% GBF. Fluorescence values were normalized to 1 ppm of quinine sulfate in 0.1 N H_2SO_4 at Ex/Em = 350/450 nm.

Figure 3. Biomass accumulation for cultures grown in Medium A$^+$, 6.25 %, 9.9 %, or 12.5 % (v/v) GBF media with 1 % CO_2 at (A) 37°C and 200 µmol photons m^{-2} s^{-1}, (B) 37°C and 100 µmol photons m^{-2} s^{-1}, (C) 27°C and 200 µmol photons m^{-2} s^{-1}, or (D) 27°C and 100 µmol photons m^{-2} s^{-1}. The values represent the mean ± SD of biological triplicates.

Figure 4. Membrane permeability of cultures grown in Medium A$^+$, 6.25 %, 9.9 %, or 12.5 % (v/v) GBF media with 1 % CO_2 at (A) 37°C and 200 µmol photons m^{-2} s^{-1}, (B) 37°C and 100 µmol photons m^{-2} s^{-1}, (C) 27°C and 200 µmol photons m^{-2} s^{-1}, or (D) 27°C and 100 µmol photons m^{-2} s^{-1}. The values represent the mean ± SD of biological triplicates.

Figure 5. Reactive oxygen species and membrane permeability assay. Cells were grown to early linear phase in Medium A$^+$ or 12.5% GBF with the appropriate treatment under continuous illumination (200 µmol photons m^{-2} s^{-1}) with 1 % CO_2 at 37°C. Fluorescence values were normalized to OD$_{730}$. The values represent the mean ± SD of biological triplicates.

Figure 6. Absorption spectra of PCC 7002 cultures exposed to GBF, at two different cultivation temperatures. Cultures were grown in Medium A$^+$, 6.25 %, or 12.5 % (v/v) GBF media under continuous illumination (200 µmol photons m^{-2} s^{-1}) with 1 % CO_2 at (A) 37°C or (B) 27°C for 72

hours. The spectra were recorded in dilute cell suspensions and normalized to an OD_{730}. The peak at 438 nm is due to carotenoids, the peak at 637 nm is due to phycobilisomes, and the peak at 683 nm is due to chlorophyll a.

Figure 7. Rates of oxygen evolution as a function of acclimation temperature and light intensity. Cells were grown to early linear phase in Medium A$^+$ or 12.5% GBF under continuous illumination (200 µmol photons m^{-2} s^{-1}) with 1 % CO_2 at (A) 37°C or (B) 27°C. Cells were pelleted, resuspended in Medium A$^+$ or 12.5% GBF, and the rate of maximal oxygen evolution was measured with 10 mM HCO_3^- as an electron acceptor at increasing light intensities. The values represent the mean ± SE of biological duplicates.

Figure 8. Photosynthetic oxygen evolution rates at saturating light conditions. Cells were grown to early linear phase in Medium A$^+$ under continuous illumination (200 µmol photons m^{-2} s^{-1}) with 1 % CO_2 at 37°C). Cells were pelleted and resuspended in Medium A$^+$, 12.5% GBF, 12.5% GBF + 310 mM NaCl (control for osmolarity), 12.5% GBF prefiltered with activated carbon, Medium A$^+$ + 10 µM DCMU (control for inhibition for electron transport), or pre-treated with 800 µg ml^{-1} Lincomycin for 1 hour (control for the inhibition of protein synthesis.) Maximal oxygen evolution rates were measured with 10 mM HCO_3 as an electron acceptor at 2700 µmol photons m^{-2} s^{-1}. The values represent the mean ± SE of biological duplicates.

TABLES

Table 1. Nutrient composition of batch of 100% GBF used for subsequent experiments

Analyte[a]	Concentration[b] (mg L^{-1}) (±SD)
NH$_3$–N	1180 ± 135
NO$_3$–N	7.5 ± 0.04
SRP	78 ± 10
COD	735 ± 4
TSS	467 ± 123
VSS	18 ± 2
TS	2350 ± 36
TS (Glass Fiber Filtered)	2030 ± 42
TS (0.45 um membrane filtered)	2000 ± 151

[a] SRP-soluble reactive phosphorus, COD-chemical oxygen demand, TSS- total suspended solids, VSS – volatile suspended solids, TS-total solids

[b] Values shown represent the mean ± standard deviation (SD) of at least three technical replicates.

Table 2. Characteristics of GBF and secondary effluent over the 6 month experimental period.

Source	Analyte	Concentration[a] (mg L^{-1}) (±COV)
GBF	NH$_3$–N	920 ± 22%
	NO$_3$–N	3.9 ± 43%
	SRP	54 ± 27%
Secondary Effluent	NH$_3$–N	n.d.
	NO$_3$–N	19.3 ± 6.6%
	SRP	n.d.

[a] Values shown represent the mean ± coefficient of variation (COV) of at least three technical replicates.

Table 3. Growth Rates with varying Temperature and Light Intensity

Media	Light Intensity (μmol photons m^{-2} s^{-1})	Temperature (°C)	Growth Rate (±SD) (OD day^{-1})
A^+	200	37	0.66 ± 0.04
6.25% GBF	200	37	0.58 ± 0.05
12.5% GBF	200	37	0.05 ± 0.01
A^+	100	37	0.28 ± 0.01
6.25% GBF	100	37	0.24 ± 0.01
9.9% GBF	100	37	0.11 ± 0.02
12.5% GBF	100	37	0.02 ± 0.00
A^+	200	27	0.73 ± 0.08
6.25% GBF	200	27	0.32 ± 0.03
9.9% GBF	200	27	0.42 ± 0.06
12.5% GBF	200	27	0.28 ± 0.04
A^+	100	27	0.25 ± 0.01
6.25% GBF	100	27	0.26 ± 0.02
9.9% GBF	100	27	0.24 ± 0.01
12.5% GBF	100	27	0.26 ± 0.02

[a] Linear growth rates of cultures grown in Medium A^+ or GBF with the appropriate treatment under continuous illumination (200 μmol photons m^{-2} s^{-1} or 100 μmol photons m^{-2} s^{-1}) with 1 % CO_2 at 37°C or 27°C. The values represent the mean ± SD of biological triplicates.

Table 4. Fatty acid content and composition of cultures grown in Medium A$^+$ or 12.5% GBF media under continuous illumination (200 µmol photons m^{-2} s^{-1}) at 27°C for 72 hours

Growth Conditions[1]		Fatty Acids Species					
		C16:0	C16:1 Δ9	C18:0	C18:1 Δ9	C18:2 Δ9,12	C18:3 Δ9,12,15
Medium A$^+$	mg FA gDCW^{-1}	6.7 ± 0.7	2.0 ± 0.1	0.2 ± 0.1	1.5 ± 0.2	3.3 ± 0.2	1.4 ± 0.1
	% FA	44 ± 1	13 ± 1	2 ± 0	10 ± 0	22 ± 1	9 ± 1
GBF	mg FA gDCW^{-1}	11.6 ± 3.1	3.5 ± 0.8	1.3 ± 0.3	2.2 ± 0.7	3.9 ± 0.8	4.4 ± 1.6
	% FA	43 ± 0	13 ± 0	5 ± 0*	8 ± 0	15 ± 1*	16 ± 1*

[1] Strains were grown in the Medium A$^+$ or 12.5% GBF media under continuous illumination (200 µmol photons m^{-2} s^{-1}) at 27°C for 72 hours. Fatty acids were extracted and derivatized as previously described. The values represent the mean ± SD of two independent experiments. * represent statistically significant differences from control (p-value<0.005 using an unpaired t-test)

Figure 1. Flow diagram and nutrient streams obtained from the Nine Springs Wastewater Treatment Plant (Dane County, Wisconsin, USA). Stars indicate sampling points.

Figure 2. Excitation Emission Matrix of (A) Medium A$^+$, (B) Secondary Effluent, (C) 6.25% GBF, and (D) 12.5% GBF. Fluorescence values were normalized to 1 ppm of quinine sulfate in 0.1 N H$_2$SO$_4$ at Ex/Em = 350/450 nm.

Figure 3. Biomass accumulation for cultures grown in Medium A[+], 6.25 %, 9.9 %, or 12.5 % (v/v) GBF media with 1 % CO_2 at (A) 37°C and 200 μmol photons m^{-2} s^{-1}, (B) 37°C and 100 μmol photons m^{-2} s^{-1}, (C) 27°C and 200 μmol photons m^{-2} s^{-1}, or (D) 27°C and 100 μmol photons m^{-2} s^{-1}. The values represent the mean ± SD of biological triplicates.

Figure 4. Membrane permeability of cultures grown in Medium A[+], 6.25 %, 9.9 %, or 12.5 % (v/v) GBF media with 1 % CO_2 at (A) 37°C and 200 μmol photons m^{-2} s^{-1}, (B) 37°C and 100 μmol photons m^{-2} s^{-1}, (C) 27°C and 200 μmol photons m^{-2} s^{-1}, or (D) 27°C and 100 μmol photons m^{-2} s^{-1}. The values represent the mean ± SD of biological triplicates.

Figure 5. Reactive oxygen species and membrane permeability assay. Cells were grown to early linear phase in Medium A^+ or 12.5% GBF with the appropriate treatment under continuous illumination (200 µmol photons m^{-2} s^{-1}) with 1 % CO_2 at 37°C. Fluorescence values were normalized to OD_{730}. The values represent the mean ± SD of biological triplicates.

Figure 6. Absorption spectra of PCC 7002 cultures exposed to GBF, at two different cultivation temperatures. Cultures were grown in Medium A$^+$, 6.25 %, or 12.5 % (v/v) GBF media under continuous illumination (200 μmol photons m^{-2} s^{-1}) with 1 % CO_2 at (A) 37°C or (B) 27°C for 72 hours. The spectra were recorded in dilute cell suspensions and normalized to an OD_{730}. The peak at 438 nm is due to carotenoids, the peak at 637 nm is due to phycobilisomes, and the peak at 683 nm is due to chlorophyll a.

Figure 7. Rates of oxygen evolution as a function of acclimation temperature and light intensity. Cells were grown to early linear phase in Medium A$^+$ or 12.5% GBF under continuous illumination (200 μmol photons m^{-2} s^{-1}) with 1 % CO_2 at (A) 37°C or (B) 27°C. Cells were pelleted, resuspended in Medium A$^+$ or 12.5% GBF, and the rate of maximal oxygen evolution was measured with 10 mM HCO_3^- as an electron acceptor at increasing light intensities. The values represent the mean ± SE of biological duplicates.

O₂ Evolution Rates at Saturating Light for 37°C Acclimated Cultures

Figure 8. Photosynthetic oxygen evolution rates at saturating light conditions. Cells were grown to early linear phase in Medium A$^+$ under continuous illumination (200 μmol photons m^{-2} s^{-1}) with 1 % CO_2 at 37°C). Cells were pelleted and resuspended in Medium A$^+$, 12.5% GBF, 12.5% GBF + 310 mM NaCl (control for osmolarity), 12.5% GBF prefiltered with activated carbon, Medium A$^+$ + 10 μM DCMU (control for inhibition for electron transport), or pre-treated with 800 μg ml^{-1} Lincomycin for 1 hour (control for the inhibition of protein synthesis.) Maximal oxygen evolution rates were measured with 10 mM HCO_3 as an electron acceptor at 2700 μmol photons m^{-2} s^{-1}. The values represent the mean ± SE of biological duplicates.