Seeing spots: Measuring, quantifying heritability, and assessing fitness consequences of

coat pattern traits in a wild population of giraffes (*Giraffa camelopardalis*)

DEREK E. LEE,[1,2,*] DOUGLAS R. CAVENER,[2] AND MONICA L. BOND[1,3]

[1] Wild Nature Institute, Concord, New Hampshire, USA

[2] Department of Biology, Pennsylvania State University, University Park, Pennsylvania, USA

[3] Department of Evolutionary Biology and Environmental Studies, University of Zürich,

Zürich, Switzerland

* Correspondent: derek@wildnatureinstitute.org

RH: Giraffe coat pattern traits

ABSTRACT

Polymorphic phenotypes of mammalian coat color have been important to the study of

genetics and evolution, but little is known about the heritability and fitness consequences of

variation in complex coat pattern traits in wild populations. Understanding the current

evolution of coat patterns requires reliably measuring traits, quantifying heritability of the

traits, and identifying the fitness consequences of specific phenotypes. Giraffe coat markings

are highly variable and it has been hypothesized that variation in coat patterns most likely

affects fitness by camouflaging neonates against predators. We quantified spot pattern traits

of wild Masai giraffes using image analysis software, determined whether spot pattern traits

were heritable, and assessed whether variation in heritable spot pattern traits was related to

fitness as measured by juvenile survival. The methods we described comprise a framework

for objective quantification of mammalian coat pattern traits based on photographic coat

pattern data. We demonstrated that characteristics of giraffe coat spot shape are heritable. We

did not find evidence for juvenile survival consequences of variation in spot traits, suggesting

that spot traits are currently not under strong directional, disruptive, or stabilizing selection

for neonate camouflage in our study population, but our sample size could not detect small

differences in survival. Spot trait variation also may be more relevant to other components of

fitness, such as adult survival or fecundity. We hope this case study will inspire further

investigations of coat pattern traits.

Key words: adaptation, coat pattern, heritability, natural selection, phenomics, phenotypic

selection, photogrammetry, quantitative genetics, remote measurement, visual animal

biometry

34   **INTRODUCTION**

Complex coat patterns are found on many mammal species and these phenotypic traits are

hypothesized to play adaptive roles in predator and parasite evasion, thermoregulation, and

communication (Cott 1940; Caro 2005). Pigmentation biology has played a prominent role in

the foundation of genetics and evolutionary biology with most work on vertebrates focused

on a few starkly different color morphs in mice (Hoekstra 2006). However, most studies to

date of the heritability and adaptation of complex animal skin pigmentation patterns have

concentrated on fishes with their multiple chromatophores (Kelsh 2004). Researchers have

proposed mathematical models for pattern-formation mechanisms in animal markings

(Murray 1981; Maini 2004; Garvie and Trenchea 2014), and hypothesized the genetic and

developmental mechanisms for the markings (Mills and Patterson 2009; Eizirik *et al.* 2010),

and although studies of heritability and adaptation of coat patterns in wild populations are

becoming more prevalent (Kruuk *et al.* 2008), we see a need for broadly applicable tools and

techniques to quantify coat pattern traits in wild populations (Willisch et al. 2013).

48   Understanding the current evolution of a trait in a wild population requires reliably

measuring individual variation in the trait, quantifying heritability as the proportion of

observed phenotypic variation of a trait that is passed from parent to offspring, and assessing

the fitness consequences of phenotypic variation (Lande and Arnold 1983; Falconer and

Mackay 1996). The fraction of variability in a phenotypic trait that is explained by genetic

factors is heritability, which can be estimated as the resemblance of the offspring to its

parents (Falconer and Mackay 1996; Roff 1997). Fitness of a trait can be assessed in many

ways, but because neonatal mortality in mammals is generally higher than any other age class

(Lee and Strauss 2016), one common proxy of phenotypic fitness is juvenile survival

(Paterson *et al.* 1998). Here we provide an example quantification and analysis of heritability

and fitness for mammalian coat pattern traits using giraffe (*Giraffa camelopardalis*) as the

organism of interest.

Giraffe skin pigmentation is uniformly dark grey (Dimond and Montagna 1976), but

the spots that make up their coat pattern markings are highly variable in traits such as color,

roundness, and perimeter tortuousness, and this variation has been used to classify subspecies

(Lydekker 1904), and to reliably identify individuals because patterns do not change with age

(Foster 1966; Bolger *et al.* 2012; Dagg 2014). Dagg (1968) first presented evidence from a

small zoo population that the shape, number, area, and color of spots in giraffe coat patterns

may be heritable, but analysis of spot traits in wild giraffes, and tools for objectively

measuring spot characteristics have been lacking. It has been hypothesized that giraffe coat

patterns evolved to camouflage neonates whose primary defense against predation is

concealment (Langman 1977; Mitchell and Skinner 2003); thus the most likely fitness effects

from variation in coat patterns should be variation in juvenile survival. Alternative

hypotheses about the adaptive value of giraffe coat markings include thermoregulation

(Skinner and Smithers 1990), and in this social species with good visual sensory perception

(Dagg 2014; VanderWaal *et al.* 2014), markings could also facilitate individual recognition

(Tibbetts and Dale 2007) and kin recognition (Beecher 1982; Tang-Martinez 2001).

The spot patterns of Masai giraffes (*G. c. tippelskirchii*) are particularly diverse

among giraffe populations (Dagg 1968; **Fig 1**). Masai giraffes spots vary in color and shape

from those that are nearly round with very smooth edges (low tortuousness), to extremely

incised or lobate (high tortuousness). Quantifying heritability and fitness consequences of

phenotypic variation in coat pattern traits of giraffes could potentially contribute to the

understanding of the evolution of mammalian coat patterns. Our purpose in this study was to

1) objectively and reliably quantify the spot pattern traits of wild Masai giraffes in northern

Tanzania using photogrammetry and image analysis software, 2) determine whether spot

pattern traits were heritable from mother to offspring, and 3) determine whether variation in

heritable spot pattern traits was related to fitness as measured by juvenile survival.

METHODS

Field Data Collection

This study used data from individually identified, wild, free-ranging Masai giraffes in a 1700

km^2 sampled area within a 4400 km^2 region of the Tarangire Ecosystem, northern Tanzania,

East Africa. We collected data during systematic road transect sampling for photographic

capture-mark-recapture (PCMR). We conducted 26 daytime surveys for giraffe PCMR data

between January 2012 and February 2016. We sampled giraffes three times per year around 1

February, 1 June, and 1 October near the end of every precipitation season (short rains, long

rains, and dry, respectively) by driving a network of fixed-route transects on single-lane dirt

tracks in the study area. We surveyed according to Pollock's robust design sampling

framework (Pollock 1982; Kendall *et al.* 1995), with three occasions per year. Each sampling

occasion was composed of two sampling events during which we surveyed all transects in the

study area with only a few days interval between events. Each sampling occasion was

separated by a 4-month interval (4.3 years $\times$ 3 occasions year^{-1} $\times$ 2 events occasion^{-1} = 26

survey events).

During PCMR sampling events, a sample of individuals were encountered and either

"sighted" or "resighted" by slowly approaching and photographing the animal's right side at

a perpendicular angle (Canon 40D and Rebel T2i cameras with Canon Ultrasonic IS 100 –

400 mm lens, Canon U.S.A., Inc., One Canon Park, Melville, New York, 11747, USA). We

identified individual giraffes using their unique and unchanging coat patterns (Foster 1966;

Dagg 2014) with the aid of pattern-recognition software Wild-ID (Bolger *et al.* 2012). We

attempted to photograph every giraffe encountered, and recorded sex and age class based on

physical characteristics. We categorized giraffes into four age classes: neonate calf (0 – 3

months old), older calf (4 – 11 months old), subadult (1 – 3 years old for females, 1 – 6 years

old for males), or adult (> 3 years for females, > 6 years for males) using a suite of physical

characteristics (Strauss *et al.* 2015), and size measured with photogrammetry (Lee *et al.*

2016a).

All animal work was conducted according to relevant national and international

guidelines. This research was carried out with permission from the Tanzania Commission for

Science and Technology (COSTECH), Tanzania National Parks (TANAPA), the Tanzania

Wildlife Research Institute (TAWIRI). No Institutional Animal Care and Use Committee

(IACUC) approval was necessary because animal subjects were observed without disturbance

or physical contact of any kind.

118   *Quantification of Spot Patterns*

We analysed spot traits of each animal within the shoulder and rib area by cropping all

images to an analysis rectangle that fit horizontally between the anterior edge of the rear leg

and the chest, and vertically between the back and where the skin folded beneath the posterior

edge of the foreleg (**Fig 1**). We quantified spot characteristics of each animal's pattern using

the Color Histogram and Analyze Particles procedures of ImageJ (Schneider *et al.* 2012). For

color analysis, we used the entire analysis rectangle and full-color photos. For spot

measurements we analysed 8-bit greyscale images that we converted to bicolor (black and

white) using the ImageJ Enhance Contrast and Threshold commands. To account for

differences in image resolution and animal size (including age-related growth), we set the

measurement unit of each image equal to the number of pixels in the height of the analysis

rectangle. Therefore all measurements are in giraffe units (GU), where 1 GU = height of the

analysis rectangle (**Fig 1**). We excluded particles cut off by the edge of the analysis rectangle

to avoid the influence of incomplete spots, and we also excluded spots whose area was

<0.00001 GU^2 to eliminate the influence of speckles.

We characterized each animal's spot pattern traits within the analysis rectangle using

the following 11 metrics available in ImageJ: number of spots; mean spot size (area); mean

spot perimeter; mean angle between the primary axis of an ellipse fit over the spot and the x-

axis of the image; mean circularity ($4\pi \times [Area] / [Perimeter]^2$ with a value of 1.0 indicating

a perfect circle and smaller values indicating an increasingly elongated shape); mean

maximum caliper (the longest distance between any two points along the spot boundary, also

known as Feret diameter); mean Feret angle (the angle [0 to 180 degrees] of the maximum

caliper); mean aspect ratio (of the spot's fitted ellipse); mean roundness ($4 \times [Area]\pi \times$

$[Major\ axis]^2$ or the inverse of *aspect ratio*); mean solidity ($[Area] / [Convex\ area]$, also

called tortuousness); and mode shade ($[65536 \times r] + [256 \times g] + [b]$ using RGB values from

color histogram from full color photos).

We quantified among-individual variation in trait values by reporting the mean, SD,

and coefficient of variation (CV) of each trait. We quantified the repeatability of our spot

pattern trait measurement technique by computing the amount of among-measurement

variation made on different photos from different dates for the same animal using a set of 30

animals with >2 images per animal.

149   *Heritability of Spot Traits*

Parent-offspring (PO) regression is one of the traditional quantitative genetics tools used to

measure heritability (h^2; Falconer and Mackay 1996). PO regression compares phenotypic

trait values in parents to those same trait values in their offspring, with the slope of the linear

regression line between the mean parent phenotype and the mean offspring phenotype

providing an estimate of the heritability of the trait. Our analysis of maternal-offspring

regression and the estimated slope represents one-half of total heritability. Advantages and

disadvantages of the parent-offspring method to estimate heritability compared to other

methods such as the half-sibling method or the animal model have been elucidated (Åkesson

et al. 2008; de Villemereuil 2013), with the primary advantage when studying wild

populations being that PO regression requires less information about family structure and

only one offspring per individual (Roff 1997; Lynch and Walsh 1998). While PO regression

can have low statistical power and inherent biases when estimating heritability due to

environmental effects shared by related individuals or unbalanced sampling (Wilson *et al.*

2010), this methodology was the most appropriate for our study design because we were

unable to identify fathers, and our sample did not include any maternal siblings or half-

siblings.

We identified mother-calf pairs by observing extended suckling behavior. Wild

female giraffes very rarely suckle a calf that is not their own (Pratt and Anderson 1979). We

examined all identification photographs for individuals in known mother-calf pairs, and

selected the best-quality photograph for each animal based on focus, clarity, perpendicularity

to the camera, and unobstructed view of the torso. We found 31 known mother-calf pairs with

high-quality photographs of both animals.

We predicted spot pattern traits of a calf would be correlated with those of its mother

but not with a random cow. For comparison of spot characteristics between known mother-

calf pairs, we created a null expectation set of random cow-calf pairs using the same

photographs by assigning a random mother to each calf (without replacement and without

pairing a mother with her own calf). We tested our prediction for each spot characteristic

using simple linear regressions of calf values versus mother values (heritability models), and

calf values versus random cow values (null expectation models). Because we examined 11

spot traits, we used the Bonferroni adjustment (α/number of tests) to account for multiple

tests and set our adjusted $\alpha = 0.0045$. We performed statistical operations using the lm

function in R (R Core Development Team 2013).

182   *Fitness of Spot Patterns from Juvenile Survival*

We assembled encounter histories for all calves first observed as neonates for survival

analysis. For each calf we selected the best-quality calf-age (age <6 mo) photograph based on

focus, clarity, perpendicularity to the camera, and unobstructed view of the torso. We

analysed survival using capture-mark-recapture apparent survival models. No mark-recapture

analyses except 'known fate' models can discriminate between mortality and permanent

emigration, therefore when we speak of survival it is technically 'apparent survival', but

during the first season of life we expect no calves to emigrate from the study area. We

estimated neonate survival during their first season of life as a function of individual spot

traits using Program MARK to analyse complete capture-mark-recapture encounter histories

of giraffes first sighted as neonates (White and Burnham 1999). We analysed our encounter

histories using Pollock's Robust Design models to estimate age-specific survival (Pollock

1982; Kendall *et al.* 1995), with and without spot covariates, and ranked models using AICc

following Burnham and Anderson (2002). We examined linear and quadratic relationships of

all 11 spot traits on juvenile survival to determine whether directional, disruptive, or

stabilizing selection was occurring. Our study design did not allow for the detection of other

forms of selection such as balancing or frequency-dependent selection. Based on previous

analyses for this population (Lee *et al.* 2016a, b), we constrained parameters for survival (S)

and temporary emigration (γ' and γ'') to be linear functions of age, and capture and

recapture (c and p) were time dependent $\{(S(A + \text{covariate}), \gamma'(A), \gamma''(A), c(t), p(t)\}$, thus

only the spot trait covariates changed among models. Giraffe calf survival does not vary by

sex (Lee *et al.* 2016b), so we analysed all calves together. We tested goodness-of-fit in

encounter history data using U-CARE (Choquet *et al.* 2009), and we found some evidence for

lack of fit ($\chi^2_{62} = 97$, $P = 0.01$), but we felt this was largely due to lack of age effects in the

goodness-of-fit tested model, whereas age effects were included in our model selection and

estimation. Additionally, because the computed $\hat{c}$ adjustment was < 3 ($\hat{c} = 1.5$), we felt our

models fit the data adequately and we did not apply a variance inflation factor (Burnham and

Anderson 2002; Choquet *et al*. 2009).

210   **RESULTS**

We were able to quantify 11 spot traits using ImageJ, and found the traits with greatest

individual heterogeneity as measured by the CV were number of spots and area of spots

(these two traits were negatively correlated), and mode shade (**Table 1**). Traits with the least

individual variation were solidity and roundness (**Table 1**). Measurements were highly

repeatable and reliable with the low variation in measurements from different photos of the

same individual (**Table 1**).

217   We found no spot pattern traits that had significant PO regression coefficients

between calves and random cows, but two characters, circularity and solidity (tortuousness)

(**Fig 2**) were significantly correlated between calves and their mothers indicating heritability

(**Table 1**). PO slope for circularity was 0.52 (SE = 0.16) and for solidity was 0.53 (SE =

0.17). We calculated post-hoc power for our significantly heritable traits of circularity and

solidity using our observed effect size (r-squared = 0.25, equivalent to Cohen's f^2 = 0.33),

alpha level (0.05), and sample size (31), and found power = 0.87. With one-half the observed

effect size (r-squared = 0.125) and our same sample size, power = 0.53. To achieve power =

0.80 with r-squared = 0.125, the sample size would need to have been 56 mother-calf pairs.

226   Our survival analysis of 258 calves first encountered as neonates indicated there was

no evidence that individual covariates of spot traits significantly affected survival during the

first season of life, but model selection uncertainty was high (**Table 2**). The top-ranked model

was the null model of no covariate effects, and all covariate beta coefficient confidence

intervals included zero. Power analysis indicated our effective sample size of 944 was far

below the 2600 necessary to detect a 0.05 difference in survival probability with a power of

0.80.

233   DISCUSSION

Our photographic capture-recapture dataset spanning multiple years enabled us to objectively

and reliably quantify coat pattern traits of wild giraffes using image analysis software, and

demonstrate that giraffe coat pattern traits of spot shape are heritable from mother to calf. The

methods we described should serve as a framework for objective quantification of

mammalian coat pattern traits, and could also be useful for taxonomic or phenotypic

classifications based on photographic coat pattern data. We did not find evidence for fitness

consequences of individual variation in spot traits on juvenile survival suggesting that spot

pattern traits are currently not under strong directional, disruptive, or stabilizing selection in

our study population, but our sample size was insufficient to detect small differences in

survival.

244   Photogrammetry to remotely measure animal traits has used geometric approaches

that estimate trait sizes using laser range finders and known focal lengths (Lyon 1994; Lee *et*

al. 2016a), photographs of the traits together with a predetermined measurement unit (Ireland

et al. 2006; Willisch et al. 2013), or lasers to project equidistant points on animals while they

are photographed (Bergeron 2007). We hope the framework we have described using ImageJ

to quantify spot characteristics from photographs will prove useful to future efforts at

quantifying animal markings, and suggest the resultant trait measurements could be useful in

a formal cluster analysis (Kaufman and Rousseeuw 2009) to classify subspecies, polymorphic

phenotypes, or other groups based on variation in markings.

253   Masai giraffe spot patterns are particularly diverse among giraffe subspecies (Dagg

1968), and there are spot patterns in northern Tanzania that bear strong similarities to other

giraffe subspecies elsewhere in Africa. Two recent genetic analyses of giraffe taxonomy both

placed Masai giraffes as their own species (Brown *et al.* 2007; Fennessy *et al.* 2016), but the

lack of quantitative tools to objectively analyze coat patterns for taxonomic classification

may underlie some of the confusion that currently exists in giraffe systematics (Bercovitch *et*

al. 2017). We expect the application of image analysis to giraffe coat patterns will provide a

new, robust dataset to address taxonomic and evolutionary hypotheses.

Our analyses highlighted two aspects of giraffe spots that were most heritable and

which may have adaptive significance. Circularity describes how close the spot is to a perfect

circle, and solidity describes how smooth and entire the edges are versus tortuous, ruffled,

lobed, or incised. These two characteristics could form the basis for quantifying spot patterns

of giraffes across Africa, and gives field workers a new quantitative lexicon for describing

spots (**Fig 2**). Our mode shade measurement was a crude metric, and color is greatly affected

by lighting conditions, so we suggest standardization of photographic methods to control for

lighting if color is to be analyzed in future studies.

One possible explanation for the lack of juvenile survival effects from spot variation

in our analysis is the recent reduction in large predator density in our study area (Packer *et al.*

2011; Bauer *et al.* 2015). If the function of the coat markings is to provide anti-predation

camouflage, reduced predator densities due to trophy hunting and pastoralist retaliatory

killings of predators may have alleviated predation pressure on giraffe calves sufficiently to

remove the selection pressure for certain spot traits (Lichtenfeld 2005; Lee *et al.* 2016b).

Alternatively, the possibility remains that spot traits may serve adaptive functions such as

thermoregulation (Skinner and Smithers 1990), social communication (VanderWaal *et al.*

2014), or indicators of individual quality (Ljetoff *et al.* 2007), rather than anti-predation

camouflage, and thus may demonstrate associations with other components of fitness, such as

survivorship in older age classes or fecundity. Individual recognition, kin recognition, and

inbreeding avoidance also could play a role in the evolution of spot patterns in giraffes

(Beecher 1982; Tibbetts and Dale 2007; Sherman *et al.* 1997). Different aspects of spot traits

may also be nonadaptive and serve no function, or spot patterns could be affected by

pleiotropic selection on a gene that influences multiple traits (Lamoreux *et al.* 2010).

Our study revealed that several spot traits are heritable and, hence, able to respond to

selection in giraffe populations. However, given that these spot characteristic traits did not

significantly affect neonatal survival, the main theory for the adaptive nature of giraffe spots

(Langman 1977; Mitchell and Skinner 2003), we conclude that spot patterns are currently of

minor importance for the adaptation of our study population, although this may be a recent

consequence of fewer predators. It may also be that some patterns affect fitness only in

certain environments, or our measurements of spot traits may not accurately reflect the

aspects of spot patterns most important to adaptation for neonatal crypsis. Other aspects of

spot variation may prove to be more relevant to fitness, such as social effects of individual

recognition or kin recognition, or thermoregulation, and deserve further investigation. We

may also have simply lacked sufficient sample sizes necessary to detect small adaptation or

fitness effects. Small and very difficult to detect differences in fitness may have profound

influence on phenotypic evolution compounded over thousands of generations (Orr 2009).

Patterned coats of mammals are hypothesized to be formed by two distinct processes:

a spatially oriented developmental mechanism that creates a species-specific pattern of skin

cell differentiation and a pigmentation-oriented mechanism that uses information from the

pre-established spatial pattern to regulate the synthesis of melanin (Eizirik *et al.* 2010). The

giraffe skin has more extensive pigmentation and wider distribution of melanocytes than most

other animals (Dimond and Montagna 1976). Coat pattern variation may reflect discrete

polymorphisms potentially related to life-history strategies, a continuous signal related to

individual quality, or a combination of both. Future work on the genetics of coat patterns will

hopefully shed light upon the mechanisms and consequences of coat pattern variation.

306   **ACKNOWLEDGEMENTS**

307   This research was carried out with permission from the Tanzania Commission for Science

308   and Technology (COSTECH), Tanzania National Parks (TANAPA), the Tanzania Wildlife

309   Research Institute (TAWIRI), African Wildlife Foundation, and Manyara Ranch

310   Conservancy.

311   **FUNDING**

312   Financial support for this work was provided by Sacramento Zoo, Columbus Zoo, Tulsa Zoo,

313   and Cincinnati Zoo.

314   **DATA AVAILABILITY**

315   We have deposited the primary data underlying these analyses as follows:

316   - Sampling locations, original data photos, and spot trait data: Dryad

317   **REFERENCES**

Åkesson, M., Bensch, S., Hasselquist, D., Tarka, M., and Hansson, B. 2008. Estimating

heritabilities and genetic correlations: comparing the 'animal model' with parent-

offspring regression using data from a natural population. PLoS One 3:e1739.

doi:10.1371/journal.pone.0001739.

Arnold, T. W, 2010. Uninformative parameters and model selection using Akaike's

Information Criterion. Journal of Wildlife Management, 74:1175–1178.

Bauer, H., *et al*. 2015. Lion (Panthera leo) populations are declining rapidly across Africa,

except in intensively managed areas. Proceedings of the National Academy of Sciences

112:14894–14899.

Beecher, M. D. 1982. Signature systems and kin recognition. American Zoologist 22:477–

490.

Bercovitch, F. B., *et al*. 2017. How many species of giraffe are there? Current Biology

27:R136–R137.

Bergeron, P., 2007. Parallel lasers for remote measurements of morphological traits. Journal

of Wildlife Management 71:289–292.

Bolger, D. T., Morrison, T. A., Vance, B., Lee, D., and Farid, H. 2012. A computer☐assisted

system for photographic mark–recapture analysis. Methods in Ecology and Evolution

3:813–822.

Brown, D. M., *et al*. 2007. Extensive population genetic structure in the giraffe. BMC

Biology 5:57.

Burnham, K. P., and Anderson, D. R. 2002. Model selection and multimodel inference: a

practical information-theoretical approach. Springer-Verlag. New York, New York.

Caro, T. 2005. The Adaptive Significance of Coloration in Mammals. BioScience 55:125–

136.

Choquet, R., Lebreton, J.-D., Gimenez, O., Reboulet, A.-M., and Pradel, R. 2009. U-CARE:

Utilities for performing goodness of fit tests and manipulating CApture-REcapture data.

Ecography 32:1071–1074.

Cott, H. B. 1940. Adaptive coloration in animals. Methuen Publishing. London, United

Kingdom.

Dagg, A. I. 1968. External features of giraffe. Mammalia 32:657–669.

Dagg, A. I. 2014. Giraffe biology, behavior and conservation. Cambridge University Press.

New York, New York.

de Villemereuil, P., Gimenez, O., and Doligez, B. 2013. Comparing parent-offspring

regression with frequentist and Bayesian animal models to estimate heritability in wild

populations: a simulation study for Gaussian and binary traits. Methods in Ecology and

Evolution 4:260–275.

Dimond, R. L., and Montagna, W. 1976. The skin of the giraffe. Anatomical Record 185:63–

75.

Eizirik, E., *et al*. 2010. Defining and mapping mammalian coat pattern genes: multiple

357   genomic regions implicated in domestic cat stripes and spots. Genetics 184:267–275.

358   Falconer, D. S., and Mackay, T. F. C. 1996. Introduction to quantitative genetics. 4th edition.

359   Pearson/Prentice Hall. New York, New York.

360   Fennessy, J., *et al*. 2016. Multi-locus analyses reveal four giraffe species instead of one.

361   Current Biology 26:2543–2549.

362   Foster, J. B. 1966. The giraffe of Nairobi National Park: home range, sex ratios, the herd, and

363   food. African Journal of Ecology 4:139–148.

364   Garvie, M. R., and Trenchea, C. 2014. A three level finite element approximation of a pattern

365   formation model in developmental biology. Numerische Mathematik 127:397–422.

366   Hoekstra, H. E. 2006. Genetics, development and evolution of adaptive pigmentation in

367   vertebrates. Heredity 97:222–234.

368   Ireland, D., Garrott, R. A., Rotella, J., Banfield, J. 2006. Development and application of a

369   mass-estimation method for Weddell seals. Marine Mammal Science 22:361–378.

370   Kaufman, L., and Rousseeuw, P. J. 2009. Finding groups in data: an introduction to cluster

371   analysis. John Wiley & Sons. Hoboken, New Jersey.

372   Kelsh, R. N. 2004. Genetics and evolution of pigment patterns in fish. Pigment Cell

373   Melanoma Research 17:326–336.

374   Kendall, W. L., Pollock, K. H., and Brownie, C. 1995. A likelihood based approach to

375   capture–recapture estimation of demographic parameters under the robust design.

376   Biometrics 51:293–308.

377   Kruuk, L. E., Slate, J., and Wilson, A. J. 2008. New answers for old questions: the

378   evolutionary quantitative genetics of wild animal populations. Annual Review of

379   Ecology, Evolution, and Systematics 39:525–548.

Lamoreux, M. L., Delmas, V., Larue, L., and Bennett, D. 2010. The colors of mice: a model

genetic network. John Wiley & Sons. Hoboken, New Jersey.

Lande, R., and Arnold, S. J. 1983. The measurement of selection on correlated characters.

Evolution 37:1210–1226.

Langman, V. A. 1977. Cow☐calf Relationships in Giraffe (Giraffa camelopardalis giraffa).

Ethology 43:264–286.

Lee, D. E., Bond, M. L., Kissui, B. M., Kiwango, Y. A., and Bolger, D. T. 2016a. Spatial

variation in giraffe demography: a test of 2 paradigms. Journal of Mammalogy 97:1015–

1025.

Lee, D. E., Kissui, B. M., Kiwango, Y. A., and Bond, M. L. 2016b. Migratory herds of

wildebeests and zebras indirectly affect calf survival of giraffes. Ecology and Evolution

6:8402–8411. doi:10.1002/ece3.2561.

Lee, D. E., and Strauss, M. K. L. 2016. Giraffe demography and population ecology.

Reference module in earth systems and environmental sciences. Elsevier. Amsterdam,

Netherlands.

Lichtenfeld, L. L. 2005. Our shared kingdom at risk: Human–lion relationships in the 21[st]

century. Ph.D. dissertation, Yale University, New Haven, Connecticut.

Ljetoff, M., Folstad, I., Skarstein, F., and Yoccoz, N. G. 2007. Zebra stripes as an amplifier

of individual quality? Annales Zoologici Fennici 44: 368–376.

Lydekker, R. 1904. On the Subspecies of Giraffa camelopardalis. Proceedings of the

Zoological Society of London 74:202–229.

Lynch, M., and Walsh, B. 1998. Genetics and analysis of quantitative traits. Sinauer.

Sunderland, Massachusetts.

Lyon, B. E. 1994. A technique for measuring precocial chicks from photographs. The Condor

86:805–809.

Maini, P. K. 2004. Using mathematical models to help understand biological pattern

formation. Comptes Rendus Biologies 327:225–234.

Mills, M. G., and Patterson, L. B. 2009. Not just black and white: pigment pattern

development and evolution in vertebrates. Seminars in Cell and Developmental Biology

20:72–81.

Mitchell, G., and Skinner, J. D. 2003. On the origin, evolution and phylogeny of giraffes

Giraffa camelopardalis. Transactions of the Royal Society of South Africa 58:51–73.

Murray, J. D. 1981. On pattern formation mechanisms for lepidopteran wing patterns and

mammalian coat markings. Philosophical Transactions of the Royal Society B: Biological

Sciences 295:473–496.

Orr, H. A. 2009. Fitness and its role in evolutionary genetics. Nature Reviews Genetics

10:531.

Packer, C., Brink, H., Kissui, B. M., Maliti, H., Kushnir, H., and Caro, T. 2011. Effects of

trophy hunting on lion and leopard populations in Tanzania. Conservation Biology

25:142–153.

Paterson, S., Wilson, K., and Pemberton, J. M. 1998. Major histocompatibility complex

variation associated with juvenile survival and parasite resistance in a large unmanaged

ungulate population (*Ovis aries* L.). Proceedings of the National Academy of Sciences

95:3714–3719.

Pollock, K. H. 1982. A capture–recapture design robust to unequal probability of capture.

Journal of Wildlife Management 46:752–757.

Pratt, D. M., and Anderson, V. H. 1979. Giraffe Cow☐Calf Relationships and Social

Development of the Calf in the Serengeti. Ethology 51:233–251.

R Core Development Team. 2013. R: a language and environment for statistical computing.

R Foundation for Statistical Computing. Vienna, Austria.

Roff, D. A. 1997. Evolutionary Quantitative Genetics. New York: Chapman and Hall.

Schneider, C. A., Rasband, W. S., and Eliceiri, K. W. 2012. NIH Image to ImageJ: 25 years

of image analysis. Nature Methods doi:10.1038/nmeth.2089[C].

Sherman, P. W., Reeve, H. K., and Pfennig, D. W. 1997. Recognition systems. Pp. 69–96 in

Behavioural ecology: an evolutionary approach (J. R. Krebs and N. B. Davies, eds.).

Blackwell Scientific. Oxford, United Kingdom.

Sikes, R. S. and Animal Care and Use Committee of the American Society of Mammalogists,

2016. 2016 Guidelines of the American Society of Mammalogists for the use of wild

mammals in research and education. Journal of Mammalogy 97:663–688.

Skinner, J. D., and Smithers, R. H. N. 1990. The mammals of the Southern African Sub-

region. 2nd ed. University of Pretoria. Pretoria, South Africa.

Strauss, M. K., Kilewo, M., Rentsch, D., and Packer, C. 2015. Food supply and poaching

limit giraffe abundance in the Serengeti. Population Ecology 57:505–516.

Tang-Martinez, Z. 2001. The mechanisms of kin discrimination and the evolution of kin

recognition in vertebrates: a critical re-evaluation. Behavioural Processes 53:21–40.

Tibbetts, E. A., and Dale, J. 2007. Individual recognition: it is good to be different. Trends in

Ecology and Evolution 22:529–537.

VanderWaal, K. L., Wang, H., McCowan, B., Fushing, H., and Isbell, L. A. 2014. Multilevel

social organization and space use in reticulated giraffe (*Giraffa camelopardalis*).

Behavioral Ecology 25:17–26.

White, G. C., and Burnham, K. P. 1999. Program MARK: survival estimation from

populations of marked animals. Bird Study 46(Supplement):120–138.

Willisch, C. S., Marreros, N., and Neuhaus, P. 2013. Long-distance photogrammetric trait

estimation in free-ranging animals: A new approach. Mammalian Biology 78:351–355.

Wilson, A. J., *et al*. 2010. An ecologist's guide to the animal model. Journal of Animal

Ecology 79:13–26.

457   FIGURE CAPTIONS

Fig 1. Representative images of spot patterns of mother-calf pairs of Masai giraffes (Giraffa

camelopardalis tippelskirchii) from the Tarangire ecosystem, Tanzania used in this study.

The blue rectangle shows the area analysed using ImageJ to characterize spot pattern

traits.

Fig 2. Representative spot outlines from Masai giraffes in northern Tanzania and their

corresponding circularity and solidity values. Ranges of spot trait values from 213 calves

are given in parentheses.

Table 1. Summary statistics for parent-offspring regressions of spot traits of Masai giraffes in northern Tanzania. Mean trait values, SD

(standard deviation), CV (among-individuals coefficient of variation), Repeatability (mean % variation in measurements from different pictures

of the same individual), PO slope coefficients (represents one-half of total heritability), F-statistics, P values, and r-squared values are provided.

Significantly heritable traits are in bold.

	Number	Area	Perimeter	Angle	**Circularity**	Maximum Caliper	Feret Angle	Aspect Ratio	Roundness	**Solidity**	Mode Shade
Mean	18.9	0.04	0.99	87.96	0.51	0.29	88.2	1.69	0.63	0.84	6924050
SD	7.5	0.01	0.25	15.39	0.08	0.06	14.5	0.15	0.04	0.04	3930565
CV	0.40	0.39	0.25	0.17	0.15	0.19	0.16	0.09	0.06	0.05	0.57
Repeatability	11	11	13	4	9	8	7	5	3	2	13
PO Slope Coefficient	0.20	0.20	0.27	0.04	**0.52**	0.21	-0.15	0.19	0.08	**0.53**	0.44
PO Coefficient SE	0.23	0.21	0.18	0.20	**0.16**	0.21	0.15	0.18	0.17	**0.17**	0.22
$F_{1,29}$	0.76	0.87	2.27	0.04	**9.97**	1.01	0.91	1.11	0.19	**9.73**	4.16
P value	0.39	0.36	0.14	0.84	**0.0037**	0.32	0.35	0.30	0.66	**0.0041**	0.051
r-squared	0.03	0.03	0.07	0	**0.26**	0.03	0.03	0.04	0.010	**0.25**	0.13

Table 2. Model selection results for giraffe calf survival as a linear or quadratic function of

spot trait covariates. The top-ranked model was the null model of no covariate effects, and all

covariate beta coefficient confidence intervals included zero, indicating no evidence for

significant spot trait effects on calf survival. Model structure in all cases was {S(A +

Covariate) g"(A) g'(A) p(t) c(t)} with covariate structure in survival. Minimum AICc =

3247.45, W = AICc weight, k = number of parameters.

Covariate Model	ΔAICc	W	k
Null (no spot trait covariate)	0	0.15	45
Aspect Ratio	0.96	0.09	46
Roundness	1.12	0.08	46
Solidity	1.50	0.07	46
Number of Spots	1.79	0.06	46
Angle	1.90	0.06	46
Perimeter	1.98	0.05	46
Circularity^2	2.06	0.05	47
Feret Angle^2	2.10	0.05	47
Mode Shade	2.11	0.05	46
Maximum Caliper	2.30	0.04	46
Feret Angle	2.31	0.04	46
Circularity	2.32	0.04	46
Solidity^2	2.49	0.04	47
Angle^2	2.84	0.03	47
Number of Spots^2	3.11	0.02	47

Aspect Ratio^2	3.18	0.02	47
Roundness^2	3.32	0.02	47
Mode Shade^2	3.41	0.02	47
Maximum Caliper^2	3.71	0.01	47
Perimeter^2	4.53	0.01	47

| Pair# | Mother | Calf |

Shape	Circularity (0.10—0.94)	Solidity (0.52—0.96)
	0.15	0.66
	0.13	0.60
	0.88	0.94
	0.37	0.76
	0.40	0.83
	0.57	0.82
	0.44	0.76
	0.12	0.53
	0.83	0.96