1 **CropQuant: An automated and scalable field phenotyping platform for crop**
2 **monitoring and trait measurements to facilitate breeding and digital agriculture**
3

4 Ji Zhou[1,2,*], Daniel Reynolds[1], Thomas Le Cornu[1,3], Danny Websdale[1,3], Simon
5 Orford[2], Clare Lister[2], Oscar Gonzalez-Navarro[1,2], Stephen Laycock[3], Graham
6 Finlayson[3], Tim Stitt[1], Matthew D. Clark[1,3], Michael W. Bevan[2], Simon Griffiths[2,*]
7

8 [1]Earlham Institute, Norwich Research Park, Norwich UK
9 [2]John Innes Centre, Norwich Research Park, Norwich UK
10 [3]University of East Anglia, Norwich Research Park, Norwich UK
11 [*]Correspondence should be addressed to: ji.zhou@earlham.ac.uk &
12 Simon.Griffiths@jic.ac.uk
13

Abstract

Automated phenotyping technologies are capable of providing continuous and precise
measurements of traits that are key to today's crop research, breeding and agronomic
practices. In additional to monitoring developmental changes, high-frequency and
high-precision phenotypic analysis can enable both accurate delineation of the
genotype-to-phenotype pathway and the identification of genetic variation influencing
environmental adaptation and yield potential. Here, we present an automated and
scalable field phenotyping platform called CropQuant, designed for easy and cost-
effective deployment in different environments. To manage infield experiments and
crop-climate data collection, we have also developed a web-based control system
called CropMonitor to provide a unified graphical user interface (GUI) to enable real-
time interactions between users and their experiments. Furthermore, we established a
high-throughput trait analysis pipeline for phenotypic analyses so that lightweight
machine-learning modelling can be executed on CropQuant workstations to study the
dynamic interactions between genotypes (G), phenotypes (P), and environmental
factors (E). We have used these technologies since 2015 and reported results
generated in 2015 and 2016 field experiments, including developmental profiles of
five wheat genotypes, performance-related traits analyses, and new biological insights
emerged from the application of the CropQuant platform.

Introduction
The great American wheat breeder and agricultural innovator Orville Vogel once
stated, the plant we are looking for is in our plots, but we have to be there when it is.
In order to select varieties with greater yield potential and enhanced environmental
adaptation, agricultural practitioners, including breeders, farmers and crop researchers,
have been optimising trait combination since the beginning of agriculture[1,2]. Four
decades after temporary success in ensuring global food security[3], we are now facing
an even bigger challenge to feed generations to come[4]. Due to a narrowing range of
available genetic diversity of modern crop germplasm collections[5] and increased
fluctuations in growing conditions[6], there is increasing emphasis placed on exploiting
new sources of genetic variation to enhance environmental adaptation and sustainable
yield in crop landraces and wild relatives[7]. To identify and assess these types of traits,
multiple regular measures of crop growth and development are required to quantify
subtle and dynamic phenotypes from many plots in different growing environments,
demanding new screening technologies to integrate field environmental datasets with
multi-scale phenotypic analyses to understand genotype-by-environment interactions
(GxE) and associate them to genetic variation[8–10].
51
In contrast to current field phenotyping methodologies, which are still involving
laborious manual scoring and relatively subjective selection, modern genetic and
genomics techniques are being rapidly deployed in breeding and crop research to
identify and utilise traits such as improved stress tolerance and disease resistance[11].
For example, quantitative trait locus (QTL) analysis and genome-wide association
studies (GWAS) are used to identify loci[12], whole genome sequencing used to reveal
gene content and genetic variation[13], and marker-assisted selection (MAS) and
genomic selection used to breed new lines with favourable alleles[14,15]. Therefore, field
phenotyping and the integration of environmental data with meaningful phenotypic
analyses are a key bottleneck that limits the potential of recent advances in crop
genetic and genomic technologies[16,17].
63
Remote sensing platforms[18] and open image-based analytics software libraries[19]
start to enable researchers, breeders and agronomists to develop new approaches to
understand and improve crop performance. For instance, unmanned aerial vehicles
(UAVs) and light aircraft are being used to study crop performance and field
variability[20,21]. Satellite imaging[22–24] and ground-based portable devices[25,26] have been
applied to take snapshots of crop growth to estimate yield-related traits using canopy
photosynthesis rate and normalised difference vegetation indices (NDVI). Field-based
agricultural vehicles have been developed to capture physiological and developmental
traits during the growing season[18,27]. Finally, large imaging platforms equipped with
3D laser scanners (e.g. near-infrared laser lines and Light Detection and Ranging,
LiDAR) and multi- or hyper-spectral sensors are applied to automate crop monitoring
of a fixed number of pots or plots, either in the field[28,29] or in greenhouses[12,30].
Although these advances are making important contributions to the research domain,
there are limitations and challenges associated with their usage such as high costs,
restricted mobility and scalability, limited frequency of screening, and inadequate
software tools for phenotypic analyses[31,32]. In particular, while satellite imagery and
UAVs are capable of screening tens of thousands of plots at multiple locations, their
applications are subject to civil aviation rules, low spatial resolution and bad weather
conditions such as heavy rainfall, strong wind and cloud coverage. Ground-based

portable devices and vehicles have shown greater mobility and high-resolution field
data between multiple sites; however, they require experienced specialists to operate,
limiting their applications to infrequent phenotyping. Still, stationary large imaging
platforms are providing key data on dynamic crop growth and GxE interactions; but
their scale of operation is restricted and they are relatively expensive for less well-
funded research laboratories to access. Furthermore, they mostly rely on proprietary
analytics software for data management and trait analysis, requiring ongoing licensing
maintenance to use software products and extra fees if tailored functions are needed.
For these reasons, it is challenging for researchers and breeding communities to adopt
new phenotyping approaches due to their expenses, lack of suitable software, limited
scope of operation and maintenance costs[33]. The ability to facilitate crop improvement
programmes at multiple scales and locations is still limited.
95

To enable the next-generation breeding and associated crop research[34], affordable
field phenotyping technologies need to be developed. New methods should exploit
up-to-date remote sensing technologies together with state-of-the-art computer vision
and software solutions, to equip researchers with diverse tools for multi-scale field
phenotyping needs. The work described here aims to address these challenges by the
development of an automated field phenotyping platform called **CropQuant**, which
integrates cost-effective hardware with open source software capable of complex
analytic solutions. We demonstrate applications of CropQuant (CQ) through multiple
growth phenotypes measurements based on defined wheat genotypes over two
growing seasons. Dynamic predictive growth models were also created to forecast the
performance of wheat genotypes under varied environment conditions.
107

Materials and Methods
An open and low-cost hardware design
In order to carry out automated phenotyping, we have deployed many low-cost crop
monitoring CQ workstations (*terminals*) operating jointly via a preinstalled or self-
operating infield wireless network. **Figure 1** shows the system architecture of the CQ
platform. A terminal is centred around a simple single-board computer (**Fig. 1a**),
running a customised Python-based analytic software package on the Linux *Debian*
operating system (OS), integrating *Pi* or USB camera sensors (e.g. red-green-blue
(RGB) or no infrared filter, NoIR), climate sensors (ambient and soil-based), sensor
circuit boards, with wired or wireless data communications (**Figs 1b&c**). The design
of the CQ platform is driven by the concept of Internet of Things (IoT)[35,36] as well as
how to utilise hardware and software resources that are widely available, so that crop
phenotyping solution can be scalable and affordable for the communities.
121

We have used these technologies in field experiments since 2015. **Figure 2** shows
an experimental scale of how CQ workstations were deployed for onsite and offsite
wheat assessment experiments in 2015 and 2016. To keep costs low so that the
technology can be adopted by the communities easily, we have developed different
CQ versions (**Figs 2a-d**). For example, an all-in-one CQ (**Fig. 2a**) uses a *Raspberry Pi*
2 single-board computer to control internal hardware (**Figs 2b**), including (1) a *Pi*
camera sensor for time-lapse crop photography, (2) a tailored circuit board (**Fig. 2c**)
to integrate climate sensors for collecting environmental data (i.e. soil temperature
and moisture, ambient temperature and humidity, and light levels), (3) a USB WiFi
dongle (or a radio transmitter) for data transfer and remote interactions, and (4) a USB

flash drive for local data storage. **Supplementary Figure 1** demonstrates how an all-
in-one CQ was used in field experiments.
134

Besides the relatively costly all-in-one version ($230-$240 to build), other more
specialised versions (**Fig. 2d**) were much cheaper to produce. For example, a business
card size CQ (version 4, $80-90) uses a *Pi* Zero computer and is tailored for crop
photography. A version 2 CQ (**Fig. 2b**, $170-180) can mount different sensor groups
(ambient or soil-based) for assessing agronomic characterisation and crop adaptation.
With hardware modularity in mind, we have tested a range of single-board computers
(e.g. the *Raspberry Pi* series, Intel's Galileo and Edison) for performing simple infield
image analysis as well as integrating modular components. Although we finally chose
Pi computers due to its performance-to-price ratio and extensive community support,
the platform can operate on other single-board computers, if *Pi* computers are not
available. For the peripheral hardware, we used off-the-shelf weatherproof containers
(IP67 rating) together with micro-USB and Ethernet couplers (IP66 rating) to ensure
environmental endurance and outdoor deployment. **Supplementary Figure 2** shows
the hardware design of a version 2 CQ. A full hardware list and a construction manual
are included in **Supplementary Note 1.**
150

Offsite self-operating mode
Figure 2e demonstrates an offsite field experiment using the CQ platform in 2015,
where 10 field workstations were deployed to monitor canopy development (**Movie 1**)
and crop growth (**Movie 2**) on one-metre wheat plots. CQ devices were powered by
lead acid batteries with trickle charging from solar panels. To operate the device with
minimal energy requirements, we have implemented a headless access mode to carry
out wired data transfer. Besides the programmed imaging task, the system was only
wakened if an Ethernet connection (i.e. local area network, LAN) was established.
Offsite CQs were self-operating and used to perform image-based phenotyping. The
infield imaging script running on CQs can be seen in **Supplementary Note 2**.
161

Onsite networking mode
For onsite experiments (**Fig. 2f**), CQs were powered by 5V/2A power supplies and
connected to a field WiFi network. There were 14 networked CQ terminals (21 at
peak time, with two dedicated for tiller abortion studies) jointly operating in 2016,
monitoring 12 six-metre wheat plots to study performance-related traits and yield
production. Although not thoroughly tested, for pre-installed WiFi network, the scale
of the CQ platform can be increased by adding more standard routers to allow more
CQ connections. We have added new functions to networked CQs including wireless
control, programmable imaging, and on-board quality control (Online Methods). For
instance, end-users can access CQ workstations remotely for real-time inspection,
either using a portable device (e.g. a tablet or a smartphone) in the field or an onsite
office computer (**Fig. 2g** and **Supplementary Fig. 3**). They can check the field in
different regions to: (1) review historical crop images, (2) initiate new experiments,
and (3) transfer crop-climate data to external computing storage.
176

These monitoring activities are administered by our web-based control system,
CropMonitor (Figs 2h-j and **Supplementary Fig. 4**), where the status of each CQ
terminal is updated constantly with information such as online or offline status,
operational mode (e.g. green for operating, amber for idle, and red for operation error
or ending tasks), representative daily images, micro-environment readings, and the

usage of computing resources (i.e. CPU and memory). Furthermore, the CropMonitor
system can support a range of tasks. For example, when deploying CQs in the field,
CropMonitor can activate live streaming between a CQ terminal and a smart device
(e.g. a smartphone or a tablet) to enable the calibration and installation of CQ devices
(**Supplementary Fig. 5**). During the experiment, the CropMonitor system can
establish a mesh network (based on all-in-one CQs) to support data communications
between the field and external servers (Online Methods and **Supplementary Note 3**).
Users can reposition CQ terminals at any time to change or initiate new monitoring
tasks. Notably, the IoT-style setting can improve the mobility of the CQ platform. For
example, **Supplementary Figure 6** shows a speed breeding experiment[37] monitored
by CQs over a 75-day period in 2017, which was accomplished by moving CQs to
indoor, i.e. a growth chamber (the speed breeding condition) and a glasshouse (the
control condition).
195
To increase the scalability of the phenotyping platform, we recently developed a
mesh network system to connect infield terminals with or without any pre-installed
network infrastructure. **Supplementary Figure 7** illustrates the network topology,
where all-in-one CQs are operating jointly as backbone *routers* (i.e. cluster servers),
running both dynamic host configuration protocol (DHCP) and virtual network
computing (VNC) servers for networking. Cheaper CQs (versions 2-4) are connected
to the *routers* as terminals (i.e. *end devices*). Depending on the number of *routers* and
the coverage of WiFi dongles (25-30 metres) or radio transmitters (several hundred
metres), the field mesh network can be expanded or downsized flexibly. We dedicate
an all-in-one CQ as a *coordinator* to control data communications between terminals
and external networks. Meanwhile, computing tasks such as image selection, quality
control and initial data annotation are distributed to terminals to reduce computational
burden of the in-depth trait analysis (**Supplementary Note 4**).
209
Additionally, to verify the outputs of low-cost remote sensors integrated in CQs, we
have utilised the meteorological outputs of a commercial weather station (**Figs 2k&l**),
including temperature, rainfall, photosynthetically active solar radiation, wind speed
and relative humidity. Phenotypic and climate datasets were saved on an onsite high-
performance computing (HPC) cluster (SGI UV2000 system with Intel Xeon cores)
for durable data storage (**Fig. 2m**). Since the application of the CQ technology, we
have successfully accomplished three tasks essential for the next-generation field
phenotyping[17]: (1) *continuous monitoring* via time-lapse crop photography, (2) *infield*
evaluation through networked terminals and the CropMonitor system, and (3) *efficient*
data transfer using distributed computing and wireless data communications through
an infield network (**Supplementary Fig. 8**).
221
The high-throughput analysis pipeline
In order to enable accurate delineation of the genotype-to-phenotype pathway and
identify genetic variation influencing environmental adaptation and yield potential,
we chose a high-frequency (two-three times per hour) and high-precision (2592x1944
pixels per image) phenotyping approach to monitor the morphological change of
crops. After the crop photography phase, we exploited open image processing and
machine learning libraries such as OpenCV[38], Scikit-learn[39] and Scikit-image[40] and
developed an automated analysis pipeline to extract biologically relevant outputs.
230

We designed the analysis pipeline to be executable on either a workstation PC or an
HPC cluster. Firstly, to arrange the collected image series, we have developed a
selection algorithm to choose representative images based on their size, clarity[41],
imaging dates and genotypes (**Fig. 3**, *Step 1*). Only high-quality images were retained
for trait analysis (Online Methods and **Supplementary Note 5**). All datasets were
archived in a central repository such as HPC clusters for future reference. Then, we
developed a referencing algorithm to define the location of a monitored plot over time
(**Fig. 3**, *Step 2*). In real-world agricultural and breeding situations for which CQs are
deployed, strong wind, heavy rainfall, irrigation and chemical spraying can lead to
modest camera movements, causing cross-reference problems when comparing trait
analyses for a plot over time (**Movie 3**). To resolve this issue, we have designed the
referencing algorithm to identify the initial plot location so that each image in the
series can be transferred to the same position for comparison. For example, the
algorithm detects 2D coordinates of white reference canes installed in the plot and
dark markers on a ranging pole for height scales using colour- and shape-based
feature selection[39]. Then, it classifies pixels into five groups to represent the canopy
space and background objects such as wheel tracks, sky, and the reference canes using
simple k-means[42] and spectral clustering[43] algorithms. Finally, a pseudo 3D reference
system is established to record important coordinates of the plot region, the canopy
space, and height markers, together with converting measurements from pixels to
metric units such as centimetres (Online Methods and **Supplementary Note 6**).
252
Following Step 2, we integrated the initial reference location into a performance-
related trait analysis algorithm. For a given image series, the algorithm applies an
adaptive intensity and gamma equalisation method[44] to minimise colour distortion
caused by varied field illumination. Then, it tracks geometric differences[38] between
the monitored plot and the reference location. If the plot location has changed in a
given image, a geometric transformation method[45] will be applied to reposition the
image, removing areas outside the plot region and may or may not generate a black
bar to the top of the image (**Fig. 3**, *Step 3*). Within the plot region, the algorithm
detects the visible part of the ranging pole (**Movie 4**) as well as the canopy space for
height measurement. For instance, to measure the height of the canopy, an entropy-
based texture analysis is used to determine whether the canopy region is changing
between two consecutive images using gray-level co-occurrence matrices (GLCM)[46].
If positional changes (moving up or down, depending on growth stages) are identified,
the canopy height is recorded and corner-featured points[47] are detected (**Fig. 3**, *Step 4*),
generating many red pseudo points casting in the canopy region for measuring canopy
height (**Movie 5** and **Supplementary Note 7**). These pseudo points can also be used
to represent the tips of erect leaves at stem elongation or jointing (the Zadoks scale[48],
growth stages, GS 32-39), reflective surfaces of curving leaves or crop heads between
booting and anthesis (GS 41-69), and corner points on spikelets during senescence
(GS 71-95). Using the trait analysis algorithm, we have computed the dynamic height
changes to present growth patterns for different wheat genotypes (**Fig. 3**, *Step 5.1*).
274
Developmental related trait measurements
In addition to the canopy height, we also developed functions to calculate other traits.
For example, vegetative greenness is calculated based on the normalised greenness
value (0-255) for a given plot over time. The output was used to assess the change of
green biomass and vegetation period. We used this trait to evaluate a Stay-Green
mutant (prolonged green leaf area duration with delayed leaf senescence; **Fig. 3**, *Step*

5.2). The main orientation of a plot (0-180^{o}) is also quantified based on edge detection
methods[49], representing the alignment of stems to estimate the change of stem rigidity
(**Fig. 3**, *Step 5.3*). Using this trait, we have identified lines with higher lodging risk
either during ripening or when interacting with heavy rainfall or strong wind (Online
Methods and **Supplementary Table 1**).
286

Results
Use case 1 – Monitoring five wheat NILs
The diverse environments for which wheat has been adapted to grow provide
opportunities for us to explore the dynamic interactions between genetic diversity and
phenotypic traits under varied environmental conditions[50]. To test the CQ platform,
we chose wheat near-isogenic lines (NILs, Online Methods) to examine a number of
key performance-related phenotypes in the same *Paragon* (a UK spring wheat variety)
genetic background[51]. **Figure 4** demonstrates five dynamic developmental profiles
generated from the experiment between May and August 2015, a 95-day period. The
experiment was conducted in plots in a field which is 2.1 miles away from Norwich
Research Park UK (see the plot layout in **Supplementary Table 2**) and all five NILs
were monitored twice per hour. The genotypes were: (1) Late-DTEM (days to ear
emergence[48], the number of days between sowing and ear emergence; *late* means
GS55 is delayed), with *Ppd-1* loss of function (lof); (2) Early-DTEM (GS55 is moved
forward), with *Ppd-D1a* photoperiod insensitivity; (3) Short stems, *Rht-D1b* semi
dwarfing; (4) Stay-Green, a stay green mutant; and (5) *Paragon* wild type (WT).
303

To compare the performance of the NILs, we used *Paragon* WT as the reference
line and highlighted six key growth stages, from stem elongation or jointing (GS 32-
39) to ripening (GS 91-95). The thermal time (degree-day, ^{o}Cd, using a 0°C base[52])
was also used as a heuristic tool[53] to normalise the crop growth. The five growth
curves (1258-2297 °Cd) approximately followed a sigmoid curve. At the beginning of
the experiment, *Ppd-D1a* NIL (Early-DTEM, coloured amber) was already at the end
of the jointing stage (GS37-39) and hence was the first to reach a maximum height;
whereas *Ppd-1* lof (Late-DTEM, coloured blue) was the last to increase in height. By
cross-referencing developmental profiles based on six growth stages, we noticed that
although *Ppd-D1a* and *Rht-D1b* (Short-Stem, coloured red) had similar maximum
heights (83.4cm and 80.6cm), the latter displayed a relatively steady rate of increase
in stature. *Ppd-1* lof's growth was the most delayed line, resulting in an extended
period of vegetation, stem extension, and overall time to ear emergence. As this
genotype has received the most thermal time units, it was the tallest line in the field
experiment. Although all NILs experienced some degree of height reduction due to a
significant storm on 24th July 2015, *Paragon* WT (coloured purple) presented a much
lower lodging risk, as it maintained its height afterwards. To verify the phenotypic
observation, we scored heading dates and canopy heights manually on the same plots
and obtained a Pearson correlation coefficient of 0.986 (**Supplementary Table 3**).
323

We summarised different temperatures and accumulated degree-days (ADDs) in
both 2015 and 2016 growing seasons (**Fig. 5a**). As the average temperature in 2015 is
much lower than in 2016, we used a fixed ADD period (1250-2300 °Cd) to segment
crop growth under different climates. Within the same ADD period, **Figure 5b** shows
dissimilar growth curves of *Paragon* WT (**Supplementary Table 4**). The 2015 curve
was much steeper during stem elongation (GS32-59, 1250-1750 °Cd), possibly

reflecting the cold spring. Flowering half complete (GS65) was reached on 24[th] June
2015 (an early drilling late maturity mode). While the 2016 curve (values are the
means of two biological replicates) had a steadier and extended development due to a
warm spring. GS65 was reached on 12[th] June 2016, 12 days ahead of 2015 (a late
drilling early maturity mode). Using the CQ platform, not only have we collected
high-frequency crop climate datasets, we also could identify dynamic developmental
variances for genotypes under different climate patterns.
337
Use case 2 – New biological insights into growth patterns
High-frequency and high-resolution deep phenotyping has already been employed in
human disease research to reveal the underling mechanisms of individual's disease[54].
In plant research, the similar approach is being adopted to characterise phenotypes of
plant responses to environmental challenges for field experiments[55]. While applying
CQs in wheat assessment experiments, we have explored new biological insights into
dynamic growth patterns using the large phenotypic data captured in the field.
345
Figure 6 presents some preliminary results of how we utilised the high-frequency
phenotypic data to extract underlying growth patterns for *Paragon* genotypes. Initially,
we calculated daily relative growth rates (RGR, comparing with the previous day) for
the five NILs monitored in 2015 and compared them with two *Paragon* WT measured
in 2016. All RGR data were aligned by the associated growth stages for comparison,
showing that all lines were active from jointing to flowering and became inactive after
grain-filling (**Fig. 6a**). After that, to study the change of RGR during the growth
stages (GS32-69), we explored the frequency and the degree of the RGR data. To be
precise, we converted the data series from its original time domain (with equal daily
readings) to the frequency domain using a fast Fourier transform algorithm (FFT)[56].
After the conversion, we separated the frequencies (x-axis, cycles per day, i.e. the
frequency of growth) and the magnitude spectrum (y-axis, normalised amplitudes, i.e.
the degree of growth) and generated underlying growth patterns of all the monitored
lines (**Supplementary Note 8**). Noticeably, for *Paragon* WT, although temperatures
and developmental profiles were significantly different between 2015 and 2016, the
underlying growth patterns for *Paragon* WT in both years were very similar (**Fig. 6b**).
We identified two distinct growth peaks: (1) around 15 days (15.3 days and 15.6 days
respectively) and (2) seven-eight days (8.4 days and 7.8 days), indicating that
Paragon WT is likely to control its underlying growth pattern based on the number of
elapsed days instead of other factors such as temperatures.
366
For Late-DTEM and Early-DTEM NILs whose genetic backgrounds only differ by
carrying alleles such as *Ppd-D1a* and *Ppd-1* lof, their growth patterns also contain
two peaks (**Fig. 6c**): (1) similar to *Paragon* WT, seven-eight days (7.1 days and 7.7
days) and (2) 23.0 days for Early-DTEM and 15.3 days for Later-DTEM. For other
Paragon NILs (e.g. Stay-Green mutant and Short-Stem), although the patterns were
slightly different, we found that at least one growth peak was close to the region of
seven-eight days (**Supplementary Note 8** and **Supplementary Fig. 9**). To verify the
FFT approach, we created a hypothetical *Paragon* growth data by combining all
Paragon NILs across two years as a technical replicate. **Figure 6d** shows that no clear
growth peaks can be detected from the hypothetical datasets. Additionally, we have
applied the FFT approach to converted the RGR series with equal degree-day
readings; similarly, no clear growth peaks can be identified (**Supplementary Note 9**).
379

As all the tested *Paragon* NILs show a growth peak at seven-eight days and only
Late-DTEM had a growth peak at 23 days, this might provide some insights into the
mechanism of *Ppd-D1a*. The cyclical 23-day peak in growth over the common 15-day
peak might reflect a changed output of the circadian clock of which *Ppd1* (PRR7 in
Arabidopsis[57]) is accelerating the development in a cyclical manner. We are currently
conducting a number of experiments, from gene expression to cell biology to advance
our understanding of this discovery generated by the CQ platform.
387
Use case 3 – GxE predictive modelling
Crop modelling is used in breeding and crop research for integrating complex external
and internal variables to understand GxE interactions and genetic systems. Many
existing models use genotypes (G) and environmental factors (E) as input parameters
to predict phenotypes (P) as the output of the models[58–62]. Similarly, we established a
light-weight GxE model to predict crop growth using continuous crop-climate data
collected by the CQ platform. Also, we used the computational power of CQ's single-
board computers to explore how to operate the GxE model on a daily basis together
with infield phenotyping tasks.
397
The key input components (environmental factors, growth stages and growth traits)
of the model and how it was utilised for predicting growth in fluctuating growing
conditions are summarised in **Figure 7**. Firstly, we selected environmental factors that
were strongly correlated with the performance-related traits such as RGR and height
at four key growth stages (from jointing to flowering) using *Pearson* correlation
(**Supplementary Table 5**). This approach has identified five out of the 12 factors
(p<0.01), including degree-day, solar radiation, rainfall, temperature, and daily light
duration. Two heat maps (**Fig. 7a**) were produced to present the selected factors at the
four stages (Online Methods and **Supplementary Note 10**). After that, we built a
stage-based predictive model using training datasets of growth stages in 2015 and
2016 (Online Methods and **Supplementary Note 11**). We employed support vector
machines (SVM)[63], a popular machine learning algorithm for classification, with
radial basis function kernels to classify growth stages. **Figure 7b** illustrates the
classified growth stages (coloured blue) benchmarked against the stages scored by
expert crop physiologists (coloured red). **Supplementary Figure 10** illustrates the
performance of the model when classifying the timing and duration for other wheat
genotypes. We found that SVMs trained on two-season *Paragon* WT data had the
highest scores using the benchmarking approach. Although the model modestly
mistimes in booting (GS41-49) and heading (GS51-59) due to their short duration, we
are adding new training data acquired in 2016 and 2017 to improve the model.
418
On the basis of the identified environmental factors and growth stage modelling, we
explored a set of linear regression models to establish a global predictive model to
forecast the continuous growth curve of *Paragon* genotypes, an approach that can be
used to help farmers and breeders to optimise crop growth and genotype selection in
the future. **Figure 7c** shows how the growth predictive model performs on the
hypothetical *Paragon* growth data (a technical replicate with mean squared error:
46.593, correlation: 0.999). The environmental factors and corresponding daily RGR
data are grouped together for each stage and a linear regression model is then
constructed to fit growth rate within the forecast growth stages, together with an
ordinary least squares method to determine model coefficients. A relative growth rate
estimate equation $y_t^s = \beta^s x_t + c^s$ is used, where x_t is the environmental data at time

point t, β^s is the model coefficients (weight vectors), and c^s are constant offsets (i.e.
intercepts for each growth stage). The super-script s denotes different growth stages
(Online Methods and **Supplementary Note 12**).

Growth estimates at each stage are concatenated to form a single vector, based on
which height values (in centimetres) and stage-based growth rates are calculated.
Using the predictive model, we produced growth estimates for four wheat NILs
(**Supplementary Fig. 11**). In this way, we compared how well the model performed
with respect to the trait analyses recordings obtained from the CQ platform. To link
the predictive model with crop agronomy, we also calculated the average standard
deviation (SD) of the predicted crop height values, so a real-time warning message
can be triggered on the CropMonitor control system. For example, if crop is growing
outside the safe region (the bounds of its estimated height region, ±1SD, **Fig. 7d**),
warning messages will be generated to inform the users that the growth is either too
quick or too slow.

Discussion
With the development of modern high-throughput and low-cost genotyping platforms,
the current bottleneck in breeding and crop research lies in phenotyping. Here, we
describe CropQuant, an automated and scalable field phenotyping system which we
believe can enable researchers with a toolkit that can fulfil multi-scale and diverse
phenotyping needs for a broader plant research and breeding communities. To deliver
the technology, we used the IoT in agriculture ethos to combine networked sensors,
distributed computing hardware, computer vision, image analysis, machine learning
and modelling to provide high-frequency field phenotyping together with GxE growth
predictions, in near real time and in a manner which closely match human scoring.

To enable the CQ system to monitor and compare developmental changes between
wheat genotypes using phenotypic data collected under different climate patterns, we
carried out an open hardware R&D strategy so that CQ devices could be built and
reproduced by other research groups in a relatively easy and cost-effective way. We
utilised widely available *Pi* single-board computers and off-the-shelf climate sensors
to facilitate the hardware design. For example, a CQ device can be equipped with a
range of imaging sensors (e.g. RGB or NoIR *Pi* cameras, or USB and IP cameras) for
varied experiments, including using *Pi* cameras to perform side-view and top-view
imaging for crop growth and canopy development studies, setting up USB endoscope
cameras below the canopy to study tiller abortion, and connecting IP cameras for field
level monitoring. Environment sensors were grouped by functions, i.e. ambient and
soil-based. So, dissimilar sensor groups could be selected for different experiments.
To increase the capability and usefulness of the low-cost CQ device, we provided the
hardware construction manual and the circuit board design (**Supplementary Fig. 12**)
so that imaging and remote sensing functions could be integrated as well as expanded.
We believe that, following the current hardware design, crop-climate data collection
in the field can be standardised, data evaluation and communication can be carried in
the field, and terminal nodes can be scalable on the CQ platform. Notably, due to the
limited manufacturing ability and R&D funding, the maximum indoor and outdoor
terminal nodes operating on the CQ platform at the same time was 21, although in
theory the system can operate at least 255 nodes simultaneously.

From the software development perspective, we created the CropMonitor control
system to provide users with a unified web-based platform which can be used to
connect phenotyping hardware with ongoing experiments in an integrated design. Not
only does it allow different users to monitor experiments in real time, but it can
incorporate different solutions in one shared web place to support experiments at
different phases, i.e. from device deployment to the completion of the experiment.
With crop-climate data collected in a standard manner, we developed a number of
open-source trait analytic algorithms to measure multiple performance-related traits to
identify genetic variation under different climate patterns. The software solutions
have been evaluated by noisy images caused by complex field environment. Still, it
can reliably execute the trait analysis tasks. To verify the results generated by the
analysis pipeline, we have scored the performance phenotypes manually on the same
plots over two growing seasons and obtained a strong correlation. Furthermore, we
established dynamic predictive growth models to forecast the performance of wheat
genotypes under varied growing conditions, which could be valuable for agronomic
practices. Although the results are promising, it is noticeable that more training
datasets are required to improve these models, ideally from varied growing conditions.
As the software solutions were implemented on open image analysis, computer vision
and machine learning libraries, they can be easily adopted and expanded for other
experiments by the communities. To support computational users to understand our
work, we have provided detailed comments in our source code.
500
From a biological perspective, the use of key performance traits generated by the
CQ platform can be an excellent tool for screening early establishment, vegetation
period, flowering, growth patterns and lodging. For example, vegetative greenness is
a useful marker to quantify senescence; utilising the side-view movie, we can closely
monitor the process of wheat aging, from the lower stem to the canopy region, a new
approach to determine physiological maturity which is important for researching grain
development and ripening. Also, continuously monitored greenness can be used in
plant pathogen interaction to analyse the activity of pathogens on the leaf surface, as
broad yellowish symptoms can be observed from susceptible plants (e.g. rust in
wheat). Moreover, crop-climate data acquired by the CQ platform can also assist us to
carry out novel biological discoveries. For instance, we are using deep-learning neural
network architectures to train a convolutional neural network classifier (CNN) to
quantify yield component traits such as spike per unit area and spike/spikelet number
(**Supplementary Fig. 13**).
515
The CQ platform, in combination with networked remote sensors, the web-based
control system, computational analytic solutions, and machine-learning based growth
modelling, has enabled a cost-effective and scalable field-scale phenotyping of wheat
germplasm. Multiple performance-related measurements were quantified in near real
time and related to growing conditions. This technology has the potential for multiple
applications in breeding and crop production, for example, to optimise the timing of
fertiliser applications, irrigation, and predicate harvest dates for maximising yields in
different agronomic scenarios. In crop breeding systems, regular field monitoring
using the CQ platform identified multiple growth and developmental variables that
provided statistically significant phenotypic analysis. These can increase the accuracy
of breeding values, particularly for environmental response factors. With more field
experimental data collected from different environments feeding into the system, the
GxE predictive model and analytics software pipeline can be continuously improved.

In particular, as the field of machine learning has progressed enormously in the last
few years, our ability to model complex nonlinear functions and extract high-level
phenotypic features is also growing. For example, we are applying deep learning (i.e.
CNN and recurrent neural networks, RNNs) to learn and extract features from
multidimensional imaging data (including visible and invisible spectrums) that are
exceptionally difficult to accomplish through traditional image analysis approaches.
Hence, we are consistently exploring deep learning to provide more accurate crop
growth and development scores as well as yield-related trait quantifications, offering
considerable value to the communities. Our future plan for the CQ platform is to
improve the hardware to enhance mobility and modularity, and work with a broader
plant research communities to jointly increase the software package for capability and
applicability in different growing conditions. So, we could finally deliver real-time
infield analysis and integrate field-based phenotyping, UAVs, and satellite into a
multi-level and multi-dimensional crop analytic system.
543

Conclusion
We believe that the CropQuant technologies described here may have a significant
impact on future crop research, breeding activities, and agronomic practices. The
reasons are: (1) the low-cost and widely available hardware centred by single-board
computers is capable of enabling tasks such as continuous crop monitoring, infield
evaluation and efficient data transfer, which are essential for the next-generation field
phenotyping; (2) automated trait analysis algorithms integrated in the CQ platform are
open-source and expandable software solutions, which are easily accessible and based
on community driven numeric and scientific libraries; (3) use cases presented in the
paper explain how to apply the CQ platform to study dynamic interactions between
genotypes, phenotypes, and environmental factors, which is capable of producing new
biological insights of growth patterns through phenotypic analyses. Moreover, our
work endeavours to address the affordability and scalability issue for the research
communities, which are independent from specific commercial hardware platforms
and proprietary or specialised software applications, allowing the utilisation of the CQ
platform to accomplish data annotation, performance-related phenotypic analysis, and
cross-referencing results freely by the academic communities. Our work confirms
previously reported results in the literature and produces novel approaches to enhance
the reproducibility of indoor and outdoor crop growth and development experiments.
Our case studies of wheat NILs are not limited. Natural variation, mineral or nutrient
stress and other crop species could also be monitored using the platform.
565

566 **Methods**
567 Methods and any associated references are available in the online version of the paper.
568 Note: Supplementary information is available in the online version of the paper.
569

Acknowledgements
We thank members of the Zhou laboratory and the Griffiths laboratory for fruitful
discussions. Prof C. Martin and Prof R. Morris at John Innes Centre for reading and
improving the manuscript. We thank S. Cossey and Prof Caccamo for supporting the
initiation of the project, M. Hewitt for helping to engineer the first prototype, C.
Mumford and her team for field trials, NBI infrastructure teams for infield network
testing, and research leaders at EI, JIC, TSL and UEA for constructive discussions.

This work was strategically funded by the BBSRC, Institute Strategic Programme
Grants (BB/J004669/1) and Core Strategic Programme Grant (BB/CSP17270/1) at the
Earlham Institute, GRO (BB/J004588/1) to M.W.B. at JIC. J.Z. was partially funded
by BBSRC's Designing Future Wheat Cross-institute Strategic Programme Grants
(BB/P016855/1) to Prof G. Moore. The biological research and related software and
hardware R&D led by J.Z. was partially supported by BBSRC's FoF award (GP105-
JZ1-B), an NRP Translational Fund (GP069-JZ1-Q), and the UK government's
Eastern Agri-Tech Initiative funding (GP080-JZ1-M).
585

Author contributions

J.Z, D.R., S.L., T.L.C., M.D.C., M.W.B. and S.G. designed research; J.Z., D.R., O.G.,
C.L. and S.O. performed the research; J.Z., D.R., T.L.C., D.W., and G.F. conducted
hardware design and the development of analytics software packages; J.Z., D.R.,
T.L.C., D.W., O.G., C.L., S.L., M.W.B. and S.G. contributed to analyse data; and J.Z.,
D.R., T.L.C., D.W., M.D.C., M.W.B and S.G. wrote the paper. All authors have read
and approved the final manuscript.
593

Competing financial interests

595 The authors declare no competing financial interests.
596

Open Access

Source code

606 Source code is freely available for academic usage, which can be downloaded at
607 https://drive.google.com/drive/folders/0B17ZL8AzLo8wNFJUVS1lOFkzb3M?usp=s
608 haring (an online Github repository is being prepared and will be updated in bioRxiv
609 as soon as possible)
610

References

1. Sauer, C. O. *Agricultural origins and dispersal*. (American Geographical Society, 1952).
2. Donald, C. M. The breeding of crop ideotypes. *Euphytica* **17,** 385–403 (1968).
3. Borlaug, N. E. Wheat breeding and its impact on world food supply. in *3rd International Wheat Genetics Symposium* 1–36 (Australian Academy of Science, 1968).
4. Pingali, P. Green Revolution:Impacts, Limits, and the path ahead. *Proc. Natl. Acad. Sci.* **109,** 12302–12308 (2012).
5. Tester, M. & Langridge, P. Breeding Technologies to Increase Crop Production in a Changing World. *Science (80-.).* **327,** 818–822 (2010).
6. Howden, S. M. *et al.* Adapting agriculture to climate change. *Proc. Natl. Acad. Sci.* **104,** 19691–19696 (2007).
7. McCouch, S. R. *et al.* Feeding the Future. *Nature* **499,** 23–24 (2013).
8. Malosetti, M., Ribaut, J. M., Vargas, M., Crossa, J. & Van Eeuwijk, F. A. A multi-trait multi-environment QTL mixed model with an application to drought and nitrogen stress trials in maize (Zea mays L.). *Euphytica* **161,** 241–257 (2008).
9. Houle, D., Govindaraju, D. R. & Omholt, S. Phenomics: the next challenge. *Nat. Rev. Genet.* **11,** 855–866 (2010).
10. Hawkesford, M. J. & Lorence, A. Plant phenotyping : increasing throughput and precision at multiple scales. *Funct. Plant Biol.* (2017). doi:10.1071/FP
11. Barabaschi, D. *et al.* Next generation breeding. *Plant Sci.* **242,** 3–13 (2015).
12. Yang, W. *et al.* Combining high-throughput phenotyping and genome-wide association studies to reveal natural genetic variation in rice. *Nat. Commun.* **5,** 1–9 (2014).
13. Brenchley, R. *et al.* Analysis of the bread wheat genome using whole-genome shotgun sequencing. *Nature* **491,** 705–10 (2012).
14. Araus, J. L. & Cairns, J. E. Field high-throughput phenotyping: the new crop breeding frontier. *Trends Plant Sci.* **19,** 52–61 (2014).
15. Bevan, M. W. *et al.* Genomic innovation for crop improvement. *Nature* **543,** 346–354 (2017).
16. Furbank, R. T. & Tester, M. Phenomics--technologies to relieve the phenotyping bottleneck. *Trends Plant Sci.* **16,** 635–44 (2011).
17. Cobb, J. N., DeClerck, G., Greenberg, A., Clark, R. & McCouch, S. Next-generation phenotyping: Requirements and strategies for enhancing our understanding of genotype-phenotype relationships and its relevance to crop improvement. *Theor. Appl. Genet.* **126,** 867–887 (2013).
18. Deery, D., Jimenez-Berni, J., Jones, H., Sirault, X. & Furbank, R. Proximal Remote Sensing Buggies and Potential Applications for Field-Based Phenotyping. *Agronomy* **4,** 349–379 (2014).
19. Eliceiri, K. *et al. Biological imaging software tools. Nature methods* **9,** (2012).
20. Sankaran, S. *et al.* Low-altitude, high-resolution aerial imaging systems for row and field crop phenotyping: A review. *Eur. J. Agron.* **70,** 112–123 (2015).
21. Duan, T. *et al.* Comparison of ground cover estimates from experiment plots in cotton, sorghum and sugarcane based on images and ortho-mosaics captured by UAV. *Funct. Plant Biol.* (2016). doi:10.1071/FP16123
22. Yang, C., Everitt, J. H., Du, Q., Luo, B. & Chanussot, J. Using high-resolution airborne and satellite imagery to assess crop growth and yield variability for precision agriculture. *Proc. IEEE* **101,** 582–592 (2013).
23. Zaman-Allah, M. *et al.* Unmanned aerial platform-based multi-spectral imaging for field phenotyping of maize. *Plant Methods* **11,** 35 (2015).
24. Lobell, D. B. The use of satellite data for crop yield gap analysis. *F. Crop. Res.* **143,** 56–64 (2013).
25. Pask, A., Pietragalla, J. & Mullan, D. *Physiological Breeding II: A Field Guide to Wheat Phenotyping. CIMMYT* (CIMMYT, 2012). doi:10.1017/CBO9781107415324.004
26. Barker, J. *et al.* Development of a field-based high-throughput mobile phenotyping platform. *Comput. Electron. Agric.* **122,** 74–85 (2016).
27. Andrade-Sanchez, P., Gore, M. & Heun, J. Development and evaluation of a field-based high-throughput phenotyping platform. *Funct. Plant Biol.* **41,** 68–79 (2014).
28. Vadez, V. *et al.* LeasyScan: A novel concept combining 3D imaging and lysimetry for high-throughput phenotyping of traits controlling plant water budget. *J. Exp. Bot.* **66,** 5581–5593 (2015).
29. Virlet, N., Sabermanesh, P., Sadeghitehran, K. & Hawkesford, M. J. Field Scanalyzer: An automated robotic field phenotyping platform for detailed crop monitoring. *Funct. Plant Biol.*

670 **44,** 143–153 (2016).

30. Chen, D. *et al.* Dissecting the Phenotypic Components of Crop Plant Growth and Drought Responses Based on High-Throughput Image Analysis. *Plant Cell Online* **26,** 4636–4655 (2014).

31. White, J. W. *et al.* Field-based phenomics for plant genetics research. *F. Crop. Res.* **133,** 101–112 (2012).

32. Kelly, D. *et al.* An opinion on imaging challenges in phenotyping field crops. *Mach. Vis. Appl.* **27,** 681–694 (2016).

33. Ribaut, J.-M., de Vicente, M. C. & Delannay, X. Molecular breeding in developing countries: challenges and perspectives. *Curr. Opin. Plant Biol.* **13,** 213–218 (2010).

34. Moose, S. P. & Mumm, R. H. Molecular plant breeding as the foundation for 21st century crop improvement. *Plant Physiol.* **147,** 969–977 (2008).

35. Gubbi, J., Buyya, R., Marusic, S. & Palaniswami, M. Internet of Things (IoT): A Vision, Architectural Elements, and Future Directions. *Futur. Gener. Comput. Syst.* **29,** 1645–1660 (2013).

36. the UK Government Chief Scientific Adviser. *The Internet of Things: making the most of the Second Digital Revolution.* (2014). doi:GS/14/1230

37. Watson, A. *et al.* Speed breeding: a powerful tool to accelerate crop research and breeding. 1–17 (2017). doi:dx.doi.org/10.1101/161182

38. Howse, J. *OpenCV Computer Vision with Python.* (Packt Publishing Ltd., 2013).

39. Pedregosa, F. *et al.* Scikit-learn: Machine Learning in Python. *J. Mach. Learn. Res.* **12,** 2825–2830 (2011).

40. van der Walt, S. *et al.* Scikit-image: image processing in Python. *PeerJ* **2,** 1–18 (2014).

41. Szeliski, R. *Computer Vision : Algorithms and Applications.* (Springer Science & Business Media, 2010). doi:10.1007/978-1-84882-935-0

42. Aloise, D., Deshpande, A., Hansen, P. & Popat, P. NP-hardness of Euclidean sum-of-squares clustering. *Mach. Learn.* **75,** 245–248 (2009).

43. Shi, J. & Malik, J. Normalized cuts and image segmentation. *IEEE Trans. Pattern Anal. Mach. Intell.* **22,** 888–905 (2000).

44. Bassiou, N. & Kotropoulos, C. Color image histogram equalization by absolute discounting back-off. *Comput. Vis. Image Underst.* **107,** 108–122 (2007).

45. Sezgin, M. & Sankur, B. Survey over image thresholding techniques and quantitative performance evaluation. *J. Electron. Imaging* **13,** 146–165 (2004).

46. Haralick, R., Shanmugan, K. & Dinstein, I. Textural features for image classification. *IEEE Transactions on Systems, Man and Cybernetics* **3,** 610–621 (1973).

47. Harris, C. & Stephens, M. A Combined Corner and Edge Detector. *Procedings Alvey Vis. Conf. 1988* 147–151 (1988). doi:10.5244/C.2.23

48. Zadocks, J. C., CHANG, T. T. & KONZAK, C. F. A decimal code for the growth stages of cereals. *Weed Res.* **14,** 415–421 (1974).

49. Kimmel, R. & Bruckstein, A. M. Regularized Laplacian zero crossings as optimal edge integrators. *Int. J. Comput. Vis.* **53,** 225–243 (2003).

50. Semenov, M. A. & Doblas-Reyes, F. J. Utility of dynamical seasonal forecasts in predicting crop yield. *Clim. Res.* **34,** 71–81 (2007).

51. Griffiths, S. *et al.* Meta-QTL analysis of the genetic control of ear emergence in elite European winter wheat germplasm. *Theor. Appl. Genet.* **119,** 383–95 (2009).

52. McMaster, G. S. & Wilhelm, W. W. Growing degree-days: One equation, two interpretations. *Agric. For. Meteorol.* **87,** 291–300 (1997).

53. Lancashire, P. D. *et al.* A uniform decimal code for growth stages of crops and weeds. *Ann. Appl. Biol.* **119,** 561–601 (1991).

54. Delude, C. M. Deep phenotyping: The details of disease. *Nature* **527,** S14–S15 (2015).

55. Fiorani, F. & Schurr, U. Future scenarios for plant phenotyping. *Annu. Rev. Plant Biol.* **64,** 267–91 (2013).

56. Heideman, M., Johnson, D. & Burrus, C. Gauss and the history of the fast fourier transform. *IEEE ASSP Mag.* **1,** 14–21 (1984).

57. Nakamichi, N., Kita, M., Ito, S., Yamashino, T. & Mizuno, T. PSEUDO-RESPONSE REGULATORS, PRR9, PRR7 and PRR5, Together play essential roles close to the circadian clock of Arabidopsis thaliana. *Plant Cell Physiol.* **46,** 686–698 (2005).

58. Chenu, K., Deihimfard, R. & Chapman, S. C. Large-scale characterization of drought pattern: A continent-wide modelling approach applied to the Australian wheatbelt - spatial and temporal trends. *New Phytol.* **198,** 801–820 (2013).

59.   Tardieu, F. Why work and discuss the basic principles of plant modelling 50 years after the
first plant models? *J. Exp. Bot.* **61,** 2039–2041 (2010).

60.   Turc, O., Bouteill, M., Fuad-Hassan, A., Welcker, C. & Tardieu, F. The growth of vegetative
and reproductive structures (leaves and silks) respond similarly to hydraulic cues in maize.
New Phytol. **212,** 377–388 (2016).

61.   Dawson, I. K. *et al.* Barley: a translational model for adaptation to climate change. *New Phytol.*
206, 913–931 (2015).

62.   Palosuo, T. *et al.* Simulation of winter wheat yield and its variability in different climates of
Europe: A comparison of eight crop growth models. *Eur. J. Agron.* **35,** 103–114 (2011).

63.   Hsu, C.-W. & Lin, C.-J. A comparison of methods for multiclass support vector machines.
IEEE Trans. Neural Networks **13,** 415–425 (2002).

64.   Shaw, L. M., Turner, A. S., Herry, L., Griffiths, S. & Laurie, D. A. Mutant alleles of
Photoperiod-1 in Wheat (Triticum aestivum L.) that confer a late flowering phenotype in long
days. *PLoS One* **8,** (2013).

65.   Shaw, L. M., Turner, A. S. & Laurie, D. A. The impact of photoperiod insensitive Ppd-1a
mutations on the photoperiod pathway across the three genomes of hexaploid wheat (Triticum
aestivum). *Plant J.* **71,** 71–84 (2012).

66.   R Development Core Team. *R : A Language and Environment for Statistical Computing.* **1,** (R
Foundation for Statistical Computing, 2008).

67.   Shannon, C. E. Editorial note on 'Communication in the presence of noise'. *Proc. IEEE* **72,**
1713–1713 (1984).

68.   Kohavi, R. A Study of Cross-Validation and Bootstrap for Accuracy Estimation and Model
Selection. *Int. Jt. Conf. Artif. Intell.* **14,** 1137–1143 (1995).

Online Methods

Five wheat NILs used in the field trial represent a range of genetic variation all with
the genetic background of the UK elite spring wheat '*Paragon*'. The development of
the Late-DTEM: Par (Norstar + Gamma 319c) 3c-11, *Ppd-1* loss of function (lof)
lines is described previously[64]. The development of the Early-DTEM NILs: Par
(GS100 2A+CS2B+Son64 2D)-T10 B10 -3b16 and *Ppd-D1a* photoperiod insensitive
has also been published[65]. The novel line Stay-Green is line 2316b selected on the
basis of stay green phenotype from a population of 7000 *Paragon* EMS mutants
carried through single seed descent up to M6 developed under the Wheat Genetic
Improvement Network of the UK Department of Food and Rural Affairs (Defra). The
semi-dwarf NILs (short) were produced by marker assisted backcrossing (to BC6)
using *Rht-B1* and *Rht-D1* KASP markers (LGC). which is available online from
http://www.cerealsdb.uk.net/cerealgenomics/CerealsDB. The sources of *Rht-D1b* and
Rht-B1b were the UK winter wheat varieties 'Alchemy' and 'Robigus' respectively.
The five wheat lines were sown in single 1 m^2 plots in autumn 2014 at Church Farm,
Norfolk UK, and grown according to standard agronomic practice. The manual score
of days to ear emergence (DTEM) was done when 50% of the plot showed 50%
emergence of the ear from the flag leaf. The manual measurement of plant height was
done from the ear tip to ground level.

The CropQuant hardware contains many components, of which the centre one is a
Raspberry Pi 2 or *Pi* 3 single-board computer (we have also used Intel® Edison in a
different version of CropQuant workstation). Based on a mobile ARM processor, the
Raspberry Pi computer features on-board external connections in the form of USB
and Ethernet to allow expansion using additional peripherals as well as an array of
digital GPIO (general purpose input and output) pins to interface with. The crop
growth image acquisition was performed using a 5MP RGB or NoIR (No InfraRed
filter) camera module connected via a CSI (Camera Serial Interface) port on the *Pi*
mother board. Digital temperature and humidity sensors are connected via
manufacturer supplied circuits to the GPIO pins of the *Pi* for interactive control. The
sensors themselves are mounted separate from the circuits, externally on the
CropQuant's housing, wired through the base of the device and sheltered by a smaller,
open housing unit. The external mounting allows for accurate sensing of ambient air
conditions while sheltering the electronics from direct water damage. The CropQuant
terminal is housed within a weatherproof (IP66 rated) plastic container, sealed around
all openings allowing operation in the field. Physical connection to the system for
data transfer via USB or Ethernet and power (12/5V DC) is facilitated by water-
resistant couplers designed to be sealed against the rain and air moisture.

The CropQuant software package runs on Linux-based operating system *Debian*. It
contains two servers, NetATalk and VNC sever, to facilitate infield data transfer and
remote systems control, which allows users to connect to each CQ terminal through a
wireless (using a tablet or a smartphone) or a wired connection (using a laptop). To
enable real-time systems interactions, a GUI-based imaging program has been
developed and added into the software package to control the RGB or NoIR camera
module for time-lapse crop monitoring. The program can automatically detect the IP
address of a given CropQuant terminal so that the terminal can be associated with its
specific experiment ID of the field trial. After that, the program requests users to

specify information such as genotype, biological replicates and imaging duration via a
GUI dialog box, where users can initiate the image acquisition. The program can
automatically adjust white balance, exposure mode and shutter speed in relation to
variable infield lighting conditions using the *picamera* package[*], a Python interface to
the *Raspberry Pi* camera hardware. Both image resolution and imaging frequency
(three times per hour in our field trials) can be changed if users want to modify their
experimental settings. The program also conducts the initial quality control and data
backup after each image is captured.
811
Besides the image acquisition, the software package contains a variety of functions
such as performing simple workstation and network diagnostics and synchronising
with the central server twice within an hour to upload sensor data and CropQuant
hardware information (see CropMonitor). Representative daily images are routinely
selected and transferred to the central server during the night, which provides a daily
snapshot of the monitored crops. Image data backups held on the SD card of the
device are routinely synchronised with the server to provide an external backup as
needed, with verification of multiple separate backups being performed by the process
before it removes the archived data to free storage space. Relying on the Linux
crontab scheduling system, we can monitor the performance of the software package
and resume it automatically in cases of software interruption or power disruption. The
SD card image running on the current version of CropQuant can be downloaded via
https://drive.google.com/drive/folders/0B17ZL8AzLo8wNFJUVS1lOFkzb3M?usp=s
haring. Source code is freely available for academic usage, which was arranged into
source trees and saved in both local and central repositories. We are also preparing an
online Github repository for the CropQuant project.
828
829   [*]https://picamera.readthedocs.io/en/release-1.13/
830
CropMonitor is an IoT-style control system developed to oversee the whole CQ
platform. It is operated through an onsite central server, logging updates received
from individual clients, i.e. CQ terminals. A Python application on each workstation
is running at regular intervals, scheduled by the native *Cron* Linux command line
utility. The application queries the terminal to determine workstation status
information such as uptime, network addresses and storage usage. Sensor data and
more variable system data such as CPU temperature and the usage of processor and
memory are sampled at a higher frequency and a median average of the readings is
recorded during the half-hourly query. Once the application has collected all
necessary data it is encoded into a JSON data object and transmitted over HTTP to the
central server which stores the data in an SQL database running on a HPC cluster.
CropQuant status is displayed and automatically updated using a web-based interface,
determining whether each node is online by the time of the most recent update. The
web interface provides information, including the location of each CropQuant
terminal in the field (a field map needs to be uploaded to the central server), graphs of
collected terminal and sensor data, and facilitates device configuration, SSH and VNC
linking to all active nodes. Nodes within the CropMonitor system are categorised into
groups and projects as defined by the user, allowing the organisation of workstations
and restriction of access to stored data. CropMonitor provides a centralised real-time
monitoring system to administer the network of infield workstations and collate
collected data for visualisation, batch processing and annotation.
852

The image selection algorithm is designed to perform speedy assessment of large image datasets captured in field trials by comparing images to a number of fixed criteria. The Python-based algorithm can be executed either on a normal computer or a HPC cluster. All images which meet the analysis standards will be collated. Over 200 GB data have been generated by ten offsite CropQuant terminals in the 2015 season during a 95-day period, with 50GB data were actually analysed after the selection procedure. In turn, an image is measured based on its brightness, sharpness, and shadow percentage, allowing all images which perform above a set of thresholds to be retained for further traits analysis. To determine the brightness of an image, the median value of pixel intensity is taken by transforming the image into HSV colour space. If the median intensity value is lower than a set threshold, the image is culled and not used from this point forward. The image clarity is determined by applying a Sobel edge detection[41] to the image. The detectable edges are calculated and then correlated with sharpness and exposure range of the image. The result of the clarity detection is also compared to a set threshold, which will disqualify images if they are out of focus or unclear with ill-defined edges. The final image test is of the percentage shadow within the visible area. Dark pixels found in an image with an illumination value of below 20% are either too dark for feature extraction or containing too much shadow in monitored plots. Once all rules have been passed, selected images are included in a result folder with a CSV file recording image metadata for further high-throughput image analysis.

The plot detection algorithm detects initial reference positions of monitored plots. The algorithm identifies the coordinates of white reference canes (the plot region) and dark height markers on a ranging pole, using an ensemble of colour-based feature selection on the basis of HSV (hue, saturation and value) and *Lab* non-linear colour space. It also classifies pixels into different groups, including sky, soil between plots, crop canopy, shadow, and plot regions using simple unsupervised machine-learning techniques such as k-means and spectral clustering. After detecting initial reference objects in the image, the algorithm establishes a pseudo 3D reference system that records the 2D coordinates of the plot area, the canopy region, and height markers through a range of feature selection approaches. The pixel-metric conversion is also computed based on height markers on the ranging pole.

The CropMeasurer algorithm employs an adaptive intensity and dynamic gamma equalisation to adjust colour and contrast to minimise colour distortion caused by diverse infield lighting. The algorithm tracks geometric differences between the plot on a given image and the initial position. If different, a geometric transformation method will be applied to recalibrate the image, which removes areas outside the plot area and could generate different sizes of black bars to the top of the given image. Within a plot, CropMeasurer tracks the crop height by detecting the visible part of the ranging pole and defines the canopy region through a combined adaptive thresholding and local Otsu threshold methods. Finally, the algorithm applies Harris and Shi-Tomasi corner detection methods[47] to locate corner-featured points within the canopy region. Red pseudo points are generated to represent the tips of erect leaves, reflective surfaces of curving leaves, heads and the corner points on ears. The main orientation of a given plot is quantified based on an optimised Canny edge detection method[49], which computes the alignment of crop stems.

902 **Data interpolation and analysis** have been used to handle minor data loss during the
903 field experiments. Four days' data gap (at the end of May 2015) has been recorded on
904 a number of offsite CropQuant workstations, which was caused by SD card crash due
905 to short-term battery failure. We used cubic spline interpolation method[66] to fill the
906 small gap in the phenotypic datasets.
907
RGR and FFT conversion the data is recorded on a daily basis, the maximum
frequency component visible is every two days (0.5 cycles-per-day) due to the
Nyquist-Shannon sampling theorem[67]. The cycles-per-day can be viewed in the same
manner as Hertz, which is known as cycles-per-second and indicates a measure of
frequency. We represent the frequency in cycles-per-day as the data were recorded at
daily intervals.
914
The growth stage predictive model is the basis of the GxExP model. The model is
produced to explore how to predict growth stages for different wheat genotypes in
relation to real-time performance traits and environment data. It employs support
vector machines (SVM), a popular machine learning technique for classification, with
radial basis function kernels to classify growth stages. The performance of the model
is tested by *Paragon* WT (G1) growth data from 2015 and 2016. For *Paragon* WT
(2015), the model is trained with all other 2015 genotype data, whereas for the 2016
Paragon WT datasets, the model is trained with all 2015 data, all models utilise K-
Fold^2 cross-validation for prediction. To simulate the real-world situation for the stage
prediction, we did *not* allow the model to obtain knowledge of the previous stages.
Hence, the model mainly modestly mistimes booting (GS41-49) and heading (GS51-
59) due to the short duration of both stages.
927
The prediction in comparison with the manually recorded growth stages suggests a
successful prediction of the timing and duration across all growth stages for both 2015
and 2016 datasets, except for the short transition period during booting (GS41-49),
where the duration of booting is two days short. Due to the limited data points for
booting across all genotypes used for training, the model cannot differentiate booting
from heading sufficiently. For this matter, we are planning to add training datasets
from other varieties such as Watkins and Chinese Spring wheat in other field trials.
The stage prediction is trained by *Paragon* WT growth data from both 2015 and 2016:
(1) to predict the 2015 *Paragon* WT, the model is trained with all other 2015 NIL
growth data; (2) whereas the 2016 *Paragon* WT was based on all 2015 growth data.
Through this approach, the model can rectify itself using previous years' training data.
After the training phase, the model utilises K-Fold[68] cross-validation for the growth
stage prediction.
941
The GxE interaction model explores the interactions between the recorded crop
growth of five wheat genotypes and a number of environmental factors. Correlations
are performed for each environmental factor grouped over three days with the
recorded growth data. The reason to group environmental factors into nested three-
day periods is to remove outliers and smooth the input data. The correlations are
determined for the first four growth stage for five genotypes. The analysis is
performed on the grouped data as particular stages (e.g. booting and heading) contain
few recorded growth data due to the short duration of both stages were present during
the growth. To determine the interactions between relative growth rates (RGR) and
environmental factors, we used the formula $(e^{RGR})^{-1}$ to convert negative correlation

values to positive counterparts, as the RGR series is a decreasing sequence in relation
to the increasing nature of growth stages.

Based on significant environmental factors, linear regression models[45] have been
explored and a single linear regression model is selected to estimate RGR of five
genotypes in relation to given infield environment conditions. Environmental factors
with insignificant correlations (where $p > 0.01$, with respect to the height over the
entire time-series) are removed from the analysis as they provide little predictive
power. Ordinary least squares are used to derive the model coefficients. The RGR
data is normalised to present percentage changes in height between two consecutive
days. To predict the canopy height for a given genotype, environment data at each
growth stage is input to the global model. To derive the height of the plant over time,
successive application $h_t = h_{t-l}(1 + y_t)$ is applied, where h_t is taken from the
above equation, h_{t-l} is the height of the plant at the previous time-point, and h_0 is
equal to the initial height.

The performance of the model is verified by estimating the growth of all five NILs,
including the overall paragon growth data (GT). The estimation is displayed with
respect to the true canopy height datasets. The mean squared error recorded for G2
(*genotype two*, Late-DTEM), G3 (*genotype three*, Early-DTEM) and G4 (*genotype*
four, Stay-Green) shows that the estimated height is close to the true growth curves.
However, the error is much larger for G1 (*genotype one*, *Paragon* WT) and G5
(*genotype five* Short). This is due to the majority of crop growth happens during the
early stages (GS32-GS59), estimation deviation during these initial stages could affect
the overall height results. As the global predictive model might not be sensitive
towards specific genotypes, we are still seeking a better approach to incorporate all
genotypes with a similar genetic background into the prediction. The stage predictions
are used in the linear regression growth model that could be run on a single-board
computer such as a *Pi* computer to give accurate quantifications. We have chosen to
establish the predictive growth model based on data produced from the CQ platform
and hence did not perform cross-validation tests to offer more rigorous evidence of
how well the model will generalise to new data. Given larger datasets containing
more biological replicates, conducting cross-validation produces more reliable growth
models with increased precision. Warning messages will be triggered via the CQ
platform, if the crop growth rate has deviated from the bounds of its estimated growth
region (±1SD).

High definition movies referred in this manuscript can be freely downloaded at
https://drive.google.com/drive/folders/0B17ZL8AzLo8wNFJUVS1lOFkzb3M?usp=s
haring (source code is freely available for academic usage and we are also preparing
an online Github repository for the CropQuant project).

Code availability. We used a *Jupyter* Notebook (i.e. the iPython Notebook) to
present and explain algorithms and software solutions associated with the CQ project.
They are freely available for academic use. Software packages running on CQs, high-
throughput trait analysis algorithms, and GxE modelling can be downloaded via
GoogleDocs for academic usage and an online Github repository is being prepared.
https://drive.google.com/drive/folders/0B17ZL8AzLo8wNFJUVS1lOFkzb3M?usp=s
haring
1001

1002 *Figure 1*
1003

1004
Figure 1. A high-level system architecture of the CropQuant platform.
(**a**) The hardware and software design for a CQ workstation, including a single-board
computer, climate sensors, a tailored circuit board to integrate sensors, an imaging
sensor, local USB storage, wired and wireless network components, *Debian* operating
system, and custom-made Python software package for crop images and climate data
collection. (**b**) Data communications between CQs and either a portable device in the
field or an onsite PC workstation. (**c**) The network setting that integrates CQ terminals,
infield wireless network, and the CropMonitor control system.
1013

1014   *Figure 2*
1015

1016
Figure 2. The CropQuant phenotyping platform used in field experiments.
(**a**) An all-in-one CQ workstation. (**b**) The internal hardware design of a version 2 CQ
workstation. (**c**) A custom-made circuit board that integrates low-cost climate sensors.
(**d**) Different versions of CQ built for dissimilar monitoring tasks. (**e**) CQs used in
offsite field experiments, powered by batteries and solar panels. (**f**) The CQ platform
established for onsite field experiments, powered by 5V/2A power supplies. (**g**) Real-
time crop inspection using either a portal device in the field or a PC in an office. (**h-j**)
the CropMonitor control system administers CQ terminals and records information
such as online or offline status, operational mode, daily representative crop image,
micro-environment for the plot region, and computational resource for each connected
CQ terminal. (**k**) An infield WIFI system installed for field trials. (**l**) A commercial
field weather station. (**m**) HPC clusters used for durable data storage and trait analysis.
1029
1030

1031 *Figure 3*

1032

1033
Figure 3. The high-throughput analysis pipeline established for batch processing
and measuring performance-related traits.
(**1**) *Step 1*, high-quality crop images were selected by the selection algorithm and
stored in both local and central repositories. (**2**) *Step 2*, initial reference positions of
monitored plots were detected by the referencing algorithm, which also calculated the
pixel-metric conversion rate. (**3**) *Steps 3* and *4*, the performance-related trait analysis
algorithm was designed to track plots of interest and conduct trait analyses to quantify
the canopy region and performance-related traits. (4) *Step 5*, traits such as dynamic
crop growth patterns in relation to thermal time (degree-day), vegetative greenness (0-
255), and the main plot orientation (0^o-180^o) were quantified and illustrated.
1044

1045 *Figure 4*

1046

1047
Figure 4. The performance of five wheat NILs monitored by the CQ platform to
identify dynamic developmental profiles.
Five wheat NILs (Late-DTEM, Early-DTEM; Short, Stay-Green, and *Paragon* WT)
and their dynamic performance in relation to environmental factors such as solar
radiation, rainfall, and temperature, during the 95-day monitoring period. Six growth
stages of *Paragon* WT were used as reference. Accumulated thermal time in degree-
day units was computed for comparison.
1055

1056   *Figure 5*

1057

1058
1059   **Figure 5. Recognising subtle and dynamic developmental variances for wheat**
1060   **genotypes under different climate patterns using the CQ platform.**
1061   (**a**) Different temperatures and accumulated degree-day patterns recorded in 2015 and
1062   2016. (**b**) Aligning and comparing growth curves of *Paragon* WT in 2015 and 2016
1063   within similar growth stages and the degree-day period (1250-2300 °Cd).
1064

1065 *Figure 6*
1066

1067
Figure 6. Extracting underlying growth patterns from continuous phenotypic
data using a fast Fourier transform (FFT).
(**a**) RGR (growth % of the previous day) is used to present the daily growth rate of all
the NILs at different growth stages. (**b**) The underlying growth patterns for *Paragon*
WT in 2015 and 2016 after the FFT conversion. (**c**) The underlying growth patterns
for Late-DTEM and Early-DTEM, after the FFT conversion. (**d**) Using a hypothetical
Paragon NILs growth data (merging all the NILs) to study the growth pattern.
1075

1076 *Figure 7*
1077

1078
Figure 7. Establishing a GxE predictive model to forecast the performance of
wheat genotypes under different climate patterns.
(**a**) Applying a correlation model to identify highly correlated environmental factors,
including thermal time, solar radiation, rainfall and growth stage duration using
Pearson correlation (p<0.01). (**b**) A growth stage-based predictive model applied to
estimate the key growth stages of *Paragon* WT in 2015 and 2016 compared with
manual scoring. (**c**) A global growth model comparing real growth curve measured by
CQ (red dotted line) with estimated growth curve (green line). (**d**) Warning messages
triggering mechanism to alert users if crop growth is outside the safe bounds (±1SD)
of the estimated growth region.
1089

1090 *Supplementary figure 1*
1091

1092
Supplementary figure 1. An all-in-one CQ device deployed in field experiments.
(**a**) Low-cost remote sensors (e.g. light levels, ambient temperature and humidity, soil
temperature and moisture) integrated by a tailored circuit board and then connected to
a *Raspberry Pi* computer via GPIO (general purpose input/output) pins. (**b**) A number
of all-in-one CQs being tested for establishing a mesh network in the Zhou laboratory.
(**c**) An all-in-one CQ device deployed in a wheat field experiment in 2017. (**d**) A soil-
based sensor installed to collect soil temperature and moisture for a six-metre wheat
plot. (**e**) Light levels and ambient temperature and humidity sensors mounted on the
top of the all-in-one CQ, together with an Ethernet coupler (black) and a micro-USB
coupler (blue).
1103

1104 *Supplementary figure 2*
1105

1106
1107
Supplementary figure 2. The hardware design of a version 2 CropQuant.
(**a**) *Side view*: (1) a *Raspberry Pi* 2, (2) digital sensor circuits, (3) a *Pi* camera module,
(4) a sensor GPIO connection, (5) an external sensor housing, (6) a digital sensor
connection, (7) an inline power fuse, (8) a voltage converter, (9) an external power
connection, (10) an external Ethernet connection coupler. (**b**) *Front view*: (1) a
camera module, (2) an external camera UV lens, (3) a camera sunlight shield, (4) an
external sensor housing. (**c**) *Base view*: (1) a digital humidity sensor, (2) a digital
temperature sensor, and (3) an external power connection.
1116

1117   *Supplementary figure 3*
1118

1119
1120   **Supplementary figure 3. Real-time infield crop monitoring via portable devices.**
1121   (**a**) A crop scientist using an Android tablet to connect to infield CQ terminals to
1122   examine the performance of wheat growth. (**b&c**) After connecting to the VNC server
1123   running on a CQ terminal, different experiments can be inspected and managed by
1124   crop scientists via a VNC viewer on the portable device, e.g. smartphones or tablets.
1125

1126 *Supplementary figure 4*
1127

1128
Supplementary figure 4. The CropMonitor control system.
(**a**) Grid view of the CropMonitor system, which provides a regular overview of all
CQ terminals in the field, including online (green) or offline (red) status, operational
modes (amber means the imaging is either finished or halted), and the duration of
crop monitoring. An experimental layout of monitored plots is also provided, showing
the location of all CQ terminals and their operational modes. (**b**) List view of the
system showing CQ's online duration, network addresses, computing storage, and
SSH/VNC tools to access CQ terminals directly from the CropMonitor control system.
(**c**) Individual view of the system which illustrates an individual CQ workstation,
containing climate sensor data and systems information during the monitoring period.
1139

1140 *Supplementary figure 5*
1141

1142
1143 **Supplementary figure 5. A real-time stream function activated for deploying CQ**
1144 **workstations in the field.**
1145 (**a-b**) A live stream function showing the location of a CQ terminal in the field as well
1146 as assisting device deployment and systems calibration via the CropMonitor system
1147 running on each CQ device.
1148

1149   *Supplementary figure 6*

1150

1151

1152   **Supplementary figure 6. Outdoor and indoor wheat experiments (e.g. wheat**

1153   **speed breeding) monitored by the CQ platform in 2017.**

1154

1155 *Supplementary figure 7*
1156

1157
1158 **Supplementary figure 7. The network topology of the mesh network system**
1159 **established by many all-in-one CQs jointly operating in the field.**
1160

1161 *Supplementary figure 8*
1162

1163
1164 **Supplementary figure 8. The systems architecture diagram of how to network**
1165 **CropQuant terminals, the CropMonitor system, and a HPC cluster.**
1166

1167 *Supplementary figure 9*
1168

1169
1170
1171 **Supplementary figure 9. The underlying growth patterns for Green Mutant NIL**
1172 **and Short-Stem NIL extracted through a fast Fourier transform (FFT) approach.**
1173

1174 ***Supplementary figure 10***

1175

1176

Supplementary figure 10. A predictive model to forecast the timing and duration of key growth stages for genotypes with *Paragon* background.

(**a-e**) The model predicts the growth stages based on all *Paragon* growth data (acquired in 2015 and 2016) and environmental factors selected by the GxE correlation model. (**f**) The model forecasts the growth stages based on overall *Paragon* data (the hypothetical growth dataset) and environmental factors.

1183

1184 *Supplementary figure 11*

1185

1186

Supplementary figure 11. A predictive model to forecast the dynamic growth
patterns of wheat genotypes with *Paragon* background in field conditions.
(**a-b**) The growth prediction based on Late-DTEM NIL and Early-DTEM NIL growth
data. Vertical dash lines indicate the manually segmented growth stages, from Stem
elongation or jointing (GS32-39) to Ripening (GS91-95). The red dotted lines stand
for real CQ measurements. If outside the safe bounds of the growth estimates, a
warning message (a triangle coloured red) will be triggered on the CropMonitor
control system. For example, the Late-DTEM line was growing too slow. (**c**) The
growth prediction based on Stay-Green NIL growth data. (**d**) The height prediction
based Short-Stem NIL growth data, which was growing too fast.
1197

1198 *Supplementary figure 12*

1199

1200
1201

1202 **Supplementary figure 12. The circuit for the all-in-one CQ workstation.**

1203

1204 **Supplementary figure 13**
1205

1206
Supplementary figure 13. Using deep-learning neural network architectures to
train algorithms to quantify yield component traits based on images acquired by
the CQ platform.
(1) *Step 1*, pre-processing and calibrate image series captured by the CQ platform in
the field. **(2)** *Step 2*, based on texture and pattern classification methods such as grey
level co-occurrence matrices (GLCM) and texture entropy, detecting spike-featured
regions, which are divided into sub-images as training data for a CNN (Convolutional
neural network) classifier. **(3)** *Step 3*, to count spikes per unit area and spike numbers
on a given image, further smaller sub-images are produced to fit a whole spikelet
region. CNN is trained to count spikelet-only sub-images. **(4)** *Step 4*, the machine-
learning based estimation are correlated with manual measurements, so that the
estimation model can be improved.
1219