# The evolutionary history of 2,658 cancers

Moritz Gerstung[1,2,#,*], Clemency Jolly[3,#], Ignaty Leshchiner[4,#], Stefan C. Dentro[2,3,5,#], Santiago Gonzalez[1,#], Daniel Rosebrock[4], Thomas J. Mitchell[2,6], Yulia Rubanova[7,8], Pavana Anur[9], Kaixian Yu[10], Maxime Tarabichi[3], Amit Deshwar[7,8], Jeff Wintersinger[7,8], Kortine Kleinheinz[11,12], Ignacio Vázquez-García[2,6], Kerstin Haase[3], Lara Jerman[1,13], Subhajit Sengupta[14], Geoff Macintyre[15], Salem Malikic[16,17], Nilgun Donmez[16,17], Dimitri G. Livitz[4], Marek Cmero[18,19], Jonas Demeulemeester[3,20], Steven Schumacher[4], Yu Fan[10], Xiaotong Yao[21,22], Juhee Lee[23], Matthias Schlesner[11], Paul C. Boutros[7,24], David D. Bowtell[25], Hongtu Zhu[10], Gad Getz[4,26,27,28], Marcin Imielinski[21,22], Rameen Beroukhim[4,29], S. Cenk Sahinalp[17,30], Yuan Ji[14,31], Martin Peifer[32], Florian Markowetz[15], Ville Mustonen[33], Ke Yuan[15,34], Wenyi Wang[10], Quaid D. Morris[7,8], Paul T. Spellman[9,#], David C. Wedge[5,35,#], Peter Van Loo[3,20,#,*], on behalf of the PCAWG Evolution and Heterogeneity Working Group[36] and the PCAWG network.

[1]European Molecular Biology Laboratory, European Bioinformatics Institute (EMBL-EBI), Cambridge CB10 1SD, United Kingdom; [2]Wellcome Trust Sanger Institute, Cambridge CB10 1SA, United Kingdom; [3]The Francis Crick Institute, London NW1 1AT, United Kingdom; [4]Broad Institute of MIT and Harvard, Cambridge, MA 02142, USA; [5]Big Data Institute, University of Oxford, Oxford OX3 7LF, United Kingdom; [6]University of Cambridge, Cambridge CB2 0QQ, United Kingdom; [7]University of Toronto, Toronto, ON M5S 3E1, Canada; [8]Vector Institute, Toronto, ON M5G 1L7, Canada; [9]Molecular and Medical Genetics, Oregon Health & Science University, Portland, OR 97231, USA; [10]The University of Texas MD Anderson Cancer Center, Houston, TX 77030, USA; [11]German Cancer Research Center (DKFZ), 69120 Heidelberg, Germany; [12]Heidelberg University, 69120 Heidelberg, Germany; [13]University of Ljubljana, 1000 Ljubljana, Slovenia; [14]NorthShore University HealthSystem, Evanston, IL 60201, USA; [15]Cancer Research UK Cambridge Institute, University of Cambridge, Cambridge CB2 0RE, United Kingdom; [16]Simon Fraser University, Burnaby, BC V5A 1S6, Canada; [17]Vancouver Prostate Centre, Vancouver, BC V6H 3Z6, Canada; [18]University of Melbourne, Melbourne, VIC 3010,

Australia; [19]Walter + Eliza Hall Institute, Melbourne, VIC 3000, Australia; [20]University of Leuven, B-3000 Leuven, Belgium; [21]Weill Cornell Medicine, New York, NY 10065, USA; [22]New York Genome Center, New York, NY 10013, USA; [23]University of California Santa Cruz, Santa Cruz, CA 95064, USA; [24]Ontario Institute for Cancer Research, Toronto, ON M5G 0A3, Canada; [25]Peter MacCallum Cancer Centre, Melbourne, VIC 3000, Australia; [26]Massachusetts General Hospital Center for Cancer Research, Charlestown, MA 02129, USA; [27]Massachusetts General Hospital, Department of Pathology, Boston, MA 02114, USA; [28]Harvard Medical School, Boston, MA 02215, USA; [29]Dana-Farber Cancer Institute, Boston, MA 02215, USA; [30]Indiana University, Bloomington, IN 47405, USA; [31]The University of Chicago, Chicago, IL 60637, USA; [32]University of Cologne, 50931 Cologne, Germany; [33]University of Helsinki, 00014 Helsinki, Finland; [34]University of Glasgow, Glasgow G12 8RZ, United Kingdom; [35]Oxford NIHR Biomedical Research Centre, Oxford OX4 2PG, United Kingdom.

[#]These authors contributed equally.

[*]To whom correspondence may be addressed:

Moritz Gerstung, European Molecular Biology Laboratory, European Bioinformatics Institute (EMBL-EBI), Hinxton, Cambridgeshire, CB10 1SD, United Kingdom. Tel: +44 (0) 1223 49 4636, email: Moritz.Gerstung@ebi.ac.uk.

Peter Van Loo, The Francis Crick Institute, 1 Midland Road, London, NW1 1AT, United Kingdom. Tel: +44 (0) 20 3796 1719, e-mail: Peter.VanLoo@crick.ac.uk.

[36]A list of members of the PCAWG Evolution and Heterogeneity Working Group can be found at the end of the manuscript.

**Summary**

Cancer develops through a process of somatic evolution. Here, we use whole-genome

sequencing of 2,778 tumour samples from 2,658 donors to reconstruct the life history,

evolution of mutational processes, and driver mutation sequences of 39 cancer types.

The early phases of oncogenesis are driven by point mutations in a small set of driver

genes, often including biallelic inactivation of tumour suppressors. Early oncogenesis

is also characterised by specific copy number gains, such as trisomy 7 in glioblastoma

or isochromosome 17q in medulloblastoma. By contrast, increased genomic

instability, a nearly four-fold diversification of driver genes, and an acceleration of

point mutation processes are features of later stages. Copy-number alterations often

occur in mitotic crises leading to simultaneous gains of multiple chromosomal

segments. Timing analysis suggests that driver mutations often precede diagnosis by

many years, and in some cases decades, providing a window of opportunity for early

cancer detection.

## Introduction

Akin to the evolution of species, the $\sim 10^{15}$ cells in the human body are subject to the forces of mutation and selection[1]. This process of somatic evolution begins before birth in the zygote and only comes to rest at death, as each cell division typically introduces between 1-10 mutations[2]. While these mutations are predominantly selectively neutral passenger mutations without transformative capacity, some are proliferatively advantageous driver mutations leading to cancer[3,4]. While the types of mutations in cancer genomes are well studied, little is known about the times when these lesions arise during somatic evolution and where the boundary between normal evolution and cancer progression should be drawn.

Sequencing of bulk tumour samples enables partial reconstruction of the evolutionary history of individual tumours, based on the catalogue of somatic mutations they have accumulated[5]. The timing of chromosomal gains during early somatic evolution and tumour progression can be estimated by counting point mutations within duplicated regions in single cancer samples[6-8]. Moreover, many studies have focused on the late stages of cancer evolution by reconstructing phylogenetic relationships between tumour samples and metastases from individual patients[9-12]. In addition, the typical ordering of driver mutations within a tumour type can be determined by aggregating pairwise timing estimates of genomic changes (for example clonal *vs*. subclonal) across many samples using preference models[13,14].

Here, we extend and apply these methods to the Pan-Cancer Analysis of Whole Genomes (PCAWG)[15] dataset, as part of the International Cancer Genome Consortium (ICGC)[16] and The Cancer Genome Atlas (TCGA)[17] to characterise the evolutionary history of 2,778 cancers from 2,658 unique donors across 39 cancer types (**Methods** and **Supplementary Information**). We determine the timing and order of mutations during cancer development to delineate patterns of chromosomal evolution within and across cancer types. We then define broad periods of tumour evolution and examine how drivers and mutational signatures vary between these stages. Finally, using clock-like mutational processes, we map mutation timing estimates into approximate real time, and create typical timelines of tumour evolution.

## Results

**Reconstructing the life history of single tumours**

A cancer cell's genome is shaped by the cumulative somatic aberrations that have arisen during its evolutionary past, and part of this history can be reconstructed from whole genome sequencing data[5] (**Fig. 1a**). Initially, each point mutation occurs on a single chromosome in a single cell, which gives rise to a lineage of cells all bearing the same mutation. If that chromosomal locus is subsequently duplicated in this lineage, and the point mutation is on the allele that is gained, the mutation will also be duplicated, and both mutated copies can be detected in sequencing data. Mutations on other chromosomes, or on the non-duplicated allele of a gained region, will be present only as a single copy. Mutations found in a subset of tumour cells have not swept through the population and must have occurred after the most recent common ancestor (MRCA) of the tumour cells in the sequenced sample. Based on these principles, we are able to approximately time individual point mutations relative to other genomic events such as large-scale chromosomal gains, and the emergence of the MRCA.

Mapping point mutations to the proportion of cells and chromosomes enables us to define broad categories, each associated with broad epochs of tumour evolution (**Fig. 1a**). *Clonal* mutations have occurred before the emergence of the MRCA and are common to all cancer cells, whilst *subclonal* mutations are only observed in a fraction of cancer cells. In regions of clonal copy number gains, clonal mutations can be further classified as *early clonal* or *late clonal. Early clonal mutations* lie within a duplicated locus and are themselves doubled, *i.e.* they occurred prior to the copy number gain. *Late clonal mutations* also reside at a gained chromosomal region, but only as one copy, *i.e.* they happened after the copy number gain. Inevitably, there will be some crossover between these categories of clonal mutations, both within and between samples, as their timing depends on the timing of the surrounding copy number gains, which may take place at different times during tumour evolution. However, the broad comparison of mutations before and after copy number gains does give additional insight into the dynamics of clonal evolution.

Importantly, the relative frequency of early and late clonal mutations provides

information about the timing of the underlying copy number segment. The ratio of

doubled to non-doubled mutations within a gained region can be used to calculate a

point estimate for when the gain happened during clonal evolution, referred to here as

*mutational timing*. For example, there would be few, if any, co-amplified early clonal

mutations if the gain had occurred right after fertilisation, whilst a gain that happened

towards the end of clonal tumour evolution would contain many duplicated

mutations[6] (**Fig. 1a** and **Online Methods**). Later on, we will show how this

mutational timing relates to real time for certain evolutionary events.

These analyses are illustrated in **Fig. 1b**. As expected, the variant allele frequency

(VAF) of somatic point mutations cluster around the values imposed by the purity of

the sample, local copy number configuration and identified subclonal population. The

depicted clear cell renal cell carcinoma has gained chromosome arm 5q at an early

time as part of an unbalanced translocation of t(3p;5q), confirming the notion that this

lesion often occurs in adolescence in this cancer type[8]. At a later time point the

sample underwent a whole genome duplication (WGD), duplicating all alleles

including the derivative chromosome in a single event, as evidenced by the mutation

time estimates of all copy number gains clustering around a single time point,

independently of the exact copy number state.

**Timing patterns of copy number gains**

To systematically explore the timing of copy number gains across cancer types, we

applied mutational timing analysis to 2,111 samples from the PCAWG dataset that

had copy number gains suitable for timing (see **Supplementary Information**). We

find that chromosomal gains typically occur after more than half of the point

mutations have been acquired (median value 0.76, inter-quartile range IQR [0.43,

0.94], with systematic differences between tumour types (**Fig. 2a**, **Extended Data**

**Fig. 1**). In glioblastoma, medulloblastoma and pancreatic neuroendocrine cancers, a

substantial fraction of gains occur early in mutational time. Conversely, in squamous

cell lung cancers and melanomas, gains arise towards the end of the mutational time

scale. Most tumour types, including breast, ovarian and colorectal cancers, show

relatively broad periods of chromosomal instability indicating a very variable timing

of gains across samples.

There are, however, certain tumour types with consistently early or late gains of
specific chromosomal regions. Most pronounced is glioblastoma, where 90% of
tumours harbour single copy gains of chromosome 7, 19 or 20 (**Fig. 2a-b**). Strikingly,
these gains are consistently timed within the first 10% of clonal mutational time,
suggesting they arise very early in a patient's lifetime. In the case of trisomy 7, this is
further supported by reports of somatic mosaicism in normal brain cells, potentially
suggesting an embryonic origin[18]. Similarly, the duplications leading to
isochromosome 17q in medulloblastoma are timed exceptionally early. Although less
pronounced, gains of chromosome 18 in B-cell non-Hodgkin lymphoma, as well as
gains of the q arm of chromosome 5 in clear cell renal cell carcinoma, often have a
distinctively early timing within the first 50% of mutation time[8]. In papillary renal
cell carcinoma, trisomy 7 is highly recurrent and occurs within the last 10% of clonal
mutational time.

Intriguingly, we observed that gains in the same tumour often appear to occur at a
similar time, pointing towards punctuated bursts of copy number gains involving the
majority of gained segments (**Fig. 2c**). While this is expected in tumours with WGD
(**Fig. 1b**), it may seem surprising to observe synchronous gains (defined as more than
80% of gained DNA in a single event) in near-diploid tumours. Still, synchronous
gains are frequent, occurring in a striking 57% (468/815) of informative near-diploid
tumours, 61% more frequently than expected by chance ($p < 0.01$, permutation test;
**Fig. 2d**). Only 6% of co-amplified chromosomal segments were linked by a direct
inter-chromosomal structural variant, indicating that the gains arise without being
necessarily physically linked. Only 10-20% of arm-level gains increment the allele-
specific copy number by more than one (**Fig. 2e**), suggesting that most of these gains
arise through mis-segregation of single copies during anaphase. This notion is further
supported by the observation that in about 85% of segments with two gained alleles,
the second gain appears with noticeable latency after the first (**Fig. 2f**). Combined, a
pattern emerges in which arm-level gains tend to arise in distinct mitotic crises that
lead to numerical and sometimes structural changes of multiple chromosomes, adding
(and possibly losing) one allelic copy at a time, in line with similar observations of
punctuated evolution in breast cancer[19].

## Timing of mutations in driver genes

As outlined above, point mutations (SNVs and indels) can be qualitatively assigned to different epochs, allowing the timing of driver mutations (**Fig. 1a, 3a**). Using a panel of 453 cancer driver genes[20], we find that pathogenic mutations in the 50 most common drivers are predominantly clonal, and often early clonal (**Fig. 3a-b**). For example, driver mutations in *TP53* and *KRAS* are 12x and 8x enriched in early clonal stages, respectively. For *TP53*, this trend is independent of tumour type (**Fig. 3c**). Mutations in *PIK3CA* are 2x more frequently clonal than subclonal, while non-coding changes near the *TERT* gene are 3x more frequently early clonal than expected.

Overall, common driver mutations predominantly occur during the early clonal period of tumour evolution. To understand how the entire landscape of all 453 driver genes changes over time, we calculated how the number of driver mutations relates to the number of driver genes in each of the evolutionary stages. This reveals an increasing diversity of driver genes mutated at later stages of tumour development: 50% of all early clonal driver mutations occur in only 9 genes, whereas the corresponding proportion of late and subclonal mutations occur in approximately 35 different genes each, a nearly 4-fold increase (**Fig. 3d**). These results are consistent with previous findings in non-small-cell lung cancers[21], and suggest that, across cancer types, the very early carcinogenic events occur in a constrained set of common drivers, while a more diverse array of drivers is involved in late tumour development.

## Relative timing of somatic driver events

Next, we sought to better understand the sequence and timing of events during tumour evolution by integrating the timing of driver point mutations and recurrent copy number changes across cancer samples. We calculated an overall probabilistic ranking of lesions (**Fig. 4a**), detailing whether each mutation occurs preferentially early or late during tumour evolution, by aggregating single patient partial order trajectories (**Fig. 4b-d**) within each cancer type (**Supplementary Information**, section 3.2, **Extended Data Fig. 2-4**).

We have confirmed through simulations that the average order of events is correctly estimated through aggregating individual patient trajectories (**Extended Data Fig. 3**),

which can be of varying level of detail depending on the sample (**Fig. 4a**,
**Supplementary Information,** section 3.2). As seen from example patient tumours
(**Fig. 4b-d**), whole genome sequencing provides insights on the actual order of driver
mutations, even on the single-sample level (*e.g.*, *APC* and *KRAS* mutations occur
before events in *TP53* in colorectal tumour SA87265; **Fig. 4b**). Through aggregation
of many such partial order relationships, we can infer probabilistic rankings of events
within each cancer type (**Fig. 4e-g**, **Extended Data Fig. 1**).

In colorectal adenocarcinoma, for example, we find *APC* mutations to have the
highest odds of occurring early, followed by *KRAS,* loss of 17p and *TP53,* and
*SMAD4* (**Fig. 4e**). Whole-genome duplications have an intermediate ranking,
indicating a variable timing, while many chromosomal gains and losses are typically
late. These results are in agreement with the classical progression *APC-KRAS-TP53*
model of Vogelstein and Fearon[22], but add considerable detail.

In other cancer types, the sequence of events in cancer progression has not previously
been studied in as much detail. For example, in pancreatic neuroendocrine cancers,
we find that many chromosomal losses, including those of chromosomes 2, 6, 11 and
16, occur early, followed by driver mutations in *MEN1* and *DAXX* (**Fig. 4f**). WGD
events occur late, after many of these tumours have reached a pseudo-haploid state
due to wide-spread chromosomal losses. In glioblastoma, we find that loss of
chromosome 10 and driver mutations in *TP53* and *EGFR* are very early, often
preceding early gains of chromosomes 7, 19 and 20 (as described above) (**Fig. 4g**).
*TERT* promoter mutations tend to occur at early to intermediate time points, while
other driver mutations and copy number changes tend to be later events.

Across cancer types, we typically find *TP53* mutations early, as well as losses of
chromosome 17 (**Extended Data Fig. 1**). WGD events usually have an intermediate
ranking and the majority of copy number changes occur after WGD. We also find that
losses typically precede gains, and consistent with the results above, common drivers
typically occur before rare drivers.

## Timing of mutational signatures

The cancer genome is shaped by various mutational processes over its lifetime, stemming from exogenous and cell-intrinsic DNA damage and error-prone DNA replication, each leaving behind characteristic mutational spectra, termed mutational signatures[23,24]. To quantify the changing activity of mutational processes operating in each cancer genome, we assessed differences of the observed mutational spectra between early clonal, late clonal and subclonal epochs (**Supplementary Information**). As shown in **Fig. 5a**, there are indeed examples of striking changes between early and late clonal single base substitution spectra, indicating that the underlying mutational processes have considerably changed during the clonal life history of these cancers. Overall, we find evidence for a changing mutational spectrum between early and late clonal time points in 29% (530/1,852) of informative samples ($p < 0.05$, Bonferroni-adjusted likelihood-ratio test), typically changing 19% of mutations (median absolute difference; range 4%–66%).

To quantify whether the observed temporal changes can be attributed to known mutational processes, we decomposed the mutational spectra at each timepoint into a catalogue of 57 mutational signatures, tailored to this cohort and fully described in Alexandrov *et al.*[24] (**Fig. 5a; Methods**). To this end, we were also able to make use of two new categories of signatures: a series of doublet base substitution signatures, and a series of indel signatures[24].

The fold-changes of mutations attributed to a given signature between early and late clonal stages of tumour evolution indeed vary by several orders of magnitude, revealing dynamic evolution of many mutational processes during the life history of many, but not all tumours (**Fig. 5b**).

In addition to this predominantly undirected temporal variability, a number of signatures demonstrate directional trends in their activity over time (**Fig. 5c**). As one may expect, we observe signatures of exogeneous mutagens to be predominant in the early clonal stages of tumourigenesis. These include tobacco smoking (single base signature SBS4), median fold change 0.54, IQR [0.34, 0.8]), which is most pronounced in lung adenocarcinoma (median change 0.32; IQR [0.2,0.37]; **Extended Data Fig. 5**) and consistent with previous reports[25,26]. Quite similarly, the signature of

UV light in melanoma contributes far fewer mutations at late clonal stages (SBS7;
mean fold change 0.17, IQR [0.09, 0.41]). Another strong decrease over time, similar
to that of tobacco smoke and UV damage, is found for a signature of unknown
aetiology, SBS12, which acts mostly in liver cancers (median fold change 0.31, IQR
[0.09, 0.49]).

Some mutational processes are preferentially active in late stages of clonal evolution.
For example, we see that APOBEC mutagenesis (SBS2 and SBS13) increases in
many cancer types from the early to late clonal stages (median fold change 2, IQR
[0.83, 3.8]), as does a newly described signature attributed to indirect damage from
UV light (SBS38; median fold 3.2, IQR [2.0,10])[24]. Lastly it is interesting to note that
chemo-resistant ovarian cancers display an increase in the mutational signature
associated with platinum-based therapy (double base signature DBS5; median fold
change 1.8, IQR [1.0, 3.3]).

**Chronological time estimates of whole genome duplications and subclonal**
**diversification**

Changes in the mutation rate of cancers influence timing estimates based upon
mutational data. Due to increased proliferation and in some cases acquired
hypermutation, one would generally expect an increase in the mutation rate over time,
yet some mutational processes appear more variable than others.

CpG>TpG mutations as the result of spontaneous deamination of methylated
cytosines (SBS1), constitute a promising candidate for a clock-like mutational
process[27], as they are ubiquitously found in all cancer and normal tissues[28,29]. The
number of CpG>TpG mutations increase constantly with age in this cohort (**Fig. 6a**),
with moderate variation between samples and across tissues (**Extended Data Fig.**
**6a**). As long as the rate of the mutational clock is constant within a sample, any inter-
sample variation can be accounted for, since the age at diagnosis is used to calibrate
the mutation clock in each tumour.

However, there may also be an increase in the CpG>TpG rate over time, which can be
estimated in two ways. First, sequencing data of matched primary and relapse samples
from the same donor suggest that there is an increased rate of mutation accumulation

between diagnosis and relapse (as evidenced here for publicly available data from ovarian and breast cancer[30,31]), **Fig. 6b**, and therefore that this acceleration likely extends to periods immediately prior to diagnosis. Formalising this argument for eight cancer types[30-33] yields a typical acceleration of around 5x [range: 1x-12x] post-diagnosis relative to pre-diagnosis (**Fig. 6c**). A second line of evidence comes from a systematic analysis of the relation between CpG>TpG mutation burden and age at diagnosis[34] in this cohort (**Extended Data Fig. 6a**). This indicates that, across cancer types, typically 20-40% of mutations are found independent of age and are thereby likely caused by an acceleration during cancer progression (**Extended Data Fig. 6b-c**). This constant offset is broadly compatible with a 5x acceleration over a period of about five years prior to a diagnosis at a typical age of 60 years.

One effect of an increasing mutation rate is that chronological time estimates appear closer to diagnosis and have higher variance, as the exact onset of acceleration is unknown (**Fig. 6d**). Applying this logic to the timing of whole-genome duplications based on CpG>TpG mutations, we find that the typical timing of WGD is about one decade before diagnosis (**Fig. 6e**). However, there is substantial variability among samples of a given tumour type, with many cases dating back more than two decades. This latency corresponds to rather different patterns in the lifetime of individuals (**Fig. 6f**): Ovarian adenocarcinomas have WGD occurring more than two decades before diagnosis in almost half of patients. For these early cases, the estimated age of WGD falls well into female reproductive life, concordant with the observation that the number of ovulations is a risk factor for ovarian cancer[35]. Conversely, for colorectal adenocarcinoma, most duplications shortly precede diagnosis. Finally, in the childhood cancer medulloblastoma, the majority of WGD events fall into the first two years of life, suggesting an early developmental onset of the disease.

Reassuringly, the earliness of WGDs is caused by depletion of early (that is co-amplified) mutations and is independent of the overall mutation burden (**Extended Data Fig. 6d-f**), making it unlikely that the observed early duplications are artefacts of acquired hypermutation. While the observation that the majority of CpG>TpG mutations occur proportionally to age places some bounds on the strength of acceleration, some uncertainty persists. We thus note that, without any acceleration, the estimated median latency between WGD and diagnosis would be 6-35 years for

the majority of cancer types analysed in this study; conversely at an acceleration of
10x the median latency would be 1.6-12 years (range for tumour types with ≥10
samples; **Fig. 6g**).

We used a similar approach to calculate the duration of subclonal diversification, *i.e.*
the time between the MRCA and the last detectable subclone. The typical timing is
considerably shorter and about 1yr, but interestingly, there are also cases where this
period is estimated to last more than ten years (**Fig. 6h**). We note, however, that
timing of the subclonal diversification period is more difficult and the results
presented here are lower bounds. This is because there may be earlier subclonal
branches missed in a single sample, there is limited power to detect very late and
deeply subclonal branches and also because the underlying phylogenetic relationships
between subclones cannot always be unambiguously resolved and were treated as
densely branching. Assuming linear instead of branching phylogenies approximately
doubles the inferred latency (**Extended Data Fig. 6g**).

While the exact timing of individual samples remains challenging due to low
CpG>TpG mutation numbers and unknown mutation rates for individual tumours,
these findings are entirely consistent with epidemiological observations: cancer
generally arises past the age of 50 (Ref. [36]), and the typical latency between
carcinogen exposure and cancer detection, e.g. in tobacco-associated cancers, is
several years to multiple decades[37]. Furthermore, the progression of most known
precancerous lesions to carcinomas usually spans multiple years, if not decades[38-45].
Our data corroborate these time scales and show that they hold in cases without
detectable premalignant conditions, raising hopes that these tumours could also be
detected in precancerous stages.

## Discussion

Taken together, the data presented here enable us to draw semi-quantitative timelines summarising the typical evolutionary history of each cancer type (**Fig. 7**). These make use of the qualitative timing of point mutations and copy number drivers, as well as signature activities, which can be interleaved with the chronological estimates of WGD and MRCA (**Fig. 7**, **Extended Data Fig. 1** for a complete list of all cancer types).

Individual tumour types show characteristic evolutionary trajectories, reflecting differences in the mutational processes driving evolution, the portfolio of driver mutations and their timing. While glioblastoma, for example, is often initiated by mutations in *EGFR* and gain of chromosome 7, *APC* and *KRAS* mutations are typically the earliest events in colorectal adenocarcinoma, followed by *TP53*, in line with classic findings[22]. This is similar to reports of frequent *KRAS* and *TP53* mutations and deletions of 17p in precursors to pancreatic adenocarcinoma[46], confirming the early role found here. *TP53* with accompanying 17p deletion is one of the most frequent initiating mutations in a variety of cancers, including ovarian cancer, where it is the hallmark of its precancerous precursor lesion[47]. CpG>TpG mutations (SBS1) are active throughout tumour development, however, their relative contribution decreases in favour of more diverse mutational processes acting at later stages of cancer development. Similarly, genomic instability, leading to frequent gains and losses, is typically seen late in cancer progression. Unlike most other cancers, high grade serous ovarian adenocarcinomas typically acquire chromosomal gains within the first half of clonal evolution (**Fig. 7d**). Our findings are consistent with these tumours having a high frequency of *TP53* and homologous recombination repair defects[48], and rendering them among the most genomically unstable of all solid cancers[49].

Across cancer types, our findings provide insight into how selection acts during cancer development, with an increased repertoire of driver mutations in late tumour evolution. Genetic canalization in early stages suggests that epistasis of fitness effects is constraining evolution to select for only a small set of mutations in genes and chromosomal aberrations that are able to initiate neoplastic transformation. Over time,

as tumours evolve, the small- and large-scale somatic changes they subsequently

accumulate propel them towards increasingly diverse paths driven by individually

rare, atypical driver mutations.

Our analysis reveals that tumour evolution is driven by crises leading to simultaneous

gains of multiple or even all chromosomal segments, likely as the result of numerical

and structural misconfiguration during a single mitotic failure. Therefore, the extent

of chromosome-scale copy number aberration observed in tumour genomes is likely

to be caused by much fewer events than the number of chromosomes affected.

Finally, our study enlightens the typical time scales of *in vivo* tumour progression,

with WGD occurring typically a decade before diagnosis and the onset of subclonal

variegation a few years before diagnosis. The presence of genetic aberrations for 10

years raises hopes that aberrant clones could be detected early, prior to reaching fully

malignant potential. Interestingly, this also shows that individual mutations, including

driver gene mutations and chromosomal gains, can occur several decades before

cancer diagnosis, demonstrating how cancer genomes have been shaped by a lifelong

process of somatic evolution.

## References

Cairns, J. Mutation selection and the natural history of cancer. *Nature* **255**, 197--200, doi:10.1038/255197a0 (1975).

Martincorena, I. & Campbell, P. J. Somatic mutation in cancer and normal cells. *Science* **349**, 1483-1489, doi:10.1126/science.aab4082 (2015).

Nowell, P. C. The clonal evolution of tumor cell populations. *Science* **194**, 23--28, doi:10.1126/science.959840 (1976).

Ahmad, A. S., Ormiston-Smith, N. & Sasieni, P. D. Trends in the lifetime risk of developing cancer in Great Britain: comparison of risk for those born from 1930 to 1960. *Br J Cancer* **112**, 943-947, doi:10.1038/bjc.2014.606 (2015).

Nik-Zainal, S. *et al.* The life history of 21 breast cancers. *Cell* **149**, 994-1007, doi:10.1016/j.cell.2012.04.023 (2012).

Durinck, S. *et al.* Temporal dissection of tumorigenesis in primary cancers. *Cancer Discov* **1**, 137-143, doi:10.1158/2159-8290.CD-11-0028 (2011).

Purdom, E. *et al.* Methods and challenges in timing chromosomal abnormalities within cancer samples. *Bioinformatics* **29**, 3113-3120, doi:10.1093/bioinformatics/btt546 (2013).

Mitchell, T. J. *et al.* Timing the Landmark Events in the Evolution of Clear Cell Renal Cell Cancer: TRACERx Renal. *Cell* **173**, 611-623.e617, doi:10.1016/j.cell.2018.02.020 (2018).

Gerlinger, M. *et al.* Intratumor Heterogeneity and Branched Evolution Revealed by Multiregion Sequencing. *New England Journal of Medicine* **366**, 883-892, doi:10.1056/NEJMoa1113205 (2012).

Gundem, G. *et al.* The evolutionary history of lethal metastatic prostate cancer. *Nature* **520**, 353-357, doi:10.1038/nature14347 (2015).

Yates, L. R. *et al.* Subclonal diversification of primary breast cancer revealed by multiregion sequencing. *Nat Med* **21**, 751-759, doi:10.1038/nm.3886 (2015).

Brastianos, P. K. *et al.* Genomic Characterization of Brain Metastases Reveals

Branched Evolution and Potential Therapeutic Targets. *Cancer Discov* **5**, 1164-1177, doi:10.1158/2159-8290.CD-15-0369 (2015).

Papaemmanuil, E. *et al.* Clinical and biological implications of driver mutations in myelodysplastic syndromes. *Blood* **122**, 3616-3627, doi:10.1182/blood-2013-08-518886 (2013).

Landau, D. A. *et al.* Mutations driving CLL and their evolution in progression and relapse. *Nature* **526**, 525-530, doi:10.1038/nature15395 (2015).

Campbell, P. J. *et al.* Pan-cancer analysis of whole genomes. *bioRxiv* (2017).

International Cancer Genome Consortium *et al.* International network of cancer genome projects. *Nature* **464**, 993-998, doi:10.1038/nature08987 (2010).

McLendon, R. *et al.* Comprehensive genomic characterization defines human glioblastoma genes and core pathways. *Nature* **455**, 1061-1068, doi:10.1038/nature07385 (2008).

Heim, S. *et al.* Trisomy 7 and sex chromosome loss in human brain tissue. *Cytogenet Cell Genet* **52**, 136-138, doi:10.1159/000132863 (1989).

Gao, R. *et al.* Punctuated copy number evolution and clonal stasis in triple-negative breast cancer. *Nat Genet* **48**, 1119-1130, doi:10.1038/ng.3641 (2016).

Sabarinathan, R. *et al.* The whole-genome panorama of cancer drivers. *bioRxiv*, doi:10.1101/190330 (2017).

Jamal-Hanjani, M. *et al.* Tracking the Evolution of Non-Small-Cell Lung Cancer. *N Engl J Med* **376**, 2109-2121, doi:10.1056/NEJMoa1616288 (2017).

Fearon, E. R. & Vogelstein, B. A genetic model for colorectal tumorigenesis. *Cell* **61**, 759--767, doi:10.1016/0092-8674(90)90186-I (1990).

Alexandrov, L. B. *et al.* Signatures of mutational processes in human cancer. *Nature* **500**, 415-421, doi:10.1038/nature12477 (2013).

Alexandrov, L. *et al.* The Repertoire of Mutational Signatures in Human Cancer. *bioRxiv* (2018).

25 McGranahan, N. *et al.* Clonal status of actionable driver events and the timing
of mutational processes in cancer evolution. *Sci Transl Med* **7**, 283ra254,
doi:10.1126/scitranslmed.aaa1408 (2015).

26 Rosenthal, R., McGranahan, N., Herrero, J., Taylor, B. S. & Swanton, C.
DeconstructSigs: delineating mutational processes in single tumors
distinguishes DNA repair deficiencies and patterns of carcinoma evolution.
*Genome Biol* **17**, 31, doi:10.1186/s13059-016-0893-4 (2016).

27 Alexandrov, L. B. *et al.* Clock-like mutational processes in human somatic
cells. *Nat Genet*, doi:10.1038/ng.3441 (2015).

28 Blokzijl, F. *et al.* Tissue-specific mutation accumulation in human adult stem
cells during life. *Nature* **538**, 260-264, doi:10.1038/nature19768 (2016).

29 Welch, J. S. *et al.* The origin and evolution of mutations in acute myeloid
leukemia. *Cell* **150**, 264-278, doi:10.1016/j.cell.2012.06.023 (2012).

30 Patch, A.-M. *et al.* Whole-genome characterization of chemoresistant ovarian
cancer. *Nature* **521**, 489-494, doi:10.1038/nature14410 (2015).

31 Yates, L. R. *et al.* Genomic Evolution of Breast Cancer Metastasis and
Relapse. *Cancer Cell* **32**, 169-184.e167, doi:10.1016/j.ccell.2017.07.005
(2017).

32 Ding, L. *et al.* Clonal evolution in relapsed acute myeloid leukaemia revealed
by whole-genome sequencing. *Nature*, doi:10.1038/nature10738 (2012).

33 Morrissy, A. S. *et al.* Divergent clonal selection dominates medulloblastoma
at recurrence. *Nature* **529**, 351-357, doi:10.1038/nature16478 (2016).

34 Tomasetti, C., Vogelstein, B. & Parmigiani, G. Half or more of the somatic
mutations in cancers of self-renewing tissues originate prior to tumor
initiation. *Proc Natl Acad Sci U S A* **110**, 1999-2004,
doi:10.1073/pnas.1221068110 (2013).

35 Purdie, D. M., Bain, C. J., Siskind, V., Webb, P. M. & Green, A. C. Ovulation
and risk of epithelial ovarian cancer. *Int J Cancer* **104**, 228-232,
doi:10.1002/ijc.10927 (2003).

36   Siegel, R. L., Miller, K. D. & Jemal, A. Cancer Statistics, 2017. *CA Cancer J*
*Clin* **67**, 7-30, doi:10.3322/caac.21387 (2017).

37   Thun, M. J., Henley, S. J. & Calle, E. E. Tobacco use and cancer: an
epidemiologic perspective for geneticists. *Oncogene* **21**, 7307-7325,
doi:10.1038/sj.onc.1205807 (2002).

38   Bostwick, D. G. & Qian, J. High-grade prostatic intraepithelial neoplasia. *Mod*
*Pathol* **17**, 360-379, doi:10.1038/modpathol.3800053 (2004).

39   Brenner, H. *et al.* Risk of progression of advanced adenomas to colorectal
cancer by age and sex: estimates based on 840,149 screening colonoscopies.
*Gut* **56**, 1585-1589, doi:10.1136/gut.2007.122739 (2007).

40   Gazdar, A. F. & Brambilla, E. Preneoplasia of lung cancer. *Cancer Biomark* **9**,
385-396, doi:10.3233/CBM-2011-0166 (2010).

41   Sanders, M. E., Schuyler, P. A., Dupont, W. D. & Page, D. L. The natural
history of low-grade ductal carcinoma in situ of the breast in women treated
by biopsy only revealed over 30 years of long-term follow-up. *Cancer* **103**,
2481-2484, doi:10.1002/cncr.21069 (2005).

42   Schlecht, N. F. *et al.* Human papillomavirus infection and time to progression
and regression of cervical intraepithelial neoplasia. *J Natl Cancer Inst* **95**,
1336-1343 (2003).

43   Whitson, M. J. & Falk, G. W. Predictors of Progression to High-Grade
Dysplasia or Adenocarcinoma in Barrett's Esophagus. *Gastroenterol Clin*
*North Am* **44**, 299-315, doi:10.1016/j.gtc.2015.02.005 (2015).

44   Xie, M. *et al.* Age-related mutations associated with clonal hematopoietic
expansion and malignancies. *Nat Med* **20**, 1472-1478, doi:10.1038/nm.3733
(2014).

45   Genovese, G. *et al.* Clonal hematopoiesis and blood-cancer risk inferred from
blood DNA sequence. *N Engl J Med* **371**, 2477-2487,
doi:10.1056/NEJMoa1409405 (2014).

46   Bardeesy, N. & DePinho, R. A. Pancreatic cancer biology and genetics. *Nat*

*Rev Cancer* **2**, 897-909, doi:10.1038/nrc949 (2002).

47    Folkins, A. K. *et al.* A candidate precursor to pelvic serous cancer (p53
signature) and its prevalence in ovaries and fallopian tubes from women with
BRCA          mutations.          *Gynecol          Oncol*          **109**,          168-173,
doi:10.1016/j.ygyno.2008.01.012 (2008).

48    Cancer Genome Atlas Research Network. Integrated genomic analyses of
ovarian carcinoma. *Nature* **474**, 609-615, doi:10.1038/nature10166 (2011).

49    Ciriello, G. *et al.* Emerging landscape of oncogenic signatures across human
cancers. *Nat Genet* **45**, 1127-1133, doi:10.1038/ng.2762 (2013).

50    Dentro, S. C. *et al.* Portraits of genetic intra-tumour heterogeneity and
subclonal selection across cancer types. *bioRxiv*, doi:10.1101/312041 (2018).

## Methods

### Data set

The PCAWG series consists of 2,778 tumour samples (2,703 white listed, 75 grey listed) that represent 2,658 cancers. All samples in this dataset underwent whole genome sequencing (minimum average coverage 30x in the tumour, 25x in the normal), and were processed with a set of project-specific pipelines for alignment, variant calling, and quality control[15]. Copy number calls were established by combining the output of six individual callers into a consensus using a multi-tier approach, resulting in a copy number profile, a purity and ploidy value and whether the tumour has undergone a whole genome duplication[50]. Consensus subclonal architectures have been obtained by integrating the output of eleven subclonal reconstruction callers[50], after which all SNVs, indels and SVs are assigned to a mutation cluster using the MutationTimer.R approach (**Supplementary Information**). Driver calls have been defined by the PCAWG Driver working group[20] while mutational signatures are defined by the PCAWG Signatures working group[24]. A more detailed description can be found in **Supplementary Information**, section 0.

### Timing of gains

We used three related approaches to calculate the timing of copy number gains (see **Supplementary Information**, section 1). In brief, the common feature is that the expected variant allele frequency of a mutation is related to the underlying number of alleles carrying a mutation according to the formula

$$E[X] = n\,m\,f\, / \,[N\,(1-\rho) + C\,\rho]$$

Here $X$ is the number of reads, $n$ denotes the coverage of the locus, the mutation copy number $m$ is the number of alleles carrying the mutation (which is usually inferred), $f$ is the frequency of the clone carrying the given mutation ($f = 1$ for clonal mutations). $N$ is the normal copy number (2 on autosomes, 1 or 2 for chromosome X and 0 or 1 for chromosome Y), $C$ the total copy number of the tumour and $\rho$ the purity of the sample.

The number of mutations at each allelic copy number then informs about the time

when the gain has occurred. The basic formulae for timing each gain are, depending

on the copy number configuration:

Copy number 2+1: $T = 3 n_2 / (2n_2 + n_1)$

Copy number 2+2: $T = 2 n_2 / (2n_2 + n_1)$

Copy number 2+0: $T = 2 n_2 / (2n_2 + n_1)$

Here 2+1 refers to major and minor copy number of 2 and 1, respectively. Methods

differ slightly in how the number of mutations present on each allele are calculated

and how uncertainty is handled (Supplementary Information).

**Timing of mutations**

The mutation copy number $m$ and the clonal frequency $f$ is calculated according to the

principles indicated above. Details can be found in **Supplementary Information**,

section 1.2. Mutations with $f = 1$ are denoted as *clonal*, and mutations with $f < 1$ as

*subclonal*. Mutations with $f = 1$ and $m > 1$ are denoted as *early clonal* (coamplified).

In cases with $f = 1$, $m = 1$ and $C > 2$, mutations were annotated as *late clonal*, if the

minor copy number was 0, otherwise *clonal* [*unspecified*] (**Supplementary**

**Information**, section 1.2.)

**Timing of driver mutations**

A catalogue of driver point mutations (SNVs and indels) was provided by the

PCAWG Drivers and Functional Interpretation Group[20]. The timing category was

calculated as above. From the four timing categories, odds ratios of early/late clonal

and clonal (early, late or unspecified clonal)/subclonal were calculated for driver

mutations against the distribution of all other mutations present in fragments with the

same copy number composition in the samples with each particular driver. The

background distribution of these odds ratios was assessed with 1,000 bootstraps

(**Supplementary Information**, section 3.1.)

**Integrative timing**

For each pair of driver point mutations and recurrent copy number alterations, an ordering was established (earlier, later or unspecified). The information underlying this decision was derived from the timing of each driver point mutation, as well as from the timing status of clonal and subclonal copy number segments. These tables were aggregated across all samples and a sports statistics model was employed to calculate the overall ranking of driver mutations. A full description is given in **Supplementary Information**, section 3.2.

**Timing of mutational signatures**

Mutational trinucleotide substitution signatures, as defined by the PCAWG Mutational Signatures Working Group[24], were fit to samples with observed signature activity, after splitting point mutations into either of the four epochs. A likelihood ratio test based on the multinomial distribution was used to test for differences in the mutation spectra between time points. Time-resolved exposures were calculated using non-negative linear least squares. Full details are given in **Supplementary Information**, section 4.

**Real-time estimation of WGD and MRCA**

CpG>TpG mutations were counted in an NpCpG context, except for Skin-Melanoma, where CpCpG and TpCpG were excluded due to the overlapping UV mutation spectrum. For visual comparison, the number of mutations was scaled to the effective genome size, defined as the $1 / \text{mean} (m_i / C_i)$ where $m_i$ is the estimated number of allelic copies of each mutation and $C_i$ the total copy number at that locus, thereby scaling to the final copy number and the time of change.

A hierarchical Bayesian linear regression was fit to relate the age at diagnosis to the scaled number of mutations, ensuring positive slope and intercept through a shared gamma distribution across cancer types.

For tumours with multiple time points, the set of mutations shared between diagnosis and relapse ($n_D$) and those specific to the relapse ($n_R$) was calculated. The rate

acceleration was calculated as $a = n_R / n_D \times t_D / t_R$. This analysis was performed

separately for all substitutions and for CpG>TpG mutations.

Based on these analyses, a typical increase of 5x for most cancer types was chosen,

with a lower value of 2.5x for brain cancers and a value of 7.5x for ovarian cancer.

The correction for transforming an estimate of a copy number gain in mutation time

into chronological time depends not only on the rate acceleration, but also on the time

at which this acceleration occurred. As this is generally unknown, we performed

Monte Carlo simulations of rate accelerations spanning an interval of 15 years prior to

diagnosis, corresponding roughly to 25% of time for a diagnosis at 60 years of age,

noting that a 5x rate increase over this duration yields an offset of about 33% of

mutations, compatible with our data. Subclonal mutations were assumed to occur at

full acceleration. The proportion of subclonal mutations was divided by the number of

identified subclones, thus conservatively assuming branching evolution. Full details

are given in **Supplementary Information**, section 5.

**Cancer timelines**

The results from each of the different timing analyses are combined in timelines of

cancer evolution for each tumour type (**Figure 7** and **Extended Data Fig. 1**). Each

timeline begins at the fertilised egg, and spans up to the median age of diagnosis

within each cohort. Real time estimates for WGD and the MRCA act as anchor

points, allowing us to roughly map the four broadly defined time periods (early

clonal, intermediate, late clonal and subclonal) to chronological time during a

patient's lifespan. Specific driver mutations or copy number alterations can be placed

within each of these time frames based on their ordering from the league model

analysis. The most active mutational signatures within each cohort are displayed, with

font sizes indicating the mean proportion of mutations attributed to each signature

during a given time period. Full details are given in **Supplementary Information**,

section 6.

## Data availability

Raw variant calls will be made available through the PCAWG data portal, https://dcc.icgc.org/pcawg. Processed data from the PCAWG Evolution and Heterogeneity working group, including timing estimates for all copy number gains, real time estimates of WGD and MRCA, as well as mutation signature activities, is available at https://www.synapse.org/#!Synapse:syn14193595. Analysis code can be downloaded via the github repository https://github.com/PCAWG-11/Evolution, containing submodules reflecting software tools and analysis workflows used in this analysis.

## Figure Legends

**Figure 1. Principles of timing mutations**

(**a**) Principles of timing mutations based on whole genome sequencing. According to the clonal evolution model of cancer, tumour cells evolve in multiple selective sweeps. During some of these sweeps, copy number gains are acquired, which can be used for timing analyses (green and purple epochs). Mutations acquired after the last clonal expansion are present in distinct subclonal populations (red epoch). The number of sequencing reads reporting point mutations can be used to discriminate variants as early or late clonal (green/purple) in cases of specific copy number gains, as well as clonal (blue) or subclonal (red) in cases without (right). The distribution of the number of early and late clonal mutations carries information about the timing of the copy number gains with the exact relation depending on the resulting copy number configuration (bottom). (**b**) Example case illustrating the annotation of point mutations based on the variant allele frequency (VAF, top) and copy number configuration including structural variants (middle). Each is shown as a function of genomic coordinate (x-axis). The resulting timing estimates for each copy number segment are shown at the bottom, indicating that, after an initial gain of chromosome arm 5q as part of an unbalanced translocation, a whole genome duplication occurred which gained all existing chromosomes in a single event.

**Figure 2. Pan-cancer timing patterns of arm-level gains**

(**a**) Overview of timing arm-level copy number gains across different cancer types. Pie charts depict the distribution of the inferred mutation time for a given copy number gain in a cancer type. Green colours denote early clonal gains prior to all point mutations (mutation time t=0) with a gradient to purple for late clonal gains after all clonal point mutations (t=1). The size of each chart is proportional to the recurrence of this event in a given tissue. (**b**) Heatmaps representing timing estimates of gains on different chromosome arms (x-axis) for individual samples (y-axis) for selected tumour types. (**c**) Two near-diploid example cases illustrating synchronous gains with a single peak in amplification activity (top) and asynchronous gains with multiple amplification periods (bottom). Secondary gains leading to three copies of

the same allele are separately indicated. (**d**) Distribution of synchronous and asynchronous gain patterns across samples, split by whole genome duplication status (left). Uninformative samples carry too few or too small gains to be timed accurately. Systematic permutation tests reveal a 61% enrichment of synchronous gains in near-diploid samples (right). (**e**) Proportion of large copy number segments with secondary gains per timing class. (**f**) Distribution of the relative latency of secondary gains, scaled to the time after the first gain. 0% = synchronous with first gain, 100% = at MRCA.

## Figure 3. Timing of driver point mutations

(**a**) Top: distribution of point mutations over different mutation periods in 2,778 samples. Middle: timing distribution of driver mutations in the 50 most recurrent lesions across 2,583 white listed samples from unique donors. Bottom: distribution of driver mutations across cancer types; colour as defined in the inset. (**b**) Relative timing of the 50 most recurrent driver lesions, calculated as the odds ratio of early versus late clonal driver mutations versus background (green, purple) or clonal versus subclonal. Odds ratios overlapping 1 in less than 5% of bootstrap samples are considered significant and have been coloured. Odds ratios have been truncated at 50. (**c**) Relative timing of *TP53* mutations across cancer types, coloured and truncated as in (**b**). The number of mutations in parenthesis represents the total number of *TP53* mutations used in the analysis for each cancer type. (**d**) Estimated number of unique lesions (genes) contributing 50% of all driver mutations in different timing epochs. Error bars denote the range between 0 and 1 pseudocounts.

## Figure 4. Relative ordering of somatic events

Preferential ordering of somatic copy number events and driver point mutations ithin tumour cohorts. (**a**) Schematic representation of the ordering process. (**b-d**) examples of individual patient trajectories (partial ordering relationships), the constituent data for the League model ordering process. Preferential ordering diagrams for (**e**) colorectal adenocarcinoma, (**f**) pancreatic neuroendocrine cancer and (**g**)

glioblastoma. Probability distributions show the uncertainty of timing for specific

events in the cohort. Events with odds above 10 (either earlier or later) are

highlighted. The prevalence of the event type in the cohort is displayed as a bar plot

on the right.

**Figure 5. Dynamic mutational processes during early and late clonal tumour**

**evolution**

(**a**) Examples of tumours with substantial changes between mutation spectra of early

and (top) late clonal time points (bottom). or each sample, the attribution of mutations

to the most characteristic single-base signatures are shown. Percentages missing from

100% are due to other, less characteristic contributions. (**b**) Fold changes between

relative proportions of early and late clonal mutations attributed to individual

mutational signatures. Points are coloured by tissue type. Data are shown for n = 530

samples with measurable changes in their overall mutation spectra and restricted to

signatures active in at least 10 samples.

**Figure 6. Real-time estimation of mutational landmarks**

(**a**) CpG>TpG mutations/Gb (scaled to the effective genome size) as a function of age

at diagnosis across cancer types. Colours are defined in panel **h**. (**b**) CpG>TpG

mutations/Gb for ovarian and breast cancer samples with matched primary and

relapse samples. (**c**) CpG>TpG mutation rate increase inferred from paired samples

for 6 cancer types. (**d**) Schematic of absolute timing under mutation rate increase. (**e**)

Time of occurrence of WGD in individual patients, split by tumour type. The error

bars at individual data points are 80% confidence intervals, reflecting uncertainty

stemming from the number of mutations per segment and onset of the rate increase,

which was chosen between 2.5x and 7.5x. Overlaid boxplots demarcate the quartiles

and median of the distribution with whiskers indicating 5% and 95% quantiles. (**f**)

Scatter plots showing the time of diagnosis (x-axis) and inferred time of WGD (y-

axis) with error bars as in **e**. (**g**) Median timing of WGD per cancer type, as a function

of CpG>TpG rate increase. Dotted lines indicate cancer types with less than 5

informative samples. Dashed and dash-dotted lines represent Lung-AdenoCa and

Lung-SCC, respectively; colours as in part **e.** (**h**) Timing of subclonal diversification

using CpG>TpG mutations in individual patients.

**Figure 7. Cancer timelines**

Typical timelines of tumour development, for (a) glioblastoma, (b) colorectal

adenocarcinoma, (c) squamous cell lung cancer, (d) ovarian adenocarcinoma, and (e)

pancreatic adenocarcinoma. Each timeline represents the length of time, in years,

between the fertilised egg and the median age of diagnosis per cancer type. Real time

estimates for major events, such as WGD and the emergence of the MRCA are used

to define early, variable, late and subclonal stages of tumour evolution approximately

in chronological time. Driver mutations and copy number aberrations are shown in

each stage according to their preferential timing, as defined by relative ordering.

Mutational signatures that, on average, change over the course of tumour evolution

(confidence intervals not overlapping 0) are shown in the epoch in which their activity

is greatest. Where applicable, lesions with a timing already defined in the literature

are annotated, * denotes events that agree with our timing, while † denotes events that

either were not found in our study, or were found to have a different timing.

## Extended Data Figure Legends

**Extended Data Figure 1. Summary of all results obtained per cancer type**

(**a**) Clustered heatmaps of mutational timing estimates for gained segments, per patient. Colours as indicated in main text: green represents early clonal events, purple represents late clonal. (**b**) Relative ordering of copy number events and driver mutations across all samples per cancer type. (**c**) Distribution of mutations across early clonal, late clonal and subclonal stages, for the most common driver genes per cancer type. A maximum of 10 driver genes are shown. (**d**) Clustered mutational signature fold changes between early clonal and late clonal stages, per patient. Green and purple indicate, respectively, a signature decrease and increase in late clonal from early clonal mutations. Inactive signatures are coloured white. (**e**) As in (**d**) but for clonal versus subclonal stages. Blue indicates a signature decrease and red an increase in subclonal from clonal mutations. (**f**) Typical timeline of tumour development, per cancer type.

**Extended Data Figure 2. Comparison of methods used for timing of individual copy-number gains**

(**a-b**) Pairwise comparison of the three approaches for timing individual copy number gains and (**c**) comparison with the simulated truth, showing high concordance.

**Extended Data Figure 3. Validation of League model relative ordering reconstruction based on simulated cohorts of whole-genome samples**

(**a**) League model results on a simulated cohort of samples from a single generalized relative order of events (with varied prevalence) showing high concordance with the true trajectory. (**b**) League model results on a simulated cohort of samples from a complex mixture of trajectories with different order of events showing high concordance with the expected average trajectory. (**c**) Estimation of accuracy of the League model reconstruction by simulation of a set of 100 cohorts with random trajectory mixtures and quantifying the distance in log odds early/late from perfect ordering. For the vast majority of events (even with low number of occurrences in the

cohort) the log odds error does not exceed 1, confirming that barely any of the events

would switch between timing categories.

**Extended Data Figure 4. Correlation between League model and Bradley-Terry**

**model ordering**

Direct comparison for each tumour type of the League and Bradley-Terry models for

determining the order of recurrent somatic mutations and copy number events. Axes

indicate the ordered events observed in the respective tumour types. Correlation is

quantified by Spearman's rank correlation coefficient.

**Extended Data Figure 5. Overview of early to late clonal signature changes**

**across tumour types**

Pie charts representing signature changes per cancer type. The colour scheme is as

above; signatures that decrease between early and late are coloured green, signatures

that increase are purple. The size of each pie chart represents the frequency of each

signature. Signatures are split into three categories: "clock-like", comprising the

putative clock signatures 1 and 5, "frequent", which are signatures present in 10 or

more cancer types, and "cancer type specific", which are in fewer than 10 cancer

types and are often limited to specific cohorts.

**Extended Data Figure 6. Absolute timing**

(**a**) CpG>TpG mutation/Gb (scaled to copy number) as a function of age at diagnosis

for 33 informative cancer types. The dotted line denotes the median mutations/yr (ie.,

not offset), while the shading denotes the 95% credible interval of a hierarchical

Bayesian linear regression model. (**b**) Maximum a posteriori estimates of rate and

offset for 33 cancer types with 95% credible intervals. (**c**) Median fraction of

mutations attributed to linear age-dependent accumulation, based on estimates from **b**

and the age at diagnosis for each sample. (**d**) Scatter plot of number of early (co-

amplified) CpG>TpG mutations (y-axis) as a function of the mutational time estimate

of WGD (x-axis). The black line denotes a non-linear loess fit with 95% confidence

interval. Colours define the cancer type as in part **a**. (**e**) Total CpG>TpG mutations (y-

axis) as a function of the mutation time estimate of WGD (x-axis). Colours and fit as

in **d**. (**f**) Inferred median absolute timing patient-specific rate increase, depending on

the observed CpG>TpG mutation burden, allowing for a higher (up to 10x) mutation

rate increase in cancers with more mutations. (**g**). Comparison of the median duration

of subclonal diversification per cancer type assuming branching and linear

phylogenies.

## Author contributions

MG, CJ, IL, SG, PA, DR, DGL, PTS and PVL performed timing of point mutations and copy number gains. SG and MG performed qualitative timing of driver point mutations and analyses of synchronous gains, LJ timed secondary copy number gains. IL, TJM, DR, DGL, DCW and GG performed relative timing of somatic driver events and implemented integrative models. CJ, YR, PVL and QDM performed timing of mutational signatures. MG performed real-time estimation of whole-genome duplication and subclonal diversification. SG assessed mutation rates in relapsed samples. CJ, MG, IL, YR, DR and PVL constructed cancer timelines. MG, CJ, IL, SCD, SG, TJM, YR, PA, JD, PCB, DDB, VM, QDM, PTS, DCW and PVL interpreted the results. SCD, IL, JW, AD, IVG, KeY, GM, MP, SM, ND, KaY, SSe, KH, MT, JD, DGL, DR, JL, MC, SCS, YJ, FM, VM, HZ, WW, QDM, DCW and PVL performed subclonal architecture analysis. SCD, IL, KK, VM, MP, XY, DGL, SSc, RB, MI, MS, DCW and PVL performed copy number analysis. JW, SCD, IL, KH, DGL, KK, DR, DCW, QDM and PVL derived a consensus of copy number analysis results. KaY, MT, AD, SCD, IL, DCW, MG, PVL, QDM and WW derived a consensus of subclonal architecture results. YF and WW contributed to subclonal mutation calls. PTS, DCW and PVL coordinated the study. MG, CJ, PTS, YR, IL, QDM, DCW and PVL wrote the manuscript, which all authors approved.

## Acknowledgements

This work was supported by the Francis Crick Institute, which receives its core funding from Cancer Research UK (FC001202), the UK Medical Research Council (FC001202), and the Wellcome Trust (FC001202). This project was enabled through access to the MRC eMedLab Medical Bioinformatics infrastructure, supported by the Medical Research Council (grant number MR/L016311/1). MT and JD are postdoctoral fellows supported by the European Union's Horizon 2020 research and innovation program (Marie Skłodowska-Curie Grant Agreement No. 747852-SIOMICS and 703594-DECODE). JD is a postdoctoral fellow of the FWO. FM, GM and KeY would like to acknowledge the support of the University of Cambridge, Cancer Research UK and Hutchison Whampoa Limited. GM, KeY and FM were funded by CRUK core grants C14303/A17197 and A19274. SSe and YJ are supported by NIH R01 CA132897. SM is supported by the Vanier Canada Graduate Scholarship. SCS is supported by the NSERC Discovery Frontiers Project, "The Cancer Genome Collaboratory" and NIH Grant GM108308. HZ is supported by grant NIMH086633 and an endowed Bao-Shan Jing Professorship in Diagnostic Imaging. WW is supported by the U.S. National Cancer Institute (1R01 CA183793 and P30 CA016672). PTS was supported by U24CA210957 and 1U24CA143799. DCW is funded by the Li Ka Shing foundation. PVL is a Winton Group Leader in recognition of the Winton Charitable Foundation's support towards the establishment of The Francis Crick Institute.

## Members of the PCAWG Evolution and Heterogeneity Working Group

Stefan C. Dentro[1,2,3,*], Ignaty Leshchiner[4,*], Moritz Gerstung[5,*], Clemency Jolly[1,*], Kerstin Haase[1,*], Maxime Tarabichi[1,2,*], Jeff Wintersinger[6,7,*], Amit G. Deshwar[6,7,*], Kaixian Yu[8,*], Santiago Gonzalez[5,*], Yulia Rubanova[6,7,*], Geoff Macintyre[9,*], David J. Adams[2], Pavana Anur[10], Rameen Beroukhim[4,11], Paul C. Boutros[6,12], David D. Bowtell[13], Peter J. Campbell[2], Shaolong Cao[8], Elizabeth L. Christie[13,14], Marek Cmero[14,15], Yupeng Cun[16], Kevin J. Dawson[2], Jonas Demeulemeester[1,17], Nilgun Donmez[18,19], Ruben M. Drews[9], Roland Eils[20,21], Yu Fan[8], Matthew Fittall[1], Dale W. Garsed[13,14], Gad Getz[4,22,23,24], Gavin Ha[4], Marcin Imielinski[25,26], Lara Jerman[5,27], Yuan Ji[28,29], Kortine Kleinheinz[20,21], Juhee Lee[30], Henry Lee-Six[2], Dimitri G. Livitz[4], Salem Malikic[18,19], Florian Markowetz[9], Inigo Martincorena[2], Thomas J. Mitchell[2,31], Ville Mustonen[32], Layla Oesper[33], Martin Peifer[16], Myron Peto[10], Benjamin J. Raphael[34], Daniel Rosebrock[4], S. Cenk Sahinalp[19,35], Adriana Salcedo[12], Matthias Schlesner[20], Steven Schumacher[4], Subhajit Sengupta[28], Ruian Shi[6], Seung Jun Shin[8,36], Lincoln D. Stein[12], Ignacio Vázquez-García[2,31], Shankar Vembu[6], David A. Wheeler[37], Tsun-Po Yang[16], Xiaotong Yao[25,26], Ke Yuan[9,38], Hongtu Zhu[8], Wenyi Wang[8,#], Quaid D. Morris[6,7,#], Paul T. Spellman[10,#], David C. Wedge[3,39,#], Peter Van Loo[1,17,#]

[1]The Francis Crick Institute, London NW1 1AT, United Kingdom; [2]Wellcome Trust Sanger Institute, Cambridge CB10 1SA, United Kingdom; [3]Big Data Institute, University of Oxford, Oxford OX3 7LF, United Kingdom; [4]Broad Institute of MIT and Harvard, Cambridge, MA 02142, USA; [5]European Molecular Biology Laboratory, European Bioinformatics Institute (EMBL-EBI), Cambridge CB10 1SD, United Kingdom; [6]University of Toronto, Toronto, ON M5S 3E1, Canada; [7]Vector Institute, Toronto, ON M5G 1L7, Canada; [8]The University of Texas MD Anderson Cancer Center, Houston, TX 77030, USA; [9]Cancer Research UK Cambridge Institute, University of Cambridge, Cambridge CB2 0RE, United Kingdom; [10]Molecular and Medical Genetics, Oregon Health & Science University, Portland, OR 97231, USA; [11]Dana-Farber Cancer Institute, Boston, MA 02215, USA; [12]Ontario Institute for Cancer Research, Toronto, ON M5G 0A3, Canada; [13]Peter MacCallum Cancer Centre, Melbourne, VIC 3000, Australia; [14]University of Melbourne, Melbourne, VIC 3010,

Australia; [15]Walter + Eliza Hall Institute, Melbourne, VIC 3000, Australia; [16]University of

Cologne, 50931 Cologne, Germany; [17]University of Leuven, B-3000 Leuven, Belgium;

[18]Simon Fraser University, Burnaby, BC V5A 1S6, Canada; [19]Vancouver Prostate Centre,

Vancouver, BC V6H 3Z6, Canada; [20]German Cancer Research Center (DKFZ), 69120

Heidelberg, Germany; [21]Heidelberg University, 69120 Heidelberg, Germany; [22]Massachusetts

General Hospital Center for Cancer Research, Charlestown, MA 02129, USA;

[23]Massachusetts General Hospital, Department of Pathology, Boston, MA 02114, USA;

[24]Harvard Medical School, Boston, MA 02215, USA; [25]Weill Cornell Medicine, New York,

NY 10065, USA; [26]New York Genome Center, New York, NY 10013, USA; [27]University of

Ljubljana, 1000 Ljubljana, Slovenia; [28]NorthShore University HealthSystem, Evanston, IL

60201, USA; [29]The University of Chicago, Chicago, IL 60637, USA; [30]University of

California Santa Cruz, Santa Cruz, CA 95064, USA; [31]University of Cambridge, Cambridge

CB2 0QQ, United Kingdom; [32]University of Helsinki, 00014 Helsinki, Finland; [33]Carleton

College, Northfield, MN 55057, USA; [34]Princeton University, Princeton, NJ 08540, USA;

[35]Indiana University, Bloomington, IN 47405, USA; [36]Korea University, Seoul,

02481, Republic of Korea; [37]Human Genome Sequencing Center, Baylor College of

Medicine, Houston, TX 77030, USA; [38]University of Glasgow, Glasgow G12 8RZ, United

Kingdom; [39]Oxford NIHR Biomedical Research Centre, Oxford OX4 2PG, United Kingdom.

*: These authors contributed equally

#: These authors jointly directed the work

# Figure 1: Principles of timing mutations

**Principles of timing mutations. (a)** Principles of timing mutations based on whole genome sequencing. According to the clonal evolution model of cancer, tumour cells evolve in multiple selective sweeps. During some of these sweeps, copy number gains are acquired, which can be used for timing analyses (green and purple epochs). Mutations acquired after the last clonal expansion are present in distinct subclonal populations (red epoch). The number of sequencing reads reporting point mutations can be used to discriminate variants as early or late clonal (green/purple) in cases of specific copy number gains, as well as clonal (blue) or subclonal (red) in cases without (right). The distribution of the number of early and late clonal mutations carries information about the timing of the copy number gains with the exact relation depending on the resulting copy number configuration (bottom). **(b)** Example case illustrating the annotation of point mutations based on the variant allele frequency (VAF, top) and copy number configuration including structural variants (middle). Each is shown as a function of genomic coordinate (x-axis). The resulting timing estimates for each copy number segment are shown at the bottom, indicating that, after an initial gain of chromosome arm 5q as part of an unbalanced translocation, a whole genome duplication occurred which gained all existing chromosomes in a single event.

# Figure 2. Pan-cancer timing patterns of arm-level gains

**a** Timing of arm-level gains, per chromosome and cancer type

**b** Timing heatmaps of individual tumour samples

**c** Temporal patterns of primary copy number gains

**Synchronous:** > 80% co-occuring
SA501385, 33yr, Liver-HCC, ploidy=2.4

**Asynchronous:** < 80% co-occuring
SA542034, 90yr, Lymph-BNHL, ploidy=2.2

**d** Distribution of primary amplification activity patterns

WGD asynchronous n=149
near diploid synchronous, n=468
WGD synchronous n=542
near diploid asynchronous n=347
WGD uninformative n=129
near diploid uninformative n=1143

$p < 0.01$

**e** Proportion of secondary copy number gains

**f** Timing of secondary copy number gains

**Pan-cancer timing patterns of arm-level gains. (a)** Overview of timing arm-level copy number gains across different cancer types. Pie charts depict the distribution of the inferred mutation time for a given copy number gain in a cancer type. Green colours denote early clonal gains prior to all point mutations (mutation time t=0) with a gradient to purple for late clonal gains after all clonal point mutations (t=1). The size of each chart is proportional to the recurrence of this event in a given tissue. **(b)** Heatmaps representing timing estimates of gains on different chromosome arms (x-axis) for individual samples (y-axis) for selected tumour types. **(c)** Two near-diploid example cases illustrating synchronous gains with a single peak in amplification activity (top) and asynchronous gains with multiple amplification periods (bottom). Secondary gains leading to three copies of the same allele are separately indicated. **(d)** Distribution of synchronous and asynchronous gain patterns across samples, split by whole genome duplication status (left). Uninformative samples carry too few or too small gains to be timed accurately. Systematic permutation tests reveal a 61% enrichment of synchronous gains in near-diploid samples (right). **(e)** Proportion of large copy number segments with secondary gains per timing class. (f) Distribution of the relative latency of secondary gains, scaled to the time after the first gain. 0% = synchronous with first gain, 100% = at MRCA.

# Figure 3: Timing of driver mutations

**Timing of driver point mutations. (a)** Top: distribution of point mutations over di¬fferent mutation periods in 2,778 samples. Middle: timing distribution of driver mutations in the 50 most recurrent lesions across 2,583 white listed samples from unique donors. Bottom: distribution of driver mutations across cancer types; colour as defined in the inset. **(b)** Relative timing of the 50 most recurrent driver lesions, calculated as the odds ratio of early versus late clonal driver mutations versus background (green, purple) or clonal versus subclonal. Odds ratios overlapping 1 in less than 5% of bootstrap samples are considered significant and have been coloured. Odds ratios have been truncated at 50. **(c)** Relative timing of TP53 mutations across cancer types, coloured and truncated as in **(b)**. The number of mutations in parenthesis represents the total number of TP53 mutations used in the analysis for each cancer type. **(d)** Estimated number of unique lesions (genes) contributing 50% of all driver mutations in different timing epochs. Error bars denote the range between 0 and 1 pseudocounts.

# Figure 4: Relative ordering of somatic events

**Relative ordering of somatic events.** Preferential ordering of somatic copy number events and driver point mutations within tumour cohorts. **(a)** Schematic representation of the ordering process. **(b-d)** examples of individual patient trajectories (partial ordering relationships), the constituent data for the League model ordering process. Preferential ordering diagrams for **(e)** colorectal adenocarcinoma, **(f)** pancreatic neuroendocrine cancer and **(g)** glioblastoma. Probability distributions show the uncertainty of timing for specific events in the cohort. Events with odds above 10 (either earlier or later) are highlighted. The prevalence of the event type in the cohort is displayed as a bar plot on the right.

**Figure 5:** Dynamic mutational processes during early and late clonal tumour evolution

Dynamic mutational processes during early and late clonal tumour evolution. **(a)** Examples of tumours with substantial changes between mutation spectra of early and (top) late clonal time points (bottom). or each sample, the attribution of mutations to the most characteristic single-base signatures are shown. Percentages missing from 100% are due to other, less characteristic contributions. **(b)** Fold changes between relative proportions of early and late clonal mutations attributed to individual mutational signatures. Points are coloured by tissue type. Data are shown for n = 530 samples with measurable changes in their overall mutation spectra and restricted to signatures active in at least 10 samples.

**Figure 6:** Real-time estimates

**Real-time estimation of mutational landmarks. (a)** CpG>TpG mutations/Gb (scaled to the effective genome size) as a function of age at diagnosis across cancer types. Colours are defined in panel **h**. **(b)** CpG>TpG mutations/Gb for ovarian and breast cancer samples with matched primary and relapse samples. **(c)** CpG>TpG mutation rate increase inferred from paired samples for 6 cancer types. **(d)** Schematic of absolute timing under mutation rate increase. **(e)** Time of occurrence of WGD in individual patients, split by tumour type. The error bars at individual data points are 80% confidence intervals, reflecting uncertainty stemming from the number of mutations per segment and onset of the rate increase, which was chosen between 2.5x and 7.5x. Overlaid boxplots demarcate the quartiles and median of the distribution with whiskers indicating 5% and 95% quantiles. **(f)** Scatter plots showing the time of diagnosis (x-axis) and inferred time of WGD (y-axis) with error bars as in e. **(g)** Median timing of WGD per cancer type, as a function of CpG>TpG rate increase. Dotted lines indicate cancer types with less than 5 informative samples. Dashed and dash-dotted lines represent Lung-AdenoCa and Lung-SCC, respectively; colours as in part e. **(h)** Timing of subclonal diversification using CpG>TpG mutations in individual patients.

# Figure 7: Cancer timelines

**Cancer timelines.** Typical timelines of tumour development, for **(a)** glioblastoma, **(b)** colorectal adenocarcinoma, **(c)** squamous cell lung cancer, **(d)** ovarian adenocarcinoma, and **(e)** pancreatic adenocarcinoma. Each timeline represents the length of time, in years, between the fertilised egg and the median age of diagnosis per cancer type. Real time estimates for major events, such as WGD and the emergence of the MRCA are used to define early, variable, late and subclonal stages of tumour evolution approximately in chronological time. Driver mutations and copy number aberrations are shown in each stage according to their preferential timing, as defined by relative ordering. Mutational signatures that, on average, change over the course of tumour evolution (confidence intervals not overlapping 0) are shown in the epoch in which their activity is greatest. Where applicable, lesions with a timing already defined in the literature are annotated, * denotes events that agree with our timing, while † denotes events that either were not found in our study, or were found to have a different timing.

**Extended Data Figure 1: Summary of all results obtained per cancer type. (a)** Clustered heatmaps of mutational timing estimates for gained segments, per patient. Colours as indicated in main text: green represents early clonal events, purple represents late clonal. **(b)** Relative ordering of copy-number events and driver mutations across all samples per cancer type. **(c)** Distribution of mutations across early clonal, late clonal and subclonal stages, for the most common driver genes per cancer type. A maximum of 20 driver genes are shown. **(d)** Clustered mutational signature fold changes between early clonal and late clonal stages, per patient. Green and purple indicate, respectively, a signature decrease and increase in late clonal from early clonal mutations. Inactive signatures are coloured white. **(e)** As in **(d)** but for clonal vs. subclonal stages. Blue indicates a signature decrease and red an increase in subclonal from clonal mutations. **(f)** Typical timeline of tumour development, per cancer type.

## Contents

# Extended Data Figure 1. Biliary-AdenoCA

**a**

Time, fraction of mutations

early clonal — 0.00 0.25 0.50 0.75 1.00 — late clonal

samples

1p 1q 2p 2q 3p 3q 4p 4q 5p 5q 6p 6q 7p 7q 8p 8q 9p 9q 10p 10q 11p 11q 12p 12q 13q 14q 15q 16p 16q 17p 17q 18p 18q 19p 19q 20p 20q 21q 22q WGD

**c**

Driver mutation frequency

- early clonal
- late clonal
- clonal [unsp.]
- subclonal

TP53 ARID1A SMAD4 KRAS ACVR2A PIK3CA PTEN

**b**

prevalence

| event | | |
|---|---|---|
| loss 9p | | 56% |
| homdel 9p21.3 | | 24% |
| loss 17p | | 50% |
| loss 14q32.33 | | 18% |
| loss ARID1A | | 38% |
| TP53 | | 28% |
| loss 6q | | 56% |
| ARID1A | | 15% |
| loss 1p36.23 | | 38% |
| gain 1q | | 26% |
| KRAS | | 12% |
| SMAD4 | | 15% |
| loss 4q | | 47% |
| PTEN | | 9% |
| gain 8q | | 32% |
| gain 7q | | 21% |
| loss 9q | | 41% |
| loss 8p | | 53% |
| loss 14q | | 50% |
| gain 7p | | 28% |
| loss RB1 | | 18% |
| WGD | | 26% |
| gain 12p | | 21% |
| loss 3p | | 56% |
| loss 13q | | 44% |
| loss 20p | | 41% |

odds late > 10

odds early > 10

relative rank (preference)

early ⟶ late/subclonal

**d** Early/late

Signature fold change, early vs. late

10+ 5 2 1 0.5 0.2 0.1–

SBS5 SBS1 ID1 ID2 SBS40 SBS2&13 DBS2 SBS18 ID8 DBS9 DBS4 SBS6-44 SBS32 SBS17 DBS7

**e** Clonal/subclonal

Signature fold change, clonal vs. subclonal

10+ 5 2 1 0.5 0.2 0.1–

SBS5 SBS1 ID1 SBS40 ID2 SBS2&13 DBS2 SBS18 ID8 DBS9 DBS4 SBS6-44 SBS32 SBS24 SBS17 SBS12 DBS7

**f**

preferentially early | variable/constant | late | subclonal

**Drivers:** *TP53*
**CNA:** -9p, --9p21.3, -17p, -14q32.33, -ARID1A

**Signatures:**
SBS: 2&13

**Drivers:** *ARID1A, KRAS, SMAD4*
**CNA:** -6q, -1p36.23, +1q, -4q, +8q, -7q, -9q, -8p, -14q, +7p, -RB1

**Signatures:**
DBS: 2, 4, 9, 11
ID: 1, 2, 8
SBS: 1, 5, 12, 18, 40

**CNA:** +12p, -3p, -13q, -20p

**Signatures:**
SBS: 1, 40

fertilised egg

-2.5yrs

WGD (26% samples) -1.2 yrs IQR [0.8, 3.9]

MRCA -0.1 yrs IQR [0, 1.2]

diagnosis 64 yrs, IQR [58, 74]

last detectable subclone

# Extended Data Figure 1. Bladder-TCC

**a** — Time, fraction of mutations

early clonal — 0.00  0.25  0.50  0.75  1.00 — late clonal

samples

1p 1q 2p 2q 3p 3q 4p 4q 5p 5q 6p 6q 7p 7q 8p 8q 9p 9q 10p 10q 11p 11q 12p 12q 13q 14q 15q 16p 16q 17p 17q 18p 18q 19p 19q 20p 20q 21q 22q WGD

**c**

Driver mutation frequency

- early clonal
- late clonal
- clonal [unsp.]
- subclonal

TERT  KDM6A  TP53  FGFR3  KMT2D  ARID1A  CDKN1A  RB1  ELF3

**b**

event (top to bottom):
loss 17p — 71%
TP53 — 43%
FGFR3 — 24%
loss 2q37.1 — 33%
CDKN1A — 19%
loss 9p — 67%
homdel 9p21.3 — 43%
promTERT — 62%
loss 9q — 81%
KDM6A — 28%
loss 11p — 67%
ARID1A — 28%
RB1 — 14%
gain 1q23.3 — 19%
loss 16p13.3 — 24%
WGD — 62%
gain 4p16.3 — 28%
loss 8p — 62%
gain 20q — 33%
loss 6q — 56%
gain 3q — 33%
loss 14q — 62%
gain 17q — 43%
gain 3p — 28%
loss 4p — 56%
gain 1q — 33%
loss 4q — 62%
loss 19p — 56%

odds late > 10
odds early > 10

early ——→ late/subclonal

relative rank (preference)

**d** Early/late

SBS5  SBS2&13  SBS1  ID2  ID1  DBS11  DBS2  SBS40  DBS4  DBS6

Signature fold change, early vs. late
10+  5  2  1  0.5  0.2  0.1−

**e** Clonal/subclonal

SBS5  SBS2&13  SBS1  ID1  ID2  DBS11  DBS2  ID8  SBS40  DBS4  DBS6

Signature fold change, clonal vs. subclonal
10+  5  2  1  0.5  0.2  0.1−

**f**

**preferentially early**

**Drivers:** *TP35, FGFR3, CDKN1A, TERT*
**CNA:** -17p, -2q37.1, -9p, --9p21.3, -9q
**Signatures:** SBS: 2&13, 5

**variable/constant**

**Drivers:** *KDM6A, RB1*
**CNA:** +1q23.3, -16p13.3
**Signatures:** DBS: 2, 4, 11; SBS: 40

**late**

**CNA:** +4p16.3, -8p, +20q, -6q, +3q, -14q, +17q, +3p, -4p, +1q, -4q, -19p
**Signatures:** ID: 1, 2, 8; SBS: 1

**subclonal**

fertilised egg

-10yrs

diagnosis 65 yrs, IQR [56, 77]

MRCA -0 yrs IQR [0, 0.5]

last detectable subclone

WGD (65% samples) -7.9yrs IQR [4.1, 18.7]

# Extended Data Figure 1. Bone-Osteosarc

# Extended Data Figure 1. Breast-AdenoCA

# Extended Data Figure 1. Cervix-SCC

**a**

**b** Early/late

**c** Clonal/subclonal

**d**

# Extended Data Figure 1. CNS-GBM

# Extended Data Figure 1. CNS-Medullo

# Extended Data Figure 1. CNS-Oligo

**a**

**b**

**c** Early/late

**d** Clonal/subclonal

**e**

# Extended Data Figure 1. CNS-PiloAstro

a

Time, fraction of mutations
early clonal   0.00  0.25  0.50  0.75  1.00   late clonal

samples

1p 1q 2p 2q 3p 3q 4p 4q 5p 5q 6p 6q 7p 7q 8p 8q 9p 9q 10p 10q 11p 11q 12p 12q 13q 14q 15q 16p 16q 17p 17q 18p 18q 19p 19q 20p 20q 21q 22q WGD

b

prevalence

gain 11 — 10%
gain 6 — 16%
gain 5 — 16%
gain 10 — 7%
FGFR1 — 6%
gain 8 — 6%
BRAF — 7%
gain 12 — 9%
gain 7 — 12%

odds
late > 10
odds
early > 10

event

1     5
early ⟶ late/subclonal
relative rank (preference)

c

Driver mutation frequency

early clonal
late clonal
clonal [unsp.]

BRAF   FGFR1

d

Early/late

Signature fold change, early vs. late
10+ 5 2 1 0.5 0.2 0.1−

SBS5  SBS1  SBS40  SBS19  ID1

e

Clonal/subclonal

Signature fold change, clonal vs. subclonal
10+ 5 2 1 0.5 0.2 0.1−

SBS5  SBS1  SBS40  ID1  SBS19  SBS23

f

clonal, variable/constant          subclonal

CNA: +11, +6, +5, +7

Signatures:
ID: 1, 2
SBS: 1, 5, 40

diagnosis
8 yrs,
IQR [4, 12]

fertilised egg     -2.5yrs

MRCA
-0.9 yrs
IQR [0.5, 3.8]

last detectable
subclone

# Extended Data Figure 1. ColoRect-AdenoCA

# Extended Data Figure 1. Eso-AdenoCA

# Extended Data Figure 1. Head-SCC

# Extended Data Figure 1. Kidney-ChRCC

**a**

**b**

**c**     **d** Early/late     **e** Clonal/subclonal

**f**

# Extended Data Figure 1. Kidney-RCC.clearcell

# Extended Data Figure 1. Kidney-RCC.papillary

# Extended Data Figure 1. Liver-HCC

# Extended Data Figure 1. Lung-AdenoCA

# Extended Data Figure 1. Lung-SCC

# Extended Data Figure 1. Lymph-BNHL

# Extended Data Figure 1. Lymph-CLL

# Extended Data Figure 1. Myeloid-AML

**a**

**b** Early/late

**c** Clonal/subclonal

**e**

# Extended Data Figure 1. Myeloid-MPN

# Extended Data Figure 1. Ovary-AdenoCA

# Extended Data Figure 1. Panc-AdenoCA

# Extended Data Figure 1. Panc-Endocrine

# Extended Data Figure 1. Prost-AdenoCA

# Extended Data Figure 1. Skin-Melanoma

# Extended Data Figure 1. SoftTissue-Leiomyo

**a**

samples

1p 1q 2p 2q 3p 3q 4p 4q 5p 5q 6p 6q 7p 7q 8p 8q 9p 9q 10p 10q 11p 11q 12p 12q 13q 14q 15q 16p 16q 17p 17q 18p 18q 19p 19q 20p 20q 21q 22q WGD

**b**

Driver mutation frequency

- early clonal
- late clonal

TP53

**c**

Early/late

SBS5 SBS40 SBS1 ID1 ID2 ID8 SBS2&13 DBS2 DBS9 DBS6 DBS4 SBS30 SBS17 DBS11

Signature fold change,
early vs. late

10+
5
2
1
0.5
0.2
0.1−

**d**

Clonal/subclonal

SBS5 SBS40 SBS1 ID1 ID2 ID8 SBS2&13 DBS2 DBS9 DBS6 DBS4 SBS30 SBS17 DBS11

Signature fold change
clonal vs. subclonal

10+
5
2
1
0.5
0.2
0.1−

**e**

preferentially early | variable/constant | late | subclonal

**Signatures:**
**SBS:** 1

**Sigs:**
**DBS:** 2, 4, 6, 9
**ID:** 1, 2, 8
**SBS:** 2&13, 5, 40

fertilised egg

WGD (47% samples)
-14.2yrs IQR [11.4, 115]

-10yrs

MRCA
-0.9 yrs
IQR [0.4, 1.9]

diagnosis
63 yrs,
IQR [56,70]

**last detectable subclone**

# Extended Data Figure 1. SoftTissue-Liposarc

**a**

**b**

Early/late

**c**

Clonal/subclonal

**d**

# Extended Data Figure 1. Stomach-AdenoCA

# Extended Data Figure 1. Thy-AdenoCA

**a**

prevalence

event

BRAF — 21%

promTERT — 21%

early ⟶ late/subclonal

relative rank (preference)

**b**

Driver mutation frequency

- early clonal
- late clonal
- clonal [unsp.]

TERT  BRAF

**c** Early/late

Signature fold change, early vs. late

10+
5
2
1
0.5
0.2
0.1–

SBS5  SBS40  SBS1  ID1

**d** Clonal/subclonal

Signature fold change, clonal vs. subclonal

10+
5
2
1
0.5
0.2
0.1–

SBS5  SBS1  SBS40  ID1  SBS2&13  ID8  ID2  DBS4  DBS2

**e**

clonal, variable/constant

Drivers: *BRAF*

Signatures:
**DBS: 2, 4, 11**
**ID: 1, 2, 8**
**SBS: 1, 2&13, 5, 40**

subclonal

Drivers: *TERT*

fertilised egg

-2.5yrs

MRCA
-2.3 yrs
IQR [1.4, 3.7]

diagnosis
51 yrs,
IQR [42, 65]

last detectable subclone

# Extended Data Figure 1. Uterus-AdenoCA

**a**

Time, fraction of mutations
early clonal — late clonal
0.00 0.25 0.50 0.75 1.00

samples

1p 1q 2p 2q 3p 3q 4p 4q 5p 5q 6p 6q 7p 7q 8p 8q 9p 9q 10p 10q 11p 11q 12p 12q 13q 14q 15q 16p 16q 17p 17q 18p 18q 19p 19q 20p 20q 21q 22q WGD

**c**

event

loss 17p
loss 19p13.3
loss 17q11.2
FBXW7
PPP2R1A
PTEN
ARHGAP35
gain 1q
PIK3R1
loss 16q
loss 9q
TP53
PIK3CA
gain 10q
gain 8q
gain 17q12
homdel 10q23.31
gain 19q12
KRAS
ARID1A
loss 22q
loss 3p
loss 8p
CTNNB1
loss 4q
gain 20q
WGD
loss 10q
homdel 13q14.2
loss 21q
gain 3q
loss 15q

odds late > 10
odds early > 10

1   5   10   15   20   25   30
early ⟶ late/subclonal
relative rank (preference)

prevalence
59%
45%
12%
16%
20%
33%
24%
33%
31%
53%
53%
61%
45%
20%
27%
12%
6%
28%
10%
28%
37%
39%
47%
14%
37%
22%
35%
37%
8%
39%
28%
47%

**b**

Driver mutation frequency

TP53 PIK3CA PTEN PIK3R1 ARID1A ARHGAP35 PPP2R1A CTNNB1 FBXW7 MAP3K1 KMT2C KRAS

early clonal
late clonal
clonal [unsp.]
subclonal

**d**

Early/late

SBS5 ID2 ID1 SBS1 SBS40 ID8 SBS9 SBS2&13 DBS4 SBS6 SBS11 DBS2 SBS7 DBS3 DBS10 SBS3 SBS28 SBS10 DBS5

Signature fold change, early vs. late
10+
5
2
1
0.5
0.2
0.1–

**e**

Clonal/subclonal

SBS5 ID2 ID1 SBS1 SBS40 ID8 SBS9 SBS2&13 SBS6-44 DBS4 SBS6 SBS11 DBS2 SBS7 DBS3 DBS10 SBS28 SBS10 DBS5

Signature fold change, clonal vs. subclonal
10+
5
2
1
0.5
0.2
0.1–

**f**

preferentially early
**Drivers:** *FBXW7, PPP2R1A, PTEN, ARHGAP35*
**CNA:** -17p, -19p13.3, -17q11.2, +1q
**Signatures:**
**ID:** 1
**SBS:** 1, 5

variable/constant
**Drivers:** *PIK3R1, TP53, KRAS, ARID1A , CTNNB1,*
**CNA:** -16q, -9q, +10q, +8q, +17q12, +19q12, -22q, -3p, -8p, -4q, +20q
**Signatures:**
**DBS:** 2, 4, 5, 7, 9, 11
**ID:** 2, 8; **SBS:** 6-44, 40

late
**CNA:** -10q, -21q, +3q, -15q
**Signatures:**
**DBS:** 6
**SBS:** 2&13

subclonal
**Sigs:**
**DBS:** 6,
**SBS:** 2&13

fertilised egg

-5yrs

WGD (33% samples)
-4.6yrs IQR [3.2, 8.7]

MRCA
-0.9 yrs
IQR [0.3, 2.1]

diagnosis
69 yrs,
IQR [62,74]

last detectable subclone

**Extended Data Figure 2:** Comparison of methods used for timing of individual copy-number gains

**Comparison of methods used for timing of individual copy number gains.** (**a-b**) Pairwise comparison of the three approaches for timing individual copy number gains and (**c**) comparison with the simulated truth, showing high concordance.

**Extended Data Figure 3:** Validation of League model relative ordering reconstruction based on simulated cohorts of whole-genome samples

**a** Single Timeline (Noisy Reconstruction)

**b** Mixed Timeline (Noisy Reconstruction)

**c** Timing of 100 cohorts with League Model (Simulated Profiles w/ Real Reconstruction from Simulation)

**Validation of League model relative ordering reconstruction based on simulated cohorts of whole-genome samples.** **(a)** League model results on a simulated cohort of samples from a single generalized relative order of events (with varied prevalence) showing high concordance with the true trajectory. **(b)** League model results on a simulated cohort of samples from a complex mixture of trajectories with different order of events showing high concordance with the expected average trajectory. **(c)** Estimation of accuracy of the League model reconstruction by simulation of a set of 100 cohorts with random trajectory mixtures and quantifying the distance in log odds early/late from perfect ordering. For the vast majority of events (even with low number of occurrences in the cohort) the log odds error does not exceed 1, confirming that barely any of the events would switch between timing categories.

**Extended Data Figure 4:** Correlation between League model and Bradley-Terry model ordering

**Correlation between League model and Bradley-Terry model ordering.** Direct comparison for each tumour type of the League and Bradley-Terry models for determining the order of recurrent somatic mutations and copy number events. Axes indicate the ordered events observed in the respective tumour types. Correlation is quantified by Spearman's rank correlation coefficient.

# Extended Data Figure 5: Overview early to late clonal signature changes across tumour types

**Overview of early to late clonal signature changes across tumour types**. Pie charts representing signature changes per cancer type. The colour scheme is as above; signatures that decrease between early and late are coloured green, signatures that increase are purple. The size of each pie chart represents the frequency of each signature. Signatures are split into three categories: "clock-like", comprising the putative clock signatures 1 and 5, "frequent", which are signatures present in 10 or more cancer types, and "cancer type specific", which are in fewer than 10 cancer types and are often limited to specific cohorts.

# Extended Data Fig. 6

**Absolute timing, related to Figure 6**. **(a)** CpG>TpG mutation/Gb (scaled to copy number) as a function of age at diagnosis for 33 informative cancer types. The dotted line denotes the median mutations/yr (ie., not offset), while the shading denotes the 95% credible interval of a hierarchical Bayesian linear regression model. **(b)** Maximum a posteriori estimates of rate and offset for 33 cancer types with 95% credible intervals. **(c)** Median fraction of mutations attributed to linear age-dependent accumulation, based on estimates from b and the age at diagnosis for each sample. **(d)** Scatter plot of number of early (co-amplified) CpG>TpG mutations (y-axis) as a function of the mutational time estimate of WGD (x-axis). The black line denotes a non-linear loess fit with 95% confidence interval. Colours define the cancer type as in part a. **(e)** Total CpG>TpG mutations (y-axis) as a function of the mutation time estimate of WGD (x-axis). Colours and fit as d. **(f)** Inferred median absolute timing patient-specific rate increase, depending on the observed CpG>TpG mutation burden, allowing for a higher (up to 10x) mutation rate increase in cancers with more mutations. **(g)** Comparison of the median duration of subclonal diversification per cancer type assuming branching and linear phylogenies.