Parallel use of a shared genomic island of speciation in clinal and mosaic hybrid zones between cryptic seahorse lineages

Florentine Riquet[1, 2], Cathy Liautard-Haag[1, 2], Lucy Woodall[3, 4], Carmen Bouza[5], Patrick Louisy[6, 7], Bojan Hamer[8], Francisco Otero-Ferrer[9], Philippe Aublanc[10], Vickie Béduneau[11], Olivier Briard[12], Tahani El Ayari[1, 2], Sandra Hochscheid[13], Khalid Belkhir[1, 2], Sophie Arnaud-Haond[1, 14], Pierre-Alexandre Gagnaire[1, 2], Nicolas Bierne[1, 2]

[1] Institut des Sciences de l'Evolution de Montpellier, Université Montpellier, Montpellier, France
[2] CNRS Institut des Sciences de l'Evolution, UMR5554 UM-CNRS-IRD-EPHE, Station Marine OREME, Sète, France
[3] Department of Zoology, University of Oxford, Oxford, OX1 3PS, UK
[4] Natural History Museum, Cromwell Road, London SW7 5BD, UK
[5] Department of Genetics, Faculty of Veterinary Science, Universidade de Santiago de Compostela, Campus de Lugo, Lugo, Spain
[6] University of Nice Sophia Antipolis, ECOMERS Laboratory, Faculty of Sciences, Parc Valrose, Nice, France
[7] Association Peau-Bleue, 46 rue des Escais, Agde, France
[8] Center for Marine Research, Ruder Boskovic Institute, Giordano Paliaga 5, 52210 Rovinj, Croatia
[9] Grupo en Biodiversidad y Conservación, IU-ECOAQUA, Universidad de Las Palmas de Gran Canaria, Crta. Taliarte s/n, 35214 Telde, Spain
[10] Institut océanographique Paul Ricard, Ile des Embiez, Six-Fours-les-Plages, France
[11] Océarium du Croisic, Avenue de Saint Goustan, Le Croisic, France
[12] Aquarium de Biarritz, Biarritz Océan, Plateau de l'Atalaye, Biarritz, France
[13] Stazione Zoologica Anton Dohrn, Department Research Infrastructures for Marine Biological Resources, Aquarium Unit, Napoli, Italy
[14] Ifremer - MARine Biodiversity, Exploitation and Conservation, UMR 9190 IRD-IFREMER-UM-CNRS, Sète, France

Abstract

Diverging semi-isolated lineages either meet in narrow clinal hybrid zones, or have a mosaic distribution associated with environmental variation. Intrinsic reproductive isolation is often emphasized in the former and local adaptation in the latter, although both can contribute to isolation. Rarely these two patterns of spatial distribution are reported in the same study system, while this could provide fundamental information on the endless debate about the relative contribution of intrinsic reproductive isolation and local adaptation on the speciation process. Here we report that the low diversity long-snouted seahorse *Hippocampus guttulatus* is genetically subdivided into five cryptic semi-isolated lineages. Along the Atlantic coasts, northern and southern lineages meet and coexist with little hybridization in the southwest of France, forming a clinal hybrid zone. In the Mediterranean Sea, two lineages have a mosaic distribution associated with lagoon-like and marine habitats. A fifth lineage was identified in the Black Sea. Genetic homogeneity over large spatial scales within each lineage, together with among-locus

variance in differentiation levels between lineages and spatial patterns of introgression
provide evidence that partial reproductive isolation is maintaining the divergence.
Surprisingly, the North Atlantic and the Mediterranean lagoon lineages are genetically
similar for a single chromosome-wide island showing parallel genetic differentiation
among regions. Since Atlantic lineages distribution lacks association with habitat
variation, genetic parallelism suggests that a shared genomic barrier contributes to
reproductive isolation in contrasted contexts -*i.e.* spatial *vs*. ecological. We discuss how a
genomic hotspot of parallel differentiation could have evolved and become associated
either with space or with a patchy environment in a single study system.
56

57 **Keywords:** clinal hybrid zone, mosaic hybrid zone, reproductive isolation, local
58 adaptation, ecological speciation, parallel evolution

Introduction

The spatial context of contact zones between partially isolated taxa and its relationship with environmental variation was long thought to offer great promises to unravel the nature and origin of species. Though each taxa may be genetically homogeneous over large distances, they meet in abrupt genetic discontinuities, called hybrid zones, in which partial reproductive isolation limits gene exchange (Barton and Hewitt 1985, Hewitt 1988). Hybrid zones are extremely informative for exploring the genetic basis of reproductive isolation (e.g. Teeter et al. 2008, Christe et al. 2016) and local adaptation (e.g. Jones et al. 2012, Larson et al. 2013, Soria-Carrasco et al. 2014), as well as identifying genomic regions involved either in increased genomic differentiation (reviewed in Ravinet et al. 2017) or adaptive introgression (Hedrick 2013). The hybrid zone literature usually contrasts two spatial patterns (Harrison 1993): (i) clinal hybrid zones, with parapatrically distributed parental forms on both sides of a cline-shaped genetic divide, and (ii) mosaic hybrid zones, when the environment consists of a mosaic of habitat patches to which taxa (ecotypes, host races, hybridizing species) are somehow specialized. Hybrid zones of both types have been recognized to be multifactorial and maintained both by exogenous and endogenous diverging mechanisms (*i.e.* local adaptation and intrinsic reproductive isolation, respectively; Barton and Hewitt 1985, Bierne et al. 2011). Nonetheless, clinal hybrid zones tend to be interpreted as being mainly maintained by intrinsic reproductive isolation evolved in allopatry before contact (the tension zone model, Barton and Hewitt 1985). Conversely, local adaptation is often emphasized to drive mosaic distributions which are interpreted as evidence that selection at local adaptation genes is acting to increase differentiation between habitats in a parallel fashion (Nosil and Feder 2012). This sketchy dichotomy is anchored by emblematic study systems for which decades of research allowed to support such interpretations. For instance, the hybrid zone between the mice *Mus musculus musculus* and *M. m. domesticus* in central Europe (Boursot et al. 1993) has been well demonstrated to be maintained by selection against hybrid genotypes (Britton-Davidian et al. 2005, Good et al. 2008) after secondary contact (Duvaux et al. 2011). Although the position of the zone was initially found associated with rainfall (Hunt and Selander 1973), local adaptation is not considered as contributing a lot to the isolation. At the other extreme, local adaptation at the same genetic variants has repeatedly allowed marine three-spined sticklebacks to evolve into a freshwater ecotype (Jones et al. 2012). Intrinsic selection is thought absent in the marine-freshwater sticklebacks system, although hybrids perform poorly in both environment and tend to reside in salinity ecotones (Vines et al. 2016). It would be misleading, however, to suggest the two alternative spatial contexts and relations to environmental variation may correspond to alternative routes to speciation (e.g. mutation-order *vs.* ecological speciation; Coyne and Orr 2004, Nosil and Feder 2012). The list is also long of hybrid zones maintained both by local adaptation and intrinsic reproductive isolation. *Bombina* toads (Szymura and Barton 1986), *Gryllus* crickets (Rand and Harrison 1989, Larson et al. 2014), or *Mytilus* mussels (Bierne et al. 2003), among others, are well-known examples of mosaic hybrid zones maintained by both exogenous and endogenous selection.

We here studied the population genetics of the long-snouted seahorse, *Hippocampus guttulatus,* across a large part of its distribution range. We developed an assay of 286 informative SNPs chosen among more than 2,500 SNPs from a population

transcriptomic study (Romiguier et al. 2014). *H. guttulatus* is thought to display poor
dispersal abilities (e.g. site-fidelity, weak swimming performance, lack of dispersive
stage) and inhabits fragmented habitats along its distribution range (from the English
Channel through the Mediterranean and Black Seas; Lourie and Vincent 2004). In
addition, small and patchy populations are often observed in this small coastal fish that is
infrequently seen. Accordingly *H. guttulatus* has a very low genetic diversity (Romiguier
et al. 2014). However, genetic differentiation proved to be very low over very large
geographic distances (Woodall et al. 2015), or across the Cape Finisterre oceanographic
barrier (López et al. 2015). Four well-differentiated genetic clusters, each distributed over
extended regions, were delineated by abrupt genetic discontinuities corresponding to
usual delimitations between vicariant marine lineages (Woodall et al. 2015) – between
the Iberian Peninsula and the Bay of Biscay in the North Eastern Atlantic, between the
Atlantic Ocean and the Mediterranean Sea, and between the Mediterranean and Black
seas. Although the differentiation matches well with oceanographic barriers, this pattern
is also in good accordance with the existence of reproductively isolated cryptic lineages,
which boundaries were trapped by exogenous barriers (*i.e.* the coupling hypothesis;
Bierne et al. 2011).
Our array of informative SNPs and a broader sampling in the Mediterranean Sea
and contact zones, allowed us to describe five cryptic semi-isolated lineages: two
paratrically distributed lineages in the Atlantic Ocean, two patchily distributed lineages
associated with lagoon and marine environments in the Mediterranean Sea, and one in the
Black Sea. Surprisingly, a shared genomic island of clustered outlier loci was involved
both in the isolation between the two parapatric lineages in the Atlantic Ocean and
between the marine and lagoon ecotypes in the Mediterranean Sea, with the North
Atlantic lineage being related to the lagoon ecotype at this genomic island. However, the
two Atlantic lineages inhabit both marine and lagoon habitats. The *H. guttulatus* complex
becomes one of a few systems where a clinal and a mosaic hybrid zone are observed
concomitantly, and a new counter-example that provides evidence of genetic parallelism
in absence of ecological convergence.
134
Materials and Methods
Sampling and DNA extraction
H. guttulatus samples were collected alive from 25 sites (Fig. 1A) using a variety
of methods (snorkeling, scuba diving, trawling nets, aquaria, donations). The dorsal fin of
each individual was clipped using a non-lethal procedure (Woodall et al. 2012), before
releasing back the individual into its natural habitat or aquaria, respectively. Each
individualized sample was preserved in 96% ethanol for subsequent genetic analyses.
Whole genomic DNA was extracted following either Woodall et al. (2015), López
et al. (2015) or using a standard CetylTrimethyl Ammonium Bromide (CTAB)
Chloroform:Isoamyl alcohol (24:1) protocol (Doyle and Doyle 1987). For samples from
Vigo (Spain), the whole genome was amplified using the GenomiPhi kit (GE HealthCare)
according to the manufacturer's protocol. Quality and quantity of DNA extraction was
checked on an agarose gel, and normalized to 35 ng.μL^{-1} using Qubit Fluorometric
Quantitation (Invitrogen). SNP-genotyping was performed (see below for SNP calling)
using the GoldenGate Genotyping Assay with VeraCode technology at ADNid company
(http://www.adnid.fr/index.html, Montpellier, France).

Data mining for SNP markers

A set of 12,613 contigs was examined to identify SNPs. This included one mitochondrial contig (GenBank accession number: AF192664) and 12,612 contigs from Romiguier et al. (2014). Romiguier et al. (2014) sequenced the transcriptomes of six *H. guttulatus* from three locations (Le Croisic, Atlantic Ocean, France; Faro, Atlantic Ocean, Portugal; Thau lagoon, Mediterranean Sea, France), and two *H. hippocampus* from two locations (Sète, Mediterranean Sea, France; Bizerte, Mediterranean Sea, Tunisia) using a combination of high-throughput sequencing methods. Based on these 12,613 contigs, SNPs were identified using the same bioinformatic pipeline as described in Bouchemousse et al. (2016). SNPs were called with Read2SNPs (Gayral et al. 2013) and filtered out according to the following criteria to exclude: 1) SNPs showing more than two alleles, 2) SNPs failing to be sequenced in at least one location, 3) SNPs present only in one individual (*i.e.* singletons), 4) SNPs identified as paralogs using the paraclean option of Read2SNPs, and 5) SNPs closer than 20 bp from a contig extremity or an exon limit when blasted against the stickleback, cod and tilapia genomes. This resulted in 2,684 selected SNPs that were screened with the Illumina Assay Design Tool (ADT) software to select 384 SNPs on the basis of their quality index (ADT score > 0.6).

To identify their chromosomal positions, the template sequences of the targeted SNP were 1) blatted (Bhagwatt et al. 2012) against the genomes of the zebrafish (*Dano rerio*, Howe et al. 2013), fugu (*Takifugu rubripes*, Kai et al. 2011), tetraodon (*Tetraodon nigroviridis*, Jaillon et al. 2004), Nile tilapia (*Oreochromis niloticus*, Brawand et al. 2014), medaka (*Oryzias latipes*, Kasahara et al. 2007), stickleback (*Gasterosteus aculeatus*, Jones et al. 2012), and European seabass (*Dicentrarchus labrax*, Tine et al. 2014), 2) blasted against the genome of the Gulf pipefish (*Syngnathus scovelli*, Small et al. 2016) and 3) blasted against the scaffolds of the tiger tail seahorse (*Hippocampus comes*, Lin et al. 2016), followed by blats and blasts of the scaffolds against the eight well-assembled fish genomes.

All SNPs were oriented using *H. hippocampus* as an outgroup species to identify the most parsimonious ancestral variant. The Joint Site-Frequency Spectrum obtained from the original transcriptome-wide SNP dataset (Romiguier et al. 2014) was compared to the JSFS obtained by the subset of 384 SNPs to investigate potential ascertainment bias due to marker choice. The JSFS represents the bivariate distribution of allelic frequencies in the two populations (Ewens 1972). It is a $(2n_1+1)$ x $(2n_2+1)$ dimension matrix, with n_1 the number of individuals in population 1 and n_2 the number of individuals in population 2, where each entry $S(i,j)$ gives the number of SNPs for which the derived allele was found i and j times in population 1 and 2, respectively. In order to compare the JSFS obtained with the two datasets, the JSFS dimension was here projected down to a 5x5 matrix, which was the dimension of the Rominguier et al. (2014) dataset.

Analyses of Genetic Diversity and Genetic Structure

The mean number of alleles per locus (N_{all}) and the allelic richness (A_r, *i.e.* the expected number of alleles corrected for sampling size based on a rarefaction method) were estimated using the function divBasic implemented in the R package diveRsity (Keenan et al. 2013). Allelic frequencies, expected heterozygosity (H_e) and fixation index (F_{IS}) were estimated using GENEPOP on the web (Raymond and Rousset 1995, Rousset

2008). The raw SNP data was visualized by a color diagram using the software
INTROGRESS (Gompert and Buerkle 2010). Alleles derived from each of the parental
populations were counted at the individual level and converted into a matrix of counts,
used to visualize the multilocus genotype of each individual.
Genetic structure among sampling sites was depicted using Principal Component
Analysis (PCA) computed on the matrix of genotypes, and with an individual-based
Bayesian clustering method. The aim of these two analyses was to identify and describe
genetic clusters using different statistical approaches. The PCA were carried out using the
R package ADEGENET 1.4–2 (Jombart 2008, Jombart et al. 2011). The individual-based
Bayesian clustering analysis was performed with the software STRUCTURE 2.3.4
(Pritchard et al. 2000, Falush et al. 2003). For each value of K (ranging from 1 to 25), 30
replicate chains of 150,000 Monte Carlo Markov Chain (MCMC) iterations were run
after discarding 50,000 burn-in iterations. An admixture model with correlated allele
frequencies was applied with *a priori* information on samples origin. Note that this
method makes the assumption of homogeneous admixture rate in the genome (neutral
admixture) and therefore return a sort of weighted average admixture rate when
introgression is heterogeneous across the genome. To determine individual ancestry
proportions (Q-values) that best matched across all replicate runs, we used CLUMPP
(Jakobsson and Rosenberg 2007) and visualized individuals' assignment under the R
software.
PCAs and STRUCTURE were useful to identify different genetic clusters and to
illustrate the lack of genetic structure among samples that belong to these clusters. Based
on this, individuals were subsequently pooled into clusters with a minimum of seven
individuals per cluster, and genetic structure was computed among clusters by calculating
global and pairwise F_{ST} (Weir and Cockerham 1984) using GENEPOP on the web
(Rousset 2008). Exact tests for population differentiation (10,000 dememorization steps,
500 batches and 5,000 iterations per batch) were carried out to test for differences in
allele frequencies. Q-values, defined as the adjusted p-values using an optimized False
Discovery Rate approach, were computed using the QVALUE package in the R software
(Storey 2002) to correct for multiple testing.
Population trees were generated using the Neighbor Joining algorithm based on
Nei's genetic distances (Nei 1972) in the software PHYLIP 6.695 (Felsentein 2005).
Hossegor (site 6 in Fig. 1A) was removed from the analysis, as in this present site,
individuals from both North and South genetic clusters were observed.
Evolutionary history of genetic clusters was also investigated under a model of
divergence and admixture events using the population graph approach implemented in
the TREEMIX software (Pickrell and Pritchard 2012). This software uses the covariance
matrix of allele frequency between pairs of populations to infer both populations' splits
and gene flow. A maximum likelihood population tree is first generated under the
hypothesis of an absence of migration, and admixture events are sequentially added,
improving (or not) the tree model. This statistical method shows the benefit of
constructing population trees while testing for gene flow between diverged populations at
the same time. Hossegor (site 6 in Fig. 1A) and Chalkida (site 24 in Fig. 1A) were
removed from these analyses, the former showing individuals from both North and South
genetic clusters in the same site, the latter generating noise, due to a small sample size
(Pickrell and Pritchard 2012). Using the total data set, *i.e.* 286 loci, five migration events

were sequentially added to look for the best tree to fit the data and the number of
migration events that reached an asymptotic likelihood was kept.
245
Outlier detection
We first measured the genome-wide distribution (measured by the Lewontin-
Krakaeur statistic) of genetic differentiation and the neutral expectation to investigate any
departure from neutrality. The Lewontin-Krakaeur statistic was measured for each locus
with LK=(d-1)(K-1)$Fst/\overline{Fst}$, where d=2 is the number of populations and K=2 is number
of alleles. We then fitted a Generalized Extreme Value Distribution using the Broyden-
Fletcher-Goldfarb-Shanno algorithm to visualize highly differentiated loci between
populations.
The use of several independent methods is recommended to improve accuracy of
outlier loci detection (de Villemereuil et al. 2014). Each method is differently impacted
by the genetic structure and/or demographic history of the study species, which is usually
unknown, leading to frequent inconsistencies across methods (Gagnaire et al. 2015). To
cope with these problems, outlier loci were detected using two different methods. First,
BAYESCAN, which implements a method developed by Foll and Gaggiotti (2008), is a
Bayesian method that uses a logistic regression model to estimate directly the posterior
probability that a given locus is under selection. Briefly, locus population F_{ST} values are
decomposed as a linear combination of a locus specific component α and a population
specific component β. A locus is assumed to be under selection when α is necessary to
explain the observed pattern of diversity. For each locus, a reversible jump MCMC
explores two alternative models (with or without α) and estimates their relative posterior
probabilities. We used default parameter values in our analyses to detect outliers among
the clusters previously identified. The second approach used was the one developed by
Duforet-Frebourg et al. (2014) and implemented in the R package PCAdapt (Luu et al.
2016). Briefly, it is based on a hierarchical model where population structure is first
depicted using K factors. No *a priori* on the genetic structure (and thus, no clustering) is
here required in advance. Loci that are atypically related to population structure,
measured by the K factors, are identified as outliers. For each value of K (ranging from 1
to 10), 10 replicate chains of 150,000 Monte Carlo Markov Chain iterations were
performed and we discarded the first 5,000 iterations as burn-in. PCA, Structure, and
Treemix analyses were repeated on sub-datasets including 1) only outlier loci (*i.e.* 14
loci), and 2) only non-outlier loci (*i.e.* 272 loci). Identical parameters as described above
were used.
We tested for genotype discordance among loci in an admixed sample by
conducting a genomic cline analysis, using Barton's concordance method as described in
Macholán et al. (2011). Basically, the method fits a quadratic function to the relationship
between a single locus hybrid index (0 or 1 if homozygous, 0.5 if heterozygous) and the
genome-wide hybrid index. The function parameters measure the deviation from x = y
(*i.e.* the expectation of homogeneous genome-wide introgression) as a function of the
expected heterozygosity. More than testing for the deviation from the diagonal, our aim
here was instead to compare genomic clines (*i.e.* regression curves) between geographic
samples. The reason was that discordance was observed in one population and not others.
Finally, we tested for gene ontology (GO) terms enrichment to determine if
outlier loci displayed functional enrichment, compared to the full dataset, using Fisher's

Exact Test with Multiple Testing Correction of FDR (Benjamini and Hochberg 1995)
implemented in the software Blast2GO (Conesa and Götz 2008).
291
Results
SNPs characterization/calling
Out of the 384 SNPs, including five SNPs (one mitochondrial and four nuclear)
that diagnostically distinguished *H. guttulatus* from *H. hippocampus*, a total of 318 SNPs
successfully amplified. Four SNPs were monomorphic resulting in 314 SNPs examined
for 292 *H. guttulatus*, among which 286 showed a minor allele frequency above 5%.
Note that the markers diagnostic of *H. hippocampus* specimens were used to control the
taxonomic status of each fish, but being monomorphic in *H. guttulatus*, were removed
from the final dataset.
In order to evaluate the extent of ascertainment bias that may be induced by our
procedure for selecting markers, we compared the Joint Site-Frequency Spectra (JSFS)
obtained with the original dataset of Romiguier et al. (2014), which is assumed freed
from ascertainment bias, and our 314-SNP dataset. JSFSs are shown in Supplementary
Information 1 (Fig. SI1). As expected, the data of Romiguier et al. (2014) is principally
composed of singletons that represented 35-40% of the SNPs (Fig. SI1A), while this
proportion was reduced twofold in our 314-SNP dataset as desired (Fig. SI1B). We
efficiently removed rare variants without biasing the frequency spectrum too much as the
deficit of singletons was homogeneously compensated by every cell of the JSFSs: the
comparison of the two JSFSs (Fig. SI1C) reveals that very few cells apart from singletons
have an excess above 5%, suggesting limited ascertainment bias in our SNP panel, except
with rare alleles as expected.
313
A strong genetic structure delineating five clusters
Estimates of mean number of alleles per locus (N_{all}), allelic richness (A_r),
expected heterozygosity (H_e), and departure from Hardy-Weinberg equilibrium (F_{IS}), for
each study site and each genetic cluster identified, are presented in Table SI1. The gene
diversity was globally similar among populations and no significant departure from
Hardy-Weinberg equilibrium (HWE) was observed.
The Principal Component Analysis (PCA) revealed a strong genetic structure
separating five clusters (Fig. 2A). Along the first two axes (Fig. 2A, left panel),
explaining 60.1% and 16.2% of the variance, the PCA showed clear differentiation
between North Atlantic (sites 1-5, in red), South Atlantic (sites 7-12, in blue),
Mediterranean Sea (sites 13-25, site 16 excluded, in green) and Thau lagoon (site 16, in
pink). Hossegor individuals (site 6, in purple) clustered either with the North Atlantic (12
individuals) or with the South Atlantic groups (3 individuals). Mediterranean sites spread
out along the third axis (9.9% of the variance explained, Fig. 2A, right panel), with
Bizerte (site 20, in gold) being closer to the Thau lagoon on the one hand, and Varna (site
25, in turquoise) standing out from the Mediterranean cloud on the other hand. This
pattern of genetic structure was also detected by the STRUCTURE analysis (Fig. 2B).
Five clusters were identified, composed by: 1- the North Atlantic sites, 2- the South
Atlantic sites, 3- the Mediterranean sites, 4- Thau lagoon and 5- Varna. A gradient of
introgression is visible along the Atlantic coasts, with a decreasing proportion of south
Atlantic cluster ancestry north to Hossegor (site 6), suggesting asymmetrical gene flow.

While most individuals from Hossegor shared the North Atlantic genetic background,
three individuals were genetically assigned to the South Atlantic genetic background.
Bizerte (site 20), and to a lesser extent Chalkida (site 24), proved to have mixed
ancestries from Thau lagoon and Mediterranean clusters.
Representing the matrix of raw genotyping data (Fig. 2C) also clearly pointed out
the delineations between the five detected clusters (highlighted by vertical dotted lines in
Fig. 2C). The deepest node of Neighbor Joining tree (Fig. 1D) separated two groups, one
composed by the North Atlantic Ocean sites only, and the second by South Atlantic and
all Mediterranean Sea sites. Concerning the Mediterranean Sea group, Thau and Bizerte
lagoons clustered together, and with Varna (site 25). Mediterranean group and Chalkida
clustered together, also separated by one internal node to Varna.
No significant departure from panmixia was observed within each cluster (Table
SI1). Further, genetic homogeneity was observed between sites within each cluster (Table
SI2). In contrast, Fisher exact tests revealed significant differences in allelic frequencies
among clusters $(p$-value < 0.001), with significant differentiation being observed for all
pairwise comparisons (F_{ST} ranging from 0.025 between the Mediterranean cluster and
Chalkida to 0.257 between the North Atlantic cluster and Varna).
352
Signature of selection, genetic parallelism and reticulated evolution
In all pairwise comparisons between clusters the genome-wide distribution of
genetic differentiation (as measured by the LK statistic) departed from the neutral
expectation. This is illustrated in Figure 3 for the comparison between the North Atlantic
and South Atlantic clusters and holds for all other pairwise comparisons of clusters (data
not shown).
Fourteen loci out of 286 (4.9%) were consistently identified to depart from
neutrality in both BayeScan and PCAdapt analyses, using a 5% False Discovery Rate
(FDR). These outlier loci were located on different *H. guttulatus* contigs, but six to eight
of them were located on a unique chromosome when mapped to one of the seven well-
assembled fish genomes -zebrafish aside-, of which scaffolds were placed (Fig. 4). Eight
of these outlier loci were consistently found in this unique chromosome in, at least, four
of the seven fish genomes. Furthermore, four of these outlier loci (SNPs 232, 312, 87 and
311) mapped to this chromosome for the seven fish genomes (colored in orange and pink
in Fig. 4). SNP 29 was also consistently located on this chromosome, except in the Gulf
pipefish, for which it mapped to an unplaced scaffold. However, there was little
consistency in their position or order along the chromosome among species (Fig. 4).
Interestingly, two non-outlier loci also mapped to this unique chromosome in, at least,
two of the seven fish genomes (Fig. 4). When blasted against the tiger tail seahorse
scaffolds, the fourteen outlier loci were located on eleven scaffolds. Two SNPs (SNPs 29
and 286) mapped to *H. comes* scaffold 8. Similarly, three SNPs (SNPs 87, 212 and 311)
mapped to scaffold 154. Out of the eleven scaffolds involving the outliers, three scaffolds
(scaffolds 8, 339, 397) consistently mapped to the unique chromosome identified here
above, when blatted and basted against the seven well-assembled fish genomes, though
with some exceptions. *H. comes* scaffold 8 was not identified in this chromosome only
for the Nile tilapia, while scaffolds 339 and 397 were not identified in the medaka and
zebrafish and the Nile tilapia and zebrafish, respectively. *H. comes* blasts and blats
suggested that SNPs 87, 311, 212, 312 and 232 also mapped to a unique chromosome in

this species. Similarly, SNP 29, the most differentiated locus (Fig.5C), mapped to the
unique chromosome in the European seabass and the stickleback, though to an unplaced
scaffold in the other well-assembled fish genomes, suggesting that this SNP (and SNP
286 blasted on the same *H. comes* scaffold) may also be located on this chromosome. A
similar pattern holds for SNP 34. Note that four outliers were also found on the same
chromosome in the genome of the zebrafish, which is the most divergent species used as
an outgroup (Fig. 4). The analysis of gene ontology terms for outlier loci did not reveal
any significant functional enrichment.
Interestingly, the visualization of raw allele frequency data at individual SNPs
(Fig. 2C) nicely revealed how the pattern observed at outlier loci departed from the one
observed at other loci, not only in the level of differentiation but also in reinforcing the
previously observed groupings of populations. Outliers were classified into five
categories based on their allele frequency patterns (Fig. 5): those distinguishing 1- North
Atlantic sites, Thau lagoon, Bizerte and, in a lesser extent Chalkida and Varna, from
South Atlantic and Mediterranean sites (Fig. 5A) or 2- only from South Atlantic (Fig.
5B); 3- those separating Thau lagoon (and in a lesser extent Bizerte, Chalkida and Varna)
from all other sites (Fig. 5C), 4- those differentiating North Atlantic sites from all other
sites, Thau lagoon included (Fig. 5E); and 5- the SNPs distinguishing Mediterranean sites
vs. Atlantic sites (Fig. 5F). Note that Hossegor and Chalkida individuals were both split
into 2 groups according to their North or South Atlantic / Thau lagoon or Mediterranean
genetic background, respectively. The first and most abundant category of outliers
pointed out the convergence of allele frequency patterns (*i.e.* genetic parallelism)
between North Atlantic and Thau lagoon individuals. Interestingly, most markers of this
category were found to map to a single chromosome in our cross-mapping approach
using different fish genomes, along with the two markers highly differentiated between
lagoon and sea ecotypes (Fig. 4 and 5). Contrasting with a consistent pattern in allelic
frequencies in each of the five outlier categories, the non-outlier loci that mapped to a
unique chromosome did not show any consistent pattern (Fig. 5D). Note that SNP 242
(Fig. 5B) that is an outlier of differentiation between the Atlantic Ocean and the
Mediterranean Sea mapped also to this unique chromosome but only in the Gulf pipefish
(Fig. 4).
In order to visualize parallelism in the extent of genetic differentiation in the
whole SNP dataset, allele frequency differences (AFDs) between Mediterranean lagoon
and sea locations (x-axis in Fig. 5G) were plotted against AFDs between North and South
Atlantic clusters (y-axis in Fig. 5G). As expected, outliers showing genetic parallelism
appear in the top right of Figure 5G, showing both elevated genetic differentiation
between Mediterranean lagoon and marine sites (x-axis) and between north and south
Atlantic sites (y-axis). The third category of outliers (Fig. 5C), also observed along the
same chromosome when compared against the other fishes (Fig. 4), revealed a higher
differentiation only between habitats in the Mediterranean Sea. The fifth category (Fig.
5E) revealed a higher genetic differentiation only between Atlantic lineages.
Figure 6 shows the result of the genomic cline analysis obtained with SNP 29, one
of the outliers that clustered on the same chromosome, and SNP 130, an outlier between
the northern and the southern lineage but that maps to another chromosome and which is
not differentiated in the Mediterranean Sea (no genetic parallelism). Similar results were
obtained with other loci from the same categories. We found that the regressions were

different at SNP 29, or other outliers that clustered on the same chromosome, when we
included the Hossegor sample in the analysis or not (Fig. 6), while regressions were
always close to the diagonal with SNP 130, or other outliers that maps to other
chromosomes. This analysis demonstrates the discordance between the clustered outlier
loci and the rest of the genome in the Hossegor sample, in which north Atlantic-like
seahorses have the south Atlantic allele at clustered outlier.

The PCA performed with the fourteen outliers differed from the PCA based on
286 SNPs. Bizerte, Varna, the North Atlantic sites and Chalkida (Greece) were found
closer to lagoon along the first axis (59.5% of the variance explained; Fig. SI2A).
STRUCTURE analysis identified three clusters (data not shown), composed by 1- the
North Atlantic sites, 2- the South Atlantic and Mediterranean sites, and 3- Thau lagoon,
Bizerte and, on a lesser extent Varna that have mixed ancestries from both North Atlantic
and Thau lagoon while Chalkida revealed ancestries of Thau lagoon and the
Mediterranean clusters. Genetic affinities between the North Atlantic sites and Thau
lagoon also emerged in the Neighbor Joining tree (Fig. 1C): Thau and Bizerte lagoons
branched together, and with the North Atlantic populations to form a group composed of
Mediterranean lagoons and North Atlantic populations.

In contrast, the PCA based on 272 neutral SNPs showed a very similar pattern to
the PCA obtained with 286 SNPs along the three first axes (10.2%, 6.0% and 2.3% of the
variance explained; Fig. SI2B). STRUCTURE analysis identified four clusters (data not
shown), with Varna clustering with the Mediterranean sites. Two groups of populations
emerged based on 272 neutral SNPs in the Neighbor Joining tree (Fig. 1B), one
composed by the Atlantic Ocean samples, the second by all Mediterranean Sea samples.
Neutral SNPs seemed reflecting geography, although Isolation By Distance was not
significant (Mantel test *p*-value = 0.23).

Finally, the population tree inferred using TREEMIX without accounting for
migration (Fig. 2D, left panel) was highly consistent with all above analyses, with
Atlantic clusters branching together on one hand, and Thau and Bizerte lagoons
branching together on the other hand. Interestingly, three admixture events significantly
improved the model as compared to a situation without migration (*p*-value < 0.001; Fig.
2D, right panel). This population tree indicated significant gene flow among four *H.*
guttulatus clusters, between the north and the south genetic clusters in the Atlantic coasts,
in agreement with the gradient of introgression observed in Figure 2B, between marine
and lagoon samples in the Mediterranean Sea, and finally between the North Atlantic and
Mediterranean lagoon samples. We suggest that this latter observation is most probably
mainly driven by parallel outlier loci (see discussion below).

Discussion
Genetic analyses of the long-snouted seahorse revealed that this species is
subdivided into several cryptic semi-isolated lineages. Strong genetic subdivisions
delineate genetic clusters that otherwise showed a remarkable genetic homogeneity over
large areas, despite limited dispersal abilities and seemingly small population sizes (but
see Curtis and Vincent 2006 in Ria Formosa, Portugal). The spatial organization of the
different genetic backgrounds proved to be an unusual combination of geographic
subdivision in the Atlantic Ocean, and genetic structure related to the sea-lagoon
ecological contrast in the Mediterranean Sea. Surprisingly, these two subdivisions partly

relied on the same genetic architecture. This was supported by the finding of genetic parallelism at some markers showing extreme levels of differentiation between habitats in the Mediterranean Sea, being also amongst the most differentiated between geographic lineages in the Atlantic Ocean. Intriguingly, these loci showing convergent allele frequency patterns all mapped to a single chromosome in seven well-assembled fish genomes. Our results thus suggest the existence of a shared evolutionary history of speciation between Atlantic parapatric lineages and Mediterranean ecotypes. Namely, the Mediterranean lagoon ecotype was anciently related to the north-Atlantic lineage, whereas the south-Atlantic lineage shared ancestral variation with the Mediterranean marine ecotype. The underlying reproductive isolation mechanisms may thus involve a combination of intrinsic and extrinsic genetic barriers, where relative contributions may differ between the two contexts. We propose that the genomic island (*i.e.* the part of the genome gathering outlier loci in a unique chromosome) shared in the two alternative spatial contexts is mainly involved in intrinsic reproductive isolation and have coupled with adaptation to a mosaic environment in the Mediterranean Sea whereas it would have been trapped by an extrinsic barrier to dispersal or have coupled with an alternative environmental heterogeneity in the Atlantic Ocean.

1- Efficient genetic analysis in a species with low polymorphism

The long-snouted seahorse has a very low genetic diversity, ranking among the animal species with the lowest synonymous nucleotide diversity ($\pi s = 0.0016$, Romiguier et al. 2014). For comparison, its level of genetic diversity is similar to that observed in less diverse primates, including humans (Perry et al. 2012), or in the giant Galapagos tortoise (Loire et al. 2013). Although RADseq (Baird et al. 2008), RNAseq (Gayral et al. 2013) or targeted capture (Jones and Good 2016) provide a tool of choice in non-model organisms for genetic analyses, this reduced genome representation typically retrieves only a few thousands of SNPs in low-diversity species (~4,000 SNPs in *H. guttulatus* with RADseq, unpublished data). Most of these markers are rare variants providing relatively little information. The strategy implemented here relied on a selection of SNPs discovered from six transcriptomes (Romiguier et al. 2014), ending up with a dataset of 286 informative SNPs. The allele frequency distributions of the newly developed SNPs were not significantly different from those derived from the original SNP dataset, suggesting a limited ascertainment bias with rare alleles being underrepresented. Selecting SNPs characterized from a preliminary population transcriptomic survey therefore proved to be a straightforward strategy for genome-wide investigation of the spatial distribution of well-characterized genetic clusters in this protected species. Since our genotyping method only requires a small amount of DNA, it allows the use of non-lethal fin-clipping sampling techniques (Woodall et al. 2012). Moreover, the approach generated no missing data compared to classical genotyping-by-sequencing approaches.

2- Genome scans in hybrid zones

Our SNP panel allowed us to define five strongly differentiated panmictic genetic clusters, two in the Atlantic Ocean, two in the Mediterranean Sea, and one in the Black Sea, which cannot be morphologically distinguished with reliability so far (*i.e.* cryptic lineages). The average genetic differentiation between clusters and its associated variance were strong ($0.1 \pm 0.02 < F_{ST} < 0.25 \pm 0.04$). In this context, identifying outlier loci is a

complex task with problems of false positives on the one hand (Lotterhos and Whitlock
2014, 2015). Indeed, the neutral model can easily underestimate the variance of
differentiation. On the other hand, it may result in misinterpretations, because many loci
of various sorts interact together to generate a semi-permeable barrier to gene flow
(Bierne et al. 2011, 2013). Although we used recent outlier tests designed to account for
complex demography and spatial population structure, we are likely to deviate from an
implicit but basic assumption of outlier test approaches - that most loci are evolving
neutrally without being affected by the indirect effect of selection through linkage
(Bierne et al. 2013). However, genomic data increasingly support the old prediction that
genetic barriers to gene flow can affect a substantial proportion of the genome (Harrison
and Larson 2016). When explicitly tested with inferential methods that account for
among-locus variation in effective migration rates, semi-permeable barriers to gene flow
prove to be widespread (Roux et al. 2016). It is also increasingly acknowledged that
signal of local adaptation should not be easily captured without a broad sampling of the
genome (Hoban et al. 2016), suggesting that the outliers easily identified with a limited
number of loci could reflect the existence of a cryptic genetic barrier to gene flow
involving many selected loci (Bierne et al. 2011). However, the most differentiated loci
can still provide useful information, if we simply consider them as the best available
markers of differentiated genetic backgrounds.
For the seahorse as for many other species however, the genetic structure
displayed by outlier loci strongly departs from the average genetic structure, and shows a
signature of genetic parallelism (Fig. 2A –C, and Fig. SI2). Therefore, the signal of
outlier loci (especially the A category in Fig. 5A - B) is not just consistent with what is
expected for the most differentiated loci of a neutral distribution. Instead, they display a
genetic structure that is not compatible with the sample covariance matrix. Besides, when
compared to well-assembled fish genomes, these outlier loci mapped to a single
chromosome in all species (Fig. 4). Although interchromosomal rearrangements are
relatively uncommon in teleosts (Naruse et al. 2004, Small et al. 2016), the order was not
so well conserved among species. Nonetheless this result suggests the existence of a large
genomic island of differentiation as already reported in other fishes (e.g. in sticklebacks:
Jones et al. 2012, in cod: Hemmer-Hansen et al. 2013, in seabass: Tine et al. 2014), and
possibly involving a chromosomal inversion or another form of recombination
suppression.
552
3- Parapatrically distributed semi-isolated cryptic lineages in the Atlantic Ocean
In the Atlantic Ocean we confirmed the existence of two well-differentiated
genetic clusters previously identified by Woodall et al. (2015) with five microsatellites
and two mitochondrial genes. Our broader sampling of the genome allowed us to capture
genome regions with very strong genetic differentiation. The most differentiated SNP is
nearly fixed between the two lineages (SNP 130 in Fig. 5, $F_{ST} = 0.9$), and many other
SNPs demonstrate an abrupt genetic shift between the two clusters (Fig. 2C and 5).
Another crucial new observation was the co-existence of the two lineages in the same
lagoon in South West of France (Hossegor lagoon, sample 6 in Fig. 1), with an absence
of early generation hybrids in the sample but some indirect evidence of hybridization and
introgression (Fig. 2B). Such a zone of co-existence, with two genetic backgrounds in
sympatry in the absence of genetic differentiation at a large spatial scale within each

lineage, suggests strong reproductive isolation (Jiggins and Mallet 2000). This may either
be due to pre-zygotic isolation or strong selection against hybrids at early life stages.
Nonetheless, and although early generation hybrids were not detected in our sample, the
decreasing proportion of south lineage ancestry from Hossegor to the English Channel
provides evidence for asymmetric introgression (Fig. 2B). The signal was genome-wide
(Fig. 2C) and was also detected by the TREEMIX analysis.
However, an interesting pattern was observed at the genomic island outliers.
North Atlantic-type individuals sampled at Hossegor hold a different genetic signal at the
genomic island loci. The frequency of the southern lineage allele was similar between
North Atlantic-type individuals from Hossegor and South Atlantic samples (Hossegor
pale pink dot (site 6A) in Fig. 5, see also Fig. 2C). Figure 6 provides a statistical test of
this discordance and evidence for hybridization, as it requires recombination between
parental backgrounds to produce North Atlantic-type individuals with the southern allele
at the genomic island. Genomic cline analysis is usually performed to test whether the
quadratic regression of a locus hybrid index to the genome-wide hybrid index deviates
from the diagonal, which is the expectation when introgression occurs at the same rate for
all loci (Macholán et al. 2011). Here we included the geographic information to contrast
the results obtained when the Hossegor sample is included or not in the analysis, as our
objective was to infer the existence of a local discordance, in the Hossegor lagoon only.
Figures 6 shows that North-Atlantic seahorses, with a high genome-wide hybrid index,
are South Atlantic-type at the genomic island in Hossegor (low hybrid index at SNP 29),
but are North Atlantic-type elsewhere (high hybrid index at SNP 130). This result is also
an indirect evidence for epistatic or coupling interactions among loci implied in
reproductive isolation. Endogenous post-zygotic selection against hybrids does not result
in a stable polymorphism in a single isolated population. Also known as bistable variants,
the maintenance of underdominant or epistatically interacting genetic incompatibities
requires a migration-selection balance in a spatially subdivided population (Barton and
Turelli 2011), or frequency-dependent selection (Barton and de Cara 2009). We can
hypothesize that the system has fixed one state of a bistable variant in the Hossegor
lagoon. One may alternatively suggest that the southern allele confers a selective
advantage in the Hossegor lagoon environment whatever the genetic background,
however this hypothesis does not seem coherent with the fact that the northern allele is
always in higher frequency in lagoon-like habitats elsewhere.
Overall, the Atlantic contact zone possesses all the characteristics of standard
clinal hybrid zones that follow the tension zone model (Barton and Hewitt 1985); *i.e.* a
secondary contact zone maintained by a balance between migration and selection against
hybrids due to intrinsic reproductive isolation. Exogenous selection is nonetheless likely
to contribute, as the environments are quite obviously different between the two
distribution ranges. However, the two Atlantic lineages both inhabit indifferently lagoon
and sea habitats (Fig. 1A) suggesting that this ecological contrast plays little role in their
genetic. In any case, only strong intrinsic reproductive isolation can guarantee a genome-
wide barrier to gene flow explaining the co-existence of the two lineages in sympatry in
the Hossegor lagoon.
608
4- Sea and lagoon ecotypes in the Mediterranean Sea

Contrasting with the Atlantic hybrid zone, the two cryptic lineages identified in
the Mediterranean Sea were associated with lagoon/sea habitat variation. Our broader
genomic and spatial sampling revealed two crucial observations: (i) although we only
obtained samples from two lagoons (Thau in France and Bizerte in Tunisia, Fig. 1A),
this was sufficient to reveal an association with the environment that was previously
misinterpreted as possible spatial differentiation, because the Thau lagoon was the only
sample from Western Mediterranean basin in Woodall et al. (2015), and (ii) we observed
fixed differences between lagoon and marine samples that are only few kilometers apart
in the Gulf of Lion (Fig. 2 and 5). While the marine lineage was surprisingly
homogeneous over the whole Mediterranean Sea, from Greece to Spain (F_{ST} = 0.0078
n.s.), lagoon-like samples showed different genetic compositions. The Thau lagoon
sample showed a strong and genome-wide genetic differentiation from sea samples (Fig.
2). In addition, a single but important seahorse was sampled on the seaside of the Thau
lagoon (site 15 in Fig. 1A), and seven others sampled on the seaside of another lagoon of
the region (Leucate, site 14 in Fig. 1A). They all proved to belong to the marine genetic
cluster, without any sign of introgression of the lagoon background. This result suggests
strong reproductive isolation as well as strong habitat specialization of the two lineages
around French Mediterranean lagoons. However, the sample from the Bizerte lagoon
(Tunisia, sample 20 in Fig. 1), which is ecologically similar to the Thau lagoon (Sakka
Hlaili et al. 2008), is genetically similar to marine Mediterranean samples at most loci
and only share the genetic composition of the Thau lagoon at the genomic island and
other sea-lagoon outlier loci (Fig. 2, 5). In addition, a subsample of the Chalkida
population (Greece, sample 24B in Fig. 5) was composed of five individuals
heterozygous at the genomic island. The environmental condition of the Chalkida
location is hypothesized to be more lagoon-like, and this sample only provides evidence
that the genomic island polymorphism and the mosaic spatial structure extends to the
eastern basin without really providing further clues about the role of the environment.
Our results in the Mediterranean Sea (*i.e.* lagoon/maritime system) resemble those
obtained in the emblematic three-spined sticklebacks marine/freshwater system (Jones et
al. 2012), or more recently in marine migratory and freshwater resident lampreys
(Rougemont et al. 2017) or coastal-marine ecotypes in European anchovies (Le Moan et
al. 2016). When a shared divergent genomic island is observed –*i.e.* genetic parallelism,
as here in long-snouted seahorse and in the examples cited above, there could be three
possible interpretations: (i) parallel gene reuse from a shared ancestral polymorphism
present in the marine supposedly ancestral population (Jones et al. 2012), (ii) the spread
of a locally adapted allele (*i.e.* the 'transporter hypothesis'; Schluter and Conte 2009) or
(iii) secondary contact followed by spatial reassortment of the divergent lineages and
extensive introgression swamping such that only selected loci and their chromosomal
neighborhood retain the history of adaptation (Bierne et al. 2013). The three scenarios
are difficult to discriminate as they converge toward a similar pattern (Johannesson et al.
2010, Bierne et al. 2013, Welch and Jiggins 2014). In addition, the debate can also be
understood as a semantic dispute between a phylogenetic view of populations defined by
the genome-wide majority tree and the genetic view of lineages defined by the tree of
adaptive/speciation genes (Mallet et al. 2016). The Bizerte lagoon population can either
be interpreted as a marine background (defined by the majority tree) introgressed by the
lagoon allele at the genomic island, or as a lagoon background (defined by

adaptive/speciation genes) massively introgressed by neutral marine alleles.
Incorporating heterogeneous migration rates in demographic inference methods allowed
Le Moan et al. (2016), Rougeux et al. (2016) and Rougemont et al. (2017) to identify the
signal of a secondary contact history carried by islands of differentiation in lampreys,
white fishes and anchovies. Unfortunately our 286-SNPs dataset does not allow
performing such historical demographic reconstruction, although the TREEMIX analysis
reveals that episodes of secondary admixtures strongly improve the fit to the sample
covariance matrix. Anyhow, demographic reconstruction does not completely refute the
'transporter hypothesis' which stipulates freshwater alleles spread from lagoon to lagoon
(or river to river) and is a scenario that produces a very similar genomic pattern of
differentiation to the one produced by a standard secondary contact (Bierne et al. 2013,
Rougemont et al. 2016). In the case of the seahorse system, however, we made the new
observation that genetic parallelism is observed despite an apparent absence of a
common selective pressure, which offers complementary arguments to the debate.
670
5- Genetic parallelism in two different spatial/ecological contexts
Now that we have discussed the cryptic differentiation observed in the Atlantic
Ocean, on the one hand, and the Mediterranean Sea, on the other hand, we address
potential links stemming from a shared history between Atlantic and Mediterranean
lineages, underlying genetic parallelism. Clearly, the most astonishing result of our
genetic analysis was genetic parallelism between the Mediterranean lagoon ecotype and
the north Atlantic lineage at a large genomic island. Parallel evolution usually implies
distinct ecological characteristics (e.g. Butlin et al. 2014). In the present study, we found
that the genomic island was associated with the sea-lagoon ecological contrast in the
Mediterranean Sea, while there was no such genetic differentiation between lagoon and
sea samples in the Atlantic Ocean. In addition we found evidence from a genomic cline
analysis that the genomic island could be involved in intrinsic reproductive isolation in
the Hossegor lagoon hybrid zone. Shared ancestral polymorphism sieved by adaptation in
a patchy environment (Bierne et al. 2013) and incipient speciation (Guerrero and Hahn
2017 BioRxiv) would better explain our data. The association with habitat in the
Mediterranean Sea and with space in the Atlantic Ocean could be explained either by (i)
the secondary evolution of locally adapted genes within the genomic island in the
Mediterranean Sea, benefiting from the barrier to gene flow imposed by intrinsic
selection (divergence hitchhiking; Via 2012, Yeaman 2013) or (ii) alternatively the island
could have coupled with local adaptation polymorphism localized elsewhere in the
genome in the Mediterranean Sea (Bierne et al. 2011) while it would have been trapped
by a barrier to dispersal in the Atlantic Ocean (Barton 1979, Barton and Hewitt 1985).
Without further data and the true genomic position of loci in the seahorse genome, it is
difficult to disentangle the two hypotheses.
Two outlier loci (SNP 87 and SNP 311, Fig. 5C) localized in the same
chromosome as convergent SNPs (Fig. 4) are differentiated between the marine and
lagoon ecotypes in the Mediterranean Sea, but are not differentiated between the northern
and southern lineages in the Atlantic Ocean. At first sight, they could be interpreted as
evidence for secondary local sweeps in Mediterranean lagoons. However, as the position
is not well inferred from available genomes and the barrier to gene flow is less effective
in the Atlantic, they could also be interpreted as being localized in the island "shoulders"

(*i.e.* loci in the vicinity of the regions harboring local adaptation and/or reproductive isolation loci; Gagnaire et al. 2015, Le Moan et al. 2016) in which a stronger introgression rate has erased the differentiation faster in the Atlantic than in the Mediterranean. According to the gene order inferred from the closest species (Gulf pipefish and Tiger tail seahorse), these two loci could potentially be in an island shoulder (Fig. 4C). Importantly, whatever the explanation -divergence hitchhiking or coupling- it requires invoking interaction between intrinsic and ecological selection and not ecological selection alone (Bierne et al. 2011, Kulmuni and Westram 2017). Furthermore, intrinsic selection has most probably evolved first in this system as no genetic parallelism was observed in outliers discriminating lagoon to marine ecotypes. This contradicts the predominant view that ecological selection is necessarily the initial catalyzer of the chain of accumulation of barriers in ecological speciation.

Acknowledgments

We are grateful to Lucas Beranger, Michel Cantou, Philippe Lenfant, Pablo Liger, CPIE Bassin de Thau, Patrick Lelong, Francesco Di Liello and Stéphane Auffret for their help in providing *H. guttulatus* fin-clippings and to Fabienne Moreau for the BeadXpress experiment. Many thanks to Nicolas Duforet-Frebourg for computational advices. This work was funded by a Languedoc-Roussillon Region "Chercheur(se)s d'avenir" grant to NB (Connect7 project), by a LabEx CeMEB postdoctoral fellowship to FR and by Chocolaterie Guylian and a Natural Environment Research Council Industrial Case studentship (NER/S/C/2005/13461) to LW.

Literature

Baird, N. A., P. D. Etter, T. S. Atwood, M. C. Currey, A. L. Shiver, Z. A. Lewis, E. U. Selker, W. A. Cresko, and E. A. Johnson. 2008. Rapid SNP Discovery and Genetic Mapping Using Sequenced RAD Markers. PLOS ONE 3:e3376

Barton, N. 1979. Heredity - Abstract of article: The dynamics of hybrid zones. Heredity 43:341–359.

Barton, N. H., and M. A. R. de Cara. 2009. The evolution of strong reproductive isolation. Evolution 63:1171–1190.

Barton, N. H., and G. M. Hewitt. 1985. Analysis of Hybrid Zones. Annu. Rev. Ecol. Syst. 16:113–148.

Barton, N. H., and M. Turelli. 2011. Spatial Waves of Advance with Bistable Dynamics: Cytoplasmic and Genetic Analogues of Allee Effects. Am. Nat. 178:E48–E75.

Bhagwat, M., L. Young, and R. R. Robison. 2012. Using BLAT to Find Sequence Similarity in Closely Related Genomes. Curr. Protoc. Bioinforma. Ed. Board Andreas Baxevanis Al 0 10:Unit10.8

748 Benjamini, Y., and Y. Hochberg. 1995. Controlling the False Discovery Rate: A
749 Practical and Powerful Approach to Multiple Testing. J. R. Stat. Soc. Ser. B Methodol.
750 57:289–300.
751
752 Bierne, N., P. Borsa, C. Daguin, D. Jollivet, F. Viard, F. Bonhomme, and P. David. 2003.
753 Introgression patterns in the mosaic hybrid zone between Mytilus edulis and M.
754 galloprovincialis. Mol. Ecol. 12:447–461.
755
756 Bierne, N., D. Roze, and J. J. Welch. 2013. Pervasive selection or is it...? why are FST
757 outliers sometimes so frequent? Mol. Ecol. 22:2061–2064.
758
759 Bierne, N., J. Welch, E. Loire, F. Bonhomme, and P. David. 2011. The coupling
760 hypothesis: why genome scans may fail to map local adaptation genes. Mol. Ecol.
761 20:2044–2072.
762
763 Bouchemousse, S., C. Liautard-Haag, N. Bierne, and F. Viard. 2016. Distinguishing
764 contemporary hybridization from past introgression with postgenomic ancestry-
765 informative SNPs in strongly differentiated Ciona species. Mol. Ecol. 25:5527–5542.
766
767 Boursot, P., J.-C. Auffray, J. Britton-Davidian, and F. Bonhomme. 1993. The Evolution
768 of House Mice. Annu. Rev. Ecol. Syst. 24:119–152.
769
Brawand, D., C. E. Wagner, Y. I. Li, M. Malinsky, I. Keller, S. Fan, O. Simakov, A. Y. Ng,
Z. W. Lim, E. Bezault, J. Turner-Maier, J. Johnson, R. Alcazar, H. J. Noh, P. Russell, B.
Aken, J. Alföldi, C. Amemiya, N. Azzouzi, J.-F. Baroiller, F. Barloy-Hubler, A. Berlin, R.
Bloomquist, K. L. Carleton, M. A. Conte, H. D'Cotta, O. Eshel, L. Gaffney, F. Galibert, H.
F. Gante, S. Gnerre, L. Greuter, R. Guyon, N. S. Haddad, W. Haerty, R. M. Harris, H. A.
Hofmann, T. Hourlier, G. Hulata, D. B. Jaffe, M. Lara, A. P. Lee, I. MacCallum, S.
Mwaiko, M. Nikaido, H. Nishihara, C. Ozouf-Costaz, D. J. Penman, D. Przybylski, M.
Rakotomanga, S. C. P. Renn, F. J. Ribeiro, M. Ron, W. Salzburger, L. Sanchez-Pulido, M.
E. Santos, S. Searle, T. Sharpe, R. Swofford, F. J. Tan, L. Williams, S. Young, S. Yin, N.
Okada, T. D. Kocher, E. A. Miska, E. S. Lander, B. Venkatesh, R. D. Fernald, A. Meyer, C.
P. Ponting, J. T. Streelman, K. Lindblad-Toh, O. Seehausen, and F. Di Palma. 2014. The
genomic substrate for adaptive radiation in African cichlid fish. Nature 513:375–
381.
783
784 Britton-Davidian, J., F. Fel-Clair, J. Lopez, P. Alibert, and P. Boursot. 2005. Postzygotic
785 isolation between the two European subspecies of the house mouse: estimates from
786 fertility patterns in wild and laboratory-bred hybrids. Biol. J. Linn. Soc. 84:379–393.
787
788 Butlin, R. K., M. Saura, G. Charrier, B. Jackson, C. André, A. Caballero, J. A. Coyne, J.
789 Galindo, J. W. Grahame, J. Hollander, P. Kemppainen, M. Martínez-Fernández, M.
790 Panova, H. Quesada, K. Johannesson, and E. Rolán-Alvarez. 2014. Parallel Evolution
791 of Local Adaptation and Reproductive Isolation in the Face of Gene Flow. Evolution
792 68:935–949.
793

794 Christe, C., K. N. Stölting, L. Bresadola, B. Fussi, B. Heinze, D. Wegmann, and C. Lexer.
795 2016. Selection against recombinant hybrids maintains reproductive isolation in
796 hybridizing Populus species despite F1 fertility and recurrent gene flow. Mol. Ecol.
797 25:2482-2498.
798
799 Conesa, A., and S. Götz. 2008. Blast2GO: A Comprehensive Suite for Functional
800 Analysis in Plant Genomics. Int. J. Plant Genomics 2008.
801
802 Coyne, J. A., and H. A. Orr. 2004. Speciation. Sinauer.
803
804 Curtis, J. M. R., and A. C. J. Vincent. 2006. Life history of an unusual marine fish:
805 survival, growth and movement patterns of *Hippocampus guttulatus* Cuvier 1829. J.
806 Fish Biol. 68:707–733.
807
808 de Villemereuil, P., É. Frichot, É. Bazin, O. François, and O. E. Gaggiotti. 2014. Genome
809 scan methods against more complex models: when and how much should we trust
810 them? Mol. Ecol. 23:2006–2019.
811
812 Doyle, J., and J. Doyle. 1987. A rapid DNA isolation procedure for small quantities of
813 fresh leaf tissue. Phytochem. Bull. 19:11–15.
814
815 Duforet-Frebourg, N., E. Bazin, and M. G. B. Blum. 2014. Genome scans for detecting
816 footprints of local adaptation using a Bayesian factor model. Mol. Biol. Evol.
817 31:2483–2495.
818
819 Duvaux, L., K. Belkhir, M. Boulesteix, and P. Boursot. 2011. Isolation and gene flow:
820 inferring the speciation history of European house mice. Mol. Ecol. 20:5248–5264.
821
822 Ewens, W. J. 1972. The sampling theory of selectively neutral alleles. Theor. Popul.
823 Biol. 3:87–112.
824
825 Falush, D., M. Stephens, and J. K. Pritchard. 2003. Inference of Population Structure
826 Using Multilocus Genotype Data: Linked Loci and Correlated Allele Frequencies.
827 Genetics 164:1567–1587.
828
829 Felsenstein, J. 2005. PHYLIP (Phylogeny Inference Package) version 3.6. Distributed
830 by the author. Department of Genome Sciences, University of Washington,
831 Seattle.
832
833 Foll, M., and O. Gaggiotti. 2008. A Genome-Scan Method to Identify Selected Loci
834 Appropriate for Both Dominant and Codominant Markers: A Bayesian Perspective.
835 Genetics 180:977–993.
836
837 Gagnaire, P.-A., T. Broquet, D. Aurelle, F. Viard, A. Souissi, F. Bonhomme, S. Arnaud-
838 Haond, and N. Bierne. 2015. Using neutral, selected, and hitchhiker loci to assess
839 connectivity of marine populations in the genomic era. Evol. Appl. 8:769:786.

840
841 Gayral, P., J. Melo-Ferreira, S. Glémin, N. Bierne, M. Carneiro, B. Nabholz, J. M.
842 Lourenco, P. C. Alves, M. Ballenghien, N. Faivre, K. Belkhir, V. Cahais, E. Loire, A.
843 Bernard, and N. Galtier. 2013. Reference-Free Population Genomics from Next-
844 Generation Transcriptome Data and the Vertebrate–Invertebrate Gap. PLOS Genet.
845 9:e1003457.
846
847 Gompert, Z., and C. Alex Buerkle. 2010. introgress: a software package for mapping
848 components of isolation in hybrids. Mol. Ecol. Resour. 10:378–384.
849
850 Good, J. M., M. A. Handel, and M. W. Nachman. 2008. Asymmetry and polymorphism
851 of hybrid male sterility during the early stages of speciation in house mice. Evol. Int.
852 J. Org. Evol. 62:50–65.
853
854 Guerrero, R., and M. W. Hahn. 2017. Speciation as a sieve for ancestral
855 polymorphism. bioRxiv 155176.
856
857 Harrison, R. G. 1993. Hybrid Zones and the Evolutionary Process. Oxford University
858 Press.
859
860 Harrison, R. G., and E. L. Larson. 2016. Heterogeneous genome divergence,
861 differential introgression, and the origin and structure of hybrid zones. Mol. Ecol.
862 25:2454:2466.
863
864 Hedrick, P. W. 2013. Adaptive introgression in animals: examples and comparison to
865 new mutation and standing variation as sources of adaptive variation. Mol. Ecol.
866 22:4606–4618.
867
868 Hemmer-Hansen, J., E. E. Nielsen, N. O. Therkildsen, M. I. Taylor, R. Ogden, A. J.
869 Geffen, D. Bekkevold, S. Helyar, C. Pampoulie, T. Johansen, FishPopTrace
870 Consortium, and G. R. Carvalho. 2013. A genomic island linked to ecotype divergence
871 in Atlantic cod. Mol. Ecol. 22:2653–2667.
872
873 Hewitt, G. M. 1988. Hybrid zones-natural laboratories for evolutionary studies.
874 Trends Ecol. Evol. 3:158–167.
875
876 Hoban, S., J. L. Kelley, K. E. Lotterhos, M. F. Antolin, G. Bradburd, D. B. Lowry, M. L.
877 Poss, L. K. Reed, A. Storfer, and M. C. Whitlock. 2016. Finding the Genomic Basis of
878 Local Adaptation: Pitfalls, Practical Solutions, and Future Directions. Am. Nat.
879 188:379–397.
880
881 Howe, K., M. D. Clark, C. F. Torroja, J. Torrance, C. Berthelot, M. Muffato, J. E. Collins,
882 S. Humphray, K. McLaren, L. Matthews, S. McLaren, I. Sealy, M. Caccamo, C. Churcher,
883 C. Scott, J. C. Barrett, R. Koch, G.-J. Rauch, S. White, W. Chow, B. Kilian, L. T. Quintais, J.
884 A. Guerra-Assunção, Y. Zhou, Y. Gu, J. Yen, J.-H. Vogel, T. Eyre, S. Redmond, R.
885 Banerjee, J. Chi, B. Fu, E. Langley, S. F. Maguire, G. K. Laird, D. Lloyd, E. Kenyon, S.

Donaldson, H. Sehra, J. Almeida-King, J. Loveland, S. Trevanion, M. Jones, M. Quail, D. Willey, A. Hunt, J. Burton, S. Sims, K. McLay, B. Plumb, J. Davis, C. Clee, K. Oliver, R. Clark, C. Riddle, D. Elliot, D. Eliott, G. Threadgold, G. Harden, D. Ware, S. Begum, B. Mortimore, B. Mortimer, G. Kerry, P. Heath, B. Phillimore, A. Tracey, N. Corby, M. Dunn, C. Johnson, J. Wood, S. Clark, S. Pelan, G. Griffiths, M. Smith, R. Glithero, P. Howden, N. Barker, C. Lloyd, C. Stevens, J. Harley, K. Holt, G. Panagiotidis, J. Lovell, H. Beasley, C. Henderson, D. Gordon, K. Auger, D. Wright, J. Collins, C. Raisen, L. Dyer, K. Leung, L. Robertson, K. Ambridge, D. Leongamornlert, S. McGuire, R. Gilderthorp, C. Griffiths, D. Manthravadi, S. Nichol, G. Barker, S. Whitehead, M. Kay, J. Brown, C. Murnane, E. Gray, M. Humphries, N. Sycamore, D. Barker, D. Saunders, J. Wallis, A. Babbage, S. Hammond, M. Mashreghi-Mohammadi, L. Barr, S. Martin, P. Wray, A. Ellington, N. Matthews, M. Ellwood, R. Woodmansey, G. Clark, J. D. Cooper, J. Cooper, A. Tromans, D. Grafham, C. Skuce, R. Pandian, R. Andrews, E. Harrison, A. Kimberley, J. Garnett, N. Fosker, R. Hall, P. Garner, D. Kelly, C. Bird, S. Palmer, I. Gehring, A. Berger, C. M. Dooley, Z. Ersan-Ürün, C. Eser, H. Geiger, M. Geisler, L. Karotki, A. Kirn, J. Konantz, M. Konantz, M. Oberländer, S. Rudolph-Geiger, M. Teucke, C. Lanz, G. Raddatz, K. Osoegawa, B. Zhu, A. Rapp, S. Widaa, C. Langford, F. Yang, S. C. Schuster, N. P. Carter, J. Harrow, Z. Ning, J. Herrero, S. M. J. Searle, A. Enright, R. Geisler, R. H. A. Plasterk, C. Lee, M. Westerfield, P. J. de Jong, L. I. Zon, J. H. Postlethwait, C. Nüsslein-Volhard, T. J. P. Hubbard, H. Roest Crollius, J. Rogers, and D. L. Stemple. 2013. The zebrafish reference genome sequence and its relationship to the human genome. Nature 496:498–503.

Hunt, W. G., and Selander, R. K. 1973. Biochemical genetics of hybridisation in European house mice. Heredity, 31, 1133

Jaillon, O., J.-M. Aury, F. Brunet, J.-L. Petit, N. Stange-Thomann, E. Mauceli, L. Bouneau, C. Fischer, C. Ozouf-Costaz, A. Bernot, S. Nicaud, D. Jaffe, S. Fisher, G. Lutfalla, C. Dossat, B. Segurens, C. Dasilva, M. Salanoubat, M. Levy, N. Boudet, S. Castellano, V. Anthouard, C. Jubin, V. Castelli, M. Katinka, B. Vacherie, C. Biémont, Z. Skalli, L. Cattolico, J. Poulain, V. de Berardinis, C. Cruaud, S. Duprat, P. Brottier, J.-P. Coutanceau, J. Gouzy, G. Parra, G. Lardier, C. Chapple, K. J. McKernan, P. McEwan, S. Bosak, M. Kellis, J.-N. Volff, R. Guigó, M. C. Zody, J. Mesirov, K. Lindblad-Toh, B. Birren, C. Nusbaum, D. Kahn, M. Robinson-Rechavi, V. Laudet, V. Schachter, F. Quétier, W. Saurin, C. Scarpelli, P. Wincker, E. S. Lander, J. Weissenbach, and H. Roest Crollius. 2004. Genome duplication in the teleost fish Tetraodon nigroviridis reveals the early vertebrate proto-karyotype. Nature 431:946–957.

Jakobsson, M., and N. A. Rosenberg. 2007. CLUMPP: a cluster matching and permutation program for dealing with label switching and multimodality in analysis of population structure. Bioinforma. Oxf. Engl. 23:1801–1806.

Jiggins, C. D., and J. Mallet. 2000. Bimodal hybrid zones and speciation. Trends Ecol. Evol. 15:250–255.

Johannesson, K., M. Panova, P. Kemppainen, C. André, E. Rolán-Alvarez, and R. K.
Butlin. 2010. Repeated evolution of reproductive isolation in a marine snail:
unveiling mechanisms of speciation. Philos. Trans. R. Soc. B Biol. Sci. 365:1735–
1747.

Jombart, T. 2008. adegenet: a R package for the multivariate analysis of genetic
markers. Bioinformatics 24:1403–1405.

Jombart, T., R. M. Eggo, P. J. Dodd, and F. Balloux. 2011. Reconstructing disease
outbreaks from genetic data: a graph approach. Heredity 106:383–390.

Jones, F. C., M. G. Grabherr, Y. F. Chan, P. Russell, E. Mauceli, J. Johnson, R. Swofford,
M. Pirun, M. C. Zody, S. White, E. Birney, S. Searle, J. Schmutz, J. Grimwood, M. C.
Dickson, R. M. Myers, C. T. Miller, B. R. Summers, A. K. Knecht, S. D. Brady, H. Zhang,
A. A. Pollen, T. Howes, C. Amemiya, Broad Institute Genome Sequencing Platform &
Whole Genome Assembly Team, E. S. Lander, F. Di Palma, K. Lindblad-Toh, and D. M.
Kingsley. 2012. The genomic basis of adaptive evolution in threespine sticklebacks.
Nature 484:55–61.

Jones, M. R., and J. M. Good. 2016. Targeted capture in evolutionary and ecological
genomics. Mol. Ecol. 25:185–202.

Kai, W., K. Kikuchi, S. Tohari, A. K. Chew, A. Tay, A. Fujiwara, S. Hosoya, H. Suetake, K.
Naruse, S. Brenner, Y. Suzuki, and B. Venkatesh. 2011. Integration of the Genetic Map
and Genome Assembly of Fugu Facilitates Insights into Distinct Features of Genome
Evolution in Teleosts and Mammals. Genome Biol. Evol. 3:424–442.

Kasahara, M., K. Naruse, S. Sasaki, Y. Nakatani, W. Qu, B. Ahsan, T. Yamada, Y.
Nagayasu, K. Doi, Y. Kasai, T. Jindo, D. Kobayashi, A. Shimada, A. Toyoda, Y. Kuroki, A.
Fujiyama, T. Sasaki, A. Shimizu, S. Asakawa, N. Shimizu, S. Hashimoto, J. Yang, Y. Lee,
K. Matsushima, S. Sugano, M. Sakaizumi, T. Narita, K. Ohishi, S. Haga, F. Ohta, H.
Nomoto, K. Nogata, T. Morishita, T. Endo, T. Shin-I, H. Takeda, S. Morishita, and Y.
Kohara. 2007. The medaka draft genome and insights into vertebrate genome
evolution. Nature 447:714–719.

Keenan, K., P. McGinnity, T. F. Cross, W. W. Crozier, and P. A. Prodöhl. 2013.
diveRsity: An R package for the estimation and exploration of population genetics
parameters and their associated errors. Methods Ecol. Evol. 4:782–788.

Kulmuni, J., and A. M. Westram. 2017. Intrinsic incompatibilities evolving as a by-
product of divergent ecological selection: Considering them in empirical studies on
divergence with gene flow. Mol. Ecol. 26:3093-3103.

Larson, E. L., J. A. Andrés, S. M. Bogdanowicz, and R. G. Harrison. 2013. Differential
introgression in a mosaic hybrid zone reveals candidate barrier genes. Evol. Int. J.
Org. Evol. 67:3653–3661.

Larson, E. L., T. A. White, C. L. Ross, and R. G. Harrison. 2014. Gene flow and the
maintenance of species boundaries. Mol. Ecol. 23:1668–1678.

Le Moan, A., P.-A. Gagnaire, and F. Bonhomme. 2016. Parallel genetic divergence
among coastal-marine ecotype pairs of European anchovy explained by differential
introgression after secondary contact. Mol. Ecol. 25:3187-3202.

Loire, E., Y. Chiari, A. Bernard, V. Cahais, J. Romiguier, B. Nabholz, J. M. Lourenço, and
N. Galtier. 2013. Population genomics of the endangered giant Galápagos tortoise.
Genome Biol. 14:R136.

López, A., M. Vera, M. Planas, and C. Bouza. 2015. Conservation Genetics of
Threatened Hippocampus guttulatus in Vulnerable Habitats in NW Spain: Temporal
and Spatial Stability of Wild Populations with Flexible Polygamous Mating System in
Captivity. PLoS ONE 10:e0117538.

Lotterhos, K. E., and M. C. Whitlock. 2014. Evaluation of demographic history and
neutral parameterization on the performance of FST outlier tests. Mol. Ecol.
23:2178–2192.

Lotterhos, K. E., and M. C. Whitlock. 2015. The relative power of genome scans to
detect local adaptation depends on sampling design and statistical method. Mol.
Ecol. 24:1031–1046.

1001

1002　Lourie, S. A., and A. C. J. Vincent. 2004. Using Biogeography to Help Set Priorities in
1003　Marine Conservation. Conserv. Biol. 18:1004–1020.

1004

1005　Luu, K., E. Bazin, and M. G. B. Blum. 2016. pcadapt: an R package to perform genome
1006　scans for selection based on principal component analysis. bioRxiv 056135.

1007

1008　Macholán, M., S. J. E. Baird, P. Dufková, P. Munclinger, B. V. Bímová, and J. Piálek.
1009　2011. Assessing multilocus introgression patterns: a case study on the mouse X
1010　chromosome in central Europe. Evol. Int. J. Org. Evol. 65:1428–1446.

1011

1012　Mallet, J., N. Besansky, and M. W. Hahn. 2016. How reticulated are species? Bioessays
1013　38:140–149.

1014

1015　Naruse, K., M. Tanaka, K. Mita, A. Shima, J. Postlethwait, and H. Mitani. 2004. A
1016　Medaka Gene Map: The Trace of Ancestral Vertebrate Proto-Chromosomes Revealed
1017　by Comparative Gene Mapping. Genome Res. 14:820–828.

1018

1019　Nei, M. 1972. Genetic Distance between Populations. Am. Nat. 106:283–292.

1020

1021　Nosil, P., and J. L. Feder. 2012. Genomic divergence during speciation: causes and
1022　consequences. Phil Trans R Soc B 367:332–342.

1023
1024 Perry, G. H., P. Melsted, J. C. Marioni, Y. Wang, R. Bainer, J. K. Pickrell, K. Michelini, S.
1025 Zehr, A. D. Yoder, M. Stephens, J. K. Pritchard, and Y. Gilad. 2012. Comparative RNA
1026 sequencing reveals substantial genetic variation in endangered primates. Genome
1027 Res. 22:602–610.
1028
1029 Pickrell, J. K., and J. K. Pritchard. 2012. Inference of Population Splits and Mixtures
1030 from Genome-Wide Allele Frequency Data. PLoS Genet 8:e1002967.
1031
1032 Pritchard, J. K., M. Stephens, and P. Donnelly. 2000. Inference of Population
1033 Structure Using Multilocus Genotype Data. Genetics 155:945–959.
1034
1035 R Development Core Team (2011) R: A language and environment for statistical
1036 computing. R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-
1037 07-0, http://www.R-project.org/.
1038
1039 Rand, D. M., and R. G. Harrison. 1989. Ecological Genetics of a Mosaic Hybrid Zone:
1040 Mitochondrial, Nuclear, and Reproductive Differentiation of Crickets by Soil Type.
1041 Evolution 43:432–449.
1042
1043 Ravinet, M., Faria, R., Butlin, R.K., Galindo, J., Bierne, N., Rafajlović, M., Noor, M.A.F.,
1044 Mehlig, B., and A.M. Westram (accepted in Journal of Evolutionary Biology).
1045 Interpreting the genomic landscape of speciation: a road map for finding barriers to
1046 gene flow
1047
1048 Raymond, M., and F. Rousset. 1995. GENEPOP (Version 1.2): Population Genetics
1049 Software for Exact Tests and Ecumenicism. J. Hered. 86:248–249.
1050
1051 Romiguier, J., P. Gayral, M. Ballenghien, A. Bernard, V. Cahais, A. Chenuil, Y. Chiari, R.
1052 Dernat, L. Duret, N. Faivre, E. Loire, J. M. Lourenco, B. Nabholz, C. Roux, G.
1053 Tsagkogeorga, A. a.-T. Weber, L. A. Weinert, K. Belkhir, N. Bierne, S. Glémin, and N.
1054 Galtier. 2014. Comparative population genomics in animals uncovers the
1055 determinants of genetic diversity. Nature 515:261–263.
1056
1057 Rougemont, Q., C. Roux, S. Neuenschwander, J. Goudet, S. Launey, and G. Evanno.
1058 2016. Reconstructing the demographic history of divergence between European
1059 river and brook lampreys using approximate Bayesian computations. PeerJ 4.
1060
1061 Rougeux, C., L. Bernatchez, and P.-A. Gagnaire. 2016. Modeling the Multiple Facets of
1062 Speciation-with-Gene-Flow Towards Inferring the Divergence History of Lake
1063 Whitefish Species Pairs (coregonus Clupeaformis). bioRxiv 068932.
1064
1065 Rousset, F. 2008. genepop'007: a complete re-implementation of the genepop
1066 software for Windows and Linux. Mol. Ecol. Resour. 8:103–106.
1067

1068   Roux, C., C. Fraïsse, J. Romiguier, Y. Anciaux, N. Galtier, and N. Bierne. 2016. Shedding
1069   Light on the Grey Zone of Speciation along a Continuum of Genomic Divergence.
1070   PLOS Biol. 14:e2000234.
1071
1072   Sakka Hlaili, A., B. Grami, N. Niquil, M. Gosselin, D. Hamel, M. Troussellier, and H.
1073   Hadj Mabrouk. 2008. The planktonic food web of the Bizerte lagoon (south-western
1074   Mediterranean) during summer: I. Spatial distribution under different
1075   anthropogenic pressures. Estuar. Coast. Shelf Sci. 78:61–77.
1076
1077   Schluter, D., and G. L. Conte. 2009. Genetics and ecological speciation. Proc. Natl.
1078   Acad. Sci. U. S. A. 106 Suppl 1:9955–9962.
1079
1080   Small, C. M., S. Bassham, J. Catchen, A. Amores, A. M. Fuiten, R. S. Brown, A. G. Jones,
1081   and W. A. Cresko. 2016. The genome of the Gulf pipefish enables understanding of
1082   evolutionary innovations. Genome Biol. 17:258.
1083
1084   Soria-Carrasco, V., Z. Gompert, A. A. Comeault, T. E. Farkas, T. L. Parchman, J. S.
1085   Johnston, C. A. Buerkle, J. L. Feder, J. Bast, T. Schwander, S. P. Egan, B. J. Crespi, and P.
1086   Nosil. 2014. Stick insect genomes reveal natural selection's role in parallel
1087   speciation. Science 344:738–742.
1088
1089   Storey, J. D. 2002. A direct approach to false discovery rates. J. R. Stat. Soc. Ser. B Stat.
1090   Methodol. 64:479–498.
1091
1092   Szymura, J. M., and N. H. Barton. 1986. Genetic Analysis of a Hybrid Zone Between
1093   the Fire-Bellied Toads, Bombina bombina and B. variegata, Near Cracow in Southern
1094   Poland. Evolution 40:1141–1159.
1095
1096   Teeter, K. C., B. A. Payseur, L. W. Harris, M. A. Bakewell, L. M. Thibodeau, J. E. O'Brien,
1097   J. G. Krenz, M. A. Sans-Fuentes, M. W. Nachman, and P. K. Tucker. 2008. Genome-
1098   wide patterns of gene flow across a house mouse hybrid zone. Genome Res. 18:67–
1099   76.
1100
1101   Tine, M., H. Kuhl, P.-A. Gagnaire, B. Louro, E. Desmarais, R. S. T. Martins, J. Hecht, F.
1102   Knaust, K. Belkhir, S. Klages, R. Dieterich, K. Stueber, F. Piferrer, B. Guinand, N.
1103   Bierne, F. A. M. Volckaert, L. Bargelloni, D. M. Power, F. Bonhomme, A. V. M. Canario,
1104   and R. Reinhardt. 2014. European sea bass genome and its variation provide
1105   insights into adaptation to euryhalinity and speciation. Nat. Commun. 5.
1106
1107   Tyner, C., G. P. Barber, J. Casper, H. Clawson, M. Diekhans, C. Eisenhart, C. M. Fischer,
1108   D. Gibson, J. N. Gonzalez, L. Guruvadoo, M. Haeussler, S. Heitner, A. S. Hinrichs, D.
1109   Karolchik, B. T. Lee, C. M. Lee, P. Nejad, B. J. Raney, K. R. Rosenbloom, M. L. Speir, C.
1110   Villarreal, J. Vivian, A. S. Zweig, D. Haussler, R. M. Kuhn, and W. J. Kent. 2017. The
1111   UCSC Genome Browser database: 2017 update. Nucleic Acids Res. 45:D626–D634.
1112

1113 Via, S. 2012. Divergence hitchhiking and the spread of genomic isolation during
1114 ecological speciation-with-gene-flow. Philos. Trans. R. Soc. Lond. B Biol. Sci.
1115 367:451–460.
1116
1117 Vines, T., A. Dalziel, A. Albert, T. Veen, P. Schulte, and D. Schluter. 2016. Cline
1118 coupling and uncoupling in a stickleback hybrid zone. Evolution 70:1023-38.
1119
1120 Weir, B. S., and C. C. Cockerham. 1984. Estimating F-Statistics for the Analysis of
1121 Population Structure. Evolution 38:1358–1370.
1122
1123 Welch, J. J., and C. D. Jiggins. 2014. Standing and flowing: the complex origins of
1124 adaptive variation. Mol. Ecol. 23:3935–3937.
1125
1126 Woodall, L. C., R. Jones, B. Zimmerman, S. Guillaume, T. Stubbington, P. Shaw, and H.
1127 J. Koldewey. 2012. Partial fin-clipping as an effective tool for tissue sampling
1128 seahorses, Hippocampus spp. J. Mar. Biol. Assoc. U. K. 92:1427–1432.
1129
1130 Woodall, L. C., H. J. Koldewey, J. T. Boehm, and P. W. Shaw. 2015. Past and present
1131 drivers of population structure in a small coastal fish, the European long snouted
1132 seahorse Hippocampus guttulatus. Conserv. Genet. 1–15.
1133
1134 Yeaman, S. 2013. Genomic rearrangements and the evolution of clusters of locally
1135 adaptive loci. Proc. Natl. Acad. Sci. 110:E1743–E1751.
1136

1137 **Data archival location**
1138 SNPs data have been deposited at DRYAD. DOI: *to be completed after the manuscript is*
1139 *accepted for publication*
1140
1141

 Figures

 Figure 1 Sampling locations of *Hippocampus guttulatus* (A-) and Neighbor Joining trees
 (B-D-). Each study site is labeled as follow: 1- Poole, United Kingdom, 2- Brest, France,
 3- Le Croisic, France, 4- Ré Island, France, 5- Arcachon, France, 6- Hossegor, France, 7-
 Coruna, Spain, 8- Vigo, Spain, 9- Portimao, Portugal, 10- Faro (maritime site), Portugal,
 11- Faro (lagoon site), Portugal, 12- Malaga, Spain, 13- Tossa de Mar, Spain, 14-
 Leucate, France, 15- Sète (maritime site), France, 16- Thau lagoon, France, 17- La Ciotat,
 France, 18- Le Brusc, France, 19- Cavalaire-sur-Mer, France, 20- Bizerte lagoon,
 Tunisia, 21- Naples, Italy, 22- Rovinj, Croatia, 23- Kalamaki, Greece, 24- Chalkida,
 Greece, and 25- Varna, Bulgaria. Red, violet, blue, green, pink, yellow and turquoise dots
 stand for North Atlantic cluster, Hossegor, South Atlantic cluster, Mediterranean cluster,
 Thau lagoon, Bizerte lagoon and Chalkida, and Varna, respectively. B-D- Neighbor
 Joining trees of *H. guttulatus* based on 272 non-outlier SNPs (B-), 14 outliers (C-) and all
 markers (D-).

 Figure 2 Genetic population structure analyses based on 286 SNP markers.
 A-: Principal Component Analyses depicting axis 1 (explaining 60.1% of the variance)
 and axis 2 (explaining 16.2% of the variance; left panel) and axes 1 and 3 (explaining
 9.9% of the variance; right panel). Samples are labeled and colored according to Fig. 1.
 Each label shows the barycenter of each study site.
 B- Individual Bayesian ancestry proportions determined using STRUCTURE with K=5
 clusters identified. Dotted black lines separate each study site, which is labeled following
 Fig. 1. The five clusters identified are distinguished by different colors: red, blue, green,
 pink and turquoise. Each individual is depicted as a vertical bar with colors distinguishing
 its ancestries to the five clusters.
 C- The raw genotype matrix for a subset of 105 most differentiated SNP markers and all
 individuals. Individuals are ordered based on their locations (x-axis). Each rectangle
 denotes an individual's genotype at a given locus: dark green (homozygote for the allele
 most frequent in the North Atlantic cluster), green (heterozygotes) and light green
 (homozygote for the minor allele in the North Atlantic cluster). The names of the outlier
 SNPs are mentioned at the right of the matrix. Outliers' names were highlighted
 according to the category defined in Fig. 5: the first category of outliers (*i.e.* convergent
 SNPs) were represented in orange (A-) and blue dots (B-), the 3rd category in pink (C-),
 the 5th category in red (E-), and in green dots the last category of outliers (F-).
 D- Population trees inferred by TREEMIX without (left panel) or with 3 migration events
 (right panel). Admixture arrows are colored according to the migration weight. The
 model including three admixture events significantly improved the fit as compared to a
 situation without migration (*p*-value < 0.001).

 Figure 3 Empirical distributions of the original Lewontin-Krakaeur statistic (grey bars)
 for each locus and fitted a theoretical generalized extreme value distribution using BFGS
 (red line) with K = 2 alleles per locus and two demes (North and South Atlantic clusters).

 Figure 4 Phylogenomic tree (A-), outlier cross-mapping (B-) and F_{ST} along the unique
 chromosome (C-). The phylogenomic tree (A-) reconstructed from Tine et al. (2014) Lin

et al. (2016) and Small et al. (2016) is illustrated with regards to the mapping location of
H. guttulatus outlier SNPs on eight well-assembled fish genomes (*Syngnathus scovelli*,
Dicentrarchus labrax, *Gasterosteus aculeatus*, *Oryzias latipes*, *Oreochromis niloticus*,
Takifugu rubripes, *Tetraodon nigroviridis* and *Danio rerio*) and unplaced genomic
scaffolds (*Hippocampus comes*), showing a single matching chromosome for each target
species. As *H. comes* scaffolds are unplaced on the genome, *H. guttulatus* SNPs were
first blasted on *H. comes* scaffolds, then each scaffold was mapped to the other well-
assembled fish genomes. Three scaffolds (339, 397 and 8) mapped to a single
chromosome for each species and a putative chromosome was here reconstructed. The
order of the scaffolds and SNPs according to the blasts and blats was conserved. Outlier
SNPs displaying parallel differentiation between Atlantic lineages and Mediterranean
ecotypes (the A and B categories in Fig. 5) are colored in orange and blue, respectively,
while outliers highly differentiated between Mediterranean ecotypes (the C category) are
colored in red.
1202
Figure 5 *H. guttulatus* allele frequencies for the 14 outliers identified, along with two
neutral SNPs mapped to the genomic island. Each study site is labeled and colored
according to Fig. 1, with Hossegor separated in 6A and 6B and Chalkida in 24A and 24B-
and colored according to Fig. 1. SNP allele frequencies are categorized in: outliers
distinguishing higher frequencies in North Atlantic samples, Thau and Bizerte lagoons
and, in a lesser extent Chalkida and Varna (A-) plus Mediterranean samples (B-); Thau
lagoon, Chalkida, Varna (C-); North Atlantic samples (E-); and Mediterranean samples to
the other sites (F-), and neutral SNPs but mapped to the genomic island (D-). Allelic
frequency differences (AFD, G-) between lagoon and sea ecotypes (x-axis) were plotted
against AFD of North *vs*. South Atlantic lineages (y-axis) for all SNPs. Outliers were
highlighted according to their category: the first two categories of outliers (*i.e.*
convergent SNPs) were represented in orange (A-) and blue dots (B-), the 3[rd] category in
pink (C-), the 5[th] category in red (E-), and in green dots the last category of outliers (F).
1216
Figure 6 Genomic cline plots for two illustrative markers showing extreme level of
differentiation (SNPs 29 and 130). Genomic clines were conducted on North and South
Atlantic lineages following polynomial regressions on all Atlantic individuals without
Hossegor (in yellow), on all Atlantic individuals (in grey) and on all Atlantic individuals,
without North Atlantic (in blue). Dotted lines represent the 95% confidence intervals.
Circles indicate the raw genotype data (ancestral homozygotes on the top line –the "1"
genotype, heterozygotes in the middle –the "0.5" genotype, and derived homozygotes on
the bottom line –the "0" genotype).

A-

C- Neighbor-joining tree based on 14 outliers

Mediterranean Sea

South Atlantic

Mediterranean lagoons

&

North Atlantic

B- Neighbor-joining tree based on neutral markers

North Atlantic   South Atlantic   Mediterranean   Mediterranean Sea
                                    lagoons

D- Neighbor-joining tree based on all markers

Mediterranean lagoons

Mediterranean Sea

South Atlantic

North Atlantic

LK statistic

Genomic Island Loci

Outlier Loci

A- SNP 29, SNP 212, SNP 286
SNP 34, SNP 232, SNP 312

C- SNP 87, SNP 311

Non Outlier Loci

D- SNP 165, SNP 178

Non Genomic Island Loci

Outlier Loci

B- SNP 140, SNP 242

E- SNP 130, SNP 169

F- SNP 26, SNP 314

G- AFD North – South Atlantic vs AFD Mediterranean Lagoon – Sea

genetic parallelism

A type SNPs, B type SNPs, C type SNPs, E type SNP, F type SNPs