Parallel pattern of differentiation at a genomic island shared between clinal and mosaic hybrid zones in a complex of cryptic seahorse lineages

Florentine Riquet[1, 2], Cathy Liautard-Haag[1, 2], Lucy Woodall[3, 4], Carmen Bouza[5], Patrick Louisy[6, 7], Bojan Hamer[8], Francisco Otero-Ferrer[9], Philippe Aublanc[10], Vickie Béduneau[11], Olivier Briard[12], Tahani El Ayari[1, 2], Sandra Hochscheid[13], Khalid Belkhir[1, 2], Sophie Arnaud-Haond[1, 14], Pierre-Alexandre Gagnaire[1, 2], Nicolas Bierne[1, 2]

[1] Institut des Sciences de l'Evolution de Montpellier, Université Montpellier, Montpellier, France
[2] CNRS Institut des Sciences de l'Evolution, UMR5554 UM-CNRS-IRD-EPHE, Station Marine OREME, Sète, France
[3] Department of Zoology, University of Oxford, John Krebs Field Station, Wytham, OX2 8QJ, UK
[4] Natural History Museum, Cromwell Road, London SW7 5BD, UK
[5] Department of Genetics, Faculty of Veterinary Science, Universidade de Santiago de Compostela, Campus de Lugo, Lugo, Spain
[6] University of Nice Sophia Antipolis, ECOMERS Laboratory, Faculty of Sciences, Parc Valrose, Nice, France
[7] Association Peau-Bleue, 46 rue des Escais, Agde, France
[8] Center for Marine Research, Ruder Boskovic Institute, Giordano Paliaga 5, 52210 Rovinj, Croatia
[9] Grupo en Biodiversidad y Conservación, IU-ECOAQUA, Universidad de Las Palmas de Gran Canaria, Crta. Taliarte s/n, 35214 Telde, Spain
[10] Institut océanographique Paul Ricard, Ile des Embiez, Six-Fours-les-Plages, France
[11] Océarium du Croisic, Avenue de Saint Goustan, Le Croisic, France
[12] Aquarium de Biarritz, Biarritz Océan, Plateau de l'Atalaye, Biarritz, France
[13] Stazione Zoologica Anton Dohrn, Department Research Infrastructures for Marine Biological Resources, Aquarium Unit, Napoli, Italy
[14] Ifremer - MARine Biodiversity, Exploitation and Conservation, UMR 9190 IRD-IFREMER-UM-CNRS, Sète, France

Abstract

Diverging semi-isolated lineages either meet in narrow clinal hybrid zones, or have a mosaic distribution associated with environmental variation. Intrinsic reproductive isolation is often emphasized in the former and local adaptation in the latter, although both can contribute to isolation. Rarely these two patterns of spatial distribution are reported in the same study system. Here we report that the long-snouted seahorse *Hippocampus guttulatus* is subdivided into discrete panmictic entities by both types of hybrid zones. Along the European Atlantic coasts, a northern and a southern lineage meet in the southwest of France where they coexist in sympatry with little hybridization. In the Mediterranean Sea, two lineages have a mosaic distribution, associated with lagoon-like and marine habitats. A fifth lineage was identified in the Black Sea. Genetic homogeneity over large spatial scales contrasts with isolation maintained in sympatry or close parapatry at a fine scale. A high variation in locus-specific introgression rates provides

additional evidence that partial reproductive isolation must be maintaining the divergence. Surprisingly, fixed differences between lagoon and marine populations in the Mediterranean Sea belong to the most differentiated SNPs between the two Atlantic lineages, against the genome-wide pattern of structure. These parallel outlier SNPs cluster on a single chromosome-wide island of differentiation. Since Atlantic lineages do not match the lagoon-sea habitat variation, genetic parallelism at the genomic island suggests a shared genetic barrier contributes to reproductive isolation in contrasting contexts -*i.e.* spatial *vs*. ecological. We discuss how a genomic hotspot of parallel differentiation could have evolved and become associated either with space or with a patchy environment in a single study system.

Keywords: clinal hybrid zone, mosaic hybrid zone, reproductive isolation, local adaptation, ecological speciation, parallel evolution

Introduction

The spatial context of contact zones between partially isolated taxa and their relationship with environmental variation was long thought to offer great promises to unravel the nature and origin of species. Though each taxon may be genetically homogeneous over large distances, they often meet in abrupt genetic discontinuities, called hybrid zones, in which partial reproductive isolation limits gene exchange (Barton and Hewitt 1985, Hewitt 1988). Hybrid zones are extremely informative for exploring the genetic basis of reproductive isolation (e.g. Teeter et al. 2008, Christe et al. 2016) and local adaptation (e.g. Jones et al. 2012, Larson et al. 2013, Soria-Carrasco et al. 2014), as well as identifying genomic regions involved either in increased genomic differentiation (Ravinet et al. 2017) or adaptive introgression (Hedrick 2013). The hybrid zone literature usually contrasts two spatial patterns (Harrison 1993): (i) clinal hybrid zones, with parapatrically distributed parental forms on both sides of a genetic divide, and (ii) mosaic hybrid zones, when the environment consists of a mosaic of habitat patches to which taxa (ecotypes, host races, hybridizing species) are somehow specialized. Contrasting with this long-standing dichotomy, hybrid zones of both types have now been recognized to be multifactorial and maintained by exogenous and endogenous diverging mechanisms (*i.e.* local adaptation and intrinsic reproductive isolation, respectively; Barton and Hewitt 1985, Bierne et al. 2011). Nonetheless, clinal hybrid zones still tend to be interpreted as being mainly maintained by intrinsic reproductive isolation evolved in allopatry before contact (the tension zone model, Barton and Hewitt 1985). Conversely, local adaptation is often emphasized to drive mosaic distributions which are interpreted as evidence that selection at local adaptation genes is acting to increase differentiation between habitats in a parallel fashion (Nosil and Feder 2012). This sketchy dichotomy is anchored by emblematic study systems for which decades of research allow support for such interpretations. For instance, the hybrid zone between the mice *Mus musculus musculus* and *M. m. domesticus* in central Europe (Boursot et al. 1993) has been well demonstrated to be maintained by selection against hybrid genotypes (Britton-Davidian et al. 2005, Good et al. 2008) after secondary contact (Duvaux et al. 2011). Although the position of the hybrid zone was initially found to be associated with rainfall (Hunt and Selander 1973), local adaptation is not considered to contribute much to the isolation. At the other extreme, local adaptation of the same genetic variants has repeatedly allowed marine three-spined sticklebacks to evolve into a freshwater ecotype (Jones et al. 2012). Intrinsic selection is thought absent in the marine-freshwater sticklebacks system, although hybrids perform poorly in both environment and tend to reside in salinity ecotones (Vines et al. 2016). It would be misleading, however, to suggest the two alternative spatial contexts and relations to environmental variation may correspond to alternative routes to speciation (e.g. mutation-order *vs.* ecological speciation). The list of hybrid zones maintained both by local adaptation and intrinsic reproductive isolation is also long. *Bombina* toads (Szymura and Barton 1986), *Gryllus* crickets (Rand and Harrison 1989, Larson et al. 2014), or *Mytilus* mussels (Bierne et al. 2003) are well-known examples of mosaic hybrid zones maintained by exogenous and endogenous selection. However, we do not usually expect intrinsic incompatibilities to be reused in a parallel fashion. Besides, parallel genetic divergence associated with contrasting environmental conditions (e.g. marine/freshwater, highland/lowland, host races) remains a strong hallmark of ecologically-driven divergence (Bierne et al. 2013). Given this -hybrid zone- context, in

this paper we aim to provide an example of a genetic parallelism with a lack of apparent
ecological convergence. We expect this counter-example could contribute to break the
inductive reasoning that genetic parallelism means ecological convergence. However, as
we discovered this pattern by serendipity in a newly studied complex of cryptic genetic
backgrounds in a non-model system, we have to describe this system first.

We studied the population genetics of the long-snouted seahorse, *Hippocampus
guttulatus,* across a large part of its geographic range. We developed an assay of 286
informative SNPs chosen from more than 2,500 SNPs identified in a population
transcriptomic study (Romiguier et al. 2014). *H. guttulatus* displays poor dispersal
abilities (e.g. site-fidelity, weak swimming performance, lack of dispersive stage) and
inhabits fragmented coastal habitats along its distribution range (from the English
Channel through the Mediterranean and Black Seas, Lourie and Vincent 2004). In
addition most populations are small and patchy. Given these biological characteristics, a
strong genetic structure could have been expected. In agreement, a very low genetic
diversity was observed in *H. guttulatus* when compared to 75 non-model animal species
(Romiguier et al. 2014). However, genetic differentiation proved to be very weak over
very large geographic distances based on the two genetic studies conducted to date with
microsatellite loci (e.g. from the United Kingdom to North of Spain, Woodall et al. 2015,
or across the Cape Finisterre oceanographic barrier, López et al. 2015). Four well-
differentiated genetic clusters, each distributed over extended regions, were delineated by
genetic discontinuities corresponding to usual delimitations between vicariant marine
lineages (Woodall et al. 2015) – between the Iberian Peninsula and the Bay of Biscay in
the North Eastern Atlantic, between the Atlantic Ocean and the Mediterranean Sea, and
between the Mediterranean and Black Seas. Although such genetic differentiation
matches well with oceanographic barriers and was interpreted as spatial differentiation,
this pattern is also concordant with the existence of reproductively isolated cryptic
lineages, which boundaries were trapped by exogenous barriers. In this latter
interpretation, although the location of genetic breaks would be due to exogenous factors
(e.g. temperature, salinity or oceanic fronts), the barrier to gene flow would mainly be
driven by barrier loci that restrict gene flow on a large fraction of the genome (*i.e.* the
coupling hypothesis; Bierne et al. 2011, Gagnaire et al. 2015, Ravinet et al. 2017). This
hypothesis is receiving increasing support (e.g. Le Moan et al. 2016, Rougeux et al. 2016,
Rougemont et al. 2016) and could well explain the genetic structure observed in the long-
snouted seahorse.

Using newly developed SNP-markers spread along the genome and a more
extensive sampling along the *H. guttulatus* distribution range, we challenged the initial
interpretation of barriers to dispersal against the alternative hypothesis of reproductive
isolation between semi-isolated genetic backgrounds. We described five cryptic semi-
isolated lineages: two lineages in the Atlantic Ocean with a parapatric distribution, two
lineages in the Mediterranean Sea with a patchy fine-grained environment association
(lagoon *vs.* marine environments), and one in the Black Sea. Surprisingly, a shared
genomic island of clustered outlier loci was involved both in the isolation between the
two parapatric lineages in the Atlantic Ocean and between the marine and lagoon
ecotypes in the Mediterranean Sea. Furthermore, the North Atlantic lineage was related to
the lagoon ecotype at this genomic island, against the genome-wide pattern of structure.
However, the two Atlantic lineages inhabit both marine and lagoon habitats. We argue

that the *H. guttulatus* complex could become one of a few systems where a clinal and a mosaic hybrid zone are observed concomitantly, and a valuable new counter-example that provides evidence of genetic parallelism in absence of ecological convergence.

Materials and Methods

Figure 1 Sampling locations of *Hippocampus guttulatus*. Each study site is labeled as follow: 1- Poole, United Kingdom, 2- Brest, France, 3- Le Croisic, France, 4- Ré Island, France, 5- Arcachon, France, 6- Hossegor, France, 7- Coruna, Spain, 8- Vigo, Spain, 9- Portimao, Portugal, 10- Faro (maritime site), Portugal, 11- Faro (lagoon site), Portugal, 12- Malaga, Spain, 13- Tossa de Mar, Spain, 14- Leucate, France, 15- Sète (maritime site), France, 16- Thau lagoon, France, 17- La Ciotat, France, 18- Le Brusc, France, 19- Cavalaire-sur-Mer, France, 20- Bizerte lagoon, Tunisia, 21- Naples, Italy, 22- Rovinj, Croatia, 23- Kalamaki, Greece, 24- Halkida, Greece, and 25- Varna, Bulgaria. Lagoon habitats are represented by triangles, lagoon-like habitats by diamonds and maritime habitats by circles. Red, blue, green, pink and turquoise symbols stand respectively for the North Atlantic, South Atlantic, Mediterranean Sea, Mediterranean lagoon, and Varna cluster, all of them showing panmixia (solid lines). Dashed lines stand for contact zones (sites 6, 20 and 23).

Sampling and DNA extraction

  H. guttulatus samples were collected alive from 25 sites (Fig. 1) using a variety of methods (snorkeling, scuba diving, trawling nets, aquarium, donations). The dorsal fin of each individual was clipped using a non-lethal procedure (Woodall et al. 2012), before

releasing back the individual into its natural habitat. In three sites (sites 14, 18 et 19 in
Fig. 1), dorsal fins were clipped from captive-bred seahorse held at the Mare Nostrum
Aquarium, France, recently sampled in their natural habitats. Each individual sample was
preserved and stored in 96% ethanol for subsequent genetic analyses.
Whole genomic DNA was extracted following either Woodall et al. (2015), López
et al. (2015) or using a standard CetylTrimethyl Ammonium Bromide (CTAB)
Chloroform:Isoamyl alcohol (24:1) protocol (Doyle and Doyle 1987). For samples from
Vigo (Spain), the whole genome was amplified using the GenomiPhi kit (GE HealthCare)
according to the manufacturer's protocol. Quality and quantity of DNA extraction was
checked on an agarose gel, and normalized to 35 ng.μL^{-1} using Qubit Fluorometric
Quantitation (Invitrogen).
188
Data mining for SNP markers
A set of 12,613 contigs was examined to identify SNPs. This included one
mitochondrial contig (GenBank accession number: AF192664) and 12,612 contigs from
Romiguier et al. (2014). Briefly, Romiguier et al. (2014) produced high-coverage
transcriptomic data (RNAseq) for six *H. guttulatus* from three locations (Le Croisic,
Atlantic Ocean, France; Faro, Atlantic Ocean, Portugal; Thau lagoon, Mediterranean Sea,
France), and two *H. hippocampus* from two locations (Sète, Mediterranean Sea, France;
Bizerte, Mediterranean Sea, Tunisia). *De novo* transcriptome assembly based on Illumina
reads was performed using a combination of the programs ABySS (Simpson et al. 2009)
and Cap3 (Huang and Madan 1999), then mapped to predicted cDNAs (contigs) with
BWA (Li and Durbin 2009). Based on these 12,613 contigs, SNPs were identified using
the bioinformatic pipeline described in Bouchemousse et al. (2016). SNPs were called
with Read2SNPs (Gayral et al. 2013) and filtered out according to the following criteria
to exclude: 1) SNPs showing more than two alleles, 2) SNPs failing to be sequenced in at
least one location, 3) SNPs present in only one individual (*i.e.* singletons), 4) SNPs
identified as paralogs using the paraclean option of Read2SNPs, and 5) SNPs closer than
20 bp from a contig extremity or an exon limit when blasted against the stickleback, cod
and tilapia genomes. This resulted in 2,684 selected SNPs screened with the Illumina
Assay Design Tool (ADT) software to select 384 SNPs on the basis of their quality index
(ADT score > 0.6). An Illumina BeadXpress® with Veracode™ technology
(GoldenGate® Genotyping Assay) was then used to genotype the 384 selected SNPs.
To identify their chromosomal positions, the template sequences of the targeted
SNP were blasted against i) the genome of the Gulf pipefish (*Syngnathus scovelli*, Small
et al. 2016) and ii) the scaffolds of the tiger tail seahorse (*Hippocampus comes*, Lin et al.
2016). *H. comes* scaffolds being unplaced, we blasted these scaffolds against *S. scovelli*
genome and, for more consistency, blatted (Bhagwatt et al. 2012) them against seven
well-assembled fish genomes; zebrafish (*Dano rerio*, Howe et al. 2013), fugu (*Takifugu
rubripes*, Kai et al. 2011), tetraodon (*Tetraodon nigroviridis*, Jaillon et al. 2004), Nile
tilapia (*Oreochromis niloticus*, Brawand et al. 2014), medaka (*Oryzias latipes*, Kasahara
et al. 2007), stickleback (*Gasterosteus aculeatus*, Jones et al. 2012), and European
seabass (*Dicentrarchus labrax*, Tine et al. 2014).
SNPs were polarized using *H. hippocampus* as an outgroup to identify the most
parsimonious ancestral variant, which allowed the derived allele state to be identified.
The Joint Site-Frequency Spectrum (JSFS is the more informative summary statistics

regarding inter-population polymorphism; Wakeley 2008) obtained from the original
transcriptome-wide SNP dataset (Romiguier et al. 2014) was compared to the JSFS
obtained by the subset of 384 SNPs to investigate the extend of ascertainment bias
potentially induced by our marker selection. In order to compare the JSFS obtained with
both datasets, the JSFS dimension was projected down to a 5x5 matrix, which was the
dimension of the Romiguier et al. (2014) dataset.
229
Analyses of Genetic Diversity and Genetic Structure
The mean number of alleles per locus (N_{all}) and the allelic richness (A_r, *i.e.* the
expected number of alleles corrected for sampling size based on a rarefaction method)
were estimated using the function divBasic implemented in the R package diveRsity
(Keenan et al. 2013). Allelic frequencies, expected heterozygosity (H_e) and fixation index
(F_{IS}) were estimated using GENEPOP on the web (Raymond and Rousset 1995, Rousset
2008). To get a genome-wide picture from the multi-locus genotype dataset summarizing
inter-population polymorphism, the raw SNP data were visualized by INTROGRESS
(Gompert and Buerkle 2010), although its primary goal is to estimate genomic clines, but
allows in a simple manner to plot raw data to investigate any genomic signal. Alleles
derived from each of the parental populations were counted at the individual level and
converted into a matrix of counts, then used to visualize the multilocus genotype of each
individual.
Genetic structure among sampling sites was depicted using both Principal
Component Analysis (PCA) computed on the matrix of genotypes, and an individual-
based Bayesian clustering method. The latter being a model-based approach with strong
priors and hypotheses (Hardy-Weinberg equilibrium, no linkage disequilibrium)
contrasted with PCA, a distant-based approach when few (nearly no) assumptions may be
violated. Comparing results from both analyses using different statistical approaches,
allows us to make solid assumptions about our data. The PCAs were carried out using the
R package ADEGENET 1.4–2 (Jombart 2008, Jombart et al. 2011). The individual-based
Bayesian clustering analysis was performed with the software STRUCTURE 2.3.4
(Pritchard et al. 2000, Falush et al. 2003). For each value of K (ranging from 1 to 25), 30
replicate chains of 150,000 Monte Carlo Markov Chain (MCMC) iterations were run
after discarding 50,000 burn-in iterations. An admixture model with correlated allele
frequencies was applied with *a priori* information on samples origin. Note that this
method makes the assumption of homogeneous admixture rate in the genome (neutral
admixture) and therefore return a sort of weighted average admixture rate when
introgression is heterogeneous across the genome. To determine individual ancestry
proportions (q-values) that best matched across all replicate runs, CLUMPP (Jakobsson
and Rosenberg 2007) was used and individuals' assignment visualized in the R software.
Once the different genetic clusters were identified (and cross-validated using both
methods), genetic homogeneity among samples belonging to a cluster was checked,
allowing subsequently pooling of samples into clusters with a minimum of seven
individuals per cluster. Genetic structure was computed among clusters by calculating
global and pairwise F_{ST} (Weir and Cockerham 1984) using GENEPOP on the web
(Rousset 2008). Exact tests for population differentiation (10,000 dememorization steps,
500 batches and 5,000 iterations per batch) were carried out to test for differences in
allele frequencies. s, defined as the adjusted p-values using an optimized False Discovery

Rate approach, were computed using the QVALUE package in the R software (Storey
2002) to correct for multiple testing.
Evolutionary history of genetic clusters was also investigated under a model of
divergence and admixture events using the population graph approach implemented in
the TREEMIX software (Pickrell and Pritchard 2012). This software uses the covariance
matrix of allele frequency between pairs of populations to infer both population splits and
gene flow. A maximum likelihood population tree is first generated under the hypothesis
of an absence of migration, and admixture events are sequentially added, improving (or
not) the tree model. This statistical method shows the benefit of constructing population
trees while testing for gene flow between diverged populations at the same time. Samples
with too small sample size (N<7) or without random mating (see Results) were removed
as they generate erroneous results (Pickrell and Pritchard 2012). Using the total data set,
i.e. 286 loci, five migration events were sequentially added to look for the best tree to fit
the data and the number of migration events that reached an asymptotic likelihood was
kept.
284
Outlier detection
The use of several independent methods is often recommended to improve
accuracy of outlier loci detection (Pérez-Figueroa et al. 2010, de Villemereuil et al.
2014). Each method is differently impacted by the genetic structure and/or demographic
history of the study species, which is usually unknown, leading to frequent
inconsistencies across methods (Gagnaire et al. 2015). To cope with these problems,
outlier loci were detected using four different methods. First, BAYESCAN (Foll and
Gaggiotti 2008) is a Bayesian method that uses a logistic regression model to estimate
directly the posterior probability that a given locus is under selection. We used default
parameter values in our analyses to detect outliers among the clusters previously
identified. The second approach (Duforet-Frebourg et al. 2014) implemented in the R
package PCAdapt (Luu et al. 2016) is based on a hierarchical model where population
structure is first depicted using K factors. No *a priori* on the genetic structure (and thus,
no clustering) is required in advance. Loci that are atypically related to population
structure, measured by the K factors, are identified as outliers. For each value of K
(ranging from one to ten), ten replicate chains of 150,000 Monte Carlo Markov Chain
iterations were performed and we discarded the first 5,000 iterations as burn-in. The third
approach uses the estimated co-ancestry matrix to compute an extension of the original
Lewontin and Krakaeur statistic that account for the history of populations under a model
of pure drift (Bonhomme et al. 2010). Finally we used the custom simulation test
described in Fraïsse et al. (2014). The idea of this test is to use simulations of the best-
supported demographic model to obtain the neutral envelope of the joint distribution of
pairwise F_{ST} in a four-population analysis. This test uses the fact that it is easy to have
false positives in each of two pairwise comparisons but that outliers in both comparisons,
against the genome-wide structure, are more likely to be true positives. Roux et al. (2016)
found that an Isolation-with-Migration (IM) model fitted the seahorse data well and that
more parameterized models did not improve the fit. These authors also found that the
time of divergence (Tsplit) was very similar in each pairwise comparison. We therefore
used a four-population IM model with the population sizes and migration rates inferred
by Roux et al. (2016) on the transcriptome data of Romiguier et al. (2014). The two latter

tests are very similar in their spirit, although one use a simple history of divergence with
drift and the other intend to explore more complex demographic histories. They were
here used to identify parallel SNPs that provide a grouping of populations that goes
against the genome-wide trend.
319
Genotype discordance among loci in admixed samples (*i.e.* genomic cline
analysis) was tested using Barton's concordance method as described in Macholán et al.
(2011). The method fits a quadratic function to the relationship between a single locus
hybrid index (0 or 1 if homozygous, 0.5 if heterozygous) and the genome-wide hybrid
index. The function parameters measure the deviation from x = y (*i.e.* the expectation of
homogeneous genome-wide introgression) as a function of the expected heterozygosity.
Instead of testing the deviation from the diagonal, our aim was to compare genomic
clines (*i.e.* regression curves) between geographic samples, as discordance was observed
in one population and not others.
Finally, we tested for gene ontology (GO) terms enrichment to determine if
outlier loci displayed functional enrichment, compared to the full dataset, using Fisher's
Exact Test with Multiple Testing Correction of FDR (Benjamini and Hochberg 1995)
implemented in the software Blast2GO (Conesa and Götz 2008).
333
Results
SNPs characterization/calling: an efficient genotyping method in a protected species with
low genetic diversity
Of the 384 SNPs, including five SNPs that diagnostically distinguished *H.*
guttulatus from *H. hippocampus* (one mitochondrial and four nuclear), a total of 318
SNPs amplified successfully. Of these 318 SNPs, 286 showed a minor allele frequency
above 5%. Two *H. hippocampus* initially identified as *H. guttulatus* were removed from
the initial dataset, resulting in a final dataset of 292 *H. guttulatus* genotyped for 286
SNPs.
In order to evaluate the extent of ascertainment bias that may be induced by our
procedure for selecting markers, we compared the Joint Site-Frequency Spectra (JSFS)
obtained with the original dataset of Romiguier et al. (2014) -assumed freed from
ascertainment bias- with our 314-SNP dataset. JSFSs are detailed in Supporting
Information 1 (Fig. SI1). Briefly, as expected, the data of Romiguier et al. (2014) is
principally composed of singletons that represented 35-40% of the SNPs (Fig. SI1A),
while this proportion was reduced twofold in our 314-SNP dataset (Fig. SI1B). We
efficiently removed rare variants without biasing the frequency spectrum too much as the
deficit of singletons was homogeneously compensated by every other cells of the JSFSs.
An even representation over the entire allele frequency range was indeed observed based
on our dataset. The comparison of the two JSFSs (Fig. SI1C) reveals that very few cells
apart from singletons have an excess above 5%, suggesting limited ascertainment bias in
our SNP panel, except with rare alleles as expected.
With a limited ascertainment bias, with rare alleles being underrepresented, no
missing data, and constraints on our model study (small amount of DNA available with
the use of non-lethal fin-clipping sampling techniques), selecting SNPs characterized
from a preliminary population transcriptomic survey proved to be a straightforward

strategy for genome-wide investigation of the spatial distribution in this species
compared to classical genotyping-by-sequencing approaches.
362
A strong genetic structure delineating five broadly distributed panmictic genetic clusters
Estimates of the mean number of alleles per locus (N_{all}), allelic richness (A_r),
expected heterozygosity (H_e), and departure from Hardy-Weinberg equilibrium (HWE;
F_{IS}), for each study site and each genetic cluster identified, are presented in Supporting
Information 2. The gene diversity was similar among populations with no significant
departure from HWE observed (with the exception of site 6 which is a hybrid population,
see below).
370
371

372
373
374   **Figure 2** Genetic population structure based on 286 SNP markers analyzed by Principal
375   Component Analyses depicting axis 1 (explaining 60.1% of the variance) and axis 2 (explaining
376   16.2% of the variance; left panel) and axes 1 and 3 (explaining 9.9% of the variance; right panel).
377   Samples are numbered and colored according to Fig. 1. Each label shows the barycenter of each
378   study site.
379
The Principal Component Analysis (PCA) revealed clear differentiation
separating five clusters (Fig. 2). Along the first two axes (left panel), explaining 60.1%
and 16.2% of the variance, the PCA showed clear differentiation between North Atlantic
(sites 1-5, in red), South Atlantic (sites 7-12, in blue), Mediterranean Sea (sites 13-25,
lagoon site 16 excluded, in green) and Mediterranean Thau lagoon (site 16, in pink).
Hossegor individuals (site 6, in purple) clustered either with the North Atlantic (12
individuals) or with the South Atlantic groups (3 individuals). Mediterranean sites spread
out along the third axis (9.9% of the variance explained, right panel), with Bizerte (site
20, in gold) being closer to the Mediterranean Thau lagoon on the one hand, and Varna
(Black Sea, site 25, in turquoise) standing out from the Mediterranean group on the other
hand.

This pattern of genetic structure was also detected by the STRUCTURE analysis,
of which the output composed of five clusters is presented in Supporting Information S3.
By selecting a K-value of five, we distinguished identical clusters composed by: 1- the
North Atlantic sites, 2- the South Atlantic sites, 3- the Mediterranean Sea sites, 4- the
Mediterranean Thau lagoon and 5- Varna. Similar to the PCA (Fig. 2), a gradient of
introgression is visible along the Atlantic coasts, with a decreasing proportion of South
Atlantic cluster ancestry North of Hossegor (site 6). While most Hossegor individuals
belong to the North Atlantic genetic background, three individuals were genetically
assigned to the South Atlantic genetic background. Bizerte (site 20), and to a lesser extent
Halkida (site 24), proved to have mixed ancestries from both Mediterranean lagoon and
Sea clusters. Increasing the K-value resulted in additional clusters composed by Bizerte
(site 20, K=6), Halkida (site 23, K=7), but then (K>7) no individuals were completely
assigned to any new cluster.
Altogether, distance-based (PCA) and model-based (Structure) analyses supported
the identification of five clusters, a pattern also showed by the visualization of raw mutli-
locus genotype data (Supporting Fig. SI3B). This representation illustrates that most
markers contribute to the signal of five genetic clusters.
Importantly, no significant departure from panmixia was observed within each
cluster (SI2). Furthermore, genetic homogeneity was observed between sites within each
cluster (SI2). In contrast, Fisher exact tests revealed significant differences in allelic
frequencies among clusters (p-value < 0.001), with significant differentiation being
observed for all pairwise comparisons (F_{ST} ranging from 0.024 between the
Mediterranean Sea cluster and Halkida to 0.260 between the North Atlantic cluster and
Varna).
415

418   **Figure 3** Population trees inferred by TREEMIX (A) without or (B) with 3 migration events.
419   Admixture arrows are colored according to the migration weight. The model including three
420   admixture events significantly improved the fit as compared to a situation without migration (p-
421   value < 0.001). Panmictic clusters are colored according to Fig. 1.
422

Finally, the population tree inferred using TREEMIX without accounting for migration
(Fig. 3A) was highly consistent with all above analyses. Atlantic clusters branching
together on one hand, and Mediterranean lagoons (Thau and Bizerte) branching together
on the other hand. Interestingly, three admixture events significantly improved the model
as compared to a scenario without migration (p-value < 0.001; Fig. 3B). This population
tree indicated significant gene flow among four *H. guttulatus* clusters, between the north

and the south genetic clusters in the Atlantic coasts, in concordance with the gradient of
introgression along the Atlantic coasts (Fig. 2 and SI3), between marine and lagoon
samples in the Mediterranean Sea, and finally between the North Atlantic and
Mediterranean lagoon samples. We suggest that this latter observation is most probably
mainly driven by parallel outlier loci (see below), either as a result of adaptive
introgression or as a consequence of a shared history of divergence retained at outlier loci
against massive secondary gene flow.
436
Signature of selection and genetic parallelism
Nine SNPs out of 286 (3.15%) were consistently identified to depart from
neutrality with the four tests (BayeScan, PCAdapt, FLK and custom simulation test).
Interestingly, six of them showed very similar allele frequency patterns, distinguishing
North Atlantic sites, Mediterranean Thau lagoon, Bizerte (site 20) and, in a lesser extent
Halkida (site 23) and Varna (site 25) from South Atlantic and Mediterranean Sea sites
(Fig. 4), and pointing out genetic parallelism (*i.e.* convergence of allele frequency
patterns) between these lineages.
445

446
447 **Figure 4** *H. guttulatus* allele frequencies (y-axis) for the six outliers with very similar
448 high allele frequency. Each study site (x-axis) is labeled and colored according to Fig. 1,
449 reminded by a simplified map on the right. Hossegor was separated in 6A and 6B and Halkida in
450 24A and 24B according to their North or South Atlantic / Mediterranean Thau lagoon or Sea
451 genetic background, respectively (see Fig. 2, SI3).
452
These six outliers located on different *H. guttulatus* contigs were located on different
Hippocampus comes scaffolds, except SNPs 29 and 286 mapping to a unique *H. comes*
scaffold (Fig. 5A). Interestingly, these scaffolds –that contain outliers- consistently
mapped to a unique chromosome in *Syngnathus scovelli* (Fig. 5A). Results were similar
when directly blasting these six *H. guttulatus* contigs against *S. scovelli* genome, but with
SNP 29 mapping to an unplaced scaffold (Fig. 5A). A unique chromosome is still
involved when blatting *H. guttulatus* outlying contigs against seven additional well-
assembled fish genomes, in agreement with a well-conserved synteny of fishes (detailed
in SI5).
462

463
464
Figure 5 Outlier cross-mapping (A-) and F_{ST} between the Mediterranean lagoon and Sea
lineages (in black), and between the North and South Atlantic lineages (in grey) along the unique
chromosome (B-). As *H. comes* scaffolds are unplaced, *H. guttulatus* SNPs were first blasted on
H. comes scaffolds, then each *H. comes* scaffold was mapped to *S. scovelli* genome. A putative *H.*
comes chromosome was hence reconstructed (A-). The order of the scaffolds and SNPs according
to the blasts was conserved. Outlier SNPs displaying parallel differentiation between Atlantic
lineages and Mediterranean ecotypes are colored in orange, while the grey outlier showed high
differentiation between the Mediterranean Thau lagoon and other sites.
473
Genetic parallelism was visualized by plotting F_{ST} of Mediterranean lagoon and
sea locations (x-axis in Fig. 6) against F_{ST} between North and South Atlantic clusters (y-
axis in Fig. 6). Outliers showing genetic parallelism appeared in the top right of Figure 6,
showing elevated genetic differentiation between Mediterranean lagoon and marine sites
(x-axis) on the one hand, and between North and South Atlantic sites (y-axis) on the
other hand. This F_{ST} - F_{ST} co-plot also showed how the parallel pattern of differentiation
observed at these six loci would be highly unlikely under the inferred demographic
history of populations.
482

483
484 **Figure 6** Genome scan of infra-specific differentiation in *H. guttulatus*. Dashed lines represent
485 the 95% and 99% quantiles of the neutral envelope of F_{ST} obtained following Fraïsse et al.
486 (2014). Loci identified by all methods as outliers are colored in orange – the six outliers that
487 displayed parallel differentiation between Atlantic lineages and Mediterranean ecotypes – and in
488 black – the three other outliers.
489
Three other outliers were also consistently identified (SNPs 87, 130 and 140; Fig.
6 and Supporting Information SI4) using all four methods. They distinguished either the
Mediterranean Thau lagoon (*i.e.* high genetic differentiation along the x-axis in Fig. 6;
SNP 87), the North Atlantic sites or the South Atlantic sites (*i.e.* high genetic
differentiation along the y-axis in Fig. 6; SNPs 130 and 140) from all other sites. By
using a similar approach for convergent outliers, we only observed SNP 87 that
consistently mapped in the unique chromosome.
497

Figure 7 shows the genomic cline analysis obtained with SNP 29, an outlier that
clustered on the unique chromosome (Fig. 5 and SI5), and SNP 130, an outlier between
the northern and the southern Atlantic lineage but that mapped to another chromosome
and not differentiated in the Mediterranean Sea (no genetic parallelism). Regressions
were found to be different at SNP 29 and other outliers that clustered on the same
chromosome, when the Hossegor sample was included in the analysis or not (Fig. 7),
while regressions were always close to the diagonal with SNP 130 and other outliers that
mapped to other chromosomes. Introgression did not occur at the same rate for the loci
mapping to the unique chromosome, with a clear discordance between the clustered
outlier loci and the rest of the genome in the Hossegor sample, in which North Atlantic-
like seahorses have the South Atlantic allele at the genomic island outliers.
518
519   Note that of all the 168 fish sexed (out of 292 fishes), comprising individuals
520 from each of the five lineage and contact zones in a balanced sex ratio, loci with extreme
521 level of differentiation could not be related to any sex differences. In addition, the
522 analysis of gene ontology terms for outlier loci did not reveal any significant functional
523 enrichment.
524
525

Discussion
Genetic analyses of the long-snouted seahorse revealed cryptic discrete panmictic
genetic lineages that meet either in a narrow contact zone in the Atlantic, or display a
mosaic distribution associated with environmental variation in the Mediterranean Sea.

Despite limited dispersal abilities and seemingly small population sizes (but see Curtis
and Vincent 2006), each lineage showed remarkable genetic homogeneity over very large
distances, with genetic panmixia observed within each lineage. This spatial structure,
with strong and sharp genetic subdivisions, is not expected if genetic drift was
predominantly responsible for the genetic differentiation between these populations.
The spatial organization of the different genetic backgrounds proved to be an
unusual combination of geographic subdivision in the Atlantic Ocean, and genetic
structure related to the sea-lagoon ecological contrast in the Mediterranean Sea.
Surprisingly, these two subdivisions partly relied on the same genetic architecture. We
observed genetic parallelism at some markers showing extreme levels of differentiation
between habitats in the Mediterranean Sea, but also between geographic lineages in the
Atlantic Ocean. Intriguingly, all the loci showing convergent allele frequency patterns
mapped to a single chromosome. This tends either to suggest genetic differences do not
accumulate evenly across the genome, or that these genomic regions are easier to
characterize with low-density genome scans.
We suggest the existence of a shared evolutionary history between Atlantic
parapatric lineages and Mediterranean ecotypes, with the Mediterranean lagoon ecotype
anciently related to the North-Atlantic lineage. The underlying reproductive isolation
mechanisms may involve a combination of intrinsic and extrinsic genetic barriers, where
relative contributions may differ between the two contexts.
550
1- Genome scans in hybrid zones
Our SNP panel allowed us to define five discrete panmictic genetic clusters, two
in the Atlantic Ocean, two in the Mediterranean Sea, and one in the Black Sea, which
cannot be morphologically distinguished with reliability so far (*i.e.* cryptic lineages). The
average genetic differentiation between clusters and its associated variance were strong
($0.09 \pm 0.02 < F_{ST} \pm$ sd $< 0.26 \pm 0.04$). In this context, identifying outlier loci is a
complex task with problems of false positives (Lotterhos and Whitlock 2014, 2015). It is
also increasingly being acknowledged that a signal of local adaptation is not easily
captured without a broad sampling of the genome in a standard infra-specific low-linkage
disequilibrium context (Hoban et al. 2016). Finding easily nine well-supported outlier
loci with a limited number of loci suggests strong variance in differentiation levels
associated with the existence of cryptic genetic barriers involving many selected loci
(Bierne et al. 2011). Extensive linkage disequilibrium is also maintained in this complex
of semi-isolated genetic backgrounds when compared to a standard infra-specific context.
However, this is not expected to be a rare situation as genomic studies have provided
accumulating evidence that semi-permeable barriers to gene flow are widespread (Roux
et al. 2016) and affect a substantial proportion of genomes (Harrison and Larson 2016).
In this study, six of the nine outliers showed a signature of genetic parallelism
(Fig. 2, Fig. SI3). These six outliers not only proved to be the most differentiated loci
using a pairwise comparison between Northern and Southern lineages in the Atlantic
Ocean and between the lagoon and marine ecotypes in the Mediterranean Sea (Fig. 6),
but they also displayed a genetic structure that is not compatible with the sample
covariance matrix inferred with the full set of loci (FLK test, Supporting Information SI6)
or with simulations under the best-supported demographic model inferred from
transcriptome data (Roux et al. 2016). Finally, these six outliers (3.15% of the SNPs

analyzed) proved to map to the same chromosome (4.5% of the genome). These six loci
mapped to the same chromosome in available fish genomes, although the order on the
chromosome was not so well conserved in distantly related species. This result suggests
the existence of a large genomic island of differentiation as already reported in other
fishes (e.g. sticklebacks: Jones et al. 2012, cod: Hemmer-Hansen et al. 2013, Berg et al.
2017, seabass: Duranton et al. 2018), and possibly involving a chromosomal inversion or
another form of recombination suppression. This chromosome-wide island of
differentiation also explains why we detected the signal of parallel outlying
differentiation with a moderate number of loci analyzed. The chromosome could have
been easily missed with a few microsatellites but our SNP panel was largely enough to
have a few SNPs on the chromosome. These preliminary results call for genome
sequence analysis in order to better characterize the genomic island and its structural
variation, and to investigate the genome localization of additional barrier loci with a
smaller chromosomal footprint.
600
2- Parapatrically distributed semi-isolated cryptic lineages in the Atlantic Ocean
In the Atlantic Ocean we confirmed the existence of two well-differentiated
genetic clusters previously identified by Woodall et al. (2015) with five microsatellites
and two mitochondrial genes. A genetic homogeneity was observed over large distances
within each lineage (Fig. 1), contrasting with a very strong and abrupt genetic
differentiation between them. Inter-lineage genetic divergence was not only captured by
the most differentiated SNP that were nearly fixed between the two lineages (SNP 130 in
SI4, F_{ST} = 0.9), but also, by many other SNPs (SI3B), suggesting isolating mechanisms
and loci with genome-wide effects. This mechanism is also suggested by the discovery of
two coexisting lineages in the same lagoon in South West of France without early
generation hybrids sampled (Hossegor lagoon, site 6 in Fig. 1). Such zone of co-existence
in sympatry contrasting with genetic homogeneity at a large spatial scale within each
lineage suggests strong reproductive isolation (Jiggins and Mallet 2000), involving either
pre-zygotic isolation or strong selection against hybrids at early life stages.
Recent migration of the two diverged genetic backgrounds in this lagoon does not
easily explain the co-existence of the two lineages with no hybridization. In addition to
the low probability of sampling the first-generation migrants without any genomic
modifications, local introgression of the North-Atlantic seahorse in this lagoon (Fig. 3)
attests hybridization occurred and that a mechanism of isolation must maintain the
divergence. The decreasing proportion of south lineage ancestry from Hossegor to the
English Channel also provides indirect evidence for asymmetric introgression (Fig. 2,
SI3A). In addition, an interesting pattern was observed at genomic island outlier loci in
Hossegor (site 6) that further support partial reproductive isolation. The North Atlantic
lineage mostly carries the southern alleles at genomic island loci (pale pink dots in Fig. 4,
see also SI3B). To produce such North Atlantic-type individuals with a South-Atlantic
island, recombination between parental backgrounds is required, suggesting hybridization
occurred in this case. In the genomic cline analyses (Fig. 7) the geographic information
was included to contrast the results obtained with or without the Hossegor sample. The
objective of this analysis was to infer the existence of a local discordance in this site. The
swamping of North Atlantic fishes by the southern allele at the genomic island suggests
epistatic or coupling interactions among loci implied in reproductive isolation.

Endogenous post-zygotic selection against hybrids does not result in a stable
polymorphism in a single isolated population, but instead in transient polymorphisms.
Also known as bistable variants, the maintenance of underdominant or epistatically
interacting genetic incompatibilities requires a migration-selection balance in a spatially
subdivided population (Barton and Turelli 2011), or frequency-dependent selection
(Barton and de Cara 2009). The system may have fixed one state of a bistable variant in
the Hossegor lagoon. Alternatively, the southern allele may confer a selective advantage
in this lagoon whatever the genetic background. However the latter hypothesis does not
seem coherent as the northern allele is always in higher frequency in lagoon-like habitats
elsewhere, though we may have missed more subtle ecological parameters that would
have driven this pattern.
Overall, the Atlantic contact zone possesses all the characteristics of standard
clinal hybrid zones that follow the tension zone model (Barton and Hewitt 1985), *i.e.* a
secondary contact zone maintained by a balance between migration and intrinsic
reproductive isolation. Exogenous selection may also contribute, although the two
Atlantic lineages both inhabit indifferently lagoon and sea habitats (Fig. 1), and co-exist
in syntopy in Hossegor, suggesting a limited role of ecological contrast in their genetics.
In any case, only strong intrinsic reproductive isolation can guarantee a genome-wide
barrier to gene flow explaining the co-existence of the two lineages in sympatry in the
Hossegor lagoon.
652
3- Sea and lagoon ecotypes in the Mediterranean Sea
Contrasting with the Atlantic hybrid zone, the two cryptic lineages identified in
the Mediterranean Sea were associated with lagoon/sea ecosystem variation. Our broad
genomic and spatial sampling revealed two crucial observations. First, while the marine
lineage was surprisingly homogeneous over the whole Mediterranean Sea, from Greece
to Spain (F_{IT} = 0.0078 n.s.), lagoon-like samples, especially the Thau lagoon, showed a
strong and genome-wide genetic differentiation from them. Samples from two lagoons
(Thau in France and Bizerte in Tunisia, sites 16 and 20 in Fig. 1) were sufficient to
reveal an association with the environment that was previously unseen, the Thau lagoon
being the only sample from Western Mediterranean basin in Woodall et al. (2015).
Second, fixed differences between lagoon and marine samples were observed, although
they were sampled only few kilometers apart (e.g. sites 14-16 in Fig. 2, 4). A single but
important seahorse sampled on the seaside of the Thau lagoon (site 15 in Fig. 1), plus
seven others sampled on the seaside of another lagoon of the region (site 14 in Fig. 1)
proved to belong to the marine genetic cluster, without any sign of introgression.
Likewise, no evidence of introgression was observed in the Thau sample (Fig. 3). Once
again, despite genetic homogeneity over large area, such strong and abrupt genetic
differentiation suggests partial reproductive isolation between these two lineages. In this
case there are obvious ecological drivers, *i.e.* habitat specialization. Indeed, the Bizerte
lagoon (site 20 in Fig. 1), which is ecologically similar to the Thau lagoon (Sakka Hlaili
et al. 2008), has a population genetically similar to marine Mediterranean samples at
most loci and only share the genetic composition of the Thau lagoon at the genomic
island (Fig. 4 and SI3B). In addition, a subsample of the Halkida population (Greece, site
24B in Fig. 4) was composed of five individuals heterozygous at the genomic island. The
environmental parameters at this location are hypothesized to be more lagoon-like, being

a secluded bay beyond the northern end of the Euipus Strait. This sample only provides
evidence that the genomic island polymorphism and the mosaic spatial structure extends
to the eastern basin without really providing further clues about the role of the
environment.
Our results from the Mediterranean Sea (*i.e.* lagoon/marine system) resemble those
obtained from the emblematic three-spined sticklebacks marine/freshwater system (Jones
et al. 2012), or more recently from the marine-migratory/freshwater-resident lampreys
(Rougemont et al. 2016) and coastal/marine ecotypes in European anchovies (Le Moan
et al. 2016). When a shared divergent genomic island is observed –*i.e.* genetic
parallelism, as here in long-snouted seahorse among the Mediterranean lagoons, or in the
examples cited above, there could be three possible interpretations: (i) parallel gene
reuse from a shared ancestral polymorphism present in the marine supposedly ancestral
population (Jones et al. 2012), (ii) the spread of a locally adapted allele (*i.e.* the
'transporter hypothesis'; Schluter and Conte 2009) or (iii) secondary contact followed by
spatial reassortment of the divergent lineages and extensive introgression swamping such
that only selected loci and their chromosomal neighborhood retain the history of
adaptation (Bierne et al. 2013). The three scenarios are difficult to discriminate as they
converge toward a similar pattern (Johannesson et al. 2010, Bierne et al. 2013, Welch
and Jiggins 2014). Here, as for the lampreys (Rougemont et al. 2016), the Thau lagoon
provides a possible support for the secondary contact model because the differentiation,
although stronger at the genomic island, is genome-wide. The Bizerte lagoon however
can either be interpreted as a marine lineage introgressed by the lagoon allele at the
genomic island, or as a lagoon lineage (defined by adaptive/speciation genes) massively
introgressed by neutral marine alleles. Incorporating heterogeneous migration rates in
demographic inference methods allowed Le Moan et al. (2016), Rougeux et al. (2016)
and Rougemont et al. (2016) to identify the signal of a secondary contact history carried
by islands of differentiation in lampreys, white fishes and anchovies. Unfortunately our
286-SNPs dataset does not allow performing such historical demographic reconstruction.
Nonetheless the TREEMIX analysis reveals that episodes of secondary admixtures
strongly improve the fit to the sample covariance matrix, but adaptive introgression or
massive introgression swamping can both explain them. Anyhow, demographic
reconstruction does not completely refute the 'transporter hypothesis' which stipulates
lagoon alleles spread from lagoon to lagoon (or freshwater allele from river to river) and
is a scenario that produces a very similar genomic pattern of differentiation to the one
produced by a standard secondary contact (Bierne et al. 2013, Rougemont et al. 2016). In
the case of the seahorse complex, however, we made the new observation that genetic
parallelism is observed with the Atlantic populations where the structure is geographic
and independent of the lagoon-sea habitats, which offers complementary arguments to
the debate.
717
4- Genetic parallelism in two different spatial/ecological contexts
The most astonishing result of our genetic analysis was genetic parallelism
between the Mediterranean lagoon ecotype and the north Atlantic lineage at a large
genomic island. Parallel evolution implies distinct but repetitive ecological characteristics
(e.g. Butlin et al. 2014). However, this claim comes from an inductive reasoning,
validated by many observations, which a single counter-example can yet alone refute. In

the present study, we found that the genomic island was associated with the sea-lagoon
ecological contrast in the Mediterranean Sea, while there was no such genetic
differentiation between lagoon and sea samples in the Atlantic Ocean. The North and
South lineages inhabit indifferently lagoons and seas, so that what seemed obvious in the
Mediterranean Sea regarding the divergence of the two lineages, *i.e.* habitat
specialization, was not observed along the Atlantic Ocean where the divergence seems
uncorrelated to the ecological contrasts that explain the two Mediterranean ecotypes.
Besides, abiotic parameters such as temperature or salinity, did not allow bounding the
North Atlantic and Mediterranean lagoon lineages genetic parallelism. Although we may
have missed putative ecological drivers of such genetic parallelism, parallel gene reuse
driven by ecological convergence seems here unlikely. Shared ancestral polymorphism
sieved by adaptation in a patchy environment (Bierne et al. 2013) and incipient speciation
(Guerrero and Hahn 2017) would better explain our data. The association with habitat in
the Mediterranean Sea and with space in the Atlantic Ocean could be explained by a
secondary evolution of locally adapted genes within the genomic island in the
Mediterranean Sea, benefiting from the barrier to gene flow imposed by intrinsic
selection (divergence hitchhiking; Via 2012, Yeaman 2013). Alternatively, the genomic
island could have coupled with local adaptation polymorphisms localized elsewhere in
the genome in the Mediterranean Sea (Bierne et al. 2011), while it would have been
trapped by a barrier to dispersal in the Atlantic Ocean (Barton 1979, Barton and Hewitt
1985). Without further data and the true genomic position of loci in the seahorse genome,
it is difficult to disentangle the two hypotheses.
One locus (SNP 87) localized in the same chromosome as parallel SNPs (Fig. 5)
is differentiated between the marine and lagoon ecotypes in the Mediterranean Sea, but is
not differentiated between the northern and southern lineages in the Atlantic Ocean. At
first sight, it could be interpreted as evidence for a possible secondary local sweep in
Mediterranean lagoons. However, according to the gene order inferred from the closest
species (Gulf pipefish and Tiger tail seahorse) along with a barrier to gene flow less
effective in the Atlantic (see SNPs 34 and 212 in Fig. 4 and 5B), it could also be
interpreted as being localized in the island "shoulders" (*i.e.* loci in the vicinity of the
regions harboring local adaptation and/or reproductive isolation loci; Gagnaire et al.
2015, Le Moan et al. 2016) in which a stronger introgression rate has erased the
differentiation faster in the Atlantic than in the Mediterranean, in favor of the alternative
interpretation. Importantly, whatever the explanation -divergence hitchhiking or
coupling- it requires invoking interaction between intrinsic and ecological selection and
not ecological selection alone (Bierne et al. 2011, Kulmuni and Westram 2017).
Furthermore, intrinsic selection has most probably evolved first in this system as no
genetic parallelism was observed in outliers discriminating lagoon to marine ecotypes,
which would contradict the predominant view that ecological selection is necessarily the
initial catalyzer of the chain of accumulation of barriers in ecological speciation.
764
765 **Conclusions**
766 Analyzing the population genetics of the long-snouted seahorse *Hippocampus guttulatus*
767 revealed a complex of panmictic genetic backgrounds subdivided by sharp semi-
768 permeable hybrid zones. This is now standard observation in marine species (Knowlton
769 1993, Pante et al. 2015, Sheets et al. 2018) where morphological stasis might be more

widespread than in terrestrial species. The subdivision of species by hybrid zones is a long-lasting observation in the terrestrial realm (Hewitt 1989) but arguably more readily detected by morphological differences. We also easily found outlier loci despite a moderate number of loci analyzed, and the clustering of these outlier loci in a single genomic region. This also tends to become a standard observation of the recent hybrid zone literature (e.g. in stickelbacks, Jones et al. 2012; jaera, Ribardière et al. 2017; cod, Hemmer-Hansen et al. 2013).

However, we also made two additional observations that are less commonly reported and deserve a broader interest outside the study of seahorse themselves, although are important to further inform captive breeding and *in-situ* management decisions. First, we found the two, usually opposed, standard spatial structures of the hybrid zone literature, namely clinal and mosaic hybrid zones, in the same study system. This result calls for further investigations with lab and fieldwork in order to better understand the mechanisms of reproductive isolation at play and their genetic architecture. Secondly, we found a parallel pattern of differentiation at the genomic island in the two spatial/ecological contexts. Although this result will also need to be substantiated by follow-up genomic studies, it nonetheless reveals that the hallmark of ecologically driven adaptive divergence can be observed in absence of obvious ecological convergence. We argue that alternative scenarios involving secondary introgression swamping and intrinsic isolation should be more seriously considered as valid alternatives and the seahorse complex could become an interesting flagship system in the debate.

Acknowledgments
We are grateful to Lucas Beranger, Michel Cantou, Philippe Lenfant, Pablo Liger, CPIE Bassin de Thau, Patrick Lelong, Francesco Di Liello and Stéphane Auffret for their help in providing *H. guttulatus* fin-clippings and to Fabienne Moreau for the BeadXpress experiment. Many thanks to Nicolas Duforet-Frebourg and Christelle Fraïsse for computational advice. This work was funded by a Languedoc-Roussillon Region "Chercheur(se)s d'avenir" grant to NB (Connect7 project), by a LabEx CeMEB postdoctoral fellowship to FR and by Chocolaterie Guylian and a Natural Environment Research Council Industrial Case studentship (NER/S/C/2005/13461) to LW.

Literature
Barton, N. 1979. The dynamics of hybrid zones. Heredity 43:341–359.

Barton, N. H., and M. A. R. de Cara. 2009. The evolution of strong reproductive isolation. Evolution 63:1171–1190.

Barton, N. H., and G. M. Hewitt. 1985. Analysis of hybrid zones. Annu. Rev. Ecol. Syst. 16:113–148.

Barton, N. H., and M. Turelli. 2011. Spatial waves of advance with bistable dynamics: cytoplasmic and genetic analogues of Allee effects. Am. Nat. 178:E48–E75.

814   Bhagwat, M., L. Young, and R. R. Robison. 2012. Using BLAT to find sequence
815   similarity in closely related genomes. Curr. Protoc. Bioinforma. Ed. Board Andreas
816   Baxevanis Al 0 10:Unit10.8
817
818   Benjamini, Y., and Y. Hochberg. 1995. Controlling the false discovery rate: a practical
819   and powerful approach to multiple testing. J. R. Stat. Soc. Ser. B Methodol. 57:289–300.
820
821   Berg, P. R., B. Star, C. Pampoulie, I. R. Bradbury, P. Bentzen, J. A. Hutchings, S. Jentoft,
822   and K. S. Jakobsen. 2017. Trans-oceanic genomic divergence of Atlantic cod ecotypes is
823   associated with large inversions. Heredity, doi: 10.1038/hdy.2017.54.
824
825   Bierne, N., P. Borsa, C. Daguin, D. Jollivet, F. Viard, F. Bonhomme, and P. David. 2003.
826   Introgression patterns in the mosaic hybrid zone between *Mytilus edulis* and *M.*
827   *galloprovincialis*. Mol. Ecol. 12:447–461.
828
829   Bierne, N., Gagnaire, P.-A., and P. David. 2013. The geography of introgression in a
830   patchy environment and the thorn in the side of ecological speciation. Curr Zool, 59: 72–
831   86.
832
833   Bierne, N., J. Welch, E. Loire, F. Bonhomme, and P. David. 2011. The coupling
834   hypothesis: why genome scans may fail to map local adaptation genes. Mol. Ecol.
835   20:2044–2072.
836
837   Bonhomme, M., C. Chevalet, B. Servin, S. Boitard, J. Abdallah, S. Blott, and M.
838   SanCristobal. 2010. Detecting selection in population trees: the Lewontin and Krakauer
839   test extended. Genetics 186:241–262.
840
841   Bouchemousse, S., C. Liautard-Haag, N. Bierne, and F. Viard. 2016. Distinguishing
842   contemporary hybridization from past introgression with postgenomic ancestry-
843   informative SNPs in strongly differentiated *Ciona* species. Mol. Ecol. 25:5527–5542.
844
845   Boursot, P., J.-C. Auffray, J. Britton-Davidian, and F. Bonhomme. 1993. The evolution
846   of house mice. Annu. Rev. Ecol. Syst. 24:119–152.
847
Brawand, D., C. E. Wagner, Y. I. Li, M. Malinsky, I. Keller, S. Fan, O. Simakov, A. Y.
Ng, Z. W. Lim, E. Bezault, J. Turner-Maier, J. Johnson, R. Alcazar, H. J. Noh, P. Russell,
B. Aken, J. Alföldi, C. Amemiya, N. Azzouzi, J.-F. Baroiller, F. Barloy-Hubler, A.
Berlin, R. Bloomquist, K. L. Carleton, M. A. Conte, H. D'Cotta, O. Eshel, L. Gaffney, F.
Galibert, H. F. Gante, S. Gnerre, L. Greuter, R. Guyon, N. S. Haddad, W. Haerty, R. M.
Harris, H. A. Hofmann, T. Hourlier, G. Hulata, D. B. Jaffe, M. Lara, A. P. Lee, I.
MacCallum, S. Mwaiko, M. Nikaido, H. Nishihara, C. Ozouf-Costaz, D. J. Penman, D.
Przybylski, M. Rakotomanga, S. C. P. Renn, F. J. Ribeiro, M. Ron, W. Salzburger, L.
Sanchez-Pulido, M. E. Santos, S. Searle, T. Sharpe, R. Swofford, F. J. Tan, L. Williams,
S. Young, S. Yin, N. Okada, T. D. Kocher, E. A. Miska, E. S. Lander, B. Venkatesh, R.
D. Fernald, A. Meyer, C. P. Ponting, J. T. Streelman, K. Lindblad-Toh, O. Seehausen,

and F. Di Palma. 2014. The genomic substrate for adaptive radiation in African cichlid
fish. Nature 513:375–381.
861
862 Britton-Davidian, J., F. Fel-Clair, J. Lopez, P. Alibert, and P. Boursot. 2005. Postzygotic
863 isolation between the two European subspecies of the house mouse: estimates from
864 fertility patterns in wild and laboratory-bred hybrids. Biol. J. Linn. Soc. 84:379–393.
865
866 Butlin, R. K., M. Saura, G. Charrier, B. Jackson, C. André, A. Caballero, J. A. Coyne, J.
867 Galindo, J. W. Grahame, J. Hollander, P. Kemppainen, M. Martínez-Fernández, M.
868 Panova, H. Quesada, K. Johannesson, and E. Rolán-Alvarez. 2014. Parallel evolution of
869 local adaptation and reproductive isolation in the face of gene flow. Evolution 68:935–
870 949.
871
872 Christe, C., K. N. Stölting, L. Bresadola, B. Fussi, B. Heinze, D. Wegmann, and C.
873 Lexer. 2016. Selection against recombinant hybrids maintains reproductive isolation in
874 hybridizing *Populus* species despite F1 fertility and recurrent gene flow. Mol. Ecol.
875 25:2482-2498.
876
877 Conesa, A., and S. Götz. 2008. Blast2GO: A comprehensive suite for functional analysis
878 in plant genomics. Int. J. Plant Genomics 2008.
879
880 Curtis, J. M. R., and A. C. J. Vincent. 2006. Life history of an unusual marine fish:
881 survival, growth and movement patterns of *Hippocampus guttulatus* Cuvier 1829. J. Fish
882 Biol. 68:707–733.
883
884 de Villemereuil, P., É. Frichot, É. Bazin, O. François, and O. E. Gaggiotti. 2014. Genome
885 scan methods against more complex models: when and how much should we trust them?
886 Mol. Ecol. 23:2006–2019.
887
888 Doyle, J., and J. Doyle. 1987. A rapid DNA isolation procedure for small quantities of
889 fresh leaf tissue. Phytochem. Bull. 19:11–15.
890
891 Duforet-Frebourg, N., E. Bazin, and M. G. B. Blum. 2014. Genome scans for detecting
892 footprints of local adaptation using a Bayesian factor model. Mol. Biol. Evol. 31:2483–
893 2495.
894
895 Duranton, M., F. Allal, C. Fraïsse, N. Bierne, F. Bonhomme, and P.-A. Gagnaire. 2018.
896 The origin and remolding of genomic islands of differentiation in the European sea bass.
897 Nat. Commun. 9:2518.
898
899 Duvaux, L., K. Belkhir, M. Boulesteix, and P. Boursot. 2011. Isolation and gene flow:
900 inferring the speciation history of European house mice. Mol. Ecol. 20:5248–5264.
901
Falush, D., M. Stephens, and J. K. Pritchard. 2003. Inference of population structure
using multilocus genotype data: linked loci and correlated allele frequencies. Genetics
164:1567–1587.

Foll, M., and O. Gaggiotti. 2008. A Genome-Scan Method to Identify Selected Loci
Appropriate for Both Dominant and Codominant Markers: A Bayesian Perspective.
Genetics 180:977–993.

Fraïsse, C., C. Roux, J. J. Welch, and N. Bierne. 2014. Gene-flow in a mosaic hybrid
zone: is local introgression adaptive? Genetics 197:939–951.

Gagnaire, P.-A., T. Broquet, D. Aurelle, F. Viard, A. Souissi, F. Bonhomme, S. Arnaud-
Haond, and N. Bierne. 2015. Using neutral, selected, and hitchhiker loci to assess
connectivity of marine populations in the genomic era. Evol. Appl. 8:769:786.

Gayral, P., J. Melo-Ferreira, S. Glémin, N. Bierne, M. Carneiro, B. Nabholz, J. M.
Lourenco, P. C. Alves, M. Ballenghien, N. Faivre, K. Belkhir, V. Cahais, E. Loire, A.
Bernard, and N. Galtier. 2013. Reference-free population genomics from next-generation
transcriptome data and the vertebrate–invertebrate gap. PLOS Genet. 9:e1003457.

Gompert, Z., and C. Alex Buerkle. 2010. INTROGRESS: a software package for
mapping components of isolation in hybrids. Mol. Ecol. Resour. 10:378–384.

Good, J. M., M. A. Handel, and M. W. Nachman. 2008. Asymmetry and polymorphism
of hybrid male sterility during the early stages of speciation in house mice. Evol. Int. J.
Org. Evol. 62:50–65.

Guerrero, R., and M. W. Hahn. 2017. Speciation as a sieve for ancestral polymorphism.
bioRxiv 155176.

Harrison, R. G. 1993. Hybrid zones and the evolutionary process. Oxford University
Press.

Harrison, R. G., and E. L. Larson. 2016. Heterogeneous genome divergence, differential
introgression, and the origin and structure of hybrid zones. Mol. Ecol. 25:2454:2466.

Hedrick, P. W. 2013. Adaptive introgression in animals: examples and comparison to
new mutation and standing variation as sources of adaptive variation. Mol. Ecol.
22:4606–4618.

Hemmer-Hansen, J., E. E. Nielsen, N. O. Therkildsen, M. I. Taylor, R. Ogden, A. J.
Geffen, D. Bekkevold, S. Helyar, C. Pampoulie, T. Johansen, FishPopTrace Consortium,
and G. R. Carvalho. 2013. A genomic island linked to ecotype divergence in Atlantic
cod. Mol. Ecol. 22:2653–2667.

Hewitt, G. M. 1988. Hybrid zones-natural laboratories for evolutionary studies. Trends
Ecol. Evol. 3:158–167.

Hewitt, G. M. 1989. The subdivision of species by hybrid zones. *Speciation and its*
Consequences, 85-110.
952
953   Hoban, S., J. L. Kelley, K. E. Lotterhos, M. F. Antolin, G. Bradburd, D. B. Lowry, M. L.
954   Poss, L. K. Reed, A. Storfer, and M. C. Whitlock. 2016. Finding the genomic basis of
955   local adaptation: pitfalls, practical solutions, and future directions. Am. Nat. 188:379–
956   397.
957
Howe, K., M. D. Clark, C. F. Torroja, J. Torrance, C. Berthelot, M. Muffato, J. E.
Collins, S. Humphray, K. McLaren, L. Matthews, S. McLaren, I. Sealy, M. Caccamo, C.
Churcher, C. Scott, J. C. Barrett, R. Koch, G.-J. Rauch, S. White, W. Chow, B. Kilian, L.
T. Quintais, J. A. Guerra-Assunção, Y. Zhou, Y. Gu, J. Yen, J.-H. Vogel, T. Eyre, S.
Redmond, R. Banerjee, J. Chi, B. Fu, E. Langley, S. F. Maguire, G. K. Laird, D. Lloyd,
E. Kenyon, S. Donaldson, H. Sehra, J. Almeida-King, J. Loveland, S. Trevanion, M.
Jones, M. Quail, D. Willey, A. Hunt, J. Burton, S. Sims, K. McLay, B. Plumb, J. Davis,
C. Clee, K. Oliver, R. Clark, C. Riddle, D. Elliot, D. Eliott, G. Threadgold, G. Harden, D.
Ware, S. Begum, B. Mortimore, B. Mortimer, G. Kerry, P. Heath, B. Phillimore, A.
Tracey, N. Corby, M. Dunn, C. Johnson, J. Wood, S. Clark, S. Pelan, G. Griffiths, M.
Smith, R. Glithero, P. Howden, N. Barker, C. Lloyd, C. Stevens, J. Harley, K. Holt, G.
Panagiotidis, J. Lovell, H. Beasley, C. Henderson, D. Gordon, K. Auger, D. Wright, J.
Collins, C. Raisen, L. Dyer, K. Leung, L. Robertson, K. Ambridge, D. Leongamornlert,
S. McGuire, R. Gilderthorp, C. Griffiths, D. Manthravadi, S. Nichol, G. Barker, S.
Whitehead, M. Kay, J. Brown, C. Murnane, E. Gray, M. Humphries, N. Sycamore, D.
Barker, D. Saunders, J. Wallis, A. Babbage, S. Hammond, M. Mashreghi-Mohammadi,
L. Barr, S. Martin, P. Wray, A. Ellington, N. Matthews, M. Ellwood, R. Woodmansey, G.
Clark, J. D. Cooper, J. Cooper, A. Tromans, D. Grafham, C. Skuce, R. Pandian, R.
Andrews, E. Harrison, A. Kimberley, J. Garnett, N. Fosker, R. Hall, P. Garner, D. Kelly,
C. Bird, S. Palmer, I. Gehring, A. Berger, C. M. Dooley, Z. Ersan-Ürün, C. Eser, H.
Geiger, M. Geisler, L. Karotki, A. Kirn, J. Konantz, M. Konantz, M. Oberländer, S.
Rudolph-Geiger, M. Teucke, C. Lanz, G. Raddatz, K. Osoegawa, B. Zhu, A. Rapp, S.
Widaa, C. Langford, F. Yang, S. C. Schuster, N. P. Carter, J. Harrow, Z. Ning, J. Herrero,
S. M. J. Searle, A. Enright, R. Geisler, R. H. A. Plasterk, C. Lee, M. Westerfield, P. J. de
Jong, L. I. Zon, J. H. Postlethwait, C. Nüsslein-Volhard, T. J. P. Hubbard, H. Roest
Crollius, J. Rogers, and D. L. Stemple. 2013. The zebrafish reference genome sequence
and its relationship to the human genome. Nature 496:498–503.
985
986   Huang, X., and A. Madan. 1999. CAP3: A DNA Sequence assembly program. Genome
987   Res. 9:868–877.
988
989   Hunt, W. G., and Selander, R. K. 1973. Biochemical genetics of hybridisation in
990   European house mice. Heredity, 31, 1133
991
Jaillon, O., J.-M. Aury, F. Brunet, J.-L. Petit, N. Stange-Thomann, E. Mauceli, L.
Bouneau, C. Fischer, C. Ozouf-Costaz, A. Bernot, S. Nicaud, D. Jaffe, S. Fisher, G.
Lutfalla, C. Dossat, B. Segurens, C. Dasilva, M. Salanoubat, M. Levy, N. Boudet, S.
Castellano, V. Anthouard, C. Jubin, V. Castelli, M. Katinka, B. Vacherie, C. Biémont, Z.

Skalli, L. Cattolico, J. Poulain, V. de Berardinis, C. Cruaud, S. Duprat, P. Brottier, J.-P.
Coutanceau, J. Gouzy, G. Parra, G. Lardier, C. Chapple, K. J. McKernan, P. McEwan, S.
Bosak, M. Kellis, J.-N. Volff, R. Guigó, M. C. Zody, J. Mesirov, K. Lindblad-Toh, B.
Birren, C. Nusbaum, D. Kahn, M. Robinson-Rechavi, V. Laudet, V. Schachter, F.
Quétier, W. Saurin, C. Scarpelli, P. Wincker, E. S. Lander, J. Weissenbach, and H. Roest
Crollius. 2004. Genome duplication in the teleost fish *Tetraodon nigroviridis* reveals the
early vertebrate proto-karyotype. Nature 431:946–957.
1003
1004 Jakobsson, M., and N. A. Rosenberg. 2007. CLUMPP: a cluster matching and
1005 permutation program for dealing with label switching and multimodality in analysis of
1006 population structure. Bioinforma. Oxf. Engl. 23:1801–1806.
1007
1008 Jiggins, C. D., and J. Mallet. 2000. Bimodal hybrid zones and speciation. Trends Ecol.
1009 Evol. 15:250–255.
1010
1011 Johannesson, K., M. Panova, P. Kemppainen, C. André, E. Rolán-Alvarez, and R. K.
1012 Butlin. 2010. Repeated evolution of reproductive isolation in a marine snail: unveiling
1013 mechanisms of speciation. Philos. Trans. R. Soc. B Biol. Sci. 365:1735–1747.
1014
1015 Jombart, T. 2008. adegenet: a R package for the multivariate analysis of genetic markers.
1016 Bioinformatics 24:1403–1405.
1017
1018 Jombart, T., R. M. Eggo, P. J. Dodd, and F. Balloux. 2011. Reconstructing disease
1019 outbreaks from genetic data: a graph approach. Heredity 106:383–390.
1020
Jones, F. C., M. G. Grabherr, Y. F. Chan, P. Russell, E. Mauceli, J. Johnson, R.
Swofford, M. Pirun, M. C. Zody, S. White, E. Birney, S. Searle, J. Schmutz, J.
Grimwood, M. C. Dickson, R. M. Myers, C. T. Miller, B. R. Summers, A. K. Knecht, S.
D. Brady, H. Zhang, A. A. Pollen, T. Howes, C. Amemiya, Broad Institute Genome
Sequencing Platform & Whole Genome Assembly Team, E. S. Lander, F. Di Palma, K.
Lindblad-Toh, and D. M. Kingsley. 2012. The genomic basis of adaptive evolution in
threespine sticklebacks. Nature 484:55–61.
1028
1029 Kai, W., K. Kikuchi, S. Tohari, A. K. Chew, A. Tay, A. Fujiwara, S. Hosoya, H. Suetake,
1030 K. Naruse, S. Brenner, Y. Suzuki, and B. Venkatesh. 2011. Integration of the genetic map
1031 and genome assembly of *fugu* facilitates insights into distinct features of genome
1032 evolution in teleosts and mammals. Genome Biol. Evol. 3:424–442.
1033
Kasahara, M., K. Naruse, S. Sasaki, Y. Nakatani, W. Qu, B. Ahsan, T. Yamada, Y.
Nagayasu, K. Doi, Y. Kasai, T. Jindo, D. Kobayashi, A. Shimada, A. Toyoda, Y. Kuroki,
A. Fujiyama, T. Sasaki, A. Shimizu, S. Asakawa, N. Shimizu, S. Hashimoto, J. Yang, Y.
Lee, K. Matsushima, S. Sugano, M. Sakaizumi, T. Narita, K. Ohishi, S. Haga, F. Ohta, H.
Nomoto, K. Nogata, T. Morishita, T. Endo, T. Shin-I, H. Takeda, S. Morishita, and Y.
Kohara. 2007. The *medaka* draft genome and insights into vertebrate genome evolution.
Nature 447:714–719.
1041

Keenan, K., P. McGinnity, T. F. Cross, W. W. Crozier, and P. A. Prodöhl. 2013. diveRsity: An R package for the estimation and exploration of population genetics parameters and their associated errors. Methods Ecol. Evol. 4:782–788.

Knowlton, N. 1993. Sibling species in the sea. Annu. Rev. Ecol. Syst. 24:189–216.

Kulmuni, J., and A. M. Westram. 2017. Intrinsic incompatibilities evolving as a by-product of divergent ecological selection: Considering them in empirical studies on divergence with gene flow. Mol. Ecol. 26:3093-3103.

Larson, E. L., J. A. Andrés, S. M. Bogdanowicz, and R. G. Harrison. 2013. Differential introgression in a mosaic hybrid zone reveals candidate barrier genes. Evol. Int. J. Org. Evol. 67:3653–3661.

Larson, E. L., T. A. White, C. L. Ross, and R. G. Harrison. 2014. Gene flow and the maintenance of species boundaries. Mol. Ecol. 23:1668–1678.

Le Moan, A., P.-A. Gagnaire, and F. Bonhomme. 2016. Parallel genetic divergence among coastal-marine ecotype pairs of European anchovy explained by differential introgression after secondary contact. Mol. Ecol. 25:3187-3202.

Li, H. and R. Durbin. 2009. Fast and accurate short read alignment with Burrows-Wheeler Transform. Bioinformatics. 25:1754-60.

Lin, Q., S. Fan, Y. Zhang, M. Xu, H. Zhang, Y. Yang, A. P. Lee, J. M. Woltering, V. Ravi, H. M. Gunter, W. Luo, Z. Gao, Z. W. Lim, G. Qin, R. F. Schneider, X. Wang, P. Xiong, G. Li, K. Wang, J. Min, C. Zhang, Y. Qiu, J. Bai, W. He, C. Bian, X. Zhang, D. Shan, H. Qu, Y. Sun, Q. Gao, L. Huang, Q. Shi, A. Meyer, and B. Venkatesh. 2016. The seahorse genome and the evolution of its specialized morphology. Nature 540:395–399.

López, A., M. Vera, M. Planas, and C. Bouza. 2015. Conservation genetics of threatened *Hippocampus guttulatus* in vulnerable habitats in NW Spain: temporal and spatial stability of wild populations with flexible polygamous mating system in captivity. PLoS ONE 10:e0117538.

Lotterhos, K. E., and M. C. Whitlock. 2014. Evaluation of demographic history and neutral parameterization on the performance of FST outlier tests. Mol. Ecol. 23:2178–2192.

Lotterhos, K. E., and M. C. Whitlock. 2015. The relative power of genome scans to detect local adaptation depends on sampling design and statistical method. Mol. Ecol. 24:1031–1046.

Lourie, S. A., and A. C. J. Vincent. 2004. Using biogeography to help set priorities in marine conservation. Conserv. Biol. 18:1004–1020.

1088 Luu, K., E. Bazin, and M. G. B. Blum. 2016. PCAdapt: an R package to perform genome
1089 scans for selection based on principal component analysis. bioRxiv 056135.
1090
1091 Macholán, M., S. J. E. Baird, P. Dufková, P. Munclinger, B. V. Bímová, and J. Piálek.
1092 2011. Assessing multilocus introgression patterns: a case study on the mouse X
1093 chromosome in central Europe. Evol. Int. J. Org. Evol. 65:1428–1446.
1094
1095 Nosil, P., and J. L. Feder. 2012. Genomic divergence during speciation: causes and
1096 consequences. Phil Trans R Soc B 367:332–342.
1097
1098 Pante, E., N. Puillandre, A. Viricel, S. Arnaud-Haond, D. Aurelle, M. Castelin, A.
1099 Chenuil, C. Destombe, D. Forcioli, M. Valero, F. Viard, and S. Samadi. 2015. Species are
1100 hypotheses: avoid connectivity assessments based on pillars of sand. Mol. Ecol. 24:525–
1101 544.
1102
1103 Pérez‑Figueroa, A., M. J. García‑Pereira, M. Saura, E. Rolán‑Alvarez, and A.
1104 Caballero. 2010. Comparing three different methods to detect selective loci using
1105 dominant markers. J. Evol. Biol. 23:2267–2276.
1106
1107 Pickrell, J. K., and J. K. Pritchard. 2012. Inference of population splits and mixtures from
1108 genome-wide allele frequency data. PLoS Genet 8:e1002967.
1109
1110 Pritchard, J. K., M. Stephens, and P. Donnelly. 2000. Inference of population structure
1111 using multilocus genotype data. Genetics 155:945–959.
1112
1113 R Development Core Team (2011) R: A language and environment for statistical
1114 computing. R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-
1115 0, http://www.R-project.org/.
1116
1117 Rand, D. M., and R. G. Harrison. 1989. Ecological genetics of a mosaic hybrid zone:
1118 mitochondrial, nuclear, and reproductive differentiation of crickets by soil type.
1119 Evolution 43:432–449.
1120
1121 Ravinet, M., R. Faria, R. K. Butlin, J. Galindo, N. Bierne, M. Rafajlović, M. a. F. Noor,
1122 B. Mehlig, and A. M. Westram. 2017. Interpreting the genomic landscape of speciation: a
1123 road map for finding barriers to gene flow. J. Evol. Biol. 30:1450–1477.
1124
1125 Raymond, M., and F. Rousset. 1995. GENEPOP (Version 1.2): Population genetics
1126 software for exact tests and ecumenicism. J. Hered. 86:248–249.
1127
1128 Ribardière, A., C. Daguin‑Thiébaut, C. Houbin, J. Coudret, C. Broudin, O. Timsit, and T.
1129 Broquet. 2017. Geographically distinct patterns of reproductive isolation and
1130 hybridization in two sympatric species of the *Jaera albifrons* complex (marine isopods).
1131 Ecol. Evol. 7:5352–5365.
1132

Romiguier, J., P. Gayral, M. Ballenghien, A. Bernard, V. Cahais, A. Chenuil, Y. Chiari, R. Dernat, L. Duret, N. Faivre, E. Loire, J. M. Lourenco, B. Nabholz, C. Roux, G. Tsagkogeorga, A. a.-T. Weber, L. A. Weinert, K. Belkhir, N. Bierne, S. Glémin, and N. Galtier. 2014. Comparative population genomics in animals uncovers the determinants of genetic diversity. Nature 515:261–263.

Rougemont, Q., P.-A. Gagnaire, C. Perrier, C. Genthon, A.-L. Besnard, S. Launey, and G. Evanno. 2016. Inferring the demographic history underlying parallel genomic divergence among pairs of parasitic and nonparasitic lamprey ecotypes. Mol. Ecol. 26:142–162.

Rougeux, C., L. Bernatchez, and P.-A. Gagnaire. 2016. Modeling the multiple facets of speciation-with-gene-flow towards inferring the divergence history of lake whitefish species pairs (*Coregonus clupeaformis*). bioRxiv 068932.

Rousset, F. 2008. genepop'007: a complete re-implementation of the genepop software for Windows and Linux. Mol. Ecol. Resour. 8:103–106.

Roux, C., C. Fraïsse, J. Romiguier, Y. Anciaux, N. Galtier, and N. Bierne. 2016. Shedding light on the grey zone of speciation along a continuum of genomic divergence. PLOS Biol. 14:e2000234.

Sakka Hlaili, A., B. Grami, N. Niquil, M. Gosselin, D. Hamel, M. Troussellier, and H. Hadj Mabrouk. 2008. The planktonic food web of the Bizerte lagoon (south-western Mediterranean) during summer: I. Spatial distribution under different anthropogenic pressures. Estuar. Coast. Shelf Sci. 78:61–77.

Schluter, D., and G. L. Conte. 2009. Genetics and ecological speciation. Proc. Natl. Acad. Sci. U. S. A. 106 Suppl 1:9955–9962.

Sheets, E. A., P. A. Warner, and S. R. Palumbi. 2018. Accurate population genetic measurements require cryptic species identification in corals. Coral Reefs 37:549–563.

Simpson, J. T., K. Wong, S. D. Jackman, J. E. Schein, S. J. M. Jones, and İ. Birol. 2009. ABySS: A parallel assembler for short read sequence data. Genome Res. 19:1117–1123.

Small, C. M., S. Bassham, J. Catchen, A. Amores, A. M. Fuiten, R. S. Brown, A. G. Jones, and W. A. Cresko. 2016. The genome of the Gulf pipefish enables understanding of evolutionary innovations. Genome Biol. 17:258.

Soria-Carrasco, V., Z. Gompert, A. A. Comeault, T. E. Farkas, T. L. Parchman, J. S. Johnston, C. A. Buerkle, J. L. Feder, J. Bast, T. Schwander, S. P. Egan, B. J. Crespi, and P. Nosil. 2014. Stick insect genomes reveal natural selection's role in parallel speciation. Science 344:738–742.

1178 Storey, J. D. 2002. A direct approach to false discovery rates. J. R. Stat. Soc. Ser. B Stat.
1179 Methodol. 64:479–498.
1180
1181 Szymura, J. M., and N. H. Barton. 1986. Genetic analysis of a hybrid zone between the
1182 fire-bellied toads, *Bombina bombina* and *B. variegata*, near Cracow in Southern Poland.
1183 Evolution 40:1141–1159.
1184
1185 Teeter, K. C., B. A. Payseur, L. W. Harris, M. A. Bakewell, L. M. Thibodeau, J. E.
1186 O'Brien, J. G. Krenz, M. A. Sans-Fuentes, M. W. Nachman, and P. K. Tucker. 2008.
1187 Genome-wide patterns of gene flow across a house mouse hybrid zone. Genome Res.
1188 18:67–76.
1189
1190 Tine, M., H. Kuhl, P.-A. Gagnaire, B. Louro, E. Desmarais, R. S. T. Martins, J. Hecht, F.
1191 Knaust, K. Belkhir, S. Klages, R. Dieterich, K. Stueber, F. Piferrer, B. Guinand, N.
1192 Bierne, F. A. M. Volckaert, L. Bargelloni, D. M. Power, F. Bonhomme, A. V. M.
1193 Canario, and R. Reinhardt. 2014. European sea bass genome and its variation provide
1194 insights into adaptation to euryhalinity and speciation. Nat. Commun. 5.
1195
1196 Via, S. 2012. Divergence hitchhiking and the spread of genomic isolation during
1197 ecological speciation-with-gene-flow. Philos. Trans. R. Soc. Lond. B Biol. Sci. 367:451–
1198 460.
1199
1200 Vines, T., A. Dalziel, A. Albert, T. Veen, P. Schulte, and D. Schluter. 2016. Cline
1201 coupling and uncoupling in a stickleback hybrid zone. Evolution 70:1023-38.
1202
1203 Wakeley, J. 2008. Coalescent theory: an introduction. Roberts & Company Publishers,
1204 Greenwood Village, Colorado.
1205
1206 Weir, B. S., and C. C. Cockerham. 1984. Estimating F-Statistics for the analysis of
1207 population structure. Evolution 38:1358–1370.
1208
1209 Welch, J. J., and C. D. Jiggins. 2014. Standing and flowing: the complex origins of
1210 adaptive variation. Mol. Ecol. 23:3935–3937.
1211
1212 Woodall, L. C., R. Jones, B. Zimmerman, S. Guillaume, T. Stubbington, P. Shaw, and H.
1213 J. Koldewey. 2012. Partial fin-clipping as an effective tool for tissue sampling seahorses,
1214 *Hippocampus* spp. J. Mar. Biol. Assoc. U. K. 92:1427–1432.
1215
1216 Woodall, L. C., H. J. Koldewey, J. T. Boehm, and P. W. Shaw. 2015. Past and present
1217 drivers of population structure in a small coastal fish, the European long snouted seahorse
1218 *Hippocampus guttulatus*. Conserv. Genet. 1–15.
1219
1220 Yeaman, S. 2013. Genomic rearrangements and the evolution of clusters of locally
1221 adaptive loci. Proc. Natl. Acad. Sci. 110:E1743–E1751.
1222

1223 **Data archival location**