HBV vaccination and PMTCT as elimination tools in the presence of HIV: insights from a clinical cohort and dynamic model

Anna L McNaughton[1*], anna.mcnaughton@ndm.ox.ac.uk

José Lourenço[2*], jose.lourenco@zoo.ox.ac.uk

Louise Hattingh[3*], loeise@gmail.com

Emily Adland[4], emily.adland@paediatrics.ox.ac.uk

Samantha Daniels[3], danies.samantha@gmail.com

Anriette Van Zyl[3], ajmvanzyl@yahoo.com

Connie S Akiror[5], connieakiror@gmail.com

Susan Wareing[6], susan.wareing@ouh.nhs.uk

Katie Jeffery[6], katie.jeffery@ouh.nhs.uk

M Azim Ansari[1], ansari.azim@gmail.com

Paul Klenerman[1,6], paul.klenerman@ndm.ox.ac.uk

Philip J R Goulder[4], philip.goulder@paediatrics.ox.ac.uk

Sunetra Gupta[2], sunetra.gupta@zoo.ox.ac.uk

Pieter Jooste[3], jpjooste@mweb.co.za

Philippa C Matthews[1,6 §], philippa.matthews@ndm.ox.ac.uk

* These three authors contributed equally to the work presented here

[1] Nuffield Department of Medicine, Peter Medawar Building for Pathogen Research, South Parks Road, Oxford OX1 3SY, UK

[2] Department of Zoology, Peter Medawar Building for Pathogen Research, South Parks Road, Oxford OX1 3SY, UK

[3] Department of Paediatrics, Kimberley Hospital, Kimberley, 8300, South Africa

[4] Department of Paediatrics, Peter Medawar Building for Pathogen Research, South Parks Road, Oxford OX1 3SY, UK

[5] Global Healthcare Public Foundation, Makindu Lane, Kololo, Kampala, Uganda

[6] Department of Infectious Diseases and Microbiology, Oxford University Hospitals NHS Foundation Trust, John Radcliffe Hospital, Headley Way, Oxford OX3 9DU, UK

RUNNING HEAD: HBV vaccination and PMTCT as elimination tools

[§] Corresponding author Philippa Matthews

Email: philippa.matthews@ndm.ox.ac.uk; Address: Medawar Building for Pathogen

Research, South Parks Road, Oxford OX1 3SY, UK; Telephone: 0044 1865 271973

ABBREVIATIONS

- HBV – Hepatitis B virus

- SDGs – Sustainable Development Goals

- HIV – Human Immunodeficiency Virus (type 1)

- WHO – World Health Organisation

- EPI – Expanded Programme on Immunisation

- PMTCT – Prevention of mother to child transmission

- COSAC – Coinfection in South African children

- HBsAg – Hepatitis B surface antigen

- Anti-HBs – Antibody to hepatitis B surface antigen (vaccine-mediated antibody)

- KReC – Kimberley Respiratory Cohort

- RTHB – Road to Health Book

- Anti-HBc – Antibody to hepatitis B core antigen (antibody mediated by exposure to
infection)

- ODE – Ordinary differential equation

- HBeAg – Hepatitis B envelope antigen

- GHSSVH - Global Health Sector Strategy on Viral Hepatitis

- ART – Anti-retroviral therapy

- 3TC - Lamivudine

- OUH – Oxford University Hospitals

ABSTRACT

Background: Sustainable Development Goals set a challenge for the elimination of hepatitis

B virus (HBV) infection as a public health concern by the year 2030. Deployment of a robust

prophylactic vaccine and enhanced interventions for prevention of mother to child

transmission (PMTCT) are cornerstones of elimination strategy. However, in light of the

estimated global burden of 290 million cases, enhanced efforts are required to underpin

optimisation of public health strategy. Robust analysis of population epidemiology is

particularly crucial for populations in Africa made vulnerable by HIV co-infection, poverty,

stigma and poor access to prevention, diagnosis and treatment.

Methods: We here set out to evaluate the current and future role of HBV vaccination and

PMTCT as tools for elimination. We first investigated the current impact of paediatric

vaccination in a cohort of children with and without HIV infection in Kimberley, South Africa.

Second, we used these data to inform a new model to simulate the ongoing impact of

preventive interventions. By applying these two approaches in parallel, we are able to

determine both the current impact of interventions, and the future projected outcome of

ongoing preventive strategies over time.

Results: Existing efforts have been successful in reducing paediatric prevalence of HBV

infection in this setting to <1%, demonstrating the success of the existing vaccine campaign.

Our model predicts that, if consistently deployed, combination efforts of vaccination and

PMTCT can significantly reduce population prevalence (HBsAg) by 2030, such that a major

public health impact is possible even without achieving elimination. However, the prevalence

of HBV e-antigen (HBeAg)-positive carriers will decline more slowly, representing a

persistent population reservoir. We show that HIV co-infection significantly reduces titres of

vaccine-mediated antibody, but has a relatively minor role in influencing the projected time to

elimination. Our model can also be applied to other settings in order to predict time to

elimination based on specific interventions.

Conclusions: Through extensive deployment of preventive strategies for HBV, significant

positive public health impact is possible, although time to HBV elimination as a public health

concern is likely to be substantially longer than that proposed by current goals.

KEY WORDS: hepatitis b virus; epidemiology; Africa; antibodies; immunisation; PMTCT;

sustainable development goals; HIV; vaccination; elimination

BACKGROUND

The vaccine against hepatitis B virus (HBV) infection is one of the cornerstone strategies underpinning progress towards Sustainable Development Goals (SDGs) targets for HBV elimination as a public health threat by the year 2030 (1,2). Populations in southern Africa are particularly vulnerable to HBV-related morbidity and mortality due to the high prevalence of infection (≥8% in many regions) (3–5), co-endemic HIV infection (6), poor access to screening and diagnostics, limited access to antiviral therapy, stigma, and chronic neglect of education, research and resources (7,8). In this region, a substantial burden of HBV transmission occurs early in life, either vertically from mother to child, or through horizontal acquisition in young children (9). The HBV vaccine has been progressively rolled out as part of the World Health Organisation (WHO) Expanded Programme on Immunisation (EPI) over the past two decades (9), but the first vaccine dose is often postponed until age six weeks, when it is given together with other routine immunizations (10).

Vaccine deployment can be difficult to measure, as many children in Africa are born outside healthcare settings, there are no robust data regarding coverage of the three dose regimen (9), and different immunological correlates of protection have been applied (11,12). To accelerate progress towards elimination goals, suggested modifications to vaccine schedules have included shifting the first dose to be given at birth (13), additional doses in the context of HIV infection (14,15), booster doses in individuals whose antibody titre fails to meet a target threshold (11), and catch-up vaccination campaigns for adolescents and adults. However, there is a lack of robust data to inform which of these measures, individually or in combination, is most effective. Given the resource limitations of many settings in which HBV represents a public health challenge, there is an urgent need to underpin interventions with an evidence base derived both from careful observation of the existing impact of vaccination and from projections regarding ongoing impact.

On these grounds, we have set out to collect a detailed dataset to provide a snapshot of a population in South Africa in which HBV and HIV infections are co-endemic, first seeking evidence of the impact of the current immunization schedule in children, and then assessing the extent to which interventions could be predicted to achieve elimination targets. Finally, we built on this framework by adding data assimilated from the wider published literature to model the effects of different HBV vaccine deployment strategies, either alone or in combination with enhanced measures for prevention of mother to child transmission (PMTCT) through screening, antiviral therapy, accelerated immunisation and HBV immunoglobulin (16).

To date, few attempts have been made to model the impact of HBV vaccination, with one

study modelling the global prevalence of current intervention efforts (17), and another that

scrutinizes the combined impact of broad HBV elimination strategies (18). In this instance,

we report a novel approach founded on primary clinical data, quantifying the individual and

combined impact of childhood vaccination and PMTCT, and addressing the specific impact

of co-endemic HIV infection. Combining output from a clinical dataset together with a

dynamic model provides a synergistic approach to characterising the problem and projecting

the effects of vaccination. Taking the evidence together, we conclude that while vaccination

is a fundamental part of global elimination strategy and is highly effective in preventing

infection in individual children, there remains an urgent need for rigorous, enhanced

deployment of parallel strategies including education, diagnostics, antiviral therapy, and the

ongoing quest for a cure.

METHODS

Ethics Approval

Ethics approval was obtained from the Ethics Committee of the Faculty of Health Science,

University of the Free State, Bloemfontein, South Africa (HIV Study Ref: ETOVS Nr 08/09

and COSAC Study Ref: ECUFS NR 80/2014). Written consent for enrollment into the study

was obtained from the child's parent/guardian.

Study cohorts

Recruitment was undertaken in Kimberley, South Africa. In this setting, a standard three-

dose HBV immunisation schedule is deployed in infants, with the first dose at six weeks. A

previous study of HBV serology in adults in the same setting found HBsAg prevalence of

9.5% (55/579) (4). Children were recruited as part of the Co-infection in South-African

Children ('COSAC') study as previously described (19,20). The lower age limit of recruitment

was 6 months in order to limit the detection of maternal anti-HBs.

Children were recruited as follows:

1. HIV-negative children age 6-60 months (n=174), recruited through the Kimberley

Respiratory Cohort (KReC) as previously described (19). These children were admitted to

hospital between July 2014 and August 2016 with a clinical diagnosis of respiratory tract

infection. KReC children were confirmed HIV-negative in 163 cases (93.7%). A further 11

children did not have an HIV test result recorded, but were assumed to be HIV-negative

based on the clinical data recorded at the time of admission to hospital.

2. HIV-positive children were recruited primarily from HIV out-patient clinics between

September 2009 and July 2016 as previously described (19,20). We recorded date of

commencement of anti-retroviral therapy (ART), CD4+ T cell count and percentage, and HIV

RNA viral load using the time point closest to the sample that was analysed for HBV

serology. For the purpose of analysis, we divided these into two groups according by age:

i.   Age 6-60 months; n=136. This group was selected to match the age range of

177   the HIV-negative group, and also included five children who were initially

178   screened for the KReC cohort but tested HIV-positive.

ii.   Age >60 months (range 64-193 months); n=92.

At the time of undertaking this study, children were immunised with three doses of a

monovalent HBV vaccine (Biovac Paed). Where possible, we recorded the number of HBV

vaccine doses received based on the Road to Health Book (RTHB). The characteristics of

the cohorts are summarised in table 2 and all metadata can be found in Suppl. data 1.

Laboratory assessment of HBV status

Testing for Hepatitis B serum markers and DNA was performed as previously described, and

in keeping with recent implementation of HBV screening in Kimberley (20). Briefly, HBsAg

testing was carried out in Kimberley Hospital, South Africa using the Magnetic parcel

chemiluminometric immunoassay (MPCI; Advia Centaur platform). Confirmatory HBsAg

testing was carried out by the clinical microbiology laboratory at Oxford University Hospitals

(OUH) NHS Foundation Trust, Oxford, UK (Architect i2000). For all samples, anti-HBs and

anti-HBc testing were carried out by the OUH laboratory (Architect i2000). Limit of detection

of the anti-HBs assay was 10 mIU/ml.

Threshold for vaccine mediated immunity

Studies variably quote anti-HBs titres of ≥10 mIU/ml or ≥100 mIU/ml as a correlate of

protection; UK recommendations for testing HBV immunity advocate the more stringent

criterion of an anti-HBs titre of ≥100 mIU/ml (11), while early vaccine studies suggest a titre

of ≥10 mIU/ml as a clinically relevant threshold for protection (12,21). We have presented

our results pertaining to both thresholds.

Statistical analysis

Data from the cohort was analysed using GraphPad Prism v.7.0. We determined significant

differences between sub-sets within the cohort using Mann-Whitney U tests for non-

parametric data, Fisher's exact test for categorical variables and Spearman's correlation

coefficient for correlation between data points.

Mathematical model of HBV transmission and prevention

We developed a dynamic model based on ordinary differential equations (ODE), for which
parameterization of HBV transmission and prevention was based both on our Kimberley
paediatric cohort and current literature estimates. This model was fitted to the HBV
transmission background (Kimberley cohort) using Bayesian Markov Chain Monte-Carlo
(bMCMC), and stochastic simulations were used to project impact interventions under
various strategic scenarios. The full modelling and fitting approaches are detailed in
Supplementary File 2.
217
Measuring impact of interventions
SDGs for the year 2030 have been set out in the WHO Global Health Sector Strategy on
Viral Hepatitis (GHSSVH) (2). Given the public health relevance of chronic infections, in
particular of HBeAg-positive infections, we measured impact of interventions based on two
targets:
i.   The WHO target for a 90% reduction in HBsAg incidence, based on the assumption
that this applies to chronic infection. (WHO goals also refer to HBsAg prevalence,
and we present this in Supplementary File 2).
ii.   An additional target for reduction of HBeAg-positive prevalence to 1 in 1000 (0.1%) in
the whole population.
228
RESULTS
Serological evidence of exposure to HBV infection
From our cohort of 402 children in Kimberley, South Africa, three were HBsAg-positive
(0.7%; Table 1), indicating active infection. This HBsAg prevalence is significantly lower than
in adults in a comparable study population (e.g. 11.1% in a previous study (4); p<0.0001).
Exposure to HBV infection was measured using anti-HBc antibody; this was detected in
three children (0.7%), one of whom was also HBsAg-positive. The other two were HBsAg-
negative, indicating previous HBV exposure and clearance.
237
Evidence of vaccination and immunity to HBV in children aged ≤60 months
We collected written evidence of immunisation from the RTHB in 90.8% HIV-negative
(KReC) subjects and 6.3% of HIV-positive subjects (total 41.3% of cohort). None of the
HBsAg-positive children attended with a written vaccination record. Among those with a
RTHB record, 81.3% of HIV-negative and 100% of HIV-positive children were recorded as
having received three HBV vaccine doses. Among all children age ≤60 months, 238/310
(77%) had an anti-HBs titre ≥10 mIU/ml suggesting some degree of vaccine-mediated
immunity. The median anti-HBs titre in HIV-negative children was significantly higher than
among the HIV-positive group (196 mIU/ml, vs. 11 mIU/ml, respectively, p<0.0001) (Fig 2A).

There was no detectable anti-HBs antibody in 3.4% of HIV-negative vs. 47.8% of HIV-

positive children (p<0.0001). Irrespective of the antibody titre used as a threshold for

immunity, anti-HBs was higher in HIV-negative compared to HIV-positive children (Fig 2B).

There was no significant difference in anti-HBs titres between male and female participants,

either with or without HIV infection (p=0.49 and 0.31 respectively, data not shown).

Waning of vaccine response with age

HIV-positive children with anti-HBs titres ≥100mIU/ml were significantly younger than those

with lower antibody titres (median age 17 months vs. 31 months, p=0.0008), while no such

difference was observed within the HIV-negative group (Fig 3A). Using the lower threshold of

≥10mIU/ml, we found no significant difference by age in either the HIV-positive or the HIV-

negative groups (p=0.17 and 4.48 respectively, data not shown). To expand our view of the

HIV-positive group, we also added analysis of an older cohort (92 children aged >60

months), and demonstrated that anti-HBs titres were significantly lower in this older group

(p<0.0001), with only 2/92 subjects (2.2%) achieving a detectable anti-HBs titre (Fig 3B).

Anti-HBs titres waned significantly with age up to age 60 months in HIV-positive children (Fig

3C; p=0.004). We observed a similar trend in the HIV-negative cohort, but this did not reach

statistical significance (Fig 3C; p=0.07). The proportion of HIV-positive subjects with a

detectable anti-HBs titre declined steadily with age in the cohort, contrasting to the trend in

HIV-negative subjects, where individuals maintained protective anti-HBs titres despite a

trend towards decreasing mean titres (Fig 3C).

Stratification of vaccine responses by anti-retroviral therapy (ART) among HIV-

positive children

For HIV-positive children aged ≤60 months, ART treatment data were available for 79% of

subjects. Within this group, 71% were receiving ART at the time we tested for anti-HBs, and

had received a median of 20 months of treatment (IQR 6-33 months). Comparing anti-HBs

titres between ART-treated vs. untreated children, we found no significant difference

(p=0.72; 76 ART-treated, median anti-HBs 13.3 mIU/ml and 31 untreated children, median

anti-HBs 14.1 mIU/ml, data not shown). There was also no difference between anti-HBs

titres of children treated for ≤12 months vs. >12 months (p=0.50, data not shown). We did

not examine the effect of ART on anti-HBs titres in children >60 months due to the low

numbers of subjects with a detectable anti-HBs titre (n=2).

Odds of developing an anti-HBs response

We used an odds ratio (OR) analysis to identify factors associated with vaccine-mediated

protection (Fig 3D). HIV-positive status was associated with lack of protection, for antibody

titres of both <10 mIU/ml (OR 26.2, 95% CI 11.2-58.6), and <100 mIU/ml (OR 11.6, 95% CI

6.7-20.4). In contrast, younger age (<24 months) was protective in HIV-positive subjects, (for

anti-HBs <10 mIU/ml OR 0.3, 95% CI 0.2-0.5 and for anti-HBs <100 mIU/ml OR 0.3, 95% CI

0.2-0.4). Gender, ART, CD4+ count, CD4+ ratio and HIV viral load were not found to be

significantly predictive of anti-HBs titres at either threshold.

Fitting of the dynamic model to local HBV epidemiology

The model closely reproduced the fitted variables: HBV prevalence (HBsAg), prevalence of

HBV exposure (anti-HBc) and relative proportion of HBeAg-negative and HBeAg-positive

among chronic carriers (Suppl Fig 1 A12). Parameters for which insufficient support was

found in the literature, the resulting posteriors were well behaved (Suppl Fig 1B), while for

parameters using informative priors the posteriors matched well (Suppl Fig 1CD). Overall,

the obtained bell-shaped posteriors highlighted a good data fit and no identifiability issues

with the bMCMC approach. Based on these results, we next simulated stochastic projections

of intervention impact.

Model projection of the impact of routine neonatal vaccination and PMTCT alone

Based on SDGs (2), Fig 4 shows the projected impact of varying coverage of neonatal

vaccination and PMTCT. Both HBsAg incidence (Fig 4 A1) and HBeAg-positive prevalence

(Fig 4 B1) reduce faster with increasing neonatal immunization coverage, resulting in shorter

times to reach SDGs (Fig 4 A2, B2). Importantly, even immunization of 100% of neonates is

predicted to take ~99 years (95% CI 61 - 186) for the HBsAg incidence target (Fig 4 A2), and

~175 years (95% CI 103 - 278) for the HBeAg prevalence target (Fig 4 B2).

With PMTCT interventions, both HBsAg incidence (Fig 4 C1) and HBeAg prevalence (Fig 4

D1) reduced faster in time for increasing efforts, resulting in shorter times to reach the

elimination targets (Fig 4 C2, D2). However, the impact of PMTCT was smaller than

neonatal vaccination, resulting in longer times to reach SDGs. In fact, for the majority of

PMTCT effort levels simulated, SDGs could not be reached within 500 years (beige areas in

Fig 4 C2, D2). For HBeAg-positive prevalence, only when PMTCT effort was 1 (i.e. no

vertical transmission), was the SDG attainable within 500 years.

For complementary results using neonatal vaccination and PMTCT with impact on total

prevalence (acute and chronic) see Support Figure 4.

Modelling progress towards HBV elimination by the year 2030 based on combinations

of neonatal vaccination and PMTCT

We projected impact of combined interventions by the year 2030 (Fig 5 A1, B1), and

predicted the year at which SDGs would be reached (Fig 5 A2, B2). Strikingly, HBsAg

incidence could already have been reduced by >90% (Fig 5 A1) if both neonatal vaccination

and PMTCT had been deployed at 100% coverage since they became available in 1995

(mean predicted year of elimination 2017; Fig 5 A2). In reality, complete coverage is not

possible, and we therefore projected outcomes based on <100% coverage. For example,

combining neonatal vaccination and PMTCT with 90% coverage of each since 1995 would

achieve the HBsAg incidence target by 2028; if this is reduced to 80% coverage then goals

would be attained by 2044. To achieve the target reduction in HBeAg prevalence, modelled

on 90% coverage and 80% coverage of interventions, the projected years are 2072 and

2096 respectively (Fig 5 B1, B2).

Overall, the highest probability of achieving elimination targets is through a combination of

100% neonatal vaccination coverage and PMTCT (Fig 6 A1, red line). Based on 90%

coverage of neonatal vaccine and PMTCT (Fig 6 A1, A2, green line) as proposed in WHO'S

GHSSVH (2), there was only 50% probability of reaching the HBsAg incidence target by

2030, and approaching 100% probability only by 2050. For the target based on HBeAg

prevalence, the probabilities of achieving the goal were pushed forward by approximately

four decades.

For complementary results on impact and time to reach SDGs when considering

combinations of PMTCT and routine vaccination at the age of 6 see Support Figure 5, and

for combinations of PMTCT and neonate routine vaccination plus a complete catch-up

campaign see Support Figure 6.

Projecting the probability of achieving elimination targets based on combinations of

neonatal vaccination, PMTCT and enhanced vaccination

We simulated the impact of combining neonatal vaccination and PMTCT with additional

vaccine deployment (Fig 6 A1, A2), through the routine vaccination of older children (≥6

years of age), and one-off catch-up vaccination of children (<6 years) and others (>6 years).

Adding catch-up vaccination campaigns makes no impact on the probability of reaching

SDGs (Fig 6 A1, A2, blue and cyan lines). Routine vaccination at 6 years of age, even when

delivered at 100% coverage, is markedly less effective than any other projected intervention

(Fig 6 A1, A2, magenta line).

Projecting the impact of HIV on the probability of achieving elimination targets

A baseline scenario was defined by the epidemiological setting fitted by our model in the

context of Kimberley, using local HIV prevalence for each of the modelled age groups (Fig 6

B1, B2, solid line). In a sensitivity exercise, alternative scenarios were considered in which

baseline HIV prevalence was altered to zero or higher prevalence. When compared with no

HIV (Fig 6 B1, B2, dotted line), the presence of HIV at the prevalence seen in Kimberley (Fig

6 B1, B2, solid line) adds an estimated four years to achieve a 50% chance of reaching the

goals (Fig 6 B1). Higher baseline HIV prevalence (x2, x3 and x4 baseline data for Kimberley)

was used to investigate the potential impact of coinfection in high-risk populations. Thus,

increasing HIV prevalence has a negative impact on the success of interventions for HBV,

but the effects are relatively modest. In particular, doubling HIV prevalence would shift the

50% probability endpoint into the future by ~4 years for the HBsAg incidence target, and ~7

years for the HBeAg prevalence target.

DISCUSSION

This is a unique study in which we capitalize on detailed clinical cohort data collected in

South Africa in order to form a robust view of the nature of vaccine-mediated immunity, and

develop a mathematical model of HBV transmission and prevention. Overall, we

demonstrate that the optimum population intervention is high coverage neonatal vaccination,

and that this can be strengthened by robust deployment of PMTCT. However, we project

long time-scales to achieve elimination targets, congruent with the large established

reservoir of chronic HBV infection, lack of curative therapy, infection that can persist for the

entire life-span of the host, and interventions that target only a small proportion of the

population. Although a high coverage of neonatal vaccination combined with robust PMCTC

shows potential promise, the projected time-frame for elimination is currently substantially

beyond the 2030 milestone. Complete extinction of infection is far beyond reach based on

currently available interventions, and current efforts should be focused on control of HBV as

public health issue rather than complete elimination.

By assimilating the results of a clinical cohort study and a model, we develop a more

complete picture than either individual approach would provide in isolation. Only by viewing

the two conclusions together can we correctly infer that vaccination is of profound

importance in protecting individual children and significantly reducing the burden of infection

in paediatric cohorts, but also that continuing to pursue this strategy alone is not sufficient to

bring about HBV elimination, or even robust control, within the desired time-scale.

Compared to published models of other vaccine-preventable diseases (22), there is a

marked deficit in the existing literature for HBV, with few other modelling efforts represented

in the literature (23,24). Reassuringly, our findings are consistent with those of another
recent simulation of HBV prevention (18); we concur in concluding that current vaccine-
based interventions will result in a modest reduction in HBV prevalence by the year 2030.
However, there are also some important differences that distinguish our work from previous
efforts:
i.   Our evaluation provides the advantages of both clinical data and a mathematical
400   model, with close links between our cohort and simulations, and strengths in
401   interpretation of data derived through different approaches. In so doing, we have also
402   been able to specifically address the impact of co-endemic HIV.
403   ii.   We focus on a particular population for which we derive unknown epidemiological
404   parameters and apply a robust data-driven approach to others. Our Bayesian
405   framework therefore stands alone (as a tool) that can be applied to any population for
406   which empirical support of key HBV epidemiological parameters is missing. By
407   supplying the model's code, we can facilitate the use of the tool by others.
408   iii.   As outputs, we have used targets for reductions in both HBsAg incidence and
409   HBeAg-positive prevalence, and have projected the impact of interventions based
410   specifically on the WHO proposal for 90% vaccination of neonates and 90% PMTCT
411   coverage by 2030. Previous studies (17,18) have focused instead on *ad hoc* control
412   thresholds or impact on the public health problem through reduction of HBV-related
413   deaths. Our results thus contribute to an ongoing discussion regarding which goals
414   should be set, and their underlying public health implications. The model suggests
415   that reaching either of the elimination targets will require different intervention
416   coverage and different time scales. In particular, the target for reducing HBsAg
417   incidence is easier to achieve than reducing HBeAg-prevalence.
418   iv.   This is an important parsimonious, data-driven tool, offering the potential to scrutinise
419   different strategies independently from one another.
420
Impact of HIV on population interventions for HBV
Our results support previous evidence that HBV vaccine-mediated immunity wanes over
time independently of HIV serostatus, but faster for HIV-positive individuals (25). Impaired
vaccine responses have previously been reported in HIV-positive individuals (15,26–29), but
it is also possible that vaccine coverage is lower in HIV-infected children (30). However,
waning of anti-HBs titres does not necessarily correlate with loss of clinical protection;
anamnestic responses are thought to occur in a proportion of those vaccinated (31),
although this memory may be attenuated by HIV (32,33).
429

ART has previously been associated with improved HBV vaccine responses (34,35),

although we did not replicate this finding in our cohort. This can potentially be explained by a

previous study in the same setting, demonstrating that immune reconstitution takes a

median of five years after ART initiation (36). Our current study is underpowered to detect

any ART effect, given both the relatively short durations of therapy, and the small number of

untreated children. Interestingly, despite the lack of direct association with ART, children with

lower HIV viral loads had significantly higher anti-HBs titres, in keeping with previous studies

(14,34).

Our cohort highlights day-to-day challenges of drug provision and monitoring in this setting:

we did not have access to detailed prospective ART treatment data, guidelines have

changed numerous times since 2002, and lamivudine (3TC) was intermittently used as a

substitute for nevirapine (NVP) due to supply issues. During the period covered by our study,

ART was only introduced in children achieving certain immunological criteria, while new

guidelines recommend that all HIV-infected children are started on ART (37). The immune

reconstitution of this population over time is likely to reduce differences between HIV-positive

and HIV-negative groups. ART treatment is relevant to outcomes in individuals with HIV/HBV

coinfection, as first line ART regimens include either 3TC or tenofovir (TDF), both of which

have activity against HBV. Alternative approaches for HBV prevention in HIV-positive

subjects, such as supplementing the current schedule with booster vaccinations and

increased vaccine doses have been trialled with variable results (14)(15).

Our projections propose that HIV does have a negative effect on HBV interventions,

although HIV prevalence only marginally increases time to reach elimination targets, which

may not be significant in light of the long overall time-frames projected, even in the absence

of HIV. The high HIV prevalences modelled can occur in specific high-risk groups including

sex workers and men who have sex with men (38) and it is likely that increased intervention

will be required in these groups to minimise HBV transmission.

Changes required to meet international goals

The model suggests long time-lines, enumerated in centuries rather than decades, before

control targets are reached using vaccination or PMTCT alone. Combinations of these

interventions show much shorter time scales. Based on currently available interventions,

major scaling up of both neonatal vaccination and PMTCT efforts will be required.

Importantly, the prevalence of HBeAg-positive carriers, who are at an elevated risk of

chronic liver disease and hepatocellular carcinoma, as well as being at higher risk of

transmitting their infection, will decline at a slower rate. Setting a control target based on

reduction in the number of new HBV cases (i.e. HBsAg incidence) can therefore lead to the
most optimistic projections but distract attention from the importance of reducing HBeAg
prevalence.
470
Our results also underscore that a major public health impact is possible, even without
achieving elimination. Careful adjusting of expectations and aims, according to the scale on
which particular changes occur, may inform the setting of realistic targets (e.g. reduction in
the prevalence of HBeAg could be the most informative outcome measure). The wrong
choice of either target or timescale may result in unnecessary abandonment of a strategy
that could have a major impact in a few decades. In addition to informing rational use of
interventions that have a positive population impact, our study is also important in cautioning
against the use of strategies that may have little or no lasting population impact. This is
illustrated by our results for catch-up HBV vaccination, which adds little in situations where
high coverage of both neonatal immunization and PMTCT can be attained.
481
482 **Caveats and limitations**
483 Different approaches to recruitment of our HIV-positive and HIV-negative cohorts may have
484 introduced unintentional bias. The KReC children may be less healthy than HIV-negative
485 children in the community, and this approach to recruitment predominantly selected younger
486 children (on average 9.4 months younger than the HIV-positive cohort).
487
We set out to focus on children aged <60 months in order to collect data from the RTHB.
However, in practice, we did not capture good RTHB data and data collection from the
RTHB is itself subject to bias, as families who attend with such records may be those who
are most likely to have immunised their children. Numerous complex social factors are also
relevant in determining immunisation status; babies born to mothers who have HIV and/or
HBV are more likely to be in disadvantaged by poverty, and by illness and death in the
family, such that they might be less likely to present for (or respond to) vaccination.
However, in this setting (and others where antenatal HBV screening is not routinely
deployed (7,39,40)), we deem it unlikely that there is a significant difference in vaccination
rates between infants born to HBV-positive versus HBV-negative mothers.
498
HBV DNA is a more sensitive screening tool than HBsAg but was not practical due to high
cost and lack of availability in this setting. The relatively small numbers in each age group
and the lack of longitudinal follow-up for individual children puts limitations on the data
showing anti-HBs waning over time, but the trends we observe here are biologically
plausible and consistent with existing literature (25,41).

Although we have estimated and parameterized the impact of HIV status on HBV vaccine-

induced protection, we have not modelled other factors related to HIV infection. Namely, we

have not included the potential for increased susceptibility to HBV infection or increased risk

of vertical transmission. These factors would have required further model classes and

specific parameterization, for which little literature support exists. We have also not

considered the influence of population migration on the success of HBV interventions to

reach the elimination targets. Migration of non-immune and/or infected individuals into an

area would delay the time to achieve the targets estimated by our modelling approach.

CONCLUSIONS

Our results affirm the success of the HBV vaccine programme in reducing the prevalence of

HBV in children, with current paediatric prevalence rates of <1%. However, we also highlight

that cases of HBV transmission persist and that a proportion of children are potentially at risk

of infection as a result of low anti-HBs titres, either as a result of missing or incomplete

immunisation, or because of poor antibody titres following vaccination. We predict that

current elimination targets, in particular when framed around reductions of HBeAg-positive

prevalence, are unlikely to be achieved by 2030. For optimum impact, we suggest that

elimination targets should be defined around HBeAg-positive carriers, which are a major

proxy for the public health burden of HBV. Our study highlights the essential need to collect

better data that can help to inform progress towards targets, to optimize deployment of

vaccination and PMTCT, and to invest substantially in education, case finding and treatment.

The prospects of control would be substantially enhanced by improvements in therapy, and

ultimately, the only route to elimination of HBV may be to develop a cure.

DECLARATIONS

Ethics approval and consent to participate

Ethics approval was obtained from the Ethics Committee of the Faculty of Health Science, University of the Free State, Bloemfontein, South Africa (HIV Study Ref: ETOVS Nr 08/09 and COSAC Study Ref: ECUFS NR 80/2014). Written consent for enrollment into the study was obtained from the child's parent/guardian.

Consent for publication

Not applicable

Availability of data and materials

The datasets generated and/or analysed during the current study are available in the Figshare repository: https://figshare.com/s/cd1e4f324606949d1680

Model code will be made available on GitHub on acceptance of the article for publication.

Competing interests

PCM is an Associate Editor for BMC Infectious Diseases and undertakes consultancy work for Immunocore.

Funding

PCM, PK and PJRG are funded by the Wellcome Trust (grant numbers 110110/Z/15/Z to PM, 109965MA to PK, and 104748MA to PJRG); https://wellcome.ac.uk. Recruitment and serological testing of the KReC cohort was covered by a project grant awarded to PCM from the Rosetrees Trust http://www.rosetreestrust.co.uk/. SG and JL received funding from the European Research Council under the European Union's Seventh Framework Programme (FP7/2007-2013)/ERC grant agreement no. 268904-DIVERSITY https://erc.europa.eu/. PK is also funded by an NIHR Senior Fellowship https://www.nihr.ac.uk/. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Authors' Contributions

- Conceived the idea: PM, SG, PJ
- Applied for ethical permission: PM, PG, PJ
- Secured funding: PM, PG, PJ, PK
- Wrote the manuscript draft: AM, JL, PM
- Recruited children and collected clinical data: LH, EA, SD, AvZ, PJ

- Analysed clinical data: EA, CA, AM
- Undertook laboratory assays for HBV serology: SW, KJ
- Developed and tested mathematical model: JL, SG with input from AM, PM, MAA
- All authors read and approved the final manuscript.

Acknowledgements

Not applicable

REFERENCES

1. World Health Organization. Combating Hepatitis B and C to Reach Elimination by 2030: Advocacy Brief. 2016.

2. World Health Organization. Global health sector strategy on viral hepatitis 2016-2021. 2016;http://www.who.int/hepatitis/strategy2016-2021/ghs.

3. Ott JJJ, Horn J, Krause G, Mikolajczyk RTT. Time trends of chronic HBV infection over prior decades – A global analysis. Journal of Hepatology. 2017;66(April 2016):48–54.

4. Matthews PC, Beloukas A, Malik A, Carlson JM, Jooste P, Ogwu A, et al. Prevalence and characteristics of hepatitis B virus (HBV) coinfection among HIV-Positive women in South Africa and Botswana. PLoS ONE. 2015;10(7):e0134037.

5. Schweitzer A, Horn J, Mikolajczyk RT, Krause G, Ott JJ. Estimations of worldwide prevalence of chronic hepatitis B virus infection: A systematic review of data published between 1965 and 2013. The Lancet. 2015;386:1546–55.

6. Statistics South Africa. Mid-year population estimates. 2016;https://www.statssa.gov.za/publications/P0302/P030.

7. O'Hara GA, McNaughton AL, Maponga T, Jooste P, Ocama P, Chilengi R, et al. Hepatitis B Virus as a Neglected Tropical Disease. PLOS Neglected Tropical Diseases. 2017;11(10):https://doi.org/10.1371/journal.pntd.0005842.

8. Mokaya J, McNaughton AL, Burbridge L, Maponga T, O'Hara G, Andersson M, et al. A blind spot? Confronting the stigma of hepatitis B virus (HBV) infection - A systematic review. Wellcome Open Research. 2018;3(0):29.

9. Burnett RJ, Kramvis A, Dochez C, Meheus A. An update after 16 years of hepatitis B vaccination in South Africa. Vaccine. 2012;30(Supplement 3):C45–51.

10. Chotun N, Nel E, Cotton MF, Preiser W, Andersson MI. Hepatitis B virus infection in HIV-exposed infants in the Western Cape, South Africa. Vaccine. 2015;33(36):4618–22.

11. Public Health England. Hepatitis B. In: Green Book: Immunisation against infectious disease. 2016. p. 161–85.

12. Jack A, Hall A, Maine N, Mendy M, Whittle H. What level of hepatitis B antibody is protective? The Journal of Infectious Diseases. 1999;179:489–92.

13. World Health Organization. Hepatitis B vaccines: WHO position paper, July 2017 – Recommendations. Vaccine. 2017;(July).

14. Catherine F-X, Piroth L. Hepatitis B virus vaccination in HIV-infected people: a review. Human Vaccines & Immunotherapeutics. 2017;DOI: 10.1080/21645515.2016.1277844.

612   15.   Lao-araya M, Puthanakit T, Aurpibul L, Taecharoenkul S, Sirisanthana T, Sirisanthana
613   V. Prevalence of protective level of hepatitis B antibody 3 years after revaccination in
614   HIV-infected children on antiretroviral therapy. Vaccine. 2011;29(23):3977–81.
615   16.   Lee C, Gong Y, Brok J, Boxall EH, Gluud C, Eh B, et al. Hepatitis B immunisation for
616   newborn infants of hepatitis B surface antigen-positive mothers (Review). Cochrane
617   Database of Systematic Reviews. 2006;(2):DOI: 10.1002/14651858.CD004790.pub2.
618   17.   Polaris T, Collaborators O. Global prevalence, treatment, and prevention of hepatitis
619   B virus infection in 2016: a modelling study. The Lancet Gastroenterology &
620   Hepatology. 2018;http://dx.doi.org/10.1016/S2468-1253(18)30056-6.
621   18.   Nayagam S, Thursz M, Sicuri E, Conteh L, Wiktor S, Low-Beer D, et al. Requirements
622   for global elimination of hepatitis B: a modelling study. The Lancet Infectious
623   Diseases. 2016;16(12):1399–408.
624   19.   Sharp CP, Gregory WF, Hattingh L, Malik A, Adland E, Daniels S, et al. PARV4
625   prevalence, phylogeny, immunology and coinfection with HIV, HBV and HCV in a
626   multicentre African cohort. Wellcome Open Research. 2017;2(0):26.
627   20.   Jooste P, van Zyl A, Adland E, Daniels S, Hattingh L, Brits A, et al. Screening,
628   characterisation and prevention of Hepatitis B virus (HBV) co-infection in HIV-positive
629   children in South Africa. Journal of Clinical Virology. 2016;85:71–4.
630   21.   Hadler SC, Francis DP, Maynard JE, Thompson SE, Judson FN, Echenberg dean F,
631   et al. Long-term Immunogenicity and Efficacy of Hepatitis B Vaccine in Homosexual
632   Men. New England Journal of Medicine. 1986;315(4):209–14.
633   22.   Hashim A, Dang V, Bolotin S, Crowcroft NS. How and why researchers use the
634   number needed to vaccinate to inform decision making-A systematic review. Vaccine.
635   2015;33(6):753–8.
636   23.   Nayagam S, Conteh L, Sicuri E, Shimakawa Y, Suso P, Tamba S, et al. Cost-
637   effectiveness of community-based screening and treatment for chronic hepatitis B in
638   The Gambia: an economic modelling analysis. The Lancet Global Health.
639   2016;4(8):e568–78.
640   24.   Anderson S, Harper LM, Dionne-Odom J, Halle-Ekane G, Tita ATN. A decision
641   analytic model for prevention of hepatitis B virus infection in Sub-Saharan Africa using
642   birth-dose vaccination. International Journal of Gynecology & Obstetrics.
643   2018;(January):1–7.
644   25.   Chaouch H, Hachfi W, Fodha I, Kallala O, Saadi S, Bousaadia A, et al. Impact and
645   long-term protection of hepatitis B vaccination: 17 years after universal hepatitis B
646   vaccination in Tunisia. Epidemiology and Infection. 2016;144(16):3365–75.
647   26.   Sokal EM, Paganelli M, Wirth S, Socha P, Vajro P, Lacaille F, et al. Management of
648   chronic hepatitis B in childhood: ESPGHAN clinical practice guidelines: Consensus of

an expert panel on behalf of the European Society of Pediatric Gastroenterology,

Hepatology and Nutrition. Journal of Hepatology. 2013;59(4):814–29.

27.   Büchner A, Omar FE, Vermeulen J, Reynders DT. Investigating hepatitis B immunity

in patients presenting to a Paediatric Haematology and oncology unit in South Africa.

South African Medical Journal. 2014;104(9):628–31.

28.   Mayaphi SH, Rossouw TM, Masemola DP, Olorunju SA, Mphahlele MJ, Martin DJ.

HBV/HIV co-infection : The dynamics of HBV in South African patients with AIDS.

South African Medical Journal. 2012;102(3):157–62.

29.   Beghin J-C, Ruelle J, Sokal E, Bachy A, Krishna M, Hall L, et al. Effectiveness of the

South African Expanded Program of Immunization Against Hepatitis B in Children

Infected With Human Immunodeficiency Virus-1 Living in a Resource-Limited Setting

of Kwazulu-Natal. Journal of Medical Virology. 2017;89:182–5.

30.   Ndirangu J, Barnighausen T, Tanser F, Tint K, Newell ML, Bärnighausen T, et al.

Levels of childhood vaccination coverage and the impact of maternal HIV status on

child vaccination status in rural KwaZulu-Natal, South Africa. Tropical Medicine and

International Health. 2009;14(11):1383–93.

31.   Banatvala J, Van Damme P, Oehen S. Lifelong protection against hepatitis B: The

role of vaccine immunogenicity in immune memory. Vaccine. 2000;19(7–8):877–85.

32.   Lao-araya M, Puthanakit T, Aurpibul L, Sirisanthana T, Sirisanthana V. Antibody

response to hepatitis B re-vaccination in HIV-infected children with immune recovery

on highly active antiretroviral therapy. Vaccine. 2007;25(29):5324–9.

33.   Abzug MJ, Warshaw MG, Rosenblatt HM, Levin MJ, Nachman S, Pelton SI, et al.

Immunogenicity and immunologic memory after hepatitis B virus booster vaccination

in HIV-infected children receiving highly active antiretroviral therapy. The Journal of

Infectious Diseases. 2009;200(6):935–46.

34.   Kim HN, Harrington RD, Van Rompaey SE, Kitahata MM. Independent clinical

predictors of impaired response to hepatitis B vaccination in HIV-infected persons.

International journal of STD & AIDS. 2008;19(9):600–4.

35.   Pippi F, Bracciale L, Stolzuoli L, Giaccherini R, Montomoli E, Gentile C, et al.

Serological response to hepatitis B virus vaccine in HIV-infected children in Tanzania.

HIV Medicine. 2008;9(7):519–25.

36.   Mori M, Adland E, Paioni P, Swordy A, Mori L, Laker L, et al. Sex Differences in

Antiretroviral Therapy Initiation in Pediatric HIV Infection. PloS one.

2015;10(7):e0131591.

37.   World Health Organization. Consolidated guidelines on the use of antiretroviral drugs

for treating and preventing HIV infection: recommendations for a public health

approach. World Health Organization.

2016;http://apps.who.int/iris/bitstream/10665/208825/1/.

38.   South African National AIDS Council. Let Our Actions Count: Reflections on NSP

2012-2016 and moving forward to NSP 2017-2022. 2016.

39.   Chotun N, Preiser W, van Rensburg CJ, Fernandez P, Theron GB, Glebe D, et al.

Point-of-care screening for hepatitis B virus infection in pregnant women at an

antenatal clinic☐: A South African experience. PLoS ONE.

2017;12(7):doi.org/10.1371/journal. pone.0181267 Editor:

40.   Diale Q, Pattinson R, Chokoe R, Masenyetse L, Mayaphi S. Antenatal screening for

hepatitis B virus in HIV-infected and uninfected pregnant women in the Tshwane

district of South Africa. South African Medical Journal. 2015;106(1):97–100.

41.   Katoonizadeh A, Sharafkhah M, Ostovaneh MR, Norouzi A, Khoshbakht N,

Mohamadkhani A, et al. Immune responses to hepatitis B immunization 10-18 years

after primary vaccination: a population-based cohort study. Journal of Viral Hepatitis.

2016;23:805–11.

 FIGURE LEGENDS

Figure 1: Diagram of HBV dynamic model. To allow for specific parameterization of
important epidemiological states, the population was divided into susceptibles (S_x) and
vaccinated (V_x) under 3 age-groups representing infants ($x=i$, <1 years of age), children
($x=c$, 1-6 years of age) and older individuals ($x=a$, >6 years of age). Individuals were
assumed to acquire infection at any age, moving with different probabilities (Ψ, ϵ, γ, with $\Psi <$
$\epsilon < \gamma$) into acute (I) or chronic (C) infection. When chronically infected, individuals transit
between HBeAg-positive (C+) and HBeAg-negative (C-) with rate θ, and may clear infection
(R) with a small rate ρ. Vaccine-induced protection is age dependent (Δi) and assumed to
lower susceptibility to infection (λ). Interventions (in blue) include routine vaccination at birth
(Z') and other ages (ωa, ωc), as well as PMTCT at birth (Z, Z') and catch-up events (not
shown). Model is used to fit prevalence rates as observed: HBV prevalence (I + C- + C+),
HBcAb+ (R) and relative prevalence of HBeAg-positive (C+) and HBeAg-negative (C-)
individuals. For a complete description on state transitions, vaccination, force of infection,
parameters and model equations please refer to Suppl Data 2; within this document,
Bayesian parameter estimations obtained when fitting the model are presented in Suppl
Figure 1.

Figure 2: Hepatitis B surface antibody (anti-HBs) titres mediated by vaccination in
HIV-positive (HIV+) and HIV-negative (HIV-) children aged 6-60 months in Kimberley,
South Africa. A: Scatter plot representing vaccine-mediated antibody titres, indicating
median and interquartile ranges (p-value by Mann Whitney U test). B: Proportion of children
with anti-HBs ≥10 mIU/ml or ≥100 mIU/ml (p-values by Fisher's Exact Test).

Figure 3: Relationship between age and vaccine-mediated Hepatitis B surface
antibody (anti-HBs) titres in HIV-positive and HIV-negative children in Kimberley,
South Africa. A: Ages of children attaining anti-HBs titres ≥100 mIU/ml for HIV-positive and
HIV-negative children age 6-60 months. Median ages, interquartile ranges and p-values by
Mann-Whitney U test are indicated. B: Relationship between age and vaccine-mediated Ab
titre among HIV-positive children including those age 6-60 months and an older cohort age
>60 months (range 64-193 months). P-value by Mann Whitney U test. C: Anti-HBs titre and
proportion of subjects with a detectable titre for HIV-positive and HIV-negative children
according to age. On the solid lines, each point represents the mean titre (with 95%
confidence intervals) for the group of children aged ≤12 months (1 yr), 13-24 months (2 yrs),
25-36 months (3 yrs), 37-48 months (4 yrs), 49-60 months (5 years). For the same groups of
children, the dotted lines represent the proportion of subjects with a detectable titre and the

95% confidence intervals. Trends within the data were assessed using linear regression
analysis. D: Odds ratios for protective response to HBV vaccination in children age 6-60
months in Kimberley, South Africa are shown for anti-HBs titre <10mIU/ml and <100mIU/ml
in the whole cohort (green) and in HIV-positive children (black). Statistically significant OR
are denoted * and significant p-values are indicated in bold. Figure 2: Relationship between
age and vaccine-mediated Hepatitis B surface antibody (anti-HBs) titres in HIV-positive and
HIV-negative children in Kimberley, South Africa.

Figure 4: Stochastic impact of neonatal vaccination and PMTCT on HBV incidence
(HBsAg) and HBeAg+ prevalence, showing time to reach sustainable development
goals when using interventions independently. (A1-A2) Impact on **HBV incidence**
(HBsAg) (A1) and time to reach sustainable development goal (SDG) (A2) for varying
routine immunization coverage of neonates. **(B1-B2)** Impact on HBeAg+ prevalence (B1)
and time to reach SDG (B2) for varying routine immunization coverage of neonates. **(C1-C2)**
Impact on **HBV incidence (HBsAg)** (C1) and time to SDG (C2) for varying PMTCT
coverage. **(D1-D2)** Impact on HBeAg+ prevalence (D1) and time to reach SDG (D2) for
varying PMTCT coverage. **(A1, B1, C1, D1)** Lines are the mean and shaded areas the
standard deviation of model output when running 50 stochastic simulations per intervention
(sampling the parameter posteriors shown in Figure 1). **(A2, B2, C2, D2) HBV incidence**
(HBsAg) SDG is set to a reduction of 90%. HBeAg+ prevalence SDG is set to 1/1000
individuals. Beige areas mark interventions reaching SDGs after 500 years on average.
Boxplots show the variation of the 50 stochastic simulations. Numbers above and below
boxplots show the 2.5% lower and 97.5% upper limits of the solutions. **(All suplots)**
Intervention coverage varies from 0.25 to 1 (as coloured and named in subplot A1).

Figure 5: Sensitivity of mean intervention impact on HBV incidence (HBsAg) and
HBeAg+ prevalence based on combinations of routine neonatal vaccination and
PMTCT. (A1-A2) Mean impact of interventions on **HBV incidence (HBsAg)** (A1) and mean
time to reach sustainable development goals (SDGs) (A2). **(B1-B2)** Mean impact of
interventions on HBeAg+ prevalence (B2) and mean time to reach SDG (B2). **(All subplots)**
Impact is shown as percent reduction in incidence or prevalence compared to pre-
intervention levels (e.g. 50 indicates a 50% reduction compared to before the start of the
intervention). **HBV incidence (HBsAg)** SDG is set to a reduction of 90%. HBeAg+
prevalence SDG is set to 1/1000 individuals. Mean results are obtained from 50 stochastic
simulations per intervention combination (vaccination, PMTCT) with parameters sampled
from the posteriors shown in Support Figure 1. Start of interventions in the stochastic
simulations is in year 1995 to simulate an appropriate time scale to address impact by 2030.

Figure 6: Yearly estimated probabilities of achieving sustainable development goals for HBV incidence (HBsAg) and HBeAg+ prevalence based on particular combinations of interventions and local HIV prevalence levels. A total of 1000 stochastic simulations are run independently for each set of particular interventions (coloured legend, subplot A2), with each using a random parameter sample from the posteriors shown in Support Figure 1. Interventions start in year 1995. For every year post-intervention start, the proportion of simulations that have achieved the sustainable development goals (SDGs) is recorded and taken to be the probability. **(A1)** Probability of reaching **HBV incidence (HBsAg)** SDG in time (goal is set to a reduction of 90%). **(A2)** Probability of reaching HBeAg+ prevalence SDG in time (goal is set to 1/1000 individuals). **(B1, B2)** Same as subplots A1-A2 but addressing sensitivity to HIV prevalence levels in the population for a particular intervention (green, $\omega n=0.9$, $\zeta=0.9$, catch-up 0% (WHO)). Solid line is the same as in subplots A1-A2 (named HIV prevalence at baseline). Other lines present results assuming zero HIV prevalence (full line with points) or higher prevalences (dotted, dashed, line with squares). **(All subplots)** The dashed horizontal lines mark 0.5 and 0.975 probability of achieving SDGs. The grey shaded area marks the time period before 2030. In the interventions, ωn is routine vaccination of neonates, ζ the PMTCT effort, ωa routine vaccination of +6 years of age, and catch-up a one-off event of vaccination in some age groups or general population.

SUPPLEMENTARY FILES

Suppl data 1 – Cohort metadata. Metadata for three paediatric cohorts recruited in Kimberley, South Africa, including longitudinal CD4+ T cell and viral load data for paediatric HIV cohort age ≤60 months in Kimberley, South Africa. This file is available on-line via the following link: https://figshare.com/s/cd1e4f324606949d1680.

Suppl data 2 – Mathematical model and supporting figures. Complete model description, including parameters, assumptions, mathematical expressions and fitting details, further including results and figures supporting the main text.

S_x: susceptibles with age x
V_x: vaccinated with age x
C^+: chronic carriers HBeAg+
C^-: chronic carriers HBeAg-
R: recovered HBcAg+
I: acute carriers

A1 dynamics of HBV incidence (HBsAg)

NEONATE ROUTINE VACCINATION

Legend:
- 0.25
- 0.5
- 0.6
- 0.7
- 0.8
- 0.9
- 1

HBV incidence (HBsAg)

years after intervention start

A2 time to reach SDG

time (years)

472
303
251 244 191 186
149 129 106 86 75 61

vaccination coverage (proportion)

B1 dynamics of HBeAg+ prevalence

HBeAg+ prevalence

years after intervention start

B2 time to reach SDG

time (years)

409 351 306 278
265 181 168 136 118 103

vaccination coverage (proportion)

C1 dynamics of HBV incidence (HBsAg)

PMTCT

HBV incidence (HBsAg)

years after intervention start

C2 time to reach SDG

time (years)

446 380
204 155 120

reduction in vertical transmission (proportion)

D1 dynamics of HBeAg+ prevalence

HBeAg+ prevalence

years after intervention start

D2 time to reach SDG

time (years)

reduction in vertical transmission (proportion)

Table 1: Profiles of five children from Kimberley, South Africa, with serological evidence of current or previous infection with HBV (based on positive HBsAg (n=3) or anti-HBc (n=2))

Subject	K306	K405	KReC51	KReC151	K093
Cohort	HIV positive age ≤60 months	HIV positive age ≤60 months	KReC	KReC	HIV positive age >60 months
Sex	F	F	F	M	F
Age (months) at time of sampling	18	37	20	15	118
HIV infection	Positive	Positive	Negative	Negative	Positive
ART[a] (if HIV positive)	Yes	Yes	NA	NA	No
Number of doses of HBV vaccine	NK	NK	NK	3	NK
HBsAg result[b]	Detected	Detected	Detected	Not detected	Not detected
Anti-HBc result[c]	Not detected	Not detected	Detected	Detected	Detected
HBeAg result[d]	Not done	Not done	Detected	Not done	Not done
Anti-HBs result[e]	Not detected	Not detected	Not detected	Detected	Not detected
Interpretation	Active infection	Active infection	Active infection	Immunised, infected and cleared	Infected and cleared

[a]ART indicates the participant was receiving anti-retroviral therapy to treat HIV infection; [b]Hepatitis B surface antigen test; [c]Hepatitis B core antibody test; [d]Hepatitis B envelope antigen test; [e]Hepatitis B surface antibody test (vaccine mediated response). KReC = Kimberley Respiratory Cohort. NA = not applicable. NK=not known. HBV viral loads were not tested in any of these children.

Table 2: Characteristics of three paediatric study cohorts, comprising 402 children, recruited from Kimberley Hospital, South Africa.

Cohort	HIV negative; KReC (age ≤60 months)	HIV positive (age ≤60 months)	HIV positive (age >60 months)
Number of subjects	174	136	92
Age range in months	8-58	6-60	64-193
Median age in months (IQR)	18 (12-26)	29 (18-40)	137 (122-154)
Sex (% male)	55.4	44.9	45.6

KReC = Kimberley Respiratory Cohort. IQR = interquartile range.