Ancestral Reconstruction and C_3 Bundle Sheath Transcript Abundance in the Paniceae Grasses Indicate the Foundations for all Three Biochemical C_4 Sub-Types Were Likely Present in the Most Recent Ancestor

Jacob D. Washburn[1], Satya S. Kothapalli[1], Julia M. Brose[1], Sarah Covshoff[2], Julian M. Hibberd[2], Gavin C. Conant[3,4], J. Chris Pires[1]

1) Division of Biological Sciences, University of Missouri, 311 Bond Life Science Center, Columbia, MO 65211 USA
2) Department of Plant Sciences, University of Cambridge, Downing Street, Cambridge CB2 3EA UK
3) Division of Animal Sciences, 920 East Campus Drive, University of Missouri-Columbia, USA 65211
4) Program in Genetics, Bioinformatics Research Center, Department of Biological Sciences, 356 Ricks Hall, North Carolina State University, Raleigh, NC, USA 27695

ABSTRACT

In C_4 plants the enzymatic machinery underpinning photosynthesis can vary, with for example,

three distinct C_4 acid decarboxylases being used to release CO_2 in the vicinity of RuBisCO. For

decades, these decarboxylases have been used to classify C_4 species into three biochemical sub-

types. However, more recently the notion that C_4 species mix and match C_4 acid decarboxylases

has increased in popularity and, as a consequence, the validity of specific biochemical sub-types

has been questioned. Using species from the grass tribe Paniceae we show that whilst

transcripts encoding multiple C_4 acid decarboxylases accumulate in bundle sheath cells in some

species, in others, transcripts encoding only one enzyme are detected. In addition, a method

that allows isolation of bundle sheath cells from a C_3 species within the Paniceae, *Sacciolepis*

indica, was developed. Deep sequencing of bundle sheath preparations from all four species

combined with ancestral state reconstruction support the notion that the three biochemical C_4

sub-types found in the Paniceae existed together in their most recent common ancestor. Thus,

these species likely inherited the functional building blocks of all three C_4 pathways. We

conclude that classification of C_4 plants into the classical biochemical sub-types is still

appropriate for some species, and that evolution of this trait has been facilitated by

characteristics of the ancestral C_3 bundle sheath and made use of multiple convergent routes

involving either one or multiple C_4 acid decarboxylases.

INTRODUCTION

C_4 photosynthesis is considered the most productive mechanism by which plants convert

sunlight into chemical energy. In all cases, the C_4 pathway leads to increased photosynthetic

efficiency because high concentrations of CO_2 are supplied to RuBisCO. Since its discovery in the

1960s (Hatch and Slack, 1966), a unified understanding of the biochemistry underpinning C_4

photosynthesis has emerged. This basic system comprises a biochemical pump that initially

fixes HCO_3^- into C_4 acids in mesophyll (MS) cells. Subsequently, diffusion of these C_4 acids into a

separate compartment, followed by their decarboxylation, generates high concentrations of

CO_2 around RuBisCO. In many plants, the release of CO_2 occurs in bundle sheath (BS) cells

(Hatch, 1992; Furbank, 2016; von Caemmerer et al., 2017). Although this pump demands

additional ATP inputs, in warm environments where RuBisCO catalyzes high rates of

oxygenation, the C_4 pathway increases photosynthetic efficiency compared with the ancestral

C_3 state.

Elucidation of the C_4 pathway was based on analysis of sugar-cane and maize which both use

the chloroplastic NADP-dependent malic enzyme (NADP-ME) to release CO_2 in BS cells.

However, it became apparent that not all species used this chloroplastic enzyme. For example,

Megathyrsus maximus (formerly *Panicum maximum*), *Urochloa texanum* (formerly *Panicum*

texanum), and *Sporobolus poiretti* used the cytosolic phospho*enol*pyruvate carboxykinase

(PCK) (Edwards et al., 1971) to release CO_2 in the BS, whereas *Atriplex spongiosa* and *Panicum*

miliaceum contained high activities of the mitochondrial NAD-dependent malic enzyme (NAD-

ME) (Hatch and Kagawa, 1974). These findings led to the consensus that different C_4 species

made preferential use of one C_4 acid decarboxylase, and the classification of C_4 plants into one

of three distinct biochemical pathways ensued (Edwards et al., 1971; Hatch et al., 1975; Hatch

and Kagawa, 1976). According to Furbank (2016), there was some early discussion about

whether the sub-types were mutually exclusive or if one species might employ two or more

sub-types together, but in general, the sub-types were described as distinct (Hatch, 1987).

For several decades this description of three sub-types has been standard practice (Sheen,

1999; Hibberd and Covshoff, 2010) and even used in taxonomic classification (Brown, 1977).

However, more recent work has provided evidence that some C_4 species use multiple C_4 acid

decarboxylases. Corn (*Zea mays*), for example, was traditionally classified as using NADP-ME

but evidence has mounted that it also has high activities of PCK (Walker et al., 1997; Wingler et

al., 1999; Majeran et al., 2010; Furbank, 2011; Pick et al., 2011; Bellasio and Griffiths, 2013;

Sharwood et al., 2014; Wang et al., 2014; Koteyeva et al., 2015; Weissmann et al., 2016). This

blurring of the NADP-ME C_4 sub-type coincided with observations that many plants with high

amounts of PCK also contained either NADP-ME of NAD-ME (Furbank, 2011). Furthermore,

computational models of the C_4 pathways suggested that BS energy requirements could not be

met in a system with only PCK decarboxylation (Wang et al., 2014). It has therefore been

suggested that PCK may never function on its own as a distinct sub-type (Furbank, 2011;

Bräutigam et al., 2014; Wang et al., 2014).

Alternatives to the three sub-type classification have since been proposed and used in a

small number of recent publications. These include a two sub-type system (based on the use of

NADP-ME or NAD-ME), as well as a four sub-type classification placing species into NADP-ME,

NAD-ME, NADP-ME + PCK, and NAD-ME + PCK sub-types (Wang et al., 2014; Washburn et al.,

2015; Rao and Dixon, 2016). At present, none of these classification schemes has been widely

adopted by the community. Moreover, convincing experimental evidence (i.e., transcriptomic,

or proteomic data from multiple species) of traditionally PCK sub-type species using another

decarboxylation enzyme at a higher level than PCK is lacking, while enzyme activity

measurements in the older literature indicate PCK predominance for several species (Gutierrez

et al., 1974; Prendergast et al., 1987; Lin et al., 1993).

Figure 1. Species used in this study as well as their phylogenetic placement within the tribe Paniceae (Poaceae) and four of its sub-tribes. The pothosynthetic type (C_3 or C_4) and C_4 sub-type of each species is labeled in the colored triangle next to it. NADP-ME = NADP-dependent malic enzyme, PCK = phosphon*enol*pyruvate carboxykinase, NAD-ME = NAD-dependent malic enzyme. Phylogenetic placement based on Vicentini et al. (2008) and Washburn et al. (In Review).

To better understand whether the PCK pathway represents a true biochemical sub-type, and more generally to investigate the extent to which C_4 species make use of mixtures of C_4 acid decarboxylases, global patterns of mRNA abundance were assessed from BS and MS enriched samples across phylogenetically-spaced C_4 plants that are traditionally defined as exclusively using one of each of the C_4 sub-types. These species belong to the grass tribe Paniceae (Poaceae), the only known group previously documented as containing all three biochemical sub-types of C_4 photosynthesis (Sage et al., 2011). The C_3 species *Sacciolepis indica*, another member of the Panicea, was also included in this analysis to provide insight into the ancestral state in the tribe and the evolutionary transition from C_3 to different C_4 sub-types.

RESULTS

Distribution of Transcripts Encoding Components of The Core C$_4$ Cycle

Four C$_4$ species from the Paniceae tribe that have been reported to use different C$_4$ acid

decarboxylases were selected for analysis (Figure 1). *Setaria italica* is a member of the

Cenchrinae subtribe within the Paniceae and

belongs to the classical NADP-ME C$_4$ sub-

type. *Urochola fusca* is in the subtribe

Melinidinae which is classically defined as

using the PCK sub-type. *Panicum hallii*

belongs to the Panicinae subtribe and is

reported to use NAD-ME, and lastly *Digitaria*

californica,   a   member   of   the

Anthephorineae subtribe, is defined as

conforming to the NADP-ME sub-type

(Gutierrez et al., 1974; Prendergast et al.,

1987; Lin et al., 1993). Although belonging to

distinct C$_4$ biochemical sub-types these

species are derived from a common ancestor

(Figure 1). *Sacciolepis indica* represents the

C$_3$ ancestral state, and is phylogenetically

sister to the other four species while still

being part of the tribe Paniceae.

Figure 2. Representative images from leaves that have been rolled to remove mesophyll (MS) sap (Left) or bundle sheath (BS) strands after isolation (Right). All species use the C$_4$ pathway except *Sacciolepis indica* which is a C$_3$ plant. The bands of cells with low chlorophyll content in images on the left-hand side represent the position of mesophyll cells that have collapsed and sap expelled during the rolling procedure. Scale bars are depicted.

Microscopic examination of leaves of *S. italica, U. fusca, P. hallii* and *D. californica* from

which MS cell sap had been extracted using leaf rolling techniques, indicated bands of cells

containing low chlorophyll content (Figure 2) a phenotype consistent with efficient removal of

MS sap (Covshoff et al., 2013; John et al., 2014). In addition, after mechanical isolation of

leaves, BS preparations of high purity for all C_4 species were generated. Separation of BS

strands was also successful for the C_3 species *Sacciolepis indica* (Figure 2). Analysis of transcripts

derived from core C_4 genes showed clear differences in abundance between MS and BS

samples. For example, transcripts derived from carbonic anhydrase (CA), phospho*enol*pyruvate

carboxylase (PEPC) and pyruvate, orthophosphate dikinase (PPDK) genes preferentially

accumulated in MS cells (Figure 3A). In contrast, transcripts derived from the rubisco small

Figure 3. Log$_2$ fold change between mesophyll (MS) and bundle sheath (BS) enriched mRNA transcripts of *Setaria italica, Urochloa fusca, Panicum hallii* and *Digitaria californica*, or between whole leaf and BS strands for C_3 *Sacciolepis indica*. Genes depicted encode proteins of the core C_4 cycle that are known to be preferentially expressed in either: A) MS, or B) BS cells. The number of asterisks in each box represents the p-value. *** p < 0.001, ** p <0.01, * p < 0.05. CA = carbonic anhydrase, PEPC = phospho*enol*pyruvate carboxylase, NADP-MDH 1 or 2 = NADP-dependent malate dehydrogenase (two different isoforms), PPDK = pyruvate,orthophosphate dikinase, NADP-ME = NADP-dependent malic enzyme, PCK = phosphon*enol*pyruvate carboxykinase, NAD-ME = NAD-dependent malic enzyme, NAD-MDH = NADP-dependent malate dehydrogenase, RCA = rubisco activase, RBCS = rubisco small subunit.

subunit (RBCS) and rubisco activase (RCA) as well as either NADP-ME, NAD-ME or PCK were

more abundant in BS strands (Figure 3B). These differences are consistent with our current

understanding of transcript distribution between MS and BS cells in C_4 grasses (Covshoff et al.,

2013; John et al., 2014). The abundance of transcripts relating to C_4 photosynthesis in the C_3

species *S. indica* is also consistent with current knowledge of metabolism in the BS of C_3

species. For example, transcripts of RBCS and RCA were less abundant in BS strands than in

whole leaf samples, which supports the notion that the C_3 BS being less photosynthetic than C_3

MS cells (Kinsman and Pyke, 1998; Leegood, 2002; Aubry et al., 2014). Furthermore, although

not to the same degree as in the C_4 species sampled, it was also the case that transcripts

encoding each of the C_4 acid decarboxylases were more abundant in the BS strands than in the

whole leaf of *S. indica* (Figure 3). This is consistent with the finding that cells near veins of C_3

species contain relatively high activities of these C_4 enzymes (Hibberd and Quick, 2002; Brown

et al., 2010).

Some Paniceae Lineages Use Classical Sub-Types and Others Mix Decarboxylases

Setaria italica has classically been defined as belonging to the NADP-ME C_4 sub-type

(Gutierrez et al., 1974; Prendergast et al., 1987; Lin et al., 1993). Accumulation of the NADP-ME

and NADP-MDH enzymes in BS and MS cells respectively is indicative of this sub-type, and their

transcript distributions confirmed this expectation. In addition, consistent with knowledge of

the NADP-ME sub-type, in BS strands of *S. italica* transcripts encoding PCK, NAD-ME, NAD-MDH,

ASP-AT, and ALA-AT were detected at low levels. The data therefore indicate that under the

conditions we used for sampling, *S. italica* was likely making near exclusive use of NADP-ME to

decarboxylate malate in the BS (Figure 4A).

Figure 4. Relative transcript abundance between core C$_4$ pathway enzymes within mesophyll (MS) and bundle sheath (BS) of: A) *Setaria italica,* B) *Urochloa fusca,* C) *Panicum hallii,* and D) *Digitaria californica.* The schematics below each histogram indicate the enzyme complement associated with each of the three biochemical sub-types. CA = carbonic anhydrase, PEPC = phospho*enol*pyruvate carboxylase, NADP-MDH 1 or 2 = NADP-dependent malate dehydrogenase (two different isoforms), PPDK = pyruvate,orthophosphate dikinase, NADP-ME = NADP-dependent malic enzyme, PCK = phosphon*enol*pyruvate carboxykinase, NAD-ME = NAD-dependent malic enzyme, NAD-MDH = NADP-dependent malate dehydrogenase, RCA = rubisco activase, RBCS = rubisco small subunit, asparagine-aminotransferase (ASP-AT) and alanine-amino transferase (ALA-AT)

Urochola fusca is thought to exclusively use PCK to release CO_2 in the BS (Gutierrez et al.,

1974; Prendergast et al., 1987; Lin et al., 1993). The patterns of transcript accumulation in MS

and BS strands of *U. fusca* are consistent with PCK functioning in this species without

supplemental decarboxylation from either NADP-ME or NAD-ME (Figure 4B). For example, BS

stands contained barely detectable levels of transcripts encoding NADP-ME and NAD-ME, but

very high levels of those encoding PCK. In addition, consistent with the cycling of aspartate and

alanine between the two cell-types, transcripts derived from genes encoding both ASP-AT and

ALA-AT were detectable in both cell-types (Figure 4B).

In contrast to the above analysis of *S. italica* and *U. fusca*, which supports the contention

that they make almost exclusive use of NADP-ME and PCK respectively, analysis of *P. hallii* and

D. californica indicates that they likely use multiple C_4 acid decarboxylases during

photosynthesis. Although *P. hallii* is classically considered as using NAD-ME (Gutierrez et al.,

1974; Prendergast et al., 1987; Lin et al., 1993) in addition to high levels of transcripts encoding

NAD-ME, NAD-MDH, ASP-AT and ALA-AT, unexpectedly high levels of transcripts encoding

NADP-ME were detected in the BS (Figure 4C). In the case of *Digitaria californica* which is

thought to belong to the NADP-ME sub-type, although transcripts encoding NADP-ME and

NADP-MDH were abundant in BS and MS samples respectively, PCK levels were more than

double those of NADP-ME in the BS. ASP-AT levels in the MS cells were also higher than

expected for an NADP-ME sub-type plant. Overall, these data are consistent with *S. italica* and

U. fusca using a single C_4 acid decarboxylase in the BS, whereas *P. hallii* and *D. californica* do

not.

Figure 5. Three hypotheses for the evolution of C_4 sub-types within the MPCA clade (Melinidinae, Panicinae, Cenchrinae, Anthephorinae) of the tribe Paniceae. The one sub-type hypothesis posits that the most recent common ancestor (MRCA) of the group utilized one sub-type exclusively, and the other types evolved from it is a step-wise fashion. The three sub-type hypothesis suggests that all three sub-types existed in the MRCA and then each has become dominant in one clade or another over time. The C_3 hypothesis is based on the idea that each of the sub-types evolved independently from a C_3 ancestor. Hypothesizes and phylogeny based on Washburn et al. (2015) and Washburn et al. (In Review). NADP-ME = NADP-dependent malic enzyme, PCK = phosphon*enol*pyruvate carboxykinase, NAD-ME = NAD-dependent malic enzyme.

Ancestral State Reconstruction of the MPCA Clade

Several hypotheses relating to how the MPCA (Melinidinae, Panicinae, Cenchrinae, and

Anthephorinae) clade may have evolved have been proposed (Figure 5). While they have been

examined using categorical data, this is the first time that ancestral state reconstructions of the

MPCA have been performed using transcript abundance levels. The results at each node of the

MPCA phylogeny show mixtures of all three primary sub-type enzymes (Figure 6). Although this

approach indicates that transcripts for NADP-ME and PCK would have been the most abundant

in the last common ancestor of these C$_4$ species, it was also the case that substantial levels of

NAD-ME are predicted.

Figure 6. Comparisons between species for transcript abundance of NADP-dependent malic enzyme (NADP-ME), phosphon*enol*pyruvate carboxykinase (PCK), and NAD-dependent malic enzyme (NAD-ME) transcripts in Bundle Sheath strands. Below that, the nuclear gene phylogenetic relationships and ancestral state reconstruction of transcript abundance levels at each node are displayed. Transcript levels are normalized to those of Rubisco activase (RCA). Phylogenetic relationships based on Vicentini et al. (2008) and Washburn et al. (In Review).

DISCUSSION

The PCK Sub-Type

The dominance of PCK transcripts over NADP-ME and NAD-ME in *U. fusca* provide evidence

for the biological relevance of the classical PCK sub-type. These data are in direct contrast to

recent proposals that PCK cannot function on its own but rather is always ancillary to one of the

other two C_4 acid decarboxylases (Furbank, 2011; Bräutigam et al., 2014; Wang et al., 2014).

While this notion may be true in some cases, the current results suggest that it is not the case

for *U. fusca*. Furthermore, our findings are supported by earlier measurements of enzyme

activity made within the sub-tribe Melinidinae where PCK was also shown to be highly

dominant over the other sub-types (Gutierrez et al., 1974; Prendergast et al., 1987; Lin et al.,

1993), and also indicate that these differences in activity are due to differences in steady-state

transcript abundance rather than post-transcriptional modifications that act to suppress

accumulation of NADP-ME and NAD-ME.

In the case of *U. fusca,* it appears that the use of alternative sub-type classification systems

where PCK is considered ancillary to either NADP-ME or NAD-ME (Wang et al., 2014; Washburn

et al., 2015; Rao and Dixon, 2016) is not useful. In fact, NAD-ME and NADP-ME transcript

abundance in *U. fusca* were statistically indistinguishable making this species impossible to

classify into the proposed two and four sub-type systems.

C_4 Sub-Type Mixing

Of the four C_4 species examined, *P. hallii* and *D. californica* show evidence of extensive sub-

type mixing. The potential for mixing has previously been considered in *Panicum virgatum*, a

close relative of *P. hallii* (Zhang et al., 2013; Meyer et al., 2014; Rao and Dixon, 2016; Rao et al.,

2016). Rao et al. (2016) suggested that the high abundance of NADP-ME transcripts may be

accounted for by post-transcriptional or translational modifications but experimental evidence

for testing that hypothesis is lacking.

Digitaria californica also showed evidence of sub-type mixing. In this case, NADP-ME and

PCK transcripts were both reasonably abundant. Although this species is classically considered

to belong to the NADP-ME sub-type, transcripts encoding PCK were more than double the

abundance of those of NADP-ME. ASP-AT levels, which are also associated with the PCK sub-

type, were high as well. This suggests that *D. californica* may use PCK as the dominant

decarboxylation enzyme and NADP-ME secondarily.

Bundle Sheath Enriched Transcript Abundance Levels in C$_3$ Species and the Evolution of Three

Sub-Types Within a Single Clade

Analysis of transcript abundance in MS and BS cells from C$_3$ species that are closely related

to C$_4$ species is critical to understanding how C$_4$ photosynthesis evolved, and how it can be

engineered for enhanced crop production. Although transcripts loaded onto ribosomes in the

BS of C$_3$ *Arabidopsis thaliana* have been assessed, and this analysis provided insight into the

role of the BS in eudicot plants (Aubry et al., 2014), to our knowledge there is no equivalent

data from a monocotyledonous lineage in which both C$_3$ and C$_4$ species are found. The

relatively high levels of C$_4$ related transcripts in the BS of *S. indica* are consistent with previous

work on cells around the veins of C$_3$ plants (Hibberd and Quick, 2002; Brown et al., 2010; Shen

et al., 2016). Together these data support the concept that C$_3$ species are in some ways pre-

adapted to adopt the C_4 mechanism (Gould, 1989; Christin et al., 2009; Christin et al., 2015;

Washburn et al., 2016).

The results from ancestral state reconstruction favor the notion that the three C_4 sub-types

in MPCA clade in the Paniceae may have existed together in their most recent common

ancestor. The result that two of the four species sampled may be mixing sub-types is further

evidence for this hypothesis. Additionally, the presence of transcripts associated with multiple

enzymes involved in each of the three C_4 sub-types in the C_3 species *S. indica* supports all three

sub-types existing together in the MRCA rather than each sub-type evolving independently, or

sub-types evolving in a step-wise fashion one from another.

MATERIALS AND METHODS

Plant Materials

Accessions from five plant species were used in this study: *Setaria italica* yugu1, *Urochloa*

fusca LBJWC-52, *Panicum hallii* FIL2, *Digitaria californica* PI 364670, and *Sacciolepis indica* PI

338609. More details on the accessions can be found in Washburn et al. (2015) with exception

of *P. hallii* FIL2 obtained from Thomas Juenger of the University of Texas at Austin with further

details at *Panicum hallii* v2.0, DOE-JGI, http://phytozome.jgi.doe.gov/.

All plant materials were grown in controlled growth chambers at the University of Missouri

in Columbia. Plants were grown under 16 hours of light (from 6:00-20:00) and 8 hours of

darkness with temperatures of 23C during the day and 20C at night.   Lights were placed

between 86-88 cm above the plants. Plantings were grown in 4 replicates in a completely

randomized design with 32 plants per replicate (except for *Sacciolepis indica* where plants were

smaller and grown with 64 plants per replicate). The third leaf was sampled between 11:00 and

15:00 using established leaf rolling and mechanical BS isolation methods with some

modifications (See Supplemental Materials S1 for complete modified protocol) (Sheen and

Bogorad, 1985; Chang et al., 2012; Covshoff et al., 2013; John et al., 2014).

The protocol used for obtaining BS strands in *S. indica* was the same as that used for the C_4

species. Variations on the amount of time for each blending step were investigated, but only

resulted in higher levels of contamination as viewed under a microscope. That said, even when

microscopic examination indicated higher contamination levels in some *S. indica* BS samples,

transcript abundance results were qualitatively similar samples with less apparent

contamination. One step that may have been key to the isolation of C_3 BS strands, was the use

of leaf rolling on the sampled leaves just prior to the BS strand isolation procedure (Furbank et

al., 1985; John et al., 2014). This enables the removal of at least some MS sap prior to BS

isolation.

MS enriched samples where not successfully obtained for *S. indica* in this study because of

the sensitivity of the C_3 leaves to rolling. Very small amounts of leaf rolling pressure resulted in

the leaves becoming damaged to the point that the purity of MS sap obtained from them was

called into question. Leaves that were rolled gently enough not to damage the BS strands and

contaminate the MS sap resulted in sap with insufficient quantities of RNA for sequencing. It is

our opinion that MS sap could be sampled using: 1) low input mRNA extraction and sequencing

procedures, 2) a more precise instrument for leaf rolling such as that described by Leegood

(1985), and/or 3) further experimentation with the developmental stage at which MS sap is

extracted.

Sequencing

RNA was extracted using the PureLink® RNA Mini Kit (Invitrogen, Carlsbad, CA, USA) and

mRNA-seq libraries were constructed and sequenced by the University of Missouri DNA Core

Facility using the TruSeq Stranded mRNA Sample Prep Kit (Illumina, Inc., San Diego, CA, USA)

and the Illumina HiSeq and NextSeq platforms.

Analysis

Each mRNA sample was quality trimmed and mapped to the *Sorghum bicolor* genome

(Paterson et al., 2009; DOE-JGI, 2017) using Trimmomatic and Trinity following the workflows

outlined on the Trinity website (Grabherr et al., 2011; Haas et al., 2013; Bolger et al., 2014). This

processing included the use of RSEM and Bowtie2 for read mapping and counting as well as

edgeR and DESeq for differential expression analysis (Robinson et al., 2010; Li and Dewey, 2011;

Langmead and Salzberg, 2012; McCarthy et al., 2012; Love et al., 2014). A list of known C_4

photosynthesis genes was compiled based on the literature; a custom script and BLAST were

then used to find the appropriate homologous genes for each species and to compare their

relative abundance levels (Camacho et al., 2009; Chang et al., 2012; Covshoff et al., 2013;

Bräutigam et al., 2014; John et al., 2014; Tausta et al., 2014; Rao et al., 2016). Ancestral state

reconstructions where performed using the "Trace Character History" command in Mesquite

(Maddison and Maddison, 2017).

Transcript Normalization and Transcriptome Size Estimation

For comparisons across all cell types and species within this study, The Trimmed Mean of M-

values (TMM) method described by Robinson and Oshlack (2010) as implemented in DESeq was

used. For comparisons of BS enhanced transcripts between the different species, transcript

abundance levels were normalized to Rubisco activase (RCA). In the current study, BS cell

analyses where tested separately using normalization to RCA and TMM and the results were

qualitatively identical.

ACCESSION NUMBERS

Sequence data are available on NCBI SRA (https://www.ncbi.nlm.nih.gov/sra) under the

following identifiers: XXX

ACKNOWLEDGMENTS

This work was supported by grants from the University of Missouri (Mizzou Advantage, MU

Research Board, and Molecular Life Science Fellowships), and the U.S. National Science

Foundation (DEB Award no. 1501406).

LITERATURE CITED

Aubry S, Smith-Unna RD, Boursnell CM, Kopriva S, Hibberd JM (2014) Transcript residency on

ribosomes reveals a key role for the Arabidopsis thaliana bundle sheath in sulfur and

glucosinolate metabolism. The Plant Journal **78:** 659-673

Bellasio C, Griffiths H (2013) The operation of two decarboxylases (NADPME and PEPCK),

transamination and partitioning of C4 metabolic processes between mesophylll and

bundle sheath cells allows light capture to be balanced for the maize C4 pathway. Plant

Physiology

Bolger AM, Lohse M, Usadel B (2014) Trimmomatic: a flexible trimmer for Illumina sequence

data. Bioinformatics **30**

Bräutigam A, Schliesky S, Külahoglu C, Osborne CP, Weber APM (2014) Towards an integrative

model of C4 photosynthetic subtypes: insights from comparative transcriptome analysis

of NAD-ME, NADP-ME, and PEP-CK C4 species. Journal of Experimental Botany **65:** 3579-

3593

**Brown NJ, Palmer BG, Stanley S, Hajaji H, Janacek SH, Astley HM, Parsley K, Kajala K, Quick

WP, Trenkamp S, Fernie AR, Maurino VG, Hibberd JM** (2010) C4 acid decarboxylases

required for C4 photosynthesis are active in the mid-vein of the C3 species Arabidopsis

thaliana, and are important in sugar and amino acid metabolism. The Plant Journal **61:**

122-133

Brown WV (1977) The Kranz syndrome and its subtypes in grass systematics. Memoirs of the

Torrey Botanical Club **23:** 1-97

Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden T (2009)

BLAST+: architecture and applications. BMC Bioinformatics **10:** 421

**Chang YM, Liu WY, Shih AC, Shen MN, Lu CH, Lu MY, Yang HW, Wang TY, Chen SC, Chen SM, Li

WH, Ku MS** (2012) Characterizing regulatory and functional differentiation between

maize mesophyll and bundle sheath cells by transcriptomic analysis. Plant Physiol **160:**

165-177

Christin P-A, Samaritani E, Petitpierre B, Salamin N, Besnard G (2009) Evolutionary insights on

C4 photosynthetic subtypes in grasses from genomics and phylogenetics. Genome

Biology and Evolution **1:** 221-230

Christin PA, Arakaki M, Osborne CP, Edwards EJ (2015) Genetic enablers underlying the

clustered evolutionary origins of C4 photosynthesis in angiosperms. Molecular Biology

and Evolution **32:** 846-858

Covshoff S, Furbank RT, Leegood RC, Hibberd JM (2013) Leaf rolling allows quantification of

mRNA abundance in mesophyll cells of sorghum. Journal of Experimental Botany **64:**

807-813

DOE-JGI (2017) *Sorghum bicolor* v3.1. *In*,

Edwards GE, Kanai R, Black CC (1971) Phosphoenolpyruvate carboxykinase in leaves of certain

plants which fix CO2 by the C4-dicarboxylic acid cycle of photosynthesis. Biochemical

and Biophysical Research Communications **45:** 278-285

Furbank RT (2011) Evolution of the C4 photosynthetic mechanism: are there really three C4

acid decarboxylation types? Journal of Experimental Botany **62:** 3103-3108

Furbank RT (2016) Walking the C 4 pathway: past, present, and future. Journal of Experimental

Botany **67:** 4057-4066

Furbank RT, Stitt M, Foyer CH (1985) Intercellular compartmentation of sucrose synthesis in

leaves of Zea mays L. Planta **164:** 172-178

Gould SJ (1989) Wonderful life: the Burgess Shale and the nature of history, Ed 1st. W.W.

Norton, New York

Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, Adiconis X, Fan L,

Raychowdhury R, Zeng Q, Chen Z, Mauceli E, Hacohen N, Gnirke A, Rhind N, di Palma

F, Birren BW, Nusbaum C, Lindblad-Toh K, Friedman N, Regev A (2011) Full-length

transcriptome assembly from RNA-Seq data without a reference genome. Nat

Biotechnol **29:** 644-652

Gutierrez M, Gracen VE, Edwards GE (1974) Biochemical and cytological relationships in C4

plants. Planta **119:** 279-300

Haas BJ, Papanicolaou A, Yassour M, Grabherr M, Blood PD, Bowden J, Couger MB, Eccles D,

Li B, Lieber M, MacManes MD, Ott M, Orvis J, Pochet N, Strozzi F, Weeks N,

Westerman R, William T, Dewey CN, Henschel R, LeDuc RD, Friedman N, Regev A

(2013) De novo transcript sequence reconstruction from RNA-seq using the Trinity

platform for reference generation and analysis. Nat. Protocols **8:** 1494-1512

Hatch M, Kagawa T, Craig S (1975) Subdivision of C4-Pathway Species Based on Differing C4

Acid Decarboxylating Systems and Ultrastructural Features. Functional Plant Biology **2:**

111-128

Hatch M, Slack C (1966) Photosynthesis by sugar-cane leaves. A new carboxylation reaction and the pathway of sugar formation. Biochemical Journal **101:** 103-111

Hatch MD (1987) C4 photosynthesis: a unique blend of modified biochemistry, anatomy and ultrastructure. Biochimica et Biophysica Acta (BBA) - Reviews on Bioenergetics **895:** 81-106

Hatch MD (1992) I can't believe my luck. Photosynthesis Research **33:** 1-14

Hatch MD, Kagawa T (1974) Activity, location and role of NAD malic enzyme in leaves with C4-pathway photosynthesis. Austral. J. Pl. Physiol **1:** 357-369

Hatch MD, Kagawa T (1976) Photosynthetic activities of isolated bundle sheath cells in relation to differing mechanisms of C4 pathway photosynthesis. Archives of Biochemistry and Biophysics **175:** 39-53

Hibberd JM, Covshoff S (2010) The regulation of gene expression required for C4 photosynthesis. Annu Rev Plant Biol **61:** 181-207

Hibberd JM, Quick WP (2002) Characteristics of C4 photosynthesis in stems and petioles of C3 flowering plants. Nature **415:** 451-454

John CR, Smith-Unna RD, Woodfield H, Covshoff S, Hibberd JM (2014) Evolutionary Convergence of Cell-Specific Gene Expression in Independent Lineages of C4 Grasses. Plant Physiology **165:** 62-75

Kinsman EA, Pyke KA (1998) Bundle sheath cells and cell-specific plastid development in Arabidopsis leaves. Development **125:** 1815-1822

Koteyeva NK, Voznesenskaya EV, Edwards GE (2015) An assessment of the capacity for phosphoenolpyruvate carboxykinase to contribute to C4 photosynthesis. Plant Science **235:** 70-80

Langmead B, Salzberg SL (2012) Fast gapped-read alignment with Bowtie 2. Nat Methods **9**

Leegood RC (1985) The intercellular compartmentation of metabolites in leaves of Zea mays L. Planta **164:** 163-171

Leegood RC (2002) C4 photosynthesis: principles of CO2 concentration and prospects for its introduction into C3 plants. Journal of Experimental Botany **53:** 581-590

Li B, Dewey CN (2011) RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. BMC Bioinformatics **12**

Lin C, Tai Y, Liu D, Ku M (1993) Photosynthetic mechanisms of weeds in taiwan. Functional Plant Biology **20:** 757-769

Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol **15**

Maddison WP, Maddison DR (2017) Mesquite: a modular system for evolutionary analysis. *In*, Ed Version 3.2

Majeran W, Friso G, Ponnala L, Connolly B, Huang M, Reidel E, Zhang C, Asakura Y, Bhuiyan NH, Sun Q, Turgeon R, van Wijk KJ (2010) Structural and Metabolic Transitions of C(4) Leaf Development and Differentiation Defined by Microscopy and Quantitative Proteomics in Maize. The Plant Cell **22:** 3509-3542

McCarthy DJ, Chen Y, Smyth GK (2012) Differential expression analysis of multifactor RNA-Seq experiments with respect to biological variation. Nucleic Acids Research **40:** 4288-4297

Meyer E, Aspinwall MJ, Lowry DB, Palacio-Mejía JD, Logan TL, Fay PA, Juenger TE (2014)

Integrating transcriptional, metabolomic, and physiological responses to drought stress

and recovery in switchgrass (Panicum virgatum L.). BMC Genomics **15:** 527

Paterson AH, Bowers JE, Bruggmann R, Dubchak I, Grimwood J, Gundlach H, Haberer G,

Hellsten U, Mitros T, Poliakov A, Schmutz J, Spannagl M, Tang H, Wang X, Wicker T,

Bharti AK, Chapman J, Feltus FA, Gowik U, Grigoriev IV, Lyons E, Maher CA, Martis M,

Narechania A, Otillar RP, Penning BW, Salamov AA, Wang Y, Zhang L, Carpita NC,

Freeling M, Gingle AR, Hash CT, Keller B, Klein P, Kresovich S, McCann MC, Ming R,

Peterson DG, Mehboob ur R, Ware D, Westhoff P, Mayer KFX, Messing J, Rokhsar DS

(2009) The Sorghum bicolor genome and the diversification of grasses. Nature **457:** 551-

Pick TR, Brautigam A, Schluter U, Denton AK, Colmsee C, Scholz U, Fahnenstich H, Pieruschka

R, Rascher U, Sonnewald U, Weber AP (2011) Systems analysis of a maize leaf

developmental gradient redefines the current C4 model and provides candidates for

regulation. Plant Cell **23:** 4208-4220

Prendergast H, Hattersley P, Stone N (1987) New structural/biochemical associations in leaf

blades of C4 grasses (Poaceae). Functional Plant Biology **14:** 403-420

Rao X, Dixon RA (2016) The Differences between NAD-ME and NADP-ME Subtypes of C4

Photosynthesis: More than Decarboxylating Enzymes. Frontiers in Plant Science **7**

Rao X, Lu N, Li G, Nakashima J, Tang Y, Dixon RA (2016) Comparative cell-specific

transcriptomics reveals differentiation of C4 photosynthesis pathways in switchgrass

and other C4 lineages. Journal of Experimental Botany **67:** 1649-1662

Robinson MD, McCarthy DJ, Smyth GK (2010) edgeR: a Bioconductor package for differential

expression analysis of digital gene expression data. Bioinformatics **26**

Robinson MD, Oshlack A (2010) A scaling normalization method for differential expression

analysis of RNA-seq data. Genome Biology **11:** R25

Sage RF, Christin PA, Edwards EJ (2011) The C4 plant lineages of planet earth. Journal of

Experimental Botany **62:** 3155-3169

Sharwood RE, Sonawane BV, Ghannoum O (2014) Photosynthetic flexibility in maize exposed

to salinity and shade. Journal of Experimental Botany **65:** 3715-3724

Sheen J (1999) C4 GENE EXPRESSION. Annual Review of Plant Physiology & Plant Molecular

Biology **50:** 187

Sheen J-Y, Bogorad L (1985) Differential Expression of the Ribulose Bisphosphate Carboxylase

Large Subunit Gene in Bundle Sheath and Mesophyll Cells of Developing Maize Leaves Is

Influenced by Light. Plant Physiology **79:** 1072-1076

Shen W, Ye L, Ma J, Yuan Z, Zheng B, LV C, Zhu Z, Chen X, Gao Z, Chen G (2016) The existence

of C4-bundle-sheath-like photosynthesis in the mid-vein of C3 rice. Rice **9:** 20

Tausta SL, Li P, Si Y, Gandotra N, Liu P, Sun Q, Brutnell TP, Nelson T (2014) Developmental

dynamics of Kranz cell transcriptional specificity in maize leaf reveals early onset of C4-

related processes. Journal of Experimental Botany

Vicentini A, Barber JC, Aliscioni SS, Giussani LM, Kellogg EA (2008) The age of the grasses and

clusters of origins of C4 photosynthesis. Global Change Biology **14:** 2963-2977

von Caemmerer S, Ghannoum O, Furbank RT (2017) C4 photosynthesis: 50 years of discovery

and innovation. Journal of Experimental Botany **68:** 97-102

Walker RP, Acheson RM, Técsi LI, Leegood RC (1997) Phosphoenolpyruvate Carboxykinase in C4 Plants: Its Role and Regulation. Functional Plant Biology **24:** 459-468

Wang Y, Bräutigam A, Weber APM, Zhu X-G (2014) Three distinct biochemical subtypes of C4 photosynthesis? A modelling analysis. Journal of Experimental Botany **65:** 3567-3578

Washburn JD, Bird KA, Conant GC, Pires JC (2016) Convergent Evolution and the Origin of Complex Phenotypes in the Age of Systems Biology. International Journal of Plant Sciences **177:** 305-318

Washburn JD, Schnable JC, Conant GC, Brutnell TP, Shao Y, Zhang Y, Ludwig M, Davidse G, Pires JC (In Review) Genome-Guided Phylo-Transcriptomic Methods and the Nuclear Phylogentic Tree of the Paniceae Grasses.

Washburn JD, Schnable JC, Davidse G, Pires JC (2015) Phylogeny and photosynthesis of the grass tribe Paniceae. American Journal of Botany **102:** 1493-1505

Weissmann S, Ma F, Furuyama K, Gierse J, Berg H, Shao Y, Taniguchi M, Allen DK, Brutnell TP (2016) Interactions of C4 Subtype Metabolic Activities and Transport in Maize Are Revealed through the Characterization of DCT2 Mutants. The Plant Cell **28:** 466-484

Wingler A, Robert PW, Zhi-Hui C, Leegood RC (1999) Phosphoenolpyruvate Carboxykinase Is Involved in the Decarboxylation of Aspartate in the Bundle Sheath of Maize. Plant Physiology **120:** 539-545

Zhang J-Y, Lee Y-C, Torres-Jerez I, Wang M, Yin Y, Chou W-C, He J, Shen H, Srivastava AC, Pennacchio C, Lindquist E, Grimwood J, Schmutz J, Xu Y, Sharma M, Sharma R, Bartley LE, Ronald PC, Saha MC, Dixon RA, Tang Y, Udvardi MK (2013) Development of an

integrated transcript sequence database and a gene expression atlas for gene discovery

and analysis in switchgrass (Panicum virgatum L.). The Plant Journal **74:** 160-173