Characterization of a set of abdominal neuroendocrine cells that regulate stress physiology using colocalized diuretic peptides in *Drosophila*

Meet Zandawala[1, *], Richard Marley[2], Shireen A. Davies[2] and Dick R. Nässel[1, *]

1. Stockholm University, Department of Zoology, Stockholm, Sweden

2. Institute of Molecular, Cell and Systems Biology, College of Medical, Veterinary and Life Sciences, University of Glasgow, Glasgow, UK

* Correspondence to: Dr. Dick R. Nässel and Dr. Meet Zandawala

Tel: +46-8-164077

E-mail: dnassel@zoologi.su.se and meet.zandawala@zoologi.su.se

ORCID for D.R.N.: 0000-0002-1147-7766

ORCID for M.Z.: 0000-0001-6498-2208

Acknowledgements

The authors would like to acknowledge Dr. Yiting Liu and Dr. Olga Kubrak for very helpful advice and valuable discussions. We are grateful to the Bloomington *Drosophila* Stock Center, the Vienna *Drosophila* Resource Center and Drs. Jan A. Veenstra, Pilar Herrero, Young Joon Kim and Michael Texada for providing flies and reagents. Stina Höglund and the Imaging Facility at Stockholm University (IFSU) are acknowledged for maintenance of the confocal microscopes. This work was supported by a European Commission Horizon 2020, Research and Innovation Grant 634361.

Abstract

Multiple neuropeptides are known to regulate water and ion balance in *Drosophila melanogaster*. Several of these peptides also have other functions in physiology and behavior. Examples are corticotropin-releasing factor-like diuretic hormone (diuretic hormone 44; DH44) and leucokinin (LK), both of which induce fluid secretion by Malpighian tubules (MTs), but also regulate stress responses, feeding, circadian activity and other behaviors. Here, we investigated the functional relations between the LK and DH44 signaling systems. DH44 and LK peptides were only colocalized in a set of abdominal neurosecretory cells (ABLKs). Targeted knockdown of each of these peptides in ABLKs lead to increased resistance to desiccation, starvation and ionic stress. Food ingestion was diminished by knockdown of DH44, but not LK, and water retention was increased by LK knockdown only. Thus, the two colocalized peptides display similar systemic actions, but differ with respect to regulation of feeding and body water retention. We also demonstrated that DH44 and LK have additive effects on fluid secretion by MTs. It is likely that the colocalized peptides are coreleased from ABLKs into the circulation and act on the tubules where they target different cell types and signaling systems to regulate diuresis and stress tolerance. Additional targets seem to be specific for each of the two peptides and subserve regulation of feeding and water retention. Our data suggest that the ABLKs and hormonal actions are sufficient for many of the known DH44 and LK functions, and that the remaining neurons in the CNS play other functional roles.

Introduction

Orchestration of physiological and behavioral processes is commonly dependent on neuropeptide and peptide hormone signaling [see 1,2-4]. For example, feeding and postprandial effects on the organism, including satiety, nutrient and energy reallocation, diuresis, and activity/sleep are regulated by multiple peptides [see 5,6,1,7-9,3]. Thus, in *Drosophila melanogaster,* several neuropeptides such as allatostatins, neuropeptide F, short neuropeptide F (sNPF), sulfakinin, and hugin-pyrokinin are known to regulate feeding, and four peptides, diuretic hormones 31 and 44, leucokinin and CAPA regulate ion and water homeostasis [see 10,11,12,1,7,3]. After a meal other hormones, like insulin-like peptides (ILPs) and adipokinetic hormone (AKH) ensure energy mobilization and storage, or signal satiety/hunger and affect organismal activity, vigor and stress tolerance [13-15,7].

The neuroendocrine cells producing the peptides mentioned above display varying degrees of diversity, from a single small set of identical cells producing AKH or ILPs to very large populations of diverse neurons expressing sNPF [16-18]. Thus, the question is whether peptidergic neurons of a large diverse population are functionally coupled and play a concerted physiological role, or if they are parts of distributed networks where the specific neuropeptide therefore serve diverse functions. To address this question we have selected a set of neuroendocrine cells producing the neuropeptide leucokinin (LK) that consists of four morphological types of cells [19,20], and is proposed to serve multiple functions in flies [21-25,6]. This set of neurons was dissected by genetic tools to enable us to isolate the functional role of a major subset, which consists of prominent neurosecretory cells in the abdominal ganglia. We show that this subset of the LK neurons, designated ABLKs, additionally produce corticotrophin-releasing factor-like diuretic hormone, also known as diuretic hormone 44 (DH44). These ABLKs are especially intriguing since they seem to be under tight neuronal and hormonal control. Receptors for several neurotransmitters and peptides have been identified on these cells in larvae [26,27,23,28-31] and adults [23]. Of these receptors, only the action of the 5-HT1B receptor on ABLK function was probed in adults [23]. We ask what function these specific neuroendocrine cells and their colocalized peptide hormones have in physiology and behavior.

Both DH44 and LK are primarily known for their roles as diuretic hormones in various insects, including *Drosophila,* by regulating secretion in Malpighian (renal) tubules (MTs) [32,12,3,33]. However, several additional functions have been assigned to these peptides from genetic experiments. LK neurons regulate food intake, play a role in desiccation stress resistance, modulate chemosensory responses, decrease postprandial sleep, and are required for starvation-induced sleep suppression [21-25,6]. DH44, which is produced by a diverse set of neurons and neurosecretory cells [32,34], plays roles in osmotic and metabolic stresses. Knockdown of DH44 in the CNS or its receptor (DH44-R2) in the MTs results in a significant increase in desiccation tolerance [22]. Genetic ablation of DH44 neurons also results in increased starvation tolerance, however, knockdown of the DH44 receptor, DH44-R2, in the MTs impairs starvation tolerance [22]. Furthermore, DH44 producing median neurosecretory cells regulate rhythmic locomotor activity with influence from clock cells [35] and in females these neurons also regulate sperm retention in the uterus [34].

The postulated functions of LK and DH44 are, with a few exceptions, not assigned to specific neurons. Using the GAL4-UAS system [36] we target interference with LK and DH44 expression to the ABLKs and analyze *in vivo* effects on tolerance to various stressors, as well as feeding and water retention. We also employ an assay to monitor the combined activity of DH44 and LK4 on secretion in MTs. These peptides act on different cell types of the *Drosophila* MTs and activate different signal pathways [32,33], yet we show that they display an additive stimulatory effect on secretion. Thus we can show that the ABLKs, and therefore hormonal actions of the two peptides, are sufficient for regulating water and ion homeostasis and associated stress functions, but can also affect food intake, perhaps by an indirect action caused by diuretic activity. This suggests that the remaining LK and DH44 neurons may play other functional roles in the fly, and it remains to be determined whether these functions are in any way linked to those of the ABLKs.

Experimental procedures

Fly lines and husbandry

All fly strains used in this study (**Table 1**) were reared and maintained at 25°C on a standard yeast, corn meal, agar medium (see http://flystocks.bio.indiana.edu/Fly_Work/media-recipes/bloomfood.htm) supplemented with 1.5 g/L nipagin and 3 mL/L propionic acid. Experimental flies were reared under uncrowded conditions and normal photoperiod (12 hours light: 12 hours dark).

Table 1: Fly strains used in this study

Fly strain	Inserted on chromosome	Source / reference	Stock number
w^{1118}	-	BDSC	-
w^{1118}; Lk-GAL4	II	[20], Pilar Herrera, Madrid, Spain	-
w^{1118}; Lk-GAL4	II	[37]	BDSC 51993
w^{1118}; Lk-GAL4	III	Young Joon Kim, Gwangju, Korea	-
w^{1118}; DH44-GAL4	III	[38], Fly Light	BDSC 39347
w^{1118}; DH44-GAL4	III	[37]	BDSC 51987
$y^1 w^*$; Actin5c-GAL4/CyO	II	-	BDSC 4414
w^{1118};UAS-Lk RNAi	III	[39]	VDRC 14091/GD
yw;UAS-DH44 RNAi	II	-	VDRC 108473/KK
yw; Sco/Cyo; UAS-mcd8-GFP	III	BDSC	-
JFRC29-10xUAS-IVS-myr::GFP-p10	-	[40], Washington, USA	-

Immunohistochemistry and imaging

Immunohistochemistry for *Drosophila* larval and adult CNS was performed as described earlier [41]. Briefly, CNS from third instar larvae or adult male flies were dissected in phosphate buffered saline (PBS). Larval samples were fixed for 2 hours in 5% ice-cold paraformaldehyde and adult samples were fixed on ice for 3.5 – 4 hours. Samples were then washed in PBS and incubated for 48 hours at 4°C in primary antibodies diluted in PBS with 0.5% Triton X (PBST) (**Table 2**). Following this incubation, samples were

washed with PBST and incubated for 48 hours at 4°C in secondary antibodies diluted in PBST (**Table 2**). Next, samples were thoroughly washed with PBST, and following a final wash in PBS, samples were mounted in 80% glycerol. For anti-DH44 staining, tissues were blocked with 5% normal goat serum (NGS) in PBST post-fixation and 5% NGS was also included in the primary antibody solution.

Table 2: Antibodies used for immunohistochemistry

Antibody	Immunogen	Source / reference	Dilution
Primary antibody			
Rabbit anti-LK	*Leucophaea maderae* kinin I	[42]	1:2000
Rabbit anti-DH44	*Drosophila melanogaster* DH44	[32] Jan Veenstra, Bordeaux, France	1:1000
Mouse anti-GFP	Jelly fish GFP	Invitrogen	1: 1000
Secondary antibody			
Goat anti-mouse Alexa 488	-	Invitrogen	1:1000
Goat anti-rabbit Alexa 546	-	Invitrogen	1:1000

All samples were imaged with a Zeiss LSM 780 confocal microscope (Jena, Germany) using 10X, 20X or 40X oil immersion objectives. Confocal images were processed with Zeiss LSM software and Fiji [43] for projection of z-stacks, contrast and brightness, and calculation of immunofluorescence levels.

Stress resistance assays

We used 5-6 days old male flies to assay for survival under various stresses and recovery from chill coma (see [41] for details of stress assays). For each technical replicate, 15 flies were kept in a vial and their survival recorded every 3 hours (for desiccation) or 6 hours (for starvation and ionic stress) until all the flies were dead. For desiccation, flies were kept in empty vials. For starvation, flies were kept in vials containing 5 ml of 0.5% aqueous agarose (A2929, Sigma-Aldrich). For ionic stress, flies were kept in vials containing 5 ml enriched medium (100 g/L sucrose, 50 g/L yeast, 12 g/L agar, 3 ml/L propionic acid and 3 g/L nipagin) supplemented with 4% NaCl. All vials were kept at 25°C under normal photoperiod conditions for the entire duration of the experiment. For chill coma recovery experiments, flies were transferred to empty vials,

which were then placed on ice to induce a chill coma. The vials were incubated on ice

(0°C) for 4 hours and afterward transferred to 25°C to induce recovery. The number of

flies recovered was assessed every 2 min. At least 3 biological replicates and 3

technical replicates for each biological replicate were performed for each experiment.

Capillary feeding assay

Capillary feeding (CAFE) assay to measure food intake for individual flies was

performed according to the method described earlier [23]. Food consumption was

measured daily and the cumulative food intake over 4 days was calculated. The

experiment consisted of 3 biological replicates and 8–10 flies per replicate for each

genotype.

Water content measurement

For measurement of water content, 10-15 flies were frozen on dry ice and their weight

(wet weight) recorded using a Mettler Toldeo MT5 microbalance (Columbus, USA). The

flies were then dried for 1 day at 60°C and their weight (dry weight) recorded again. The

water content of the flies was determined by subtracting dry weight from wet weight

(see [44]).

Malpighian tubule secretion assay

Drosophila MT fluid secretion assays were performed as described previously [45].

Briefly, MTs from six day old *Drosophila* were dissected in Schneider's medium and

transferred to 9 μl of 50% Schneider's medium and 50% *Drosophila* saline [32]. Tubules

were left to secret for 30 minutes and non-secreting tubules were replaced to form a

data set of 10-15 secreting tubules. Basal secretion was measured for 30 min at 10 min

intervals. Following this initial incubation, 1 μl of *Drosophila* DH44 (final concentration

10^{-7} M or 10^{-9} M), LK (final concentration 10^{-9} M or 10^{-10} M) or both (Genosphere

Biotechnologies, Paris, France) were added to the incubation medium. Stimulated

secretion was measured for 30 min at 10 min intervals. Data are presented as the

secretion rate at every time point, the percentage change in secretion rate following

peptide application, and total fluid secreted over 60 min.

 Statistical analyses

The experimental data are presented as means ± s.e.m. Unless stated otherwise, one-way analysis of variance (ANOVA) followed by Tukey's multiple comparisons test was used for comparisons between three genotypes and an unpaired t-test was used for comparisons between two genotypes. For fluid secretion assays, a Mann-Whitney U test was used as some data were non-normally distributed. Stress curves were compared using Mantel-Cox log-rank test. All statistical analyses were performed using GraphPad Prism with a 95% confidence limit ($p < 0.05$).

Results

LK expression in Drosophila CNS

Several studies have previously examined the distribution of LK in *Drosophila* CNS and peripheral tissues [19,20,23,46,22]. Here, we verify the expression of *Lk-GAL4* driven GFP in both larval (Figure S1A-E) and adult *Drosophila* (Figure 1), using a GAL4 line from De Haro et al. [20]. In the larval CNS, *Lk-GAL4* drives expression in five pairs of neurons in the brain (Figure S1A, E), three pairs of neurons in the SOG (Figure S1C, E) and seven pairs of neurons in the VNC (Figure S1D, E). However, four out of the five pairs in the brain do no display any LK-immunoreactivity, but in fact react with an antiserum to ion transport peptide (ITP) (Figure S1B). These lateral neurosecretory cells are found both in larval and adult brains and are known as ipc-1 neurons [47]. Similar to the larval CNS, *Lk-GAL4* drives GFP expression in four distinct neuronal populations in the adult CNS (Figure 1). Hence, GFP expression was detected in one pair of neurons in the lateral horn and one pair in the SOG (Figure 1A, D), another four pairs in the brain, which display ITP-immunoreactivity (Figure 1B, D and S2) and eleven pairs in the VNC (Figure 1C, D). In adult flies, *Lk-GAL4* driven GFP expression in the ITP-producing cells is weak and variable (Figure S2). Interestingly, seven pairs of neurons in the VNC start expressing LK in the embryonic stage, whereas the additional four pairs (this number varies between individuals) begin to express LK during pupal development [46]. Moreover, *Lk-GAL4* also drives ectopic expression in salivary glands (Figure 1E). Although we tested two additional *Lk-GAL4* driver lines (Table 1), we utilized this *Lk-GAL4* for subsequent knockdown experiments because its expression was stronger than, or more specific than the other *Lk-GAL4* lines (Table 1; data not shown).

DH44 is expressed in the Drosophila brain and ventral nerve cord

DH44 expression in *Drosophila* has also been examined and mapped previously [32,34]. However, different GAL4 lines result in differing expression patterns. Thus, we validated the expression of two previously generated *DH44-GAL4* lines [37,38]. One of these GAL4 lines has a minimal expression pattern and drives GFP expression in only the six MNCs in the brain (data not shown) [37]. The other *DH44-GAL4* line, which was obtained from the FlyLight collection [38], resulted in a good overlap between *DH44>GFP* and DH44-immunoreactivity. This *DH44-GAL4* drives GFP expression broadly in larval (Figure S1F, G) and adult CNS (Figure 2). In larvae, predominant expression was detected in the six MNCs in the brain (Figure S1F) and seven pairs of neurosecretory cells in the VNC (Figure S1G); however, some of these cells in the VNC were not visible in both hemiganglia. In adults, expression was again detected in the six MNCs (Figure 2A) and at least four cells in the anterior (Figure 2B) and at least six cells in the posterior VNC (Figure 2C). Since there was a good overlap in GFP expression and DH44-immunoreactivity, we utilized this *DH44-GAL4* for subsequent knockdown manipulations.

LK and DH44 are co-expressed in larval and adult ABLK neurons

The functional overlap between LK and DH44 signaling systems mentioned earlier coupled with the expression of both LK and DH44 in the VNC neurons prompted us to examine if these two neuropeptides are co-expressed in subsets of neurons. Our expression data show that there is no overlap in LK and DH44 expression in the brain of larvae (Figure S3A, C and Figure 4A) or adults (Figure 3A, C and Figure 4B), but these neuropeptides are co-expressed in the larval (Figure S3B, D and Figure 4A) and adult ABLKs (Figure 3B, D and Figure 4B). Interestingly, all the larval ABLKs also express DH44 but in adults only four to eight ABLKs express DH44 (Figure 4). Furthermore, the majority of these ABLKs appear to be the ones that are generated during postembryonic neurogenesis. However, we cannot rule out the possibility that DH44 has a very low expression in other ABLKs.

Knockdown of Lk with Lk-GAL4 impacts stress response and water content

Having shown that LK and DH44 are co-expressed in ABLKs, it now becomes apparent that the previous studies employing genetic ablation, and activation or inactivation of LK

neurons could be confounded by effects of diminishing signaling with two colocalized peptides [24,23,22]. Thus, it is not only timely to dissect the behavioral phenotypes from these previous studies using RNAi-based knockdown, but we can also study the specific functions of ABLKs using intersectional crosses. Consequently, we utilized *Lk-GAL4* and *DH44-GAL4* to knock down both *Lk* and *DH44* and assayed for effects on stress tolerance, feeding and water content.

Lk-GAL4 driven *Lk-RNAi* results in a significant decrease in LK-immunoreactivity in both the brain (Figure 5A, C) and VNC (Figure 5B). These flies with *Lk* knockdown display increased survival under desiccation (Figure 6A), starvation (Figure 6B), and ionic stress (Figure 6C), but there is no difference in chill coma recovery between experimental and control flies (Figure 6D). Moreover, food intake in CAFE assay is not affected by *Lk* knockdown (Figure 6E) (see also [21]), but these flies retain more water (Figure 6F) as demonstrated earlier [48,23]. Previous work had shown that inactivation of LK neurons resulted in increased survival under desiccation, but had no impact on starvation resistance [23]. Furthermore, in that study, both the activation and inactivation of LK neurons caused the flies to feed less in CAFE assay. This was in contrast to another study employing LK and LKR mutants, which did not find altered overall food intake, but rather an increase in meal size [21]. Hence, our data on desiccation is in agreement with previous findings and clarifies the discrepancies in previous results on food intake as possibly caused by the presence of another neuroactive compound in the LK neurons that affects feeding.

Knockdown of DH44 with Lk-GAL4 impacts stress response and feeding

We verified the efficiency of *DH44* knockdown using a ubiquitous driver (*Actin 5c-GAL4*) and a specific driver (*DH44-GAL4*). Knockdown of DH44 with both these drivers results in a significant decrease in DH44-immunoreactivity (Figure 5F, G and S4). We then utilized *Lk-GAL4* to knock down *DH44* specifically in ABLKs, which resulted in increased survival under desiccation (Figure 7A), starvation (Figure 7B), ionic stress (Figure 7C), and these flies showed delayed recovery from chill coma compared to control flies (Figure 7D). Interestingly, *DH44* knockdown flies feed less in the CAFE assay (Figure 7E), but display no difference in water content compared to controls (Figure 7F). Hence, it appears that the effect of decreased feeding following LK neuron inactivation can be attributed to the presence of DH44 in ABLKs. Furthermore, LK but not DH44 has an

effect on water content. Perhaps this could be due to the fact that LK is a more potent diuretic than DH44 in *Drosophila* and hence DH44 cannot fully compensate for the lack of LK [32,33]. Alternatively, LK could also impact water retention via actions on the hindgut [49,50].

Knockdown of Lk with DH44-GAL4 impacts stress response and water content

Next, we wanted to see the effects of knocking down *Lk* in adult-specific ABLKs. We first confirmed that *DH44-GAL4* driven *Lk-RNAi* results in an efficient knockdown in adult-specific ABLKs by counting the number of cells positive for LK-immunoreactivity (Figure 5D and S5). The average number of cells stained for LK-immunoreactivity in the control flies was 21, whereas the knockdown flies only had an average of 16 cells. Moreover, the larger adult-specific ABLKs are not stained in the knockdown flies validating that the knockdown is efficient. Knockdown of *Lk* with *DH44-GAL4* resulted in increased survival during desiccation (Figure 8A), starvation (Figure 8B), ionic stress (Figure 8C), as well as a significant delay in recovery from chill coma (Figure 8D). Similar to the global *Lk* knockdown with *Lk-GAL4, Lk* knockdown in ABLKs has no effect on feeding (Figure 8E) but results in a significant increase in the water content of the flies (Figure 8F). This suggests that the effects of LK on stress response and water content could be attributed to ABLKs, and perhaps the LHLKs and SELKs of the brain play little to no part in these processes.

Knockdown of DH44 with DH44-GAL4 impacts stress response and feeding

Knockdown of DH44 with DH44-GAL4 results in an efficient knockdown in the brain (Figure 5F) and VNC (Figure 5G). Staining is abolished in MNCs but not in the other cells in the brain suggesting that staining in those cells is not specific for DH44 (Figure 5F). In order to determine if there is any interaction between LK and DH44 signaling, we measured LK peptide levels in ABLKs of *DH44* knockdown flies. Interestingly, flies with *DH44* knockdown have higher LK levels suggesting that the flies may compensate for the lack of DH44 with increased LK expression (Figure 5D, E). Moreover, flies with global *DH44* knockdown displayed no effects on survival during desiccation (Figure S6A), but showed increased resistance to starvation (Figure S6B), ionic stress (Figure S6C), and a small but significant delay in their recovery from chill coma (Figure S6D).

Furthermore, flies with *DH44* knockdown displayed no difference in feeding (Figure
S6E) and water content (Figure S6F) compared to control flies.

LK and DH44 act additively on Malpighian tubules to stimulate fluid secretion

Since LK and DH44 are coexpressed in ABLKs, they could potentially be coreleased in

to the hemolymph and result in functional interaction at the target tissue. One such site

of interactions is the MTs, since both these peptides stimulate MT secretion albeit by

action on different cell types [32,33]. Hence, we were interested in examining the

secretion rates by MTs and the volume of secreted fluid in the presence of either

peptide alone or in the presence of both (Figure 9). Our results show that the addition of

both LK and DH44 results in a secretion rate that is a sum of the secretions rates

obtained following the addition of each of those peptides separately (Figure 9A-D). This

effect was more prominent when a higher dose of DH44 (10^{-7} M instead of 10^{-9} M) was

used (Figure 9C, D). The amount of fluid secreted with peptide stimulation is also a

reflection of these increased secretion rates (Figure 9E, F). Hence, 10^{-7} M DH44 and

10^{-10} M LK result in almost identical volumes of fluid secreted (Figure 9E), whereas a

combination of both those peptides doubles the volume of fluid secreted suggesting an

additive response.

Table 3: Comparison of secretion rates between various treatments and time points presented
in Figure 9A and B. (NS = not significant, * $p < 0.05$, ** $p < 0.01$, **** $p < 0.0001$; Mann-Whitney
U test).

Treatments compared	Time (min)		
	40	50	60
10^{-7}M DH44 vs 10^{-7}M DH44 + 10^{-10}M LK	****	****	****
10^{-10}M LK vs 10^{-7}M DH44 + 10^{-10}M LK	****	****	****
10^{-9}M DH44 vs 10^{-9}M DH44 + 10^{-10}M LK	NS	**	NS
10^{-10}M LK vs 10^{-9}M DH44 + 10^{-10}M LK	*	*	NS

Knockdown of Lk in ABLKs does not influence LK stimulated Malpighian tubule secretion

Previous studies have shown that knockdown of peptides could influence the expression of their receptors and vice versa (see [41]). We wanted to determine whether knockdown of *Lk* in ABLKs, the only source of hormonal LK, affects the expression of LKR in MTs, thus influencing LK-stimulated secretion by MTs. Our results indicate that there is no significant difference in LK-stimulated (10^{-9}M and 10^{-10}M) secretion rates of MTs isolated from *DH44 > Lk RNAi* and control flies (Figure S7). This is similar to previous work where DH44 (10^{-7}M) secretion rates were similar in tubules isolated from *DH44 > DH44 RNAi* and control flies [22]. These results are in agreement with the *in vivo* experiments where flies with *Lk* knockdown display increased survival under desiccation. This phenotype would not be so apparent if the flies were able to compensate for the lack of LK.

Discussion

Our study revealed that a portion of the LK expressing neurosecretory cells (ABLKs) in abdominal ganglia coexpress DH44, similar to earlier findings in the moth *Manduca sexta* [51], the locust *Locusta migratoria* [52] and blood sucking bug *Rhodnius prolixus* [53]. Colocalization of these peptides in multiple insect orders, including basal orders, suggests that this colocalization and the subsequent functional interaction between these signaling systems evolved early on during insect evolution. Since ABLKs are the sole neurons producing both peptides in *Drosophila* we were able to use GAL4 lines to knock down each of the two peptides in these cells only and thereby isolate the contribution of the ABLKs to physiology. This enabled us to establish that these neuroendocrine cells are sufficient for many of the functions assigned to DH44 and LK and therefore these functions are hormonally mediated. In contrast, earlier studies were based upon altering peptide levels or activity in entire populations of DH44 and LK neurons [21-25,6]. Also, we showed here that the *LK-GAL4* driver includes a set of ectopic brain cells (ipc-1) that do not express LK, but another peptide ITP. The ipc-1 neurons produce sNPF and tachykinin in addition to ITP and have been found to regulate stress responses [44]. This means that in earlier studies where the *LK-GAL4* line is used to drive inactivation or activation of neurons [see e.g. [23,24]] additional phenotypes are likely to have arisen. Using our approach, where we target only ABLK

neurons, we find that both *DH44-RNAi* and *Lk-RNAi* in these cells increases resistance to desiccation, starvation and ionic stress. This suggests that diminishing the release of these two peptides from ABLKs is sufficient for this phenotype to occur. However, food intake is not affected by LK-knockdown in ABLKs, whereas DH44 knockdown diminishes feeding, and conversely knockdown of LK in ABLKs result in increased body water content, that is not seen after *DH44-RNAi*. Thus, the two colocalized peptides appear to display similar systemic actions, but differ with respect to feeding and water retention. When knocking down LK in all LK neurons we obtained a very similar set of effects as when we targeted only the ABLKs, indicating that in the assays we performed in our study the other two sets of LK neurons (LHLK and SELK) played a minimal role in the assays performed in this study.

We also demonstrated that DH44 and LK have additive effects on fluid secretion in MTs. It is likely that these two colocalized peptides are released together and act on the MTs where they target different cell types and signaling systems in order to regulate fluid secretion [32,33]. The action of these peptides on the MTs may also in part be responsible for the regulation of stress responses seen in our assays, as shown earlier for CAPA peptide and DH44 [54,22]. It is, however, not clear whether the altered food intake and water retention after DH44 and LK knockdown, respectively, are direct actions on target tissues or indirect effects caused by altered water and ion regulation in the fly.

Not only do the ABLKs produce two diuretic hormones, they also seem to be under tight neuronal and hormonal control. Receptors for several neurotransmitters and peptides have been identified on these cells in adults: the serotonin receptor 5-HT1B, LK receptor (LkR) and the insulin receptor, dInR [23,28]. Knockdown of the 5-HT1B receptor in ABLK neurons diminished LK expression, increased desiccation resistance, and diminished food intake, but manipulations of dInR expression in these cells generated no changes in physiology in the tests performed [23]. In larvae, all ABLKs colocalize LK and DH44, and several receptors have been detected in addition to 5-HT1B [23,29] and dInR [28], namely RYamide receptor [31], SIFamide receptor [30], and the ecdysis-triggering hormone (ETH) receptor, ETHR-A [27]. However, the expression of these receptors on adult ABLKs has so far not been investigated. Interestingly the functions of ABLKs in larvae, studied so far, seem to be primarily related to regulating muscle activity and ecdysis motor patterns. The 5-HT-1B receptor

on ABLKs was shown to modulate locomotor turning behavior [29], whereas ETH mediated activation of ETHR-A on ABLKs initiates the pre-ecdysis motor activity [26,27]. In this context, it is worth noting that during metamorphosis 6-8 novel ABLKs differentiate anteriorly in the abdominal ganglia [46,28], and these are the ones that display the strongest expression of DH44. In adult flies the ABLKs are neurosecretory cells with restricted arborizations in the CNS, but widespread axon terminations along the abdominal body wall and in the lateral heart nerves, whereas in larvae the same cells send axons that terminate on segmental abdominal muscles, muscle 8 [19]. It is not yet known whether larval ABLKs are involved in the regulation of diuresis and other related physiological functions *in vivo*, but certainly larval functions in locomotion and ecdysis behavior are specific to that developmental stage. Thus, it seems that there is a developmental switch of function in this set of peptidergic neuroendocrine cells.

In summary, we show that a set of abdominal neuroendocrine cells, ABLKs, coexpressing DH44 and LK are sufficient for regulation of resistance to desiccation, starvation and ionic stress, as well as modulating feeding and water content in the body. These ABLKs represent a subset of neurons that express DH44 and LK, and the functions of the remaining neurons have yet to be determined.

Conflict of Interest Statement

The authors declare that they have no conflict of interest.

Author contributions

M.Z., S.A.D. and D.R.N.: designed the research; M.Z. and R.M.: performed experiments and analyzed data; M.Z. and D.R.N.: wrote the manuscript with input from the other authors; D.R.N. and S.A.D.: obtained funding; D.R.N.: supervised the study.

References

1. Nässel DR, Winther AM (2010) *Drosophila* neuropeptides in regulation of physiology and behavior. Prog Neurobiol 92 (1):42-104. doi:10.1016/j.pneurobio.2010.04.010
2. Schoofs L, De Loof A, Van Hiel MB (2017) Neuropeptides as Regulators of Behavior in Insects. Annual Review of Entomology, Vol 62 62:35-52. doi:10.1146/annurev-ento-031616-035500
3. Schooley DA, Horodyski FM, Coast GM (2012) Hormones Controlling Homeostasis in Insects In: Gilbert LI (ed) Insect Endocrinology. Academic Press, San Diego, pp 366-429. doi:http://dx.doi.org/10.1016/B978-0-12-384749-2.10009-3
4. Taghert PH, Nitabach MN (2012) Peptide neuromodulation in invertebrate model systems. Neuron 76 (1):82-97. doi:10.1016/j.neuron.2012.08.035
5. Itskov PM, Ribeiro C (2013) The dilemmas of the gourmet fly: the molecular and neuronal mechanisms of feeding and nutrient decision making in *Drosophila*. Front Neurosci 7:12. doi:10.3389/fnins.2013.00012
6. Murphy KR, Deshpande SA, Yurgel ME, Quinn JP, Weissbach JL, Keene AC, Dawson-Scully K, Huber R, Tomchik SM, Ja WW (2016) Postprandial sleep mechanics in *Drosophila*. Elife 5. doi:10.7554/eLife.19334
7. Pool AH, Scott K (2014) Feeding regulation in *Drosophila*. Curr Opin Neurobiol 29:57-63. doi:10.1016/j.conb.2014.05.008
8. Saper CB, Chou TC, Elmquist JK (2002) The need to feed: homeostatic and hedonic control of eating. Neuron 36 (2):199-211
9. Zeltser LM, Seeley RJ, Tschop MH (2012) Synaptic plasticity in neuronal circuits regulating energy balance. Nat Neurosci 15 (10):1336-1342. doi:10.1038/nn.3219
10. Chen J, Reiher W, Hermann-Luibl C, Sellami A, Cognigni P, Kondo S, Helfrich-Forster C, Veenstra JA, Wegener C (2016) Correction: Allatostatin A Signalling in *Drosophila* Regulates Feeding and Sleep and Is Modulated by PDF. PLoS Genet 12 (12):e1006492. doi:10.1371/journal.pgen.1006492
11. Dow JA (2009) Insights into the Malpighian tubule from functional genomics. J Exp Biol 212 (Pt 3):435-445. doi:10.1242/jeb.024224
12. Halberg KA, Terhzaz S, Cabrero P, Davies SA, Dow JA (2015) Tracing the evolutionary origins of insect renal function. Nat Commun 6:6800. doi:10.1038/ncomms7800
13. Alfa RW, Kim SK (2016) Using *Drosophila* to discover mechanisms underlying type 2 diabetes. Dis Model Mech 9 (4):365-376. doi:10.1242/dmm.023887
14. Owusu-Ansah E, Perrimon N (2014) Modeling metabolic homeostasis and nutrient sensing in *Drosophila*: implications for aging and metabolic diseases. Dis Model Mech 7 (3):343-350. doi:10.1242/dmm.012989
15. Padmanabha D, Baker KD (2014) *Drosophila* gains traction as a repurposed tool to investigate metabolism. Trends Endocrinol Metab 25 (10):518-527. doi:10.1016/j.tem.2014.03.011
16. Brogiolo W, Stocker H, Ikeya T, Rintelen F, Fernandez R, Hafen E (2001) An evolutionarily conserved function of the *Drosophila* insulin receptor and insulin-like peptides in growth control. Curr Biol 11 (4):213-221
17. Kim SK, Rulifson EJ (2004) Conserved mechanisms of glucose sensing and regulation by *Drosophila* corpora cardiaca cells. Nature 431 (7006):316-320. doi:10.1038/nature02897
18. Nässel DR, Enell LE, Santos JG, Wegener C, Johard HA (2008) A large population of diverse neurons in the *Drosophila* central nervous system expresses short neuropeptide F, suggesting multiple distributed peptide functions. BMC Neurosci 9:90. doi:10.1186/1471-2202-9-90
19. Cantera R, Nässel DR (1992) Segmental peptidergic innervation of abdominal targets in larval and adult dipteran insects revealed with an antiserum against leucokinin I. Cell Tissue Res 269 (3):459-471

20. De Haro M, Al-Ramahi I, Benito-Sipos J, Lopez-Arias B, Dorado B, Veenstra JA, Herrero P
(2010) Detailed analysis of leucokinin-expressing neurons and their candidate functions in
the *Drosophila* nervous system. Cell Tissue Res 339 (2):321-336. doi:10.1007/s00441-009-
0890-y
21. Al-Anzi B, Armand E, Nagamei P, Olszewski M, Sapin V, Waters C, Zinn K, Wyman RJ,
Benzer S (2010) The leucokinin pathway and its neurons regulate meal size in *Drosophila*.
Curr Biol 20 (11):969-978. doi:10.1016/j.cub.2010.04.039
22. Cannell E, Dornan AJ, Halberg KA, Terhzaz S, Dow JA, Davies SA (2016) The corticotropin-
releasing factor-like diuretic hormone 44 (DH44) and kinin neuropeptides modulate
desiccation and starvation tolerance in *Drosophila melanogaste*r. Peptides 80:96-107.
doi:10.1016/j.peptides.2016.02.004
23. Liu Y, Luo J, Carlsson MA, Nässel DR (2015) Serotonin and insulin-like peptides modulate
leucokinin-producing neurons that affect feeding and water homeostasis in *Drosophila*. J
Comp Neurol 523 (12):1840-1863. doi:10.1002/cne.23768
24. Lopez-Arias B, Dorado B, Herrero P (2011) Blockade of the release of the neuropeptide
leucokinin to determine its possible functions in fly behavior: chemoreception assays.
Peptides 32 (3):545-552. doi:10.1016/j.peptides.2010.07.002
25. Murakami K, Yurgel ME, Stahl BA, Masek P, Mehta A, Heidker R, Bollinger W, Gingras RM,
Kim YJ, Ja WW, Suter B, DiAngelo JR, Keene AC (2016) translin Is Required for Metabolic
Regulation of Sleep. Curr Biol 26 (7):972-980. doi:10.1016/j.cub.2016.02.013
26. Kim DH, Han MR, Lee G, Lee SS, Kim YJ, Adams ME (2015) Rescheduling Behavioral
Subunits of a Fixed Action Pattern by Genetic Manipulation of Peptidergic Signaling. PLoS
Genet 11 (9):e1005513. doi:10.1371/journal.pgen.1005513
27. Kim YJ, Zitnan D, Galizia CG, Cho KH, Adams ME (2006) A command chemical triggers an
innate behavior by sequential activation of multiple peptidergic ensembles. Curr Biol 16
(14):1395-1407. doi:10.1016/j.cub.2006.06.027
28. Luo J, Liu Y, Nässel DR (2013) Insulin/IGF-regulated size scaling of neuroendocrine cells
expressing the bHLH transcription factor Dimmed in *Drosophila*. PLoS Genet 9
(12):e1004052. doi:10.1371/journal.pgen.1004052
29. Okusawa S, Kohsaka H, Nose A (2014) Serotonin and downstream leucokinin neurons
modulate larval turning behavior in *Drosophila*. J Neurosci 34 (7):2544-2558.
doi:10.1523/JNEUROSCI.3500-13.2014
30. Sellami A, Veenstra JA (2015) SIFamide acts on fruitless neurons to modulate sexual
behavior in *Drosophila melanogaster*. Peptides 74:50-56.
doi:10.1016/j.peptides.2015.10.003
31. Veenstra JA, Khammassi H (2017) Rudimentary expression of RYamide in *Drosophila*
melanogaster relative to other *Drosophila* species points to a functional decline of this
neuropeptide gene. Insect Biochem Mol Biol 83:68-79. doi:10.1016/j.ibmb.2017.03.001
32. Cabrero P, Radford JC, Broderick KE, Costes L, Veenstra JA, Spana EP, Davies SA, Dow
JA (2002) The *Dh* gene of *Drosophila melanogaster* encodes a diuretic peptide that acts
through cyclic AMP. J Exp Biol 205 (Pt 24):3799-3807
33. Terhzaz S, O'Connell FC, Pollock VP, Kean L, Davies SA, Veenstra JA, Dow JA (1999)
Isolation and characterization of a leucokinin-like peptide of *Drosophila melanogaster*. J
Exp Biol 202 (Pt 24):3667-3676
34. Lee KM, Daubnerova I, Isaac RE, Zhang C, Choi S, Chung J, Kim YJ (2015) A neuronal
pathway that controls sperm ejection and storage in female *Drosophila*. Curr Biol 25
(6):790-797. doi:10.1016/j.cub.2015.01.050
35. Cavanaugh DJ, Geratowski JD, Wooltorton JR, Spaethling JM, Hector CE, Zheng X,
Johnson EC, Eberwine JH, Sehgal A (2014) Identification of a circadian output circuit for
rest:activity rhythms in *Drosophila*. Cell 157 (3):689-701. doi:10.1016/j.cell.2014.02.024
36. Brand AH, Perrimon N (1993) Targeted gene expression as a means of altering cell fates
and generating dominant phenotypes. Development 118 (2):401-415

37. Asahina K, Watanabe K, Duistermars BJ, Hoopfer E, Gonzalez CR, Eyjolfsdottir EA, Perona P, Anderson DJ (2014) Tachykinin-expressing neurons control male-specific aggressive arousal in *Drosophila*. Cell 156 (1-2):221-235. doi:10.1016/j.cell.2013.11.045

38. Jenett A, Rubin GM, Ngo TT, Shepherd D, Murphy C, Dionne H, Pfeiffer BD, Cavallaro A, Hall D, Jeter J, Iyer N, Fetter D, Hausenfluck JH, Peng H, Trautman ET, Svirskas RR, Myers EW, Iwinski ZR, Aso Y, DePasquale GM, Enos A, Hulamm P, Lam SC, Li HH, Laverty TR, Long F, Qu L, Murphy SD, Rokicki K, Safford T, Shaw K, Simpson JH, Sowell A, Tae S, Yu Y, Zugates CT (2012) A GAL4-driver line resource for *Drosophila* neurobiology. Cell Rep 2 (4):991-1001. doi:10.1016/j.celrep.2012.09.011

39. Dietzl G, Chen D, Schnorrer F, Su KC, Barinova Y, Fellner M, Gasser B, Kinsey K, Oppel S, Scheiblauer S, Couto A, Marra V, Keleman K, Dickson BJ (2007) A genome-wide transgenic RNAi library for conditional gene inactivation in *Drosophila*. Nature 448 (7150):151-156. doi:10.1038/nature05954

40. Pfeiffer BD, Truman JW, Rubin GM (2012) Using translational enhancers to increase transgene expression in *Drosophila*. Proc Natl Acad Sci U S A 109 (17):6626-6631. doi:10.1073/pnas.1204520109

41. Kubrak OI, Lushchak OV, Zandawala M, Nässel DR (2016) Systemic corazonin signalling modulates stress responses and metabolism in *Drosophila*. Open Biol 6 (11). doi:10.1098/rsob.160152

42. Nässel DR, Cantera R, Karlsson A (1992) Neurons in the cockroach nervous system reacting with antisera to the neuropeptide leucokinin I. J Comp Neurol 322 (1):45-67. doi:10.1002/cne.903220105

43. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C, Saalfeld S, Schmid B, Tinevez JY, White DJ, Hartenstein V, Eliceiri K, Tomancak P, Cardona A (2012) Fiji: an open-source platform for biological-image analysis. Nat Methods 9 (7):676-682. doi:10.1038/nmeth.2019

44. Kahsai L, Kapan N, Dircksen H, Winther AM, Nässel DR (2010) Metabolic stress responses in *Drosophila* are modulated by brain neurosecretory cells that produce multiple neuropeptides. PLoS One 5 (7):e11480. doi:10.1371/journal.pone.0011480

45. Dow JA, Maddrell SH, Gortz A, Skaer NJ, Brogan S, Kaiser K (1994) The Malpighian tubules of *Drosophila melanogaster*: a novel phenotype for studies of fluid secretion and its control. J Exp Biol 197:421-428

46. Alvarez-Rivero J, Moris-Sanz M, Estacio-Gomez A, Montoliu-Nerin M, Diaz-Benjumea FJ, Herrero P (2017) Variability in the number of abdominal leucokinergic neurons in adult *Drosophila melanogaster*. J Comp Neurol 525 (3):639-660. doi:10.1002/cne.24093

47. Dircksen H, Tesfai LK, Albus C, Nässel DR (2008) Ion transport peptide splice forms in central and peripheral neurons throughout postembryogenesis of *Drosophila melanogaster*. J Comp Neurol 509 (1):23-41. doi:10.1002/cne.21715

48. Cognigni P, Bailey AP, Miguel-Aliaga I (2011) Enteric neurons and systemic signals couple nutritional and reproductive status with intestinal homeostasis. Cell Metab 13 (1):92-104. doi:10.1016/j.cmet.2010.12.010

49. Chintapalli VR, Wang J, Dow JA (2007) Using FlyAtlas to identify better *Drosophila melanogaster* models of human disease. Nat Genet 39 (6):715-720. doi:10.1038/ng2049

50. Bhatt G, da Silva R, Nachman RJ, Orchard I (2014) The molecular characterization of the kinin transcript and the physiological effects of kinins in the blood-gorging insect, *Rhodnius prolixus*. Peptides 53:148-158. doi:10.1016/j.peptides.2013.04.009

51. Chen Y, Veenstra JA, Hagedorn H, Davis NT (1994) Leucokinin and diuretic hormone immunoreactivity of neurons in the tobacco hornworm, *Manduca sexta*, and co-localization of this immunoreactivity in lateral neurosecretory cells of abdominal ganglia. Cell Tissue Res 278 (3):493-507

52. Thompson KS, Rayne RC, Gibbon CR, May ST, Patel M, Coast GM, Bacon JP (1995) Cellular colocalization of diuretic peptides in locusts: a potent control mechanism. Peptides 16 (1):95-104

53. Te Brugge VA, Nässel DR, Coast GM, Schooley DA, Orchard I (2001) The distribution of a
kinin-like peptide and its co-localization with a CRF-like peptide in the blood-feeding bug,
Rhodnius prolixus. Peptides 22 (2):161-173
54. Terhzaz S, Teets NM, Cabrero P, Henderson L, Ritchie MG, Nachman RJ, Dow JA,
Denlinger DL, Davies SA (2015) Insect capa neuropeptides impact desiccation and cold
tolerance. Proc Natl Acad Sci U S A 112 (9):2882-2887. doi:10.1073/pnas.1501518112

Figure captions

Figure 1: The *Lk-GAL4* drives GFP expression in four distinct neuronal populations in the adult *Drosophila* CNS. (A) One pair of neurons in the lateral horn (lateral horn LK neurons; LHLKs) and another pair of neurons in the subesophageal ganglion (subesophageal ganglion LK neurons; SELKs) express LK in adult brain of *Drosophila. Lk-GAL4* also drives weak and variable expression in four pairs of neurons in the brain (approximate location of these cells is indicated by the white box; see figure S2 for an alternate preparation where these cells are weakly stained). **(B)** These four pairs of neurons do not display any LK-immunoreactivity but are positive for ITP-immunoreactivity [20]. **(C)** In the ventral nerve cord (VNC), LK is expressed in eleven pairs of neurons (abdominal LK neurons; ABLKs). Seven pairs of smaller neurons in the posterior region (p) persist from the larval stages and the other four pairs (the number of pairs can vary between individuals) of larger neurons in the anterior region (a) are adult-specific [46]. **(D)** A schematic depiction of LK-expressing neurons in the adult brain and VNC of *Drosophila.* T1 – T3, thoracic neuromeres. **(E)** *Lk-GAL4* also drives ectopic expression in the salivary glands of adult *Drosophila.* **Note:** *JFRC29-10xUAS-IVS-myr::GFP-p10* was utilized in **A, C** and **E** whereas *UAS-mcd8-GFP* was utilized in **B**.

Figure 2: DH44 expression in the adult *Drosophila* CNS. (A) DH44 is expressed in three pairs of median neurosecretory cells (MNCs) in pars intercerebralis of adult *Drosophila.* Antiserum to *Drosophila* DH44 labels the same neurons identified by *DH44-GAL4*-driven GFP. **(B)** In the anterior region of the ventral nerve cord (VNC), DH44 is expressed in two pairs of neurons. **(C)** In the posterior region of the VNC, DH44 is expressed in at least four pairs of neurons. In both B and C, there are some neurons that display DH44-immunoreactivity but do not express GFP.

Figure 3: LK and DH44 are coexpressed in the ventral nerve cord, but not in the brain of adult *Drosophila*. (A) *DH44-GAL4* driven GFP is not colocalized with LK-immunoreactivity in the adult brain. **(B)** *DH44-GAL4* driven GFP is colocalized with LK-immunoreactivity in a subset of the abdominal LK neurons (ABLKs) in the ventral nerve cord (VNC) **(C)** *Lk-GAL4* driven GFP is not colocalized with DH44-immunoreactivity in the adult brain. **(D)** *Lk-GAL4* driven GFP is colocalized with DH44-immunoreactivity in a subset of ABLKs in the adult VNC.

Figure 4: Schematics of LK- and DH44-expressing neurons in the larval and adult CNS of *Drosophila*. (A) A schematic of the larval CNS showing the location of neurons expressing LK, DH44 or both LK and DH44. **(B)** A schematic of the adult CNS showing the location of neurons expressing LK, DH44 or both LK and DH44. LHLK, lateral horn LK neuron; SELK, subesophageal ganglion LK neuron; ABLK, abdominal LK neuron, T1 – T3, thoracic neuromeres.

Figure 5: *Lk*- and *DH44-RNAi* knockdown efficiency was tested using immunolabelling. (A, B) Knock down of *Lk* with *Lk-GAL4* driven *Lk-RNAi* causes a significant decrease in LK-immunoreactivity in the adult brain and ventral nerve cord (VNC). **(C)** Fluorescence intensity measurement of lateral horn LK neurons shows a significant decrease in LK-immunoreactivity in *Lk* knock down flies compared to control flies. **(D)** *DH44-GAL4* driven *Lk-RNAi* causes a significant decrease in LK-immunoreactivity in the adult VNC as determined by the number of immunoreactive neurons (the average number of neurons is indicated in each panel; see figure S6) that could be detected. Whereas, *DH44-GAL4* driven *DH44-RNAi* causes a significant increase in LK-immunoreactivity in adult ABLKs **(E)** and a complete abolishment of DH44-immunoreactivity in the adult brain **(F)** and VNC **(G)**.

Figure 6: Knockdown of *Lk* using *Lk-GAL4* impacts stress resistance and water content of *Drosophila*. *Lk-GAL4* driven *Lk* knock down results in a significant increase in survival compared to control flies under **(A)** desiccation, **(B)** starvation and **(C)** ionic stress (artificial food supplemented with 4% NaCl). Data are presented in survival curves and the error bars represent standard error (**** $p < 0.001$, as assessed by Log-

rank (Mantel-Cox) test) **(D)** *Lk* knock down has no impact on chill coma recovery. **(E)** There is no significant difference (One-way ANOVA) in feeding as measured by capillary feeding (CAFE) assay between *Lk* knock down and control flies. Results are presented as cumulative food intake over four days. **(F)** Flies with *Lk* knock down have a higher wet weight and dry weight and retain more water (wet weight minus dry weight) compared to control flies. (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.005$, as assessed by One-way ANOVA). Legend for B-F is the same as the one in A.

Figure 7: Knockdown of *DH44* using *Lk-GAL4* impacts stress resistance and feeding in *Drosophila*. *Lk-GAL4* driven *DH44* knock down results in a significant increase in survival compared to control flies under **(A)** desiccation, **(B)** starvation and **(C)** ionic stress (artificial food supplemented with 4% NaCl). Data are presented in survival curves and the error bars represent standard error (**** $p < 0.001$, as assessed by Log-rank (Mantel-Cox) test) **(D)** *DH44* knock down causes a small delay in chill coma recovery. (* $p < 0.05$, as assessed by Log-rank (Mantel-Cox) test) **(E)** Flies with *DH44* knockdown feed less compared to control flies in capillary feeding (CAFE) assay. Results are presented as cumulative food intake over four days. (*** $p < 0.005$, **** $p < 0.001$, as assessed by One-way ANOVA). **(F)** There is no significant difference in wet weight, dry weight and water content of *DH44*-knockdown and control flies. Legend for B-F is the same as the one in A.

Figure 8: Knockdown of *Lk* using *DH44-GAL4* impacts stress resistance and water content of *Drosophila*. *DH44-GAL4* driven *Lk* knock down results in a significant increase in survival compared to control flies under **(A)** desiccation, **(B)** starvation and **(C)** ionic stress (artificial food supplemented with 4% NaCl). Data are presented in survival curves and the error bars represent standard error (**** $p < 0.001$, as assessed by Log-rank (Mantel-Cox) test) **(D)** *Lk* knock down results in a delayed recovery from chill coma. (* $p < 0.05$, as assessed by Log-rank (Mantel-Cox) test) **(E)** There is no significant difference (One-way ANOVA) in feeding as measured by capillary feeding (CAFE) assay between *Lk* knock down and control flies. Results are presented as cumulative food intake over four days. **(F)** Flies with *Lk* knock down in ABLKs have a higher wet weight, dry weight and retain more water (wet weight minus dry weight)

compared to control flies. (** $p < 0.01$, *** $p < 0.005$, **** $p < 0.001$, as assessed by

One-way ANOVA). Legend for B-F is the same as the one in A.

Figure 9: LK and DH44 peptide application results in an additive response on fluid

secretion by Malpighian tubules (MTs) *ex vivo*. (A) Secretion rates of MTs incubated

with 10^{-7} M DH44 ($n = 28$), 10^{-10} M LK ($n = 25$), a combination of both 10^{-7} M DH44 and

10^{-10} M LK ($n = 23$), or no treatment/basal ($n = 14$). **(B)** Secretion rates of MTs

incubated with 10^{-9} M DH44 ($n = 14$), 10^{-10} M LK ($n = 25$), a combination of both 10^{-9} M

DH44 and 10^{-10} M LK ($n = 31$), or no treatment/basal ($n = 13$). For both A and B,

secretion rates were measured at 10 min intervals for 30 min before and after the

addition of peptide (indicated with an arrow). **(C, D)** Change (%) in secretion determined

by comparing the secretion rate over the first 30 min to the maximum secretion rate

following peptide application. The legend and sample size for C and D are the same as

the one in A and B, respectively. **(E, F)** Total fluid secreted for 30 min following peptide

application or no treatment (basal). Note that the amount of total fluid secreted following

the addition of both LK (10^{-10} M) and DH44 (10^{-7} M) is a sum of the total fluid secreted

following the addition of each of those peptides separately. Asterisk indicates

significantly different secretion rate compared to basal secretion rate (secretion rate

prior to the addition of peptide). (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$;

Mann-Whitney U test)

Figure S1: LK and DH44 expression in the larval *Drosophila* CNS. (A) *Lk-GAL4* drives expression in five pairs of neurons in the brain; however, four of these pairs do not display any LK-immunoreactivity [20]. **(B)** These four pairs of neurons display ITP-immunoreactivity. **(C)** Three pairs of neurons in the subesophageal ganglion express *Lk* in larval *Drosophila*. **(D)** Seven pairs of neurons in the larval ventral nerve cord (VNC) express *Lk*. **(E)** A schematic of LK-expressing neurons in the larval brain and VNC of *Drosophila.* Neurons displaying LK-immunoreactivity are labeled in red and neurons displaying ITP-immunoreactivity are labeled in black. **(F)** *DH44-GAL4* driven GFP and DH44-immunoreactivity is present in three pairs of median neurosecretory cells in the larval brain. **(G)** DH44 is expressed in several neurons, with strong expression seen in seven pairs of neurosecretory cells in the larval VNC. In both E and F, there are some neurons that contain GFP but do not contain DH44-immunoreactivity.

Figure S2: LK expression in adult *Drosophila* brain. *Lk-GAL4* drives weak GFP expression in four pairs on neurons in the adult brain. The location of these cells is indicated by white boxes.

Figure S3: LK and DH44 are coexpressed in neurons of the ventral nerve cord but not in the brain of larval *Drosophila.* (A) *DH44-GAL4* driven GFP is not colocalized with LK-immunoreactivity in the larval brain. **(B)** *DH44-GAL4* driven GFP is colocalized with LK-immunoreactivity in all seven pairs of abdominal LK neurons (ABLKs) in the ventral nerve cord (VNC) **(C)** *Lk-GAL4* driven GFP is not colocalized with DH44-immunoreactivity in the larval brain. **(D)** *Lk-GAL4* driven GFP is colocalized with DH44-immunoreactivity in ABLKs in the larval VNC.

Figure S4: Knockdown of *DH44* using *Actin5c-GAL4*. *Actin5c-GAL4* driven *DH44* knockdown results in a complete abolishment of DH44-immunoreactivity in the six neurons in pars intercerebralis of adult *Drosophila*.

Figure S5: Number of LK-immunoreactive neurons in the VNC following knockdown of *DH44* and *Lk* using *DH44-GAL4*. *Lk* knockdown but not *DH44* knockdown causes a significant decrease in the number of LK-immunoreactive neurons that could be detected in the adult VNC. (**** $p < 0.001$, as assessed by One-way ANOVA).

Figure S6: Knockdown of *DH44* using *DH44-GAL4* impacts stress resistance and feeding in *Drosophila*. *DH44-GAL4* driven *DH44* knock down results in a significant increase in survival compared to control flies under **(A)** desiccation (compared to the GAL4 control), **(B)** starvation and **(C)** ionic stress (artificial food supplemented with 4% NaCl). Data are presented in survival curves and the error bars represent standard error (**** $p < 0.001$, as assessed by Log-rank (Mantel-Cox) test) **(D)** *Lk* knock down results in a delayed recovery from chill coma. (* $p < 0.001$, as assessed by Log-rank (Mantel-Cox) test) **(E)** There is a significant decrease (One-way ANOVA) in feeding as measured by capillary feeding (CAFE) assay in *DH44* knock down flies (compared to the GAL4 control). Results are presented as cumulative food intake over four days. **(F)** There is no significant difference in wet weight, dry weight and water content of *DH44-* knockdown and control flies. Legend for B-F is the same as the one in A.

Figure S7: Knockdown of *Lk* in ABLKs with *DH44-GAL4* does not influence LK-stimulated Malpighian tubule secretion *ex vivo*. (A) Secretion rates of 10^{-9} M LK stimulated MTs isolated from *DH44 > w^{1118}* ($n = 14$) or *DH44 > Lk RNAi* flies ($n = 25$). **(B)** Secretion rates of 10^{-10} M LK stimulated MTs isolated from *DH44 > w^{1118}* ($n = 10$) or *DH44 > Lk RNAi* flies ($n = 12$). For both A and B, secretion rates were measured at 10 min intervals for 30 min before and after the addition of peptide (indicated with an arrow). **(C, D)** Change (%) in secretion determined by comparing the secretion rate over the first 30 min to the maximum secretion rate following peptide application. The legend and sample size for C and D are the same as the one in A and B, respectively. Asterisk indicates significantly different secretion rate compared to basal secretion rate (secretion rate prior to the addition of peptide). (NS = not significant, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$; Mann-Whitney U test)

A

LHLK

SELK

50 μM

C

a

p

25 μM

D

LHLK

SELK

● anti-LK positive
● anti-LK negative

T1

T2

T3

ABLKs

E

Salivary gland

20 μM

B

Lk > mcd8 GFP

25 μM

Merge

Anti-ITP

	DH44 > mcd8 GFP	Merge	Anti-DH44
Brain	A		
VNC (anterior)	B		
VNC (posterior)	C		

Scale bars: A = 50 μM; B = 25 μM; C = 25 μM

DH44 > mcd8 GFP	Merge	Anti-LK

Brain — A — 50 μM

VNC — B — 25 μM

Lk > mcd8 GFP	Merge	Anti-DH44

Brain — C — 50 μM

VNC — D — 25 μM