Quantifying movement behavior, the first step of sexual selection, in the lizard *Anolis sagrei*.

Ambika Kamath[1,2,3] and Jonathan B. Losos[1]

1. Department of Organismic and Evolutionary Biology and Museum of Comparative Zoology, Harvard University, 26 Oxford Street, Cambridge MA 02138
2. *Present Address*: Department of Ecology, Evolution and Marine Biology, University of California Santa Barbara, Santa Barbara, CA 93106
3. Author for correspondence. Email: ambikamath@gmail.com

Abstract

How individuals move through their environment dictates which other individuals they encounter, determining their social and reproductive interactions and the extent to which they experience sexual selection. Specifically, females rarely have the option of mating with all males in a population—they can only choose among the males they encounter. Further, quantifying phenotypic differences between the males that females encounter and those that sire females' offspring lends insight into how social and reproductive interactions shape male phenotypes. We used an explicitly spatiotemporal Markov chain model to estimate the number of potential mates of *Anolis sagrei* lizards from their movement behavior, and used genetic paternity assignments to quantify sexual selection on males. Females frequently encountered and mated with multiple males, offering ample opportunity for female mate choice. Sexual selection favored males that were bigger and moved over larger areas, though the effect of body size cannot be disentangled from last-male precedence. Our approach corroborates some patterns of sexual selection previously hypothesized in anoles based on describing them as territorial, whereas other results, including female multiple mating itself, are at odds with territorial polygyny, offering insight into discrepancies in other taxa between behavioral and genetic descriptions of mating systems.

Keywords: mating system, territory, territoriality, polygyny, anole, encounter rates

Introduction

Sexual selection is a layered process, with animals sequentially having to overcome

intrasexual competition, intersexual mating preferences, and, for males, post-copulatory competition

and choice before achieving reproductive success [1, 2]. Decades of research have spawned a vast

literature on each of these aspects of sexual selection. However, a crucial but little-appreciated

component of measuring sexual selection in natural populations involves quantifying the very first

step of the mating sequence, namely encountering potential mates. How individuals move across

space through time directly influences the number and phenotypic distribution of potential mates

they encounter [2–4]. Moreover, by bringing about encounters between potential mates as well as

between potential competitors, individuals' movement patterns set the stage for subsequent sexual

selection through male-male competition and female choice [5, 6]. Documenting animals' movement

patterns, to understand how often and which members of the opposite sex are encountered by

individuals, is thus fundamental to discovering the extent to which sexual selection can act in the

wild. Concurrently, quantifying phenotypic differences between potential mates (individuals

encountered) and actual mates (individuals whose offspring are borne), yields insight into the nature

of selection imposed by social and reproductive interactions.

In particular, individuals' movement patterns determine the potential for female mate choice

to drive sexual selection [7, 8]. Female mate choice has been studied extensively, yielding vigorous

debate surrounding the precise mechanisms by which it arises, acts, and is maintained across a range

of taxa (reviewed in [9–11]). Common to all models of female choice, however, is the idea that

females can choose among males. But whether and to what extent individual females in fact have

such a choice, and therefore the extent to which female choice can drive sexual selection, depends in

large part on how many males they encounter [3, 12, 13].

In studies of sexual selection, examinations of movement behavior are often restricted to

considering how females sample among males in taxa where female mate choice is already

acknowledged to be important. For example, searching behavior is often thought to be pertinent to

sexual selection in species where females must visit and sample from among males in leks or at fixed

display sites (e.g. [8, 14]). However, similar measurements of movement behavior, and of encounters

between potential mates, are equally relevant to understanding the opportunity for sexual selection

in other animal species as well, including those where female choice is not usually considered a

major selective pressure [13, 15].

Unexpected opportunities for female choice are often uncovered in species in which earlier

behavioral descriptions of mating systems, based on movement patterns as well as social

interactions, are found to be inconsistent with more recent genetic descriptions of mating patterns

[16]. For example, most birds were widely regarded as monogamous prior to the advent of genetic

tools that revealed frequent extrapair copulation [17]. Occasionally, these inconsistencies have

prompted researchers to reexamine animals' movement patterns to reconcile behavioral and genetic

descriptions of mating patterns (e.g. [18, 19]). For example, tracking the movement behavior of red

deer revealed that females move long distances between harems unexpectedly often, demonstrating

the possibility of female choice in a system where sexual selection was thought to be dominated by

male-male competition [20]. In general, though, discrepancies between behavioral and genetic

descriptions of mating systems remain common—consider how often species are described as

"socially monogamous," for example, but "genetically promiscuous." These discrepancies imply that

we remain without complete or correct descriptions of movement and social behaviors that account

for polygynandry in many species, and are thus unable to fully understand how sexual selection has

shaped and is shaped by these behaviors.

In this paper, we develop an explicitly spatiotemporal approach to estimate encounters

between potential mates from observations of the movement behavior of male and female *Anolis*

sagrei lizards. Our first goal is to investigate if females encounter multiple males, which could offer

females the possibility of mate choice. This possibility has previously been considered unlikely in

most anoles, which have widely been described as territorial and polygynous despite genetic data

revealing widespread multiple mating (reviewed in [21]). Our second goal is to characterize sexual

selection in this population by examining the predictors of male reproductive success at two levels.

First we ask if the number of potential mates encountered by males is associated with their

phenotype (the spatial extent of their movement and body size). Second, we test three hypotheses to

understand the phenotypic differences between potential mates (males encountered by females) and

actual mates (males that sire females' offspring). We first examine if females bear offspring sired by

the males they encounter more often [22]. Then, we ask if males encountered later in the breeding

season are more likely to sire offspring than males encountered earlier ("last-male precedence"; [23]).

Finally, given widespread sexual selection in animals for larger males [24] as well as pronounced

male-biased sexual size dimorphism in *A. sagrei*, we ask if females disproportionately bear offspring

sired by larger males. We find evidence that females both encounter and mate with multiple males,

and that sexual selection acts on movement behavior, male-female interactions, and body size in

male *A. sagrei*, suggesting the possibility for complex polygynandrous reproductive dynamics

including both pre- and post-copulatory female mate choice. We discuss how some of our results are

consistent with previous descriptions of *A. sagrei* as territorial while others are unlikely to have been

considered from within the framework of territoriality. Finally, we consider how paying attention to

movement behavior can allow for the reconciliation of diverse types of behavioral and genetic

studies of animal mating systems in many taxa.

Methods

Field sampling and egg collection

Anolis sagrei is a low-perching arboreal lizard native to Cuba and the Bahamas that has been

established in Florida for nearly a century [25, 26]. It co-occurs with the native, higher-perching

Anolis carolinensis throughout most of its range in the US, including in our sampled population;

however, few interactions between the two species were observed.

Anolis sagrei lizards were captured, marked, and monitored to estimate their movement

patterns in the University Gardens on the University of Florida campus in Gainesville, FL, from

March 4, 2015 to May 25, 2015 between 0900 and 1800 hours. The start date of sampling coincided

approximately with the increased activity of these lizards after the winter and the beginning of the

breeding season; we concluded sampling at about the time at which female *A. sagrei* began laying

eggs (based on our observations in 2014 of hatchlings appearing at the end of June, after an

approximately month long incubation period [27]). We caught almost every lizard within a 7140 m^2

area and marked each captured individual with a unique bead tag [28], which allowed us to

subsequently observe and identify individuals from a distance without disturbing them (in total, 4%

of observations were of unmarked individuals). When captured, we measured each individual's

snout-vent length (SVL) as a measure of body size, and removed ~2-3 cm of tail tissue for genetic

analysis. At each subsequent observation of a lizard, we noted its identity as well as the time and

location of the observation. We avoided observing the same individual more frequently than once

106 per hour, allowing ample time for lizards to resume normal behavior if disturbed by us.

107 Approximately once a month, we recaptured and re-measured males to estimate the average growth

108 rate of males in this population.

109  We measured distances between the locations at which lizards were observed to triangulate

110 their relative positions. In areas of continuous vegetation, locations more than 1m apart were

111 considered distinct. We also mapped the location of all trees within the site at which lizards were not

112 observed, to include all trees to which a lizard could potentially have moved in our estimations of

113 their movement patterns (Figure S1).

114  At the end of the observation period (which, based on 2014 observations, was approximately

115 when females began laying eggs), we captured 36 marked females and brought them into captivity.

116 Females were housed singly under established anole husbandry conditions [29] until mid-November.

117 Each cage contained a pot of soil in which the resident female laid eggs fertilized by sperm stored

118 from her copulations in the field. Eggs were incubated for two to ten days, after which embryos

119 were dissected from them for genetic analysis.

120  *Movement pattern analysis*

121  Analyses were carried out in R v. 3.3.2 [30]. We used a discrete-time Markov chain to model

122 lizards' movements between the mapped locations in the site. We divided daytime hours (0800 to

123 2000 hours; anoles are diurnal, so we assumed that the lizards did not move at night) over the

124 sampling period (83 days) into 996 hour-long blocks. Observations were assigned to the bin closest

125 to the time of the observation. Transition probabilities (P_{ij}) between locations i and j were modelled

126 as exponentially declining with the distance between the locations (d_{ij}), with rows of the transition

127 matrix then normalized to sum to one (N is the total number of locations):

128     $$P_{ij} = \frac{e^{\lambda d ij}}{\sum_{j=1}^{N} e^{\lambda d ij}}$$     (Equation 1)

129  In other words, lizards were modelled as less likely to move to distant locations than to close

130 locations, with a total probability of 1 of moving from each location to *some* location within the site,

131 including staying at the same location. The value of the exponential decline parameter λ was

132 estimated by maximizing the likelihood of the observed data (including only pairs of consecutive

133 observations of lizards, i.e. no assumptions were made while fitting the model regarding mortality or

emigration after the last observation of a lizard) using the *bbmle* package [31]; separate models were

fit for males and females.

The probability that a lizard was at a particular location at a particular hour was inferred

using the Markov chain model described above. Methodological details can be found in the

Supplementary Information, but briefly, this probability depended not only on where that lizard was

seen previously but also where it was seen next. We thus calculated, for each lizard, a matrix of

probabilities that it occupied a particular location at a particular hour, for all locations and hours.

Then, we performed element-wise multiplication of pairs of these matrices to calculate the

probability of co-occurrence at each one hour-long time bin, for every possible pair of lizards.

Encounters were categorized for each pair at each hour ("yes/no") from these probabilities by

setting cutoffs based on the co-occurrence probabilities calculated for pairs of lizards observed in

the same location within an hour of one another. For each hour, we inferred an encounter between

a pair of lizards if their co-occurrence probability was larger than the cutoff.

To quantify potential mating opportunities for each individual, we calculated the number of

females encountered by each male and the number of males encountered by each female, as

estimated by our model. The proportion of females that encounter multiple males and the mean

number of males encountered by females reveal the extent to which multiple mating by females is

possible in this population.

We quantified the spatial extent of an individual's movement by calculating the mean of the

distances from each observation of the individual to the centroid of all observations of the

individual (mean distance from centroid). Lower mean distance from the centroid indicates smaller

spatial extent. We jittered points randomly within a 0.5 m radius along both the X and Y axis of our

site before calculating mean distance from the centroid, to account for the 1 m resolution at which

locations were mapped (Figure S1).

We estimated a growth curve for males by fitting a logistic equation to males' SVL measured

initially and at recaptures, using nonlinear least squares regression [32] (see Supplementary

Information). We used this logistic growth curve to estimate the SVL of each male on the day of

each of his inferred encounters, based on his SVL at the nearest capture, to test for sexual selection

on male body size and for male avoidance of size-matched males (see below).

Parentage analysis

DNA was extracted from the 36 females housed in captivity, all 161 sampled males, and 383

offspring using an AutogenPrep 965. Six microsatellite regions were amplified for these individuals

(see Table S1 for primer and amplification protocol details; [33, 34]), with fragment analysis

performed at the DNA Analysis Facility at Yale University. Alleles were scored manually after

examining chromatogram peaks in Geneious v10.0.9 [35].

Parentage analyses were performed in CERVUS v3.0.7 [36]. High proportions of null alleles

were estimated at three loci (Table S1); following [37, 38], we retained these loci in the analysis but

typed apparent homozygotes at only one allele, with the other allele coded as missing. All offspring

had known mothers, and we provided as candidate sires for each offspring the list of males that its

mother was estimated to have encountered, as described above. In the simulation to determine

LOD score cutoffs for assignment, we provided a genotyping error rate of 0.01 (based on mother-

offspring mismatch across all loci); the proportion of loci typed was 0.81. Simulations were run with

23 candidate sires and the proportion of sires typed set to 0.75, based on the maximum number of

males encountered by any female in the population (17 males). These parameters were chosen after

running preliminary analyses with simulations in which the proportion of sires typed was set to 0.25,

0.5, or 0.75 and the number of candidate sires set correspondingly to 68, 34, or 23; we chose the

parameter combination with the closest match between the proportion of sires typed and the

observed assignment rate [39]. We performed further analyses to test the effects of restricting the

number of candidate sires relative to the whole population, and of providing all sampled males as

candidate sires for all offspring; we found that simply restricting the number of sires did not inflate

paternity assignments and that results downstream analyses were unaffected by excluding the 18

offspring whose paternity assignments differed in the analysis where all males were provided as

candidate fathers (see Supplementary Information).

Hypothesis Testing

The number of potential mates encountered and spatial extent (mean distance to the

centroid) had right-skewed distributions, and were therefore log-transformed before parametric

analyses; SVL was analyzed untransformed. We compared spatial extent between males and females

using a t-test weighted by the number of observations per individual, and investigated if variation in

males' spatial extent was related to body size, using a linear regression of SVL at first capture vs.

mean distance to the centroid, weighted by the number of observations per individual.

Next, we examined if the number of females encountered by males varied with the spatial

extent of males' movement (mean distance from the centroid) and with mean male body size at their

encounters with females, using a multiple linear regression weighted by the number of observations

per individual.

To assess if males avoided size-matched males, we examined the differences in estimated

SVL between pairs of males estimated to encounter one another, and compared these differences to

the differences in estimated SVL between randomly chosen pairs of males. For males that were

estimated to encounter one another, we used the logistic growth curve to estimate their SVL on the

day of the encounter. We initially sampled five random pairs per pair of males estimated to

encounter one another, estimated their SVLs on the same day as the corresponding encounter, and

eliminated random pairs in which either individual had an estimated SVL less than the minimum

observed SVL. We then repeated this random sampling a total of 3000 times to assess if the number

of male-male encounters among size-matched males (estimated SVL difference of 0-2 mm) was

significantly lower than expected by chance.

We used a resampling approach to examine (1) if males who sired individual females'

offspring encountered the mother significantly more often than males who encountered the same

females but did not sire offspring (encounter rate hypothesis), (2) if males who sired individual

females' offspring encountered those females later than males who encountered the same females

but did not sire offspring (last-male precedence hypothesis) and (3) males who sired individual

females' offspring were bigger than males who encountered the same females but did not sire

offspring (body size hypothesis).

We first calculated the difference between means of the number of encounters between

male-female pairs for sires and non-sires across all offspring. We also calculated, for each male-

female pair, the last hour at which the pair encountered one another and maximum SVL estimated

for the male across all encounters between the pair as an estimate of male body size, and then

calculated the difference between mean hour of last encounter and mean body size between sires

and non-sires. We then recalculated these differences between means after randomly assigning each

offspring a sire from the set of males that encountered the mother of that offspring. Random sire

assignments were performed in two ways. To address hypothesis (1), we sampled uniformly from

the list of males encountered by each mother. Then, to address hypotheses (2) and (3), we sampled

in proportion to the rate at which each male encountered each mother. The former allowed us to

test if sires encountered mothers more often than did non-sires, and the latter allowed us to test if

later-encountered males and bigger males sired offspring more often than earlier-encountered males

and smaller males, after accounting for variation across males in encounter rates. Each resampling

was repeated 10000 times.

Results

A total of 253 individuals (161 males, 92 females) were caught and marked during the

sampling period, and were observed a total of 5629 times. The number of observations per

individual ranged from one to 128; the median number of observations per individual for males and

females was 11 and 15 respectively. An example of two individuals' locations through time is shown

in Figure 1.

Figure 1. Examples of the locations through time for two individuals, U12 and U26 (left), and the probability of their co-occurrence as estimated by the Markov chain model (above). The dashed circles have a diameter of 10 m, which matches previous estimates of the territory size of male *Anolis sagrei*.

We used a Markov chain to model lizards' movements between locations in the site, where

transition probabilities were modelled as exponentially declining with the distance between locations.

We estimated λ values of -0.78 and -1.27 for males and females respectively (see Equation 1),

indicating that males were more likely than females to move longer distances. Using this Markov

chain model to estimate individuals' movement patterns and thereafter the probabilities of their co-

occurrence (see Figure 1 for an example), we calculated that females encountered 5.1 ± 3.7 males

(mean ± S.D.) and males encountered 2.9 ± 3.0 females; 78% of females and 60% of males

encountered multiple individuals of the opposite sex (Figure 2). Males encountered an average of 4.5

± 3.6 other males.

Figure 2. Histograms of the number of males encountered by females (left) and females encountered by males (right).

The mean distance from the centroid of all of an individual's locations ranged from 0.2 m to

41.3 m for males and from 0.2 m to 20.8 m for females (mean ± standard deviation for males vs.

females: 6.8 ± 7.0 m vs. 2.7 ± 3.3 m; $t = 8.1$, $df = 208.7$, $P < 0.001$). This measure of spatial extent

was weakly associated with SVL at initial capture for males ($r^2 = 0.04$, $F_{1, 135} = 4.91$, $P = 0.03$).

To estimate male body size at their encounters with females using a logistic growth curve, we

recaptured 68 males and re-measured their SVLs a total of 94 times, with 32 ± 15 (mean ± SD) days

elapsed between measurements. The mean difference in estimated SVL between pairs of males estimated to encounter one another (7.1 ± 4.5 mm) was comparable to the mean difference between randomly chosen pairs of males (7.4 ± 5.7 mm). However, observed size differences were underrepresented in the smallest (0-2 mm) category compared to random pairwise size differences (0.11 vs. 0.18 ± 0.002; $P <$ 0.0003; Figure 3).

Figure 3. Estimated SVL differences between male pairs estimated to co-occur (blue bars), compared with pairwise SVL differences between randomly chosen pairs of males, with random males' sizes estimated on the same days at the observed co-occurrences (white bars).

Males that encountered more females had a greater spatial extent ($r^2 = 0.10$, $F_{1, 113} = 13.0$, P

<0.001; Figure 4) and were larger in size on average at their encounters with females ($r^2 = 0.09$, $F_{1, 113}$

= 13.0, $P <$0.001; Figure 4); the interaction between spatial extent and SVL was not significant ($r^2 =$

0.0, $F_{1, 113} = 0.03$, $P = 0.87$)

Under the parentage analysis in which we set the potential sires for each offspring to be the

males estimated to have encountered their mother, paternity was assigned to 84% of all offspring

(323 individuals) at an 80% confidence level. We found that 64% of mothers bore offspring sired by

more than one male; including offspring with unassigned sires, this proportion rose to 81%. Using a

resampling approach to calculate p-values, we found that sires of offspring encountered mothers

significantly more often than did non-sires (mean number of encounters between mothers and sires:

102 ± 140; non-sires: 40 ± 65; $P < 0.0001$). Accounting for variation across males in how often they

encounter mothers, we found that sires encountered females significantly later than non-sires (mean

± SD of the last hour of encounter for sires: 892 ± 110; non-sires: 605 ± 258; $P < 0.0001$) and were

significantly bigger than non-sires (mean ± SD of the maximum male SVL across encounters for

male-female pairs, for sires: 57.8 ± 3.0 mm; non-sires: 53.2 ± 5.6 mm; $P < 0.0001$; Figure 5).

Figure 4. Relationship between the number of females encountered by males and the males' spatial extent (top; measured as the mean distance to the centroid) and males' mean estimated SVL across encounters (bottom).

Figure 5. Maximum male snout-vent length (SVL) and the hour of last encounter for male-female pairs, colored by whether or not the male sired any of the female's offspring.

Discussion

How animals move through space determines how many and which other individuals they

encounter, setting the stage for all of their subsequent social and reproductive interactions and

ultimately determining their reproductive success. Understanding animals' movement patterns and

the encounters they bring about is thus a key step in characterizing the mating system of a

population or species, and is essential for determining how behavior both facilitates and is subject to

sexual selection. In particular, female mate choice can contribute to sexual selection in natural

populations only if individuals' movement patterns give females the opportunity to choose among

mates. Developing a spatiotemporal characterization of the movement patterns of a population of

A. sagrei lizards, we found not only that a majority of males (60%) encountered multiple females but

also that a majority of females (78%) encountered multiple males over the first three months of the

breeding season, indicating the potential in *A. sagrei* for complex polygynandrous mating patterns

with ample scope for female choice.

Consistent with previous genetic descriptions of anole mating systems [40–42], we found

that a majority of females (64%–81%) bore offspring sired by more than one male. However, our

results are at odds with most previous behavioral descriptions of movement patterns and mating

systems in *Anolis* lizards, which leaned heavily on, and were constrained by, the framework of

territoriality. These behavioral descriptions were often coupled with an implicit expectation that

anoles mate strictly polygynously, i.e., males mate with multiple females, but females mate with just

the one male in whose territory they reside (reviewed in [21]). Consequently, field studies have often

implied that the opportunity for female choice in anoles is limited because it is precluded by

territoriality (e.g. [43–45], but see [46]). Concurrently, despite varying evidence as to whether female

choice is even possible in natural populations, laboratory studies have offered females the choice

between males to assess precopulatory mate preference [47–49] and have mated females with

multiple males to assess postcopulatory sexual selection [50, 51]. Our demonstration that individual

A. sagrei lizards in a natural population move such that females frequently encounter and mate with

multiple males indicates that female anoles definitely have the opportunity to exercise post-

copulatory mate choice and might also exercise pre-copulatory mate choice. This calls into question

the utility of territorial polygyny as a description of these lizards' mating system. More generally,

discrepancies between behavioral and genetic descriptions of mating systems are widespread across

animals (e.g. [16, 17]), and our results suggest that the re-examination of animals' movement

behavior may help to reconcile these discrepancies.

In territorial species, such re-examinations could further prompt us to revisit explanations of

aspects of animals' behavior that have long been interpreted as characteristic of territorial polygyny,

to discern if they might also be consistent with female choice. For example, we found that males

encounter size-matched males less often than expected at random, which is consistent with male

body size in anoles determining the outcome of male-male fights over territory ownership and

access to mates [52, 53] and larger males subsequently excluding other large males from their

territories [54, 55]. In this context of territoriality, smaller males are hypothesized to evade detection

by larger territorial males, residing in their territories and attempting to 'sneak' copulations with

resident females. But in taxa where females choose mates based on male body size (as may be the

case in this species; see below), larger males that retain smaller neighbors can accrue a mating

advantage compared to males with neighbors of equal size (e.g. [56, 57]. Thus, males may engage in

agonistic interactions to exclude size-matched but not smaller males from their vicinity, arranging

themselves spatially relative to other males in a manner conducive to their success in the arena of

female choice [58].

That said, territoriality is a multi-faceted concept, pertinent not only to reproductive

interactions but also to habitat selection and space use behavior [59]. Our results support long-

standing views in some of these facets. We found that males are more likely to move greater

distances and had a greater spatial extent than did females, which is consistent with previous

estimates of sex differences in territory size (e.g. [60, 61]), and with evidence for male-biased

dispersal in anoles [62, 63]. But the spatial extent of individuals in this population is substantially

higher than previous estimates of territory size in this and ecologically similar species (e.g. ~3 m^2 for

females and ~10 m^2 to ~14 m^2 for males [60, 64, 65], compared with minimum convex polygons

with mean areas of 36 m^2 for females and 225 m^2 for males in this study). Although it is always

possible that lizards in our site behaved unusually or, more generally, that movement behavior varies

across populations and species, the discrepancy may also be due in part to limited spatial and

temporal sampling in previous studies leading to underestimates of anole space use. Limited

sampling may also partially underlie why female multiple mating has rarely been considered

important in past research on anole territoriality—studies with limited spatial or temporal sampling

extents will be constrained in what behaviors they can detect (reviewed in [21]). Indeed, subsampling

from our dataset shows that if we had limited our spatial or temporal sampling extent to match the

median sampling of previous studies, we would have detected a greatly reduced number of male-

female pairs whose home ranges overlap with one another (see Supplementary Information). In

sum, we posit that while certain tenets of territoriality are well-supported in anoles, previous studies

have likely underestimated the complexity of *Anolis* lizards' movement patterns and social

interactions by being constrained by a territorial framework. Future studies of movement patterns

that explicitly do not depend on a territorial framework, conducted in a diversity of populations and

species across a range of habitats and densities, will be necessary to determine if we are correct.

Moreover, it remains unknown if similar problems afflict other species that have long been

described as territorial.

Larger males not only encountered more females but were also more likely to sire offspring

than smaller males, suggesting strong sexual selection for larger body size in male *A. sagrei*. These

results are consistent with conventional wisdom in anoles [66], and, more generally, with evidence

from across animals that body size in males predicts reproductive success [24]. However, most

previous evidence in favor of this hypothesis in anoles is based on the estimation of mating patterns

from home range area and overlap within the framework of territoriality (e.g. [60, 67]. More recent

genetic evidence includes both marginal [51, 68] and strong [41] trends towards larger males siring

more offspring. All results from our study, which integrates behavioral and genetic data, point to

sexual selection favoring larger body size in males, helping to explain the marked male-biased sexual

size dimorphism seen in this species. As a species with largely indeterminate growth, body size in

these lizards can be an indicator of age [43, 69], or the ability to survive and thrive, suggesting an

adaptive reason for females to choose to mate with, or bear offspring sired by, larger males [70].

Selection on body size is difficult to disentangle from last-male precedence—because males

are smaller at earlier times in the breeding season, it is possible that large males sire more offspring

simply because they have encountered and mated with females more recently. Last-male precedence

is found in many animals [23], including reptiles (e.g. [71]), but previous evidence from laboratory

experiments in *A. sagrei* suggests instead that the first male to mate with a female is more likely to

sire offspring than a later male [72]. More targeted laboratory experiments could readily separate size

effects from mating order effects, lending insight into the mechanisms by which post-copulatory

mate choice operates in these animals.

Additionally, sexual selection may act on movement behavior—males with larger spatial

extents encountered more females than males with smaller spatial extents. Because male body size

and spatial extent were only very weakly correlated, the results do not indicate strong ontogenetic

shifts in movement behavior. Thus, it appears that there are multiple ways for males to achieve

reproductive success—they can grow large, they can traverse large areas, or both [73].

However, understanding how the effects of body size, spatial extent, and last-male

precedence integrate with the processes of males first encountering females and then siring their

offspring is not necessarily straightforward. In particular, it remains unknown, but is unlikely, that a

single movement or mating strategy can be adaptive in the face of such social complexity. Instead,

animals may make decisions about movement depending on their social and environmental context,

as opposed to adopting fixed patterns of space-use [5, 7, 8, 74]. Such context-dependent decision

making is often referred to as the maintenance of "alternative mating strategies" (reviewed in [75])

though this variation need not be strictly discrete (e.g. [73, 76]). Individual-based models that

incorporate the various sequential, compounding influences on reproductive success could reveal if

males can make adaptive, context-dependent decisions to move or stay at particular locations based

on their phenotype and the social and environmental situations they find themselves in.

Discerning if the decision rules used by individuals in a species are consistent across variable

habitats and populations likely represents a promising way of understanding animal mating systems

[77, 78]. The Markov chain approach to modelling movement patterns and estimating interactions

presented here could be modified in future studies of anoles as well as other taxa to incorporate

both social and environmental variation influencing these decision rules—transition probabilities

between locations could be scaled according to habitat preferences or the occupancy of a location by

specific individuals, for example. Finer spatial and temporal scales of location sampling, made

possible with the advancement of automated methods of tracking animals in the wild [79], could also

allow for individual-level parameterization of any movement model, potentially improving the

accuracy with which encounters are estimated. For example, 26 individual offspring (7% of total)

were assigned sires not among the males estimated to encounter their mothers (see Supplementary

Information)—it is possible that a greater temporal resolution of location observations would have

detected the movements bringing about these encounters and subsequent matings. At present,

however, the size and cost of tracking or receiving devices and the spatial resolution of GPS data

limit the range of taxa for which such methods can be employed.

Ultimately, even finer scale location sampling will be insufficient for determining which

encounters in fact lead to matings—discovering this crucial aspect of animals' behavior will depend

on focal animal observations of encounters in natural or naturalistic conditions. For a male, the gap

between encountering a female and siring her offspring can include the sequential gauntlets of male-

male competition, female mating preferences, and post-copulatory competition and choice [1, 2].

Disentangling effects on mating success at these various levels will be essential to fully

understanding how sexual selection shapes behavior, and will require close observation of animal

interactions prior to and including mating [80–83].

However, movement behavior is a precursor to bringing about any of these interactions—it

comprises the first step of sexual selection and its quantification is therefore necessary for

understanding the shape that sexual selection can take. In this paper, we develop a spatiotemporal

framework to quantify movement behavior to estimate encounters between potential mates. This

framework reveals an infrequently-recognized opportunity for female mate choice in *Anolis sagrei*,

demonstrates that larger males are favored across multiple levels of sexual selection, and shows

evidence for sexual selection on movement behavior and the timing of male-female encounters. We

hope that similar frameworks centered on movement behavior can help to organize disparate studies

that approach sexual selection at different levels in a variety of animal taxa.

Acknowledgements

This work was carried out under permit EXOT-15-09 from the U.S. Fish and Wildlife

Service, and all procedures were approved under IACUC Protocol 26-11. Jonathan Suh, Claire

Dufour, Rachel Moon, Bonnie Kircher, and Barbara Tamires de Silviera assisted with data

collection. Thom Sanger, Tom Wichman, Steve Johnson, and Stuart McDaniel provided logistical

support. Thom Sanger additionally advised on lizard husbandry and embryo collection. Ben de

Bivort, Edward Burnell and Colin Donihue advised on the statistical analyses and, along with Yoel

Stuart and Pavitra Muralidhar, provided comments that improved the manuscript. Hanna Wegener,

Jason Kolbe, Anthony Geneva, and Pavitra Muralidhar advised on the genetic methods and analyses.

This project was funded by a Young Explorers Grant from the National Geographic Society, a

student research grant from the Animal Behavior Society, the Robert A. Chapman Memorial

Scholarship from Harvard University, and the John Templeton Foundation.

Statement of Authorship:

AK and JBL designed the study, AK collected data and performed analyses, AK and JBL wrote the

manuscript.

Data Accessibility Statement:

Data and code are currently available on GitHub (https://github.com/ambikamath/anolismovt).

References

1. Wiley RH, Poston J. 1996 Indirect mate choice, competition for mates, and coevolution of

the sexes. *Evolution*, **50**, 1371–1381. (doi: 10.1111/j.1558-5646.1996.tb03911.x)

2. Bonduriansky, R. 2003 Layered sexual selection: a comparative analysis of sexual behaviour

within an assemblage of piophilid flies. *Can. J. Zool.*, **81**, 479–491.

3. Wagner Jr. WE. 1998 Measuring female mating preferences. *Anim. Behav.*, **55**, 1029–1042.

(doi: 10.1006/anbe.1997.0635)

4. Kvarnemo C, Simmons L. 2013. Polyandry as a mediator of sexual selection before and after

mating. *Phil. Trans. R. Soc. B*, **368**, 20120042. (doi: 10.1098/rstb.2012.0042)

5. Uy JAC, Patricelli GL, Borgia G. 2000. Dynamic mate-searching tactic allows female satin

bowerbirds *Ptilonorhynchus violaceus* to reduce searching. *Proc. R. Soc. Lond. B*, **267**, 251–256.

6. Jones AG. 2009 On the opportunity for sexual selection, the Bateman gradient and the

maximum intensity of sexual selection. *Evolution*, **63**, 1673–1684. (doi: 10.1111/j.1558-

5646.2009.00664.x)

7. Dunn PO, Whittingham LA. 2006 Search costs influence the spatial distribution, but not the

level, of extra-pair mating in tree swallows. *Behav. Ecol. Sociobiol.,* **61**, 449–454. (doi:

10.1007/s00265-006-0272-3)

8. DuVal EH, Kapoor JA. 2015 Causes and consequences of variation in female mate search

investment in a lekking bird. *Behav. Ecol.*, **26**, 1537–1547. (doi: 10.1093/beheco/arv110)

9. Jones AG, Ratterman NL. 2009 Mate choice and sexual selection: What have we learned

since Darwin? *Proc. Nat. Acad. Sci.*, **106**, 10001–10008. (doi: 10.1073/pnas.0901129106)

10. Dougherty LR, Shuker DM. 2015 The effect of experimental design on the measurement of

mate choice: a meta-analysis. *Behav. Ecol.*, **26**, 311–319. (doi:10.1093/beheco/aru125)

11. Firman RC, Gasparini C, Manier MK, Pizzari T. 2017 Postmating female control: 20 years of

cryptic female choice. *Trends Ecol. Evol.* **32**, 368–382. (doi: 10.1016/j.tree.2017.02.010)

12. Morris MR. 1989 Female choice of large males in the treefrog *Hyla chrysoscelis*: the importance

of identifying the scale of choice. *Behav. Ecol. Sociobiol.*, **25**, 275–281.

13. Roff DA. 2015 The evolution of mate choice: a dialogue between theory and experiment.

Ann. N.Y. Acad. Sci., **1360**, 1–15. (doi: 10.1111/nyas.12743)

14. Real L. 1990 Search theory and mate choice. I. Models of single-sex discrimination. *Am.

Nat.*, **136**, 376–405. (doi: abs/10.1086/285103)

15. Shuker DM, Day TH. 2002 Mate sampling and the sexual conflict over mating in seaweed

flies. *Behav. Ecol.*, **13**, 83–86. (doi: 10.1093/beheco/13.1.83)

16. Hughes C. 1998 Integrating molecular techniques with field methods in studies of social

behavior: A revolution results. *Ecology*, **79**, 383–399. (doi: 10.1890/0012-

9658(1998)079[0383:IMTWFM]2.0.CO;2)

17. Griffith SC, Owens IPF, Thuman KA. 2002 Extra pair paternity in birds: a review of

interspecific variation and adaptive function. *Mol. Ecol.*, **11**, 2195–2212.

18. Double M, Cockburn A. 2000 Pre-dawn infidelity: females control extra-pair mating in

superb fairy-wrens. *Proc. R. Soc. Lond. B*, **267**, 465–470. (doi: 10.1098/rspb.2000.1023)

19. Thom MD, MacDonald DW, Mason GJ, Pedersen V, Johnson PJ. 2004 Female American

mink, *Mustela vison*, mate multiply in a few choice environment. *Anim. Behav.*, **67**, 975–984.

(doi:10.1016/j.anbehav.2003.09.008)

20. Stopher KV, Nussey DH, Clutton-Brock TH, Guinness F, Morris A, Pemberton JM. 2011.

The red deer rut revisited: female excursions but no evidence females move to mate with

preferred males. *Behav. Ecol.*, **22**, 808–818. (doi: 10.1093/beheco/arr052)

21. Kamath A, Losos JB. 2017 The erratic and contingent progression of research on

territoriality: a case study. *Behav. Ecol. Sociobiol.*, **71**, 89. (doi: 10.1007/s00265-017-2319-z)

22. Kokko H, Mappes J. 2012 Multiple mating by females is a natural outcome of a null model

of mate encounters. *Entomol. Exp. Appl.* **146**, 26–37. (doi: 10.1111/j.1570-7458.2012.01296.x)

23. Birkhead TR, Hunter FM. 1990 Mechanisms of sperm competition. *Trends Ecol. Evol.*, **5**, 48–

52. (doi: 10.1016/0169-5347(90)90047-H)

24. Andersson M. 1994 *Sexual Selection*. Princeton University Press, Princeton, USA.

25. Kolbe JJ, Glor RE, Rodríguez–Schettino L, Chamizo Lara AR, Larson A, Losos JB. 2004

Genetic variation increases during biological invasion by a Cuban lizard. *Nature*, **431**, 177–

181. (doi: 10.1038/nature02807)

26. Kolbe JJ, Larson A, Losos JB, de Queiroz K. 2008 Admixture determines genetic diversity

and population differentiation in the biological invasion of a lizard species. *Biol. Lett.* **4**, 434–

437. (doi: 10.1098/rsbl.2008.0205)

27. Sanger TJ, Losos JB, Gibson-Brown JJ. 2007 A developmental staging series for the lizard

genus *Anolis*: A new system for the integration of evolution, development, and ecology. *J.*

Morph., **269**, 129–137. (doi: 10.1002/jmor.10563)

28. Fisher M, Muth, A. 1989 A technique for permanently marking lizards. *Herpetol. Rev.*, **20**, 45–

46.

29. Sanger TJ, Hime PM, Johnson MJ, Diani J, Losos JB. 2008 Laboratory protocols for

husbandry and embryo collection of *Anolis* lizards. *Herpetol. Rev.*, **39**, 58–63.

30. R Core Team. 2016 R: A language and environment for statistical computing. R Foundation

for Statistical Computing, Vienna.

31. Bolker B, R Development Core Team. 2016 bbmle: Tools for General Maximum Likelihood

Estimation. R package version 1.0.18. https://CRAN.R-project.org/package=bbmle

32. Schoener TW, Schoener A. 1978 Estimating and interpreting body-size growth in some
Anolis lizards. *Copeia*, **1978**, 390–405.

33. Bardeleben C, Palchevskiy V, Calsbeek R, Wayne RK. 2004 Isolation of polymorphic
tetranucleotide microsatellite markers for the brown anole (*Anolis sagrei*). *Mol. Ecol. Notes*, **4**,
176–178. (doi: 10.1111/j.1471-8286.2004.00602.x)

34. Wordley C, Slate J, Stapley J. 2011 Mining online genomic resources in *Anolis carolinensis*
facilitates rapid and inexpensive development of cross-species microsatellite markers for the
Anolis lizard genus. *Mol. Ecol. Resour.*, **11**, 126–133. (doi:10.1111/j.1755-0998.2010.02863.x)

35. Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A,
Markowitz S, Duran C, *et al.* 2012 Geneious Basic: an integrated and extendable desktop
software platform for the organization and analysis of sequence data. *Bioinformatics*, **28**, 1647–
1649. (doi: 10.1093/bioinformatics/bts199)

36. Kalinowski ST, Taper ML, Marshall TC. 2007 Revising how the computer program
CERVUS accommodates genotyping error increases success in paternity assignment. *Mol.*
Ecol., **16**, 1099–1106. (doi: 10.1111/j.1365-294X.2007.03089.x)

37. Le Bas NR. 2001 Microsatellite determination of male reproductive success in a natural
population of the territorial ornate dragon lizard, *Ctenophorus ornatus*. *Mol. Ecol.*, **10**, 193–203.
(doi: 10.1046/j.1365-294X.2001.01174.x)

38. Husak JF, Fox SF, Lovern MB, van den Bussche RA. 2006 Faster lizards sire more
offspring: sexual selection on whole-animal performance. *Evolution*, **60**, 2122–2130. (doi:
10.1111/j.0014-3820.2006.tb01849.x)

39. Rasmussen HB, Okello JBA, Wittemyer G, Siegismund HR, Arctander P, Vollrath F,
Douglas-Hamilton I. 2008 Age- and tactic-related paternity success in male African
elephants. *Behav. Ecol.*, **19**, 9–15. (doi:10.1093/beheco/arm093)

40. Passek KM. 2002 *Extra-pair paternity within the female-defense polygyny of the lizard,* Anolis
carolinensis*: Evidence of alternative mating strategies*. Dissertation, Virginia Polytechnic Institute,
Blacksburg, USA.

41. Johnson MA. 2007 *Behavioral ecology of Caribbean* Anolis *lizards: A comparative approach*.
Dissertation, Washington University, St. Louis, USA.

42. Calsbeek R, Bonneaud C, Prabhu S, Manoukis N, Smith TB. 2007 Multiple paternity and
sperm storage lead to increased genetic diversity in *Anolis* lizards. *Evol. Ecol. Res.* **9**, 495–503.

43. Schoener TW, Schoener A. 1980 Densities, sex ratios, and population structure in four
species of Bahamian *Anolis* lizards. *J. Anim. Ecol.*, **49**, 19–53.

44. Stamps JA. 1983 Sexual selection, sexual dimorphism, and territoriality. In: *Lizard Ecology*.
[eds.] Huey R, Pianka ER, Schoener TW. Harvard University Press, Cambridge, USA, pp
169–204.

45. Jenssen TA, Lovern MB, Congdon JD. 2001 Field-testing the protandry based mating
system for the lizard, *Anolis carolinensis*: does the model organism have the right model? *Behav.*
Ecol. Sociobiol., **50**, 162–172. (doi: 10.1007/s002650100349)

46. Tokarz RR. 1998 Mating pattern in the lizard *Anolis sagrei*: Implications for mate choice and
sperm competition. *Herpetologica*, **54**, 388–394.

47. Andrews RM. 1985 Male choice by females of the lizard, *Anolis carolinensis*. *J. Herpetol.*, **19**,
284–289.

48. Lailvaux SP, Irschick DJ. 2006 No evidence for female association with high-performance
males in the green anole lizard, *Anolis carolinensis*. *Ethology*, **112**, 707–715. (doi:
10.1111/j.1439-0310.2006.01210.x)

49. Flanagan SP, Bevier CR. 2014 Do male activity level and territory quality affect female
association time in the brown anole, *Anolis sagrei*? *Ethology*, **120**, 365–374. (doi:
10.1111/eth.12213)

50. Calsbeek R, Bonneaud C. 2008 Postcopulatory fertilization bias as a form of cryptic sexual
selection. *Evolution*, **62**, 1137–1148. (doi:10.1111/j.1558-5646.2008.00356.x)

51. Cox RM, Calsbeek R. 2010 Cryptic sex-ratio bias provides indirect genetic benefits despite
sexual conflict. *Science*, **328**, 92–94. (doi: 10.1126/science.1185550)

52. Tokarz RR. 1985 Body size as a factor determining dominance in staged agonistic
encounters between male brown anoles (*Anolis sagrei*). *Anim. Behav.* **33**, 746–753. (doi:
10.1016/S0003-3472(85)80006-3)

53. Jenssen TA, Decourcy DA, Congdon JD. 2005 Assessment in contests of male lizards
(*Anolis carolinensis*): how should smaller males respond when size matters? *Anim. Behav.*, **69**,
1325–1336. (doi:10.1016/j.anbehav.2004.07.023)

54. Trivers RL. 1976 Sexual selection and resource-accruing abilities in *Anolis garmani*. *Evolution*,
30, 253–269. (doi: 10.1111/j.1558-5646.1976.tb00908.x)

55. Fleishman LJ. 1988 The social behavior of *Anolis auratus*, a grass anole from Panama. *J.*
Herpetol., **22**, 13–23.

56. Callander S, Jennions MD, Backwell PRY. 2011 Female choice over short and long
distances: neighbour effects. *Behav. Ecol. Sociobiol.*, **65**, 2071–2078. (doi: 10.1007/s00265-011-
1216-0)

57. Callander S, Hayes CL, Jennions MD, Backwell PRY. 2012 Experimental evidence that
immediate neighbors affect male attractiveness. *Behav. Ecol.*, **24**, 730–733. (doi:
10.1093/beheco/ars208)

58. Höglund J, Johansson T, Pelabon C. 1997. Behaviourally mediated sexual selection:
characteristics of successful male black grouse. *Anim. Behav.*, **54**, 255–264. (doi:
10.1006/anbe.1996.0459)

59. Maher CR, Lott DF. 1995 Definitions of territoriality used in the study of variation in
vertebrate spacing systems. *Anim. Behav.*, **49**, 1581–1597. (doi: 10.1016/0003-3472(95)90080-
2)

60. Schoener TW, Schoener A. 1982 Intraspecific variation in home-range size in some *Anolis*
lizards. *Ecology*, **63**, 809–823. (doi: 10.2307/1936801)

61. Jenssen TA, Nunez SC. 1998 Spatial and breeding relationships of the lizard, *Anolis*
carolinensis: Evidence of intrasexual selection. *Behaviour*, **135**, 981–1003. (doi:
10.1163/156853998792640341)

62. Johansson H, Surget-Groba Y, Thorpe RS. 2008 Microsatellite data show evidence for male-
biased dispersal in the Caribbean lizard *Anolis roquet. Mol. Ecol.*, **17**, 4425–4432. (doi:
10.1111/j.1365-294X.2008.03923.x)

63. Calsbeek R. 2009 Sex-specific adult dispersal and its selective consequences in the brown
anole, *Anolis sagrei. J. Anim. Ecol.*, **78**, 617–624. (doi: 10.1111/j.1365-2656.2009.01527.x)

64. Rand AS. 1967 Ecology and social organization in the iguanid lizard *Anolis lineatopus. Proc.*
U.S. Nat. Mus., **122**, 1–79.

65. Calsbeek R, Marnocha E. 2006 Context dependent territory defense: The importance of
habitat structure in *Anolis sagrei. Ethology*, **112**, 537–543. (doi: 10.1111/j.1439-
0310.2006.01194.x)

66. Losos JB. 2009 *Lizards in an evolutionary tree.* University of California Press, Berkeley,
California, USA.

67. Ruby DE. 1984 Male breeding success and differential access to females in *Anolis carolinensis.*
Herpetologica, **40**, 272–280.

68. Harrison AS. 2014 *The evolution and diversity of the* Anolis *dewlap*. Dissertation, Harvard
University, Cambridge, USA.

69. Stamps JA. 1993 Sexual size dimorphism in species with asymptotic growth after maturity.
Biol. J. Linn. Soc., **50**, 123–145. (doi: 10.1006/bijl.1993.1050)

70. Cooper Jr WE, Vitt LJ. 1993 Female mate choice of large male broad-headed skinks. *Anim.*
Behav., **45**, 683–693. (doi: 10.1006/anbe.1993.1083)

71. Friesen CR, Mason RT, Arnold SJ, Estes S. 2014 Patterns of sperm use in two populations
of Red-sided Garter Snake (*Thamnophis sirtalis parietalis*) with long-term female sperm storage.
Can. J. Zool., **92**, 33–40. (doi: 10.1139/cjz-2013-0195)

72. Duryea MC, Kern AD, Cox RM, Calsbeek R. 2013 A novel application of approximate
Bayesian computation for detecting male reproductive advantages due to mating order.
Behav. Ecol. Sociobiol. **67**, 1867–1875. (doi: 10.1007/s00265-013-1612-8)

73. Noble DWA, Wechmann K, Keogh JS, Whiting MJ. 2013 Behavioral and morphological
traits interact to promote the evolution of alternative reproductive tactics in a lizard. *Am.*
Nat., **182**, 726–742. (doi: 10.1086/673535)

74. Jordan LA, Brooks RC. 2011 Recent social history alters male courtship preferences.
Evolution, **66**, 280–287. (doi: 10.1111/j.1558-5646.2011.01421.x)

75. Shuster SM. 2010 Alternative mating strategies. In: *Evolutionary Behavioral Ecology*. [eds.]
Westneat, D.F. & Fox, C.W. Oxford University Press, New York, USA, pp. 435–450.

76. Shine R, Langkilde T, Wall M, Mason RT. 2005 Alternative male mating tactics in garter
snakes, *Thamnophis sirtalis parietalis*. *Anim. Behav.*, **70**, 387–396. (doi:
10.1016/j.anbehav.2005.04.001)

77. Kirkpatrick M, Rand AS, Ryan MJ. 2006 Mate choice rules in animals. *Anim. Behav.*, **71**,
1215–1225. (doi: 10.1016/j.anbehav.2005.11.010)

78. Patricelli GL, Krakauer AH, McElreath R. 2011 Assets and tactics in a mating market:
Economic models of negotiation offer insights into animal courtship dynamics on the lek.
Curr. Zool., **57**, 225–236.

79. Kays R, Crofoot MC, Jetz W, Wikelski M. 2015 Terrestrial animal tracking as an eye on life
and planet. *Science*, **348**, aaa2478. (doi: 10.1126/science.aaa2478)

80. Wong BB, Candolin U. 2005 How is female mate choice affected by male competition. *Biol.*
Rev. Camb. Philos. Soc., **80**, 559–571. (doi: 10.1017/S1464793105006809)

81. Hunt J, Breuker CJ, Sadowski JA, Moore AJ. 200. Male-male competition, female mate

choice and their interaction: determining total sexual selection. *J. Evol. Biol.*, **22**, 13–26. (doi:

10.1111/j.1420-9101.2008.01633.x)

82. Stapley J. 2008 Female mountain log skinks are more likely to mate with males that court

more, not males that are dominant. *Anim. Behav.*, **75**, 529–538. (doi:

10.1016/j.anbehav.2007.05.017)

83. Engqvist L. 2009 Should I stay or should I go? Condition- and status-dependent courtship

decisions in the scorpionfly *Panorpa cognata*. *Anim. Behav.* **78**, 491–497. (doi:

10.1016/j.anbehav.2009.05.021)

Supplementary Information

Modelling lizard locations.

As described in the main text, we modelled lizards' movements between locations using a

Markov chain model, where transition probabilities were fitted as exponentially declining with the

distance between locations (Equation 1 in main text). If **M** is the transition matrix of the Markov

chain model, and an individual lizard was observed at location A_m within hour T_m and then later at

location A_n within hour T_n, then we calculated the probability that that individual was found at a

particular location (L) within the hour $T \in (T_m, \ldots, T_{n-1})$ as follows:

$$P(L, T | A_m, T_m, A_n, T_n) = \frac{\mathbf{M}^{T-(Tm)+1}[A_m, L] \times \mathbf{M}^{(Tn)-T+1}[L, A_n]}{\sum_{i=1}^{N} \mathbf{M}^{T-(Tm)+1}[A_m, A_i] \times \mathbf{M}^{(Tn)-T+1}[A_i, A_n]}$$

N is the total number of locations mapped; in our site, we mapped a total of 318 locations,

which included not only locations at which lizards were observed but also trees within the site on

which lizards were not observed but to which they might have moved (Figure S1). For the hours

preceding the first observation and following the last observation for each individual, lizards were

assumed to be equally likely to be anywhere in the site (i.e. the appropriate row or column of the

transition matrix in equation 1 was replaced by a unit vector). In sum, after this procedure, we

obtained, for each lizard, a vector of probabilities of the lizard being in each of the 318 locations

within each hour during the sampling period.

Figure S1. Map of all locations at which lizards were observed. In areas of continuous vegetation, locations more than 1m apart were considered distinct. We also mapped the location of all trees within the site at which lizards were not observed, to include all trees to which a lizard could potentially have moved in our estimations of their movement patterns, amounting to a total of 318 locations.

Estimating Encounters

To estimate whether or not a pair of lizards encountered one another, we first performed an

element-wise multiplication of the two matrices of the individuals' probabilities of being at each

location within each hour, and then summed probabilities across all locations for each hour in the

resultant matrix for the pair. Probabilities were summed across locations so that two individuals that

may be moderately likely to co-occur at multiple locations could still be estimated as encountering

one another.

The calculated probabilities of observed co-occurrences, defined as a pair of lizards observed

in the same location within one hour of each other, were used to set the probability cutoff by which

we defined whether or not other pairs of lizards encountered one another. These cutoffs depended

on location. The 318 locations in this site were not uniformly distributed, with some locations

clustered more closely together than others (Figure S1). When the probability of movement declines

with distance, as modeled here, it is easier to leave a focal location that is close to other locations

than it is to leave a relatively isolated location. Consequently, the estimated probability of remaining

at the former location will be lower compared to the probability of leaving the latter location. We

therefore set probability cutoffs to vary by the locations at which lizards in the pair were seen, on

the basis of the focal locations' proximity to nearby locations First, for the location of each observed

co-occurrence, we calculated the mean distance to the ten closest locations to it, and saw that, as

expected, the calculated probability of an encounter at that location between two lizards observed to

co-occur increased with increasing distance to the ten closest points (Figure S2). We therefore

binned locations on the basis of these distances, into 1 m increments. The minimum probability

calculated for observed co-occurrences in each bin, denoted by the black horizontal lines in Figure

S2, was set as a cutoff for estimating whether or not an encounter took place, for all pairs of lizards

for all hours in the sampling period. Of course, using such a distance-based probability cutoff to

estimate encounters meant that we needed to assign a distance to all pairs of lizards for all hours. To

do so, at each hour we considered the previous and next locations at which the lizards in the pair

were observed, and found the minimum of the means of the ten closest distances for these

locations. For defining encounters (as a binary "yes/no" variable) for each pair of lizards at each

hour, we assessed if the probability of their encounter at that hour was higher than the appropriate

probability cutoff based on the assigned distance.

Figure S2 Cutoffs (black horizontal lines) for defining encounters, based on the mean distance from each location to the ten closest locations. Points indicate the sum of probabilities of co-occurrences across all locations, for pairs of individuals at hours where the two were observed at the same location within an hour of one another.

To assess the sensitivity of the number of male-female encounters to the probability cutoff described above, we repeated analyses using the 25% quartile instead of the minimum probability in each bin; a higher cutoff is more stringent, and will recover fewer encounters, but we were curious whether females would still be estimated as encountering multiple males across the sampling period. Not surprisingly, conservatively increasing the probability threshold for defining encounters from the minimum probability of observed co-occurrences to the 25% quartile reduced estimates of the number of potential mates encountered by both females (3.4

± 3.1) and males (2.0 ± 2.1), but the potential for females to mate with multiple males remained.

Because boundary effects of a closed site might artificially increase the number of encounters, we

also assessed if the number of male-female encounters was sensitive to treating the site as closed, by

repeating the analyses described above with a simulated buffer of 50 locations, placed randomly

within 20 m around the perimeter of the site. Allowing individuals to move to these locations did

not alter estimates of the average number of males encountered by females (5.1 ± 3.7) or females

encountered by males (2.9 ± 3.0).

Growth Curve for Males

We estimated the growth curve of males in this population by fitting the following logistic

equation (with parameters *a* and *r*) to the observed data, using nonlinear least squares regression

(Schoener and Schoener 1978):

$$L2 = \frac{aL_1}{L_1 + (a - L_1)e^{rD}}$$

where L_1 and L_2 are the SVL of an individual measured at two successive captures, and D is

the number of days between the two captures. We used this logistic growth curve model to estimate

the SVL of each male on the day of each of his encounters, based on his SVL at the nearest capture.

Parameters for the logistic growth curve for male SVL (Equation 3) were estimated to be a = 63.7

and r = -0.016. Snout-vent lengths predicted by the model were highly correlated with measured

SVLs ($r^2 = 0.92$).

Paternity analysis

Tests for sexual selection were performed on the paternity assignments in which candidate

sires for offspring were the list of males their mother was estimated to have encountered (described

in main text). However, we ran two further analyses. First, to assess if high paternity assignment

rates in the main analysis were a consequence of restricting the number of candidate sires relative to

the whole population, we also ran ten replicates of the analysis in which we sampled potential sires

randomly from all 161 males, preserving the number of potential sires for each offspring individual

and supplying the same group of potential sires for all offspring of a single female. Randomizing the

identity but not the number of males encountered by each mother yielded 18% paternity assignment

on average (13% to 24%, at an 80% confidence level), suggesting that the four-fold difference in

assignment rates was due to identifying potential sires with greater accuracy based on their

spatiotemporal movement patterns (assignment rate of 84% in the main analysis).

Second, we ran an analysis in which all sampled males were provided as candidate sires for

all offspring. Simulation parameters specified for this analysis were 161 candidate sires and 0.96

proportion of males typed, based on the proportion of observations in the field that were of marked

males. Paternity assignments were concordant for 80% of the individuals that were assigned

paternity by both analyses. Of the remaining 39 individuals, the LOD scores for 21 mother-sire pairs

differed between analyses by less than 1; these were considered false positives from the second

analysis, and the assignments of the first paternity analysis were retained for downstream analysis.

For the remaining 18 offspring, with mother-sire pairs where the LOD scores between the two

analyses differed by more than 1, we found that results were not altered if we conducted

downstream hypothesis testing both including and excluding their paternity assignments from the

first analysis.

Twenty-six individuals that were assigned paternity only when all males were provided as

candidate sires. For these individuals, we calculated the minimum distance between any locations at

which the mother and putative sire were observed, to assess the minimum distances that these

individuals would have had to move to mate with one another. The minimum distance between

locations at which mothers and sires of these 26 offspring were observed (regardless of *when* these

lizards were found at those locations) ranged from 0 m to 56 m, with a median of 8.7 m.

Subsampling in Time and Space

To understand the effect of limiting sampling in space or time on male-female home range

overlap (i.e. how mating patterns have often been assessed in previous studies of anoles), we asked if

a purely spatial predictor— the overlap in minimum convex polygons (MCPs) between pairs of

individuals—is affected by subsampling from our dataset to match either the median area (400 m^2)

or the median duration of sampling (4 weeks) or both of previous studies (see Table S2 for studies

and sample areas and durations included in the calculation of the medians).

On average, in the whole dataset, female MCPs overlap with those of 12.8 ± 8.7 males,

whereas male MCPs overlap with those of 8.1 ± 6.7 females. In subsamples of a randomly selected

area of 400 m^2 (repeated 404 times), we calculated that females overlapped a mean of 5.5 ± 3.1 males

and males overlapped a mean of 2.3 ± 1.2 females. In subsamples of a duration of four weeks (for

each of a possible 55 start dates), we calculated that females overlapped a mean of 4.5 ± 1.0 males

and males overlapped a mean of 3.4 ± 0.8 females. In subsamples with a randomly selected area of

400 m^2 and duration of four weeks, we calculated that females overlapped a mean of 2.4 ± 1.9 males

and males overlapped a mean of 1.5 ± 1.0 females. Thus, limited sampling in space or time decreases

the number of mates inferred from spatial overlap of MCP estimates of home range. That said, at all

scales of subsampling, we recovered that females overlap with multiple males, possibly suggesting

that our study population may be dissimilar to many of those studied previously, and also hinting

that limited sampling in space and time is not the full explanation for why previous studies of *Anolis*

territoriality have largely ignored the potential for females to encounter and mate with multiple

males (but see Tokarz 1998).

Table S1

Locus	Reference	Primers (5' – 3')	Amplification Conditions*	Null allele frequency (for bolded loci, apparent homozygotes typed at only one allele)	Number of alleles
S70	Bardeleben et al. 2004	GCAAGAGGTAGCCTCAGC GTTATCAGTACGAGGCACTG	TD1	-0.002	18
S77	Bardeleben et al. 2004	GAGTAAAGGTCTGGGTCAGG GCAGTACAAATACCACAGAGC	TD1	-0.006	16
AC8	Wordley et al. 2011	CCCAATAGAGGAAAGGGACC AGAATCACGCCTTCTGCTTT	57.7°C	**0.053**	10
AC19	Wordley et al. 2011	GAAAAGTAGTGGGGCATTGG AGTTTCCCAAGAAAACCCGT	TD1	0.007	16
AC45	Wordley et al. 2011	GCATCACCAGTCCCTTTG GCTTTGGTTGAGTTGTATTGT TTG	57.7°C	**0.11**	6
G11	J. Kolbe (pers. comm.)	GGTGCCACTGGGTATTCTGT CTCAATCCTGCTGTCACCTG	TD2	**0.16**	19

*Amplification conditions:

1. 57.7°C: 95°C for 2', 35 cycles of 94°C for 45s, 57.7°C for 45s, 72°C for 1', 72°C for 10'.
2. TD1: 95°C for 5', 10 touchdown cycles of 95°C for 30s, 55°C – 50°C for 30s, 72°C for 45s, 25 cycles of 95°C for 30s, 50°C for 30s, 72°C for 45s, 72°C for 5'.
3. TD2: 95°C for 5', 10 touchdown cycles of 95°C for 30s, 64°C – 55°C for 30s, 72°C for 45s, 25 cycles of 95°C for 30s, 55°C for 30s, 72°C for 45s, 72°C for 5'.

Table S2. Sampling area and duration of previous studies on *Anolis* home ranges

Note that though the median duration is 3.5 weeks, a more conservative value of 4 weeks was used in the subsampling analysis above.

Study	Sampling Area (in m²)	Sampling Duration (in weeks)
Evans 1938	150000	3
Greenberg and Noble 1944	25	47
Gordon 1956	400	52
Sexton et al. 1963	200	4
Rand 1967	425	5
Jenssen 1970	930	8
Jackson 1973	14000	1
Philibosian 1975	420	0.2
Stamps 1977	9.5	5
	34	13
	47	13
	37	15
Hicks and Trivers 1983	12000	9
Ruby 1984	441	28
Fleishman 1988	169	10
Jenssen and Nunez 1998	36	1
Tokarz 1998	137	5
Pereira et al. 2002	12	1
	70	1
Paterson 2002	400	1
McMann and Paterson 2003	400	1
Calsbeek 2009	1500	2
Johnson et al. 2010	500	3
Nicholson and Richards 2011	14000	52
Bush et al. 2016	875	3
Schoener and Schoener 1982	100	–
Fitch and Henderson 1976	–	1
Fitch and Henderson 1987	–	2
Fobes et al. 1992.	–	1.5
Median	**400**	**3.5**

References:

Bardeleben, C., Palchevskiy, V., Calsbeek, R., & Wayne, R.K. 2004. Isolation of polymorphic

tetranucleotide microsatellite markers for the brown anole (*Anolis sagrei*). *Mol. Ecol. Notes*, 4, 176–178.

Birkhead, T.R., & Hunter, F.M. (1990). Mechanisms of sperm competition. Trends in Ecology and

Evolution 5: 48–52.

Bush, J.M., M.M. Quinn, E.C. Balreira, M.A. Johnson. 2016. How do lizards determine dominance? Applying ranking algorithms to animal social behaviour. Animal Behaviour 118: 65–74.

Calsbeek, R. 2009. Sex-specific adult dispersal and its selective consequences in the brown anole, *Anolis sagrei*. Journal of Animal Ecology 78: 617–624.

Evans, L.T. 1938. Cuban field studies on the territoriality of the lizard *Anolis sagrei*. Journal of Comparative Psychology 25: 97–125.

Fitch, H.S., and R.W. Henderson. 1976. A field study of the rock anoles (Reptilia, Lacertilia, Iguanidae) of Southern Mexico. Journal of Herpetology 10: 303–311.

Fitch, H.S., and R.W. Henderson. 1987. Ecological and ethological parameters in *Anolis bahorucoensis*, a species having rudimentary development of the dewlap. Amphibia-Reptilia 8: 69–80.

Fleishman, L.J. 1988. The social behavior of *Anolis auratus*, a grass anole from Panama. Journal of Herpetology 22: 13–23.

Fobes, T.M., R. Powell, J.S. Parmerlee Jr., A. Lathrop, and D.D. Smith. 1992. Natural history of *Anolis cybotes* (Sauria: Polychridae) from an altered habitat in Barahona, Dominican Republic. Caribbean Journal of Science 28: 200–207.

Gordon, R.E. 1956. The biology and biodemography of *Anolis carolinensis carolinensis* Voight. Ph.D. Dissertation, Tulane University.

Greenberg, B., and G.K. Noble. 1944. Social behavior of the American chameleon (Anolis carolinensis Voight). Physiological Zoology 17: 392 – 439.

Hicks, R.M., and R.L. Trivers. 1983. The social behavior of *Anolis valencienni*. Pp. 570 – 595 *In* G.J. Rhodin, and K. Miyata. *Recent Advances in Herpetology and Evolutionary Biology.* Museum of Comparative Zoology, Cambridge, MA, USA.

Jackson, J.F. 1973. Notes on the population biology of *Anolis tropidonotus* in a Honduran highland pine forest. Journal of Herpetology 7: 309–311.

Jenssen, T.A. 1970. The ethoecology of *Anolis nebulosus* (Sauria, Iguanidae). Journal of Herpetology, 4: 1–38.

Jenssen, T.A., and S.C. Nunez. 1998. Spatial and breeding relationships of the lizard, *Anolis carolinensis*: Evidence of intrasexual selection. Behaviour 135: 981–1003.

Johnson, M.A., L.J. Revell, and J.B. Losos. 2010. Behavioral convergence and adaptive radiation: effects of habitat use on territorial behavior in *Anolis* lizards. Evolution 64: 1151–1159.

McMann, S., and A.V. Paterson. 2003. The relationship between location and displays in a territorial lizard. Journal of Herpetology 37: 414–416.

Nicholson, K.E., and P.M. Richards. 2011. Home-range size and overlap within an introduced population of the Cuban knight anole, *Anolis equestris* (Squamata: Iguanidae). Phyllomedusa 10: 65–73.

Paterson, A.V. 2002. Effects of an individual's removal on space use and behavior in territorial neighborhoods of brown anoles (*Anolis sagrei*). Herpetologica 58: 382–393.

Pereira, H.M., S.R. Loarie, and J. Roughgarden. 2002. Monogamy, polygyny and interspecific interactions in the lizards *Anolis pogus* and *Anolis gingivinus*. Caribbean Journal of Science 38: 132–136.

Philibosian, R. 1975. Territorial behavior and population regulation in the lizards, *Anolis acutus* and *A. cristatellus*. Copeia 1975: 428–444.

Rand, A.S. 1967. Ecology and social organization in the iguanid lizard *Anolis lineatopus*. Proceedings of the United States National Museum 122: 1–79.

Ruby, D.E. 1984. Male breeding success and differential access to females in *Anolis carolinensis*. Herpetologica 40: 272–280.

Schoener, T.W., and A. Schoener. 1982. Intraspecific variation in home-range size in some *Anolis* lizards. Ecology 63: 809–823.

Sexton, O.J., H.F. Heatwole, and E.H. Meseth. 1963. Seasonal population changes in the lizard, *Anolis limifrons*, in Panama. The American Midland Naturalist 69: 482–491.

Stamps, J.A. 1977a. The relationship between resource competition, risk, and aggression in a tropical territorial lizard. Ecology 58: 349–358.

Tokarz, R.R. 1998. Mating pattern in the lizard *Anolis sagrei*: Implications for mate choice and sperm competition. Herpetologica 54: 388–394.

Wordley, C., Slate, J., & Stapley, J. 2011. Mining online genomic resources in *Anolis carolinensis* facilitates rapid and inexpensive development of cross-species microsatellite markers for the *Anolis* lizard genus. Molecular Ecology Resources, 11: 126–133.