1 **Title:** Correcting for batch effects in case-control microbiome studies.
2

3 **Authors:** Sean M. Gibbons[1,2,3], Claire Duvallet[1,2], and Eric J. Alm[1,2,3,*]
4
5 **Affiliations:** [1] Department of Biological Engineering, Massachusetts Institute of
6 Technology, Cambridge, MA, USA; [2] Center for Microbiome Informatics and
7 Therapeutics, Cambridge, MA, USA; [3] The Broad Institute of MIT and Harvard,
8 Cambridge, MA, USA
9
10 [*] corresponding author: ejalm@mit.edu

11
12 **Abstract**
13

High-throughput data generation platforms, like mass-spectrometry, microarrays,
and second-generation sequencing are susceptible to batch effects due to run-to-
run variation in reagents, equipment, protocols, or personnel. Currently, batch
correction methods are not commonly applied to microbiome sequencing
datasets. In this paper, we compare different batch-correction methods applied to
microbiome case-control studies. We introduce a model-free normalization
procedure where features (i.e. bacterial taxa) in case samples are converted to
percentiles of the equivalent features in control samples within a study prior to
pooling data across studies. We look at how this percentile-normalization method
compares to traditional meta-analysis methods for combining independent p-
values and to limma and ComBat, widely used batch-correction models
developed for RNA microarray data. Overall, we show that percentile-
normalization is a simple, non-parametric approach for correcting batch effects
and improving sensitivity in case-control meta-analyses.

28
29 **Author Summary**
30
Batch effects are obstacles to comparing results across studies. Traditional
meta-analysis techniques for combining p-values from independent studies, like
Fisher's method, are effective but statistically conservative. If batch-effects can
be corrected, then statistical tests can be performed on data pooled across
studies, increasing sensitivity to detect differences between treatment groups.
Here, we show how a simple, model-free approach corrects for batch effects in
case-control microbiome datasets.
38
Introduction

Data generated by high throughput methods like mass-spectrometry, second-

generation sequencing, or microarrays are sensitive to experimental and

computational processing [1, 2]. This sensitivity gives rise to 'batch effects'

between independent runs of an experiment. Even when different research

groups adhere to the same methodologies, these effects can arise due to slight

differences in hardware, reagents, or personnel [3]. Thus, it is inappropriate to

make direct, quantitative comparisons of uncorrected data across studies.

Several tools for reducing batch effects in RNA microarray data have been

developed. For example, surrogate variable analysis (SVA) estimates a set of

inferred variables (eigenvectors) that explain variance associated with putative

batch effects [4]. These inferred variables are then incorporated into a linear

model to correct downstream significance tests. The limma package employs a

similar linear correction to account for batch effects prior to statistical analysis [5].

SVA and limma are part of a family of linear batch-correction methods that use

different varieties of factor analysis, singular value decomposition, or regression

[4-7]. The most relied upon method to date [8], called ComBat, uses a Bayesian

approach to estimate location and scale parameters for each feature within a

batch [9]. All of these models are most effective when batch effects are not

conflated with the true biological effects [1]. Furthermore, most batch correction

methods make certain parametric assumptions.

Unfortunately, models that often work well for many types of 'omics data

may not be appropriate for microbiome datasets. In microbiome studies, batch

effects are often diffuse and conflated with biological signals [10-12]. The

microbiome field has also struggled with finding appropriate parametric models

for bacterial abundance distributions and for dealing with zeros. This is especially

true for low-biomass samples in microbiome sequencing studies, like samples

taken from the built environment [13], where populations are under-sampled, the

biological signal is relatively weak, and batch effects can be quite large [14]. One

way to get around this issue is to calculate statistics within a given batch, and

then compare significant features across batches using classic meta-analysis

techniques for combining p-values, like Fisher's and Stouffer's methods [15, 16].

These meta-analysis techniques are robust to batch effects across independent

studies, but have less statistical power and ability to detect subtle differences

than directly pooling data across studies.

Here, we describe a model-free data-normalization procedure for

controlling batch effects in case-control microbiome studies that enables pooling

data across studies. Case-control studies include a built-in population of control

samples (e.g. healthy subjects) that can be used to normalize the case samples

(e.g. diseased subjects). For every feature (i.e. bacterial taxon), the case

abundance distributions can be converted to percentiles of the equivalent control

abundance distributions (Fig. 1). Study-specific batch effects present in the case

samples will also be present in the control samples, and by converting the case

data into percentiles of the control distribution these effects are mitigated. Upon

conversion to percentiles of the within-study controls, percentile-normalized

samples from multiple studies with similar case-control definitions can be more

appropriately pooled for statistical testing (Fig. 1). We show that this approach

effectively controls batch effects in microbiome case-control studies and we

compare this method to pooling ComBat- or limma-corrected data, and to

Fisher's and Stouffer's methods for combining independent p-values.

Figure 1. Percentile-normalization procedure converts case and control values into percentiles of the control distribution, which allows for pooling of normalized data across studies. Conceptual plot shows theoretical feature (OTU 1) abundance distributions for control samples and case samples from two independent studies. Converting a control distribution into percentiles of itself naturally gives rise to a uniform distribution (represented by flat blue distributions in central panels), while converting the case distribution into percentiles of the control distribution produces a non-uniform distribution when these two distributions differ (represented by skewed orange distributions in central panels). The right-most panel shows the result of pooling percentile distributions from study 1 and study 2. Percentile-normalization places data from separate studies onto a standardized axis that allows for cross-study comparison. Each simulated case and control distribution was produced by randomly sampling 100 times from a lognormal distribution. Study 1 control parameters: $\mu = 0.1$ and $\sigma = 0.7$. Study 1 case parameters: $\mu = 0.8$ and $\sigma = 0.5$. Study 2 control parameters: $\mu = 1.5$ and $\sigma = 0.2$. Study 2 case parameters: $\mu = 1.75$ and $\sigma = 0.13$.

Methods

Datasets

We used a collection of case-control datasets obtained from the MicrobiomeHD database [17] to validate our batch-normalization method. We focused our analyses on studies spanning five diseases: colorectal cancer (CRC) [18-21], Crohn's Disease (CD) [22-25], Ulcerative Colitis (UC) [23-25], obesity (OB) [18, 26-35], and *Clostridium difficile* induced diarrhea (CDI) [33, 36]. For a subset of three CRC studies [18-20], we were able to obtain sequence data from the same

region of the 16S gene (V4) so that these data could be processed together. The

remaining MicrobiomeHD case-control datasets were previously processed using

the same pipeline (see below), and then Operational Taxonomic Units (OTUs)

were summarized at the genus level for comparison across studies.

Sequence Data Processing

To perform OTU-level analyses across the CRC studies, we downloaded the raw

data from all of the MicrobiomeHD datasets that sequenced the V4 region of the

16S gene. We quality filtered and length trimmed each V4 dataset as described

in Duvallet et al. (2017) and concatenated these raw, trimmed FASTQ files into

one file. We removed any unique sequences that did not appear more than 20

times and clustered the remaining reads with USEARCH [37] at 97% similarity.

We assigned these OTUs taxonomic identifiers using the RDP classifier [38] with

a cutoff of 0.5.

For genus-level analyses, OTU tables and metadata were acquired from

the MicrobiomeHD database (https://doi.org/10.5281/zenodo.569601). Raw data

were downloaded from the original studies and processed through our in-house

16S-processing pipeline

(https://github.com/thomasgurry/amplicon_sequencing_pipeline) as described

previously [17]. Each study's OTU table was converted to relative abundance by

dividing each sample by its total number of reads and collapsed to genus level by

summing all OTUs with the same genus-level annotation.

To plot data in ordination space, Bray-Curtis distances were calculated

from relative abundance data using Scikit-learn

(sklearn.metrics.pairwise.pairwise_distances; metric='braycurtis') [39]. Non-

metric multidimensional scaling (NMDS) coordinates were calculated for two

axes based on Bray-Curtis distances using Scikit-learn (sklearn.manifold.MDS;

n_components=2, metric=False, max_iter=500, eps=1e-12,

dissimilarity='precomputed').

Percentile Normalization

In this procedure, control feature distributions are percentile-normalized against

themselves (resulting in a uniform distribution between 0 and 100) and case

feature distributions are converted into percentiles of their equivalent control

features. Treating our controls as null-hypotheses is motivated by the idea that

healthy patients should be treated as similar across datasets, even though we

understand that they will differ due to biological as well as technical batch effects.

Relative abundance distributions were converted to percentiles using the SciPy v

0.19.0 [40] stats.percentileofscore method (kind='mean'). In order to avoid rank

pile-ups due to the presence of many zeros, we replaced zeros with pseudo

relative abundances drawn from a uniform distribution between 0.0 and 10^{-9} (i.e.

a set of random values smaller than the lowest possible relative abundance in

any dataset). Due to the zero-replacement step, p-values can shift slightly upon

re-analysis with a different random draw, which can lead to the loss or gain of

significance for features very near the significance threshold. Within each study,

control distributions for each individual OTU or genus were converted into

percentiles of themselves and case distributions were converted into percentiles

of their corresponding control distribution. We have written a python script that

performs percentile-normalization given an OTU table, a list of case sample IDs,

and a list of control sample IDs as inputs

(https://github.com/seangibbons/percentile_normalization). A QIIME 2

(https://qiime2.org) plugin for running percentile-normalization is also available

(https://github.com/cduvallet/q2-perc-norm).

ComBat

For each disease, we applied ComBat [8] to the case-control datasets analyzed

in this study. Relative abundances (OTUs in the CRC analysis or OTUs

collapsed to the genus level in the genus-level analysis) were log-transformed

prior to running ComBat (default settings), adding a pseudo relative abundance

of half the minimal frequency (across the entire feature table) to replace zeros.

ComBat-corrected data were then transformed back from log-space (i.e.

exponential transformation) prior to downstream analyses.

limma

In addition to ComBat, we applied a linear batch correction method from the

limma package in R [5]. Relative abundances (zeros replaced with pseudo

relative abundances equal to half the minimal frequency across the entire feature

table) were log-transformed as described above and then a linear model was fit

183 to subtract batch effects using the removeBatchEffect function (default settings).

184 The limma-corrected data were then transformed back from log-space (i.e.

185 exponential transformation) prior to downstream analysis.

187 *Statistical Analysis*

188 To calculate statistical significance, we restricted our statistical tests to

189 OTUs/genera that occurred in at least one third of control or one third of case

190 samples in order to reduce our multi-test correction penalty. We used the

191 Wilcoxon rank-sum test, as implemented in SciPy v0.19.0 (sicipy.stats.ranksums)

192 [40], to determine significant differences between independent groups of

193 samples. Wilcoxon tests were run either within or across studies. In order to

194 calculate statistics across studies, normalized case and control samples from

195 multiple studies of the same disease were combined together into the same OTU

196 table. Hereafter, combining datasets is referred to as 'pooling.' P-values were

197 multiple-test corrected using the Benjamini-Hochberg False Discovery Rate

198 (FDR) procedure, as implemented in StatsModels v 0.8.0

199 (statsmodels.sandbox.stats.multicomp.multipletests) [41]. Differences in overall

200 community structure were assessed using the Permutational Multivariate

201 Analysis of Variance (PERMANOVA) test in R's vegan package [42] as

202 implemented in scikit-bio (skbio.stats.distance.permanova). Fisher's and

203 Stouffer's methods for combining p-values were performed using SciPy v0.19.0

204 (scipy.stats.combine_pvalues; method='fisher' or method='stouffer'). For

205 Stouffer's method, weights for each study were calculated as the square root of

the number of cases plus the number of controls. OTUs/genera with significant

responses in opposing directions across studies were excluded from Fisher and

Stouffer analyses.

in silico experiments

We ran an *in silico* titration experiment using the OTU-level data to

simulate pooling of control samples from different datasets before calculating

significant differences. Healthy samples from one study were mixed with healthy

samples from another study at different proportions prior to calculating significant

differences in OTU frequencies between cases and controls. Case and control

groups were subsampled to 40 samples each. Control samples were substituted

by randomly selected samples from another study along a fractional gradient (0-

100% control samples from another study). We calculated significant differences

between case and control groups using the Wilcoxon rank-sum test and applied

an FDR correction. OTUs with q-values ≤ 0.05 were considered significant. The

titration experiment was rerun 20 times, and the results were averaged.

Similar to the titration experiment, we ran an OTU-level analysis of how

batch-correction methods might impact false-positive rates by randomly selecting

40 control samples from the Baxter et al. (2016) study as artificial 'controls' and

40 control samples from the Zeller et al. (2014) study as artificial 'cases' (across

20 iterations) for each data type (i.e. raw, percentile-normalized, limma-

corrected, and ComBat-corrected). We then calculated significant differences

between these artificial 'case' and 'control' groups as outlined above to generate

p-values for each OTU.

Results

Batch effects at OTU-level resolution

To minimize possible biases across data sets, we identified three colorectal

cancer (CRC) studies that sequenced the same region of the 16S gene (V4). We

reprocessed the raw sequence data from each study in the same quality filtering

and OTU picking pipeline to obtain bioinformatically-standardized results. OTUs

that occurred in at least one third of case or one third of control samples (i.e.

within individual studies) were retained for all downstream statistical analyses.

Despite standardizing the bioinformatic processing of these data, we saw

significant batch effects in healthy patients across studies (PERMANOVA p <

0.001; Fig. 2). The similarity between samples from the Baxter et al. (2016) and

Zackular et al. (2014) studies is due to the fact that they were sourced from the

same patient cohort (although samples were processed separately), making this

comparison a good pseudo-negative control for batch effects [18, 20]. There was

an apparent reduction in the batch effect after applying ComBat, although

differences between batches remained weakly significant (PERMANOVA p =

0.008, Fig. 2) [8]. Due to the non-independence between the Baxter and Zackular

patient cohorts, we removed the smaller of the two studies (Zackular) from all

downstream analyses. Out of a total of 1,021 OTUs that passed our abundance

filter, 681 differed significantly in uncorrected relative abundance between the

Baxter and Zeller healthy controls (FDR q ≤ 0.05).

Figure 2. Batch effects between healthy controls from different studies can
be reduced by ComBat and percentile-normalization. Non-metric
multidimensional scaling (NMDS) plot showing the distribution of healthy controls
from three colorectal cancer studies in ordination space (Bray-Curtis distances of
relative abundance OTU-level data). Despite standardized bioinformatic
processing, healthy patients differed significantly in their gut microbiomes across
studies (PERMANOVA p < 0.001; batch accounts for 6.342% of the total
variance). Studies were still significantly different after applying ComBat, an
established batch-correction method (PERMANOVA p < 0.01). However,
percentile-normalization did a better job of stabilizing the variance across studies
and removed any apparent batch effect (PERMANOVA p > 0.5).
264

We ran an *in silico* titration experiment to simulate pooling of control

samples from different datasets before calculating significant differences. Healthy

samples from one study were mixed with healthy samples from another study at

different proportions prior to calculating significant differences in OTU

frequencies between cases and controls (see conceptual outline in Fig. 3). For

non-normalized data, the number of significant OTUs greatly increased due to

batch effects as more control samples were substituted in from another study.

This result highlights the danger of pooling raw data across batches. ComBat-

and limma-corrected data performed better than uncorrected data, but still

showed many spurious results as the proportion of control samples from another

study increased (Fig. 3). Percentile-normalization showed no increase in

spurious results over the titration gradient (Fig. 3). Although we do see several

significant CRC-associated OTUs in the full dataset (see below), these were not

detected in the titration experiment due to the loss of statistical power when

reducing case and control groups to only 40 samples each.

Figure 3. Pooling non-normalized samples from different studies can give
rise to many spurious associations. The control group from one study is
gradually substituted with randomly chosen control samples from another study
(non-normalized, percentile-normalized, limma-corrected, and ComBat-
corrected), keeping the total number of case and control samples fixed at n=40
(see conceptual illustration on the left). Mixing in non-normalized control samples
from another study gave rise to spurious results due to batch effects (blue lines).
ComBat- and limma-corrected data showed fewer spurious associations (green
and red lines). Percentile-normalization showed no increase in spurious results
along the titration gradient (orange lines).
292

We ran a second *in silico* experiment to determine whether the false-

positive rate was impacted by our different batch-correction methods. We

randomly selected 40 samples from the Baxter healthy controls as artificial

'controls' and 40 samples from the Zeller healthy controls as artificial 'cases' for

each data type (i.e. non-normalized, percentile-normalized, limma-corrected, and

ComBat-corrected) and calculated significant OTU-level differences between

these groups. We repeated this process twenty times to generate a set of p-value

distributions. We found that the fraction of p-values ≤ 0.05 can be as high as

~70% for the non-normalized data (Fig. 4). This result matches with our finding

that 681 out of the 1,021 OTUs in this dataset differed significantly across Baxter

and Zeller controls (q ≤ 0.05). Each normalization technique drastically reduced

the number of false positives, but percentile-normalization gave the best results

(Fig. 4). When low-abundance OTUs were included in the analysis, ComBat and

limma showed highly skewed p-value distributions, giving rise to a larger number

of false positives than the non-normalized data (Fig. S1).

Figure 4. False positive rates are reduced by batch-correction methods.
Random sets of 40 Baxter controls and random sets of 40 Zeller controls were
selected for null case-control comparisons (20 iterations). Smaller points show
the fraction of p-values ≤ 0.05 within a given iteration, while larger dots show the
average value across all 20 iterations. Within each category, smaller points are
randomly jittered along the x-axis for better visualization. The fraction of p-values
≤ 0.05 is highly inflated for non-normalized data (red dashed line shows the null-
expectation for p-values). Only abundant OTUs (detected in at least a third of
case or control samples) were included in this analysis.
318
319

We next assessed the performance each cross-study analysis method by

comparing OTU-level results across two independent CRC datasets. In the

Baxter study, there were 172 healthy (control) samples and 120 CRC (case)

samples, with 14 OTUs (from *Fusobacterium*, *Coprococcus*, *Butyricicoccus*,

Gemmiger, *Faecalibacterium*, *Roseburia*, *Parvimonas*, *Haemophilus*,

Porphyromonas, *Peptostreptococcus*, *Streptophyta*, *Bacteroides* and *Clostridium*

XIVa genera) showing significant differences in abundance between cases and

controls (FDR q ≤ 0.05). For Zeller, there were 71 control and 40 case samples,

with 18 OTUs (from *Butyricicoccus*, *Butyricimonas*, *Fusobacterium*, *Closridium*

XIVa, *Streptococcus*, *Parabacteroides*, *Alistipes*, *Anaerostipes*, *Parvimonas*,

Peptostreptococcus, *Blautia*, *Dialister,* and *Bacteroides* genera) that differed

significantly across cases and controls (FDR q ≤ 0.05).

In the absence of batch effects, pooling data across datasets of the same

disease should increase sensitivity to detect significant cross-study associations.

We pooled percentile-normalized, limma-corrected, and ComBat-corrected data,

respectively, across Baxter and Zeller studies to look for OTUs that differed

significantly across cases and controls. These pooled results were then

compared to classic methods for combining p-values from each dataset's

individual results (above). For the percentile-normalized data, we found 39 OTUs

that differed significantly between cases and controls (FDR q ≤ 0.05), 21 of which

overlapped with the within-study results. The pooled limma-corrected and

ComBat-corrected data resulted in 37 and 36 significant OTUs, respectively. 35

of the OTUs identified as significant by ComBat were also significant in the limma

results. 30 of the limma results and 29 of the ComBat results were also

significant in the percentile-normalization results, respectively. Fisher's method

identified seven significant OTUs from *Clostridium XIVa*, *Streptococcus*,

Fusobacterium, *Parvimonas*, *Peptostreptococcus*, and *Anaerostipes* genera,

which were also found in the percentile-normalized results. Stouffer's method

identified the same seven OTUs found using Fisher's method. Overall, the

pooling methods improve statistical power to detect significant OTUs over

traditional meta-analysis methods. For example, particular OTUs from

Desulfovibrio and *Parabacteroides* genera were identified as significantly

enriched in CRC patients in the pooled results (ComBat, limma, and percentile-

normalized), but not in the within-study results or in the Fisher and Stouffer

results. Pooled analysis of percentile-normalized data also identified on

Enterobacter OTU enriched in cancer patients, two OTUs from the

Lachnospiraceae family that were enriched in controls and one *Lachnospiraceae*

that was enriched in cases, which were missed by the within-study analyses. In

all, 18 OTUs were identified in the pooled, percentile-normalized results that

were missed by the within-study analyses (Fig. 5). These additional taxonomic

associations (e.g. *Desulfovibrio*, *Costridium XIVa*, and *Lachnospiraceae*) are

consistent with prior meta-analyses of CRC microbiome studies [17, 43]. It is

important to visualize the data being fed into statistical tests to determine

whether significant associations are being driven by outlier studies or by other

artifacts. The associations identified in Figure 5 appear to be biologically

meaningful due to the overall consistency of the effect directions across studies.

**Figure 5. OTUs significant across CRC studies, but not within a given
study.** Pooling data provides greater statistical power to detect subtle, yet
consistent differences in OTU abundances across sample groups. 18 OTUs are
labeled by their most resolved taxonomic annotation. Each OTU in this plot was
not found to be significant within either Baxter or Zeller studies, but became
significant after pooling the percentile-normalized datasets (q ≤ 0.05).

Batch effects at genus-level resolution across multiple diseases

In order to assess the performance of different meta-analysis techniques across

a larger set of studies and diseases, we summarized OTU abundances at the

genus level for five diseases across 18 studies - *Clostridium difficile induced*

diarrhea (CDI), Crohn's disease (CD), ulcerative colitis (UC), obesity (OB), and

CRC. There were a total of 306 unique genera detected across studies. There

were two CDI case-control studies: Schubert et al. (2014) had 154 control and 93

case samples [33]; Vincent et al. (2013) had 25 control and 25 case samples

[36]. There were four inflammatory bowel disease (IBD) studies that included CD

patients and three that also included UC patients: Papa et al. (2012) had 24 non-

IBD control samples, 23 CD samples, and 43 UC samples [23]; Morgan et al.

(2012) had 18 control, 61 CD and 47 UC samples [24]; Willing et al. (2010) had

35 control, 16 UC and 29 CD samples [25]; Gevers et al. (2014) had 16 non-IBD

control and 146 CD samples, with no UC samples [22]. There were eleven

studies with lean and obese (OB) cohorts: Turnbaugh et al. (2009) had 33

controls and 102 cases [26]; Goodrich et al. (2014) had 428 controls and 185

cases [30]; Escobar et al. (2014) had 10 controls and 10 cases [27]; Zhu et al.

(2014) had 16 controls and 25 cases [44]; Jumpertz et al. (2011) had 12 controls

and 9 cases [34]; Ross et al. (2015) had 26 controls and 37 cases [29]; Zupancic

et al. (2012) had 96 controls and 101 cases [28]; Baxter et al. (2016) had 125

controls and 47 OB cases [18]; Schubert et al. (2014) had 68 controls and 34 OB

cases [33]; Wu et al. (2011) had 59 controls and 9 cases [32]; and Zeevi et al.

(2015) had 567 controls and 151 cases [31]. There were four independent CRC

studies, including the Baxter and Zeller studies listed in the OTU-level analysis

(see above for sample sizes). The remaining two CRC studies are Wang et al.

(2012), which had 54 control and 44 case samples [45], and Chen et al. (2012),

which had 22 controls and 21 cases [21].

Most genera that differed significantly within a given study were not

significant in other studies of the same disease. For example, of the 36 unique

genera that showed significant differences within any given OB study, none were

found to be significant in all studies (I = 0; Table 1). Indeed, there were no genera

that were significant across all studies in the majority of diseases studied (I = 0;

Table 1). CDI only had two studies, and of the 38 significant results, only six were

shared across both datasets. Overall, few genera were significant within two or

more studies (2N ≤ 6; Table 1).

The number of genera that differed significantly across pooled cases and

controls changed depending on how the data were batch-corrected (Table 1 and

Table S1). For every disease, percentile-normalization yielded the largest

number of significant genera when compared to other methods. Overall,

ComBat- and limma-corrections resulted in many fewer significant genera,

especially for CD, UC, and CRC (Table 1 and Table S1). Half of the IBD (CD and

UC) studies included non-IBD patients with inflammatory symptoms as controls

rather than clinically healthy patients. These biologically relevant differences in

inflammatory symptoms between control cohorts were conflated with batches

and were likely smoothed out by ComBat and limma corrections. Fisher's and

Stouffer's methods consistently identified fewer significant associations than

percentile-normalization (Table 1 and Table S1). Pooling data prior to running a

statistical test is a more sensitive technique than combining independently

calculated p-values [46]. Thus, percentile-normalization increases the statistical

power to detect differences across studies while controlling for false positives

and batch effects.

disease	method	pooled	within
CDI (N=2)	percentile	37	U = 38 / 2N = 6
	Fisher	12	
	Stouffer	12	
	ComBat	36	
	limma	36	
CD (N = 4)	percentile	19	U = 13 / I = 0 / 2N = 2
	Fisher	6	
	Stouffer	6	
	ComBat	2	
	limma	1	
UC (N = 3)	percentile	10	U = 17 / I = 0 / 2N = 1
	Fisher	4	
	Stouffer	4	
	ComBat	5	
	limma	5	
CRC (N = 4)	percentile	12	U = 20 / I = 0 / 2N = 3
	Fisher	9	
	Stouffer	7	
	ComBat	5	
	limma	5	
OB (N = 11)	percentile	18	U = 36 / I = 0 / 2N = 6
	Fisher	4	
	Stouffer	6	
	ComBat	13	
	limma	15	

Table 1. Normalization methods impact the number of significant genus-
level associations between cases and controls across multiple diseases.
Numbers of genera that differ significantly between cases and controls for five
diseases. In the 'disease' column, CDI = *Clostridium difficile* induced diarrhea,
CD = Crohn's Disease, UC = Ulcerative Colitis, CRC = Colorectal Cancer, and
OB = obesity. 'N =' shows the number of studies included in each meta-analysis.
The method column indicates how the data were processed prior to running
significance tests (percentile-normalized, Fisher's method for combining p-
values, Stouffer's method for combining p-values, ComBat-corrected, or limma-
corrected). The significance threshold used was $q \leq 0.05$ (FDR). The 'pooled'
column shows the total number of genera that were found to be significantly
different *across* pooled studies for a given disease. The 'within' column shows
the total number of unique (non-redundant) genera that were identified as
significantly different *within* each study (U = union), the number of genera that
were significant in all individual studies (I = intersection), and the number of

significant genera that were consistently significant in at least two studies (2N;
2N == I for CDI).
445

To better assess how percentile normalization impacted the pooled

results, we looked at genera that were significant within a single-study but not

across studies after pooling. There were 12 genera that were significant within a

subset of CRC studies, but not after pooling (Fig. 6). *Gemmiger*, *Bacteroides*,

and *Roseburia* showed variable responses across studies, sometimes enriched

in controls and other times enriched in cases. The remaining genera showed

weak associations within one or two studies, but did not differ significantly across

studies (i.e. $q \leq 0.05$). These genera that show weak or inconsistent responses

across batches may not be reliable disease biomarkers. However, by including

larger numbers of CRC studies in future meta-analyses it is likely that some of

these genera could pass the significance threshold. Two genera - *Lactobacillus*

and *Desulfovibrio* - were not significantly different between cases and controls

within an individual study, but became significant after pooling (Fig. 7). These

genera showed weak, but largely consistent enrichment in cancer patients and

demonstrate the utility of pooling datasets to detect subtle differences.

Figure 6. Genera that show a significant difference between CRC cases and controls within a given study, but not after pooling. 12 genera showed significant differences between cases and controls within a study (q ≤ 0.05), but not after pooling across CRC studies.

Figure 7. Genera that do not show a significant difference between CRC cases and controls within a given study, but do after pooling. Two genera did not show significant differences between cases and controls within a study, but became significant after pooling across CRC studies (q ≤ 0.05).

While prior work has suggested that there may not be consistent

associations between the gut microbiome and obesity [35], we observed six

genera in a recent meta-analysis that differed significantly across two or more

(out of five) independent obesity studies [17]. Of these six genera, four

(*Roseburia*, *Clostridium IV*, *Oscillobacter*, and *Pseudoflavonifractor*) were also

found to be significant in the pooled, percentile-normalized results (Table S1; Fig.

S2). The two remaining genera not found to be significant in the percentile-

normalized analysis (*Mogibacterium* and *Anaerovorax*) showed highly irregular

responses across the 11 obesity studies analyzed in this study (Fig. S3). Despite

the irregular behavior of these genera, Fisher's and Stouffer's methods both

identified *Mogibacterium* as significantly associated with obesity (Table S1).

Discussion

Batch effects are unavoidable when working with high-throughput data

generation platforms. The RNA microarray community has been proactive in the

development of tools for dealing with these effects [1, 8]. However, these tools

are not as effective when batch effects are confounded with biological signals or

when parametric assumptions do not apply, which is often the case in

microbiome case-control studies. Therefore, model-free methods are needed for

correcting batch effects across microbiome datasets. Fortunately, case-control

studies can be internally normalized by their own control samples. Any study-

specific batch effects in the case samples will be present in the control samples,

and by converting the case data into percentiles of the control distribution these

effects are attenuated without making parametric assumptions.

Relative abundance, limma-corrected, and ComBat-corrected data - but

not percentile-normalized data - quickly yielded a large number of spurious

results when cases from one study were tested against controls from another

(Fig. 3). Additionally, when control populations from different batches were

compared to one another, non-normalized data yielded a much larger number of

false positives than batch-corrected data (Fig. 4). Our percentile-normalization

approach was much more effective than limma and ComBat in controlling false

positives (Fig. 4), especially in the presence of low-abundance taxa (Fig. S1).

Because pooling datasets increases statistical power, it is tempting to pool

these data even in the absence of suitable batch-correction methods.

Consequently, pooling non-normalized data from different batches has been

common practice in the microbiome field [27, 29, 35, 47-55]. In this paper, we

demonstrate why this practice is highly inadvisable. Pooling batch-corrected data

from multiple studies allowed us to detect significant differences that were not

found within a given study (Figs. 5 and 7), while removing associations that were

weak or inconsistent across studies (Fig. 6 and S3). Percentile-normalized

results often identified significant differences between cases and controls that

were missed by other normalization methods (Table 1). For CDI, percentile-

normalized results identified about the same number of significant hits as the

other batch-correction methods (Table 1), which was likely due to the fact that

the biological signal associated with diarrhea is very strong [17]. In cases where

the biological signal is strong, results should be robust to the types of analyses

employed. For UC and CD studies (IBD), percentile-normalization identified

several significant genera that limma and ComBat did not (Table 1). The reduced

number of significant hits from limma- and ComBat-corrected data for IBD was

likely due to heterogeneous control cohorts across these studies (i.e. healthy

patients vs. non-IBD patients), which likely smoothed-out inflammation-

associated signals. This result highlights the importance of having consistent

definitions for case and control cohorts across studies.

We compared percentile-normalization and pooling to Fisher's and

Stouffer's methods for combining independent p-values. Stouffer's method is

similar to Fisher's, but includes weights for each p-value based on the number of

samples in a study. Percentile-normalization consistently identified a larger

number of significant hits than Stouffer's and Fisher's methods, confirming that

pooling data increases sensitivity (i.e. reducing putative false negatives).

Methods for combining p-values from independent studies are quite robust and

should probably be considered as a safe alternative to pooling (i.e. lower chance

of false positives). However, in the case of our obesity analysis, Fisher's and

Stouffer's methods identified *Mogibacterium* as significant despite its apparent

inconsistency across studies (Fig. S3).

In conclusion, we present a robust, model-free procedure for transforming

each feature in a microbiome case-control dataset into percentiles of its control

distribution (Fig. 1). The main conditions for applying this method are that 1) each

batch must have a sizeable number of control samples (i.e. the density of the

control distribution limits the resolution of the percentile-transformation of the

case samples), and 2) case and control populations should be consistently

defined across batches (i.e. same definition of 'healthy' or 'diseased' groups).

Given these caveats, percentile-normalized features can be pooled across

studies for univariate statistical testing (whichever test a researcher prefers -

ideally non-parametric), alleviating the batch effect problem. This model-free

procedure could also be applied to other types of 'omics datasets with

consistently defined internal controls. We find that this procedure allows us to

identify differences between cases and controls that are often missed by more

conservative meta-analysis techniques. Methods developed for batch-correction

in microarray data, like limma and ComBat, can partially reduce batch effects in

microbiome studies (Figs. 2-4), but appear to obscure real patterns if batch

effects are not independent of biological signals or if the parametric assumptions

of these models are not valid. We suggest that methods like limma and ComBat

are useful for studies lacking case and control groups. However, when studies

have consistently defined internal controls, percentile-normalization should be

the preferred batch correction approach. Future work should focus on developing

parametric models specifically for batch correction in microbiome datasets, which

could further improve sensitivity to detect subtle biological differences across

studies.

Acknowledgements

We thank Sean Kearney and members of the Alm and Irizarry labs for helpful

feedback and Marc Sze and Patrick Schloss for providing online code to

download the Wu et al. (2011) and Zeevi et al. (2015) obesity datasets.

References

1.   Leek JT, Scharpf RB, Bravo HC, Simcha D, Langmead B, Johnson WE, et
al. Tackling the widespread and critical impact of batch effects in high-throughput
data. Nat Rev Genet. 2010;11(10):733-9.

2.   Goh WWB, Wang W, Wong L. Why batch effects matter in omics data,
and how to avoid them. Trends Biotechnol. 2017;35(6):498-507.
3.   Schloss PD, Gevers D, Westcott SL. Reducing the Effects of PCR
Amplification and Sequencing Artifacts on 16S rRNA-Based Studies. PLoS One.
2011;6(12):e27310. doi: 10.1371/journal.pone.0027310.
4.   Leek JT, Storey JD. Capturing heterogeneity in gene expression studies
by surrogate variable analysis. PLoS Genet. 2007;3(9):e161.
5.   Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, et al. limma powers
differential expression analyses for RNA-sequencing and microarray studies.
Nucleic Acids Res. 2015;43(7):e47-e.
6.   Alter O, Brown PO, Botstein D. Singular value decomposition for genome-
wide expression data processing and modeling. Proc Natl Acad Sci USA.
2000;97(18):10101-6.
7.   Benito M, Parker J, Du Q, Wu J, Xiang D, Perou CM, et al. Adjustment of
systematic microarray data biases. Bioinformatics. 2004;20(1):105-14.
8.   Chen C, Grennan K, Badner J, Zhang D, Gershon E, Jin L, et al.
Removing Batch Effects in Analysis of Expression Microarray Data: An
Evaluation of Six Batch Adjustment Methods. PLoS One. 2011;6(2):e17238. doi:
10.1371/journal.pone.0017238.
9.   Johnson WE, Li C, Rabinovic A. Adjusting batch effects in microarray
expression data using empirical Bayes methods. Biostatistics. 2007;8(1):118-27.
10.   Weiss S, Amir A, Hyde ER, Metcalf JL, Song SJ, Knight R. Tracking down
the sources of experimental contamination in microbiome studies. Genome Biol.
2014;15(12):564. doi: 10.1186/s13059-014-0564-2.
11.   Shen H, Rogelj S, Kieft TL. Sensitive, real-time PCR detects low-levels of
contamination by Legionella pneumophila in commercial reagents. Mol Cell
Probes. 2006;20. doi: 10.1016/j.mcp.2005.09.007.
12.   Nguyen NH, Smith D, Peay K, Kennedy P. Parsing ecological signal from
noise in next generation amplicon sequencing. New Phytol. 2015;205(4):1389-
93. doi: 10.1111/nph.12923.
13.   Gibbons SM. The Built Environment Is a Microbial Wasteland. mSystems.
2016;1(2):e00033-16.
14.   Chase J, Fouquier J, Zare M, Sonderegger DL, Knight R, Kelley ST, et al.
Geography and Location Are the Primary Drivers of Office Microbiome
Composition. mSystems. 2016;1(2). doi: 10.1128/mSystems.00022-16.
15.   Fisher RA. Statistical methods for research workers: Genesis Publishing
Pvt Ltd; 1925.
16.   Stouffer SA. Adjustment during army life: Princeton University Press;
1949.
17.   Duvallet C, Gibbons SM, Gurry T, Irizarry RA, Alm EJ. Meta-analysis of
gut microbiome studies identifies disease-specific and shared responses. Nat
Comm. 2017;8(1):1784.
18.   Baxter NT, Ruffin MT, Rogers MA, Schloss PD. Microbiota-based model
improves the sensitivity of fecal immunochemical test for detecting colonic
lesions. Genom Med. 2016;8(1):37.

19.   Zeller G, Tap J, Voigt AY, Sunagawa S, Kultima JR, Costea PI, et al.
Potential of fecal microbiota for early-stage detection of colorectal cancer. Mol
Sys Biol. 2014;10(11):766.
20.   Zackular JP, Rogers MA, Ruffin MT, Schloss PD. The human gut
microbiome as a screening tool for colorectal cancer. Cancer Prev Res.
2014;7(11):1112-21.
21.   Chen W, Liu F, Ling Z, Tong X, Xiang C. Human intestinal lumen and
mucosa-associated microbiota in patients with colorectal cancer. PLoS One.
2012;7(6):e39743.
22.   Gevers D, Kugathasan S, Denson LA, Vázquez-Baeza Y, Van Treuren W,
Ren B, et al. The treatment-naive microbiome in new-onset Crohn's disease. Cell
Host Microbe. 2014;15(3):382-92.
23.   Papa E, Docktor M, Smillie C, Weber S, Preheim SP, Gevers D, et al.
Non-invasive mapping of the gastrointestinal microbiota identifies children with
inflammatory bowel disease. PLoS One. 2012;7(6):e39242.
24.   Morgan XC, Tickle TL, Sokol H, Gevers D, Devaney KL, Ward DV, et al.
Dysfunction of the intestinal microbiome in inflammatory bowel disease and
treatment. Genome Biol. 2012;13(9):R79.
25.   Willing BP, Dicksved J, Halfvarson J, Andersson AF, Lucio M, Zheng Z, et
al. A pyrosequencing study in twins shows that gastrointestinal microbial profiles
vary with inflammatory bowel disease phenotypes. Gastroenterology.
2010;139(6):1844-54. e1.
26.   Turnbaugh PJ, Hamady M, Yatsunenko T, Cantarel BL, Duncan A, Ley
RE, et al. A core gut microbiome in obese and lean twins. Nature.
2009;457(7228):480-4.
27.   Escobar JS, Klotz B, Valdes BE, Agudelo GM. The gut microbiota of
Colombians differs from that of Americans, Europeans and Asians. BMC
Microbiol. 2014;14(1):311.
28.   Zupancic ML, Cantarel BL, Liu Z, Drabek EF, Ryan KA, Cirimotich S, et al.
Analysis of the gut microbiota in the old order Amish and its relation to the
metabolic syndrome. PLoS One. 2012;7(8):e43052.
29.   Ross MC, Muzny DM, McCormick JB, Gibbs RA, Fisher-Hoch SP,
Petrosino JF. 16S gut community of the Cameron County Hispanic Cohort.
Microbiome. 2015;3(1):7.
30.   Goodrich JK, Waters JL, Poole AC, Sutter JL, Koren O, Blekhman R, et al.
Human genetics shape the gut microbiome. Cell. 2014;159(4):789-99.
31.   Zeevi D, Korem T, Zmora N, Israeli D, Rothschild D, Weinberger A, et al.
Personalized nutrition by prediction of glycemic responses. Cell.
2015;163(5):1079-94.
32.   Wu GD, Chen J, Hoffmann C, Bittinger K, Chen Y-Y, Keilbaugh SA, et al.
Linking long-term dietary patterns with gut microbial enterotypes. Science.
2011;334(6052):105-8.
33.   Schubert AM, Rogers MA, Ring C, Mogle J, Petrosino JP, Young VB, et
al. Microbiome data distinguish patients with Clostridium difficile infection and
non-C. difficile-associated diarrhea from healthy controls. mBio.
2014;5(3):e01021-14.

34. Jumpertz R, Le DS, Turnbaugh PJ, Trinidad C, Bogardus C, Gordon JI, et
al. Energy-balance studies reveal associations between gut microbes, caloric
load, and nutrient absorption in humans. Am J Clin Nutr. 2011;94(1):58-65.
35. Sze MA, Schloss PD. Looking for a Signal in the Noise: Revisiting Obesity
and the Microbiome. MBio. 2016;7(4):e01018-16.
36. Vincent C, Stephens DA, Loo VG, Edens TJ, Behr MA, Dewar K, et al.
Reductions in intestinal Clostridiales precede the development of nosocomial
Clostridium difficile infection. Microbiome. 2013;1(1):18.
37. Edgar RC. Search and clustering orders of magnitude faster than BLAST.
Bioinformatics. 2010;26(19):2460-1.
38. Wang Q, Garrity GM, Tiedje JM, Cole JR. Naïve Bayesian Classifier for
Rapid Assignment of rRNA Sequences into the New Bacterial Taxonomy. Appl
Environ Microbiol. 2007;73(16):5261-7. doi: 10.1128/aem.00062-07.
39. Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O, et
al. Scikit-learn: Machine learning in Python. J Mach Learning Res.
2011;12(Oct):2825-30.
40. Jones E, Oliphant T, Peterson P. SciPy: Open source scientific tools for
Python, 2001–. URL http://www.scipy org. 2007;73:86.
41. Seabold S, Perktold J, editors. Statsmodels: Econometric and statistical
modeling with python. Proceedings of the 9th Python in Science Conference;
2010.
42. Oksanen J. Multivariate analysis of ecological communities in R: vegan
tutorial. R package version. 2011;1(7).
43. Tjalsma H, Boleij A, Marchesi JR, Dutilh BE. A bacterial driver–passenger
model for colorectal cancer: beyond the usual suspects. Nat Rev Microbiol.
2012;10(8):575-82.
44. Zhu L, Baker RD, Baker SS. Gut microbiome and nonalcoholic fatty liver
diseases. Pediatric Res. 2014;77(1-2):245-51.
45. Wang T, Cai G, Qiu Y, Fei N, Zhang M, Pang X, et al. Structural
segregation of gut microbiota between colorectal cancer patients and healthy
volunteers. ISME J. 2012;6(2):320-9.
46. Glass GV. Primary, secondary, and meta-analysis of research. Educ Res.
1976;5(10):3-8.
47. Oliveira FS, Brestelli J, Cade S, Zheng J, Iodice J, Fischer S, et al.
MicrobiomeDB: a systems biology platform for integrating, mining and analyzing
microbiome experiments. Nucleic Acids Res. 2017;46(D1):D684-D91.
48. Huse SM, Dethlefsen L, Huber JA, Welch DM, Relman DA, Sogin ML.
Exploring microbial diversity and taxonomy using SSU rRNA hypervariable tag
sequencing. PLoS Genet. 2008;4(11):e1000255.
49. Koren O, Knights D, Gonzalez A, Waldron L, Segata N, Knight R, et al. A
guide to enterotypes across the human body: meta-analysis of microbial
community structures in human microbiome datasets. PLoS Comput Biol.
2013;9(1):e1002863.
50. Krych L, Hansen CH, Hansen AK, Van Den Berg FW, Nielsen DS.
Quantitatively different, yet qualitatively alike: a meta-analysis of the mouse core

gut microbiome with a view towards the human gut microbiome. PLoS One.
2013;8(5):e62578.
51.   Lozupone CA, Li M, Campbell TB, Flores SC, Linderman D, Gebert MJ, et
al. Alterations in the gut microbiota associated with HIV-1 infection. Cell Host
Microbe. 2013;14(3):329-39.
52.   Lozupone CA, Stombaugh J, Gonzalez A, Ackermann G, Wendel D,
Vazquez-Baeza Y, et al. Meta-analyses of studies of the human microbiota.
Genome Res. 2013;23. doi: 10.1101/gr.151803.112.

53.   Walters WA, Xu Z, Knight R. Meta-analyses of human gut microbes
associated with obesity and IBD. FEBS Lett. 2014;588(22):4223-33.
54.   Pasolli E, Truong DT, Malik F, Waldron L, Segata N. Machine learning
meta-analysis of large metagenomic datasets: tools and biological insights. PLoS
Comput Biol. 2016;12(7):e1004977.
55.   Vogtmann E, Hua X, Zeller G, Sunagawa S, Voigt AY, Hercog R, et al.
Colorectal cancer and the human gut microbiome: reproducibility with whole-
genome shotgun sequencing. PLoS One. 2016;11(5):e0155362.

Supplemental Materials

Table S1. Excel file containing tabs for each disease analyzed in this study: each

tab contains information on significant genera from Table 1.

Figure S1. False positive rates are reduced by percentile-normalization, but
not by ComBat or limma, in the presence of low-abundance OTUs. Random
sets of 40 Baxter controls and random sets of 40 Zeller controls were selected for
null case-control comparisons (20 iterations). Smaller points show the fraction of
p-values ≤ 0.05 within a given iteration, while larger dots show the average value
across all 20 iterations. Within each category, smaller points are randomly jittered
along the x-axis for better visualization. The fraction of p-values ≤ 0.05 is highly
inflated for all methods except percentile-normalization (red dashed line shows
the null-expectation for p-values). All OTUs were included in this analysis (i.e. no
abundance filter prior to running tests). ComBat and limma show highly skewed
p-value distributions when including low-abundance OTUs.
753

Fig. S2. 18 genera significantly different (q ≤ 0.05) between obese and healthy patients based on pooled percentile-normalized analysis (from Table 1). Gray points show data pooled from all 11 obesity studies. Other colors show individual obesity studies. Studies listed in order from left to right: Zhu et al. (2014), Zeevi et al. (2015), Wu et al. (2011), Baxter et al. (2016), Schubert et al. (2014), Zupancic et al. (2012), Ross et al. (2015), Jumpertz et al. (2011), Turnbaugh et al. (2009), Goodrich et al. (2014), Escobar et al. (2014). X symbols indicate the average control percentiles, while + symbols indicate the average case percentiles.

778
779
Fig. S3. Two genera that were significant (q ≤ 0.05) in at least two obesity
studies in Duvallet et al. (2017), but were not significant in the pooled,
percentile-normalized analysis. Gray points show data pooled from all 11
obesity studies. Other colors show individual obesity studies. Studies listed in
order from left to right: Zhu et al. (2014), Zeevi et al. (2015), Wu et al. (2011),
Baxter et al. (2016), Schubert et al. (2014), Zupancic et al. (2012), Ross et al.
(2015), Jumpertz et al. (2011), Turnbaugh et al. (2009), Goodrich et al. (2014),
Escobar et al. (2014). X symbols indicate the average control percentiles, while +
symbols indicate the average case percentiles.
789