Fantastic beasts and how to sequence them:
genomic approaches for obscure model organisms

Mikhail V. Matz

Department of Integrative Biology, University of Texas at Austin, Austin, TX 78712, USA

matz@utexas.edu

Summary:

Application of genomic approaches to "obscure model organisms" (OMOs), meaning species with little or no genomic resources, enables increasingly sophisticated studies of genomic basis of evolution, acclimatization and adaptation in real ecological contexts. Here, I highlight sequencing solutions and data handling techniques most suited for genomic analysis of OMOs.

Glossary:

- **Allele Frequency Spectrum, AFS** (same as Site Frequency Spectrum, SFS): histogram of the number of segregating variants depending on their frequency in one or more populations.
- **Restriction site-Associated DNA (RAD) sequencing**: family of diverse genotyping methods that sequence short fragments of the genome adjacent to recognition site(s) for specific restriction endonuclease(s).
- **Linkage Disequilibrium (LD)**: in this review, correlation of genotypes at a pair of markers across individuals.
- **LD block**: typical distance between markers in the genome across which their genotypes remain correlated.
- **Genome scan:** profiling of genotypes along the genome looking for unusual patterns. Often used to look for signatures of natural selection or introgression.
- **"Denser-than-LD" genotyping**: genotyping of several polymorphic markers per LD block.
- **Highly contiguous reference**: genome or transcriptome reference sequence containing the least amount of fragmentation.
- **Phased data**: data showing which SNP alleles belong to the same homologous chromosome copy.
- **Cross-tissue gene expression analysis**: looking for individual-specific shifts in gene expression detectable across multiple tissues. Such shifts are predominantly genetic in nature.

The focus of this review is mainly on the type of sequencing data required and how to obtain it in the most cost-efficient way rather than on analytical approaches. That said, I could not help but mention some highly promising analytical methods that are not yet broadly adopted by OMO researchers, such as demographic inference based on allele frequency spectra and annotation-independent analyses of gene expression data.

I will start with the summary of general types of questions in OMO studies and corresponding
data types required. We might be interested in the following four layers of genomic information,
each requiring a specific type of experimental and reference data:

1. Genome-wide patterns of neutral variation. This data can elucidate population structure,
population sizes, and migration rates, as well as changes of these parameters through time. This
analysis benefits from high quality genotype calls but does not require dense genome coverage; it
can even be performed in the absence of reference genome.

2. Regions in the genome particularly affected by non-drift processes (natural selection,
introgression, etc). This type of analysis, typically referred to as "genome scanning", takes
genome-wide neutral variation as baseline and looks for regions in the genome exhibiting highly
dissimilar patterns. It requires "denser-than-LD" genotyping and a highly contiguous reference
(see Glossary) to make sure no signal is overlooked.

3. Genome-wide gene expression, an extremely information-rich resource reflecting both
environmental and genetic variation. Streamlined transcript counting methods represent a cost-
efficient alternative to the industry-standard RNA-seq for generating quantitative data. Analysis
of gene expression does not require a genome reference, although a transcriptome reference must
be generated at some point. The reference does not have to be highly contiguous.

4. Epigenetics, here limited to DNA methylation. A variety of methods have been recently
developed that can generate data for DNA methylation analysis. For vertebrates, genome
reference is needed, but for other animals or plants, in which DNA methylation is much less
prevalent and predominantly occurs in exons, transcriptome or exome presents a good cost-
efficient alternative. The reference does not have to be highly contiguous.

Genome-wide neutral variation

Allele Frequency Spectrum analysis

Neutral genetic markers are traditionally analyzed using summaries of allele frequency
differences between populations, such as F_{ST}. The large amount of markers accessible through
next-generation sequencing opened up the possibility to dramatically enhance this approach by
modeling the evolution of the whole allele frequency spectrum (AFS, see Glossary). AFS
represents a rich source of information to fit alternative models with time-resolved population
sizes and migration rates as parameters (Box 1) based on coalescent simulations (*fastsimcoal2*,
[1]), diffusion approximation (*dadi*, [2]), or ordinary differential equations (*moments*, [3]). Model
selection is then based on likelihood ratio tests or Akaike information criterion. The new *moments*
method is particularly promising, as it is substantially faster than its predecessors and includes
built-in bootstrap, demographic model plotting, and capacity to analyze up to five populations
simultaneously. It is also very helpful that *moments* inherits the python code structure well
familiar to *dadi* practitioners.

Experimental data

The data required for AFS analysis is several thousand biallelic neutral single nucleotide
polymorphisms (SNPs). Ideally, SNPs must not be closely physically linked in the genome to
represent independent data points, although it is fully appropriate to analyze linked SNPs with
AFS methods. The lack of requirement for contiguous SNP coverage makes various flavors of
restriction site-associated DNA (RAD) sequencing (recently reviewed in [4,5], see Glossary) well
suited for this analysis. In our experience, *dadi* [2] and *moments* [3] work robustly with 5-10
thousand SNPs (a typical RAD output) when analyzing populations individually or in pairs.
Fitting models with three (*dadi*) or more (*moments, fastsimcoal2*) populations might be
problematic with this relatively low number of SNPs but is usually not required for OMOs (Box
1). Recent population size changes are often of special interest in OMOs; since they
predominantly affect rare alleles, their robust detection requires 20 or more high-quality
genotypes per population [6]. This preference for more individuals rather than more SNPs per
individual is an additional factor that makes cost-efficient RAD the approach of choice for AFS-
based analysis. That said, relatively low number of independent (unlinked) SNPs generated by
some RAD protocols might limit the power of the AFS analysis, and a good subject for a future
study would be the effect of the number of unlinked SNPs on AFS model selection and
uncertainties of parameter estimates. In this regard it is worth noting that RAD flavors differ
considerably in the number of unlinked loci in the genome that they interrogate [4,5].

For demographic inference, the AFS data must be filtered to exclude potential sites under
selection. Whichever test is used to identify such sites (for example, Bayescan, [7]), for their
removal the false discovery rate should be set as high as 0.5 to ensure purging of the majority of
non-neutral sites. Although under this setting half of the removed sites would be neutral, their
removal will not affect the overall AFS as long as the removed fraction does not comprise more
than 1-2% of the total number of sites.

Genotyping quality

In diploids, the most common genotyping error is missing one of the alleles in a heterozygote (i.e.,
a false homozygote call); and the next most common error is missing the whole SNP locus
entirely. Both these "missing data" errors are due to insufficient sequencing coverage, the
problem that is pervasive in today's OMO studies. Such errors strongly affect AFS in the region
of rare alleles, which is unfortunate since rare alleles are the most informative about recent
population history [6,8]. A telltale sign of poor heterozygote calling is under-representation of
singletons, but frequencies of doubletons and higher-order frequency bins are also distorted,
which has strong effect on AFS itself and inferred demographic parameters until mean
sequencing coverage approaches ~10x [9]. When coverage is 10x or higher a good way to filter
data is to select SNPs genotyped in >90-95% of samples [10]; importantly for RAD approach,
this would select SNPs that are unlikely to be affected by null alleles due to mutation in the
restriction endonuclease recognition site [4]. For obvious reasons, for AFS analysis genotype
calls should never be quality-filtered based on allele frequencies (for example, retaining only

variants that are detected in a minimum of two individuals or requiring minor allele frequency to
exceed some cutoff). A robust empirical way to evaluate the consistency of genotype calls is to
compare results for independently processed biological samples of the same genotype [11]. Such
genotyping replicates are quite feasible in RAD and are also useful to identify true SNPs for
training variant quality score recalibration model of the GATK pipeline [12]. For low-coverage
data (<10x), a general solution is provided by the *ANGSD* package [13], which generates AFS as
well as other population genetic statistics based on genotype likelihoods without actually calling
genotypes [14]. This method generates unbiased single-population AFS even with 2x coverage
[9]. Still, there is a concern that high variation in coverage across samples and populations might
affect *ANGSD* statistics; to avoid this potential issue it is recommended to discard the lowest-
coverage outliers and down-sample reads from highest-coverage outliers (J. Ross-Ibarra, pers.
comm.).
144
145 *PCR duplicates*
146
Presence of PCR duplicates in many early RAD applications has been repeatedly highlighted as a
source of genotyping errors [4,15] due to induced over-dispersion of read counts among alleles
and loci. Interestingly, the proportion of PCR duplicates does not depend on the number of PCR
cycles performed during library preparation. Instead, it depends on the ratio between the number
of reads sequenced (N_r) and the number of unique fragments present in the sample prior to PCR
(N_o): the fraction of duplicates is the same as expected when sampling N_r from N_o with
replacement. Fortunately, PCR duplicates are easy to identify and remove using degenerate tags
ligated to RAD fragments prior to amplification [16]. Most present-day RAD protocols now
implement this simple deduplication procedure, including the current version of 2bRAD [11].
156
157 *Genome reference for AFS analysis*
158
A great advantage of RAD-based AFS analysis for OMOs is that SNPs can be called based on
RAD reads themselves, without the need for genome reference. Several *de novo* RAD genotyping
pipelines have been developed, such as STACKS, pyRAD, and UNEAK (see references in [4])
that work for most RAD flavors, plus a similarly structured pipeline for 2bRAD
(https://github.com/z0on/2bRAD_denovo) that takes into account the fact that in 2bRAD either
strand of the locus can be sequenced. Still, using a reference genome to call RAD genotypes
provides three important advantages. First, it identifies physically linked (and thus potentially
non-independent) groups of SNPs, to be resampled as units during AFS bootstrap. The second
advantage is particularly important for OMOs sampled in the field: mapping to reference genome
automatically discards reads from contaminant DNA sources (viruses, bacteria, ingested food,
symbionts etc). To be able to discard such contaminants in *de novo* RAD pipeline the experiment
must include at least one sample generated from a clean source and consider only the RAD loci
observed in that sample.
172
The third advantage of reference-based genotyping is the possibility to discriminate between
ancestral and derived SNP alleles, to attain the best power of AFS-based inference. Counter-
intuitively, the best reference for AFS analysis is not a genome of the species under investigation

but a genome of a related outgroup species, separated from the focal one by a few million years
of evolution, because the SNP state as in the outgroup can be assumed to represent the ancestral
state (e.g., [17]). If the reads are mapped to the same-species genome, to identify ancestral states
of the variants a single well-sequenced RAD sample of an outgroup taxon could be included. The
analysis will then be limited to sites that can be successfully genotyped both in ingroup and
outgroup; in effect, the result is going to be the same as when mapping the reads from whole
project to an outgroup's genome. Although some proportion of ancestral states will be
misidentified due to incomplete lineage sorting, convergence or technical artifacts, this error is
easy to account for by including a single additional parameter into the model, specifying the
proportion of the AFS that needs to be flipped when predicting the data (e.g., [18]). The reference
for AFS does not have to be highly contiguous; the contigs should be just long enough to cover a
typical LD block for meaningful bootstrapping.
188
189   **Genome scanning**
190
Since outlier regions by definition occupy only a small portion of the genome and typically do not
form a single cluster, their confident detection requires "denser-than-LD" genotyping (see
Glossary). It has been argued that in most situations, RAD-like approaches would sample the
genome too sparsely to satisfy this requirement [19,20]. Although many successful genome scans
based on RAD have been published [21], RAD cannot be recommended for genome scanning
since it inevitably leaves considerable fraction of the genome unexplored. Even when LD is
known to be extensive enough for RAD to produce "denser-than-LD" genotyping, a better
solution might be to take full advantage of the extended LD and go instead for low coverage
whole-genome sequencing (WGS) followed by imputation, to obtain full-genome phased data
(Table 1).
201
The types of sequencing approaches for genome scanning with their pros and cons are
summarized in Table 1. Importantly, all of them require highly accurate reads mapped to a
reference for confident SNP detection, making short Illumina reads the genotyping data type of
choice. Some of the very promising approaches that have not yet been fully adopted for OMOs
are exome-seq and ultra-low whole genome sequencing (WGS) with imputation. Exome-seq used
to be a prerogative of model organisms because of the need for exome-capture platform
development, but it has recently been shown that OMO exome can be captured just as efficiently
using bead-bound normalized cDNA obtained from the OMO itself (EecSeq Puritz 2017). Such
"home-made exome" sequencing could become an excellent alternative to RAD since it would
interrogate essentially all the interpretable genetic variation for a comparably low cost. Ultra-low
WGS with imputation used to require extensive reference haplotype panels available only for
well-established model organisms. However, several methods have been recently developed
(most notably STITCH, [22]) that can impute phased genotypes and correct genotyping errors in
ultra-low coverage data without relying on reference panels. Still, their applicability for each new
OMO must be experimentally confirmed because the success of imputation critically depends on
multiple polymorphisms occurring within a typical LD block, and whether this is so is not known
for OMOs *a priori*. Demographic events such as strong recent bottleneck, domestication, or
recent colonization would make imputation more efficient because of more extensive LD and

small number of founding haplotypes [22], and conversely, in large outbred populations
imputation will be less accurate and might require sequencing of a very large number (thousands)
of individuals. The accuracy of imputation can be evaluated by sequencing a few individuals at
high coverage (>10x) to generate high-confidence genotype calls and then attempting to impute
them based on sub-sampled read sets to emulate low coverage. It must be noted that it is
inappropriate to measure imputation accuracy by imputing genotype calls masked in high-quality
datasets (as in, for example, [23]): masked data do not contain false homozygote calls and
therefore do not correctly represent the real-life situation.
228
229 **Gene expression**
230
231   There are many aspects to gene expression, of which I here focus on just one: abundance
232 or protein-coding (polyadenylated) transcripts. The reason is that transcript abundance is by far
233 the most interpretable and it can be very easily analyzed in OMOs.
234
235 *Counting transcripts instead of resequencing them*
236
Typical RNA-seq [24] resequences the whole transcriptome in each sample, but there is a
much more economic way to count abundances of protein-coding transcripts: sequence just a
single fragment per each transcript molecule and count reads corresponding to each gene. TagSeq
[25], for example, sequences a single randomly generated fragment near the 3'-end of the
transcript, which is the most economic use of sequencing effort and removes bias towards longer
transcripts. In a recent benchmarking study TagSeq was actually more accurate than the standard
RNA-seq in measuring transcript abundances, despite nearly tenfold lower cost [26]. More
recently introduced QuantSeq [27] is conceptually similar to TagsSeq: it also sequences a single
randomly generated fragment near the 3'-end of each transcript but has a different library
preparation procedure, implemented as a kit from Lexogen (https://www.lexogen.com/quantseq-
3mrna-sequencing/). Bioinformatics analysis for both TagSeq ad QuantSeq is highly simplified
compared to typical RNA-seq. TagSeq was originally designed for OMOs and so its pipeline uses
transcriptome rather than genome as a reference to attribute reads to genes
(https://github.com/z0on/tag-based_RNAseq). One notable feature of the current version of
TagSeq pipeline is that it includes removal of PCR duplicates based on adaptor-derived
degenerate tags [11], similarly to 2bRAD and for the same reason – to avoid PCR-associated
over-dispersion or read counts.
254
255 *Analysis of gene expression "beyond gene lists"*
256
The unfortunate tradition that OMO research inherits from the biomedical field is putting
too much emphasis on possible functional implications of expression changes of specific genes.
For OMOs, this is bound to remain inconclusive because gene annotations are often absent,
tentative or based predominantly on similarity to human genes, which may or may not serve the
same function in the OMO. Even greater problem is interpretation bias: too often researchers
focus primarily on genes that "make sense" and ignore the rest. This leads to conclusions

reflecting predominantly the researchers' idea of what *should* be going on rather than what is actually happening.

Table 2 lists alternative ways of objective analysis of gene expression data that are enabled by the large sample sizes feasible with TagSeq or QuantSeq. They either do not require gene annotations or rely sufficiently general functional summaries to be robust to occasional missing or mis-annotations. Particularly useful for OMOs are analyses that use gene expression patterns as anonymous multivariate readouts to compare and classify samples, such as principal coordinate analysis (PCoA) or differential analysis of principal components (DAPC). Related multivariate analyses to visualize and classify genome-wide gene expression data, recently reviewed in [28], have become the mainstream tool of single-cell RNA-seq, where they are used to discover cell types and quantify differences between them. With appropriate experimental design, in OMOs these analyses can lead to much more definitive biological conclusions than studies scrutinizing long lists of differentially expressed genes passing a certain significance cutoff.

Gene expression as functional summary of genotype

Gene expression is best known for its context-dependence reflecting phenotypic plasticity, which is the view inherited from biomedical research dealing with genetically uniform models. In natural populations, one of the most important sources of gene expression variation is genetic difference among individuals, manifested as context-*in*dependent, individual-specific deviations in gene expression. This is easy to demonstrate in clonally replicated organisms such as corals. In two reciprocal transplantation experiments performed on different coral species in different oceans, stable between-genotype differences accounted for more than 50% of total gene expression variation despite transplantation of clonal fragments for up to a year to highly dissimilar sites [29,30]. In non-clonal model organisms such as mice or humans, the best demonstration of the effect of genetic variation on gene expression are abundant differences in expression between alleles of the same gene [31,32]. In humans, fixed between-population differences are exemplified by hundreds of genes that are differentially expressed between African and European Americans [33]. All this suggests that gene expression can be a proxy of not only phenotypic plasticity and acclimatization, but of genetic variation and adaptation. A major advantage of the use of gene expression for these types of studies is that gene expression integrates over many functionally relevant variants in the genome and thus represents a condensed functional summary of the genotype [34].

In humans, nearly half of all genetic variants affecting gene expression have detectable effects in all tissues [32], and so one feasible way to separate genotype-specific gene expression from context-dependent variation might be to perform "cross-tissue" comparison (see Glossary) to isolate body-wide expression shifts [35]. In the coming years, cross-tissue or similar analysis is likely to become a major approach to study functional genetic variation in natural populations.

Epigenetics

Among many covalent chromatin modifications I will discuss DNA methylation since it currently receives the most attention in OMOs. Still it must be mentioned that in plants histone methylation appears be no less and perhaps even more involved in acclimatization and transgenerational plasticity [36]. While vertebrates show high methylation throughout the genome, invertebrates and plants methylate their genomes sparsely and mostly in protein-coding regions (so-called gene body methylation, GBM, [37]). The function of his ubiquitous and evolutionarily ancient DNA modification remains unclear [38,39] and the greatest challenge in the next few years will be to decipher it. The most important questions are: (i) Does GBM affect gene expression? (ii) Can it be modified on ecological timescale, to achieve acclimatization to a novel environment? (iii) Can acquired changes in GBM be transmitted across generations? If the answers to all three questions are "yes", then we have a mechanism for transgenerational inheritance of acquired traits, which is an exciting (albeit tentative, [40]) possibility. Table 3 summarizes the methods for generating DNA methylation data. If every gene in the genome has to be interrogated, MBD-seq and meDIP provide the best resolution for sequencing effort [38]. If the goal of the study is to characterize general methylation patterns rather than identify specific genes, highly cost-efficient solutions are provided by RRBS-seq and methylRAD. For studies requiring single base resolution, the best approach appears to be direct detection by PacBio or ONT – however, these exciting developments still require validation in complex genomes.

Generating a reference sequence

For all approaches described here, the accuracy of the reference sequence in terms of per-base error rate must only be high enough to allow unambiguous mapping of high-accuracy (Illumina) reads. The gold standard of genome sequence quality, Q30 or 99.9% accuracy, would not provide any benefit compared to a rough draft accuracy of 99%. Occasional errors in the reference would manifest themselves as SNPs that are not polymorphic in the analyzed samples and therefore irrelevant for analysis. This is the same reason why it is possible to use a genome of a related species as a reference.

For AFS analysis, which does not require highly contiguous reference, even a rough genome draft that can be assembled from a single lane worth of 150b paired-end reads from Illumina HiSeq would be suitable. However, substantially better options are now becoming available for a comparably low price tag. The technology offered by 10x Genomics [41] attaches specific barcodes to short reads originating from the same long DNA fragment, which allows assembling Illumina HiSeq data into very long haplotypes. The two single-molecule long-read "third-generation sequencing" methods, Single Molecule Real Time (SMRT) sequencing by PacBio and nanopore sequencing by ONT, produce reads with broad length distribution, including exceedingly long ones (tens to hundreds of kilobases) resulting in a qualitatively more contiguous genome assemblies [42–45] (Table 4, see [43] for recent benchmarking study of assembly pipelines). At the moment of this writing, read accuracy and cost of data for PacBio (Sequel system) and ONT (R9 flow cell) were equivalent; PacBio generated higher proportion of long reads than ONT; however, PacBio's library prep required ten fold more high-quality DNA than

350 ONT. Both for PacBio and ONT it is critically important to obtain high molecular weight DNA in
351 fragments exceeding 20kb in length. For new OMOs, it is also essential to confirm that the DNA
352 is accessible to enzymatic modifications by trying to digest it with a frequent-cutting restriction
353 endonuclease.
354
For genome scanning, gene expression, or invertebrate DNA methylation analyses targeting
protein-coding sequences (exome) genome sequence might not be the best reference; instead, a
highly contiguous transcriptome assembly would be preferable. Until now the standard way to
generate a *de novo* transcriptome was to perform high-coverage RNA-seq and assemble the
results with Trinity [46]. In the coming years, it is expected that even higher-quality and lower-
cost OMO transcriptomes would be generated by PacBio or ONT sequencing of full-length
cDNA (or, for ONT, direct mRNA sequencing). The long-read capacity of these technologies
would essentially obviate the need for assembly, leaving only the sequence correction procedure
to be performed.
364
Finally, which tissue or body part to sample for sequencing? For genome sequencing it does not
matter much as long as contamination by other DNA sources can be kept to a minimum, but for
de novo transcriptomics it is not a trivial question, as gene expression varies dramatically across
tissues and life cycle stages. In mammals, there is definitely an organ of choice that expresses
nearly all genes in the genome: testis. Rather unexpected transcriptome complexity in the testis is
putatively due to chromatin re-packaging during spermatogenesis, which results in genome-wide
transcription leakage [47]. If so, testis might be a good choice for *de novo* transcriptomics not
only for mammals but for any organism that produces compact sperm.
373
374 **Note on data sharing**
375
As we have seen, the best power of ecological genomics in OMOs is achieved using a genome or
transcriptome reference. Every new reference dataset enables new biological questions, and the
whole OMO field will get a great boost if these resources are promptly shared. Please consider
rapidly sharing your reference data, at least as soon as the initial preprint of your paper is posted
to bioRxiv and ideally sooner, by distributing the link to data through research-related email list
or professional twitter feed.
382

References
1   Excoffier, L. *et al.* (2013) Robust demographic inference from genomic and SNP data.
PLoS Genet. 9, e1003905
2   Gutenkunst, R.N. *et al.* (2009) Inferring the joint demographic history of multiple
populations from multidimensional SNP frequency data. *PLoS Genet.* 5, e1000695
3   Jouganous, J. *et al.* (2017) Inferring the Joint Demographic History of Multiple
Populations: Beyond the Diffusion Approximation. *Genetics* DOI:
10.1534/genetics.117.200493
4   Andrews, K.R. *et al.* (2016) Harnessing the power of RADseq for ecological and
evolutionary genomics. *Nat. Rev. Genet.* 17, 81–92
5   Puritz, J.B. *et al.* (2014) Demystifying the RAD fad. *Mol. Ecol.* 23, 5937–5942
6   Robinson, J.D. *et al.* (2014) Sampling strategies for frequency spectrum-based population
genomic inference. *Bmc Evol. Biol.* 14, 254

7   Günther, T. and Coop, G. (2013) Robust identification of local adaptation from allele
frequencies. *Genetics* 195, 205–20

8   Schiffels, S. *et al.* (2016) Iron Age and Anglo-Saxon genomes from East England reveal
British migration history. *Nat. Commun.* 7, 1–9

9   Han, E. *et al.* (2014) Characterizing bias in population genetic inferences from low-
coverage sequencing data. *Mol. Biol. Evol.* 31, 723–35

10   Matz, M. V *et al.* (2017) Potential for rapid genetic adaptation to warming in a Great
Barrier Reef coral. *bioRxiv* at <http://www.biorxiv.org/content/early/2017/06/18/114173>

11   Dixon, G.B. *et al.* (2015) Genomic determinants of coral heat tolerance across latitudes.
Science (80-.). 348, 1460–1462

12   Mckenna, A. *et al.* (2010) The Genome Analysis Toolkit: A MapReduce framework for
analyzing next-generation DNA sequencing data. *Genome Res.* 20, 1297–1303

13   Korneliussen, T. *et al.* (2014) ANGSD: Analysis of Next Generation Sequencing Data.
BMC Bioinformatics 15, 356

14   Nielsen, R. *et al.* (2012) SNP calling, genotype calling, and sample allele frequency
estimation from new-generation sequencing data. *PLoS One* 7, e37558

15   Andrews, K.R. and Luikart, G. (2014) Recent novel approaches for population genomics
data analysis. *Mol. Ecol.* 23, 1661–1667

16   Casbon, J.A. *et al.* (2011) A method for counting PCR template molecules with
application to next-generation sequencing. *Nucleic Acids Res.* 39, e81

17   Gojobori, J. *et al.* (2007) Adaptive evolution in humans revealed by the negative
correlation between the polymorphism and fixation phases of evolution. *Proc. Natl. Acad.*
Sci. U. S. A. 104, 3907–12

18   Tine, M. *et al.* (2014) European sea bass genome and its variation provide insights into
adaptation to euryhalinity and speciation. *Nat. Commun.* 5, 5770

19   Lowry, D.B. *et al.* (2017) Breaking RAD: An evaluation of the utility of restriction site
associated DNA sequencing for genome scans of adaptation. *Mol. Ecol. Resour.* 17, 142–
152

20   Tiffin, P. and Ross-Ibarra, J. (2014) Advances and limits of using population genetics to
understand local adaptation. *Trends Ecol. Evol.* 29, 673–680

21   McKinney, G.J. *et al.* (2017) RADseq provides unprecedented insights into molecular
ecology and evolutionary genetics: comment on Breaking RAD by Lowry *et al* . (2016).
Mol. Ecol. Resour. 17, 356–361

22   Davies, R.W. *et al.* (2016) Rapid genotype imputation from sequence without reference
panels. *Nat. Genet.* 48, 965–969

23   Money, D. *et al.* (2015) LinkImpute: Fast and Accurate Genotype Imputation for
Nonmodel Organisms. *G3 Genes, Genomes, Genet.* 5, 2383–2390

24   Morin, R.D. *et al.* (2008) Profiling the HeLa S3 transcriptome using randomly primed
cDNA and massively parallel short-read sequencing. *Biotechniques* 45, 81–94

25   Meyer, E. *et al.* (2011) Profiling gene expression responses of coral larvae (Acropora
millepora) to elevated temperature and settlement inducers using a novel RNA-Seq
procedure. *Mol. Ecol.* 20, 3599–3616

26   Lohman, B.K. *et al.* (2016) Evaluation of TagSeq, a reliable low-cost alternative for
RNAseq. *Mol. Ecol. Resour.* 16, 1315–1321

27   Moll, P. *et al.* (2014) QuantSeq 3 ' mRNA sequencing for RNA quantification. *Nat.*
Methods 11, 25

28   Rostom, R. *et al.* (2017) Computational approaches for interpreting scRNA-seq data.
FEBS Lett. DOI: 10.1002/1873-3468.12684

29   Kenkel, C.D. and Matz, M. V (2016) Gene expression plasticity as a mechanism of coral
adaptation to a variable environment. *Nat. Ecol. Evol.* 1, 14

30   Dixon, G.B. *et al.* (2014) Bimodal signatures of germline methylation are linked with gene

447		expression plasticity in the coral Acropora millepora. *BMC Genomics* 15, 1109
448	31	Crowley, J.J. *et al.* (2015) Analyses of allele-specific gene expression in highly divergent
449		mouse crosses identifies pervasive allelic imbalance. *Nat. Genet.* 47, 353–60
450	32	The GTEx Consortium (2015) The Genotype-Tissue Expression (GTEx) pilot analysis:
451		Multitissue gene regulation in humans. *Science (80-.).* 348, 648–660
452	33	Melé, M. *et al.* (2015) The human transcriptome across tissues and individuals. *Science*
453		*(80-.).* 348, 660–665
454	34	Gamazon, E.R. *et al.* (2015) A gene-based association method for mapping traits using
455		reference transcriptome data. *Nat. Genet.* 47, 1091–1098
456	35	Wheeler, H.E. *et al.* (2016) Survey of the Heritability and Sparse Architecture of Gene
457		Expression Traits across Human Tissues. *PLOS Genet.* 12, e1006423
458	36	Lämke, J. and Bäurle, I. (2017) Epigenetic and chromatin-based mechanisms in
459		environmental stress adaptation and stress memory in plants. *Genome Biol.* 18, 124
460	37	Feng, S. *et al.* (2010) Conservation and divergence of methylation patterning in plants and
461		animals. *Proc. Natl. Acad. Sci. U. S. A.* 107, 8689–94
462	38	Dixon, G.B.. *et al.* (2016) Evolutionary Consequences of DNA Methylation in a Basal
463		Metazoan. *Mol. Biol. Evol.* 33, msw100
464	39	Sarda, S. *et al.* (2012) The Evolution of Invertebrate Gene Body Methylation. *Mol Biol*
465		*Evol* 29, 1907–1916
466	40	Uller, T. *et al.* (2013) Weak evidence for anticipatory parental effects in plants and
467		animals. *J. Evol. Biol.* 26, 2161–2170
468	41	Kitzman, J.O. (2016) Haplotypes drop by drop. *Nat. Biotechnol.* 34, 296–298
469	42	Gordon, D. *et al.* (2016) Long-read sequence assembly of the gorilla genome. *Science (80-*
470		*.).* 352, aae0344
471	43	Michael, T.P. *et al.* (2017) High contiguity Arabidopsis thaliana genome assembly with a
472		single nanopore flow cell. *bioRxiv* at
473		<http://www.biorxiv.org/content/early/2017/06/14/149997>
474	44	Jansen, H.J. *et al.* (2017) Rapid de novo assembly of the European eel genome from
475		nanopore sequencing reads. *bioRxiv* at
476		<http://www.biorxiv.org/content/early/2017/01/20/101907>
477	45	Berlin, K. *et al.* (2015) Assembling large genomes with single-molecule sequencing and
478		locality-sensitive hashing. *Nat. Biotechnol.* 33, 623–630
479	46	Haas, B.J. *et al.* (2013) De novo transcript sequence reconstruction from RNA-seq using
480		the Trinity platform for reference generation and analysis. *Nat Protoc* 8, 1494–1512
481	47	Soumillon, M. *et al.* (2013) Cellular Source and Mechanisms of High Transcriptome
482		Complexity in the Mammalian Testis. *Cell Rep.* 3, 2179–2190
483	48	Cosart, T. *et al.* (2011) Exome-wide DNA capture and next generation sequencing in
484		domestic and wild species. *BMC Genomics* 12, 347
485	49	Bundock, P.C. *et al.* (2012) Enrichment of genomic DNA for polymorphism detection in a
486		non-model highly polyploid crop plant. *Plant Biotechnol. J.* 10, 657–667
487	50	De Wit, P. *et al.* (2015) SNP genotyping and population genomics from expressed
488		sequences - current advances and future possibilities. *Mol. Ecol.* 24, 2310–2323
489	51	Bay, R.A. and Palumbi, S.R. (2014) Multilocus adaptation associated with heat resistance
490		in reef-building corals. *Curr. Biol.* 24, 2952–6
491	52	Futschik, A. and Schlötterer, C. (2010) The Next Generation of Molecular Markers From
492		Massively Parallel Sequencing of Pooled DNA Samples. *Genetics* 186, 207–218
493	53	Kofler, R. *et al.* (2011) PoPoolation2: identifying differentiation between populations
494		using sequencing of pooled DNA samples (Pool-Seq). *Bioinformatics* 27, 3435–3436
495	54	Therkildsen, N.O. and Palumbi, S.R. (2017) Practical low-coverage genomewide
496		sequencing of hundreds of individually barcoded samples for population and evolutionary
497		genomics in nonmodel species. *Mol. Ecol. Resour.* 17, 194–208

55 Paradis, E. *et al.* (2004) APE: Analyses of phylogenetics and evolution in R language.
Bioinformatics 20, 289–290

56 Oksanen, J. *et al.* vegan: Community Ecology Package. , *R package ver. 2.4–3*. (2017)

57 Wright, R.M. *et al.* (2015) Gene expression associated with white syndromes in a reef
building coral, Acropora hyacinthus. *BMC Genomics* 16, 371

58 Strader, M.E. *et al.* (2016) Red fluorescence in coral larvae is associated with a diapause-
like state. *Mol. Ecol.* 25, 559–569

59 Jombart, T. (2008) adegenet: a R package for the multivariate analysis of genetic markers.
Bioinformatics 24, 1403–5

60 Langfelder, P. and Horvath, S. (2008) WGCNA: an R package for weighted correlation
network analysis. *BMC Bioinformatics* 9, 559

61 Bay, R.A. and Palumbi, S.R. (2017) Transcriptome predictors of coral survival and growth
in a highly variable environment. *Ecol. Evol.* 7, 4794–4803

62 Rose, N.H. *et al.* (2016) Gene Networks in the Wild: Identifying Transcriptional Modules
that Mediate Coral Resistance to Experimental Heat Stress. *Genome Biol. Evol.* 8, 243–
252

63 Lister, R. *et al.* (2009) Human DNA methylomes at base resolution show widespread
epigenomic differences. *Nature* 462, 315–22

64 Meissner, A. *et al.* (2008) Genome-scale DNA methylation maps of pluripotent and
differentiated cells. *Nature* 454, 766–770

65 Serre, D. *et al.* (2010) MBD-isolated Genome Sequencing provides a high-throughput and
comprehensive survey of DNA methylation in the human genome. *Nucleic Acids Res.* 38,
391–9

66 Jacinto, F. V. *et al.* Methyl-DNA immunoprecipitation (MeDIP): Hunting down the DNA
methylome. , *BioTechniques*, 44. (2008) , 35–43

67 Wang, S. *et al.* (2015) MethylRAD: a simple and scalable method for genome-wide DNA
methylation profiling using methylation-dependent restriction enzymes. *Open Biol.* 5,
150130

68 Flusberg, B.A. *et al.* (2010) Direct detection of DNA methylation during single-molecule,
real-time sequencing. *Nat. Methods* 7, 461–5

69 Feng, Z. *et al.* (2013) Detecting DNA Modifications from SMRT Sequencing Data by
Modeling Sequence Context Dependence of Polymerase Kinetic. *PLoS Comput. Biol.* 9,
e1002935

70 Stoiber, M.H. *et al.* (2017) De novo Identification of DNA Modifications Enabled by
Genome-Guided Nanopore Signal Processing. *bioRxiv* at
<http://www.biorxiv.org/content/early/2017/04/10/094672>

71 Koren, S. *et al.* (2017) Canu: scalable and accurate long-read assembly via adaptive k-mer
weighting and repeat separation. *Genome Res.* 27, 722–736

72 Chin, C.-S. *et al.* (2016) Phased diploid genome assembly with single-molecule real-time
sequencing. *Nat. Methods* 13, 1050–1054

73 Li, H. (2016) Minimap and miniasm: Fast mapping and de novo assembly for noisy long
sequences. *Bioinformatics* 32, 2103–2110

74 Vaser, R. *et al.* (2017) Fast and accurate de novo genome assembly from long uncorrected
reads. *Genome Res.* 27, 737–746

75 Walker, B.J. *et al.* (2014) Pilon: An integrated tool for comprehensive microbial variant
detection and genome assembly improvement. *PLoS One* 9, e112963

544
545

Box 1: AFS models.
In the world of OMOs we are usually dealing with samples from many populations, which would be hard or impossible to model simultaneously; moreover, there are usually many populations left unsampled. To infer meaningful demographic parameters in a sparsely sampled system of many populations, a practical solution is to perform two-dimensional AFS analysis of all population pairs [10]. Typical hypotheses and corresponding tests are:
- Are the two populations demographically separate?
  - o compare model with split to model without split, under which the two compared populations are regarded as independent samples from the same population.
- If yes, is there still gene flow between them?
  - o compare split models with and without migration.
- If yes, is the gene flow symmetric or asymmetric?
  - o compare split model with two potentially different migration rates to a split model with a single symmetrical migration rate.
- Was population size stable or went through changes in the past?
  - o compare single-population model involving population size change in the past to a standard neutral model.

Simple command-line scripts for AFS plotting and running basic pairwise models in *moments* can be found here: https://github.com/z0on/AFS-analysis-with-moments. To access the full potential of *moments*, however, the user is expected to compose python scripts of their own.

547 **Table 1. Genotyping approaches for genome scanning.**

Approach	Features	Pros	Cons
Exome-seq [48,49]	Isolates and sequences only the protein-coding portion of genome.	Dense coverage of genes guarantees that coding variants and variants linked to cis-regulatory mutations are discovered.	Other (arguably less important) types of variation are not profiled (e.g., distant enhancers).
RNA-seq [50,51]	Sequences RNA.	Same as exome sequencing.	Genotyping quality of a gene depends on expression level. Allele-specific expression affects accuracy of heterozygote calls.
Pool-seq [52,53]	Sequences pooled DNA from multiple individuals from each population.	Dense whole-genome coverage with confident determination of allele frequencies in populations.	No possibility for individual–based analysis (such as STRUCTURE) or validation based on genotype-phenotype association across individuals. Must be confident in *a priori* population designations.
Low-coverage whole-genome sequencing (WGS) [54]	Sequences individual genomes at ~1-4x coverage.	Dense whole genome coverage at individual level.	Per-site genotypes are unreliable because of missing data; must use uncertainty-aware analysis such as *ANGSD*.
Ultra-low coverage WGS with imputation [22]	Sequences individual genomes at <2x coverage, imputes missing genotypes and corrects false homozygote calls	Dense whole genome coverage at individual level, phased data enables haplotype-based analysis	Rare alleles (minor allele frequency<0.05) are missed. Requires large sample sizes (depending on LD, hundreds or thousands of individuals). Accuracy of imputation must be experimentally validated for every new OMO.

548
549

550 **Table 2. Gene expression analyses not relying on accurate gene annotations**

Analysis	What does it do	Software	Applications
Principal coordinate analysis based on Manhattan distances (sum of all log-fold changes across genes)	Characterizes overall transcriptome differences across experimental groups. Measures fraction of variation attributable to each experimental factor.	R: package *ape*, function *pcoa* [55] package *vegan,* function *adonis* [56]	[57,58]
Differential analysis of principal components (DAPC)	Quantifies transcriptome differences between samples with respect to specified multivariate axis. Good for quantifying overall gene expression plasticity.	R: package *adegenet* [59]	[29]
Weighted gene co-expression network analysis (WGCNA)	Identifies co-regulated groups of genes, which are linked to experimental factors and traits *post hoc*. Method of choice for complex experimental designs (>20 samples) with many quantitative traits measured.	R: package WGCNA [60]	[29,61,62]
Rank-based functional summaries of KOG (euKaryotic Orthologous Groups) classes	Reveals broad functional trends in gene expression. Particularly useful for OMOs since it tolerates sparse and inaccurate annotations. Its main use is for statistical comparison of highly diverse datasets, even from different species.	R: package *KOGMWU* [11].	[11,58]

553 **Table 3. Methods for interrogating DNA methylation**

Method	Features	Pros	Cons
Whole-Genome Bisulfite Sequencing (WGBS) [63]	Sequences complete genome after bisulfite conversion	Complete characterization of 5me-cythosine methylation at single-base resolution	High coverage is required to obtain quantitative data. In non-vertebrate OMOs, much sequencing effort is wasted since most of genome is not methylated.
RRBS-seq [64]	Bisulfite sequencing of genome fragments adjacent to all (methylated and un-methylated) CCGG sites	Saves costs dramatically compared to WGBS.	Only a fraction of all CpG sites is interrogated. Complicated library preparation protocol. Sequencing effort is wasted on non-methylated sites.
MBD-seq [65], meDIP [66]	Pull-down and sequencing of methylated DNA.	Optimizes sequencing effort by focusing on methylated DNA.	Complicated library preparation protocol. Resolution equals the length of pulled-down fragments (~300-500b). Pull-down procedure is not absolutely efficient, many reads still correspond to un-methylated genome regions.
methylRAD [67]	Direct sequencing of genomic fragments adjacent only to the methylated CCGG and CCWGG sites.	Very simple library prep protocol. Highly cost-efficient due to focus on methylated sites only.	Only a fraction of all CpG sites is interrogated. New method, requires further benchmarking.
PacBio [68,69]	Direct detection of modified DNA bases during normal SMRT sequencing, based on polymerase lags.	Robust detection of 4-methylcytosine, 8-oxoguanine, and N6-methyladenine. Single-base resolution.	Same as WGBS. 5-methylcytosine, the most common methylation mark in animals, is not reliably detected.
ONT [70]	Direct detection of modified DNA bases during normal nanopore sequencing, based on conductivity changes.	Detects all marks, including 5-methylcytosine. Single-base resolution.	Same as WGBS.

554
555

557 **Table 4. Assembly pipelines for PacBio and ONT reads**

Pipeline	Required coverage	Features	Pros	Cons
Canu +Quiver* [71]	>30x	Correct and trims reads before assembly.	Best accuracy at base, indel and assembly level.	Very computationally demanding for large genomes. Generates incomplete assemblies at low coverage.
Falcon + Quiver* [72]	>50x	Similar to Canu.	Standard for PacBio.	Very computationally demanding for large genomes. High reliance on reads >20kb. Highly incomplete assemblies at low coverage.
minimap + miniasm + racon [73,74]	<30x	Raw reads are assembled, correction is done post-assembly	Very fast even for large genomes. Works with lower coverage, shorter reads than Canu and Falcon.	The resulting accuracy is lower than with Canu + Quiver.
pilon [75]	NA (error correction method)	Performs additional correction post-assembly.	Boosts accuracy for any assembly.	Requires high-quality Illumina reads.

558 *Quiver is a consensus polishing software that is now replaced by Arrow to handle PacBio
559 Sequel data (https://github.com/PacificBiosciences/GenomicConsensus). Racon [74] can be used
560 instead of Quiver/Arrow [43].
561