1

Multinucleotide mutations cause false inferences of lineage-specific positive selection

Aarti Venkat[1], Matthew W. Hahn[2], Joseph W. Thornton*[1,3]

(1) Department of Human Genetics, University of Chicago, Chicago IL 60637, USA

(2) Department of Biology and Department of Computer Science, Indiana University, Bloomington IN 47405, USA

(3) Department of Ecology & Evolution, University of Chicago, Chicago IL 60637, USA

*Correspondence: Joseph Thornton, joet1@uchicago.edu

Keywords: adaptation, adaptive evolution, branch-site test, codon models, transversions

ABSTRACT

Phylogenetic tests of adaptive evolution, which infer positive selection from an excess of

nonsynonymous changes, assume that nucleotide substitutions occur singly and independently.

But recent research has shown that multiple errors at adjacent sites often occur in single events

during DNA replication. These multinucleotide mutations (MNMs) are overwhelmingly likely

to be nonsynonymous. We therefore evaluated whether phylogenetic tests of adaptive evolution,

such as the widely used branch-site test, might misinterpret sequence patterns produced by

MNMs as false support for positive selection. We explored two genome-wide datasets

comprising thousands of coding alignments – one from mammals and one from flies – and found

that codons with multiple differences (CMDs) account for virtually all the support for lineage-

specific positive selection inferred by the branch-site test. Simulations under genome-wide,

empirically derived conditions without positive selection show that realistic rates of MNMs

cause a strong and systematic bias in the branch-site and related tests; the bias is sufficient to

produce false positive inferences approximately as often as the branch-site test infers positive

selection from the empirical data. Our analysis indicates that genes may often be inferred to be

under positive selection simply because they stochastically accumulated one or a few MNMs.

Because these tests do not reliably distinguish sequence patterns produced by authentic positive

selection from those caused by neutral fixation of MNMs, many published inferences of adaptive

evolution using these techniques may therefore be artifacts of model violation caused by

unincorporated neutral mutational processes. We develop an alternative model that incorporates

MNMs and may be helpful in reducing this bias.

INTRODUCTION

Identifying genes that evolved under the influence of positive natural selection is a central goal in molecular evolutionary biology. During recent decades, likelihood-based phylogenetic methods have been developed to identify gene sequences that retain putative signatures of past positive selection [1-10]. Perhaps the most widely used of these is the branch-site test (BST) of episodic selection, which allows positive selection to affect only some codons on one or a few specified branches of a phylogeny, and therefore has relatively high power compared to methods that detect selection across an entire sequence or an entire phylogenetic tree [5,6,11]. The BST has been the basis for published claims of lineage-specific adaptive evolution in many thousands of individual genes [12-16].

The BST and related methods use a likelihood ratio test to compare how well two mixture models of sequence evolution on a phylogeny fit an alignment of coding sequence data. The null model constrains all codons to evolve with rates of nonsynonymous substitution (d_N) less than or equal to the rate of synonymous substitution (d_S), as expected under purifying selection and drift. In the positive selection model, some sites are allowed to have $d_N > d_S$ on a branch or branches of interest. If the increase in likelihood of this model given the data is greater than expected due to chance alone, the null model is rejected and adaptive evolution is inferred. The BST has been shown to be conservative, with a low rate of false positive inferences, when sequences are generated under an evolutionary process corresponding to the null model [6,11]. It is widely appreciated that likelihood ratio tests can become biased if the underlying probabilistic model is incorrect [17]. The effect on the BST of a few forms of model violation—such as an unequal distribution of selective effects among sites, positive selection on non-foreground lineages, high sequence divergence, and non-allelic gene conversion—have been previously studied [18-22] and the test has been found to be reasonably robust to most but not all forms of violation examined [6,23,24].

Recent research in molecular genetics and genomics suggests a potentially important phenomenon that has not been incorporated into models used in tests of positive selection: the propensity of DNA polymerases to produce mutations at neighboring sites. All implementations of the BST and other likelihood-based tests of adaptive evolution use models in which mutations occur only at individual nucleotide sites and are fixed singly and independently. Codons with multiple differences between them can be interconverted only by serial single-nucleotide

substitutions, the probability of which is the product of the probabilities of each independent

event. Recent molecular studies have shown, however, that mutations affecting adjacent

nucleotide sites often occur during replication, apparently because certain DNA microstructures

recruit error-prone polymerases that lack proofreading activity and therefore make multiple

errors close together [25–33]. Consistent with these mechanisms, genetic studies of human trios and

mutation-accumulation experiments in laboratory organisms indicate that *de novo* mutations

occur in tandem or at nearby sites more frequently than expected if each occurred independently

[25,32–36], and these multinucleotide mutations (MNMs) are enriched in transversions [35,37,39]. The

precise frequency at which MNMs occur is difficult to estimate, but a recent compilation of

genetic studies in humans concluded that about 0.4% of mutations, polymorphisms, and

substitutions are at directly adjacent sites (counting each tandem pair as one event) [34]. In

Drosophila melanogaster genomes, analysis of rare polymorphisms and mutation-accumulation

experiments estimated that 1.3% of all mutations are at adjacent sites [38]. Although the methods

and data sources in these studies differ, these findings suggest that tandem MNMs probably

account for on the order of 1% of mutational events.

We hypothesized that these mutational processes might lead to false signatures of

positive selection in the BST. Because of the structure of the genetic code, virtually all MNMs

in coding sequences are nonsynonymous, and most would comprise multiple nonsynonymous

nucleotide changes if they were to occur by single nucleotide steps (**Supplementary Table 1**).

The enrichment of transversions in MNMs further increases the propensity for MNMs to produce

nonsynonymous changes, because transversions are more likely than transitions to be

nonsynonymous. MNMs are therefore likely to produce codons with multiple differences

(CMDs) that contain an apparent excess of nonsynonymous substitutions. When these CMDs are

assessed using a method that treats all substitutions as independent events, a model that allows

d_N to exceed d_S at some sites may have a higher likelihood than one that restricts d_N/d_S to values

≤ 1. Further, the assumption that all mutations have the same transversion-transition rate might

exacerbate the tendency to misinterpret MNM-produced nonsynonymous changes as evidence

for positive selection. Of course, CMDs can also be driven to fixation by positive selection [11,40–

42]—and the same is true of transversion-rich substitutions—but these considerations suggest that

failing to incorporate MNMs in likelihood models might make tests of adaptive evolution

susceptible to false positive inferences. The BST and other lineage-specific tests might be

particularly sensitive to this problem because they seek signatures of positive selection acting on

small numbers of codons on one or a few specified branches of the tree [43]. Simulation studies

suggest that MNMs may elevate false positive rates in some selection tests [44], but there has been

no comprehensive analysis of the effect of MNMs, particularly on the branch-site test or under

realistic, genome-scale conditions.

RESULTS

To understand the effect of MNMs on the accuracy of the branch-site test, we analyzed in

detail two previously published genome-wide datasets, which represent classic examples of the

application of the test [13,15,45]. The mammalian dataset consists of coding sequences of 16,541

genes from six eutherian mammals; we retained for analysis only the 6,868 genes with complete

species coverage. The fly dataset consists of 8,564 genes from six species in the melanogaster

subgroup clade, all of which had complete coverage (**Supplementary Fig. 1**). The fly genes

have higher sequence divergence than those in the mammalian dataset, allowing us to examine

the performance of the BST under different evolutionary conditions.

We used the classic BST to identify genes putatively under positive selection (P<0.05) on

the human lineage in the mammalian dataset and on each of the six terminal lineages in flies. 82

genes in humans and 3,938 in flies yielded significant tests (**Supplementary Table 2**). To

facilitate robust further analysis of CMDs, we filtered out genes in which CMDs occurr at sites

with indels or in which the ancestral states of CMDs are reconstructed differently between the

null and positive selection models; we also applied a multiple testing correction (FDR <0.20). In

flies, 443 genes were retained after these steps. Thirty human genes passed the CMD alignment

and reconstruction filter, but none met the FDR threshold, consistent with previous analyses of

these data; [15] nevertheless, we included the 30 initially significant human genes because this

lineage is the object of intense interest and because its short length contrasts with the fly

branches. These two groups constitute the "BST-significant" sets of genes in flies and humans.

CMDs provide virtually all support for positive selection

We sought to determine how much of the evidence for positive selection comes from CMDs. We first observed that CMDs were dramatically enriched in BST-significant genes compared to non-BST-significant genes (**Fig. 1a**). In humans, BST-significant genes contain one CMD on average, while BST-nonsignificant genes contain none (**Supplementary Fig. 2**). The pattern is similar but less extreme in flies, with the average number of CMDs per BST-significant gene greater than that in non-significant genes (**Supplementary Fig. 2**). When CMD-containing codons are excluded from the alignments, the vast majority of genes that were BST-significant lose their signature of selection in both datasets (**Fig. 1b**).

We next calculated the fraction of statistical support for positive selection that comes from CMDs. The total support for positive selection in an alignment is defined as the difference between the log-likelihood of the positive selection model and that of the null model, summed across all codons in the alignment. The fraction of support from CMDs is the support from CMD-containing codons divided by the total support across the entire alignment. CMDs account for >95% of the support for positive selection in virtually all BST-significant genes in both datasets; in about 70% of genes, CMDs provide all the support (**Fig. 1c**).

Finally, we examined the BST's *a posteriori* identification of sites under positive selection. We found that CMDs were far more likely to be classified as positively selected than non-CMDs. Among genes that were BST-significant on the human lineage, every CMD was inferred to be under positive selection using a Bayes Empirical Bayes posterior probability (PP) cutoff > 0.5. Using a more stringent cutoff of PP>0.9, 66 percent of CMDs were classified as positively selected, compared to 0.07% of non-CMDs. In the fly dataset, CMDs accounted for 90% of codons with BEB>0.9, although they represent less than 1% of all codons (**Fig. 1d**).

CMDs are therefore the primary drivers of the signature of selection identified in the BST. A single CMD provides sufficient statistical support to yield a signature of positive selection on the human lineage, and only a few CMDs in a gene are enough to do the same in flies.

Incorporating MNMs eliminates the signature of positive selection in many genes

CMDs might be enriched in BST-positive genes because of an MNM-induced bias or because they were fixed by positive selection. To incorporate both neutral and selection-driven

fixation of MNMs into a BST framework, we developed a codon model in which double-

nucleotide changes are allowed, with the parameter δ serving as a multiplier that modifies the

rate of each double-nucleotide substitution relative to single-nucleotide substitutions. We

implemented a version of the BST (BS+MNM) that is identical to the classic version, except that

both the null and positive selection models allow MNMs. Simulations under conditions derived

from a sample of genes in the mammalian dataset show that the method estimates the parameters

used to generate the sequences with reasonable accuracy (**Supplementary Fig. 3**).

We first fit the BS+MNM null model to all alignments in the mammalian and fly

datasets. The average estimate of δ across all genes was 0.026 in mammals and 0.062 in flies,

with δ in both cases about twice as high in the subset of BST-significant genes as in BST-

nonsignificant genes (**Fig. 2a)**. Using a likelihood-ratio test, we found significant support for the

BS+MNM null model (compared to the classic BST null model) in 22% of human genes and

>50% of fly genes (**Supplementary Table 3**); simulations without MNMs showed that this

comparison has a very low false-positive rate (**Supplementary Table 4).**

We then used this BS+MNM test to evaluate the empirical sequences for positive

selection. We found that 96% of the BST-significant genes on the human lineage lost

significance in the BS+MNM test (**Figs. 2b, Supplementary Table 5**). In flies, 38% of the

BST-significant genes lost significance; a substantial fraction of those that retained significance

were enriched in triple substitutions, a process not accounted for in our model (**Figs. 2b,**

Supplementary Table 5).

MNMs cause false positive inferences on a genome-wide scale

That the BS+MNM test eliminates the signature of positive selection from many genes

could have several causes, including: 1) the more complex BS+MNM model may have reduced

power to identify authentic positive selection compared to the BST, 2) incorporating MNMs may

ameliorate a bias towards false positive inference in the classic BST that is caused by MNMs,

and 3) the additional δ parameter in the BS+MNM test may allow it to incorporate other forms of

sequence complexity, potentially ameliorating a bias caused by other model violations.

We addressed these possibilities in two ways. First, we performed power analyses of the

BS+MNM test using simulations in which positive selection is present in the generating model.

We simulated sequence data on the mammalian and fly phylogenies using genome-wide

averages for all parameters of the BST positive selection model, but we varied the strength of

positive selection (ω_2) and the proportion of sites under positive selection. We then applied the

BS+MNM test to these data and found that it can reliably detect strong positive selection ($\omega_2 >$

20) when it affects more than 10% of sites in a typical gene, or moderate positive selection ($10 <$

$\omega_2 < 20$) that affects a larger fraction of sites (**Supplementary Fig. 4a**). Under parameters

derived from both datasets, the test's power is similar to that of the classic BST, with slight

reductions under only a few conditions on the fly lineage. Thus, although some genes may have

lost their signature of selection because of reduced power in the BS+MNM test, it appears

unlikely that a difference in power is the primary cause of the dramatic reduction in the number

of positive results when the test is used.

Second, we used simulations under null conditions to directly evaluate the frequency of

false positive inferences by the classic BST when sequences are generated with realistic rates of

multinucleotide mutation. For every gene in the mammalian and fly datasets, we simulated

sequence evolution under the null BS+MNM model without positive selection using parameters

derived from the alignments, including δ. These parameters generate sequences with an

observed frequency of tandem substitutions of 1.6% in humans and 3.2% in the *D. melanogaster*

lineage in flies, similar to or slightly higher than the observed frequencies in the empirical

datasets (1.3% and 1.6%, respectively), presumably because the BS+MNM model captures some

but not all aspects of real sequence evolution (**Supplemental Table 6**) [34, 38].

We then analyzed these positive-selection-free simulated data using the classic BST. In

both humans and flies, the number of genes with significant results—all of which are false

positive inferences—was greater than the number of genes that the BST had concluded were

under positive selection using the empirical data (**Fig. 3a**). In flies, almost 9 percent of tests

were false positives ($P<0.05$), despite the conservative approach the method uses to calculate P-

values [6,11], compared to just 1 percent under control simulations without MNMs. Further, more

than 1,700 of these false positive tests survived FDR adjustment, compared to just 4 in the

control simulations (**Supplementary Table 2)**. In humans, the fraction of false positive

inferences is lower, consistent with the test's reduced power in this dataset, but still about three

times greater than in the control simulations.

These false inferences are caused primarily by MNM-induced bias, because simulating

data under identical control conditions without MNMs ($\delta = 0$) produced few positive tests. All

other parameters were identical between the generating model and analysis models, so other

forms of model violation do not contribute to the bias observed in the simulation experiments.

Taken together, these findings indicate that MNMs under realistic evolutionary conditions

produce a strong and widespread bias in the BST toward false inferences of positive selection.

This bias is strong enough to cause the BST to make false inferences of positive selection at

about the same rate as it infers selection in the real genomes of humans and flies. In the

simulations, every positive result is false; in the tests of real sequences, the fraction of positive

results that are true is unknown.

Systematic bias caused by chance MNMs in longer genes

We next sought to identify the causal factors that determine whether a gene yields a false

positive result in the BST because of MNM-induced bias. Most genes are only several hundred

codons long, and only a few percent of mutations are MNMs, so on phylogenetic branches of

short to moderate length many genes will contain no CMDs caused by multinucleotide

mutations. The hypothesis that neutral fixation of MNMs contributes to inferences of positive

selection in the BST predicts that a gene's propensity to produce a BST-significant result should

depend on factors that increase the probability it will contain one or more fixed MNMs by

chance, including its length and the gene-specific rate at which MNMs occur within it.

We first tested for an effect of gene length on the results of the branch-site test. As

predicted, we observed that BST-significant empirical genes were on average 100 and 16 codons

longer than non-significant genes in the human and fly empirical datasets, respectively (**Fig. 3b**).

The relationship between length and propensity to yield a BST positive result could arise because

genes that present a larger "target" are more likely to undergo MNMs than shorter genes;

alternatively, longer genes, by including more sites for analysis, might increase the power of the

BST to detect authentic positive selection. However, in genome-wide simulations under the null

model with no positive selection (but with $\delta > 0$), genes with false positive BSTs are longer than

the non-significant genes by an average of 26 and 31 codons using the human and fly

parameters, respectively (**Supplementary Fig. 5**). This result cannot be attributed to increased

power to detect true positive selection and supports the conclusion that mutational target size

contributes to a gene's propensity to manifest MNM-induced bias by chance alone.

To directly test the causal relationship between sequence length and false-positive bias in

the BST, we simulated sequence evolution at increasing sequence lengths, using evolutionary

parameters derived from each of the BST-significant genes in the mammalian and fly datasets.

For each gene's parameters, we simulated 50 replicate alignments under the BS+MNM null

model and then analyzed them using the classic BST (**Supplementary Fig. 6a**). The false

positive rate for any gene's simulations is defined as the fraction of replicates with a significant

LRT in the classic BST, using a P-value cutoff of 0.05. When sequences 5,000 codons long

were simulated, 96% of BST-significant genes had an unacceptable FPR (>0.05), with a median

FPR of 0.39: increasing sequence length to 10,000 codons exacerbated the bias, with 100% of

genes yielding an unacceptable FPR and a median FPR of 0.56 (**Fig. 3c**). In flies, a similar

pattern was evident, and the false positive rates were even higher (median FPR=0.74 and 0.90 at

5,000 and 10,000 codons, respectively). Control simulations under identical conditions but with

δ=0 led to very low FPRs (median 0.02 to 0.03 in both datasets), even with very long sequences

(grey dots in **Fig. 3c**). A similar systematic and length-dependent bias also resulted when

sequences were simulated under gene-specific conditions, but with δ fixed to its average across

the thousands of BST-nonsignificant genes in each dataset (**Supplementary Fig. 6b**). Although

the sequence lengths tested are longer than most real genes, these experiments directly establish

that a gene's probability of returning a significant BST result in the absence of positive selection

is directly related to the target size it presents for chance fixation of MNMs.

We next evaluated whether the gene-specific rate of multinucleotide mutation affects a

gene's propensity to yield a positive result in the BST. As predicted, we observed that BST-

significant genes in the empirical datasets had higher estimated δ than nonsignificant genes **(Fig.**

2a). Genes producing false positive results in the genome-wide null simulations under empirical

conditions also tended to have higher δ (**Fig 3d**); this result that cannot be attributed to the

possibility that δ might be fitting CMDs fixed by positive selection, because positive selection

was absent from the generating model.

To directly test the effect of the neutral MNM substitution rate on the BST, we simulated

sequences 5,000 codons long under the null BS+MNM model, with a variable δ and all other

parameters fixed to their averages across all genes. We found that increasing δ led to a

monotonic increase in the frequency of false positive inferences. The FPR was >0.05 when δ

was only 0.001 and 0.013 on the human and fly lineages, respectively. When δ was equal to its

genome-wide average (0.026 and 0.062 in mammals and flies), false positive inferences occurred

at rates of 22 and 17 percent, respectively (**Fig. 3e**). As δ increased, so too did the inferred value

of the parameter ω_2, which represents the inferred intensity of positive selection in the model

(**Fig. 3f**).

Typical evolutionary conditions are therefore sufficient to cause a strong and systemic

bias in the BST.   MNMs are rare, however, so longer genes and those with higher rates of

multinucleotide mutation are more likely to undergo this process and manifest the bias. This

view is further supported by the fact that fewer genes are BST-positive on the human branch –

which is so short that substitutions of any type are rare, and MNMs even more so – than on the

fly phylogeny, where branches are longer, more CMDs are apparent, and hundreds of genes have

BST signatures of selection.   Taken together, these findings suggest that although some genes

with BST-significant results in empirical datasets could have evolved adaptively, many may

simply be those that happened to fix multinucleotide substitutions by chance alone.

Transversion-enrichment in CMDs exacerbates bias in the branch-site test

MNMs tend to produce more transversions than classical single-site mutational

processes, so if CMDs are produced by MNMs, they should be transversion-rich [35, 37, 39]. As

predicted, we found that transversion:transition ratio is elevated in CMDs relative to that in non-

CMDs by factors of three and two in mammals and flies, respectively (**Fig. 4a**). In the subset of

BST-significant genes, CMDs have an even more elevated transversion:transition ratio, as

expected if transversion-rich MNMs bias the test (**Fig. 4a**). These data are consistent with the

hypothesis that a transversion-rich MNM process produced many of the CMDs in BST-

significant genes, but it is also possible that positive selection could have enriched for

transversions.

To test whether transversion-enrichment in MNMs exacerbates the BST's bias, we

developed an elaboration of the BS+MNM model in which an additional parameter allows

MNMs to have a different transversion:transition rate ratio (κ_2) than single-site substitutions do

(κ_1). We estimated the maximum likelihood estimates of the model's parameters for every gene

in the mammalian and fly datasets and simulated sequences using genome-wide median values

for all model parameters and branch lengths, except for κ_2, which we varied. Sequences 10,000

codons long were used, because simulating shorter sequences resulted in a high variance in the

realized transversion:transition ratio. We analyzed these data using the classic BST and

calculated the fraction of replicates in which positive selection was inferred. We found that

increasing κ_2 caused a rapid and monotonic increase in the false positive rate, indicating that

transversion enrichment in MNMs exacerbates the test's bias. The effect is strong: when κ_2/κ_1 is

increased from 1 to 2, the FPR approximately doubles (**Fig. 4b**). Thus, realistic rates of MNM

generation and transversion-enrichment together cause an even stronger bias in the BST than

MNMs alone. This result cannot be accounted for by positive selection driving fixation of

transversions, because no positive selection was present in the simulations.

MNMs affect a newer test of positive selection

In recent years, newer likelihood-based methods have been introduced to test for episodic

site-specific positive selection [2,3,7]. All these methods are based on models of sequence

evolution that, like the BST, do not allow MNMs but instead model CMDs as the result of serial

site-specific substitutions. We therefore hypothesized that these methods might also be biased

by MNMs. We chose a more recent method, BUSTED [2], which was developed primarily to test

for episodic site-specific selection events across an entire tree. We tested its performance on

alignments 5,000 codons long that were simulated using the BS+MNM null model and

parameters estimated from the BST-significant gene alignments in humans and flies. To test for

MNM-induced bias, we compared results when δ was assigned to three different values: zero, its

average across all alignments in the mammalian or fly datasets, or its gene-specific value in each

of the BST-significant genes (**Supplementary Fig. 6a**).

We found that BUSTED was also sensitive to MNM-induced bias. When $\delta=0$, virtually

no genes' parameters led to frequent false positive inferences, with a median FPR <0.03 across

genes (**Fig. 5**). But when δ was assigned to its empirically estimated gene-specific value, the

parameters from every gene in humans and the majority in flies yielded false positive rates

>0.05, with median FPRs of 0.29 and 0.5, respectively (**Fig. 5**). Frequent false positive

inferences were evident when sequences were simulated using genome-wide average estimates

of δ, as well.

CMDs that invoke multiple nonsynonymous steps drive the signature of positive selection

Finally, we sought further insight into the reasons why CMDs yield a false signature of

positive selection in the BST and related tests. In standard models of codon evolution, CMDs

are interpreted as the result of two or more serial independent substitutions, even though they can

be produced by MNMs in a single mutational event. We hypothesized that CMDs that imply

multiple nonsynonymous nucleotide substitutions under these models would provide the

strongest support for the positive selection model. We therefore classified CMDs in the

empirical datasets by the minimum number of nonsynonymous single-nucleotide substitutions

required from the ancestral to derived codon state under standard codon models. As predicted,

we found that CMDs that imply more than one nonsynonymous step are dramatically enriched in

BST-significant genes (**Fig 6a**).

We also examined the statistical support provided by different kinds of CMDs. As the

number of nonsynonymous steps increased, the statistical support provided for the positive

selection model also increased (**Fig. 6b**). CMDs that imply one nonsynonymous and one

synonymous step typically provide weak to moderate support for the positive selection model,

but CMDs that imply two nonsynonymous steps provide very strong support. In many cases, a

single CMD in this latter category is sufficient to yield a statistically significant signature of

positive selection.

DISCUSSION
379

Our results demonstrate that the branch-site test suffers from a strong and systematic bias

toward false positive inferences. This bias is caused by a mismatch between the method's

underlying codon model of evolution – which assumes that a codon with multiple differences can

be produced only by two or more independent substitution events – and the recently discovered

phenomenon of multinucleotide mutation, which produces such codons in a single event.

Because of the structure of the genetic code and the high transversion rates that characterize

MNMs, most codons produced by this mechanism cause more than one nonsynonymous single-

nucleotide change. Confronted with this kind of codon data, the likelihood calculated by the

BST is determined by the product of the probabilities of the individual mutations. Under the null

model, the probability of such compound events is extremely small, but it can increase

dramatically when d_N/d_S exceeds one, as the positive selection model allows. This increase in

likelihood afforded by the positive selection model is much greater than it would be if the

substitution were interpreted as the result of a single multinucleotide event. Indeed, our results

show that a single codon comprising two nonsynonymous substitutions is often sufficient to

yield a statistically significant signature of positive selection in the BST for an entire gene.

As a result, CMDs are the primary drivers of positive results by the BST. Virtually all

statistical support for positive selection in real alignments comes from CMD-containing sites;

removing them from the alignment or incorporating MNMs into the BST's model eliminates the

signature of selection from the majority of genes. CMDs can be produced by either positive

selection or by neutral evolution under multinucleotide mutation. In the former case, the BST

will be correct; however, the test cannot reliably distinguish CMDs that represent authentic

evidence of positive selection from those caused by MNM-induced bias.

The bias is strong and pervasive under realistic conditions. Indeed, when sequences were

simulated under the null model using parameters estimated from the fly and mammalian datasets,

the number of genes with false positive BSTs was approximately the same as the number of

positive BST results when the empirical data were analyzed. There is therefore no excess of

BST-positive results in these genomes beyond that potentially attributable to MNM-induced bias.

Worse, these null simulations did not include the elevated transversion rate that characterizes

MNMs, which exacerbates the test's bias. Taken together, these results suggest the possibility

that MNM-induced bias could explain many of the BST's inferences of positive selection in

these datasets.

Are our findings from these datasets generalizable? MNMs appear to be a property of all

eukaryotic replication processes, and the MNM rates that we observed in mammals and flies are

in the same range as those previously identified in genetic and molecular studies in a variety of

eukaryotic species [25,34,38]. Both datasets comprise a small number of taxa, but the BST seeks

evidence of selection on individual branches, so it seems unlikely that larger trees will somehow

inoculate the test against MNM-induced bias. We observed strong bias on lineages with

divergence levels ranging from very low (on the human terminal branch) to moderate (the fly

branches), so this problem does not appear to be unique to highly diverged sequences or

phylogenies with long branches. We must therefore consider the possibility that many of the

thousands of previously published reports of positive selection based on the BST could simply be

the ones that happened by chance to neutrally fix one or more multinucleotide mutations.

We do not contend that the BST is always wrong or that molecular adaptive evolution

does not occur. The existence of a bias, even a strong one, towards false positive inferences does

not mean that all positive inferences are false: some of the CMDs in BST-significant genes may

have evolved because of authentic positive selection, either by repeated substitution of single

nucleotides in a codon or selection on MNMs. But because the BST test cannot distinguish the

kinds of sequence data produced by positive selection from those produced by neutral evolution

of MNMs, it does not provide reliable evidence that a gene evolved adaptively; nor does it

provide a reliable estimate of the fraction of genes in a large set that evolved under positive

selection. There are numerous cases of strongly supported adaptive evolution, such as those

involving host-parasite and intracellular genetic conflicts, that have produced sequence

signatures of positive selection in the BST and related tests that are likely to be authentic [46]. The

persuasive evidence in these cases, however, comes from sources other than the sequence

signature.

If the BST and other lineage-specific tests based on the single-step codon model are

unreliable in the face of multinucleotide mutation, what should researchers do? The BS+MNM

test could be used to accommodate multinucleotide mutation; our results suggest this may be a

promising approach. But there are many forms of evolutionary complexity that are not

incorporated in this model, such as MNMs that affect three consecutive nucleotides in a codon,

elevated transversion probability within MNMs, and many other kinds of heterogeneity that

might bias the BS+MNM test [47–49]. Other models are also available to incorporate MNMs [9], but

their accuracy and robustness are not well characterized, either. More work is therefore required

before the BS+MNM or similar models can be used with confidence in the branch-site or similar

tests.

A complementary approach is to use functional experiments to explicitly test hypotheses

that specific historical changes in molecular sequence caused changes in function or phenotype

thought to have mediated adaptation [50,51]. Indeed, the bias we observed may help to explain why

some molecular experiments have shown that codons with a high posterior probability of

positive selection in the BST do not contribute to putative adaptive functions, whereas the codon

changes that do confer those functions have low or moderate PPs [52]. Experimental tests provide

the most convincing evidence of a gene's putative adaptive history, but they require time-

consuming laboratory and fieldwork [53,54] , so it is not clear how to implement them on a genome-

wide scale. Future research may develop and validate more robust models to detect positive

selection, and these may help to identify candidate genes for which specific, testable hypotheses

of past molecular adaptation on specific phylogenetic lineages can be formulated. The test

primarily used for this purpose till now, however, is unreliable.

ACKNOWLEDGEMENTS

We are grateful to the members of the Thornton lab for discussion and helpful comments. We

thank the Beagle2, Midway2, and Tarbell supercomputing clusters at the University of Chicago.

We also thank the developers of HyPhy for presenting an open source platform that allows

limitless customization of standard analyses. Funding was provided by NIH R01GM104397 and

R01GM121931 (JWT), NSF DEB-1601781 (JWT and AV), NSF DBI-1564611 (MWH), and the

Precision Health Initiative of Indiana University (MWH).

AUTHOR CONTRIBUTIONS

Analyses were designed by all authors, performed by AV, and interpreted by all authors. The

manuscript was written by AV and JWT with contributions from MWH.

COMPETING FINANCIAL INTERESTS

The authors declare no competing financial interests.

METHODS

Datasets, quality control, and inference of BST-significant genes. We analyzed two
previously published comprehensive datasets of protein-coding alignments on a genomic scale,
one in six mammals, the other in six *Drosophila* species (**Supplementary Table 2**) [13,15,45]. We
aimed to apply the branch-site test on every terminal lineage in the *Drosophila* dataset, and on
the human lineage in the mammal dataset. We only retained gene alignments without gross
misalignments, possessing complete coverage in all fly species, and minimally all primate
species. We then applied the branch-site test as implemented in CODEML 4.7 to each alignment,
assuming the phylogenetic relationships reported in the published studies (**Supplementary Fig.**
2) [13,15]. Branch lengths and model parameters were estimated for each alignment by maximum
likelihood (ML), and the F3x4 model was used for codon frequencies. We tested each gene in
mammals for selection on the terminal branch leading to humans; in flies, each gene was tested
separately for selection on each of the six terminal branches, and we express the fraction of
positive inferences across genes as the proportion of all tests conducted [6]. As is standard
practice, we calculated P-values using a likelihood ratio test with 1 df (χ_1^2) which makes the test
conservative under the null hypothesis [6]. Genes were initially identified as having a putative BST
signature of selection at P<0.05. We then applied a correction for multiple testing to a false
discovery rate (FDR) <0.20 using the *q-value* package in R (available at
http://github.com/jdstorey/qvalue).

493   To facilitate unambiguous analysis of CMDs, we removed genes containing CMDs
falling in gaps. We also removed genes for which the ML ancestral reconstructions reported by
CODEML at the base of the tested branch differed between the null and positive selection
models, yielding a set of genes with CMDs that do not depend upon which model is chosen. In
flies, 443 gene-tests ("genes") were retained after these filters and constitute the BST-significant
set of genes from this dataset. No genes on the human lineage were significant after FDR
correction, so we retained as the BST-significant set from this dataset those genes that passed the
ancestral reconstruction filter and had P<0.05 (**Supplementary Table 2**). The BST-
nonsignificant set of genes comprises all genes that pass the alignment and ancestral
reconstruction filter that are not in the BST-significant set (*n*=6757, humans; *n*=6883, flies). We
also repeated our analysis of CMD enrichment (see below) using a gene set that had not been
filtered for reconstruction consistency and found that our conclusions were unchanged
(**Supplementary Table 7**)

506   We only considered genes where the ancestral codons (both CMD and non-CMD codons)
have the same reconstruction under the BST null and BST alternate models. In doing so, we have
also excluded CMDs in codons with gaps in the alignment. For example, in the human dataset, of
the 82 genes that initially provided support for positive selection, 30 genes consist of
unambiguously reconstructed codons under the null and alternate model (the BST-significant

gene set). In 49 genes, CMDs fall in gaps. We did not consider the ancestral codon
reconstructions at these sites, and excluded these from our analyses due to alignment
ambiguities. The remaining 3 genes have CMDs that do not fall in gaps, for which the ancestral
codons were reconstructed differently under the null and alternate models. If we re-consider
these 3 'positively selected' genes that were excluded, we find 3 additional CMDs, one in each
of the genes. Including these genes made little to no difference to our CMD enrichment results.
517

Support for positive selection. CMDs were identified in BST-significant and BST-
nonsignificant genes as codons with 2 or 3 observed nucleotide differences between the ML
states at the ancestral and extant nodes for the branch being tested; non-CMDs are codons with 0
or 1 differences on the branch tested.   CMDs were not assessed on branches not tested.

To determine the role of CMDs in significant results from the BST, we excluded codon
positions in BST-significant genes containing CMDs, reanalyzed the data using the BST, and
calculated the fraction of tests that retained a significant result ($P<0.05$).

We quantified the proportion of statistical support for positive selection in BST-
significant genes that comes from CMDs as follows. The site-specific support provided by one
codon site in an alignment is the difference between the log-likelihoods of the positive selection
model and the null model given the data at that site. Support for positive selection provided by
all CMDs in a gene (*support$_{CMD}$*) is the support summed over all CMD sites in the alignment.
The proportion of support provided by CMDs is *support$_{CMD}$ / (support$_{CMD}$ + support$_{nonCMD}$)*. This
proportion can be greater than 1 if support by non-CMDs is negative, as occurs if the likelihood
of the null model at non-CMD sites is higher than that of the positive selection model, given the
parameters of each model estimated by ML over all sites.

Sites were classified *a posteriori* as under positive selection if their Bayes Empirical
Bayes posterior probability of being in class 2 ($\omega_2>1$) under the positive selection model in
CODEML was >0.5 (moderate support) or >0.9 (strong support).

We categorized observed CMDs by the minimum number of nonsynonymous single-
nucleotide steps implied under the Goldman-Yang model between the ancestral and derived
states. For each CMD comprising two nucleotide differences, there are two paths by which they
can be interconverted by two single nucleotide steps. We determined whether the steps on these
paths would be nonsynonymous or synonymous using the standard genetic code and then
calculated the mean number of nonsynonymous steps averaged over the two paths. Paths
involving stop-codons were not included. We conducted a similar analysis for all possible
CMDs in the universal genetic code table.
545

546 BS+MNM codon substitution
547 model and test. The codon
548 substitution model of the
549 classic BST is based on the
550 Goldman-Yang (GY) model [5].

$$q_{ij} = \begin{cases} \kappa\pi_j & \text{synonymous transversion} \\ \pi_j & \text{synonymous transition} \\ \kappa\omega\pi_j & \text{non-synonymous transversion} \\ \omega\pi_j & \text{non-synonymous transition} \\ 0 & \text{two or more differences} \end{cases} \quad \text{.........} \quad (1)$$

551 Sequence evolution is modeled as a Markov process, where the matrix element q_{ij}, the
552 instantaneous rate of change from ancestral codon i to derived codon j, is defined for four types
553 of changes: synonymous transitions and transversions, and nonsynonymous transitions and
554 transversions (see q_{ij} equation 1). Three parameters are estimated from the data by maximum-
555 likelihood: ω, the ratio of nonsynonymous substitution rate to the synonymous substitution rate
556 (d_N/d_S); π_j, the equilibrium frequency of codon j; and κ, the transversion:transition rate ratio.

557 Element q_{ij} is zero for substitutions involving more than one difference, so codons with multiple
558 differences can only evolve through intermediate codons that are a single change away. A
559 scaling factor applied to the matrix ensures that branch lengths are interpreted as the expected
560 number of substitutions per codon.

561 We developed a
562 modification of the GY
563 model that incorporates
564 MNMs using the
565 parameter, δ, which
566 represents the relative
567 instantaneous rate of double
568 substitutions to that of
569 single substitutions (see q_{ij}

$$q_{ij} = \begin{cases} \kappa\pi_j & \text{synonymous transversion} \\ \pi_j & \text{synonymous transition} \\ \kappa\omega\pi_j & \text{non-synonymous transversion} \\ \omega\pi_j & \text{non-synonymous transition} \\ \omega\delta\kappa^2\pi_j & \text{non-synonymous, 2 transversions} \\ \omega\delta\pi_j & \text{non-synonymous, 2 transitions} \\ \omega\delta\kappa\pi_j & \text{non-synonymous, 1 transversion, 1 transition} \\ \delta\pi_j & \text{synonymous, 2 transitions} \\ \delta\kappa^2\pi_j & \text{synonymous, 2 transversions} \\ \delta\kappa\pi_j & \text{synonymous, 1 transversion, 1 transition} \\ 0 & \text{otherwise} \end{cases} \quad \text{.........} \quad (2)$$

570 equation 2). When $\delta = 0$, the BS+MNM model reduces to the classic BST model that does not
571 incorporate MNMs (q_{ij} equation 1). Triple substitutions have an instantaneous rate of zero.

572 The BS+MNM test of positive selection is identical to the BST, except it utilizes this
573 MNM codon model. We implemented this test by modifying the branch-site test batch file
574 (YangNielsenBranchSite2005.bf) in Hyphy 2.2.6 software by declaring δ a global variable,
575 incorporating it into the codon table, and allowing it to be optimized by ML as it other model
576 parameters are.

577 We validated the BS+MNM implementation by simulating 50 replicate alignments using
578 the BS+MNM null model in Hyphy under genome-median parameters (see below). We then
579 used the BS+MNM procedure to find the ML estimate of each parameter, including branch
580 lengths, given each alignment and the topology of the phylogeny used to generate the sequences.
581 We compared the distribution of estimates over replicates to the "true" values used to generate
582 the sequences (**Supplementary Fig. 3**).

To test if there is statistical support in the data for the BS+MNM null model relative to
the standard BST null model, we performed an LRT with 1 df, comparing the fit of the
BS+MNM null model and the BST null model on our empirical genes. Briefly, for each of the
6868 human genes, we tested if the BS+MNM null model fit the data better than the BST null
model at P<0.05 and also applied and adjustment for multiple testing (FDR<0.2). We performed
similar LRTs for each of the six terminal lineages in flies. To determine whether this test might
be prone to falsely infer support for the BS+MNM model, we simulated control sequences under
the null BST model with parameters derived from the empirical sequences and performed the
LRT as described above. Only 2 percent of genes in humans and 2.6 percent in flies yielded
significant support for BS+MNM at P < 0.05. Zero human genes and 0.006 percent of fly genes
retained significance after multiple testing adjustment (FDR <0.2). (**Supplementary Table 4**).
594
595

Simulations and analysis of false-positive bias. To characterize bias in the BST and other tests
of selection, we conducted sequence simulations in the absence of positive selection under
empirically derived conditions. We used the BS+MNM method we implemented in Hyphy to
estimate by maximum likelihood (ML) the gene-specific branch lengths and parameters of the
null BS+MNM model for every gene in the mammalian and fly datasets. We also calculated the
genome-wide median of each parameter over all genes in each dataset (the "genome-average"
parameter value). Probability density characterizations for parameters δ and gene length were
performed using the *density* function in R.

We simulated sequence evolution under the BS+MNM null model using either gene-
specific or genome-median parameters. First, we simulated a "pseudo-genome" without positive
selection by simulating one replicate of each of the 6868 and 8564 mammalian and fly
alignment, each at its empirical length, using the BS+MNM null model and the ML parameter
estimates inferred for that gene from the empirical data. We then ran the BST on these
sequences, testing for signatures of positive selection on the human lineage and each terminal fly
lineage (**Supplementary Table 2**). Control simulations were conducted under identical
conditions but with δ=0.

To test the effect of gene length on bias in the BST, we focused on genes in the BST-
significant set. For each gene's gene-specific parameters, we simulated 50 replicates alignments
of length 5,000 or 10,000 codons. We analyzed these alignments using the BST, assigning the
human branch as foreground for mammalian genes or, for flies, the same branch that produced a
significant result when the empirical data were analyzed. The false positive rate (FPR) for any
gene's parameters is the fraction of replicates yielding a positive test (P<0.05). We also repeated
these simulations and analyses using the genome-median value of δ. For control experiments
without MNMs, we set δ =0 in the simulations.

To test the effect of the rate at which MNM substitutions are produced on false positive
inference rates, we simulated evolution of alignments 5,000 codons long under the BS+MNM
null model, using genome-median estimates for all parameters except δ, which we varied. At
each value of δ, we simulated 50 replicates. We analyzed each replicate using the BST for
selection on the human or *D. simulans* lineages and calculated the proportion of replicates for
each value of δ that yielded a false positive inference ($P<0.05$).

We computed the observed proportion of tandem substitutions as a fraction of all
substitutions on the human and *D. melanogaster* lineages in both empirical and simulated
datasets. For each of the 6868 genes in the curated mammalian dataset, we aligned the human
gene to the inferred sequence of the human-chimp ancestor, identified all substitutions as
differences between these sequences, and calculated the proportion of tandem substitutions, T, as
the number of substitutions at adjacent sites divided by the sum of substitutions at adjacent sites
and those at non-adjacent sites across all sites in the dataset. Differences at adjacent sites were
counted as a single tandem substitution. For each of the 8564 genes in the fly dataset, we aligned
the *D. melanogaster* sequence to the *D. melanogaster/D. simulans* ancestor and followed the
procedure described above. For simulated sequences, we repeated this procedure using the
sequences simulated under the BS+MNM null model and parameters estimated from each gene
in the empirical datasets.
638

BUSTED. To examine the accuracy of BUSTED, we used Hyphy software 2.2.6 (batch files
BUSTED.bf and QuickSelectionDetection.bf). We analyzed the 5,000 codon-long alignments
simulated under the BS+MNM null model, using parameters estimated by ML for each BST-
significant gene, with δ assigned either to its gene-specific estimate, its genome-average, or to
zero. We applied BUSTED to the replicate alignments to test for selection ($P<0.05$) on the
human lineage or the same fly lineage that was significant for that gene in the BST of the
empirical data.
646

Power analyses. To characterize the statistical power of the BST and BS+MNM tests, we
simulated sequence evolution with positive selection of variable intensity and pervasiveness
(**Supplementary Fig. 4**). Specifically, we used the BS positive model in Hyphy to simulate
sequence evolution with the human and *D. simulans* terminal branches as the foreground
branches. We used genome-average estimates of all parameters, including gene length (418 and
510 codons for mammals and flies, respectively), but we varied ω_2 and p_2. 20 replicate
alignments were simulated under each set of conditions and then analyzed using the BST, the
BS+MNM test, or BUSTED. For each set of conditions, the true positive rate was calculated as
the fraction of replicates yielding a significant test of positive selection ($P<0.05$ for BST and
BS+MNM, FDR<0.20 for at least one site in the alignment for BUSTED).
657

BS+MNM+ κ_2 model:

We developed the

BS+MNM+ κ_2 model,

which incorporates into

the BS+MNM model (q_{ij}

equation 2) two different

transversion:transition

rate ratio parameters, κ_1

$$q_{ij} = \begin{cases} \kappa_1 \pi_j & \text{synonymous transversion} \\ \pi_j & \text{synonymous transition} \\ \omega \kappa_1 \pi_j & \text{non-synonymous transversion} \\ \omega \pi_j & \text{non-synonymous transition} \\ \omega \delta \kappa_2^2 \pi_j & \text{non-synonymous, 2 transversions} \\ \omega \delta \pi_j & \text{non-synonymous, 2 transitions} \\ \omega \delta \kappa_2 \pi_j & \text{non-synonymous, 1 transversion, 1 transition} \\ \delta \pi_j & \text{synonymous, 2 transitions} \\ \delta \kappa_2^2 \pi_j & \text{synonymous, 2 transversions} \\ \delta \kappa_2 \pi_j & \text{synonymous, 1 transversion, 1 transition} \\ 0 & \text{otherwise} \end{cases} \qquad \cdots\cdots\cdots \quad (3)$$

for single-site substitutions and κ_2 for MNMs (see q_{ij} equation 3). All free parameters of the

model are estimated by ML given a sequence alignment. This model was implemented by

further modifying our BS+MNM batchfile in Hyphy 2.2.6 software by declaring κ_2 a global

variable, incorporating it into the codon table, and allowing it to be optimized by ML as other

parameters are in the batch file.

For validation, we estimated the parameters of the BS+MNM+ κ_2 null model by ML for

every alignment in each dataset and calculated the genome-average median estimate of each

parameter (Fig. S7). We then simulated 50 replicate alignments of length 418 and 510 codons in

the mammalian and fly datasets respectively, under the BS+MNM+ κ_2 null model with all model

parameters set to their genome-wide median. We then estimated each parameter by ML under

the null model given each alignment and compared the distribution of estimates to the parameters

used to generate the alignments. We found that most parameters were estimated accurately, but

estimates of κ_2 had high variance (**Supplementary Fig. S7**), presumably because the quantity of

data in a single gene, in which CMDs are typically rare, is inadequate to support a robust

estimate of this parameter. We therefore limited our use of this model to generating sequences

by simulation rather than making inferences from sequence data.

To determine the effect of the MNM-specific transversion:transition rate on false-positive

bias in the BST, we simulated sequences 10,000 codons long under the BS+MNM+κ_2 null

model, using genome-median parameters except κ_2, which we varied. For each value of κ_2, we

simulated 50 replicates, applied the BST, and calculated the FPR as the fraction of replicates

yielding a positive inference (P<0.05).

Data availability. The empirical alignments reanalyzed in this study are available in the

supplementary information of the original publications that generated these data [12, 16, 45].

Code availability. The custom HYPHY batch codes for the BS+MNM and BS+MNM+κ_2 tests

are available as supplementary files and at

https://github.com/JoeThorntonLab/MNM_SelectionTests.

REFERENCES

1.   Goldman, N. & Yang, Z. A codon-based model of nucleotide substitution for protein-
coding DNA sequences. *Mol Biol Evol* **11**, 725-736 (1994).
2.   Murrell, B. et al. Gene-wide identification of episodic selection. *Mol Biol Evol* **32**, 1365-
1371 (2015).
3.   Murrell, B. et al. Detecting individual sites subject to episodic diversifying selection. *PLoS
Genet* **8**, e1002764 (2012).
4.   Smith, M. D. et al. Less is more: an adaptive branch-site random effects model for efficient
detection of episodic diversifying selection. *Mol Biol Evol* **32**, 1342-1353 (2015).
5.   Yang, Z. & Nielsen, R. Codon-substitution models for detecting molecular adaptation at
individual sites along specific lineages. *Mol Biol Evol* **19**, 908-917 (2002).
6.   Zhang, J., Nielsen, R. & Yang, Z. Evaluation of an improved branch-site likelihood method
for detecting positive selection at the molecular level. *Mol Biol Evol* **22**, 2472-2479 (2005).
7.   Pond, S. L., Frost, S. D. & Muse, S. V. HyPhy: hypothesis testing using phylogenies.
Bioinformatics **21**, 676-679 (2005).
8.   Kosiol, C., Holmes, I. & Goldman, N. An empirical codon model for protein sequence
evolution. *Mol Biol Evol* **24**, 1464-1479 (2007).
9.   Whelan, S. & Goldman, N. Estimating the frequency of events that cause multiple-
nucleotide changes. *Genetics* **167**, 2027-2043 (2004).
10.   Muse, S. V. & Gaut, B. S. A likelihood approach for comparing synonymous and
nonsynonymous nucleotide substitution rates, with application to the chloroplast genome.
Mol Biol Evol **11**, 715-724 (1994).
11.   Yang, Z. & dos Reis, M. Statistical properties of the branch-site test of positive selection.
Mol Biol Evol **28**, 1217-1228 (2011).
12.   Han, M. V., Demuth, J. P., McGrath, C. L., Casola, C. & Hahn, M. W. Adaptive evolution
of young gene duplicates in mammals. *Genome Res* **19**, 859-867 (2009).
13.   Drosophila, G. C. et al. Evolution of genes and genomes on the Drosophila phylogeny.
Nature **450**, 203-218 (2007).
14.   Foote, A. D. et al. Convergent evolution of the genomes of marine mammals. *Nat Genet*
47, 272-275 (2015).
15.   Kosiol, C. et al. Patterns of positive selection in six Mammalian genomes. *PLoS Genet* **4**,
e1000144 (2008).
16.   Roux, J. et al. Patterns of positive selection in seven ant genomes. *Mol Biol Evol* **31**, 1661-
1685 (2014).
17.   Zhang, J. Performance of likelihood ratio tests of evolutionary hypotheses under inadequate
substitution models. *Mol Biol Evol* **16**, 868-875 (1999).
18.   Nozawa, M., Suzuki, Y. & Nei, M. Reliabilities of identifying positive selection by the
branch-site and the site-prediction methods. *Proc Natl Acad Sci U S A* **106**, 6700-6705
(2009).
19.   Casola, C. & Hahn, M. W. Gene conversion among paralogs results in moderate false
detection of positive selection using likelihood methods. *J Mol Evol* **68**, 679-687 (2009).
20.   Anisimova, M. & Yang, Z. Multiple hypothesis testing to detect lineages under positive
selection that affects only a few sites. *Mol Biol Evol* **24**, 1219-1228 (2007).

21. Kosakovsky Pond, S. L. et al. A random effects branch-site model for detecting episodic
diversifying selection. *Mol Biol Evol* **28**, 3033-3043 (2011).
22. Zhang, J. Frequent false detection of positive selection by the likelihood method with
branch-site models. *Mol Biol Evol* **21**, 1332-1339 (2004).
23. Gharib, W. H. & Robinson-Rechavi, M. The branch-site test of positive selection is
surprisingly robust but lacks power under synonymous substitution saturation and variation
in GC. *Mol Biol Evol* **30**, 1675-1686 (2013).
24. Zhai, W., Nielsen, R., Goldman, N. & Yang, Z. Looking for Darwin in genomic sequences-
-validity and success of statistical methods. *Mol Biol Evol* **29**, 2889-2893 (2012).
25. Schrider, D. R., Hourmozdi, J. N. & Hahn, M. W. Pervasive multinucleotide mutational
events in eukaryotes. *Curr Biol* **21**, 1051-1054 (2011).
26. Saribasak, H. et al. DNA polymerase ζ generates tandem mutations in immunoglobulin
variable regions. *J Exp Med* **209**, 1075-1081 (2012).
27. Loeb, L. A. & Monnat, R. J. DNA polymerases and human disease. *Nat Rev Genet* **9**, 594-
604 (2008).
28. Matsuda, T., Bebenek, K., Masutani, C., Hanaoka, F. & Kunkel, T. A. Low fidelity DNA
synthesis by human DNA polymerase-eta. *Nature* **404**, 1011-1013 (2000).
29. Seplyarskiy, V. B., Bazykin, G. A. & Soldatov, R. A. Polymerase ζ Activity Is Linked to
Replication Timing in Humans: Evidence from Mutational Signatures. *Mol Biol Evol* **32**,
3158-3172 (2015).
30. Stone, J. E., Lujan, S. A., Kunkel, T. A. & Kunkel, T. A. DNA polymerase zeta generates
clustered mutations during bypass of endogenous DNA lesions in Saccharomyces
cerevisiae. *Environ Mol Mutagen* **53**, 777-786 (2012).
31. Arana, M. E., Seki, M., Wood, R. D., Rogozin, I. B. & Kunkel, T. A. Low-fidelity DNA
synthesis by human DNA polymerase theta. *Nucleic Acids Res* **36**, 3847-3856 (2008).
32. Besenbacher, S. et al. Multi-nucleotide de novo Mutations in Humans. *PLoS Genet* **12**,
e1006315 (2016).
33. Chen, J. M., Férec, C. & Cooper, D. N. Complex Multiple-Nucleotide Substitution
Mutations Causing Human Inherited Disease Reveal Novel Insights into the Action of
Translesion Synthesis DNA Polymerases. *Hum Mutat* **36**, 1034-1038 (2015).
34. Chen, J. M., Cooper, D. N. & Férec, C. A new and more accurate estimate of the rate of
concurrent tandem-base substitution mutations in the human germline: ~0.4% of the single-
nucleotide substitution mutation rate. *Hum Mutat* **35**, 392-394 (2014).
35. Harris, K. & Nielsen, R. Error-prone polymerase activity causes multinucleotide mutations
in humans. *Genome Res* **24**, 1445-1454 (2014).
36. Hodgkinson, A. & Eyre-Walker, A. Variation in the mutation rate across mammalian
genomes. *Nat Rev Genet* **12**, 756-766 (2011).
37. Francioli, L. C. et al. Genome-wide patterns and properties of de novo mutations in
humans. *Nat Genet* **47**, 822-826 (2015).
38. Assaf, Z. J., Tilk, S., Park, J., Siegal, M. L. & Petrov, D. A. Deep sequencing of natural and
experimental populations of Drosophila melanogaster reveals biases in the spectrum of new
mutations. *Genome Res* **27**, 1988-2000 (2017).
39. Zhu, W. et al. Concurrent nucleotide substitution mutations in the human genome are
characterized by a significantly decreased transition/transversion ratio. *Hum Mutat* **36**, 333-
341 (2015).
40. Averof, M., Rokas, A., Wolfe, K. H. & Sharp, P. M. Evidence for a high frequency of

simultaneous double-nucleotide substitutions. *Science* **287**, 1283-1286 (2000).

41. Bazykin, G. A., Kondrashov, F. A., Ogurtsov, A. Y., Sunyaev, S. & Kondrashov, A. S.
Positive selection at sites of multiple amino acid replacements since rat-mouse divergence.
Nature **429**, 558-562 (2004).

42. Rogozin, I. B. et al. Evolutionary switches between two serine codon sets are driven by
selection. *Proc Natl Acad Sci U S A* **113**, 13109-13113 (2016).

43. Suzuki, Y. False-positive results obtained from the branch-site test of positive selection.
Genes Genet Syst **83**, 331-338 (2008).

44. De Maio, N., Holmes, I., Schlötterer, C. & Kosiol, C. Estimating empirical codon hidden
Markov models. *Mol Biol Evol* **30**, 725-736 (2013).

45. Larracuente, A. M. et al. Evolution of protein-coding genes in Drosophila. *Trends Genet*
24, 114-123 (2008).

46. Sironi, M., Cagliani, R., Forni, D. & Clerici, M. Evolutionary insights into host-pathogen
interactions from mammalian sequence data. *Nat Rev Genet* **16**, 224-236 (2015).

47. Bloom, J. D. An experimentally determined evolutionary model dramatically improves
phylogenetic fit. *Mol Biol Evol* **31**, 1956-1978 (2014).

48. Lopez, P., Casane, D. & Philippe, H. Heterotachy, an important process of protein
evolution. *Mol Biol Evol* **19**, 1-7 (2002).

49. Pond, S. K. & Muse, S. V. Site-to-site variation of synonymous substitution rates. *Mol Biol*
Evol **22**, 2375-2385 (2005).

50. Barber, M. F. & Elde, N. C. Nutritional immunity. Escape from bacterial iron piracy
through rapid evolution of transferrin. *Science* **346**, 1362-1366 (2014).

51. Chan, Y. F. et al. Adaptive evolution of pelvic reduction in sticklebacks by recurrent
deletion of a Pitx1 enhancer. *Science* **327**, 302-305 (2010).

52. Field, S. F., Bulina, M. Y., Kelmanson, I. V., Bielawski, J. P. & Matz, M. V. Adaptive
evolution of multicolored fluorescent proteins in reef-building corals. *J Mol Evol* **62**, 332-
339 (2006).

53. Barrett, R. D. & Hoekstra, H. E. Molecular spandrels: tests of adaptation at the genetic
level. *Nat Rev Genet* **12**, 767-780 (2011).

54. Siddiq, M.A, Loehlin, D.W., Montooth, K.L., Thornton J.W. Experimental test and
refutation of a classic case of molecular adaptation in *Drosophila melanogaster*. Nature
Ecology and Evolution 1, doi:10.1038/s41559-016-0025 (2017)

Figure 1 Codons with multiple nucleotide differences (CMDs) drive branch-site signatures of
selection.

(a) CMDs are enriched in genes with a signature of positive selection. Codons were
classified by the number of nucleotide differences between the ancestral and terminal
states on branches tested for positive selection. CMDs have $\geq$2 differences; non-CMDs
have $\leq$1 difference. The CMD/non-CMD ratio is shown for genes with a significant
signature of selection in the BST (BST-sig) and those without (BST-ns). Fold-enrichment is
shown as the odds ratio. *, P=4e-4 by χ^2 test; **, P=1e-41 by Fisher's exact test.
(b) Percentage of genes that retain a signature of positive selection when CMDs are excluded
from the branch-site test analysis.
(c) Distribution across BST-significant genes of the proportion of total support for the positive
selection model that is provided by CMDs. Total support is the difference in log-likelihood
between the positive selection and null models, summed over all codons in the alignment.
Support from CMDs is summed over codons with multiple differences. The proportion of
support from CMDs can be greater than 1 if the log-likelihood difference between models is
negative at non-CMDs.
(d) Most codons classified as positively selected are CMDs. The number of CMDs and non-
CMDs in BST-significant genes are shown according to their Bayes Empirical Bayes
posterior probability (PP) of being in the positively selected class.

Figure 2 Incorporating MNMs into the branch-site model eliminates the signature of positive selection in many genes. The mammalian and fly datasets were reanalyzed using a version of the BST that allows MNMs (BS+MNM) by including a parameter δ, a multiplier on the rate of each double substitution relative to single substitutions.

(a) The distribution of ML estimates of δ across genes with (white) and without (black) a significant result in the classic BST is shown for empirical alignments. Median estimates of δ in BST-significant and BST-nonsignificant genes are 0.047 and 0.026 in humans, respectively, and 0.107 and 0.062 in flies. *, P=6.7e-4; **, P=1e-8 by Mann-Whitney U Test.

(b) Proportion of genes with a significant result in the BST that lose or retain that signature using the BS+MNM test. Genes that remain significant but contain CMDs with three differences, which are not incorporated into BS+MNM, are also shown.

857
Figure 3 MNMs cause a strong bias in the branch-site test under realistic conditions.   For each
gene in the mammalian and fly datasets, the parameters of the BS+MNM null model were
estimated by maximum likelihood. We then simulated sequence evolution under each gene's
inferred null parameters and used the classic BST on the simulated alignments to test for positive
selection on the human and terminal fly lineages.
(a) The fraction of all tests that are BST-significant (P<0.05) is shown for the data simulated
under the BS+MNM null model, the original empirical sequence alignments, and a control
dataset simulated with $\delta = 0$. Each gene's length in the simulation was identical to its
empirical length.
(b) BST-significant genes are longer than BST non-significant genes. The probability density of
gene lengths in the two categories is shown for the empirical mammalian and fly datasets.
Median lengths in BS-significant and non-significant genes, respectively, were 642 and 343
bp in humans; in flies, 448 and 399 bp. The difference between the two distributions was
evaluated using a Mann-Whitney U test. *, P=8e-4; **, P=8e-5;
(c) Systematic bias in the BST. For each gene with a significant result in the BST using the
empirical data, we simulated 50 replicates using the BS+MNM null model and the ML
parameter estimates for that gene at lengths of 5,000 and 10,000 codons; these data were then
analyzed using the BST. The false positive rate (FPR) for any gene's simulation (black
points) is the proportion of replicates with P<0.05. Gray points show FPR for control
simulations with $\delta = 0$. Dashed lines, FPR of 0.05. The solid diagonal line has a slope of 1.
(d) The distribution of ML estimates of δ across genes with (white) and without (black) a
signature of positive selection in the classic BST is shown for data simulated under the

BS+MNM null model. Median δ in BST-significant and BST-nonsignificant genes = 0.03
and 0.0009 in humans, 0.04 and 0.08 in flies. Difference between the distributions was
evaluated using a Mann-Whitney U Test. ***, P=1e-12; ****, P=1e-199.

(e) Increasing the MNM rate increases bias in the BST. Sequences 5,000 codons long were
883   simulated using the BS+MNM model and the median value of each model parameter and
884   branch length across all genes in each dataset, but δ was allowed to vary. The rate of false
885   positives (P<0.05) in 50 replicates at each value of δ is shown. Solid line, hyperbolic fit to
886   the data; dotted line, FPR level of 5%.   Arrowhead, median δ across all genes.

(f) Relationship between δ and inferred ω_2. Sequences simulated in (e) were used to infer
888   the ω_2 estimated by BST under the positive selection model, and the relationship
889   plotted on a log-log scale. The best-fit linear regression line is shown along with the
890   coefficient of determination.

894
Figure 4 Transversion-enrichment in CMDs biases the BST.
(a) The ratio of transversions:transitions observed in CMDs and in non-CMDs is shown for
BST-significant and BST-nonsignificant genes. Fold-enrichment is shown as the odds ratio.
*, P=5e-4; **, P=3e-25 by Fisher's exact test.
(b) Increasing the transversion rate in MNMs increases bias of the BST. Sequences 10,000
codons long were simulated using an elaboration of the BS+MNM model that allows MNMs
to have a transversion:transition rate (κ_2) different from that in single-nucleotide substitutions
(κ_1).   50 replicate alignments were simulated under the null model using the average value
of every model parameter and branch length across all genes in each dataset, except κ_2 was
allowed to vary. The rate of false positives (P<0.05) at each value of κ_2 is shown.
Arrowheads show the false positive rate when sequences were simulated with κ_2 equal to κ_1.
Dotted line, FPR of 5%.
907

908
Figure 5 MNMs bias a newer test of positive selection. False positive inferences under realistic
conditions using BUSTED. For every BST-significant gene in each dataset, 50 replicate
alignments 5,000 codons long were simulated using the BS+MNM null model and parameter
values estimated from the empirical sequences. These alignments were then analyzed for a
signature of positive selection ($P<0.05$) using BUSTED. δ was assigned to its gene-specific
estimate, to its average across all genes in each dataset, or to zero. FPR is the proportion of
replicate alignments for each gene with $P<0.05$. Each dot represents the FPR for one gene; black
bars are the median across genes.
917
918

Figure 6 CMDs implying multiple nonsynonymous steps drive the BST.

(a) For every CMD, the mean of the number of nonsynonymous single-nucleotide steps on the two direct paths between the ancestral and derived states was calculated.   In BST-significant and BST-nonsignificant genes, the ratio of CMDs invoking more than one nonsynonymous step to those invoking one or fewer such steps is shown. Fold-enrichment is shown as the odds ratio. *, P=9e-04; **P= 1.6e-67 by Fisher's exact test.

(b) Support for the positive selection model provided by CMDs depends on the number of implied nonsynonymous single-nucleotide steps. Support is the log-likelihood difference between the positive selection and null models of the BST given the data at a single codon site. Box plots show the distribution of support by CMDs in BST significant genes categorized according to the mean number of implied nonsynonymous steps. Dotted line, support of 1.92, at which the BST yields a significant result for an entire gene (P<0.05). In human BST-significant genes, no CMDs imply zero non-synonymous changes.