Selection on non-antigenic gene segments of seasonal influenza A virus and its

impact on adaptive evolution

Jayna Raghwani[1*], Robin Thompson[1], Katia Koelle[2]

[1] Department of Zoology, University of Oxford, Oxford, United Kingdom

[2] Department of Biology, Duke University, Durham, United States

* Corresponding author: jayna.raghwani@zoo.ox.ac.uk

Keywords: Seasonal influenza A/H3N2, virus adaptation, reassortment, linkage

effects

ABSTRACT

Most studies on seasonal influenza A/H3N2 virus adaptation have focused on the

main antigenic gene, haemagglutinin. However, there is increasing evidence that the

genome-wide genetic background of novel antigenic variants can influence these

variants' emergence probabilities and impact their patterns of dominance in the

population. This suggests that non-antigenic genes may be important in shaping the

viral evolutionary dynamics. To better understand the role of selection on non-

antigenic genes in the adaptive evolution of seasonal influenza viruses, we here

develop a simple population genetic model that considers a virus with one antigenic

and one non-antigenic gene segment. By simulating this model under different

regimes of selection and reassortment, we find that the empirical patterns of lineage

turnover for the antigenic and non-antigenic gene segments are best captured when

there is both limited viral coinfection and selection operating on both gene segments.

In contrast, under a scenario of only neutral evolution in the non-antigenic gene

segment, we see persistence of multiple lineages for long periods of time in that

segment, which is not compatible with the observed molecular evolutionary patterns.

Further, we find that reassortment, occurring in coinfected individuals, can increase

the speed of viral adaptive evolution by primarily reducing selective interference and

genetic linkage effects mediated by the non-antigenic gene segment. Together, these

findings suggest that, for influenza, with 6 internal or non-antigenic gene segments,

the evolutionary dynamics of novel antigenic variants are likely to be influenced by

the genome-wide genetic background as a result of linked selection among both

beneficial and deleterious mutations.

INTRODUCTION

Seasonal influenza is a major infectious disease that causes 3 to 5 million worldwide cases of severe illness and 250,000 to 500,000 deaths each year in humans (1). Of the currently circulating flu viruses, influenza A subtype H3N2 is the predominant virus contributing to these morbidity and mortality estimates. This virus is known to rapidly evolve, particularly antigenically (2), enabling it to perpetually evade herd immunity and re-infect individuals in the population. Consequently, there has been great interest in understanding how this virus evolves antigenically, especially with respect to its main antigenic gene, haemagglutinin (HA). In particular, these investigations have focused on identifying key sites involved in viral antigenicity (3-6), which has provided compelling evidence of immune-mediated selection acting upon HA.

However, the limited standing genetic diversity observed for HA has been difficult to reconcile based on recurrent positive selection alone, since the high virus mutation rate and the presence of strong diversifying selection predicts a large antigenic repertoire over time (7). The observed low-level genetic diversity of the HA is reflected in its spindly, ladder-like phylogeny, which indicates that only a single viral lineage persists over time. Genetic variants belonging to this persisting lineage have been characterized antigenically, indicating that every two to eight years a major antigenic change occurs that necessitates the updating of components of the seasonal influenza vaccine (6, 8, 9). Phylodynamic models have proven to be invaluable to understanding how host immunity and viral evolution can lead to these interesting phenomena of a spindly phylogeny and a single major circulating antigenic variant dominating global infection dynamics (7, 10-12). While these models differ in their

specific explanations of what processes shape this restricted antigenic evolution of

influenza A/H3N2, they in general have had to either impose strong among-strain

competition for susceptible hosts (7, 11, 12) and/or limit the antigenic mutation rate

(10, 12). More recent work on the molecular evolution of the HA indicates that clonal

interference and background selection are also important determinants of the adaptive

dynamics of the HA (13-17).

While it is clear that the evolution of HA is a key component of influenza A/H3N2's

adaptive evolution, the role of other gene segments, in particular those that encode

internal proteins, is less well understood. There is a small but growing number of

studies that indicate that selection also acts on viral phenotypes beyond antibody-

mediated immune escape. For example, the appearance and dominance of the CA04

antigenic lineage is attributed in part to the increased replicative fitness and virulence

conferred by two amino-acid substitutions in the polymerase acidic (PA) gene

segment (18). There is also evidence that cytotoxic T-lymphocyte (CTL) immune

pressure can exert selection pressure on influenza A virus. Specifically, recent work

has shown that adaptive substitutions in the nucleoprotein (NP) gene predominantly

occur at T-cell epitopes (19, 20).

Interestingly, the genetic diversity of internal or non-antigenic genes in influenza

A/H3N2 virus is also limited, although to a lesser extent than for HA (21). One

explanation for this observation is that these gene segments are in strong linkage with

HA, which means that any evolutionary force that reduces genetic diversity of the HA

(e.g., selective sweeps and genetic bottlenecks) will also similarly impact the rest of

the virus genome. However, whole-genome analyses of seasonal influenza A viruses

indicate that reassortment is relatively frequent, with each gene segment having

somewhat of a distinctive evolutionary history (21-25). Estimated differences in the

times to most recent common ancestor (TMRCAs) across the genome can also exceed

six years (21), which is inconsistent with strong linkage effects solely shaping the

genetic diversity patterns of this virus. An alternative explanation for the limited

genetic diversity of non-antigenic gene segments is selection. Although there are

several distinct models that can generate the restricted diversity of HA by invoking

selection (10-12, 14, 15, 26), there has been very little consideration of whether

selection also contributes to shaping the evolutionary dynamics of non-antigenic

genes.

Here we evaluate the importance of selection on non-antigenic gene segments in the

adaptive evolution of seasonal influenza A/H3N2 by analyzing the evolutionary

dynamics of the viral genome and using a population genetic model to determine the

critical processes that can reproduce features of these observed evolutionary

dynamics. The main questions we address are whether selection on non-antigenic

gene segments impact the evolutionary dynamics of the non-antigenic gene segments

themselves, and through linkage effects, the antigenic gene segments. Instead of

examining the complexity of 8 distinct gene segments, we simplify our model by

considering a virus that contains only two gene segments, corresponding to one

antigenic gene (e.g., HA) and one non-antigenic gene (e.g., PA). By simulating the

model such that lineages can be traced back in time, we examine the patterns of

genetic diversity of the virus across different assumptions of selection and

reassortment. We find that selective effects on both gene segments and limited

reassortment (via limited coinfection rates) are necessary to capture the key TMRCA

patterns of influenza A/H3N2 virus genome. Furthermore, we find that the rate of

adaptive evolution of the virus increases under this evolutionary regime, which is

predominantly a result of reassortment reducing interference effects contributed by

the non-antigenic gene segment.

MATERIALS AND METHODS

A) Evolutionary dynamics of seasonal influenza A/H3N2 virus genome

To characterize the evolutionary dynamics of A/H3N2, we used a published global

whole-genome dataset of viruses sampled from 1977 to 2009 ($n = 676$) (27). Time-

scaled trees were estimated with BEAST v1.8 (28) by employing a relaxed

uncorrelated log-normal distributed molecular clock (29), a codon-structured

nucleotide substitutional model (30), and a Bayesian Skygrid coalescent prior (31).

Two independent chains of 200 million steps were executed for each of the eight gene

segments to ensure that adequate mixing and stationarity had been achieved. The

posterior tree distribution for each segment was further examined with PACT (32),

which infers the times to the most recent common ancestor (TMRCAs) across the

entire evolutionary history at regular intervals. To quantify and visualize patterns of

genetic diversity in each segment, mean TMRCAs over time were plotted using the R

package ggplot2 (33) and genealogical trees were plotted with ggtree (34).

B) Phylodynamic model of infection and coinfection

To explore the evolutionary processes underlying the empirical patterns of TMRCA

observed for the influenza A/H3N2 virus genome, we formulated a simple population

genetic model with a constant number of $N = 1000$ infected individuals. In the model,

individuals were either infected with a single virus (I_s) or coinfected with two viruses

(I_{co}). We did not consider coinfection with more than two viruses. The virus genome

consisted of one antigenic segment and one non-antigenic segment.

We simulated the infected population of hosts over time using a modified Moran

model. Specifically, we allowed for two types of infection events: 'infection/recovery'

events and coinfection events. When an 'infection/recovery' event occurred, an

infected individual (I_s or I_{co}) was chosen to generate a new singly-infected individual.

If it was a coinfected individual generating the new infection, that individual

transmitted either of the viruses he was infected with or a reassortant virus (Figure

1A). We assumed an equal probability of each viral gene segment being transmitted,

such that a reassortant strain was transmitted 50% of the time. At the same time as the

generation of the new infection occurred, recovery of a randomly chosen infected

individual (I_s or I_{co}) also occurred. If a coinfected individual was chosen to recover, he

cleared both infecting viral strains. Because infection events were always offset by

recovery events, as is traditional in Moran models where a 'birth' is always offset by a

'death', the total number of infected individuals in the population remained constant

(Figure 1B). Infection/recovery events occurred at a rate of $\alpha = 0.25$ per capita per

day, reflecting a typical duration of influenza infection of approximately 4 days (35).

Coinfection events were marked by singly-infected individuals infecting other singly-

infected individuals (Figure 1B). Coinfection events occurred from singly infected

individuals at a per capita rate of $\beta = 0.0125$ per day. This corresponds to a

coinfection level of approximately 5% of the total infected population at equilibrium

(see Text S1). Ascertaining an empirical coinfection rate for influenza A/H3N2

viruses in general, or at the within-subtype level, is very difficult, since the low

circulating viral diversity is likely to limit our ability to distinguish between

independent infecting viral strains. Nevertheless, the number of influenza coinfections

can be estimated when viral strains involved belong to either different subtypes or

types (e.g. A/H3N2 and A/H1N1 or influenza A and B viruses, respectively) (36, 37).

These types of coinfection have been known to occur between 1-2% in sampled

influenza A viral infections (36-38). We set the level of coinfection in our model

slightly higher than these empirical estimates, at ~5%, to reflect that these empirical

estimates between different subtypes or types are likely underestimates.

185
Figure 1: Schematic of the events in the population genetic model
(A) Infection transmission by a coinfected individual. When a coinfected individual transmits,
each gene segment is randomly chosen from the two viral strains present in that individual.
Consequently, there is an equal probability of transmitting a non-reassortant strain (two
strains on the left) as there is of transmitting a reassortant strain (two strains on the right). (B)
Schematic of the main events: infection/recovery and coinfection. Infection and recovery are
coupled, such that the infected population remains constant. Upon infection (indicated by
curly arrows), singly-infected individuals (blue circles) and coinfected individuals (yellow
circles) generate new singly-infected individuals. Recovery of singly-infected and coinfected
individuals removes them from the population (denoted by dashed arrows). Here, two
infection/recovery events are shown that occur in the same τ time step. Coinfection events
occur when a singly-infected individual infects another singly-infected individual. This results
in a new coinfected individual in the infected population, carrying two viral strains.

Coinfection events result in an increase in the number of coinfected individuals in the
population and a decrease in the number of singly-infected individuals.
201

Evolution of the antigenic and non-antigenic gene segments

We let mutations occur at transmission events, which consist of both 'infection'

events and 'coinfection' events. We let the number of new mutations present in the

transmitting virus be Poisson-distributed with mean $U = 0.1$, with each mutation

being equally likely to land on the antigenic or the non-antigenic gene segment. We

allow the distribution of mutational fitness effects to differ between the two gene

segments. Specifically, we assume that 30% of mutations are beneficial and 70% of

mutations are deleterious on the antigenic gene segment. On the non-antigenic gene

segment, we assume that 5% of mutations are beneficial, 30% of mutations are

deleterious, and the remaining 65% of mutations are neutral. A higher proportion of

beneficial mutations are assumed in the antigenic gene segment to capture the

selective advantage that antigenic mutations are likely to have through evasion of herd

immunity. The non-antigenic gene segment is assumed to have a greater proportion of

neutral mutations to reflect the observation that internal genes undergo greater neutral

evolution than external genes (25). We assume that the fitness effects for beneficial

mutations are exponentially distributed with mean 0.03 and that the fitness effects for

deleterious mutations are exponentially distributed with mean 0.09. We do not

consider lethal mutations. Importantly, the distributions of mutational fitness effects

on the antigenic and non-antigenic gene segment capture the salient features of

recently determined mutational fitness effects for seasonal influenza A virus (39) (see

Figure S1).

Viral fitness is calculated by multiplying fitness values at each site across the genome.

Multinomial sampling based on viral fitness is applied at each transmission event to

determine which individual will infect (or coinfect) next. For coinfected individuals,

we initially determine which virus is transmitted from the two infecting parental viral

strains (see Figure 1A) and compute the viral fitness accordingly.

Tracking lineages over time

The model is implemented in Java using a Gillespie tau-leap algorithm (40) for

computational efficiency with a time step τ of 0.25 days. Starting from an equilibrium

number of singly- and coinfected individuals (Text S1), we run each simulation for 60

years, analysing results only from the last 20 years.

To be able to infer the genealogical history of the viral population, we track in our

model who-infected-whom at the level of infected individuals and for each gene

segment. A random sample of 100 singly-infected individuals is used to infer the

TMRCA of each gene segment at yearly intervals. Viral gene genealogies are

reconstructed from the last twenty years of simulation using a random sample of 300

singly-infected individuals. The tracked infection histories are used to determine the

first 'coalescent' event, which corresponds to finding the two sampled individuals that

shared the most recent common ancestor for a given gene segment. Specifically, this

process involves tracing back the transmission events from the sampled infections,

and establishing the parental virus in common with the most recent transmission time.

This procedure is repeated until all sampled and ancestral lineages reach the parental

viral infection that represents the most recent common ancestor of the entire sample.

RESULTS

Genealogical diversity of seasonal influenza A/H3N2 virus

Figure 2 shows how the genealogical diversity of seasonal influenza A/H3N2 varies

over time for each gene segment. We observe that the mean TMRCA of the HA gene

segment (1.90 years) is 0.5-1.1 years younger than the other gene segments, indicating

that HA experiences the fastest lineage turnover in the virus genome. The maximum

TMRCA for this gene segment also does not exceed 5 years. These TMRCA patterns

reflect that the HA gene genealogy has a single viral lineage dominating over time

(Figure S2). NA is found to have the second lowest mean TMRCA (2.4 years),

indicative of slightly longer lineage persistence than HA (Figure S2). The non-

antigenic gene segments of A/H3N2 are marked by larger mean TMRCAs and by

more extensive variation in genealogical diversity over time, indicating that multiple

lineages can co-exist for significant periods, e.g. up to ~7 years in M1 (Figure S2).

Together, these observations are compatible with positive selection predominantly

acting upon the antigenic genes, most notably the HA.

Figure 2: TMRCA through time plots for individual seasonal influenza A/H3N2 gene
segments
The mean TMRCA over time is estimated from a posterior tree distribution for each gene
segment at 6-month intervals. The black dashed lines indicate the overall mean TMRCA for
the focal gene segment in each subplot. The gray dashed lines in the non-HA gene segment
subplots show the overall mean TMRCA for the HA.

Evolutionary dynamics when only antigenic gene segment is under selection

To better understand the patterns of genealogical diversity of the influenza A/H3N2

virus genome, we first simulated the described model under the assumption that the

adaptive evolution of the virus is restricted to the antigenic gene segment, with all

mutations on the non-antigenic gene segment assumed to be neutral. Further,

coinfection was not permitted ($\beta = 0$). Figure 3 shows results from a representative

simulation. Viruses with higher fitness constantly emerge and become dominant in the

population over time (Figure 3A). At any given time, significant fitness variation is

present in the population, with lower fitness viruses able to persist in the population

over extended periods (Figure 3A).

Figure 3: Adaptive evolution in the absence of coinfection and when selection acts only
on the antigenic gene segment
(A) The distribution of fitness in the viral population over time. (B) Mean (log) fitness of the
virus population over time. (C) Gene genealogy reconstructed from model simulation by
sampling 300 singly-infected individuals over 20 years, following a burn-in of 40 years.
291

Interestingly, the simulated viral population evolves in a punctuated manner, as

indicated by the change in mean (log) population fitness over time (Figure 3B). This

suggests that the tempo of adaptive evolution varies over time. If we consider only

beneficial mutations on the antigenic gene segment, we instead observe a smooth and

continual increase in the mean population fitness over time (Figure S3). These results

indicate that more complex evolutionary dynamics can emerge with a broader

distribution of fitness effects. Lastly, consistent with previous studies (15-17), this

evolutionary regime where clonal interference and background selection are present

reproduces HA's spindly phylogeny (Figure 3C).

Under the model with both positive and negative fitness effects on only the antigenic

gene segment, no significant changes in the mean TMRCA of the antigenic gene

segment occur with increasing levels of coinfection (Figure 4A). In contrast, the mean

TMRCA of the non-antigenic gene segment becomes notably greater as the rate of

coinfection increases. These results are consistent with reassortment reducing the

hitchhiking of the neutrally evolving non-antigenic gene segment with the non-

neutrally evolving antigenic gene segment. As a consequence, the non-antigenic gene

segment is able to explore more genetic backgrounds, which leads to an increase in its

genetic diversity. Expectedly, the rate of adaptive evolution is unaffected by changes

in the rate of coinfection (Figure 4B).

Figure 4: Genealogical diversity and the rate of adaptive evolution at different levels of coinfection when selection acts only on the antigenic gene segment
(A) Distribution of TMRCAs of the antigenic (red) and non-antigenic (blue) gene segments at different levels of coinfection. Three different coinfection levels were considered: 0% ($\beta = 0$), 5% ($\beta = 0.0125$), and 9% ($\beta = 0.025$). 500 simulations were used to obtain the TMRCA distribution at each of the three coinfection levels. The dashed lines show the mean TMRCA for the focal gene segment in each subplot. (B) Mean (log) fitness of the virus population over time under different coinfection rates.

Reassortment increases the rate of adaptive evolution when a non-antigenic gene segment is under selection

Next, we examined the behaviour of the model when selection occurs on both gene segments. First, we looked at the changes in mean population fitness and fitness variation over time under increasing coinfection rates ($\beta = 0$, 0.0125, and 0.025 per day), for the whole virus (Figure 5A), the antigenic gene segment (Figure 5B), and the non-antigenic gene segment (Figure 5C). Strikingly, the rate of virus adaptation is

significantly greater in the presence of coinfection than when it is absent (Figure 5A).

This phenomenon appears to be primarily driven by the non-antigenic gene segment,

which also experiences a notably higher rate of adaptive evolution when coinfection

occurs in the population (Figure 5C). In contrast, although coinfection increases the

fitness variation of both gene segments (Figure 5B and C), the difference in the rate of

adaptive evolution of the antigenic gene segment in the absence versus in the presence

of coinfection appears to be slight (Figure 5B).

Figure 5: Adaptive evolution of the virus when both gene segments undergo selection at
varying levels of coinfection
The mean (log) population fitness and population fitness variation are shown for (A) the
whole virus, (B) the antigenic gene segment, and (C) the non-antigenic gene segment for
three different coinfection rates, corresponding to $\beta = 0$, $\beta = 0.0125$, and $\beta = 0.025$.

Together, these results indicate that when coinfection is absent the non-antigenic gene

segment experiences greater selective interference (both among beneficial and

deleterious mutations) and genetic hitchhiking than the antigenic gene segment, for

the simple reason that there are significantly more mutations with selective effects on

the latter. In other words, when there is strong linkage between the two segments,

selection on the antigenic gene segment will have a larger impact on the non-antigenic

gene segment than the non-antigenic gene segment will have on the antigenic gene

segment. As a consequence, while reassortment is expected to reduce linkage effects

between both gene segments, larger gains in fitness are more likely for the non-

antigenic gene segment as it can explore comparatively more advantageous genetic

backgrounds.

These evolutionary dynamics also have an impact on the mean TMRCAs of both gene

segments (Figure 6). Although this pattern is more discernible for the non-antigenic

gene segment, it does indicate that the antigenic gene segment is influenced to some

degree by linkage effects from the non-antigenic gene segment. The larger increase in

the mean TMRCA for the non-antigenic gene segment is also consistent with the non-

antigenic gene segment experiencing comparatively greater linkage effects than the

antigenic gene segment.

Figure 6: Genealogical diversity of the virus when selection acts on both gene segment at
varying levels of coinfection
The distribution of TMRCAs for antigenic (red) and non-antigenic (blue) gene segments for
three different coinfection rates: $\beta = 0$, $\beta = 0.0125$, and $\beta = 0.025$. As in Figure 4, 500
simulations were used to obtain each TMRCA distribution at each of the three coinfection
levels. The dashed lines show the mean TMRCAs for the each of the two gene segments in
each subplot. The solid black lines in the subplots indicates the mean TMRCA of the gene
segment when there is no coinfection.
369

Empirical genealogical patterns are compatible with selection on both antigenic

and non-antigenic gene segments

Figure 7 shows representative gene genealogies for the antigenic and non-antigenic

gene segments in the presence of coinfection ($\beta = 0.0125$) and, in both cases, when

selection acts on the antigenic gene segments. The genealogies of the antigenic gene

segment when the non-antigenic gene segment evolves either selectively (Figure 7A)

or neutrally (Figure 7D) are topologically similar, with a single lineage persisting over

time in both cases. In contrast, quite different gene genealogies are observed for the

non-antigenic gene segment (Figures 7B and E). Specifically, the non-antigenic gene

segment is associated with significantly lower genealogical diversity when it is under

selection (Figure 7B) compared to when it is not (Figure 7E). In both cases, however,

there is still greater persistence of multiple lineages relative to the antigenic gene

segment, indicative of slower population turnover.

Figure 7: Representative gene genealogies and TMRCA dynamics from simulations when the non-antigenic gene segment evolves either neutrally or under selection
(A-C) Results obtained from model simulations with coinfection and selective effects occurring on both gene segments. (D-F) Results obtained from model simulations with coinfection but no selection on the non-antigenic gene segment. Panels A and D depict the gene genealogies of the antigenic gene segment. Panels B and E depict the gene genealogies of the non-antigenic gene segment. Panels C and F show the TMRCA dynamics of both gene segments over time. The red and blue lines correspond to the antigenic and non-antigenic gene segments, respectively. (G-I) Inferred influenza A/H3N2 MCC phylogenies for the HA gene segment and the PA gene segment, along with their TMRCA dynamics. HA is the dominant antigenic gene segment. PA is a non-antigenic gene segment. The TMRCA dynamics for HA and PA are shown in panel I in orange and light blue lines, respectively.

When comparing TMRCA patterns between the antigenic gene segment and the non-antigenic gene segment, it is notable that when the non-antigenic gene segment evolves neutrally, the common ancestor of the non-antigenic gene segment is consistently older than the antigenic gene segment (Figure 7F). However, when

selection affects both gene segments, we note a closer correspondence with the

empirical TMRCA dynamics (Figure 7C, compared to Figure 7I)). Specifically, in

addition to the antigenic gene segment undergoing more frequent fluctuations in the

TMRCA over time compared to the non-antigenic gene segment, the TMRCA of both

gene segments can occasionally coincide, which likely indicates a shared common

ancestor, perhaps as a result of a genome-wide selective sweep. We further examined

this observation by comparing the differences in TMRCA between the gene segments

(Figure S4). The higher density around zero years of difference in the TMRCA

suggests that the likelihood of sharing a common ancestor is greater when selective

effects occur on both gene segments (Figure S4).

Sensitivity of results to model parameters

a) Infected population size

While it is well established that human influenza A/H3N2 virus has a strong seasonal

transmission pattern in some populations, we decided to model a constant infected

population. This decision was motivated largely by undertaking a simple and standard

approach to examine the patterns of viral diversity due to selection, mutation, and

reassortment alone. However, given that regions with low-level, constant disease

transmission (e.g. the tropics) frequently seed seasonal outbreaks in temperate locales

(21, 41-43), the effective population size of global influenza A/H3N2 viruses is

expected to be relatively small and constant over time (21, 42). Consequently, the

assumption of a constant infected population size is not unreasonable since regions

with year-round influenza infection are expected to ultimately shape the overall

evolutionary dynamics of the virus. We tested the effects of population size on the

evolutionary behavior of the model (Figure S5). Specifically, we ran 100 simulations

at each of three population sizes (N=1000, N = 5000, and N=10000), under the model

parameterization with both gene segments experiencing positive and negative

selection. Notably, similar TMRCA patterns were observed regardless of population

size, such that the antigenic gene segment typically had a younger TMRCA compared

to the non-antigenic gene segment (Figure S5). However, as we increase the

population size, the TMRCA of both gene segments increases, indicating greater

lineage persistence in the population. This corroborates a standard expectation from

coalescent theory: smaller populations have comparatively more recent common

ancestors than larger populations due to stronger effects of genetic drift. Thus, when

the infected population size is fixed at N=1000, deterministic and stochastic forces

will shape the population's genetic diversity, both of which have been implicated in

the evolutionary dynamics of seasonal influenza A viruses (21). At larger population

sizes, we can, however, recover lower TMRCAs when we increase the mean effect

size of mutations (results not shown).

b) Mutation and coinfection rates

As it is difficult to ascertain the per-genome, per transmission, mutation rate for a

two-segment virus, we varied the per-genome per-transmission mutation rate U

between 0.05 and 0.2 (Figure S6). Overall, these simulations yielded qualitatively

similar results: the antigenic gene segment had a younger TMRCA than the non-

antigenic gene segment. Interestingly, at higher mutation rates, mean TMRCAs for

the non-antigenic gene segment were appreciably smaller and mean TMRCAs for the

antigenic gene segment were slightly smaller. Further, the difference in the mean

TMRCAs for the antigenic and non-antigenic gene segment was smaller at higher

mutations, most likely reflecting a concomitant increase in interference effects

between the gene segments.

We also looked at the sensitivity of the coinfection rate by varying β from 0.0025 to

0.25 per day (Figure S7). When the frequency of the coinfected individuals was set at

1% in the total infected population (i.e. $\beta = 0.0025$ per day) the TMRCA of the two

gene segments was found to be very similar (Figure S7: mean difference in TMRCA

is <0.5 years). While this level of coinfection corresponds well with empirical

estimates (36-38), the difference in TMRCAs between antigenic and non-antigenic

gene segments is not consistent with the observed evolutionary dynamics in Figure 2.

DISCUSSION

We have developed a simple population genetic model to examine the role of non-

antigenic gene segments in the adaptive evolution of seasonal influenza A viruses. In

contrast to previous phylodynamic and predictive models of HA evolution, which

have exclusively focused on HA (10, 12, 14, 15, 44), our approach allows us to

evaluate the importance of selection on non-antigenic gene segments and

intrasubtypic reassortment to the molecular and adaptive evolutionary dynamics of the

virus genome. We find that the limited genetic diversity of non-antigenic gene

segments and differences in TMRCAs between non-antigenic and antigenic gene

segments are principally captured when selection on both antigenic and non-antigenic

gene segments occurs and in the presence of low-level reassortment. Furthermore,

while our results indicate that selection on the non-antigenic gene segment can

slightly influence the evolutionary dynamics of the antigenic gene segment,

reassortment increases viral adaptation in our model primarily by decreasing selective

interference acting upon the non-antigenic gene segment, rather than the antigenic

gene segment.

Given that only two segments are modeled in this study, it would be interesting to see

if these results still hold when additional non-antigenic gene segments are considered.

One prediction is that since linkage effects are expected to increase with additional

gene segments, we are more likely to see the cumulative effect of selection acting on

the non-antigenic segments on the antigenic gene segment. Furthermore, the fitness

variation of each gene segment (and the overall virus) is also likely to increase, thus

enabling selection to be more efficacious. Consequently, in light of this hypothesis,

our finding that non-antigenic gene segment has minimal impact on the antigenic gene

segment is likely to be overly conservative.

Since the coinfection level assumed in our model is ~5%, around 2.5% of the infected

population is expected to carry a first-generation reassortant virus. Interestingly, this

low level of reassortment is consistent with a recently estimated frequency of

reassortment events observed among sampled virus genomes over time, at around

3.35% (25). Further evidence that the intrasubtypic reassortment is restricted at the

between-host level comes from a recent finding that even at the within-host scale the

effective reassortment rate is very limited (45). This indicates that the difference in

the TMRCA across the seasonal influenza A virus genome is likely to arise from a

low-level of reassortment in the virus population. Importantly, this has strong

implications for the adaptive evolution of the virus, since it suggests that selective

interference among gene segments has the potential to influence the fate of beneficial

mutations in the genome.

Although reassortment is notoriously associated with pandemic influenza (46), there

are several historical events in both seasonal influenza A/H3N2 and in seasonal

influenza A/H1N1 where intrasubtypic reassortment has been implicated in antigenic

cluster transitions (16, 22, 47). Furthermore, given that these instances are often

associated with greater disease severity and incidence, akin to pandemic influenza, it

also indicates that intrasubtypic reassortment can facilitate significant improvements

in viral fitness. Consequently, this suggests that reassortment predominantly increases

the rate of virus adaptive evolution by reducing selective interference effects across

the genome.

We did not explicitly consider epistasis in our simulation model. There is evidence

that epistatic interactions both within and between gene segments can drive the

adaptive evolution of seasonal influenza A viruses. For example, T-cell immune

escape mutations in NP have been enabled by stability-mediated epistasis (19, 20) and

functional mismatches between the activities of the HA and the NA are known to

decrease viral fitness considerably (48). However, to effectively model epistasis, a

detailed knowledge about the fitness landscape of the virus genome, which is

currently lacking, is necessary. Elucidating the epistatic interactions in influenza A

viruses should be a focus of future work, since it could help explain the role that

intrasubtypic reassortment plays in contributing to the adaptive evolution of seasonal

influenza (49), and more broadly, it could help understand the epidemic (and even

pandemic) potential of reassortant viruses.

Our findings that selection is likely to act upon both antigenic and non-antigenic gene

segments and that reassortment can influence the rate of virus adaptive evolution have

important implications for predicting future influenza strains. In particular, our study

indicates that viral mutations are subjected to linkage effects within and to a

somewhat lesser extent between gene segments, consistent with the conclusions of

(37). As a consequence, we anticipate better forecasting can be achieved if the virus

genetic background is considered as a whole, and is not just restricted to HA. This

will be largely dependent on obtaining a more comprehensive understanding of the

phenotypic variation in other gene segments, which we recommend should be a

priority for future research.

REFERENCES

1.	WHO. Influenza Fact Sheet. wwwwhoint/mediacentre/factsheets/fs211/en/. 2014.

2.	Hay AJ, Gregory V, Douglas AR, Lin YP. The evolution of human influenza viruses. Philos Trans R Soc Lond B Biol Sci. 2001;356(1416):1861-70.

3.	Wiley DC, Wilson IA, Skehel JJ. Structural Identification of the Antibody-Binding Sites of Hong-Kong Influenza Hemagglutinin and Their Involvement in Antigenic Variation. Nature. 1981;289(5796):373-8.

4.	Wilson IA, Cox NJ. Structural Basis of Immune Recognition of Influenza-Virus Hemagglutinin. Annu Rev Immunol. 1990;8:737-&.

5.	Bush RM, Bender CA, Subbarao K, Cox NJ, Fitch WM. Predicting the evolution of human influenza A. Science. 1999;286(5446):1921-5.

6.	Koel BF, Burke DF, Bestebroer TM, van der Vliet S, Zondag GC, Vervaet G, et al. Substitutions near the receptor binding site determine major antigenic change during influenza virus evolution. Science. 2013;342(6161):976-9.

7.	Ferguson NM, Galvani AP, Bush RM. Ecological and immunological determinants of influenza evolution. Nature. 2003;422(6930):428-33.

8.	Plotkin JB, Dushoff J, Levin SA. Hemagglutinin sequence clusters and the antigenic evolution of influenza A virus. Proc Natl Acad Sci U S A. 2002;99(9):6263-8.

9.	Smith DJ, Lapedes AS, de Jong JC, Bestebroer TM, Rimmelzwaan GF, Osterhaus AD, et al. Mapping the antigenic and genetic evolution of influenza virus. Science. 2004;305(5682):371-6.

10.	Koelle K, Cobey S, Grenfell B, Pascual M. Epochal evolution shapes the phylodynamics of interpandemic influenza A (H3N2) in humans. Science. 2006;314(5807):1898-903.

11.	Bedford T, Rambaut A, Pascual M. Canalization of the evolutionary trajectory of the human influenza virus. BMC Biol. 2012;10:38.

12.	Zinder D, Bedford T, Gupta S, Pascual M. The roles of competition and mutation in shaping antigenic and genetic diversity in influenza. PLoS Pathog. 2013;9(1):e1003104.

13.	Illingworth CJ, Mustonen V. Components of selection in the evolution of the influenza virus: linkage effects beat inherent selection. PLoS Pathog. 2012;8(12):e1003091.

14.	Strelkowa N, Lassig M. Clonal interference in the evolution of influenza. Genetics. 2012;192(2):671-82.

15.	Luksza M, Lassig M. A predictive fitness model for influenza. Nature. 2014;507(7490):57-61.

16.	Koelle K, Rasmussen DA. The effects of a deleterious mutation load on patterns of influenza A/H3N2's antigenic evolution in humans. Elife. 2015;4:e07361.

17.	Kim K, Kim Y. Population genetic processes affecting the mode of selective sweeps and effective population size in influenza virus H3N2. BMC Evol Biol. 2016;16:156.

18.	Memoli MJ, Jagger BW, Dugan VG, Qi L, Jackson JP, Taubenberger JK. Recent human influenza A/H3N2 virus evolution driven by novel selection factors in addition to antigenic drift. J Infect Dis. 2009;200(8):1232-41.

19.	Gong LI, Bloom JD. Epistatically interacting substitutions are enriched during adaptive protein evolution. PLoS Genet. 2014;10(5):e1004328.

20.	Gong LI, Suchard MA, Bloom JD. Stability-mediated epistasis constrains the evolution of an influenza protein. Elife. 2013;2:e00631.

21.   Rambaut A, Pybus OG, Nelson MI, Viboud C, Taubenberger JK, Holmes EC.
The genomic and epidemiological dynamics of human influenza A virus. Nature.
2008;453(7195):615-9.
22.   Holmes EC, Ghedin E, Miller N, Taylor J, Bao Y, St George K, et al. Whole-
genome analysis of human influenza A virus reveals multiple persistent lineages and
reassortment among recent H3N2 viruses. PLoS Biol. 2005;3(9):e300.
23.   Nelson MI, Viboud C, Simonsen L, Bennett RT, Griesemer SB, St George K,
et al. Multiple reassortment events in the evolutionary history of H1N1 influenza A
virus since 1918. PLoS Pathog. 2008;4(2):e1000012.
24.   Westgeest KB, Russell CA, Lin X, Spronken MI, Bestebroer TM, Bahl J, et al.
Genomewide analysis of reassortment and evolution of human influenza A(H3N2)
viruses circulating between 1968 and 2011. J Virol. 2014;88(5):2844-57.
25.   Berry IM, Melendrez MC, Li T, Hawksworth AW, Brice GT, Blair PJ, et al.
Frequency of influenza H3N2 intra-subtype reassortment: attributes and implications
of reassortant spread. Bmc Biology. 2016;14.
26.   Recker M, Pybus OG, Nee S, Gupta S. The generation of influenza outbreaks
by a network of host immune responses against a limited set of antigenic types. Proc
Natl Acad Sci U S A. 2007;104(18):7711-6.
27.   Bhatt S, Holmes EC, Pybus OG. The genomic rate of molecular adaptation of
the human influenza A virus. Mol Biol Evol. 2011;28(9):2443-51.
28.   Drummond AJ, Suchard MA, Xie D, Rambaut A. Bayesian phylogenetics with
BEAUti and the BEAST 1.7. Mol Biol Evol. 2012;29(8):1969-73.
29.   Drummond AJ, Ho SY, Phillips MJ, Rambaut A. Relaxed phylogenetics and
dating with confidence. PLoS Biol. 2006;4(5):e88.
30.   Shapiro B, Rambaut A, Drummond AJ. Choosing appropriate substitution
models for the phylogenetic analysis of protein-coding sequences. Mol Biol Evol.
2006;23(1):7-9.
31.   Gill MS, Lemey P, Faria NR, Rambaut A, Shapiro B, Suchard MA. Improving
Bayesian population dynamics inference: a coalescent-based model for multiple loci.
Mol Biol Evol. 2013;30(3):713-24.
32.   Bedford T. PACT: Posterior Analysis of Coalescent Trees 2015 [Available
from: http://bedford.io/projects/PACT/.
33.   Wickham H. ggplot2: Elegant Graphics for Data Analysis. Use R. 2009:1-212.
34.   Yu GC, Smith DK, Zhu HC, Guan Y, Lam TTY. GGTREE: an R package for
visualization and annotation of phylogenetic trees with their covariates and other
associated data. Methods Ecol Evol. 2017;8(1):28-36.
35.   Carrat F, Vergu E, Ferguson NM, Lemaitre M, Cauchemez S, Leach S, et al.
Time lines of infection and disease in human influenza: a review of volunteer
challenge studies. Am J Epidemiol. 2008;167(7):775-85.
36.   Goka E, Vallely P, Mutton K, Klapper P. Influenza A viruses dual and
multiple infections with other respiratory viruses and risk of hospitalisation and
mortality. Influenza Other Respir Viruses. 2013;7(6):1079-87.
37.   Perez-Garcia F, Vasquez V, de Egea V, Catalan P, Rodriguez-Sanchez B,
Bouza E. Influenza A and B co-infection: a case-control study and review of the
literature. Eur J Clin Microbiol Infect Dis. 2016;35(6):941-6.
38.   Poon LL, Song T, Rosenfeld R, Lin X, Rogers MB, Zhou B, et al. Quantifying
influenza virus diversity and transmission in humans. Nat Genet. 2016;48(2):195-200.
39.   Visher E, Whitefield SE, McCrone JT, Fitzsimmons W, Lauring AS. The
Mutational Robustness of Influenza A Virus. PLoS Pathog. 2016;12(8):e1005856.

40. Gillespie DT. Approximate accelerated stochastic simulation of chemically reacting systems. J Chem Phys. 2001;115(4):1716-33.

41. Russell CA, Jones TC, Barr IG, Cox NJ, Garten RJ, Gregory V, et al. The global circulation of seasonal influenza A (H3N2) viruses. Science. 2008;320(5874):340-6.

42. Bahl J, Nelson MI, Chan KH, Chen R, Vijaykrishna D, Halpin RA, et al. Temporally structured metapopulation dynamics and persistence of influenza A H3N2 virus in humans. Proc Natl Acad Sci U S A. 2011;108(48):19359-64.

43. Lemey P, Rambaut A, Bedford T, Faria N, Bielejec F, Baele G, et al. Unifying viral genetics and human transportation data to predict the global transmission dynamics of human influenza H3N2. PLoS Pathog. 2014;10(2):e1003932.

44. Bedford T, Cobey S, Pascual M. Strength and tempo of selection revealed in viral gene genealogies. BMC Evol Biol. 2011;11:220.

45. Sobel Leonard A, McClain MT, Smith GJ, Wentworth DE, Halpin RA, Lin X, et al. The effective rate of influenza reassortment is limited during human infection. PLoS Pathog. 2017;13(2):e1006203.

46. Morens DM, Taubenberger JK, Fauci AS. The persistent legacy of the 1918 influenza virus. N Engl J Med. 2009;361(3):225-9.

47. Nelson MI, Edelman L, Spiro DJ, Boyne AR, Bera J, Halpin R, et al. Molecular epidemiology of A/H3N2 and A/H1N1 influenza virus during a single epidemic season in the United States. PLoS Pathog. 2008;4(8):e1000133.

48. Neverov AD, Kryazhimskiy S, Plotkin JB, Bazykin GA. Coordinated Evolution of Influenza A Surface Proteins. PLoS Genet. 2015;11(8):e1005404.

49. Neverov AD, Lezhnina KV, Kondrashov AS, Bazykin GA. Intrasubtype reassortments cause adaptive amino acid replacements in H3N2 influenza genes. PLoS Genet. 2014;10(1):e1004037.

A

Coinfected Transmission

0.25   0.25   0.25   0.25

 Singly-infected (I_s)

Coinfected (I_{co})

Parental virus 1

Parental virus 2

B

Before | Process | After

Infection/Recovery

infected population

Coinfection