SOD1^{A4V} aggregation alters ubiquitin homeostasis in a cell model of ALS

Natalie E. Farrawell[1,2], Isabella Lambert-Smith[1,2], Kristen Mitchell[1,2], Jessie
McKenna[3], Luke McAlary[1,2,4], Prajwal Ciryam[5,6,7], Kara L. Vine[1,2], Darren N.
Saunders[3], Justin J. Yerbury[1,2*]

[1]Illawarra Health and Medical Research Institute, Wollongong, NSW, Australia
[2]School of Biological Sciences, Centre of Medicine and Molecular Bioscience,
University of Wollongong, Northfields Ave, Wollongong, NSW, Australia, 2522
[3]School of Medical Sciences, Faculty of Medicine, UNSW Australia
[4]Department of Physics & Astronomy, University of British Columbia, Vancouver,
British Columbia, Canada.
[5]Department of Chemistry, University of Cambridge, Cambridge CB2 1EW, UK
[6]Department of Molecular Biosciences, Rice Institute for Biomedical Research,
Northwestern University, Evanston, IL 60208-3500, USA
[7]Department of Neurology, Columbia University College of Physicians & Surgeons,
New York, NY 10032-3784
15
16
17
18 * Correspondence:
19 Dr Justin John Yerbury
20 Illawarra Health and Medical Research Institute
21 Northfields Ave, Wollongong, NSW, 2522, Australia.
22 Ph: 61-2-42981534; Email: jyerbury@uow.edu.au
23

Keywords: protein aggregation, ubiquitin, SOD1, ALS, supersaturation, proteostasis,
neurodegeneration, ubiquitomics, degron, proteasome.

Abstract

Amyotrophic lateral sclerosis (ALS) is a fatal neurodegenerative disease involving the

selective death of upper and lower motor neurons in the primary motor cortex and

spinal cord. A hallmark of ALS pathology is the accumulation of ubiquitinated

protein inclusions within motor neurons. Previous studies suggest the sequestration of

ubiquitin (Ub) into inclusions reduces the availability of free Ub, which is essential

for cellular function and survival. However, the dynamics of the Ub landscape in ALS

have not yet been described. Here we show that Ub homeostasis is altered in a SOD1

cell model of ALS. Utilising fluorescently tagged Ub, we followed the distribution of

Ub in living cells expressing SOD1 and show that Ub is present at the earliest stages

of SOD1 aggregation. We also report that cells containing aggregates of mutant

SOD1 have greater ubiquitin-proteasome system (UPS) dysfunction as measured by

the accumulation of the fluorescent proteasome reporter tdTomato[CL1]. Furthermore,

SOD1 aggregation is associated with the redistribution of Ub and depletion of the free

Ub pool. Ubiquitomics analysis indicates that mutant SOD1 is associated with a shift

of Ub to a pool of supersaturated proteins including those associated with oxidative

phosphorylation and metabolism, corresponding with altered mitochondrial

morphology and function. Taken together, these results suggest misfolded SOD1

contributes to UPS dysfunction and that Ub homeostasis is an important target for

monitoring pathological changes in ALS.

48
49

Background

Amyotrophic lateral sclerosis (ALS, also known as motor neuron disease, MND) is a progressive neurodegenerative disease leading to paralysis of voluntary muscles due to the death of motor neurons in the brain and spinal cord. The prognosis of ALS is poor, usually leading to death within 2 to 5 years of first symptoms. A fraction of patients also develop clinical or subclinical frontotemporal dementia (FTD) [1]. In most cases of ALS, the cause remains unknown (sporadic ALS; sALS). However, approximately 10% of cases are inherited (familial ALS; fALS). A large proportion (20%) of fALS cases can be attributed to mutations in the gene encoding superoxide dismutase 1 (SOD1) [2]. SOD1 was the first gene discovered to cause fALS and is also the most widely studied. There are now over 20 genes known to cause ALS [2] including a growing list of genes associated with dysregulation of ubiquitin (Ub) signalling. Genetic mutations in VCP, SQSTM1, UBQLN2, OPTN have all been associated with ALS, and are all part of the cell's protein degradation machinery. In addition, recently discovered mutations in TBK1 [3] and CCNF [4] add to this growing list of degradation machinery associated with ALS. The precise role of each of these genes in the homeostasis of Ub is unknown, but Ub sequestration into insoluble inclusions is common to all forms of ALS [5].

Abnormal accumulation of proteins into insoluble aggregates is a hallmark of many neurodegenerative diseases, including Alzheimer's disease, Parkinson's disease Huntington's disease and ALS [6]. In the context of ALS, there is growing evidence that a correlation exists between protein aggregate load and neuronal loss in ALS spinal cord [7-11]. Our previous work showed a correlation between *in vitro*

aggregation propensity and rate of disease progression [12], suggesting that protein

aggregates are intimately linked with motor neuron cell death. Recent work also

indicates that protein misfolding and aggregation may be responsible for disease

progression through a prion like propagation throughout the nervous system [11, 13-

16]. It is unlikely that misfolding alone is responsible for the disease and that post

translational modifications play an important role [17]. One crucial post translational

modification is ubiquitination, necessary for protein degradation. Degradation defects

that lead to inclusion formation are associated with a tendency for cells to be

dysfunctional and undergo apoptosis [18-20].

Inclusions associated with neurodegeneration consist of a variety of proteins including

proteins specific to the disease (e.g. Aβ and tau in Alzheimer's disease [21]), proteins

associated with cellular quality control machinery (e.g. molecular chaperones [22, 23]

and the proteasome [24]) and other unrelated aggregation prone proteins [25, 26].

Based on analysis of human tissue, it has been shown that a large number of proteins

are supersaturated in the cell, with cellular concentrations under wild-type conditions

that exceed their predicted solubility [25, 26]. These supersaturated proteins are

associated with the biochemical pathways underpinning a variety of

neurodegenerative diseases. Most recently, we have shown that proteins co-

aggregating with SOD1, TDP-43, and FUS inclusions are supersaturated [5],

consistent with a collapse of motor neuron protein homeostasis in ALS. Others have

found that the proteins that co-aggregate with c9orf72 dipeptide repeats in cell models

are also supersaturated [27]. The composition of inclusions found in ALS varies

considerably depending on whether the disease is sporadic or familial, and the

genetics of the familial forms.

Ub is a pervasive feature of inclusions in ALS, regardless of underlying genetic

aetiology. Ub is a versatile signalling molecule responsible for controlling an array of

cellular pathways including transcription, translation, vehicle transport and apoptosis

[28]. Ub labels substrate proteins via a highly ordered multi-step enzymatic cascade

with specific differences in the length and topology of poly-ubiquitin chains

determining a range of signalling outcomes, including proteolytic degradation via the

proteasome [29, 30]. Inside cells, Ub exists in a dynamic equilibrium between free Ub

and Ub conjugates and is controlled by the opposing actions of Ub ligases and

deubiquitinating enzymes (DUBS) [31, 32]. It has been proposed that the

sequestration of Ub into insoluble aggregates may deplete the free Ub pool required

by many essential cellular processes [32].

Although mounting genetic and functional evidence suggests an important role for the

UPS in the development of ALS pathology, the distribution and availability of Ub in

ALS models has not yet been described. In the work reported here, we sought to

characterise the Ub landscape in a cell-based SOD1 model of ALS **(Figure 1)**. We

show that Ub homeostasis is disrupted in cells containing SOD1 aggregates by

following the distribution of fluorescently labelled Ub in live cells expressing SOD1.

The aggregation of mutant SOD1 leads to an accumulation of the proteasome reporter

tdTomato^{CL1}, indicative of UPS dysfunction. This dysfunction was further supported

by the redistribution of the Ub pool and decrease in free Ub levels observed in cells

with SOD1 aggregates. Moreover, ubiquitome analysis confirmed misfolded SOD1

was associated with Ub redistribution and subsequent alterations to mitochondrial

morphology. This report highlights that disruption to Ub homeostasis is associated

with aggregation of misfolded proteins and may play an important role in the

pathogenesis of ALS.

Methods

Plasmids. pEGFP-N1 vectors containing human SOD1^{WT} and SOD1^{A4V} were generated as described [33]. GFP-Ub and mRFP-Ub (Addgene plasmids 11928 and 11935, provided by Nico Dantuma [31, 34]) were acquired from Addgene (USA). The mcherry-Ub construct was created by replacing the mRFP sequence in mRFP-Ub with mcherry fluorescent protein. The tdTomato^{CL1} construct was obtained by cloning the CL1 sequence (ACKNWFSSLSHFVIHL) into pcDNA3.1(+)tdTomato.

Cell Culture and Transfection. Neuroblastoma × Spinal cord hybrid NSC-34 cells [35] were maintained in Dulbecco's Modified Eagles Medium/ Ham's Nutrient Mixture F12 (DMEM/F12) supplemented with 10% foetal bovine serum (FBS, Bovogen Biologicals, Australia). Cells were maintained at 37 °C in a humidified incubator with 5% atmospheric CO_2. For confocal microscopy, cells were grown on 13 mm round coverslips in 24-well plates or on 8-well μ-slides (Ibidi, Germany). Cells were grown in 6-well plates for cell lysate and cell sorting experiments. Cells were transfected using Lipofectamine 3000 (Invitrogen, USA) according to manufacturer's instructions with 0.5 μg DNA per well for a 24-well plate, 0.2 μg DNA per well for 8-well μ-slides and 2.5 μg DNA per well for 6-well plates. For co-transfections the amount of DNA was divided equally between constructs.

Measurement of UPS function. To quantify UPS activity, NSC-34 cells were co-transfected with the fluorescent proteasome reporter tdTomato^{CL1} and treated overnight (~18 h) with 0-30μM of the proteasome inhibitor MG132. The level of tdTomato^{CL1} fluorescence was determined using flow cytometry on a Becton Dickinson flow cytometer 48 h post transfection as in [36].

Live cell imaging. Imaging of NSC-34 cells co-transfected with SOD1^{A4V}-GFP and

mRFP-Ub was performed 24 h post transfection on a Leica TCS SP5 confocal

microscope. Cells were imaged every 15 min over 17 h in a CO_2 chamber maintained

at 37˚C with 5% CO_2. For fluorescence recovery after photobleaching (FRAP) and

fluorescence recovery after nucleus photobleaching (FRANP) experiments, analysis

was performed on transfected NSC-34 cells 48 h post transfection using the LAS AF

FRAP Application Wizard on the Leica TCS SP5 confocal Microscope. Images were

acquired using the 63× objective with two line averages and a scan speed of 700 Hz.

Five pre-bleach images were acquired over 7.5 s with the 561 nm laser set at 20%

power. The region of interest (ROI) was then bleached using the 'zoom in ROI'

method over five frames of 1.5 s at 100% laser power. For FRANP analysis, the entire

nucleus was bleached. Fluorescence recovery or loss was monitored for 120 s with the

laser power set back at 20%.

Frequency distribution analysis. GFP fluorescence in NSC-34 cells expressing

soluble and insoluble (aggregated) SOD1^{A4V}-GFP was quantified from confocal

images taken 48 h post transfection using ImageJ 1.48v [37]. Frequency distribution

analysis was subsequently performed in GraphPad Prism version 5.00 for Windows

(GraphPad software, USA).

Immunofluorescence. NSC-34 cells grown on coverslips were transfected with GFP-

tagged mutant SOD1 and fixed for 20 min at room temperature (RT) with 4%

paraformaldehyde (PFA) (Merck Millipore, USA) in phosphate buffered saline (PBS)

48 h post transfection. Cells were permeabilized in 1% tritonX-100 (TX-100) in PBS

for 30 min on ice before blocking for 1 h at RT with 5% FBS, 1% Bovine serum

albumin (BSA) 0.3% TX-100 in PBS. Cells were incubated with rabbit primary

antibodies against K48 or K63 Ub chain linkages (05-1307/05-1308, Merck

Millipore) overnight at 4 °C followed by Alexa Fluor 647-conjugated anti-rabbit IgG

secondary antibody for 5 h at RT. All antibodies were diluted in 1% BSA, 0.1% TX-

100 in PBS and cells were washed with PBS between each incubation step.

Spinal cord sections from the SOD1^{G93A} mouse were also stained for SOD1 and K48

or K63 Ub chain linkages. A pap pen (Daido Sangyo, Tokyo, Japan) was used to

separate tissue sections mounted onto the same slide. Sections were then fixed with

4% PFA for 15 min at RT before permeabilization at RT for 10 min with 0.1% TX-

100 in PBS containing 2% (v/v) normal horse serum (NHS). Sections were then

blocked for 20 min at RT with 20% NHS and 2% BSA in PBS followed by staining

overnight at 4 °C in a humidified chamber with sheep anti-superoxide dismustase-1

(ab8866, Abcam, UK), mouse neuron-specific beta III tubulin antibody (ab78078,

Abcam) and rabbit anti-ubiquitin K48 or K63 (Merck, Millipore). The following day,

sections were incubated with Alexa Fluor-conjugated secondary antibodies reactive to

sheep, mouse and rabbit IgG for 1 h at RT. Following staining, coverslips were

mounted onto slides using ProLong Diamond Antifade Mountant (Molecular probes,

USA). All imaging experiments were analysed on a Leica TCS SP5 confocal

microscope.

Cell lysis. NSC-34 cells grown in 6-well plates and transfected with SOD1-GFP or

mRFP-Ub were harvested 48 h post transfection with trypsin/EDTA (Gibco, USA).

Cells were washed with PBS before being resuspended in RIPA buffer (50 mM

TrisHCl pH 7.4, 1% (w/v) sodium deoxycholate, 150 mM NaCl, 1 mM EDTA, 1%

TX-100, 0.1% SDS, 10mM NEM, 1 mM sodium orthovanadate, Halt™ Protease

Inhibitor Cocktail (Thermo Scientific, USA)). Protein concentration was determined

by BCA assay.

Western blot. Cell lysates with a total protein concentration of 100 μg were reduced

with β-mercaptoethanol and heated for 10 min at 70 °C before being analysed on a

Mini-PROTEAN TGX Stain-Free Gel (Bio-Rad, USA) for 2 h at 100 V. Proteins

were transferred onto a nitrocellulose membrane (Pall Corporation, USA) using the

standard protocol on the Bio-Rad Trans-Blot Turbo Transfer System. The membrane

was blocked in 5% skim milk powder in Tris buffered saline, 0.2% (v/v) Tween-20

(TBST) for 1 h at RT before probing for Ub with rabbit anti-ubiquitin antibody

(ab137025, Abcam) overnight at 4 °C. The following day the membrane was washed

three times with TBST over 30 min before incubating with horseradish peroxidise-

conjugated anti-rabbit antibody (1706515, Bio-Rad) for 1 h at RT. The membrane was

visualised with chemiluminescent substrate (Thermo Scientific) on the Amersham

Imager 6600RGB.

Cell sorting. NSC-34 cells transfected with SOD1-GFP were harvested 48 h post-

transfection with trypsin-EDTA (Gibco) and washed with PBS before being

resuspended in resuspension buffer (PBS containing 25 mM HEPES, 1 mM EDTA

and 1% FBS) and filtered through a 40 μm nylon mesh strainer (Greiner Bio-One,

Austria). Enrichment of GFP positive cells was performed on a Bio-Rad S3e cell

sorter. Cells were sorted based on GFP fluorescence (excitation 488 nm, emission

525/30nm) and collected into 5 mL FACS tubes containing DMEM/F12

supplemented with 10% FBS and 2 × penicillin/streptomycin (Gibco). Following

enrichment, cells were lysed in RIPA buffer (as above) for ubiquitomics analysis.

Ubiquitomics. Proteomics to identify ubiquitin-modified proteome (ubiquitome) in

NSC-34 cells was performed as described in [38]. Cells were lysed in lysis buffer (50

mM Tris-HCl (pH 7.5), 150 mM NaCl, 0.5% NP-40, 1 mM DTT, 1× EDTA-free

protease inhibitor cocktail, 10 mM N-ethylmaleimide, 1 mM sodium orthovanadate).

Protein concentration was determined using a BCA assay and 300 μg of total protein

was used per sample to immunopurify mono- and poly-ubiquitinated proteins using

30 µL of VIVAbind Ub affinity matrix (Ubiquitin Kit, VIVA Bioscience). Samples

were incubated for 2 h at 4 °C with end-to-end mixing then matrix was collected by

centrifugation (1 min; 4 °C; 1000 × g) and washed six times in detergent free wash

buffer (500 µL). After washing, beads were digested for 30 min at 27 °C, then

reduced with 1 mM DTT and left to digest overnight at room temperature with

sequencing-grade trypsin (5 μg/mL, Promega), as described previously [39]. Samples

were alkylated with 5 mg/mL iodoacetamide and protease digestion terminated with

trifluoroacetic acid. Trypsinized eluents were collected after brief centrifugation then

purified and desalted using self-packed tips with 6 layers of C18 Empore disks

(Pacific Laboratory Products), then dried in a SpeedVac. Samples were then

resuspended in 12 µL 5% formic acid, 2% acetonitrile and stored at -80 °C.

For nLC-MS/MS analysis, 5 uL of each of the peptide samples were loaded and

separated along a C18 column (400 mm, 75 μm ID, 3 μm silica beads) and introduced

by nanoelectrospray into an LTQ Orbitrap Velos Pro coupled to an Easy-nLC HPLC

(Thermo Fisher). Tandem mass spectrometry data was collected for the top 10 most

abundant ions per scan over a 140-minute time gradient. The order of data collection

was randomized to interchange between biological conditions with BSA run between

each sample to minimize temporal bias. MS/MS raw files were analyzed using the

Andromeda search engine integrated into MaxQuant (v1.2.7.4) [40] against the

Uniprot Mouse database. A false discovery rate of 1% was tolerated for protein,

peptide, and sites, and one missed cleavage was allowed. MaxQuant output data were

filtered to remove contaminants, reverse hits, proteins only identified by site and

proteins with < 2 unique peptides [41]. An individual protein was defined as present

under a particular condition if it was detected in a minimum of two replicates.

Protein-protein interactions and KEGG pathway analysis among the resulting protein

list was analyzed using STRING (v10) [42] (with a confidence score of 0.700) and

Cytoscape (v3.1.1) [43].

Analysis of Mitochondrial Morphology and Function. NSC-34 cells transiently

transfected with SOD1-GFP were incubated with 100nM Mitotracker Deep Red FM

(Molecular probes) or Mitotracker CMXRos (Molecular probes) for 30 min at 37 °C

prior to analysis by confocal microscopy or flow cytometry 48 h post transfection.

Mitochondrial morphology was examined using a mitochondrial morphology macro

[44] on ImageJ 1.48v. Mitotracker Red and Mitotracker CMXRos accumulation in

cells expressing high levels of SOD1-GFP was also assessed by flow cytometry

(Becton Dickinson).

Results

SOD1 aggregates are ubiquitylated early

We previously showed that all cellular SOD1 aggregates contained Ub at the time

points tested [45]. Here we used real-time imaging using a Ub fusion protein to follow

inclusion formation in a single cell from its genesis until it had taken up a large

proportion of the cytoplasm. Previous work has demonstrated that the behaviour of

GFP-Ub fusions in cells is indistinguishable from endogenous Ub and provides a

robust fluorescent indicator of Ub distribution [31]. We created a mCherry-Ub fusion

protein that behaved in a similar manner to the GFP fusion protein (supplementary

Figure S1). Ub was observed in foci at the earliest detectable stages of SOD1

aggregation (**Figure 2A**, see insert). Ub was continuously added to inclusions

throughout their formation, as was SOD1 (**Figure 2A**). To ensure that the Ub

accumulation in SOD1 inclusions was not an artefact of Ub over expression, we next

probed for endogenous Ub using antibodies. SOD1^{A4V} expression causes inclusions in

approximately 15% of NSC-34 cells (**Figure S2)** and in those cells that contain

inclusions we observed a high degree of overlap between SOD1 aggregates and Ub

when using both K48 and K63 chain specific antibodies (**Figure 2B**). We also found

that SOD1 aggregates contained K48 chains in SOD1^{G93A} mouse spinal motor

neurons (**Figure 2C**).

Cells with SOD1 aggregates have increased UPS dysfunction

We previously showed that a UPS reporter containing a CL1 degron peptide for rapid

degradation accumulates to significantly higher amounts in ALS patient fibroblasts

compared to controls, suggesting an overwhelmed UPS [36]. Work from others

suggests that SOD1 aggregates are toxic because they interfere with the quality

control function of the juxtanuclear quality control compartment (JUNQ) [20]. To

examine the relationship between SOD1 aggregation and UPS dysfunction we used

transiently transfected NSC-34 cells that contained SOD1^{A4V}-GFP and tdTomato^{CL1}.

We initially used flow cytometry to examine the effect of proteasome inhibition on

reporter accumulation. While there was a significant difference in reporter signal

between SOD1 and SOD1^{A4V} expressing cells, both cell lines showed a similar dose-

dependent increase in reporter signal with MG132 treatment (**Figure 3A**). This

suggests that, while SOD1^{A4V} expression causes significant UPS disruption, there is

no specific vulnerability to proteasome inhibition of cells expressing mutant SOD1

compared to wild-type expressing cells.

However, this analysis examined the entire population of SOD1^{WT} or SOD1^{A4V}

expressing cells and previous work using a similar CL1-containing fluorescent

reporter suggests that the aggregation of Huntingtin and CFTR cause UPS

dysfunction [46]. We hypothesized that inclusions formed by mutant SOD1 might

influence UPS activity in a similar fashion, a phenomenon that may not be observed

when analysing the entire population of cells, given that only 15% of NSC-34 cells

expressing SOD1^{A4V} produce inclusions (**Figure S2**). In order to identify cells with

SOD1 inclusions in our flow cytometry data we tested the relationship between

aggregation and total cellular fluorescence (**Figure 3B**). Cells were transiently

transfected with SOD1^{A4V}-GFP and imaged 48 h post transfection using confocal

microscopy. The total cellular fluorescence for individual cells was plotted for both

cells with and without inclusions. The fluorescence of both populations was normally

distributed (**Figure 3B**). The mean fluorescence of cells containing aggregates was

significantly greater than that of cells without aggregates (**Figure 3C**), presumably

due to continued accumulation of SOD1 into inclusions. However, there was

significant overlap between the two populations so we could not distinguish

populations entirely. We decided to analyze the most highly fluorescent cells (top

10%) across the entire population. In this subset of cells, approximately 80%

contained SOD1^{A4V} inclusions (**Figure 3D**). We reasoned that analyzing the top 10%

fluorescent cells would sufficiently enrich for cells containing inclusions. Using flow

cytometry, we observed a 4× increase in reporter fluorescence (P< 0.001) - indicating

UPS dysfunction - in the top 10% of SOD1^{A4V}–GFP expressing cells compared to

control cells in the same fluorescence range. That is, UPS dysfunction in SOD1^{A4V}

cells is largely restricted to cells containing aggregates. By comparison, only a small

increase (1.2×) in reporter fluorescence was observed in the entire population of

SOD1^{A4V}–GFP expressing cells compared with the entire SOD1^{WT}-GFP expressing

population (P< 0.01) (**Figure 3E**). This suggests that SOD1 aggregation is associated

with compromised UPS function.

Ubiquitin pools are disturbed in cells containing SOD1 aggregates

Ub exists in a dynamic equilibrium in the cell, partitioning into four major pools; i)

immobile in the nucleus, ii) immobile in the cytoplasm, iii) soluble polyUb chains and

iv) a small fraction as free monomeric Ub [31]. Mobile Ub is a combination of free

monomeric Ub, free Ub chains and Ub attached to diffusible proteins (**Figure 1**).

Immobile Ub is primarily bound to histones in the nucleus and bound to organelles

and cytoskeleton in the cytoplasm (**Figure 1**). To determine if the observed UPS

dysfunction in cells with mutant SOD1 aggregates was associated with altered Ub

homeostasis we examined the relative distribution of Ub into different cellular pools

following expression of wild-type or mutant SOD1 genes by FRAP (**Figure 4A-E**).

The diffusion rate back into the bleached area provides information on the mobility of

the Ub species present in solution and the amount of immobile fluorophore in the

compartment of interest (i.e. bound to membrane vesicles or cytoskeleton in the

cytosol, and bound to histones in the nucleus). We selected ROIs in both the nucleus

and cytoplasm (**Figure 4B**) in cells co-expressing mCherry-Ub and either GFP,

SOD1^{WT}-GFP or SOD^{A4V}-GFP. In the case of SOD^{A4V}-GFP expressing cells, we

performed the analysis on two subpopulations, with and without aggregates. As

previously reported, we find that Ub fluorescence recovery after bleaching (indicating

Ub mobility) is higher in the cytoplasm than in the nucleus, likely due to a large

proportion of nuclear Ub being attached to histones. Patterns of recovery appeared

similar in all treatments, with the exception of a lower cytoplasmic recovery in the

cells containing SOD^{A4V}-GFP aggregates (**Figure 4A**). After calculating the mean

half-life (T$_{1/2}$) of recovery in each case, we find no significant differences between

any of the cell populations (**Figure 4D**). This suggests that within the mobile

population of Ub, the presence of SOD^{A4V}-GFP aggregates does not significantly alter

the kinetics of Ub diffusion.

This observation prompted us to investigate whether SOD^{A4V}-GFP aggregates alter

the amount of mobile Ub available in the cell. We find that on average, 51% of the

cellular Ub signal is detected in aggregates, indicating that the presence of insoluble

aggregates alters the distribution of the Ub pool (**Figure 4C**). The significance of this

aggregate-associated fraction of Ub is clear in the context of our analysis of the

relative distribution of mobile Ub pools. Based on our FRAP analysis, which does not

consider the aggregate-associated Ub, there is no significant difference in the mobility

of Ub between SOD^{A4V}-GFP cells and controls (**Figure 4E**). However, if we correct

for the 51% of Ub we know to be immobile in cells with inclusions, we find that the

actual mobile Ub pool in SOD^{A4V}-GFP cells is significantly smaller than in controls

(**Figure 4E**).

Free monomeric ubiquitin is lowered in cells with SOD1 aggregates

To test whether the depletion of mobile Ub in cells containing SOD^{A4V}-GFP

aggregates may diminish free monomeric Ub, we next examined the relative amounts

of monomeric Ub in cells expressing GFP, SOD1^{WT}-GFP, or SOD^{A4V}-GFP for 48 h,

by Western blot (**Figure 5A**). We did not observe any significant differences using

this method, likely due to the fact that this analysis represents a total measure of all

cells in the culture, including non-transfected cells. We therefore turned to

fluorescence recovery after nucleus photobleaching (FRANP) analysis to monitor free

Ub in single cells. Monomeric mCherry-Ub will diffuse through the nuclear pore (<

60 kDa), while mCherry-Ub incorporated into Ub chains will not [31]. Hence, relative

free monomeric Ub can be measured by monitoring diffusion through the nuclear pore

following fluorescence bleaching solely in the nucleus or cytosol [31]. We transfected

cells with GFP, SOD1^{WT}-GFP, or SOD^{A4V}-GFP (**Figure 5B**) for 48 h, then bleached

the entire nucleus and quantified the amount of relative monomeric Ub diffusing

through the nuclear pore back into the nucleus (**Figure 5C**). We observed a

significant drop in free monomeric Ub in cells containing SOD^{A4V}-GFP aggregates,

but no difference in the level of monomeric Ub between cells expressing SOD1^{WT} and

SOD^{A4V} in the absence of inclusions (**Figure 5D**).

SOD^{A4V}-GFP induced changes in the ubiquitome of NSC34 cells

Free Ub exists in complex equilibrium with multiple conjugated forms and ALS mutations may induce redistribution of Ub through altered activity in various cellular pathways. To investigate changes in the Ub-modified proteome (ubiquitome) of cells expressing SOD^{WT} compared with SOD^{A4V}-GFP, we performed proteomics following enrichment of cell lysates for ubiquitylated proteins (**Figure 6A**). We identified 316 ubiquitylated proteins common to cells expressing either wild-type or mutant SOD1. 55 proteins were uniquely present in the ubiquitome of cells expressing mutant SOD1 and 12 unique proteins were identified in the ubiquitome of cells expressing wild-type SOD1 (**Figure 6B, and Table 1**). Network analysis of known protein-protein interactions and ontology in the set of ubiquitylated proteins unique to cell expressing mutant SOD1 using the String database [42] identified a number of enriched pathways, including metabolic pathways, ubiquitin-proteasome system (UPS), ribosome and mRNA processing and transport (**Figure 6C**). Expression of a chronic misfolded protein has previously been observed to increase proteome misfolding and we have shown that ALS aggregates are composed of supersaturated proteins - that is, proteins which have expression levels higher than one might predict given their solubility [5].

We compared the supersaturation scores for both the unfolded and native states of ubiquitylated proteins found uniquely in SOD^{A4V}-GFP expressing cells to the whole proteome. The median supersaturation score of these 55 ubiquitylated proteins is 10 × higher than the whole proteome in the unfolded state (σ_u) and 25 x higher than the whole proteome in the native state (σ_f) (**Figure 6D**). These data are consistent with proteome instability and resulting redeployment of Ub driving altered Ub distribution

and subsequent impairment of Ub homeostasis upon expression of a chronically

misfolded protein.

Cells with SOD1 aggregates have altered mitochondrial morphology and

function

Pathway analysis of our ubiquitome data (above) indicated specific enrichment of

mitochondrial/metabolic proteins in the ubiquitome of SOD1^{A4V}-expressing cells

(**Table 1**), suggesting that Ub redistribution/homeostasis might be associated with

compromised mitochondrial function in these cells. Indeed, previous work has shown

that mutations in SOD1 lead to mitochondrial dysfunction [47-49]. In order to assess

mitochondrial function in SOD1^{A4V}-expressing cells, we first examined mitochondrial

morphology using Mitotracker and confocal microscopy (**Figure 7A-C**). We found

that SOD1^{A4V}-expressing cells contained a significantly higher number of

mitochondria per cell compared to controls (**Figure 7B**). In addition, we these

mitochondria are significantly more circular than those in control cells (**Figure 7C**).

We quantified mitotracker fluorescence in cells using flow cytometry, as above

focusing on the top 10% most fluorescent cells (based on SOD-GFP expression) to

enrich for cells with aggregates (i.e. when expressing SOD1^{A4V}). This subset of

SOD1^{A4V}-expressing cells has higher mitotracker fluorescence compared to SOD1^{WT}

expressing cells exhibiting the same level of fluorescence - consistent with

accumulation of mitochondria with SOD1^{A4V} expression (**Figure 7D**). In the same

subset of aggregate-containing cells we observed increased fluorescence of the

functional reporter CMXRos, suggesting alterations in mitochondrial membrane

potential (**Figure 7E**).

Discussion

A unifying feature of neurodegenerative diseases such as ALS is the presence of Ub within insoluble protein aggregates [10, 50, 51]. Beyond labelling substrates for degradation via the proteasome, Ub is an important regulator of cellular processes such as transcription, translation, endocytosis and DNA repair. The sequestration of Ub into inclusions may therefore reduce the availability of free Ub essential for these processes, compromising cellular function and survival. To gain insight into the regulation of Ub homeostasis in ALS, we followed the dynamic distribution of Ub at a single cell level in a well-established SOD1 cell model of ALS. Our results confirm that the expression of mutant SOD1 leads to UPS dysfunction and corresponding disruption of Ub homeostasis, suggesting these processes plays a key role in the development of ALS pathology.

In cell models, Ub regulates the concentrations of TDP-43 and SOD1 [52, 53], and proteasome inhibition can trigger their abnormal accumulation. While our previous work established that SOD1 aggregates contain Ub [45], results here confirm that SOD1 co-aggregates with Ub, and that Ub is present at the earliest stage of aggregation. Furthermore, SOD1 aggregates were found to contain both K48 and K63 polyubiquitin chains, signalling degradation via the proteasome and autophagy pathways, respectively. Interestingly, both K48 and K63-linked ubiquitylation has been associated with the formation of inclusions [54], with previous studies showing that K63 polyubiquitination directs misfolded SOD1 to the ubiquitin-aggresome route when the UPS is inhibited [55]. These data are consistent with the need for tightly regulated Ub homeostasis in neurons, and suggest any perturbation in this homeostasis could cause Ub depletion and subsequent toxicity. It is likely that protein

aggregation causes Ub depletion, which could then drive further aggregation in a

positive feedback loop (**Figure 8**).

In the nervous system, the UPS contributes to the regulation of many aspects of

synaptic function, such as neuronal growth and development, neuronal excitability,

neurotransmission, long-term potentiation (LTP) and synapse formation and

elimination [56, 57]. UPS dysfunction is therefore central to neuronal health and

neurodegenerative disease [6]. Impairment of the UPS has been implicated strongly in

the pathogenesis of ALS [52, 58, 59]. Here, we reveal that the aggregation of SOD1

compromises UPS function, decreasing cellular capacity to degrade the proteasome

reporter tdTomato^{CL1}. Our findings are consistent with the previous demonstration of

UPS dysfunction in ALS, with accumulation of the fluorescent UPS reporter Ub^{G76V}-

GFP observed in the spinal cord and cranial motor neurons of SOD1^{G93A} mice [59].

Moreover, a motor neuron specific knockdown of proteasome subunit Rpt3 in the

absence of ALS genetic background results in an ALS phenotype in mice - including

locomotor dysfunction, progressive motor neuron loss and mislocalization of ALS

markers TDP-43, FUS, ubiquilin 2, and optineurin [58]. These data support a model

where a reduction in UPS capacity is sufficient to drive ALS pathology in mice. Not

even overexpression of human mutant TDP-43 gives such an accurate reproduction of

a human ALS-like phenotype in mice. In ALS patient spinal motor neurons, 85%

(38/40) of cytoplasmic TDP-43 foci or inclusions were positive for Ub. In fact, almost

all cellular Ub in these neurons was sequestered within large, skein-like inclusions

[45]. Interestingly, Ub has been found to accumulate in inclusions without the

aggregation of TDP-43 in sporadic ALS [7], suggesting that aggregation of proteins

such as TDP-43, FUS and SOD1 may not be necessary for Ub depletion-induced

toxicity. In human skin fibroblasts it has been shown that the UPS reporter GFP^{CL1}

accumulates significantly more in ALS patient fibroblasts compared to controls,

suggesting a compromised UPS [36]. This raises the question of whether there are

alternate means to control protein concentration when protein aggregates disrupt

degradation mechanisms. It was recently proposed that widespread transcriptional

repression may serve this purpose in Alzheimer's disease [60].

The work presented here suggests that the aggregation of misfolded SOD1 alters Ub

homeostasis and subsequently depletes the free Ub pool in cells. Cellular Ub exists in

a complex equilibrium between free and conjugated Ub [31]. Neurons are vulnerable

to free Ub deficiency, which if prolonged may lead to cell death [61, 62]. Many

factors can influence Ub homeostasis. For example, proteasome inhibition depletes

free Ub to as low as 5% of basal levels in less than 2 h [63, 64]. Inhibition of

translation also depletes free Ub through reduced production, while toxicity can be

rescued by overexpression of Ub [65]. Accumulation of ubiquitylated proteins in

inclusions is an important potential mechanism for depletion of free Ub, and in this

context free Ub levels can be partially restored by overexpression or removal of Ub

from the aggregated protein through ubiquitin-specific proteases. It was recently

shown that while the ataxia associated mutation in Usp14 causes reduction in free Ub

and neuromuscular junction dysfunction in mice, overexpression of Ub restored free

Ub levels in motor neurons and improved NMJ structure [66]. Additionally, free Ub

could be increased in cells containing huntingtin aggregates by overexpression of the

de-ubiquitylation enzyme USP14 [67]. Not only did this protect cells from aggregate-

induced toxicity, it also reduced ER stress, which is thought to precede inclusion

formation in ALS models [18]. These data further support the concept that modulation

of cellular Ub pools is an important factor in the pathogenesis of neurodegenerative

disease.

The pathways responsible for modulating Ub homeostasis in ALS are not yet well

understood. Our ubiquitomics analysis of cells expressing SOD1 reveals that the

unique ubiquitome of cells expressing mutant SOD1 is enriched for proteins prone to

aggregation (i.e. supersaturated). This is consistent with findings that proteins

associated with neurodegeneration are supersaturated [5, 25, 26]. Furthermore, a

substantial proportion of the ubiquitylated proteins identified are from pathways

known to be dysfunctional in ALS, including RNA processing, metabolic and

mitochondrial pathways. This is not surprising given that both RNA binding proteins

and oxidative phosphorylation and metabolic pathways are also prone to aggregation

in neurodegeneration [26]. These results suggest that mutations in SOD1 may lead to

mitochondrial dysfunction by disrupting Ub homeostasis. Interestingly, we find that

cells expressing mutant SOD1 had altered mitochondrial morphology and function.

Disturbances to mitochondrial morphology have been previously reported in human

motor neurons carrying the SOD1^{A4V} mutation [68] and recent studies have revealed

that mitochondrial impairment occurs soon after proteasome inhibition [69]. In fact,

continued proteasome dysfunction in mouse brain cortical neurons inhibited the

degradation of ubiquitylated mitochondrial proteins and led to the accumulation of

dysfunctional mitochondria [70]. Misfolded proteins such as mutant SOD1 have also

been shown to interact with mitochondrial proteins and translocate into the

mitochondrial matrix, where they accumulate and induce mitochondrial dysfunction

[71-73]. However, this cytotoxicity can be attenuated through the ubiquitylation of

misfolded but non-aggregated SOD1 - which promotes its degradation via the UPS

[74]. Collectively, these studies demonstrate an integral functional relationship

between impairment of the UPS and mitochondrial dysfunction, and that modulation

of cellular Ub pools may rescue mitochondrial dysfunction caused by the

accumulation of SOD1.

Conclusions

In conclusion, we observe that the aggregation of mutant SOD1 leads to altered UPS

activity and redistribution of Ub, causing disrupted Ub homeostasis. Given that Ub

controls many essential cellular pathways that are also dysfunctional in ALS -

including transcription, translation, vesicle transport, mitochondrial function and

apoptosis - these findings suggest that Ub homeostasis is a central feature of ALS

pathogenesis. Further understanding of the contribution of Ub homeostasis to ALS

pathology may be imperative to understanding the molecular pathways underpinning

neurodegenerative disease more broadly.

564
565
566

List of abbreviations

List of abbreviations
Amyotrophic lateral sclerosis (ALS)
Bovine serum albumin (BSA)
Deubiquitinating enzymes (DUBS)
Familial ALS (fALS)
Fluorescence loss in photobleaching (FLIP)
Fluorescence recovery after photobleaching (FRAP)
Frontotemporal dementia (FTD)
Juxtanuclear quality control compartment (JUNQ)
Long-term potentiation (LTP)
Motor neuron disease (MND)
Neuromuscular junction (NMJ)
Paraformaldehyde (PFA)
Phosphate buffered saline (PBS)
Region of interest (ROI)
Room temperature (RT)
Sporadic ALS (sALS)
Superoxide dismutase 1 (SOD1)
TritonX-100 (TX-100)
Ubiquitin (Ub)
Ubiquitin-proteasome system (UPS)

593   **References**
594
[1] M.R. Turner, O. Hardiman, M. Benatar, B.R. Brooks, A. Chio, M. de Carvalho,
P.G. Ince, C. Lin, R.G. Miller, H. Mitsumoto, G. Nicholson, J. Ravits, P.J. Shaw, M.
Swash, K. Talbot, B.J. Traynor, L.H. Van den Berg, J.H. Veldink, S. Vucic, M.C.
Kiernan, Controversies and priorities in amyotrophic lateral sclerosis, Lancet
Neurol 12(3) (2013) 310-22.
[2] S. Chen, P. Sayana, X. Zhang, W. Le, Genetics of amyotrophic lateral sclerosis:
an update, Mol Neurodegener 8 (2013) 28.
[3] E.T. Cirulli, B.N. Lasseigne, S. Petrovski, P.C. Sapp, P.A. Dion, C.S. Leblond, J.
Couthouis, Y.F. Lu, Q. Wang, B.J. Krueger, Z. Ren, J. Keebler, Y. Han, S.E. Levy, B.E.
Boone, J.R. Wimbish, L.L. Waite, A.L. Jones, J.P. Carulli, A.G. Day-Williams, J.F.
Staropoli, W.W. Xin, A. Chesi, A.R. Raphael, D. McKenna-Yasek, J. Cady, J.M.
Vianney de Jong, K.P. Kenna, B.N. Smith, S. Topp, J. Miller, A. Gkazi, F.S.
Consortium, A. Al-Chalabi, L.H. van den Berg, J. Veldink, V. Silani, N. Ticozzi, C.E.
Shaw, R.H. Baloh, S. Appel, E. Simpson, C. Lagier-Tourenne, S.M. Pulst, S. Gibson,
J.Q. Trojanowski, L. Elman, L. McCluskey, M. Grossman, N.A. Shneider, W.K.
Chung, J.M. Ravits, J.D. Glass, K.B. Sims, V.M. Van Deerlin, T. Maniatis, S.D. Hayes,
A. Ordureau, S. Swarup, J. Landers, F. Baas, A.S. Allen, R.S. Bedlack, J.W. Harper,
A.D. Gitler, G.A. Rouleau, R. Brown, M.B. Harms, G.M. Cooper, T. Harris, R.M.
Myers, D.B. Goldstein, Exome sequencing in amyotrophic lateral sclerosis
identifies risk genes and pathways, Science (2015).
[4] K.L. Williams, S. Topp, S. Yang, B. Smith, J.A. Fifita, S.T. Warraich, K.Y. Zhang, N.
Farrawell, C. Vance, X. Hu, A. Chesi, C.S. Leblond, A. Lee, S.L. Rayner, V.
Sundaramoorthy, C. Dobson-Stone, M.P. Molloy, M. van Blitterswijk, D.W.
Dickson, R.C. Petersen, N.R. Graff-Radford, B.F. Boeve, M.E. Murray, C. Pottier, E.
Don, C. Winnick, E.P. McCann, A. Hogan, H. Daoud, A. Levert, P.A. Dion, J. Mitsui, H.
Ishiura, Y. Takahashi, J. Goto, J. Kost, C. Gellera, A.S. Gkazi, J. Miller, J. Stockton,
W.S. Brooks, K. Boundy, M. Polak, J.L. Munoz-Blanco, J. Esteban-Perez, A. Rabano,
O. Hardiman, K.E. Morrison, N. Ticozzi, V. Silani, J. de Belleroche, J.D. Glass, J.B.
Kwok, G.J. Guillemin, R.S. Chung, S. Tsuji, R.H. Brown, Jr., A. Garcia-Redondo, R.
Rademakers, J.E. Landers, A.D. Gitler, G.A. Rouleau, N.J. Cole, J.J. Yerbury, J.D.
Atkin, C.E. Shaw, G.A. Nicholson, I.P. Blair, CCNF mutations in amyotrophic lateral
sclerosis and frontotemporal dementia, Nat Commun 7 (2016) 11253.
[5] P. Ciryam, I.A. Lambert-Smith, D.M. Bean, R. Freer, F. Cid, G.G. Tartaglia, D.N.
Saunders, M.R. Wilson, S.G. Oliver, R.I. Morimoto, C.M. Dobson, M. Vendruscolo, G.
Favrin, J.J. Yerbury, Spinal motor neuron protein supersaturation patterns are
associated with inclusion body formation in ALS, Proc Natl Acad Sci U S A
(2017).
[6] J.J. Yerbury, L. Ooi, A. Dillin, D.N. Saunders, D.M. Hatters, P.M. Beart, N.R.
Cashman, M.R. Wilson, H. Ecroyd, Walking the tightrope: proteostasis and
neurodegenerative disease, J Neurochem 137(4) (2016) 489-505.
[7] M.T. Giordana, M. Piccinini, S. Grifoni, G. De Marco, M. Vercellino, M.
Magistrello, A. Pellerino, B. Buccinna, E. Lupino, M.T. Rinaudo, TDP-43
redistribution is an early event in sporadic amyotrophic lateral sclerosis, Brain
Pathol 20(2) (2010) 351-60.
[8] J. Brettschneider, K. Arai, K. Del Tredici, J.B. Toledo, J.L. Robinson, E.B. Lee, S.
Kuwabara, K. Shibuya, D.J. Irwin, L. Fang, V.M. Van Deerlin, L. Elman, L.
McCluskey, A.C. Ludolph, V.M. Lee, H. Braak, J.Q. Trojanowski, TDP-43 pathology

and neuronal loss in amyotrophic lateral sclerosis spinal cord, Acta Neuropathol
128(3) (2014) 423-37.
[9] N. Ticozzi, A. Ratti, V. Silani, Protein aggregation and defective RNA
metabolism as mechanisms for motor neuron damage, CNS and Neurological
Disorders - Drug Targets 9(3) (2010) 285-296.
[10] P.N. Leigh, H. Whitwell, O. Garofalo, J. Buller, M. Swash, J.E. Martin, J.M. Gallo,
R.O. Weller, B.H. Anderton, Ubiquitin-immunoreactive intraneuronal inclusions
in amyotrophic lateral sclerosis. Morphology, distribution, and specificity, Brain
114(2) (1991) 775-788.
[11] M.J. Strong, S. Kesavapany, H.C. Pant, The pathobiology of amyotrophic
lateral sclerosis: A proteinopathy?, Journal of Neuropathology and Experimental
Neurology 64(8) (2005) 649-664.
[12] L. McAlary, J.A. Aquilina, J.J. Yerbury, Susceptibility of Mutant SOD1 to Form
a Destabilized Monomer Predicts Cellular Aggregation and Toxicity but Not In
vitro Aggregation Propensity, Front Neurosci 10 (2016) 499.
[13] R. Zeineddine, J.F. Pundavela, L. Corcoran, E.M. Stewart, D. Do-Ha, M. Bax, G.
Guillemin, K.L. Vine, D.M. Hatters, H. Ecroyd, C.M. Dobson, B.J. Turner, L. Ooi, M.R.
Wilson, N.R. Cashman, J.J. Yerbury, SOD1 protein aggregates stimulate
macropinocytosis in neurons to facilitate their propagation, Mol Neurodegener
10 (2015) 57.
[14] C. Munch, J. O'Brien, A. Bertolotti, Prion-like propagation of mutant
superoxide dismutase-1 misfolding in neuronal cells, Proc Natl Acad Sci U S A
108(9) (2011) 3548-53.
[15] V. Sundaramoorthy, A.K. Walker, J. Yerbury, K.Y. Soo, M.A. Farg, V. Hoang, R.
Zeineddine, D. Spencer, J.D. Atkin, Extracellular wildtype and mutant SOD1
induces ER-Golgi pathology characteristic of amyotrophic lateral sclerosis in
neuronal cells, Cell Mol Life Sci 70(21) (2013) 4181-95.
[16] L.I. Grad, J.J. Yerbury, B.J. Turner, W.C. Guest, E. Pokrishevsky, M.A. O'Neill, A.
Yanai, J.M. Silverman, R. Zeineddine, L. Corcoran, J.R. Kumita, L.M. Luheshi, M.
Yousefi, B.M. Coleman, A.F. Hill, S.S. Plotkin, I.R. Mackenzie, N.R. Cashman,
Intercellular propagated misfolding of wild-type Cu/Zn superoxide dismutase
occurs via exosome-dependent and -independent mechanisms, Proc Natl Acad
Sci U S A 111(9) (2014) 3620-5.
[17] L. McAlary, J.J. Yerbury, J.A. Aquilina, Glutathionylation potentiates benign
superoxide dismutase 1 variants to the toxic forms associated with amyotrophic
lateral sclerosis, Sci Rep 3 (2013) 3275.
[18] J.D. Atkin, M.A. Farg, K.Y. Soo, A.K. Walker, M. Halloran, B.J. Turner, P. Nagley,
M.K. Horne, Mutant SOD1 inhibits ER-Golgi transport in amyotrophic lateral
sclerosis, J Neurochem 129(1) (2014) 190-204.
[19] A.S. Tsvetkov, M. Arrasate, S. Barmada, D.M. Ando, P. Sharma, B.A. Shaby, S.
Finkbeiner, Proteostasis of polyglutamine varies among neurons and predicts
neurodegeneration, Nat Chem Biol 9(9) (2013) 586-92.
[20] S.J. Weisberg, R. Lyakhovetsky, A.C. Werdiger, A.D. Gitler, Y. Soen, D.
Kaganovich, Compartmentalization of superoxide dismutase 1 (SOD1G93A)
aggregates determines their toxicity, Proc Natl Acad Sci U S A 109(39) (2012)
15811-6.
[21] F. Chiti, C.M. Dobson, Protein misfolding, functional amyloid, and human
disease, Annu Rev Biochem 75 (2006) 333-66.

690 [22] M.Y. Sherman, A.L. Goldberg, Cellular defenses against unfolded proteins: a
691 cell biologist thinks about neurodegenerative diseases, Neuron 29(1) (2001) 15-
692 32.
693 [23] J.J. Yerbury, J.R. Kumita, Protein chemistry of amyloid fibrils and
694 chaperones: Implications for amyloid formation and disease, Current Chemical
695 Biology 4(2) (2010) 89-98.
696 [24] Q. Huang, M.E. Figueiredo-Pereira, Ubiquitin/proteasome pathway
697 impairment in neurodegeneration: Therapeutic implications, Apoptosis 15(11)
698 (2010) 1292-1311.
699 [25] P. Ciryam, G.G. Tartaglia, R.I. Morimoto, C.M. Dobson, M. Vendruscolo,
700 Widespread aggregation and neurodegenerative diseases are associated with
701 supersaturated proteins, Cell Rep 5(3) (2013) 781-90.
702 [26] P. Ciryam, R. Kundra, R.I. Morimoto, C.M. Dobson, M. Vendruscolo,
703 Supersaturation is a major driving force for protein aggregation in
704 neurodegenerative diseases, Trends Pharmacol Sci 36(2) (2015) 72-7.
705 [27] S. Boeynaems, E. Bogaert, D. Kovacs, A. Konijnenberg, E. Timmerman, A.
706 Volkov, M. Guharoy, M. De Decker, T. Jaspers, V.H. Ryan, A.M. Janke, P. Baatsen, T.
707 Vercruysse, R.M. Kolaitis, D. Daelemans, J.P. Taylor, N. Kedersha, P. Anderson, F.
708 Impens, F. Sobott, J. Schymkowitz, F. Rousseau, N.L. Fawzi, W. Robberecht, P. Van
709 Damme, P. Tompa, L. Van Den Bosch, Phase Separation of C9orf72 Dipeptide
710 Repeats Perturbs Stress Granule Dynamics, Mol Cell 65(6) (2017) 1044-1055 e5.
711 [28] A. Hershko, A. Ciechanover, The ubiquitin system, Annu Rev Biochem 67
712 (1998) 425-79.
713 [29] A. Ciechanover, P. Brundin, The ubiquitin proteasome system in
714 neurodegenerative diseases: sometimes the chicken, sometimes the egg, Neuron
715 40(2) (2003) 427-46.
716 [30] C.M. Pickart, Mechanisms underlying ubiquitination, Annu Rev Biochem 70
717 (2001) 503-33.
718 [31] N.P. Dantuma, T.A. Groothuis, F.A. Salomons, J. Neefjes, A dynamic ubiquitin
719 equilibrium couples proteasomal activity to chromatin remodeling, J Cell Biol
720 173(1) (2006) 19-26.
721 [32] T.A. Groothuis, N.P. Dantuma, J. Neefjes, F.A. Salomons, Ubiquitin crosstalk
722 connecting cellular processes, Cell Div 1 (2006) 21.
723 [33] B.J. Turner, J.D. Atkin, M.A. Farg, D.W. Zang, A. Rembach, E.C. Lopes, J.D.
724 Patch, A.F. Hill, S.S. Cheema, Impaired extracellular secretion of mutant
725 superoxide dismutase 1 associates with neurotoxicity in familial amyotrophic
726 lateral sclerosis, J Neurosci 25(1) (2005) 108-17.
727 [34] S. Berginck, F.A. Salomons, D. Hoogstraten, T.A. Groothuis, H. de Waard, J. Wu,
728 L. Yuan, E. Citterio, A.B. Houtsmuller, J. Neefjes, J.H. Hoeijmakers, W. Vermeulen,
729 N.P. Dantuma, DNA damage triggers nucleotide excision repair-dependent
730 monoubiquitylation of histone H2A, Genes Dev 20(10) (2006) 1343-52.
731 [35] N.R. Cashman, H.D. Durham, J.K. Blusztajn, K. Oda, T. Tabira, I.T. Shaw, S.
732 Dahrouge, J.P. Antel, Neuroblastoma x spinal cord (NSC) hybrid cell lines
733 resemble developing motor neurons, Dev Dyn 194(3) (1992) 209-21.
734 [36] S. Yang, K.Y. Zhang, R. Kariawasam, M. Bax, J.A. Fifita, L. Ooi, J.J. Yerbury, G.A.
735 Nicholson, I.P. Blair, Evaluation of Skin Fibroblasts from Amyotrophic Lateral
736 Sclerosis Patients for the Rapid Study of Pathological Features, Neurotox Res
737 28(2) (2015) 138-46.

[37] C.A. Schneider, W.S. Rasband, K.W. Eliceiri, NIH Image to ImageJ: 25 years of
image analysis, Nat Methods 9(7) (2012) 671-5.
[38] S.R. Nagarajan, A.E. Brandon, J.A. McKenna, H.C. Shtein, T.Q. Nguyen, E.
Suryana, P. Poronnik, G.J. Cooney, D.N. Saunders, A.J. Hoy, Insulin and diet-
induced changes in the ubiquitin-modified proteome of rat liver, PLoS One 12(3)
(2017) e0174431.
[39] B. Turriziani, A. Garcia-Munoz, R. Pilkington, C. Raso, W. Kolch, A. von
Kriegsheim, On-beads digestion in conjunction with data-dependent mass
spectrometry: a shortcut to quantitative and dynamic interaction proteomics,
Biology (Basel) 3(2) (2014) 320-32.
[40] J. Cox, M. Mann, MaxQuant enables high peptide identification rates,
individualized p.p.b.-range mass accuracies and proteome-wide protein
quantification, Nat Biotechnol 26(12) (2008) 1367-72.
[41] D.R. Croucher, M. Iconomou, J.F. Hastings, S.P. Kennedy, J.Z. Han, R.F.
Shearer, J. McKenna, A. Wan, J. Lau, S. Aparicio, D.N. Saunders, Bimolecular
complementation affinity purification (BiCAP) reveals dimer-specific protein
interactions for ERBB2 dimers, Sci Signal 9(436) (2016) ra69.
[42] D. Szklarczyk, A. Franceschini, S. Wyder, K. Forslund, D. Heller, J. Huerta-
Cepas, M. Simonovic, A. Roth, A. Santos, K.P. Tsafou, M. Kuhn, P. Bork, L.J. Jensen,
C. von Mering, STRING v10: protein-protein interaction networks, integrated
over the tree of life, Nucleic Acids Res 43(Database issue) (2015) D447-52.
[43] M.E. Smoot, K. Ono, J. Ruscheinski, P.L. Wang, T. Ideker, Cytoscape 2.8: new
features for data integration and network visualization, Bioinformatics 27(3)
(2011) 431-2.
[44] R.K. Dagda, S.J. Cherra, 3rd, S.M. Kulich, A. Tandon, D. Park, C.T. Chu, Loss of
PINK1 function promotes mitophagy through effects on oxidative stress and
mitochondrial fission, J Biol Chem 284(20) (2009) 13843-55.
[45] N.E. Farrawell, I.A. Lambert-Smith, S.T. Warraich, I.P. Blair, D.N. Saunders,
D.M. Hatters, J.J. Yerbury, Distinct partitioning of ALS associated TDP-43, FUS and
SOD1 mutants into cellular inclusions, Sci Rep 5 (2015) 13416.
[46] N.F. Bence, R.M. Sampat, R.R. Kopito, Impairment of the ubiquitin-
proteasome system by protein aggregation, Science 292(5521) (2001) 1552-5.
[47] M. Mattiazzi, M. D'Aurelio, C.D. Gajewski, K. Martushova, M. Kiaei, M.F. Beal,
G. Manfredi, Mutated human SOD1 causes dysfunction of oxidative
phosphorylation in mitochondria of transgenic mice, J Biol Chem 277(33) (2002)
29626-33.
[48] C. Vande Velde, K.K. McDonald, Y. Boukhedimi, M. McAlonis-Downes, C.S.
Lobsiger, S. Bel Hadj, A. Zandona, J.P. Julien, S.B. Shah, D.W. Cleveland, Misfolded
SOD1 associated with motor neuron mitochondria alters mitochondrial shape
and distribution prior to clinical onset, PLoS One 6(7) (2011) e22031.
[49] W. Song, Y. Song, B. Kincaid, B. Bossy, E. Bossy-Wetzel, Mutant SOD1G93A
triggers mitochondrial fragmentation in spinal cord motor neurons:
neuroprotection by SIRT3 and PGC-1alpha, Neurobiol Dis 51 (2013) 72-81.
[50] J. Lowe, A. Blanchard, K. Morrell, G. Lennox, L. Reynolds, M. Billett, M.
Landon, R.J. Mayer, Ubiquitin is a common factor in intermediate filament
inclusion bodies of diverse type in man, including those of Parkinson's disease,
Pick's disease, and Alzheimer's disease, as well as Rosenthal fibres in cerebellar
astrocytomas, cytoplasmic bodies in muscle, and mallory bodies in alcoholic liver
disease, J Pathol 155(1) (1988) 9-15.

[51] H. Mori, J. Kondo, Y. Ihara, Ubiquitin is a component of paired helical
filaments in Alzheimer's disease, Science 235(4796) (1987) 1641-4.
[52] E.L. Scotter, C. Vance, A.L. Nishimura, Y.B. Lee, H.J. Chen, H. Urwin, V.
Sardone, J.C. Mitchell, B. Rogelj, D.C. Rubinsztein, C.E. Shaw, Differential roles of
the ubiquitin proteasome system and autophagy in the clearance of soluble and
aggregated TDP-43 species, J Cell Sci 127(Pt 6) (2014) 1263-78.
[53] K. Miyazaki, T. Fujita, T. Ozaki, C. Kato, Y. Kurose, M. Sakamoto, S. Kato, T.
Goto, Y. Itoyama, M. Aoki, A. Nakagawara, NEDL1, a novel ubiquitin-protein
isopeptide ligase for dishevelled-1, targets mutant superoxide dismutase-1, J Biol
Chem 279(12) (2004) 11327-35.
[54] J.M. Tan, E.S. Wong, D.S. Kirkpatrick, O. Pletnikova, H.S. Ko, S.P. Tay, M.W. Ho,
J. Troncoso, S.P. Gygi, M.K. Lee, V.L. Dawson, T.M. Dawson, K.L. Lim, Lysine 63-
linked ubiquitination promotes the formation and autophagic clearance of
protein inclusions associated with neurodegenerative diseases, Hum Mol Genet
17(3) (2008) 431-9.
[55] H. Wang, Z. Ying, G. Wang, Ataxin-3 regulates aggresome formation of
copper-zinc superoxide dismutase (SOD1) by editing K63-linked polyubiquitin
chains, J Biol Chem 287(34) (2012) 28576-85.
[56] A.M. Mabb, M.D. Ehlers, Ubiquitination in postsynaptic function and
plasticity, Annu Rev Cell Dev Biol 26 (2010) 179-210.
[57] H. Kawabe, N. Brose, The role of ubiquitylation in nerve cell development,
Nat Rev Neurosci 12(5) (2011) 251-68.
[58] Y. Tashiro, M. Urushitani, H. Inoue, M. Koike, Y. Uchiyama, M. Komatsu, K.
Tanaka, M. Yamazaki, M. Abe, H. Misawa, K. Sakimura, H. Ito, R. Takahashi, Motor
neuron-specific disruption of proteasomes, but not autophagy, replicates
amyotrophic lateral sclerosis, J Biol Chem 287(51) (2012) 42984-94.
[59] C. Cheroni, M. Marino, M. Tortarolo, P. Veglianese, S. De Biasi, E. Fontana,
L.V. Zuccarello, C.J. Maynard, N.P. Dantuma, C. Bendotti, Functional alterations of
the ubiquitin-proteasome system in motor neurons of a mouse model of familial
amyotrophic lateral sclerosis, Hum Mol Genet 18(1) (2009) 82-96.
[60] P. Ciryam, R. Kundra, R. Freer, R.I. Morimoto, C.M. Dobson, M. Vendruscolo,
A transcriptional signature of Alzheimer's disease is associated with a
metastable subproteome at risk for aggregation, Proc Natl Acad Sci U S A
113(17) (2016) 4753-8.
[61] Z. Tan, W. Qu, W. Tu, W. Liu, M. Baudry, S.S. Schreiber, p53 accumulation due
to down-regulation of ubiquitin: relevance for neuronal apoptosis, Cell Death
Differ 7(7) (2000) 675-81.
[62] Z. Tan, W. Tu, S.S. Schreiber, Downregulation of free ubiquitin: a novel
mechanism of p53 stabilization and neuronal cell death, Brain Res Mol Brain Res
91(1-2) (2001) 179-88.
[63] E.G. Mimnaugh, H.Y. Chen, J.R. Davie, J.E. Celis, L. Neckers, Rapid
deubiquitination of nucleosomal histones in human tumor cells caused by
proteasome inhibitors and stress response inducers: effects on replication,
transcription, translation, and the cellular stress response, Biochemistry 36(47)
(1997) 14418-29.
[64] A. Patnaik, V. Chau, J.W. Wills, Ubiquitin is part of the retrovirus budding
machinery, Proc Natl Acad Sci U S A 97(24) (2000) 13069-74.
[65] J. Hanna, D.S. Leggett, D. Finley, Ubiquitin depletion as a key mediator of
toxicity by translational inhibitors, Mol Cell Biol 23(24) (2003) 9251-61.

[66] J.H. Vaden, J.A. Watson, A.D. Howard, P.C. Chen, J.A. Wilson, S.M. Wilson,
Distinct effects of ubiquitin overexpression on NMJ structure and motor
performance in mice expressing catalytically inactive USP14, Front Mol Neurosci
8 (2015) 11.
[67] A. Hyrskyluoto, C. Bruelle, S.H. Lundh, H.T. Do, J. Kivinen, E. Rappou, S.
Reijonen, T. Waltimo, A. Petersen, D. Lindholm, L. Korhonen, Ubiquitin-specific
protease-14 reduces cellular aggregates and protects against mutant huntingtin-
induced cell degeneration: involvement of the proteasome and ER stress-
activated kinase IRE1alpha, Hum Mol Genet (2014).
[68] E. Kiskinis, J. Sandoe, L.A. Williams, G.L. Boulting, R. Moccia, B.J. Wainger, S.
Han, T. Peng, S. Thams, S. Mikkilineni, C. Mellin, F.T. Merkle, B.N. Davis-
Dusenbery, M. Ziller, D. Oakley, J. Ichida, S. Di Costanzo, N. Atwater, M.L. Maeder,
M.J. Goodwin, J. Nemesh, R.E. Handsaker, D. Paull, S. Noggle, S.A. McCarroll, J.K.
Joung, C.J. Woolf, R.H. Brown, K. Eggan, Pathways disrupted in human ALS motor
neurons identified through genetic correction of mutant SOD1, Cell Stem Cell
14(6) (2014) 781-95.
[69] S. Maharjan, M. Oku, M. Tsuda, J. Hoseki, Y. Sakai, Mitochondrial impairment
triggers cytosolic oxidative stress and cell death following proteasome inhibition,
Sci Rep 4 (2014) 5896.
[70] A. Ugun-Klusek, M.H. Tatham, J. Elkharaz, D. Constantin-Teodosiu, K. Lawler,
H. Mohamed, S.M. Paine, G. Anderson, R. John Mayer, J. Lowe, E. Ellen Billett, L.
Bedford, Continued 26S proteasome dysfunction in mouse brain cortical neurons
impairs autophagy and the Keap1-Nrf2 oxidative defence pathway, Cell Death
Dis 8(1) (2017) e2531.
[71] C. Vijayvergiya, M.F. Beal, J. Buck, G. Manfredi, Mutant superoxide dismutase
1 forms aggregates in the brain mitochondrial matrix of amyotrophic lateral
sclerosis mice, J Neurosci 25(10) (2005) 2463-70.
[72] D. Jaarsma, F. Rognoni, W. van Duijn, H.W. Verspaget, E.D. Haasdijk, J.C.
Holstege, CuZn superoxide dismutase (SOD1) accumulates in vacuolated
mitochondria in transgenic mice expressing amyotrophic lateral sclerosis-linked
SOD1 mutations, Acta Neuropathol 102(4) (2001) 293-305.
[73] L. Ruan, C. Zhou, E. Jin, A. Kucharavy, Y. Zhang, Z. Wen, L. Florens, R. Li,
Cytosolic proteostasis through importing of misfolded proteins into
mitochondria, Nature (2017).
[74] R. Yonashiro, A. Sugiura, M. Miyachi, T. Fukuda, N. Matsushita, R. Inatome, Y.
Ogata, T. Suzuki, N. Dohmae, S. Yanagi, Mitochondrial ubiquitin ligase MITOL
ubiquitinates mutant SOD1 and attenuates mutant SOD1-induced reactive
oxygen species generation, Mol Biol Cell 20(21) (2009) 4524-30.

Declarations

Ethics approval and consent to participate
Not applicable

Consent for publication
Not applicable

Availability of data and material
The datasets used and/or analysed during the current study available from the corresponding author on reasonable request.

Competing interests
The authors declare that they have no competing interests

Funding
I.A.L.S was supported by Rotary Health Australia. P.C. was supported by grants from the US-UK Fulbright Commission, St. John's College, University of Cambridge, and the National Institutes of Health (Northwestern University Medical Scientist Training Program Grant T32 GM8152-28). D.N.S, K.L.V and J.J.Y were supported by US Department of Defence (AL150057) and the Motor Neuron Disease Research Institute of Australia (Cunningham Family MND Research Grant, GIA1656).
 J.J.Y was also supported by grants from the National Health and Medical Research Council (1095215, 1084144).

Authors' contributions
N.E.F, P.C, L.M, K.L.V, D.N.S, and J.J.Y designed research; N.E.F, P.C, I.A.L.S, L.M, J.M, and K.M performed research; N.E.F, I.A.L.S, P.C, L.M, K.L.V, D.N.S, and J.J.Y analyzed data; and N.E.F, P.C, D.N.S, and J.J.Y wrote the paper. All authors read and approved the final manuscript.

Acknowledgements
Not applicable

Authors' information (optional)

923
924
Figure 1: Ubiquitin exists in a dynamic equilibrium in the cell, partitioning into
mobile and immobile fractions. Schematic representation of Ub homeostasis,
showing the different methods used to interrogate various aspects of this system in
this study. Ub is synthesised from several genes, generating a pool of free Ub that
consists of monomeric and polymeric chains of Ub. Ub can be conjugated with
substrate proteins via the enzymatic activity of a large number of Ub ligases. The
cellular fate or outcome of such ubiquitylation is dictated by the chain type formed by
Ub. The mobile fraction of Ub measured by fluorescence recovery after
photobleaching (FRAP) is the sum of monomeric and polymeric free Ub and
diffusible ubiquitylated protein (blue box). The immobile pool of Ub is that
conjugated to large slowly diffusing structures such as vesicles, organelles, histones
and cytoskeleton (green box). Ubiquitin proteasome system (UPS) degradation
function can be compared using fluorescent reporters. Generally accumulation of
reporters suggests increased competition, or decreased rates of degradation. The
ubiquitin-modified proteome can be quantified using enrichment capture and mass
spectrometry (ubiqitomics). A subset of mobile Ub is free monomeric Ub can be
measured using fluorescence recovery after nucleus photobleaching (FRANP).
Monomeric Ub is the only form of tagged Ub that can diffuse through the nuclear
pore, so by bleaching the entire nucleus we can measure monomeric Ub as it diffuses
back in. In the context of protein aggregation a proportion of Ub is conjugated to
substrates within the inclusions, we can quantify the proportion of Ub incorporated
into inclusions using image analysis.
947

948
949
Figure 2: SOD1 co-aggregates with ubiquitin. (A) NSC-34 cells co-transfected with
SOD1^{A4V}-GFP and mRFP-Ub were imaged every 15 min for 17 h. (B) NSC-34 cells
overexpressing SOD1^{A4V}-GFP were fixed, permeabilized and stained for Ub$_{K48}$ and
Ub$_{K63}$ polyubiquitin chains 48 h post transfection. (C) Spinal cord from the SOD1^{G93A}
mouse was fixed, permeabilized and stained for neuron-specific beta III tubulin,
SOD1 and Ub$_{K48}$. Co-localization of SOD1 and Ub is indicated with arrows.
956

957
958
Figure 3: Mutant SOD1^{A4V} alters UPS activity. (A) Dose dependant response of
UPS activity (tdTomato^{CL1} fluorescence) in NSC-34 cells co-transfected with SOD1-
GFP after overnight treatment with the proteasome inhibitor MG132. Data represent
mean CL1-Degron fluorescence ± SEM (n=3), ** P< 0.01; *** P<0.001. (B)
Frequency distribution analysis of SOD1-GFP fluorescence was performed on cells
expressing soluble (diffuse) SOD1^{A4V} and aggregated SOD1^{A4V}. (C) Cells containing
SOD1^{A4V} aggregates exhibited significantly higher fluorescence than cells that did not
contain aggregates, **** P< 0.0001. (D) 80% of cells expressing the highest SOD1-
GFP signal (top 10%) contained aggregates. (E) Cells expressing the highest GFP
signal typically contained aggregates and revealed greater significant differences in
tdTomato^{CL1} fluorescence between cells expressing SOD1^{WT} and SOD1^{A4V}.
970

971
Figure 4: Ubiquitin distribution is altered in cells containing SOD1^{A4V}
aggregates. (A) NSC-34 cells co-transfected with SOD1-GFP and mCherry-Ub were
photobleached in either the nucleus or cytoplasm and recovery of Ub fluorescence
was monitored for 120 s. Data shown are means ± SEM (n=3) and are representative
of 3 independent experiments. (B) Representative confocal images of pre-bleach,
post-bleach and recovery endpoint are shown with the ROI marked in yellow. (C) The
amount of Ub incorporated into SOD1^{A4V} inclusions was quantified using ImageJ. (D)
Diffusion rates of mCherry-Ub measured in both the nucleus and cytoplasm of co-
transfected NSC-34 cells. Error bars are SEM (n ≥ 6, combined from 3 independent
experiments). (E) Quantification of the proportion of mobile Ub in the nucleus and
cytoplasm of cells expressing either SOD1^{WT}, soluble SOD1^{A4V} or insoluble
SOD1^{A4V}. SOD1^{A4V} insoluble (TOTAL) represents total Ub levels taking into account
the proportion of Ub incorporated into SOD1 inclusions (determined in (C)). Data
shown are means ± SEM combined from 3 independent experiments (n ≥ 7). One way
Anova was used to compare differences. * and † indicate significant difference to
SOD1^{A4V} insoluble (TOTAL) in the cytoplasm and nucleus respectively. * † P < 0.05,
** † † P < 0.01 .
989

Figure 5: Reduced levels of free monomeric ubiquitin in NSC-34 cells containing
SOD1^{A4V} aggregates. (A) Western blot analysis of cell lysates of NSC-34 cells
transiently transfected with SOD1-GFP, mRFP-Ub or eGFP control. Samples were
separated under reducing conditions and probed with anti-ubiquitin antibody. (B) The
entire nucleus of co-transfected NSC-34 cells was photobleached and the recovery of
nuclear Ub was monitored as a proportion of cytoplasmic fluorescence for 120 s. Data
shown are means ± SEM (n ≥ 17) combined from 3 independent experiments. (C)
Representative confocal images of pre-bleach, post-bleach and final read. (D). The
percentage of mobile Ub in the nucleus at the final read was quantified as a proportion
of cytoplasmic fluorescence. Data represent mean ± SEM (n ≥ 17, combined from 3
independent experiments)* P < 0.05, ** P < 0.01, *** P<0.001.

1002
1003
1004
1005

1006
1007
Figure 6: The ubiquitylated proteome of transfected NSC-34 cells. (A) NSC-34
cells expressing GFP fusions of SOD1^{WT} or SOD1^{A4V} were subjected to high affinity
purification to isolate ubiquitylated proteins which were subsequently identified by
LC-MS/MS. (B) Venn analysis comparing the number of ubiquitylated proteins in
NSC-34 cells transfected with either SOD1^{A4V} (n=2) or SOD1^{WT} (n=3). (C) STRING
analysis of protein-protein interaction network of ubiquitylated proteins unique to
either SOD1^{A4V} or SOD1^{WT}. Proteins were identified as present in each condition if
they were present in at least two replicates for that condition. Additionally proteins
were only considered "unique" to either SOD1^{WT} or SOD1^{A4V} if they were not
identified in any of the replicates for any of the other conditions. (D) The median
supersaturation score calculated for the unfolded (σ_u) and native (σ_f) states of proteins
unique to the SOD1^{A4V} ubiquitome. Fold Δ refers to the increase in supersaturation
score from the known mouse proteome.
1021

1022
1023
Figure 7: Cells containing SOD1 aggregates display altered mitochondrial
morphology and dysfunction. (A) NSC-34 cells transiently transfected with SOD1-
GFP were stained for mitochondria 48 h post transfection with Mitotracker Deep Red.
The number of mitochondria (B) and circularity (C) was determined using a
mitochondrial morphology macro in ImageJ. Data represent mean ± SEM (n ≥ 23) * P
< 0.05, ** P < 0.01. (D) Mitotracker fluorescence was also quantified by flow
cytometry in cells expressing the highest levels of SOD1-GFP. (E) Mitochondrial
membrane potential was examined through the accumulation of Mitotracker Red
CMXRos. Data represent mean ± SEM (n = 3) *** P<0.001.
1033

1034

1035

1036

1037 **Figure 8: Disrupted Ub homeostasis in ALS**. Mutations in SOD1-associated ALS
1038 disrupt Ub homeostasis, either directly or through sequestration of Ub into protein
1039 aggregates. These changes result in altered Ub distribution and subsequent depletion
1040 of free Ub, eventually reaching a threshold below which vital cellular functions are
1041 severely compromised, and ultimately result in cell death.

1042

1043

1044

1045

1046

1047

1048

Table 1. Proteins identified unique to A4V or WT SOD1:

Unique to SOD[1A4V]

Entry	Entry name	Protein names	Gene names	Unique peptides
P14206	RSSA_MOUSE	40S ribosomal protein SA	Rpsa Lamr1 P40-8	5
P47753	CAZA1_MOUSE	F-actin-capping protein subunit alpha-1	Capza1 Cappa1	3
Q91V12	BACH_MOUSE	Cytosolic acyl coenzyme A thioester hydrolase	Acot7 Bach	3
Q9QUR6	PPCE_MOUSE	Prolyl endopeptidase	Prep Pep	4
Q922D8	C1TC_MOUSE	C-1-tetrahydrofolate synthase	Mthfd1	2
P45376	ALDR_MOUSE	Aldose reductase	Akr1b1 Akr1b3 Aldor1 Aldr1	4
Q8CGZ0	CHERP_MOUSE	Calcium homeostasis endoplasmic reticulum protein	Cherp Scaf6	3
Q8R081	HNRPL_MOUSE	Heterogeneous nuclear ribonucleoprotein L	Hnrnpl Hnrpl	2
O08784	TCOF_MOUSE	Treacle protein	Tcof1	2
O09061	PSB1_MOUSE	Proteasome subunit beta type-1	Psmb1	2
O54734	OST48_MOUSE	Dolichyl-diphosphooligosaccharide--protein glycosyltransferase 48 kDa subunit	Ddost	4
P29341	PABP1_MOUSE	Polyadenylate-binding protein 1	Pabpc1 Pabp1	4
P35564	CALX_MOUSE	Calnexin	Canx	5
P62492	RB11A_MOUSE	Ras-related protein Rab-11A	Rab11a Rab11	2
P54729	NUB1_MOUSE	NEDD8 ultimate buster 1	Nub1 Nyren18	2
P56135	ATPK_MOUSE	ATP synthase subunit f, mitochondrial	Atp5j2	2
P56959	FUS_MOUSE	RNA-binding protein FUS	Fus	3
Q7TPR4	ACTN1_MOUSE	Alpha-actinin-1	Actn1	5
P61957	SUMO2_MOUSE	Small ubiquitin-related modifier 2	Sumo2 Smt3b Smt3h2	2
P62717	RL18A_MOUSE	60S ribosomal protein L18a	Rpl18a	4

Q3THS6	METK2_MOUSE	S-adenosylmethionine synthase isoform type-2 (AdoMet synthase 2)	Mat2a	2
Q3TIU4	PDE12_MOUSE	2',5'-phosphodiesterase 12 (2'-PDE) (2-PDE)	Pde12	2
Q3U2C5	RN149_MOUSE	E3 ubiquitin-protein ligase RNF149	Rnf149 Greul4	2
Q60692	PSB6_MOUSE	Proteasome subunit beta type-6	Psmb6 Lmp19	3
Q61074	PPM1G_MOUSE	Protein phosphatase 1G	Ppm1g Fin13 Ppm1c	3
Q61081	CDC37_MOUSE	Hsp90 co-chaperone Cdc37	Cdc37	2
Q61990	PCBP2_MOUSE	Poly(rC)-binding protein 2 (Alpha-CP2)	Pcbp2 Cbp Hnrnpx Hnrpx	2
Q62318	TIF1B_MOUSE	Transcription intermediary factor 1-beta (TIF1-beta) (E3 SUMO-protein ligase TRIM28)	Trim28 Kap1 Krip1 Tif1b	4
Q64514	TPP2_MOUSE	Tripeptidyl-peptidase 2 (TPP-2)	Tpp2	2
Q8QZT1	THIL_MOUSE	Acetyl-CoA acetyltransferase, mitochondrial	Acat1	2
Q8VCN5	CGL_MOUSE	Cystathionine gamma-lyase	Cth	3
Q8VDW0	DX39A_MOUSE	ATP-dependent RNA helicase DDX39A	Ddx39a Ddx39	2
Q7TMK9	HNRPQ_MOUSE	Heterogeneous nuclear ribonucleoprotein Q (hnRNP Q)	Syncrip Hnrpq Nsap1 Nsap1l	3
Q91YQ5	RPN1_MOUSE	Dolichyl-diphosphooligosaccharide--protein glycosyltransferase subunit 1	Rpn1	5
P17225	PTBP1_MOUSE	Polypyrimidine tract-binding protein 1 (PTB)	Ptbp1 Ptb	4
Q99LX5	MMTA2_MOUSE	Multiple myeloma tumor-associated protein 2 homolog	Mmtag2	2
Q9CQ60	6PGL_MOUSE	6-phosphogluconolactonase (6PGL)	Pgls	5
Q9CRB9	MIC19_MOUSE	MICOS complex subunit Mic19	Chchd3 Mic19	3
Q9CWF2	TBB2B_MOUSE	Tubulin beta-2B chain	Tubb2b	3
Q9CWJ9	PUR9_MOUSE	Bifunctional purine biosynthesis protein PURH	Atic Purh	4
Q9CZ13	QCR1_MOUSE	Cytochrome b-c1 complex subunit 1, mitochondrial (Complex III subunit 1)	Uqcrc1	4
Q9D0E1	HNRPM_MOUSE	Heterogeneous nuclear ribonucleoprotein M	Hnrpm Hnrpm	3
Q9D0M3	CY1_MOUSE	Cytochrome c1, heme protein, mitochondrial (Complex III subunit 4)	Cyc1	3
Q9DCH4	EIF3F_MOUSE	Eukaryotic translation initiation factor 3 subunit F	Eif3f Eif3s5	4
Q9DCL9	PUR6_MOUSE	Multifunctional protein ADE2	Paics	5
Q9JJI8	RL38_MOUSE	60S ribosomal protein L38	Rpl38	3
Q9JM13	RABX5_MOUSE	Rab5 GDP/GTP exchange factor	Rabgef1 Rabex5	3

Q9JMH6	TRXR1_MOUSE	Thioredoxin reductase 1, cytoplasmic	Txnrd1 Trxr1	2
Q9QZD9	EIF3I_MOUSE	Eukaryotic translation initiation factor 3 subunit I	Eif3i Eif3s2 Trip1	4
Q9WU78	PDC6I_MOUSE	Programmed cell death 6-interacting protein	Pdcd6ip Aip1 Alix	4
Q9Z1Z2	STRAP_MOUSE	Serine-threonine kinase receptor-associated protein	Strap Unrip	2
Q9Z2X1	HNRPF_MOUSE	Heterogeneous nuclear ribonucleoprotein F	Hnrnpf Hnrpf	3

Unique to SOD^{WT}

B2RY56	RBM25_MOUSE	RNA-binding protein 25	Rbm25	2
P51410	RL9_MOUSE	60S ribosomal protein L9	Rpl9	5
P04370	MBP_MOUSE	Myelin basic protein	Mbp Shi	4
P43276	H15_MOUSE	Histone H1.5	Hist1h1b H1f5	3
P62911	RL32_MOUSE	60S ribosomal protein L32	Rpl32	3
P63166	SUMO1_MOUSE	Small ubiquitin-related modifier 1 (SUMO-1)	Sumo1 Smt3c Smt3h3 Ubl1	2
P97350	PKP1_MOUSE	Plakophilin-1	Pkp1	17
Q60932	VDAC1_MOUSE	Voltage-dependent anion-selective channel protein 1 (VDAC-1)	Vdac1 Vdac5	8
Q6ZQ38	CAND1_MOUSE	Cullin-associated NEDD8-dissociated protein 1	Cand1 D10Ertd516e Kiaa0829	2
Q7TSF1	DSG1B_MOUSE	Desmoglein-1-beta	Dsg1b Dsg5	6
Q9Z315	SNUT1_MOUSE	U4/U6.U5 tri-snRNP-associated protein 1	Sart1 Haf	2

SOD1^{A4V} aggregation alters ubiquitin homeostasis in a cell model of ALS

Natalie E. Farrawell[1,2], Isabella Lambert-Smith[1,2], Kristen Mitchell[1,2], Jessie McKenna[3], Luke McAlary[1,2,4], Prajwal Ciryam[5,6,7], Kara L. Vine[1,2], Darren N. Saunders[3], Justin J. Yerbury[1,2*]

[1]Illawarra Health and Medical Research Institute, Wollongong, NSW, Australia
[2]School of Biological Sciences, Centre of Medicine and Molecular Bioscience, University of Wollongong, Northfields Ave, Wollongong, NSW, Australia, 2522
[3]School of Medical Sciences, Faculty of Medicine, UNSW Australia
[4]Department of Physics & Astronomy, University of British Columbia, Vancouver, British Columbia, Canada.
[5]Department of Chemistry, University of Cambridge, Cambridge CB2 1EW, UK
[6]Department of Molecular Biosciences, Rice Institute for Biomedical Research, Northwestern University, Evanston, IL 60208-3500, USA
[7]Department of Neurology, Columbia University College of Physicians & Surgeons, New York, NY 10032-3784

Supplementary Methods

To quantify SOD1 inclusion formation, images were acquired as described previously [1]. Briefly, z-stack images of NSC-34 cells transfected with SOD1-GFP constructs were acquired 48 h post transfection using the 63X objective, 512 x 512 pixels, 1 line and frame average and a scan speed of 400 Hz. Z-stacks were processed into a single image using LAS-AF Lite software (Leica) and at least 100 transfected cells were analysed for the presence of inclusions.

Toxicity Assay. The toxicity of cells expressing SOD1 was monitored over 68 h in an Incucyte automated fluorescent microscope (Essen BioScience, USA) as described in [1]. Cells were dissociated 24 h post transfection and replated in 96-well plates at a confluency of 20% in phenol red free DMEM/F12 containing 10% FBS. Images were acquired every 2 h and analysed using a processing definition trained to select GFP positive cells. The number of GFP positive cells was normalised to time zero before SOD1^{A4V} numbers were adjusted to SOD1^{WT} values.

Supplementary Figures

Figure S1: mcherry-Ub behaves in a similar manner to GFP-Ub. (A) NSC-34 cells transfected with either GFP-Ub (A) or mcherry-Ub (B) were photobleached in the cytoplasm or nucleus and recovery of Ub fluorescence was monitored for up to 170 s. Representative confocal images of prebleach, postbleach and recovery endpoint are shown with the ROI marked in yellow.

Figure S2: SOD1^{A4V} expression induces inclusion formation and is associated with increased cell death in NSC-34 cells. (A) Inclusion formation in NSC-34 cells expressing SOD1-GFP was determined 48 h post transfection. (B) The toxicity of SOD1^{A4V} was monitored in comparison to SOD1^{WT} over a 68 h period.

[1] L. McAlary, J.A. Aquilina, J.J. Yerbury, Susceptibility of Mutant SOD1 to Form a Destabilized Monomer Predicts Cellular Aggregation and Toxicity but Not In vitro Aggregation Propensity, Front Neurosci 10 (2016) 499.