ISG15 induces IL-10 production in human monocytes and is a biomarker of

disease severity during active tuberculosis

Paula Fernandes dos Santos[*], Johan Van Weyenbergh[¶], Murilo Delgobo[*], Daniel de

Oliveira Patricio[*], Brian J. Ferguson[§], Rodrigo Guabiraba[#], Tim Dierckx[¶], Soraya

Maria Menezes[¶], André Báfica[*★] and Daniel Santos Mansur[*★1]

*Laboratory of Immunobiology, Department of Microbiology, Immunology and

Parasitology, Universidade Federal de Santa Catarina, Campus Trindade, Centro de

Ciências Biológicas, Bloco A, sala 213, Santa Catarina, Brasil. CEP 88040-900

§Department of Pathology, University of Cambridge

#ISP, INRA, Université François Rabelais de Tours, 37380 Nouzilly, France

¶Department of Microbiology and Immunology, Rega Institute for Medical Research,

Laboratory for Clinical and Epidemiological Virology, KU Leuven - University of

Leuven, Leuven, Belgium

[1] Lead Contact

★Correspondence:

daniel.mansur@ufsc.br (D.S.M)

andre.bafica@ufsc.br (A.B.)

Running title

Human ISG15 drives a monocyte/IL-10 axis that is disrupted in active tuberculosis

Abstract (150 words)

Interferon stimulated gene 15 (ISG15) deficiency in humans leads to severe

interferonopathies and mycobacterial disease, the latter being previously associated to

its extracellular cytokine-like activity. Here, we demonstrate a novel role for secreted

ISG15 as an IL-10 inducer, unique to human primary monocytes. Employing *ex vivo*

systems analysis of human transcriptome datasets, we observed a significant

correlation of ISG15-induced monocyte IL-10 and lymphocyte IFNγ balanced

expression. This effect was associated with p38 MAPK and PI3K signalling in

healthy volunteers. The specificity and MAPK/PI3K-dependence of ISG15-induced

monocyte IL-10 production was confirmed *in vitro* using CRISPR/Cas9 knockout and

pharmacological inhibitors. Moreover, this *ISG15/IL10* axis was amplified in leprosy

but disrupted in human active tuberculosis (TB) patients. Importantly, ISG15 strongly

correlated with inflammation and disease severity during active TB. In conclusion,

this study identifies a novel anti-inflammatory ISG15/IL-10 myeloid axis that is

disrupted in active TB, revealing a potential biomarker for disease severity in this

major human disease.

Introduction

Type one interferons (IFN-I) exert most of their functions by inducing the expression of interferon-stimulated genes (ISGs). To date, over 300 ISGs have been described (de Veer et al., 2001; Der et al., 1998) and interferon stimulated gene of 15 KDa (ISG15) is prominently expressed in response to infection, in autoimmune diseases, cancer and physiological processes such as pregnancy (Dos Santos and Mansur, 2017; Hansen and Pru, 2014; Henkes et al., 2015; Hermann and Bogunovic, 2017; Tecalco Cruz and Mejia-Barreto, 2017; Wang et al., 2017a). ISG15 is synthesized as a 17 KDa precursor that is cleaved in the C-terminal region producing a mature form of 15 KDa. Also called ubiquitin cross-reactive protein (UCRP), ISG15 was the first ubiquitin-like protein to be described and it can be covalently linked to other proteins in a process called ISGylation (Dos Santos and Mansur, 2017; Haas et al., 1987; Loeb and Haas, 1992; Skaug and Chen, 2010). ISGylation is important for host defence against several viruses such as Influenza A, Vaccinia, Ebola, HIV and Hepatitis C virus in both human and mouse models (Morales and Lenschow, 2013; Schoggins and Rice, 2011; Skaug and Chen, 2010).

In addition to its intracellular ISGylation-mediated processes, the mature form of ISG15 can be secreted and possesses cytokine-like activities modulating leukocyte functions (Bogunovic et al., 2013; Dos Santos and Mansur, 2017). For instance, soluble ISG15 was found to enhance production of IFNγ by lymphocytes and NK cells (Bogunovic et al., 2012; D'Cunha et al., 1996) and to stimulate NK cell proliferation (D'Cunha et al., 1996) as well as neutrophil migration (Owhashi et al., 2003). Importantly, ISG15 deficiency in humans is associated with a severe Mendelian susceptibility to mycobacterial disease (Bogunovic et al., 2012) and cells from patients with a nonsense mutation or a frame-shift in *isg15* are deficient in IFNγ-

mediated immunity. This activity is attributed to the effects of extracellular ISG15 in

NK cells and possibly occurs through an unknown receptor (Bogunovic et al., 2012).

Furthermore, humans lacking ISG15 also develop exacerbated IFN-I-induced

immunopathology (Zhang et al., 2015). This evidence suggests that extracellular or

free ISG15 may regulate multiple aspects of the host immune response to pathogens

and implicate this protein as an important component induced during infection and/or

inflammatory processes involving IFN-I signalling. However, despite its ability to

induce pro-inflammatory mediators, IFN-I may also display opposing anti-

inflammatory effects (Billiau, 2006; Borden et al., 2007; McNab et al., 2015).

Nevertheless, whether soluble extracellular ISG15 modulates anti-inflammatory

responses has not been reported.

The present study demonstrates that ISG15 induces IL-10 synthesis in human primary

monocytes through MAPK- and PI3K-dependent pathways. Additionally, analysis of

human transcriptome data sets identified a myeloid *ISG15*/*IL10* axis present in

homeostasis. In contrast, the *ISG15*/*IL10* axis is disrupted during active TB and

ISG15 mRNA levels strongly correlate with inflammatory and disease severity

markers. These data suggest ISG15 may play a role in the crosstalk between Type

I/Type II IFNs and IL-10 and reveal *ISG15* mRNA levels as potentially useful

biomarker in human active TB.

Results and Discussion

ISG15 induces IL-10 production in human PBMC

Extracellular ISG15 stimulates IFNγ production by human NK cells (Bogunovic et al., 2012), so to investigate whether ISG15 regulates synthesis of other inflammatory cytokines, PBMCs were exposed to soluble ISG15 and 24h cell culture supernatants assayed for several cytokines by cytometric bead array (CBA). Out of this panel, only IL-10 and IL-6 were induced by ISG15 (Supplementary figure 1A). IL-10 is a key immune-regulatory cytokine that exerts opposing effects to IFNγ, hence this result was further assessed by treating PBMCs with different concentrations of pro- or mature ISG15 indicating a concentration-dependent response (Figure 1A). Following intracellular processing of pro-ISG15, its C-terminal LRLRGG domain is exposed and the protein becomes mature, a necessary requirement for ISGylation (Knight et al., 1988; Loeb and Haas, 1992; Narasimhan et al., 1996; Potter et al., 1999). Both pro- and mature ISG15 induced IL-10 secretion in human PBMCs in a similar manner (Figure 1A), indicating that LRLRGG sequence does not need to be exposed for ISG15-mediated IL-10 production. Exogenous ISG15 stimulated IL-10 synthesis by PBMCs from most of the healthy donors tested (Figure 1B). Control experiments showed that heat denatured ISG15 did not promote IL-10 synthesis demonstrating this protein requires its folded structure to induce cell signalling (Figure 1B). Kinetic analysis of ISG15-stimulation in human PBMCs showed a peak of IL-10 mRNA and protein synthesis after 6 and 12 hours respectively (Figure 1C and D). Interestingly, this response was found to be specific for primary cells as a library of human cell lines (NKL, NK92, THP-1, Karpas, U937 and Jurkat) treated with ISG15 did not produce IL-10 (data not shown). Additionally, ISG15 treatment did not induce cell death by means of annexin V expression and propidium iodide (PI) incorporation

(Figure 1E-G) suggesting IL-10 was actively secreted, not released from nor induced

by apoptotic or necrotic cells. Together, these data show that ISG15 induces IL-10

synthesis and secretion by primary human PBMCs, independent of cell death.

CD14$^+$ cells are the main producers of ISG15-induced IL-10

ISG15 can act on different cell types (Bogunovic et al., 2012; D'Cunha et al., 1996;

Owhashi et al., 2003; Recht et al., 1991) hence intracellular cytokine staining was

used to identify the source of ISG15-induced IL-10 in PBMCs subpopulations. These

experiments indicated that a CD14$^+$ population is the main source of IL-10 (Figure

2A) with an average 2.5 fold-increase of CD14$^+$IL-10$^+$ cells as compared to

unstimulated cultures (Figure 2B). Next, PBMCs were separated into CD14$^+$ and

CD14$^-$ populations and both groups were exposed to soluble ISG15. Quantification of

IL-10 and IFNγ 24 hours post stimulation confirmed CD14$^+$ population to be the main

producers of IL-10 (Figure 2C) whilst we corroborated previous work showing the

CD14$^-$ population to be the source of ISG15-induced IFNγ (Figure 2D) (Bogunovic et

al., 2012; D'Cunha et al., 1996). Additionally, these data indicate that recombinant

ISG15-induced IL-10 synthesis by CD14$^+$ populations does not require the presence

of CD14$^-$ cells.

To test whether endogenously produced ISG15 stimulates IL-10 synthesis, a co-

culture experiment was set up using a lung epithelial cell line, A549, as a source of

ISG15 (Wang et al., 2017b). For these assays, an *ISG15*-knockout (KO) A549 cell

line was generated using CRISPR/Cas9 technology (Supplementary Figure 1B, clone

3). Wild type (WT) or *ISG15*-KO A549 cells were then co-cultured with purified

human primary CD14$^+$ cells or stimulated with LPS as a positive control. In this

setup, A549-monocyte co-cultures led to a consistent production of IL-10, an outcome

completely abrogated when *ISG15*-KO A549 cells were used. This effect could be

rescued by re-introduction of the *ISG15* gene into the knockout cells (Figure 2E) thus

demonstrating the specificity of epithelial cell-derived ISG15 for the induction IL-10.

To study co-regulation of *ISG15/IL10/IFNG* pathways in different cell types *ex vivo*,

we next examined transcriptome datasets of purified major human leukocyte subsets

(ImmuCo, ImmuSort) (Wang et al., 2015a; Wang et al., 2015b). Expression levels of

ISG15 and *IL10* are positively correlated in total PBMCs, purified monocytes and

macrophages, but not neutrophils and T-cells (Figures 2F-J). Although neutrophils

display the highest *ISG15/IL10* expression ratio, monocytes are the main *ex vivo IL10*

expressing cell type (60.19% of cells with IL10 transcripts above detection limit, vs.

17.33% in PBMCs and 5.97% in neutrophils, Fig. 2K), thus corroborating with earlier

in vitro results (Fig. 2A-D). Consistent with a previous study (Tamassia et al., 2013),

low or undetectable *IL10* transcripts in human neutrophils (Fig. 2K) are explained by

the inactive chromatin configuration of the *IL10* locus in these cells. Together, this set

of results suggests a role for extracellular rather than intracellular ISG15 as inducer of

monocyte-derived IL-10 (this study) and NK-derived IFNγ (Bogunovic et al., 2012).

Since the unique susceptibility of ISG15-deficient children to low virulence

mycobacteria has underscored a role for extracellular ISG15 (Bogunovic et al., 2012),

we performed a systems analysis approach to gain insights on the possible influence

of ISG15/IL-10 axis during mycobacterial exposure in humans.

IL-10 production in response to ISG15 requires MAPK-PI3K signalling pathways

Mitogen-activated protein kinase (MAPK) and phosphatidylinositol-4,5-bisphosphate

3-kinase (PI3K) signalling pathways have been shown to participate in *IL10*

transcription in human monocytes and macrophages. For instance, p38, ERK1/2 and

PI3K are crucial for IL-10 synthesis during microbial stimuli such as LPS and

Mycobacterium (Ma et al., 2001; Nair et al., 2009). Thus, we analysed a published

transcriptome dataset of latent TB using WebGEstalt and Ingenuity Pathway Analysis

(IPA). As shown in Supplementary Table, MAPK and PI3K signalling pathways were

significantly enriched in latent TB transcriptomes, as compared to healthy, uninfected

controls. We next investigated whether members of MAPK and PI3K signalling

families displayed divergent expression patterns between latent and active TB. MAPK

family members were up-regulated in both latent and active TB (Figure 3A).

However, *PIK3CA* (PI3Kalpha) levels were up-regulated in active TB and down-

regulated in latent TB. In this scenario *MAPK14* (p38) expression levels were

significantly and positively correlated to *ISG15* levels *ex vivo* (Figure 3B).

Additionally, *MAPK3* (MAP3K/ERK1) levels were positively correlated to *IL10*

(Figure 3C) and negatively correlated to *IFNG* transcript levels (Figure 3D). Finally,

PIK3CA and *PIK3CB* (PI3Kbeta) transcripts were negatively correlated to *IL10*

expression levels (Figure 3E). These data suggested that ISG15/IL-10 as well as

MAPK/PI3K-associated transcripts are co-regulated during mycobacterial stimulation

in vivo and raised the possibility that MAPK/PI3K pathway is involved in ISG15-

induced IL-10 responses in monocytes. Indeed, following exposure of CD14[+] cells to

ISG15, increased phosphorylation of p38 MAPK was observed (Figure 3F). More

importantly, the use of two distinct inhibitors for p38 (Figures 3G-H) as well as

inhibitors for MEK1/2 (Figure 3I) and PI3K (Figure 3J) abrogated IL-10 production

in ISG15-stimulated monocytes. In contrast, chloroquine, an inhibitor targeting DNA-

PKCs/TLR9/endosome signalling pathways, did not affect IL-10 synthesis induced by

ISG15 (Figure 3J). These results suggest a central role for p38 activation and MAPK

as well as PI3K signalling in ISG15-induced IL-10 production by primary monocytes.

An ISG15/IL10/IFNG cluster in healthy controls is disrupted during active TB

Altogether, our data indicated that ISG15 is associated with immunoregulatory

responses and it could have an important role in mycobacterial-induced inflammation.

To test this concept, publicly available transcriptome data sets from established

cohorts of healthy controls and patients with leprosy as well as latent or active

tuberculosis were examined. Positive and negative correlations (Spearman Rho) were

calculated between normalized transcript levels of MAPK/PI3K/STAT signalling

family members, established myeloid lineage markers (*CD14*,

CD16=*FCGR3A/FCGR3B*, *ITGAM*, *ITGAX*) and lymphoid lineage markers (*CD4*,

CD8, CD56=*NCAM1*, *ITGAL*) plus *ISG15*, *IL10* and *IFNG* transcripts. Unsupervised

hierarchical clustering of these transcripts was then performed, based on the resulting

correlation matrices. In healthy controls (Figure 4A), *ISG15* mRNA strongly clusters

with *IL10*, and to a lesser extent with *IFNG*, indicating the existence of a regulatory

balance between pro-and anti-inflammatory effects under homeostatic conditions.

Leprosy lesions have been shown to express both type I IFN and IL-10, a scenario

that leads to suppression of IFNγ effector activities (Teles et al., 2013). An

unsupervised hierarchical cluster analysis of the cohort published by Teles and

colleagues shows the myeloid anti-inflammatory *ISG15*/*IL10* axis maintained in

leprosy lesions (Figure 4B), demonstrated by a single expression cluster comprised of,

ISG15, *IL10* and monocyte (*CD14*) and myeloid markers (CD64=*FCGR1*,

CD11c=*ITGAX*, PU.1=*SPI1*, CD16=*FCGR3*). Consistent with previous data (Teles et

al., 2013), this disease cluster was negatively correlated to a "protective"

CD8/*IFNG*/*STAT4* cluster, which is associated with milder (borderline

tuberculoid/paucibacillary) clinical form, whereas the *ISG15*/*IL10*/*CD14* cluster was

associated with the severe (lepromatous/multibacillary) disease form. Surprisingly,

the *ISG15*/*IL10* axis was disrupted in whole blood transcriptomes of active TB

(Figure 4C). However, ISG15 (but not *IL10* or *IFNG*) retained its association with

disease status and monocyte/myeloid markers (*CD14, FCGR1*). Since TB disease

signature in the whole blood is predominated by neutrophils (Berry et al., 2010) and

monocytes only make up a minor fraction in these samples, we next investigated

whether components of the *ISG15*/*IL10* signalling pathway might be overexpressed in

purified monocytes from TB patients, as compared to control monocytes. Indeed,

Ingenuity Pathway Analysis identified MAPK signalling as significantly enriched in

monocytes from TB patients (Supplementary Table), and p38 MAPK was

significantly interconnected with several TB signature genes (*FCGR1, IL27,*

SIGLEC6) in a disease network (Figure 4D). Together, these results show the

ISG15/IL10 axis is disrupted during active TB.

To further investigate the possible connection between ISG15 and *M. tuberculosis-*

mediated immunopathology, we examined the expression of this gene in an

independent large cohort in which both detailed clinical parameters and

corresponding transcriptome data are available (Berry et al., 2010). Expression levels

of *ISG15* were significantly correlated with established inflammatory biomarkers such

as erythrocyte sedimentation rate (ESR), C-reactive protein (CRP), tissue damage

(Modal X-ray grade) as well as systemic clinical parameters (neutrophil count,

haemoglobin, and globulin serum concentration) (Figure 4 G-L). These results

suggest ISG15 is a sensitive biomarker of disease severity in active TB patients.

ISG15 is critical for the production of IFNγ in cells from vaccine strain *BCG*-infected

patients (Bogunovic et al., 2012). Moreover, ISG15 has a synergistic effect when

combined with IL-12 (Bogunovic et al., 2012), an important inducer of IFNγ (Chan et

al., 1992; Chan et al., 1991). Furthermore, IL-10 inhibits production of IL-12 and,

consequently, of IFNγ by PBMCs (D'Andrea et al., 1993), pointing to a pleiotropic

effect for ISG15. Cell type and context dependent effects of ISG15 could explain

these diverse activities. This work and others (Bogunovic et al., 2012) suggest that

despite ubiquitous expression in different cell types, neutrophils are the major source

of secreted ISG15. Although we have not tested this directly, we speculate that

soluble neutrophil-derived ISG15 places neutrophils as pivotal to the orchestration of

immune responses *in vivo*, driving the production of at least two major cytokines, IL-

10 and IFNγ. Interestingly, the intra and extracellular location of ISG15 and its

ability to induce a plethora of effects in distinct cells, resembles the function of an

alarmin (Rider et al., 2017). As shown in figure 4, ISG15's function is context

dependent, varying from a driver of an anti-inflammatory monocytic/IL-10 axis in

homeostasis and the less severe *M. leprae* infection to a strong pro-inflammatory

IFNγ-biased scenario during active TB. Whether virulent *M. tuberculosis* hijacks the

ISG15/IL-10 axis contributing to induction of tissue pathology remains to be

determined.

In conclusion, these findings confirm and extend previous work characterizing soluble

extracellular ISG15 as a pleiotropic cytokine (or alarmin) that induces both pro- and

anti-inflammatory effects in a variety of cell types. Moreover, the combined *ex vivo*

and *in vitro* approach uncovers a novel myeloid *ISG15/IL10* p38-mediated anti-

inflammatory signalling cascade, which is preserved in human leprosy but disrupted

in active TB. Strikingly, our data indicate *ISG15* mRNA as a novel biomarker of

disease severity during acute TB that merits further investigation.

Figure Legends

FIGURE 1 – ISG15 induces the production of IL-10 in human PBMCs. (**A**) Dose-dependent IL-10 production as measured by ELISA 24 hours post stimulation of PBMCs with both pro- and mature ISG15 ([ISG15]: 0.15; 0.45; 1.5, 4.5 and 15 μg/mL). (**B**) Induction of IL-10 by recombinant, but not heat-treated, ISG15 using PBMCs from a total of 5 different donors in 8 independent experiments. (**C**) *IL10* mRNA expression in PBMCs treated with ISG15 at 6, 12, 24 and 48 hours post stimulation. (**D**) Quantification of IL-10 in the supernatant of human PBMCs at 6, 12, 24 and 48 hours after treatment with of ISG15 (**E**) Representative dot-plot evaluating cell-death in human PBMCs by annexin V and PI staining after treatment with ISG15 (2.0 μg/m) or Staurosporine (1 μM). (**F, G**) Quantification of cell death from the experiment described in (**E**). Unless stated otherwise, ISG15 concentration was 1.5 μg/mL. Error bars indicate SEM for biological replicates in each experiment. In each experiment PBMCs from 3 or more different donors were used. * P-value<0.05, ** p-value<0.01 and **** p-value<0.0001. ISG15HT, ISG15 heat-treated; STA, Staurosporine; PI, Propidium Iodide; Uns, Unstimulated.

FIGURE 2 – CD14$^+$ cells are the main source of IL-10 upon ISG15 stimulation. (**A**) Representative dot plot of intracellular staining of IL-10 in CD14$^+$, CD56$^+$, CD4$^+$ and CD8$^+$ in ISG15-treated PBMCs. (**B**) PBMCs from 6 different individuals showing fold increase in IL-10 production from CD14$^+$, CD56$^+$, CD4$^+$ and CD8$^+$ populations after ISG15 stimulation. (**C-D**) ELISA quantification of IL-10 and IFNγ in the supernatants of CD14$^+$ and CD14$^-$ separated populations treated with ISG15 (**E**) A549 WT or ISG15 KO cells were co-cultured with primary CD14$^+$ cells magnetically separated from PBMCs and IL-10 production was measured by ELISA

24 hours later. *ISG15* KO cells were also transfected with a plasmid expressing *ISG15*

in order to rescue its function. LPS was used as a positive control for IL-10

stimulation. Error bars indicate SEM for biological replicates in each experiment. All

experiments were repeated at least two times. In every experiment PBMCs from 3 or

more donors were used. (**F-J**) Transcriptome datasets of healthy controls (ImmuCo,

ImmuSort) confirm *ISG15* and *IL10 ex vivo* expression levels are strongly and

positively correlated in total PBMCs (**F**), purified primary monocytes (G) and

macrophages (**H**), but not neutrophils (**I**) or T cells (**J**). Red lines indicate the

approximate threshold for *IL10* mRNA detection (determined for each individual

microarray). (**J**) Neutrophils display the largest *ISG15*/*IL10* ratio *ex vivo*, whereas

monocytes are the major *IL10* expressing leukocyte population

(>PBMCs>neutrophils) under homeostatic conditions. **** p-value<0.0001.

FIGURE 3: ISG15 induces monocyte derived IL-10 via p38, MEK1/2 and PI3K

signalling pathways, which are deregulated in human mycobacterial infections. (**A**)

MAPK family members expression in both latent and active TB. *Ex vivo* expression

levels correlation of *MAPK14* and *ISG15* (**B**)*, MAPK3* and *IL-10* (**C**)*, MAPK3* and

IFNG (**D**) and of *PIK3CA* or *PIK3CB* with *IL-10* (**E**) during latent TB infection (**F**)

Representative immunoblot showing the phosphorylation of p38 MAPK 15 min after

the stimulation of ISG15 in CD14[+] cells. (**G-J**) CD14[+] cells were treated for 1h with

p38 (10 μM), MEK1/2 (50 μM) and PI3K inhibitors (50 μM) (B-E respectively) prior

to addition of ISG15 (1μg/mL) or LPS (100 ng/mL). (**K**) Chloroquine (5μg/mL) was

used as an unrelated control drug. 24 hours after treatment, supernatant was harvested

and used for IL-10 quantification by ELISA. Error bars indicate SEM for biological

replicates in two independent experiments. * P-value<0.05, ** p-value<0.01 and ***

p-value<0.001. Uns, Unstimulated.

FIGURE 4: An anti-inflammatory *ISG15/IL10* myeloid axis is amplified in human

leprosy and disrupted in human tuberculosis, revealing a novel clinical biomarker. (**A-**

C) Heatmaps representing positive (red) and negative (blue) correlation matrix of

selected genes (see text) classified by unsupervised hierarchical clustering (Eucledian

distance). (**A**) Healthy controls (GSE80008) (**B**) Leprosy patients (GSE82160) (**C**)

TB cohort (GSE85487) (**D**) Significantly enriched network (Ingenuity Pathway

Analysis) showing p38 MAPK as highly interconnected in the monocyte

transcriptome of TB patients. (**E-F**) *ISG15* transcript correlates with (**G, H**)

established inflammatory metrics (erythrocyte sedimentation rate (ESR), C-reactive

protein (CRP), (**I**) tissue damage (Modal X-ray grade) as well as (**J-L**) systemic

clinical parameters (neutrophil count, haemoglobin, and globulin serum

concentration).

Supplementary figure

(**A**) ISG15 induces the production inflammatory cytokines. PBMCs from healthy

donors were treated with ISG15 (15 µg/mL) and supernatant was harvested 24 hours

for inflammatory cytokines quantification by cytometric Bead Array (CBA). ISG15

induced the production of IL-10, IL-6 and IL-1β. (**B**) Generation of ISG15 deficient

A549 cell line. ISG15 deficient A549 cell line was produced using CRISPR/Cas9.

After clone selection, cells were stimulated with IFNβ (1000 IU/mL), proteins

extracted after 24 hours and immunoblotted with anti-ISG15 and anti-tubulin

antibodies. Clone 3 (ISG15 KO) was used for further experiments.

Supplementary Table

Tab 1. PI3K gene expression enriched during latent tuberculosis.

Tab 2. MAPK gene expression enriched during latent tuberculosis.

Tab 3. Ingenuity Pathway Analysis for MAPK in monocytes during latent

tuberculosis.

Tab 4. p38 network in monocytes during active tuberculosis.

Material and methods

Reagents

ISG15 was purchased from Boston Biochem and tested for endotoxins by R&D

Systems (endotoxin value for lot #DBHF0614021 is <0.00394 EU/μg). LAL assay

(Lonza) was performed according to the manufacturer's instruction and the endotoxin

level of recombinant ISG15 was below the detection threshold. Pro-ISG15 (UL-615)

was also purchased from Boston Biochem. *E. coli* LPS (strain O111:B4) (Invivogen)

was used as a positive control for IL-10 production in human PBMCs and monocytes.

P38 kinase inhibitors SB203580 and SB220025 (Calbiochem) were used at 10 μM,

MEK1/2 inhibitor U0126 (Cell Signaling) at 50 μM and PI3K inhibitor Ly294002

(Cell Signaling) was at 50 μM. Solvent (DMSO, medium) was used as negative

control and chloroquine (Sigma, 5μg/mL), a DNA-PKC/TLR endosomal signalling

inhibitor, was used as an additional negative control.

Primary human cells

Human PBMCs were separated from healthy individuals using Ficoll-paque (GE)

according to manufacturer's instructions. Briefly, blood was collected in heparin-

containing tubes, and gently mixed 1:1 with saline solution and gently mixed before

being added over one volume of Ficoll-paque reagent. The gradient was centrifuged

for 35 minutes at 400 x g, 18˚C. PBMCs were harvested and washed once with 45 mL

of saline solution for 10 min at 400 x g, 18˚C. Subsequently, cell pellet was

suspended and washed twice with 5 mL of saline solution for 10 min at 200 x g, 18˚C

to remove platelets. The remaining cell pellet was suspended to the desired density in

RPMI 1640 (Gibco) supplemented with 5% foetal calf serum (Hyclone), 2mM L-

glutamine (Gibco), 1 mM sodium pyruvate (Gibco), 25 mM HEPES (Gibco), 100

U/mL penicillin and 100 μg/mL streptomycin (Gibco). Cells were plated as described

in each experiment. Human primary monocytes (CD14[+] cells) were separated from

PBMCs using CD14 microbeads (Miltenyi Biotec) according to manufacturer's

instructions with the exception of the MACS buffer, which was prepared using 3%

foetal calf serum. Monocyte enrichment varied between 73 to 92% between

experiments. The use of PBMCs from healthy donors was previously approved by

UFSC ethical committee (IRB#283/08).

Generation of isg15 knockout cell lines

A549 lung epithelial cells were co-transfected with three gRNA/Cas9/GFP plasmids

(provided by Horizon) targeting the *ISG15* locus using JetPEI (PolyPlus

Transfection). The guide RNAs used were 5' GGCTGTGGGCTGTGGGCTGT 3', 5'

GGTAAGGCAGATGTCACAGG 3' and 5' TGGAGCTGCTCAGGGACACC 3'.72

hours after transfection, cells sorted for GFP fluorescence and then separated by

limiting dilution. Single-cell derived clones were selected for ISG15 expression

(Supplementary Figure 1B).

A549 – CD14+ co-culture

A549 WT or ISG15 KO cells were seeded at 2 x 10^5 cells/ml in 24 well-plates. Cells

rested in the incubator for 6 hours before ISG15 KO cells were transfected with

ISG15-pCEP4 plasmid using FugeneHD reagent (Promega) according to

manufacturer's instructions. Cells were then washed and LPS was added 18 hours

after transfection and immediately prior to the addition of a 2 x 10^5 CD14$^+$ cells

overlay. Following 24 hours of co-culture, supernatants were harvested for IL-10

quantification.

Immunoblotting

1 x 10^5 CD14$^+$ cells were added to a 96-well plate and left to rest overnight. Cells

were stimulated with ISG15 (1μg/ml) and after 15 minutes cells were spun at 4˚C,

supernatant was removed and M-PER lysis buffer (Thermo Scientific) containing

protease inhibitors (Complete, Mini Protease Inhibitor Tablets, Roche) and

phosphatase inhibitors (#524625, Calbiochem) was added to the cells. Protein

separation was performed according to M-PER manufacturer's instructions.

Antibodies concentrations for detection of p38 (Cell Signaling #9212) and p-p38 (Cell

Signaling #9211), ISG15 (Cat: A600, R&D Systems) and anti-α-Tubulin (clone

DM1A, Millipore) were those suggested by the manufacturers. For Western blots, at

least 20 μg of total protein were separated and transferred to a PVDF 0.22 μm

blotting membrane. Membrane was blocked for at least 1 hour with 1X Tris Buffered

Saline-0.1% Tween20 (TBST) with 5% w/v non-fat dry milk and subsequently

washed 3 times with TBST for 5 minutes each wash. Membrane was incubated with

primary antibodies diluted in 5% w/v BSA, 1X TBS, 0.1% Tween20 at 4˚C with

gentle shaking overnight. Membrane was washed 3 times for 5 min each with TBST

and then incubated with the appropriate secondary HRP-linked antibody for 1 hour at

room temperature. Membrane was washed 3 times of 5 min each with TBST before

detection with ECL chemiluminescent substrate (Pierce).

p38 MAPK and PI3K signalling pathway inhibition

1×10^5 CD14$^+$ cells were added to a 96-well plate and left to rest overnight. Inhibitors

were added to cells for 1 hour prior to ISG15 treatment. 24 hours after treatment, cells

were spun at 4°C; supernatant was harvested and IL-10 was quantified by ELISA.

Cytokine quantification

For exploratory experiments, IL-1β, IL-6, IL-10, IL-12p70 and TNF were quantified

in supernatants by human inflammatory cytokine cytometric beads array kit (CBA,

BD Biosciences). IL-10 and IFNγ were quantified using Human IL-10 DuoSet ELISA

kit (R&D Systems) or Human IFNγ mini kit (Thermo Scientific) according to

manufacturer's instructions.

Flow Cytometry Assays

PBMCs were seeded at a density of 5×10^5 cells per well in 150 μL of medium. After

8 hours of resting at 37°C with 5% CO_2, cells were treated with ISG15 (2 μg/mL) or

LPS (1 μg/mL), unless indicated otherwise. Golgi Plug protein transport inhibitor

(BD Biosciences) was added 1-hour post treatment, according to manufacturer's

instructions. Then, 12 hours post treatment; growth medium was removed and cold

1X HBSS (Gibco) with 2.5 mM EDTA was added to each well. The tissue culture

dish was kept at 4°C for 30 minutes and cells were suspended and transferred to 1.5

mL tubes. Cells were washed in a final volume of 1mL cold 1X HBSS (Gibco) with

2.5 mM EDTA at 300 x g and 4°C for 5 minutes. Supernatant was removed and cells

were suspended in FACS buffer (1% BSA, 1% sodium azide in 1X PBS). Anti-human

antibody mix containing anti-CD4 APC-Cy7 (clone OKT1) (BioLegend), anti-CD8

PE-Cy7 (clone SK1) (BioLegend), anti-CD14 PerCP-Cy5.5 (clone M5E2)

(BioLegend), anti-CD56 FITC (clone NCAM 16.2) (BD Biosciences) was added to

the cell suspension for 40 min at 4ºC in the presence of 10% AB blood-type human

serum to block Fc receptors. Afterwards, cells were washed once with 1 mL of 1X

PBS at 300 x g, 4°C, for 5 minutes and 1 mL of fixation buffer (1% paraformaldehyde

in 1X PBS) was added to the cells. Tubes were kept in the dark at room temperature

for 15 min and then centrifuged at 300 x g, 4°C, for 10 min to remove supernatant. 1

mL of permeabilization buffer (0.5% saponin in FACS buffer) was added to the cells

and tubes were centrifuged at 300 x g, 4 C for 10 min. Intracellular stain with anti-IL-

10-PE (clone JES3-9D7) (BioLegend) was carried out for 30 min in the dark at room

temperature. Cells were washed with permeabilization buffer at 300 x g, 4°C, 5 min,

supernatant was removed and cells were suspended in FACS buffer prior to

acquisition of 1×10^5 events or more. For analysis, all acquired events displayed as

forward scatter (FSC) and side scatter (SSC) parameters were selected. After that,

single cells events were selected using FSC area and height parameters (FSC-A x

FCS-H) and auto-fluorescence was excluded using APC as an open channel.

Intracellular IL-10 was then quantified in monocytes ($CD14^+IL\text{-}10^+$), NK cells

($CD56^+IL\text{-}10^+$), CD4 ($CD4^{high}IL\text{-}10^+$) and CD8 T cells ($CD8^{high}IL\text{-}10^+$). Gates were

set according to unstained PBMC sample and controls. All samples were acquired on

a Becton-Dickinson Verse flow cytometer using BD FACSuite^{TM} software. In order

to analyse cell death, 5×10^5 PBMC/well were treated with ISG15 2 µg/mL or

Staurosporine (Sigma) 1 µM for 24 hours. Cells were then harvest, washed with 1 mL

of PBS at 300 x g, room temperature, 5 min, supernatant was removed and cells were

washed once in 1 mL of 1x Annexin biding buffer (eBioscience). Cell were

resuspended at 10^6 cells/mL in 1x Annexin biding buffer and FITC conjugated

Annexin V (eBioscience) was added to the cell suspension for 15 min, room

temperature, according to manufacture's instruction. Following incubation period,

cells were washed with 1 mL of 1x Annexin biding buffer, 300 x g, room

temperature, 5 min and resuspended in 200 uL of 1x annexin biding buffer. Propidium

iodide (BD Pharmingen) was added at 0.25 μg/mL to the cell suspension prior to

sample acquisition. Samples were acquired on a Becton-Dickinson Canto II flow

cytometer using BD FACSDiva^{TM} software.

Real time quantitative PCR (qPCR)

For relative quantification of *IL10* gene expression, total RNA was extracted from

PBMCs treated or not with ISG15. RNA was extracted after 6, 12, 24 or 48 hours of

treatment using RNeasy RNA extraction kit (Qiagen). Using 400 ng of RNA, cDNA

was produced with High-Capacity cDNA Reverse Transcription Kit (Applied

Biosystems) and 2μL of the product was used for the qPCR reaction in a final volume

of 10 μL. qPCR reactions were performed using the primers forward 5'GAG ATC

TCC GAG ATG CCT TCA G 3'and reverse 5'CAA GGA CTC CTT TAA CAA

CAA GTT GT 3' (Skrzeczynska-Moncznik et al., 2008). Fold-increase in *IL10* gene

expression was determined by relative quantification using hypoxanthine

phosphoribosyltransferase (*HPRT*) as endogenous control. Primers forward and

reverse for *HPRT* were 5' CCTGCTGGATTACATCAAAGCACTG 3' and 5'

TCCAACACTTCGTGGGGTCCT 3', respectively, and were used at 250 nM each.

Microarray analysis

Curated and annotated publicly available data-sets (Berry et al., 2010; Novais et al.,

2015; Speake et al., 2015; Teles et al., 2013; Wang et al., 2015a; Wang et al.,

2015b)(GXB, ImmuCo, ImmuSort, BioGPS, GEO) were obtained from large,

established cohorts of healthy controls, latent and active tuberculosis patients,

comprising *ex vivo* and *in vitro* whole blood, total PBMCs, purified leukocyte subsets,

non-leukocyte human primary cells and skin biopsies (leprosy patients, healthy

controls and cutaneous leishmaniasis as non-mycobacterial infectious control). Novel

datasets were generated for both whole blood and PBMCs from healthy controls and

individuals infected with other non-mycobacterial intracellular pathogens

(*Leishmania*, HIV-1, HTLV-1). PBMCs were isolated as above and immediately

frozen in Trizol to preserve RNA integrity. Following Trizol extraction, total RNA

was further purified using an RNeasy kit according to the manufacturer's protocol

(QIAGEN, Venlo, Netherlands). Affymetrix Whole Genome microarray analysis was

performed by the VIB Nucleomics Facility (Leuven, Belgium) using a GeneChip$^{®}$

Human Gene 1.0 ST Array with the WT PLUS reagent kit (Affymetrix, Santa Clara,

CA, USA) according to the manufacturer's specifications. Data preprocessing (RMA)

was performed using the Bioconductor xps package. All microarray raw data are

available   at   Gene   Expression   Omnibus   database   (GEO,

http://www.ncbi.nlm.nih.gov/geo/) under series accession numbers GSE80008,

GSE82160, GSE85487.

Enrichment analysis

The Ingenuity Pathway Analysis (IPA) program was used to perform the initial

pathway/function level analysis on genes determined to be differentially expressed in

the microarray analysis (Ingenuity Systems, Red Wood City, CA). Uncorrected p-

values and absolute fold-changes were used with cut-offs of p<0.05. Based on a

scientific literature database, the genes were sorted into gene networks and canonical

pathways, and significantly overrepresented pathways were identified. Further

enrichment analysis was performed, including Gene Ontology (GO) term enrichment

using the WEB-based GEne SeT AnaLysis Toolkit (WebGestalt), KEGG pathway

enrichment using the pathway database from the Kyoto Encyclopedia of Genes and

Genomes and transcription factor target enrichment using data from the Broad

Institute Molecular Signatures Database (MSigDB). Genesets from the GO, KEGG

pathways, WikiPathways and Pathway Commons databases, as well as transcription

factors, were considered overrepresented if their corrected p-value was smaller than

0.05. Principal component analysis, correlation matrices (Spearman), unsupervised

hierarchical (Eucledian distance) clustering were performed using XLSTAT and

visualized using MORPHEUS (https://software.broadinstitute.org/morpheus/).

Data processing and statistical analyses

Data derived from *in vitro* experiments was processed using Graphpad Prism 6 and

analysed using unpaired Student's T test unless stated otherwise. Statistical

significance is expressed as follows: * P-value<0.05, ** p-value<0.01, *** p-

value<0.001 and **** p-value<0.0001. In all cases, data shown are representative

from at least two independent experiments. Data from experiments performed in

triplicate are expressed as mean ± SEM.

Acknowledgements

DSM received support from CAPES Computational Biology (23038.010048/2013-

27), CNPq Universal (473897/2013-0) and the Academy of Medical Sciences/UK

(NAF004/1005). PFS and MD received CAPES and CNPq student fellowships

respectively. AB received financial support from NIH-GRIP (TW008276), HHMI-

ECS (55007412). AB is CNPq-PQ scholar and CAPES/ESE. BJF received support

from an Isaac Newton Trust/Wellcome Trust ISSF/University of Cambridge research

grant and a Wellcome Trust Seed Award (201946/Z/16/Z). TD received grant support

from VLAIO (IWT141614). JVW received grant support from CAPES (PVE) and

FWO (G0D6817N).

We would like to thank Prof. Aristobolo Mendes Silva from UFMG for providing

reagents, suggestions and critical evaluation of these results.

The authors declare no conflict of interest.

Author contributions

547 PFS, JVW and RG designed, performed experiments, analysed the data and wrote the

548 manuscript; MD, DOP, TD and SMM performed experiments and analysed the data,

549 BF, AB and DSM designed experiments analysed the data and wrote the manuscript.

References

Berry, M.P., C.M. Graham, F.W. McNab, Z. Xu, S.A. Bloch, T. Oni, K.A. Wilkinson, R.
Banchereau, J. Skinner, R.J. Wilkinson, C. Quinn, D. Blankenship, R.
Dhawan, J.J. Cush, A. Mejias, O. Ramilo, O.M. Kon, V. Pascual, J. Banchereau,
D. Chaussabel, and A. O'Garra. 2010. An interferon-inducible neutrophil-
driven blood transcriptional signature in human tuberculosis. *Nature*
466:973-977.
Billiau, A. 2006. Anti-inflammatory properties of Type I interferons. *Antiviral*
research 71:108-116.
Bogunovic, D., S. Boisson-Dupuis, and J.L. Casanova. 2013. ISG15: leading a
double life as a secreted molecule. *Exp Mol Med* 45:e18.
Bogunovic, D., M. Byun, L.A. Durfee, A. Abhyankar, O. Sanal, D. Mansouri, S. Salem,
I. Radovanovic, A.V. Grant, P. Adimi, N. Mansouri, S. Okada, V.L. Bryant,
X.F. Kong, A. Kreins, M.M. Velez, B. Boisson, S. Khalilzadeh, U. Ozcelik, I.A.
Darazam, J.W. Schoggins, C.M. Rice, S. Al-Muhsen, M. Behr, G. Vogt, A. Puel,
J. Bustamante, P. Gros, J.M. Huibregtse, L. Abel, S. Boisson-Dupuis, and J.L.
Casanova. 2012. Mycobacterial disease and impaired IFN-gamma

immunity in humans with inherited ISG15 deficiency. *Science* 337:1684-
1688.
Borden, E.C., G.C. Sen, G. Uze, R.H. Silverman, R.M. Ransohoff, G.R. Foster, and G.R.
Stark. 2007. Interferons at age 50: past, current and future impact on
biomedicine. *Nat Rev Drug Discov* 6:975-990.
Chan, S.H., M. Kobayashi, D. Santoli, B. Perussia, and G. Trinchieri. 1992.
Mechanisms of IFN-gamma induction by natural killer cell stimulatory
factor (NKSF/IL-12). Role of transcription and mRNA stability in the
synergistic interaction between NKSF and IL-2. *Journal of immunology*
148:92-98.
Chan, S.H., B. Perussia, J.W. Gupta, M. Kobayashi, M. Pospisil, H.A. Young, S.F.
Wolf, D. Young, S.C. Clark, and G. Trinchieri. 1991. Induction of interferon
gamma production by natural killer cell stimulatory factor:
characterization of the responder cells and synergy with other inducers.
The Journal of experimental medicine 173:869-879.
D'Cunha, J., E. Knight, Jr., A.L. Haas, R.L. Truitt, and E.C. Borden. 1996.
Immunoregulatory properties of ISG15, an interferon-induced cytokine.
Proceedings of the National Academy of Sciences of the United States of
America 93:211-215.
de Veer, M.J., M. Holko, M. Frevel, E. Walker, S. Der, J.M. Paranjape, R.H.
Silverman, and B.R. Williams. 2001. Functional classification of interferon-
stimulated genes identified using microarrays. *Journal of leukocyte biology*
69:912-920.
Der, S.D., A. Zhou, B.R. Williams, and R.H. Silverman. 1998. Identification of genes
differentially regulated by interferon alpha, beta, or gamma using
oligonucleotide arrays. *Proceedings of the National Academy of Sciences of*
the United States of America 95:15623-15628.
Dos Santos, P.F., and D.S. Mansur. 2017. Beyond ISGlylation: Functions of Free
Intracellular and Extracellular ISG15. *Journal of interferon & cytokine*
research : the official journal of the International Society for Interferon and
Cytokine Research
Haas, A.L., P. Ahrens, P.M. Bright, and H. Ankel. 1987. Interferon induces a 15-
kilodalton protein exhibiting marked homology to ubiquitin. *The Journal*
of biological chemistry 262:11315-11323.
Hansen, T.R., and J.K. Pru. 2014. ISGylation: a conserved pathway in mammalian
pregnancy. *Adv Exp Med Biol* 759:13-31.
Henkes, L.E., J.K. Pru, R.L. Ashley, R.V. Anthony, D.N. Veeramachaneni, K.C. Gates,
and T.R. Hansen. 2015. Embryo mortality in Isg15-/- mice is exacerbated
by environmental stress. *Biol Reprod* 92:36.
Hermann, M., and D. Bogunovic. 2017. ISG15: In Sickness and in Health. *Trends*
Immunol 38:79-93.
Knight, E., Jr., D. Fahey, B. Cordova, M. Hillman, R. Kutny, N. Reich, and D.
Blomstrom. 1988. A 15-kDa interferon-induced protein is derived by
COOH-terminal processing of a 17-kDa precursor. *The Journal of biological*
chemistry 263:4520-4522.
Loeb, K.R., and A.L. Haas. 1992. The interferon-inducible 15-kDa ubiquitin
homolog conjugates to intracellular proteins. *The Journal of biological*
chemistry 267:7806-7813.

Ma, W., W. Lim, K. Gee, S. Aucoin, D. Nandan, M. Kozlowski, F. Diaz-Mitoma, and A.
Kumar. 2001. The p38 mitogen-activated kinase pathway regulates the
human interleukin-10 promoter via the activation of Sp1 transcription
factor in lipopolysaccharide-stimulated human macrophages. *The Journal*
of biological chemistry 276:13664-13674.
McNab, F., K. Mayer-Barber, A. Sher, A. Wack, and A. O'Garra. 2015. Type I
interferons in infectious disease. *Nature reviews. Immunology* 15:87-103.
Morales, D.J., and D.J. Lenschow. 2013. The antiviral activities of ISG15. *Journal of*
molecular biology 425:4995-5008.
Nair, S., P.A. Ramaswamy, S. Ghosh, D.C. Joshi, N. Pathak, I. Siddiqui, P. Sharma,
S.E. Hasnain, S.C. Mande, and S. Mukhopadhyay. 2009. The PPE18 of
Mycobacterium tuberculosis interacts with TLR2 and activates IL-10
induction in macrophage. *Journal of immunology* 183:6269-6281.
Narasimhan, J., J.L. Potter, and A.L. Haas. 1996. Conjugation of the 15-kDa
interferon-induced ubiquitin homolog is distinct from that of ubiquitin.
The Journal of biological chemistry 271:324-330.
Novais, F.O., L.P. Carvalho, S. Passos, D.S. Roos, E.M. Carvalho, P. Scott, and D.P.
Beiting. 2015. Genomic profiling of human Leishmania braziliensis lesions
identifies transcriptional modules associated with cutaneous
immunopathology. *The Journal of investigative dermatology* 135:94-101.
Owhashi, M., Y. Taoka, K. Ishii, S. Nakazawa, H. Uemura, and H. Kambara. 2003.
Identification of a ubiquitin family protein as a novel neutrophil
chemotactic factor. *Biochemical and biophysical research communications*
309:533-539.
Potter, J.L., J. Narasimhan, L. Mende-Mueller, and A.L. Haas. 1999. Precursor
processing of pro-ISG15/UCRP, an interferon-beta-induced ubiquitin-like
protein. *The Journal of biological chemistry* 274:25061-25068.
Recht, M., E.C. Borden, and E. Knight, Jr. 1991. A human 15-kDa IFN-induced
protein induces the secretion of IFN-gamma. *Journal of immunology*
147:2617-2623.
Rider, P., E. Voronov, C.A. Dinarello, R.N. Apte, and I. Cohen. 2017. Alarmins: Feel
the Stress. *Journal of immunology* 198:1395-1402.
Schoggins, J.W., and C.M. Rice. 2011. Interferon-stimulated genes and their
antiviral effector functions. *Current opinion in virology* 1:519-525.
Skaug, B., and Z.J. Chen. 2010. Emerging role of ISG15 in antiviral immunity. *Cell*
143:187-190.
Skrzeczynska-Moncznik, J., M. Bzowska, S. Loseke, E. Grage-Griebenow, M.
Zembala, and J. Pryjma. 2008. Peripheral blood CD14high CD16+
monocytes are main producers of IL-10. *Scandinavian journal of*
immunology 67:152-159.
Speake, C., S. Presnell, K. Domico, B. Zeitner, A. Bjork, D. Anderson, M.J. Mason, E.
Whalen, O. Vargas, D. Popov, D. Rinchai, N. Jourde-Chiche, L. Chiche, C.
Quinn, and D. Chaussabel. 2015. An interactive web application for the
dissemination of human systems immunology data. *Journal of*
translational medicine 13:196.
Tamassia, N., M. Zimmermann, M. Castellucci, R. Ostuni, K. Bruderek, B. Schilling,
S. Brandau, F. Bazzoni, G. Natoli, and M.A. Cassatella. 2013. Cutting edge:
An inactive chromatin configuration at the IL-10 locus in human
neutrophils. *Journal of immunology* 190:1921-1925.

Tecalco Cruz, A.C., and K. Mejia-Barreto. 2017. Cell type-dependent regulation of
free ISG15 levels and ISGylation. *J Cell Commun Signal*
Teles, R.M., T.G. Graeber, S.R. Krutzik, D. Montoya, M. Schenk, D.J. Lee, E.
Komisopoulou, K. Kelly-Scumpia, R. Chun, S.S. Iyer, E.N. Sarno, T.H. Rea, M.
Hewison, J.S. Adams, S.J. Popper, D.A. Relman, S. Stenger, B.R. Bloom, G.
Cheng, and R.L. Modlin. 2013. Type I interferon suppresses type II
interferon-triggered human anti-mycobacterial responses. *Science*
339:1448-1453.
Wang, B.X., S.A. Grover, P. Kannu, G. Yoon, R.M. Laxer, E.A. Yeh, and E.N. Fish.
2017a. Interferon-Stimulated Gene Expression as a Preferred Biomarker
for Disease Activity in Aicardi-Goutieres Syndrome. *Journal of interferon &*
cytokine research : the official journal of the International Society for
Interferon and Cytokine Research 37:147-152.
Wang, P., H. Qi, S. Song, S. Li, N. Huang, W. Han, and D. Ma. 2015a. ImmuCo: a
database of gene co-expression in immune cells. *Nucleic acids research*
43:D1133-1139.
Wang, P., Y. Yang, W. Han, and D. Ma. 2015b. ImmuSort, a database on gene
plasticity and electronic sorting for immune cells. *Scientific reports*
5:10370.
Wang, W., Y. Yin, L. Xu, J. Su, F. Huang, Y. Wang, P.P.C. Boor, K. Chen, W. Wang, W.
Cao, X. Zhou, P. Liu, L.J.W. van der Laan, J. Kwekkeboom, M.P.
Peppelenbosch, and Q. Pan. 2017b. Unphosphorylated ISGF3 drives
constitutive expression of interferon-stimulated genes to protect against
viral infections. *Science signaling* 10:
Zhang, X., D. Bogunovic, B. Payelle-Brogard, V. Francois-Newton, S.D. Speer, C.
Yuan, S. Volpi, Z. Li, O. Sanal, D. Mansouri, I. Tezcan, G.I. Rice, C. Chen, N.
Mansouri, S.A. Mahdaviani, Y. Itan, B. Boisson, S. Okada, L. Zeng, X. Wang,
H. Jiang, W. Liu, T. Han, D. Liu, T. Ma, B. Wang, M. Liu, J.Y. Liu, Q.K. Wang, D.
Yalnizoglu, L. Radoshevich, G. Uze, P. Gros, F. Rozenberg, S.Y. Zhang, E.
Jouanguy, J. Bustamante, A. Garcia-Sastre, L. Abel, P. Lebon, L.D.
Notarangelo, Y.J. Crow, S. Boisson-Dupuis, J.L. Casanova, and S. Pellegrini.
2015. Human intracellular ISG15 prevents interferon-alpha/beta over-
amplification and auto-inflammation. *Nature* 517:89-93.
698
699
700