RESEARCH ARTICLE

Dynamic clustering of dynamin-amphiphysin rings regulates membrane constriction and

fission coupled with GTP hydrolysis

Tetsuya Takeda[1*], Toshiya Kozai[2], Huiran Yang[1], Daiki Ishikuro[2], Kaho Seyama[1], Yusuke

Kumagai[2], Tadashi Abe[1], Hiroshi Yamada[1], Takayuki Uchihashi[4], Toshio Ando[3*], Kohji Takei[1*]

[1]Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, Okayama University,

Okayama, Japan; [2]Department of Physics, College of Science and Engineering, Kanazawa

University, Kanazawa, Japan; [3]Bio-AFM Frontier Research Center, Institute of Science and

Engineering, Kanazawa University, Kanazawa, Japan; [4]Department of Physics, School of

Science, Nagoya University

*For correspondence: ttakeda@okayama-u.ac.jp (T.T.), tando@staff.kanazawa-u.ac.jp (T.A.)

and kohji@md.okayama-u.ac.jp (K.T.).

ABSTRACT

Dynamin is a mechanochemical GTPase essential for membrane fission during clathrin

mediated endocytosis. Dynamin forms washer ring-shaped/helical complexes at the neck of

clathrin-coated pits and their structural changes coupled with GTP hydrolysis drive membrane

fission. Dynamin and its binding protein amphiphysin cooperatively regulates membrane

remodeling during fission, but its precise mechanism remains elusive. In this study, we analyze

structural changes of dynamin-amphiphysin complexes during membrane fission using electron

microscopy (EM) and high-speed atomic force microscopy (HS-AFM). Interestingly, HS-AFM

analyses show that the dynamin-amphiphysin rings are rearranged to form clusters upon GTP

hydrolysis and membrane constriction occurs at protein-uncoated regions flanking the clusters.

We also show a novel function of amphiphysin in size control of the clusters to enhance

biogenesis of endocytic vesicles. Our new approaches using combination of EM and HS-AFM

clearly demonstrates dynamics of dynamin-amphiphysin complexes during membrane fission

suggesting a novel "clusterase" model of dynamin-mediated membrane fission.

INTRODUCTION

Clathrin mediated endocytosis (CME) is the best characterized endocytic pathway by which

cells incorporate extracellular molecules into cells with the aid of clathrin coat (Kirchhausen,

Owen, & Harrison, 2014; McMahon & Boucrot, 2011). CME is required for various essential

processes including neuronal transmission, signal transduction and other cell membrane activities

such as cell adhesion and migration. For precise progression of membrane invagination and

fission during CME, various proteins need to be assembled in a temporally and spatially

coordinated manner at the site of endocytosis.

One of those endocytic proteins, dynamin, is a GTPase essential for membrane fission in CME

(Antonny et al., 2016; Ferguson & De Camilli, 2012; Schmid & Frolov, 2011). There are three

dynamin isoforms in mammals: dynamin 1 and dynamin 3, two tissue specific isoforms which are

highly expressed in neurons, and dynamin 2, a ubiquitously expressed isoform (Cao, Garcia, &

McNiven, 1998; T. Cook, Mesa, & Urrutia, 1996; T. A. Cook, Urrutia, & McNiven, 1994).

Structural studies from several groups demonstrated that dynamin consists of five structurally

distinct domains: a GTPase domain, a bundle signaling element (BSE), a stalk, a pleckstrin

homology (PH) domain and proline-rich domain (PRD) from N-terminus to C-terminus (Faelber

et al., 2011; Ford, Jenni, & Nunnari, 2011; Reubold et al., 2015). The GTPase domain is

responsible for hydrolysis of GTP (guanosine triphosphate) and PH domain is required for

membrane association by binding to negatively charged phospholipids such as PI (4,5) P_2

(phosphatidylinositol 4,5-bisphosphate). The stalk structure serves as a binding interface for

dimerization and oligomerization of dynamin, and BSE, which is located between the stalk and

GTPase domain, functions as a flexible hinge required for structural changes of dynamin upon

GTP hydrolysis. Dynamin forms washer ring-like or helical oligomers which were first observed

in presynaptic terminals of *shibire* mutant flies at restrictive temperature (Koenig & Ikeda, 1989).

Dynamin also assembles into helices at the neck of endocytic pits in the isolated presynaptic

nerve terminals treated with slowly-hydrolyzable GTP analogue GTPγS (guanosine

5'-O-[gamma-thio]triphosphate) (Takei, McPherson, Schmid, & De Camilli, 1995). Similar

dynamin rings/helix were reconstituted *in vitro* either with liposomes (Sweitzer & Hinshaw,

1998; Takei et al., 1998) or without liposomes in a low salt condition (Hinshaw & Schmid, 1995).

There is a consensus view about the dynamin-mediated membrane constriction and fission

which is well supported by previous studies from different groups: membrane constriction is

required, but not sufficient, for fission (Antonny et al., 2016; Faelber et al., 2012; Schmid &

Frolov, 2011). However, it is still controversial how constriction is achieved, and what GTP

energy is used for. For example, membrane constriction could be achieved by assembly into the

highly-constricted state when dynamin is bound to GTP (Chen, Zhang, Egelman, & Hinshaw,

2004; Mattila et al., 2015; Mears, Ray, & Hinshaw, 2007; Zhang & Hinshaw, 2001).

Alternatively, membrane constriction could be achieved by hydrolysis of GTP that induces a

conformational change leading to constriction (Cocucci, Gaudin, & Kirchhausen, 2014; Marks et

al., 2001; Roux, Uyhazi, Frost, & De Camilli, 2006). However, precise mechanisms involved in

dynamin-mediated membrane constriction and fission remain unclear.

Amphiphysin is a BAR domain protein required for membrane invagination in CME (Wigge

et al., 1997). Amphiphysin has a lipid interacting BAR (Bin–AMPH–Rvs) domain in its

N-terminal, a medial clathrin/AP-2 binding (CLAP) domain and C-terminal Src homology 3

(SH3) domain. The BAR domain of amphiphysin forms crescent-shaped dimer and its concave

surface serves as a platform for bending membrane or sensing membrane curvature (Peter et al.,

2004). The CLAP domain binds to clathrin and AP-2, major components of clathrin coated pits,

and helps to recruit amphiphysin to the sites of CME. In addition, the C-terminal SH3domain of

amphiphysin binds directly to the PRD of dynamin 1 (David, McPherson, Mundigl, & de Camilli,

1996; Takei, Slepnev, Haucke, & De Camilli, 1999) and enhances dynamin's GTPase activity in

the presence of liposomes (Takei et al., 1999; Yoshida et al., 2004). Amphiphysin copolymerizes

with dynamin 1 into washer ring-shaped complexes, which form membrane tubules *in vitro*

(Takei et al., 1999; Yoshida et al., 2004) similar to those formed from synaptic plasma

membranes (Takei et al., 1995). Furthermore, injection of specific antibodies against

amphiphysin into the giant synapse in lampreys (Evergren et al., 2004) or amphiphysin KO in

mice (Di Paolo et al., 2002) causes suppressed endocytosis in synaptic vesicle recycling. These

results suggest that dynamin mediates membrane fission in CME in collaboration with

amphiphysin *in vivo*. However, precise contribution of amphiphysin in the dynamin-mediated

membrane fission remains elusive.

In this study, we analyze dynamics of dynamin-amphiphysin ring complexes using a novel

approach combining EM and HS-AFM. Firstly, we show that the dynamin-amphiphysin rings

are rearranged to form clusters upon GTP hydrolysis, and membrane constriction occurs at

protein-uncoated regions between the clusters. Secondly, we reveal that GTP hydrolysis is

required and sufficient for the cluster formation by dynamin-amphiphysin complexes by EM

analyses. Finally, we show a novel function of amphiphysin in controlling cluster size, which in

turn regulates biogenesis of endocytic vesicles. These findings provide new insights into the

mechanism of membrane constriction and fission by dynamin-amphiphysin complexes.

RESULTS

GTP hydrolysis is required and sufficient for membrane constriction by dynamin-amphiphysin

complexes

To elucidate the mechanisms of dynamin-mediated membrane fission, we reconstituted the

minimum system *in vitro* and analyzed the time course of its structural changes using EM.

Human dynamin 1 and amphiphysin were purified (Supplementary Figure S1A) and their activity

to form ring-shaped complexes in a buffer of physiological ionic strength and pH condition was

confirmed (Supplementary Figure S1B). As previously described (Sweitzer & Hinshaw, 1998;

Takei et al., 1998; Takei et al., 1999), the dynamin-amphiphysin complexes induced tubulation

of large unilamellar vesicles (LUVs) in the absence of GTP (Figure 1A, No GTP). Immediately

after the addition of 1 mM GTP, the appearance of lipid tubules was not affected (Figure 1A, GTP

1 s), but they started to form multiple constriction sites over time (Figure 1A, GTP 5 s, 10 s and 30

s) and membrane fission occurred finally and numerous vesicles were generated within 1 min

(Figure 1A, GTP 1 min).

Next, we tried to clarify how the membrane constriction and fission by dynamin-amphiphysin

ring complexes are correlated with guanine nucleotide conditions during GTP hydrolysis. The

appearance of lipid tubules (Figure 1B, No GTP) was not affected in the presence of either

slowly-hydrolyzable GTP analogue GTPγS (Figure 1B, GTPγS) or non-hydrolyzable GTP

analogue GMP-PNP (guanosine 5′-[β,γ-imido]triphosphate) (Figure 1B, GMP-PNP). In contrast,

in the presence of GDP (guanosine diphosphate) and vanadate, the complex which mimics the

GDP·Pi transition state, lipid tubules were constricted at multiple sites (Figure 1B, GDP +

vanadate). Addition of only GDP did not cause membrane constriction or fission, but membrane

tubules were deformed (Figure 1B, GDP). Finally, numerous vesicles were generated 10 min after

the addition of 1 mM GTP, in which multiple rounds of GTP hydrolysis were likely to have taken

place (Figure 1B, GTP 10 min). Taken these results together, GTP hydrolysis is essential for both

membrane constriction and fission by the dynamin-amphiphysin complexes, but subsequent

dissociation of GTP hydrolytic products (GDP and/or phosphate) is required for completing

membrane fission.

GTP hydrolysis induces clustering of dynamin-amphiphysin ring complexes

Although we determined the requirement of GTP hydrolysis in membrane constriction and

fission by the dynamin-amphiphysin complexes, structural changes of the complexes were not

clearly resolved in the *in vitro* assay system using LUVs. To improve the resolution, we used

rigid lipid nanotubes containing glycolipid galactosylceramide (GalCer) (Wilson-Kubalek,

Brown, Celia, & Milligan, 1998), instead of using LUVs in the *in vitro* assay system. Lipid

nanotubes are rod-shaped liposomes and similar in size to the unconstricted necks of

clathrin-coated pits observed *in vivo* (Figure 2A, Nanotube). Dynamin-amphiphysin complexes

assembled into helices on the lipid nanotubes (Figure 2A, No GTP), which is similar to those

formed by dynamin alone (Stowell, Marks, Wigge, & McMahon, 1999). Interestingly, the

dynamin-amphiphysin rings transiently formed clusters after the addition of GTP (Figure 2A,

GTP 1 s and 20 s, brackets). The dynamin-amphiphysin clusters were disorganized over time and

partially dissociated from the nanotubes (Figure 2A, GTP 30 s and 1 min).

To correlate the dynamics of dynamin-amphiphysin complexes with GTP hydrolysis, we

examined structural changes of the complexes on lipid nanotubes at different transition states of

GTP hydrolysis. The appearance of dynamin-amphiphysin helices was unchanged even in the

presence of GTPγS or GMP-PNP (Figure 2B, No GTP, GTPγS and GMP-PNP). Interestingly,

addition of GDP and vanadate induced rearrangement of the dynamin-amphiphysin ring

complexes to form clusters similar to those observed after the addition of GTP (Figure 2B, GDP +

vanadate). The average pitch between washer-rings in the clusters were shorter (15.00 nm ± 2.2,

mean pitch ± s.e.m.) compared to the average pitch of the ring complexes in No GTP control

(19.96 nm ± 0.47, mean pitch ± s.e.m.). In contrast, GDP alone did not affect the distribution of

dynamin-amphiphysin rings (Figure 2B, GDP). Finally, the dynamin-amphiphysin ring

complexes were disorganized and eventually dissociated from the lipid nanotubes 10 min after

the addition of 1 mM GTP (Figure 2B, GTP 10 min). Taken these results together, the

dynamin-amphiphysin ring complexes transiently form clusters in the GTP hydrolysis transition

state of GDP·Pi during which membrane tubules are constricted.

HS-AFM revealed dynamic clustering of dynamin-amphiphysin ring complexes upon GTP

hydrolysis

To elucidate the dynamics of dynamin-amphiphysin ring complexes during the membrane

constriction and fission, we analyzed the clustering process of the complexes using HS-AFM

(Ando, Uchihashi, & Kodera, 2013). LUVs were stably immobilized on the carbon-coated and

glow-discharged mica substrate (Supplementary Figure S2A; Supplementary Movie S1), and

they were successfully tubulated in the presence of dynamin and amphiphysin (Supplementary

Figure S2B; Supplementary Movie S2). The dynamin-amphiphysin rings on the lipid tubules

were aligned with an almost regular pitch (22.04 nm ± 0.68, mean pitch ± s.e.m.) and they were

immobile before GTP addition (Figure 3A, 0 s and 21 s; Supplementary Movie S3). Interestingly, the dynamin-amphiphysin rings became mobile after GTP addition and eventually formed clusters consisting of a few rings with shorter pitch (15.72 nm ± 0.30, mean pitch ± s.e.m.) (Figure 3A, from 42 s to 131 s; Supplementary Movie S3). Particle tracking analyses of the individual dynamin-amphiphysin rings showed that the dynamin-amphiphysin complexes were static before GTP addition (Figure 3B, 5-21 sec; Supplementary Movie S4), but addition of 1mM GTP stimulated longitudinal movement of the ring complexes, leading to the cluster formation (Figure 3B, 38-54 sec, 38-86 sec and 38-118 sec; Supplementary Movie S5). Although membrane fission was not observed in this sample probably due to a strong attachment of the lipid tubule to the substrate, the rings had a tendency to constrict during cluster formation (Figure 3C; Supplementary Figure S3). These results suggest that dynamin-amphiphysin ring complexes undergo two modes of structural changes, longitudinal clustering and radial constriction, during GTP hydrolysis.

Membrane fission occurs at protein-uncoated regions flanking dynamin-amphiphysin clusters

We next tried to correlate the cluster formation of dynamin-amphiphysin ring complexes with membrane constriction and fission. In the representative sample in which membrane constriction and fission occurred, a few dynamin-amphiphysin rings merged to form a cluster over time after GTP addition (Figure 4A, 0 s, 125.3 s, 185.5 s and 227.5 s; Supplementary Movie S6). Interestingly, membrane constriction occurred at flanking regions of the cluster where membrane was bare of dynamin-amphiphysin complexes (Figure 4A, fission point (FP).1 and FP.2). The heights at sites marked with FP.1 and FP.2 were not changed before constriction (Figure 4B, before constriction), but they became lower in a stepwise manner from a pre-constriction height of around 30 nm down to 20-25 nm or below (Figure 4B, after constriction). Similar longitudinal redistribution of the dynamin-amphiphysin rings before membrane constriction was also observed in another sample, in which constriction occurred at one end of clustered dynamin-amphiphysin complexes (Figure.4c, arrow；Supplementary Movie S8). These results strongly suggest that membrane constriction and fission occur at the protein-uncoated regions created as a result of the clustering of dynamin-amphiphysin ring complexes.

Amphiphysin contributes to efficient vesicle formation by controlling cluster formation

We previously demonstrated that amphiphysin stimulates the GTPase activity of dynamin and thus enhances vesicle biogenesis (Yoshida et al., 2004). In this study, we also noticed that the

average size of vesicles formed by dynamin-amphiphysin complexes (69.99 ± 2.93 nm, mean

diameter ± s.e.m.) was significantly smaller compared to those formed by dynamin alone (204.58

± 12.25 nm, mean diameter ± s.e.m.) after GTP addition (Figure 5A). Consistently,

dynamin-amphiphysin complex formed constriction sites with shorter intervals (150.28 ± 9.75

nm, mean intervals ± s.e.m.) compared to those formed by dynamin alone (193.49 ± 15.82 nm,

mean intervals ± s.e.m.) in the presence of GDP and vanadate (Figure 5B). To further elucidate

roles of amphiphysin in the membrane constriction and fission, the cluster formation by dynamin

alone was compared to that by dynamin-amphiphysin complexes, using lipid nanotubes. As

already described, dynamin-amphiphysin complexes formed clusters with a few rings in the

presence of GDP and vanadate (34.22 ± 1.66 nm, mean cluster size ± s.e.m.) (Figure 5C,

Dynamin + Amphiphysin). In contrast, dynamin alone formed larger-sized clusters consist of

more ring complexes (59.27 ± 4.69 nm, mean cluster size ± s.e.m.) (Figure 5C, Dynamin). These

results suggest that amphiphysin contributes to the effective generation of properly sized vesicles

by controlling the cluster formation of dynamin-amphiphysin ring complexes.

DISCUSSION

1   In this study, we analyzed dynamics of dynamin-amphiphysin ring complexes during
membrane constriction and fission using EM and HS-AFM. EM analyses showed that GTP
hydrolysis is required for both membrane constriction and fission, but dissociation of hydrolytic
products (GDP and/or phosphate) seems necessary for the completion of membrane fission
(Figure 1). In the presence of GTP or GDP and vanadate, dynamin-amphiphysin ring complexes
are reorganized, resulting in the formation of clusters consisting of a few dynamin-amphiphysin
rings (Figure 2). HS-AFM analyses directly demonstrated that GTP hydrolysis induces dynamic
longitudinal movement of the dynamin-amphiphysin rings as well as constriction during cluster
formation (Figure 3). Interestingly, HS-AFM analyses also demonstrated that membrane
constriction and fission occur at the "protein-uncoated" regions created as a result of cluster
formation of dynamin-amphiphysin complexes (Figure 4). Finally, we found that amphiphysin
contributes to effective biogenesis of endocytic vesicles by regulating size of the clusters formed
by dynamin-amphiphysin ring complexes (Figure 5).

There is a consensus view about the requirement of GTP hydrolysis in membrane fission, but
the requirement of GTP hydrolysis in membrane constriction is still controversial (Antonny et al.,
2016). Membrane tubules are constricted in the presence of non-hydrolyzable GTP analogue
(Chen et al., 2004; Mears et al., 2007; Zhang & Hinshaw, 2001) and more constricted with a
GTP-loaded GTPase defective K44A mutant (Sundborger et al., 2014). In both cases,
membrane tubules are evenly constricted and periodical membrane constriction sites which lead
to membrane fission is not created. In the present study, we showed that membrane constriction
sites are created in the presence of GDP and vanadate, which mimicked a transition state of
GTP hydrolysis (GDP $\cdot$ Pi), suggesting that complete hydrolysis of GTP is required for the
formation of constriction sites leading to membrane fission (Figure 1B). Membrane fission has
never been observed in the presence of GDP and vanadate, suggesting that release of GTP
hydrolytic products (GDP and/or phosphate) is a prerequisite for membrane fission. Further
analyses will more precisely reveal which intermediate state in the GTPase reaction is
responsible for the membrane fission or how many GTPase cycles are required for it.

In this study, we revealed that dynamin-amphiphysin ring complexes are rearranged to form
their clusters upon GTP hydrolysis (Figure 2 and Figure 3) and membrane fission occurs at the
flanking "protein-uncoated" membrane regions (Figure 4). In the "constrictase" model, dynamin
constricts membrane until the membrane neck reaches to the hemi-fission state, which leads to
spontaneous membrane fission (Chen et al., 2004; Hinshaw & Schmid, 1995; Mears et al., 2007).

However, several lines of evidences are apparently inconsistent with this simple model. For

instance, the super-constricted state of dynamin does not constrict the membrane sufficiently

enough to reach the hemi-fission state (Sundborger et al., 2014) and membrane tension and/or

torsion is required to overcome the energy barrier to fission (Bashkirov et al., 2008; Morlot et

al., 2012; Roux et al., 2006). Based on these results and our results in this study, we propose a

novel model of membrane fission by dynamin-amphiphysin complexes, for which we term

"clusterase" model (Figure 6). First, dynamin-amphiphysin are assembled to form washer

ring-like or helical polymers to induce membrane tubulation (Figure 6A). GTP hydrolysis

induces dynamic longitudinal rearrangement of dynamin-amphiphysin rings to form clusters

(Figure 6B). The clustered dynamin-amphiphysin rings also constrict to give local tension

and/or torsion to the membrane tube at the edge of the clusters (Figure 6C). Alternatively, the

dynamin-amphiphysin clusters may serve as a lipid diffusion barrier that causes friction leading

to membrane scission (Simunovic et al., 2017). Finally, membrane fission occurs along with

disassembly of the dynamin-amphiphysin polymers into oligomers (Figure 6D). Longitudinal

rearrangement upon GTP hydrolysis similar to the cluster formation by the

dynamin-amphiphysin complexes was also observed in an EM study on the dynamics of

dynamin with lipid nanotubes (Stowell et al., 1999) and more recently by HS-AFM analyses on

dynamics of ΔPRD dynamin (Colom, Redondo-Morata, Chiaruttini, Roux, & Scheuring, 2017),

suggesting that the longitudinal rearrangement is an intrinsic property of dynamin during

membrane fission.

We revealed that amphiphysin possibly contributes to effective vesicle biogenesis by

controlling the number of constriction sites in a long membrane tubule formed *in vitro* (Figure

5).Ttubular structures have long been known to be present in various synapses, and they are

described as "membrane tubules" (J. Heuser & Miledi, 1971) , "cisternae" (J. E. Heuser &

Reese, 1973), "synaptic tubules" (Samorajski, Ordy, & Keefe, 1966) or "anastomosing tubules"

(Ekstrom von Lubitz, 1981). The tubules are enriched in endocytic proteins including dynamin,

synaptojanin, amphiphysin, and endophilin (Fuchs, Brandstatter, & Regus-Leidig, 2014; Takei

et al., 1998), and the presence of the tubules becomes more prominent when synapses are

stimulated (Fuchs et al., 2014; Takei et al., 1998), or when membrane fission is blocked in

dynamin 1 K.O. mice (Ferguson et al., 2007). These findings strongly suggest that the tubular

structures represent endocytic intermediate at which dynamin-amphiphysin-dependent synergic

vesicle formation takes place in the synapse. Besides amphiphysin, other BAR domain proteins,

endophilin and syndapin, are also implicated in synaptic vesicle recycling (Dittman & Ryan,

2009; Koch et al., 2011; Milosevic et al., 2011). One of the important future goals of dynamin

study would be to clarify regulatory mechanisms by which dynamin alters its interactions with

various BAR proteins during synaptic vesicle recycling process.

In conclusion, live imaging analyses using HS-AFM in this study and a study from another

group (Colom et al., 2017) gave new mechanistic insights into the dynamin-mediated membrane

fission. Combinatory approaches using high temporal resolution imaging with HS-AFM and

high spatial resolution structural analyses with X-ray crystallography or Cryo-EM will be the

most powerful approach in resolving various dynamic membrane remodeling processes in the

future.

MATERIALS AND METHODS

Purification of dynamin1 and amphiphysin

Human dynamin1 was purified using the method of Warnock et al with some modification

(Warnock, Hinshaw, & Schmid, 1996). Sf9 cells grown in 600 ml of SF-900II SFM (Life

Technologies) to the cell density of 1×10^6 cells/ml and the cells were infected with

baculoviruses expressing dynamin1. After cultivation of cells at 28 ℃ for 69 hours, the infected

Sf9 cells were harvested by centrifugation at $500 \times g$ for 10 min. The cell pellet was

resuspended by 1/20 of the culture volume (30 ml) of HCB (Hepes column buffer)100 (20 mM

Hepes, 100 mM NaCl, 2 mM EGTA, 1 mM $MgCl_2$, 1 mM DTT, 1 mM PMSF, 1 μg/ml

Pepstatin A, 40 μM ALLN, pH 7.2) and cells were sonicated using a sonicator

(Advanced-Digital SONIFIER model 250, BRANSON). The cell lysate was mixed with equal

volume of HCB0 (20 mM Hepes, 2 mM EGTA, 1 mM $MgCl_2$, 1 mM DTT, 1 mM PMSF, 1

μg/ml Pepstatin A, 40 μM ALLN, pH 7.2) to make HCB50 (20 mM Hepes, 50 mM NaCl, 2

mM EGTA, 1 mM $MgCl_2$, 1 mM DTT, 1 mM PMSF, 1 μg/ml Pepstatin A, 40 μM ALLN,

pH7.2) and centrifuged at $210,000 \times g$ for 1h at 4℃. Ammonium sulfate was added to the

cleared lysate to the 30% saturation and incubated at 4℃ for 30min and centrifuged at $10,000 \times$

g for 10min to recover the dynamin1 containing fraction in the pellet. The dynamin1 pellet was

resuspended with 20 ml of HCB50 and dialyzed against 2L of HCB50 for total 4 hours (2 hours,

2 times) using dialysis membrane (Spectra/Por®Dialysis Membrane MWCO: 3500). The

dialyzed dynamin1 fraction was applied to Mono Q5/50GL column (GE healthcare) and bound

proteins were eluted stepwise using HCB50, HCB100, HCB250 and HCB1000 buffers. Purified

dynamin1 was recovered in HCB250 fraction and purity was determined by SDS-PAGE

(Supplementary Figure S1A, Dynamin).

Human amphiphysin was purified following manufacture's instruction (GE Healthcare) with

slight modifications. Host bacteria BL21 (DE3) transformed with an expression construct for

GST fusions of human amphiphysin (pGEX6P2-HsAMPH) were grown in 1 L of LB medium

to the cell density of 0.6-0.8 (OD 600 nm) at 37 ℃ and then protein expression was induced at

18 ℃ for 12 hours in the presence of 0.1 mM IPTG. The bacterial cells were harvested by

centrifugation at $7,000 \times g$ for 10 min and cell pellet was resuspended by 1/10 culture volume

(100 ml) of Elution/Wash 300 buffer (50 mM Tris-HCl, pH 8.0, 300 mM NaCl). The

resuspended cells were sonicated using Advanced-Digital SONIFIER model 250D (Branson)

and centrifuged at $261,000 \times g$ for 30 min at 4 ℃ and cleared lysate was recovered in

supernatant. To the cleared lysate, 1/100 culture volume (1 ml in bed volume) of Glutathione

Sepharose 4B Beads (GE Healthcare) was added and they are mixed using rotating mixer for 1h

at 4°C. The beads were washed with the Elution/Wash 300 buffer for 5 times in a repeated cycle

of centrifugation at 420 × g for 5 min at 4 ℃ followed by mixing with rotator for 5 min at 4℃.

The beads with purified GST fusions of amphiphsyin were equilibrated with PreScission Buffer

(50 mM Tris-HCl, 150 mM NaCl, 1 mM EDTA, 1 mM DTT, pH 7.0) and GST-tag was

removed by PreScission Protease (GE Healthcare) by incubating for 12 h at 4 ℃. The purified

amphiphysin was recovered by centrifuge (12,000 × g, 5 min at 4 °C) using spin column

(Ultrafree-Mc, GV 0.22 μm, Millipore) and purity was determined by SDS-PAGE

(Supplementary Figure S1A, Amphiphysin).

Preparation of LUVs and lipid nanotubes

Large unilamellar vesicles (LUVs) and lipid nanotubes were prepared as previously described

(Takei, Slepnev, & De Camilli, 2001). For LUVs, 70% PS (Cat. No 840032C, Avanti), 10%

biotinPE (Avanti) and 10 % cholesterol (Avanti) were mixed and, for lipid nanotubes 40% NFA

Galactocerebrosides (Sigma C1516), 40% PC (Avanti), 10% PI(4,5)P_2 (Calbiochem) and 10%

cholesterol (Avanti) were mixed in 250 μl of chloroform in a glass vial (Mighty Vial No.01 4 ml,

Maruemu Cat. No 5-115-03). Then chloroform was evaporated using slow-flow nitrogen gas to

produce lipid a lipid film on the glass and then completely dried in a vacuum desiccator for

30min. The dried lipid was rehydrated by water-saturated nitrogen gas followed by addition of

250 μl of filtered 0.3M sucrose for 2h at 37 ℃. The resultant LUVs and lipid nanotubes were

passed through 0.4 μm- and 0.2 μm- polycarbonate filters respectively 11 times using Avanti

Mini extruder. The LUVs and lipid nanotubes (1 mg/ml of final concentration) were stored in

dark at 4 ℃ avoiding photooxidation.

EM imaging of in vitro assay with liposomes and dynamin-amphiphysin complexes

LUVs and lipid nanotubes were diluted to 0.17 mg/ml in cytosolic buffer (25 mM

Hepes-KOH, pH 7.2, 25 mM KCl, 2.5 mM Magnesium acetate, 0.1 M K-glutamate, pH 7.4).

Dynamin-amphiphysin complexes (1:1 in molar ratio) were diluted to 0.6 μM in the cytosolic

buffer. Formvar filmed EM grids were carbon-coated, then glow-discharged. Droplets of the

diluted lipids (10μl each) were prepared on Parafilm and adsorbed on EM grids for 5 min at

room temperature. Then the EM grids with lipids were transferred to other droplets of the

diluted dynamin-amphiphysin complexes and incubated for 30 min at room temperature in a

humid chamber. To see the temporal effect of GTP hydrolysis, the EM grids were transferred to

1mM of GTP and incubated for various time periods (from 1s to 10 min) at room temperature.

Alternatively, the EM grids were incubated either GTP, GTPγS, GMP-PNP, GDP plus

Vanadate and GDP to analyze GTP hydrolysis transition state structures. The EM grids were

negatively stained with filtered 2% uranyl acetate and observed with transmission electron

microscope (HITACHI H-7650).

HS-AFM imaging

All AFM images shown in this article were capture by a laboratory-built HS-AFM in which

the amplitude-modulation mode was used. For the HS-AFM imaging, a small cantilever with

dimensions of 7-μm long, 2-μm wide, and 90-nm thick was used (Olympus). Its nominal spring

constant and resonant frequency were ~0.1 N/m and ~800 kHz in an aqueous solution,

respectively. To obtain a sharp tip, an amorphous carbon pillar was grown on the original

bird-beak tip of the cantilever by electron beam deposition (EBD) and then sharpened by a

plasma etching in an argon environment. The typical radius of the EBD tip was approximately 2

nm after sharpening. For the amplitude-modulation imaging, the cantilever was oscillated with

amplitude less than 10 nm under free oscillation condition and the set-point was set at ~90% of

the free oscillation amplitude. For HS-AFM imaging of liposomes and dynamin-amphiphysin

complexes with lipid tubules or nanotubes, we used mica covered with carbon film. After

coating a freshly cleaved mica surface with carbon film, hydrophilic treatment was carried out

by a grow discharge. The liposomes (0.17 mg/ml) were deposited on the hydrophilic mica

surface and incubated for 5 min at room temperature followed by deposition of proteins (0.6 μM

of dynamin1 and amphiphysin) for 30 min at room temperature. After the incubation, the

sample was thoroughly washed by cytosolic buffer to remove excess liposomes and proteins.

After the washing, the cantilever tip was approached and the imaging was performed under the

buffer.

Quantitative data analyses of EM and HS-AFM images

The EM and HS-AFM images were randomly captured to avoid data manipulation and

representative images were shown in all the figures. The average pitch between the

dynamin-amphiphysin washer rings in EM images (Figure 2) and HS-AFM images (Figure 3),

diameter of vesicles, intervals between constriction sites and size of clusters generated by either

dynamin-amphiphysin complex or dynamin (Figure 5), were all measured by FIJI (Schindelin et

al., 2012). Experimental data was statistically analyzed using Excel (Microsoft) or Prism 7

(GraphPad software).

ACKNOWLEDGEMENTS

The authors thank Dr. Aurélien Roux and Dr. Adai Colom (University of Geneva), Dr. Lorena

Redondo-Morata (INSERM U1006/Aix-Marseille Université) and Dr. Masatoshi Ichikawa

(Kyoto University) for their critical reading of the manuscript. This work was supported by

JST/CREST program (#JPMJCR13M1) (to T.A. and K.T.). This work was also supported in part

by JSPS KAKENHI Grant Numbers JP15H03540 (to T.U.), MEXT KAKENHI Grant Numbers

JP16H00830 and JP16H00758 (to T.U.).

1   *COMPETING INTERESTS*

2   No competing interests declared.

REFERENCES

Ando, T., Uchihashi, T., & Kodera, N. (2013). High-speed AFM and applications to

biomolecular systems. *Annu Rev Biophys, 42*, 393-414.

doi:10.1146/annurev-biophys-083012-130324

Antonny, B., Burd, C., De Camilli, P., Chen, E., Daumke, O., Faelber, K., . . . Schmid, S. (2016).

Membrane fission by dynamin: what we know and what we need to know. *EMBO J,*

35(21), 2270-2284. doi:10.15252/embj.201694613

Bashkirov, P. V., Akimov, S. A., Evseev, A. I., Schmid, S. L., Zimmerberg, J., & Frolov, V. A.

(2008). GTPase cycle of dynamin is coupled to membrane squeeze and release, leading

to spontaneous fission. *Cell, 135*(7), 1276-1286. doi:10.1016/j.cell.2008.11.028

Cao, H., Garcia, F., & McNiven, M. A. (1998). Differential distribution of dynamin isoforms in

mammalian cells. *Mol Biol Cell, 9*(9), 2595-2609.

Chen, Y. J., Zhang, P., Egelman, E. H., & Hinshaw, J. E. (2004). The stalk region of dynamin

drives the constriction of dynamin tubes. *Nat Struct Mol Biol, 11*(6), 574-575.

doi:10.1038/nsmb762

Cocucci, E., Gaudin, R., & Kirchhausen, T. (2014). Dynamin recruitment and membrane

scission at the neck of a clathrin-coated pit. *Mol Biol Cell, 25*(22), 3595-3609.

doi:10.1091/mbc.E14-07-1240

Colom, A., Redondo-Morata, L., Chiaruttini, N., Roux, A., & Scheuring, S. (2017). Dynamic

remodeling of the dynamin helix during membrane constriction. *Proc Natl Acad Sci U S*

A, 114(21), 5449-5454. doi:10.1073/pnas.1619578114

Cook, T., Mesa, K., & Urrutia, R. (1996). Three dynamin-encoding genes are differentially

expressed in developing rat brain. *J Neurochem, 67*(3), 927-931.

Cook, T. A., Urrutia, R., & McNiven, M. A. (1994). Identification of dynamin 2, an isoform

ubiquitously expressed in rat tissues. *Proc Natl Acad Sci U S A, 91*(2), 644-648.

David, C., McPherson, P. S., Mundigl, O., & de Camilli, P. (1996). A role of amphiphysin in

synaptic vesicle endocytosis suggested by its binding to dynamin in nerve terminals.

Proc Natl Acad Sci U S A, 93(1), 331-335.

Di Paolo, G., Sankaranarayanan, S., Wenk, M. R., Daniell, L., Perucco, E., Caldarone, B. J., . . .

De Camilli, P. (2002). Decreased synaptic vesicle recycling efficiency and cognitive

deficits in amphiphysin 1 knockout mice. *Neuron, 33*(5), 789-804.

Dittman, J., & Ryan, T. A. (2009). Molecular circuitry of endocytosis at nerve terminals. *Annu*

Rev Cell Dev Biol, 25, 133-160. doi:10.1146/annurev.cellbio.042308.113302

Ekstrom von Lubitz, D. K. (1981). Ultrastructure of the lateral-line sense organs of the ratfish, Chimaera monstrosa. *Cell Tissue Res, 215*(3), 651-665.

Evergren, E., Marcucci, M., Tomilin, N., Low, P., Slepnev, V., Andersson, F., . . . Shupliakov, O. (2004). Amphiphysin is a component of clathrin coats formed during synaptic vesicle recycling at the lamprey giant synapse. *Traffic, 5*(7), 514-528. doi:10.1111/j.1398-9219.2004.00198.x

Faelber, K., Held, M., Gao, S., Posor, Y., Haucke, V., Noe, F., & Daumke, O. (2012). Structural insights into dynamin-mediated membrane fission. *Structure, 20*(10), 1621-1628. doi:10.1016/j.str.2012.08.028

Faelber, K., Posor, Y., Gao, S., Held, M., Roske, Y., Schulze, D., . . . Daumke, O. (2011). Crystal structure of nucleotide-free dynamin. *Nature, 477*(7366), 556-560. doi:10.1038/nature10369

Ferguson, S. M., Brasnjo, G., Hayashi, M., Wolfel, M., Collesi, C., Giovedi, S., . . . De Camilli, P. (2007). A selective activity-dependent requirement for dynamin 1 in synaptic vesicle endocytosis. *Science, 316*(5824), 570-574. doi:10.1126/science.1140621

Ferguson, S. M., & De Camilli, P. (2012). Dynamin, a membrane-remodelling GTPase. *Nat Rev Mol Cell Biol, 13*(2), 75-88. doi:10.1038/nrm3266

Ford, M. G., Jenni, S., & Nunnari, J. (2011). The crystal structure of dynamin. *Nature, 477*(7366), 561-566. doi:10.1038/nature10441

Fuchs, M., Brandstatter, J. H., & Regus-Leidig, H. (2014). Evidence for a Clathrin-independent mode of endocytosis at a continuously active sensory synapse. *Front Cell Neurosci, 8*, 60. doi:10.3389/fncel.2014.00060

Heuser, J., & Miledi, R. (1971). Effects of lanthanum ions on function and structure of frog neuromuscular junctions. *Proc R Soc Lond B Biol Sci, 179*(1056), 247-260.

Heuser, J. E., & Reese, T. S. (1973). Evidence for recycling of synaptic vesicle membrane during transmitter release at the frog neuromuscular junction. *J Cell Biol, 57*(2), 315-344.

Hinshaw, J. E., & Schmid, S. L. (1995). Dynamin self-assembles into rings suggesting a mechanism for coated vesicle budding. *Nature, 374*(6518), 190-192. doi:10.1038/374190a0

Kirchhausen, T., Owen, D., & Harrison, S. C. (2014). Molecular structure, function, and dynamics of clathrin-mediated membrane traffic. *Cold Spring Harb Perspect Biol, 6*(5), a016725. doi:10.1101/cshperspect.a016725

Koch, D., Spiwoks-Becker, I., Sabanov, V., Sinning, A., Dugladze, T., Stellmacher, A., . . .
Qualmann, B. (2011). Proper synaptic vesicle formation and neuronal network activity
critically rely on syndapin I. *EMBO J, 30*(24), 4955-4969. doi:10.1038/emboj.2011.339

Koenig, J. H., & Ikeda, K. (1989). Disappearance and reformation of synaptic vesicle
membrane upon transmitter release observed under reversible blockage of membrane
retrieval. *J Neurosci, 9*(11), 3844-3860.

Marks, B., Stowell, M. H., Vallis, Y., Mills, I. G., Gibson, A., Hopkins, C. R., & McMahon, H. T.
(2001). GTPase activity of dynamin and resulting conformation change are essential for
endocytosis. *Nature, 410*(6825), 231-235. doi:10.1038/35065645

Mattila, J. P., Shnyrova, A. V., Sundborger, A. C., Hortelano, E. R., Fuhrmans, M., Neumann,
S., . . . Frolov, V. A. (2015). A hemi-fission intermediate links two mechanistically
distinct stages of membrane fission. *Nature, 524*(7563), 109-113.
doi:10.1038/nature14509

McMahon, H. T., & Boucrot, E. (2011). Molecular mechanism and physiological functions of
clathrin-mediated endocytosis. *Nat Rev Mol Cell Biol, 12*(8), 517-533.
doi:10.1038/nrm3151

Mears, J. A., Ray, P., & Hinshaw, J. E. (2007). A corkscrew model for dynamin constriction.
Structure, 15(10), 1190-1202. doi:10.1016/j.str.2007.08.012

Milosevic, I., Giovedi, S., Lou, X., Raimondi, A., Collesi, C., Shen, H., . . . De Camilli, P.
(2011). Recruitment of endophilin to clathrin-coated pit necks is required for efficient
vesicle uncoating after fission. *Neuron, 72*(4), 587-601.
doi:10.1016/j.neuron.2011.08.029

Morlot, S., Galli, V., Klein, M., Chiaruttini, N., Manzi, J., Humbert, F., . . . Roux, A. (2012).
Membrane shape at the edge of the dynamin helix sets location and duration of the
fission reaction. *Cell, 151*(3), 619-629. doi:10.1016/j.cell.2012.09.017

Peter, B. J., Kent, H. M., Mills, I. G., Vallis, Y., Butler, P. J., Evans, P. R., & McMahon, H. T.
(2004). BAR domains as sensors of membrane curvature: the amphiphysin BAR
structure. *Science, 303*(5657), 495-499. doi:10.1126/science.1092586

Reubold, T. F., Faelber, K., Plattner, N., Posor, Y., Ketel, K., Curth, U., . . . Eschenburg, S.
(2015). Crystal structure of the dynamin tetramer. *Nature, 525*(7569), 404-408.
doi:10.1038/nature14880

Roux, A., Uyhazi, K., Frost, A., & De Camilli, P. (2006). GTP-dependent twisting of dynamin
implicates constriction and tension in membrane fission. *Nature, 441*(7092), 528-531.

doi:10.1038/nature04718

Samorajski, T., Ordy, J. M., & Keefe, J. R. (1966). Structural organization of the retina in the tree shrew (Tupaia glis). *J Cell Biol, 28*(3), 489-504.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., . . . Cardona, A. (2012). Fiji: an open-source platform for biological-image analysis. *Nat Methods, 9*(7), 676-682. doi:10.1038/nmeth.2019

Schmid, S. L., & Frolov, V. A. (2011). Dynamin: functional design of a membrane fission catalyst. *Annu Rev Cell Dev Biol, 27*, 79-105. doi:10.1146/annurev-cellbio-100109-104016

Simunovic, M., Manneville, J. B., Renard, H. F., Evergren, E., Raghunathan, K., Bhatia, D., . . . Callan-Jones, A. (2017). Friction Mediates Scission of Tubular Membranes Scaffolded by BAR Proteins. *Cell, 170*(1), 172-184 e111. doi:10.1016/j.cell.2017.05.047

Stowell, M. H., Marks, B., Wigge, P., & McMahon, H. T. (1999). Nucleotide-dependent conformational changes in dynamin: evidence for a mechanochemical molecular spring. *Nat Cell Biol, 1*(1), 27-32. doi:10.1038/8997

Sundborger, A. C., Fang, S., Heymann, J. A., Ray, P., Chappie, J. S., & Hinshaw, J. E. (2014). A dynamin mutant defines a superconstricted prefission state. *Cell Rep, 8*(3), 734-742. doi:10.1016/j.celrep.2014.06.054

Sweitzer, S. M., & Hinshaw, J. E. (1998). Dynamin undergoes a GTP-dependent conformational change causing vesiculation. *Cell, 93*(6), 1021-1029.

Takei, K., Haucke, V., Slepnev, V., Farsad, K., Salazar, M., Chen, H., & De Camilli, P. (1998). Generation of coated intermediates of clathrin-mediated endocytosis on protein-free liposomes. *Cell, 94*(1), 131-141.

Takei, K., McPherson, P. S., Schmid, S. L., & De Camilli, P. (1995). Tubular membrane invaginations coated by dynamin rings are induced by GTP-gamma S in nerve terminals. *Nature, 374*(6518), 186-190. doi:10.1038/374186a0

Takei, K., Slepnev, V. I., & De Camilli, P. (2001). Interactions of dynamin and amphiphysin with liposomes. *Methods Enzymol, 329*, 478-486.

Takei, K., Slepnev, V. I., Haucke, V., & De Camilli, P. (1999). Functional partnership between amphiphysin and dynamin in clathrin-mediated endocytosis. *Nat Cell Biol, 1*(1), 33-39. doi:10.1038/9004

Warnock, D. E., Hinshaw, J. E., & Schmid, S. L. (1996). Dynamin self-assembly stimulates its GTPase activity. *J Biol Chem, 271*(37), 22310-22314.

Wigge, P., Kohler, K., Vallis, Y., Doyle, C. A., Owen, D., Hunt, S. P., & McMahon, H. T. (1997).
Amphiphysin heterodimers: potential role in clathrin-mediated endocytosis. *Mol Biol*
Cell, 8(10), 2003-2015.
Wilson-Kubalek, E. M., Brown, R. E., Celia, H., & Milligan, R. A. (1998). Lipid nanotubes as
substrates for helical crystallization of macromolecules. *Proc Natl Acad Sci U S A,*
95(14), 8040-8045.
Yoshida, Y., Kinuta, M., Abe, T., Liang, S., Araki, K., Cremona, O., . . . Takei, K. (2004). The
stimulatory action of amphiphysin on dynamin function is dependent on lipid bilayer
curvature. *EMBO J, 23*(17), 3483-3491. doi:10.1038/sj.emboj.7600355
Zhang, P., & Hinshaw, J. E. (2001). Three-dimensional reconstruction of dynamin in the
constricted state. *Nat Cell Biol, 3*(10), 922-926. doi:10.1038/ncb1001-922
12
13

FIGURE LEGENDS

Figure 1. GTP hydrolysis is required and sufficient for membrane constriction by

dynamin-amphiphysin ring complexes (A) Electron micrographs of lipid tubules induced by

dynamin-amphiphysin ring complex before GTP addition (No GTP) and at different time points

after addition of 1 mM GTP (GTP 1 s, GTP 5 s, GTP 10 s, GTP 30 s and GTP 1 min). More than

thirty samples from three individual experiments were examined and representative images are

shown. Scale bar is 200 nm. (B) Electron micrographs of lipid tubules induced by

dynamin-amphiphysin ring complex without guanine nucleotide (No GTP) or with a transition

states analogue of GTPase reaction, by adding 1 mM each of slowly hydrolyzable GTP analogue

(GTPγS), nonhydrolyzable GTP analogue (GMP-PNP), GDP combined with vanadate (GDP +

vanadate), GDP (GDP), or GTP (GTP) for 10 min. More than thirty samples from three individual

experiments were examined and representative images are shown. Scale bar is 200 nm.

Figure 2. GTP hydrolysis induces clustering of dynamin-amphiphysin complexes on lipid

nanotubes (A) Electron micrographs of a lipid nanotube (Nanotube) and those with

dynamin-amphiphysin complexes before GTP addition (No GTP) and at different time points

after GTP addition (GTP 1 s, GTP 20 s, GTP 30 s, and GTP 1 min). Clusters of

dynamin-amphiphysin ring complexes are indicated (white brackets). More than thirty samples

from three individual experiments were examined and representative images are shown. Scale bar

is 100 nm. (B) Electron micrographs of lipid nanotubes after addition of dynamin-amphiphysin

complexes without guanine nucleotide (No GTP) or with a transition states analogue of GTPase

reaction, by adding 1 mM each of slowly hydrolysable GTP analogue (GTPγS), nonhydrolyzable

GTP analogue (GMP-PNP), GDP combined with vanadate (GDP + vanadate), GDP (GDP) and

GTP (GTP) for 10 min. More than thirty samples from three individual experiments were

examined and representative images are shown. Clusters of dynamin-amphiphysin ring

complexes are indicated (white brackets). The average pitch of washer-rings in the clusters is

15.00 ± 2.2 nm (mean pitch $\pm$ s.e.m., n=63 from 7 nanotubes) in GDP + vanadate, while the

average pitch of the ring complexes is 19.96 ± 0.47 nm (mean pitch $\pm$ s.e.m., n=81 from 9

nanotubes) in No GTP control. Scale bar is 100 nm.

Figure 3. Dynamic clustering of dynamin-amphiphysin rings during GTP hydrolysis (A)

HS-AFM images captured at 1 frame/s of dynamin-amphiphysin ring complexes on membrane

tubules before (0 s and 21 s) and after GTP addition at different time points (42 s, 63 s, 84 s, 100

s, 115 s and 131 s). Dynamin-amphiphysin rings (arrowheads) are assembled into three distinct

clusters (1, 2 and 3 at 131 s). The pitch of dynamin-amphiphysin rings on the lipid tubule was

22.04 ± 0.68 nm (mean pitch $\pm$ s.e.m., n=36 from 3 time points) before GTP addition and $15.72 \pm$

0.30 nm (mean pitch $\pm$ s.e.m., n=36 from 9 time points) after GTP addition. (B) Particle tracking

of dynamin-amphiphysin rings before (5-21 sec) and after addition of 1mM GTP (38-54 sec,

38-86 sec and 38-118 sec) from Supplementary Movie S4 and S5, respectively. Particle tracking

of the complexes in the cluster 2 (light blue, dark blue and magenta) and cluster 3 (red, yellow and

green) are shown. (C) Dynamin-amphiphysin rings tend to constrict during clustering. Average

heights before ($0 \leq t \leq 100$ sec) and after clustering (101 sec $\leq t$) are 38.38 ± 0.17 nm and $36.48 \pm$

0.18 nm for cluster 1, 32.62 ± 0.15 nm and 32.56 ± 0.18 nm for cluster 2, 27.90 ± 0.13 nm and

26.75 ± 0.18 nm for cluster 3, respectively. The heights were measured from the substrate

surface. The marks *** indicate $p < 0.001$ and n.s. is not significant, respectively.

Figure 4. Membrane fission occurs at the protein-uncoated regions flanking

dynamin-amphiphysin clusters (A) Clips of HS-AFM images captured at 0.42 frames/s

showing membrane fission by dynamin-amphiphysin complexes (0 s, 125.3 s, 185.5 s and 227.5

s) in Supplementary Movie S6. Membrane fission occurred at flanking regions of a

dynamin-amphiphysin cluster. Corresponding height profiles along the red line (shown in the 0 s

image) passing through the two fission points (arrows marked with FP.1 and FP.2) are shown

below, together with clustered dynamin-amphiphysin ring complexes (red arrowheads). (B)

Height profiles at fission points (FP.1 and FP.2) over time before (Supplementary Movie S7) and

after constriction (Supplementary Movie S6). The heights of the lipid tubules from the substrate

surface were measured at the fission points. (C) Clips of HS-AFM images showing clustering

dynamin-amphiphysin complexes and membrane constriction at flanking regions of the cluster

(arrow). HS-AFM images are shown in pseudo color. Scale bar is 40 nm.

Figure 5. Amphiphysin contributes to generation of uniformly-sized vesicles by controlling

dynamin-amphiphysin clusters (A) Representative EM images of membrane vesicles generated

by dynamin-amphiphysin complexes (Dynamin + Amphiphysin) or dynamin alone (Dynamin)

after addition of GTP. Size distribution of generated vesicles are shown in the right panel. The

average sizes of vesicles were 69.98 ± 0.61 nm (mean diameter $\pm$ s.e.m., n>30, N=3) for

dynamin-amphiphysin complexes and 204.57 ± 1.10 nm (mean diameter $\pm$ s.e.m., n>45, N=3) for

dynamin alone. Scale bar is 200 nm. (B) Representative EM images of membrane constriction

induced by dynamin-amphiphysin complexes (Dynamin + Amphiphysin) and dynamin alone

(Dynamin) in the presence of GDP and vanadate. Distribution of intervals between constriction

sites are quantified in the right panel. The average intervals of constriction sites induced are

150.28 ± 9.75 nm (mean intervals ± s.e.m., n=25 from 7 tubes) by dynamin-amphiphysin

complexes and 193.5 ± 15.8 nm (mean intervals ± s.e.m., n=46 from 15 tubes) by dynamin alone.

Scale bar is 200 nm. (C) Clustering of dynamin-amphiphysin complexes (Dynamin +

Amphiphysin) and dynamin alone (Dynamin) on lipid nanotubes in the presence of GDP and

vanadate. Clusters of Dynamin-amphiphysin rings are indicated (white brackets). Distribution of

cluster size were shown as scattered plot in the right panel. Average size of the clusters formed by

dynamin-amphiphysin complexes and dynamin alone are 34.22 ± 1.66 nm (mean cluster size ±

s.e.m., n=36 from 7 tubes) and 59.27 ± 4.69 nm (mean cluster size ± s.e.m., n=30 from 5 tubes)

respectively.   Scale bar is 100 nm.

Figure 6. Clusterase model of membrane constriction and fission mediated by

dynamin-amphiphysin complexes. (A) Assembly and alignment of dynamin-amphiphysin

washer rings-like or helical polymers on membrane tubules in the presence of GTP. (B)

Dynamic longitudinal rearrangement of dynamin-amphiphysin rings to form clusters induced by

GTP hydrolysis. (C) Constriction of the clustered dynamin-amphiphysin which may provide

local tension and/or torsion and/or friction at the edge of the clusters to induce membrane

constriction. (D) Membrane fission caused by disassembly of dynamin-amphiphysin complexes

into oligomers upon release of GTP hydrolytic products.

Figure 1.Takeda et al.

A

No GTP | GTP 1 s | GTP 5 s

GTP 10 s | GTP 30 s | GTP 1 min

B

No GTP | GTPγS | GMP-PNP

GDP+vanadate | GDP | GTP

Figure 2.Takeda et al.

A

Nano-tube | No GTP | GTP 1s | GTP 20s | GTP 30s | GTP 1min

B

No GTP | GTPγS | GMP-PNP | GDP +vanadate | GDP | GTP