Paleo-Oscillomics: Reconstructing Language-Relevant Computational Capacities in Neanderthals from the Molecular Basis of Neural Oscillations

Elliot Murphy[1] and Antonio Benítez-Burraco[2]
1. Division of Psychology and Language Sciences, University College London, London, UK
2. Faculty of Philology, University of Seville, Seville, Spain

Abstract: Language seemingly evolved from changes in brain anatomy and wiring. We argue that language evolution can be better understood if particular changes in phasal and cross-frequency coupling properties of neural oscillations, resulting in core features of language, are considered. Because we cannot track the oscillatory activity of the brain from extinct hominins, we used our current understanding of the language oscillogenome (that is, the set of genes responsible for basic aspects of the oscillatory activity relevant for language) to infer some properties of the Neanderthal oscillome. We have found that several candidates for the language oscillogenome show differences in their methylation patterns between Neanderthals and humans. We argue that differences in their expression levels could be informative of differences in cognitive functions important for language.

Keywords: Language evolution, language disorders, Neanderthals, candidate-gene approach, neural oscillations, methylation patterns

1. Introduction

Language is the most distinctive human ability. Since it does not fossilize, language scientists instead need to rely on indirect evidence to gain some insight about how it evolved in our clade. Language is ultimately the product of neural activity and so skulls and brain endocasts are objects of particular interest. Because of the intimate cross-talk between the skull and the brain, we expect that the analysis of the hominin skull shape and growth pattern can inform how the brains of extinct hominin species were organized and how they developed during growth; factors which impact cognition (Zollikofer and Ponce de León 2013). Language abilities in Neanderthals are a controversial issue (Johansson 2015). The skull of anatomically-modern humans (AMH) differs both in size and shape from the skull of Neanderthals (Bruner 2004). However, it is not still clear if calvarian differences correlate with endocast differences, and hence with brain function (Mounier et al. 2016), or if there is a AMH-specific brain developmental path (see Ponce de León et al. 2016 and Williams and Cofran 2016 for discussion). Comparisons with extant primates are helping to refine our understanding of these differences (Pearce et al. 2013). Overall, gross differences in brain structure and function between hominin species have been used to infer differences in cognitive abilities important for language processing (Boeckx and Benítez-Burraco, 2014a).

In neighboring domains, neuronal oscillatory activity has been acknowledged as the basis of multiple cognitive and behavioural processes (Buzsáki 2006). In our previous work we have showed that computational operations of language can be decomposed into generic processes that can be implemented via neural oscillations (Murphy 2015a,b) and that certain multiplexing algorithms responsible for the construction of hierarchical phrase structures appear to be species-specific (Murphy 2016a). We have also showed that neural oscillations provide a more reliable explanatory level of language than the examination of nerve tracks and brain regions involved in language processing; not just in neurotypical language processing, but in cognitive conditions that are human-specific and entail language dysfunction (Benítez-Burraco and Murphy 2016, Murphy and Benítez-Burraco 2016).

While we cannot track the oscillatory activity of extinct hominins, we believe that the next best solution to moving beyond this shortcoming consists in examining the coding regions and the expression patterns of the genes responsible for the brain's oscillatory activity putatively responsible for human language. We have recently identified and functionally characterized a set of 48 genes that comprises the core of our *language oscillogenome*; that is, the set of genes responsible for basic aspects of the oscillatory activity relevant for language (Murphy and Benítez-Burraco 2017). In this paper we have looked for differences between Neanderthals and us in the coding regions of these genes, and in their methylation patterns, as well as signals of positive selection in our species. Our ultimate aim is inferring, from these genomic differences, differences in the brain activity important for language processing.

2. Evolutionary changes in the language oscillogenome

In order to achieve our objective, we first gathered via systematic literature review and database searches the available information concerning fixed changes in AMH proteins compared to Neanderthal proteins, genomic regions positively selected in AMHs after our split from Neanderthals, and differentially methylated regions (DMRs) in AMHs compared to Neanderthals. We have extensively built on Green et al's (2010) seminal characterization of the Neanderthal genome, Prüfer et al's (2013) description of the Altai Neanderthal genome, and Gokhman et al's (2014, 2017) DNA methylation maps of modern humans and extinct hominins. We then looked for our candidates in these papers and databases. As noted in Murphy and Benítez-Burraco (2017), our candidates for the language oscillogenome fulfil three criteria: i) they are associated with language disorders (developmental dyslexia and/or specific language impairment) and/or language dysfunction in cognitive disorders entailing language deficits (schizophrenia and/or autism spectrum disorder), ii) they play a role in brain rhythmicity and/or are candidates for conditions entailing brain dysrhythmias, like epilepsy; and iii) gene-oscillations-language links can be confidently established for them.

Our findings are summarized in Table 1.

Gene	Fixed aa change in AMHs	Positively selected in AMHs	Differentially methylated in AMH skeletal samples	Enriched in AMH DMRs	Oscillomic/oscillopathic features
AUTS2		✓	↑ (body gene)		Epilepsy
CACNA1C			↑ (body gene)		β, γ
CNTNAP2	✓				α
COMT			↑ (downstream the gene)		α
DYRK1A		✓			Inhibition of neural activity
EGR1				✓	Epilepsy
ELP4		✓			High amplitude centrotemporal sharp waves
FMR1	✓				θ, γ
FOXP1		✓	↑ (body gene)		Epileptiform discharges
PDGFRB			↓ (body gene)		γ
ROBO1		✓			Epilepsy
SHANK3			↓ (downstream the gene)		Seizures, abnormal EEG

Table 1. AMH-specific changes in genes involved in brain oscillations important for languages processing. The information about fixed aminoacid changes in the protein (column 1) derives for

several sources, as noted in the main text. Evidence of positive selection (column 2) is from Green et al. (2010). The data about DMRs in AMH bone (column 3) are from Gokhman et al. 2017 (extended data table 2). The data about transcription factors enriched in AMH DMRs (column 4) is from Gokhman et al. 2014 (table S5). The information about the oscillomic/oscillopathic features of the gene is from Murphy and Benítez-Burraco 2017.

Among the 48 genes we highlighted in our previous work as part of the shared signature of abnormal brain oscillations associated with language deficits and core components of the language oscillogenome, 12 of these exhibit some difference between AMHs and Neanderthals. Two proteins bear fixed changes in one position in AMHs, although these changes are not expected to impact significantly on their structure and function: CNTNAP2 exhibits a Ile345Val change (reported as benign according to PolyPhen, and as tolerated (0.79) according to SIFT), whereas FMR1 shows a nearly fixed Ser145Ala change (frequency of 0.98 in AMHs) (classified as benign according to PolyPhen, and as tolerated (1) according to SIFT). More interestingly, 5 of our candidates are found within the regions that exhibit the strongest signals of positive selection in AMHs compared to Neanderthals, according to Green et al. (2010): *AUTS2*, *DYRK1A*, *ELP4*, *FOXP1*, and *ROBO1*. These positively selected regions contain numerous binding sites for transcription factors (Supplementary file, S1), which might have been affected by AMH-specific changes. Nonetheless, the most direct evidence of putative differences in the expression pattern of genes important for language under an oscillomic lens derive from the methylation maps of the AMH and the Neanderthal genomes obtained by Gokhman and colleagues (2014, 2017). We have found that several regions within the body gene of 3 of our candidates are hypermethylated in skeletal samples of AMHs compared to Neanderthals (*AUTS2*, *CACNA1C*, and *FOXP1*), whereas one region within the body of *PDGFRB* is found hypomethylated in our species. Likewise, one hypermethylated region is found downstream one of our candidates (*COMT*), whereas one hypomethylated region is located downstream *SHANK3* (Table 1 and supplementary file, S2). All these regions contain binding sites for several transcription factors of interest, like FOXA1, FOXA2, EP300, CBPB, MEF2A, RUNX3, FOXP2, FOXA2, CEBPB (Supplementary file, S2), which we have highlighted as important for language evolution, particularly, for changes related to the externalization of our linguistic thoughts (speech) (Boeckx and Benítez-Burraco 2014b, Benítez-Burraco and Boeckx 2015). Finally, we have found that EGR1 is one of the enriched transcription factors in AMH DMRs.

Although methylation is commonly associated to promoter regulation, it is also a regulatory mechanism of gene expression when affecting to splice sites, coding regions, binding sites for transcription factors, and distal regulatory elements, like enhancers, silencers or insulators (Jones et al. 2012, Miller and Grant 2013, Lev Maor et al. 2015). Accordingly, from the above differences in their methylation patterns, we might expect differences between Neanderthals and AMHs in the expression levels/patterns of several candidates for the language oscillogenome, namely, *AUTS2*, *CACNA1C*, *COMT*, *FOXP1*, *PDGFRB*, and *SHANK3*. Obviously, it is still to be determined whether the identified DMRs in skeletal methylation maps exist between the brains of AMHs and Neanderthals too. However, this might be the case. For example, the methylation levels of *PDGFRB* are similar across various human tissues, including the brain (Gokhman et al. 2014).

Interestingly, duplications and deletions of all these genes, purportedly resulting in abnormal high or low levels of the protein, respectively, result in cognitive and language deficits (Table 2).

Gene	Abnormal brain/cognitive/behavioural features associated to the mutation of the gene in humans	Abnormal brain/cognitive/behavioural features associated to the mutation of orthologues of the gene	Abnormal brain/cognitive/behavioural features associated to deletions of the full gene or the promoter region	Abnormal brain/cognitive/behavioural features associated to duplication of the full gene
AUTS2	Autism Mental retardation Intellectual disability Global developmental delay Abnormal response to serotonin reuptake inhibitors in non-psychotic unipolar depression	Alterations of forebrain neuron development Increased cell proliferation in the midbrain and the hindbrain Decreased size of the anterior region of the head Decreased number of cerebellar neurons Generalized epilepsy with febrile seizures	Global brain atrophy Moderate intellectual disability (DECIPHER patient 279043)	N/A
CACNA1C	Timothy syndrome Autism Schizophrenia Bipolar disorder Intellectual deficiency	Abnormal central nervous system synaptic transmission Abnormal nervous system physiology Abnormal learning/memory/conditioning	Autistic behavior (DECIPHER patient 290015)	N/A
COMT	22q11.2 Deletion Syndrome Susceptibility to schizophrenia	Abnormal increased prepulse inhibition Increased anxiety-related response Enhanced contextual conditioning behavior Impaired cued conditioning behavior Increased dopamine level Hyperactivity Abnormal spatial working memory	N/A	N/A
EGR1	Schizophrenia	Abnormal forebrain, midbrain, and hindbrain development Decreased dendrite morphogenesis Generalized epilepsy with febrile seizures	N/A	Abnormal emotion/affect behavior Global developmental delay (DECIPHER patient 304609)
FMR1	Fragile X syndrome Intellectual disability Mental Competency	Abnormal cranial nerve position Disrupted sensory neuron axon guidance Abnormal axons of the lateral longitudinal fasciculus Disorganized hindbrain axons Increased synapse assembly of cholinergic neurons	Global developmental delay (DECIPHER patient 284912)	N/A

	Abnormal dendritic spine morphology Decreased long-term synaptic potentiation Abnormal long term depression Decreased β wave amplitude Impaired learning and memory Impaired turning behavior			
FOXP1	Intellectual disability Severe speech delay Mental retardation with language impairment and with or without autistic features Attention deficit hyperactivity disorder	Cerebral visual impairment Intellectual disability Malformations of cortical development Autistic features	Abnormality of the face Abnormality of the nervous system Aggressive behavior Autism Hyperactivity Intellectual disability Short attention span (DECIPHER patient 264639)	N/A
PDGFRB	Basal ganglia calcification	Basal ganglia calcification and impairment Brain injuries	N/A	N/A
SHANK3	Phelan-McDermid syndrome Schizophrenia Intellectual disability	Reduced size of the anterior part of the head Delayed brain development Abnormal striatum morphology Abnormal dendritic spine and synapse morphology Reduced amount of glutamatergic and GABAergic neurons in the hindbrain Reduced NMDA-mediated synaptic currents Abnormal glutamate-mediated receptor currents Abnormal excitatory postsynaptic currents Reduced long term potentiation Decreased post-tetanic potentiation Abnormal paired-pulse inhibition Environmentally induced seizures Impaired synaptic plasticity Impaired coordination Abnormal motor learning and coordination Impaired spatial learning and memory	Sleep disturbance (DECIPHER patient 252486) Absent speech Delayed speech and language development Febrile seizures Global developmental delay Seizures (DECIPHER patient 285792)	Intellectual disability (DECIPHER patient 251542)

Abnormal vocalization
Abnormal social investigation
Increased grooming behavior
Abnormal response to novelty
Abnormal social/conspecific interaction

Table 2. Abnormal brain, cognitive, and/or behavioural features associated to mutations in genes involved in brain oscillations important for languages processing. The abnormal features associated to the genes were retrieved from the Phenotypes tool of Ensembl (http://www.ensembl.org/index.html). Abnormal features observed in humans (column 1) derive from several databases of diseases and phenotypes directly associated with a gene, including the Online Mendelian Inheritance in Man (OMIM) compendium, Orphanet, and DDG2P, as well as from several sources of diseases and traits associated with variants of interest, including COSMIC, ClinVar, dbGaP, EGA, GIANT, HGMD-Public, MAGIC, NHGRI-EBI GWAS catalog, and UniProt. The abnormal phenotypes linked to mutations of the orthologe genes in other species come from several databases, including the International Mouse Phenotyping Consortium and Europhenome (for mouse), ZFIN (for zebrafish), Rat Genome Database (for rat), Animal_QTLdb and Online Mendelian Inhertiance in Animals. Because of biological differences between species, some animal phenotypes will not be easy to translate to our species. Columns 3 contains the available information about abnormal brain, cognitive, and/or behavioural features associated in humans to deletions of the full gene or the promoter region of the gene, as provided by DECIPHER (https://decipher.sanger.ac.uk/). Column 4 displays the abnormal brain, cognitive, and/or behavioural traits associated to duplications of the full sequence of the genes of interest, as also provided by DECIPHER.

AUTS2 is involved in cytoskeletal regulation to control neuronal migration and neurite extension, but nuclear AUTS2 also controls transcription of several genes (Oksenberg et al. 2014, Hori and Hoshino 2017). The gene has been associated with differential processing speeds (Luciano et al. 2011) and is a candidate for epilepsy (Mefford et al. 2010). *CACNA1C* encodes the alpha 1C subunit of the Cav1.2 voltage-dependent L-type calcium channel, involved in β and γ generation (Kumar et al. 2015). Pathogenic variants of *CACNA1C* have been associated with low performance scores in semantic verbal fluency tasks in schizophrenics (Krug et al. 2010), as well as to childhood-onset epilepsy, executive dysfunction, intellectual disability, and/or autism spectrum disorder (Damaj et al. 2015). *COMT*, which encodes a catechol-O-methyltransferase that catalyzes the O-methylation of neurotransmitters like dopamine, epinephrine, and norepinephrine, has been regularly associated to schizophrenia, and particularly, to language dysfunction in this condition (Egan et al. 2001, Malhotra et al. 2002). The gene has been also associated to reading abilities in the neurotypical population (Landi et al. 2013), as well as to language performance and processing, and language acquisition, particularly to verbal fluency (Krug et al. 2009, Soeiro-De-Souza et al. 2013, Sugiura et al. 2017). Low activity levels in COMT has been associated to low voltage α (Enoch et al. 2003). Intriguingly, AMHs exhibit a hypomethylated region inside the body of one gene important for COMT function, namely *MSRA*, compared to Neanderthals (Gokhman et al. 2014, table S2). *EGR1* encodes a transcription factor that contributes to regulate neural plasticity and memory consolidation (Veyrac et al. 2014). In human epileptic foci, *EGR1* expression correlates with the frequency, amplitude and area of the interictal spikes, a hallmark of epileptic neocortex (Rakhade et al. 2007). Interestingly, *EGR1* is a target of FOXP2, the renowned "language gene" (Konopka et al. 2009), whereas EGR1 downregulates *PLAUR* (Matsunoshita et al. 2011), another of FOXP2's targets (Roll et al. 2010). *FMRP1* is the main candidate for Fragile X syndrome (Kaufmann et al. 2004, Smith et al. 2012). *Fmr1*-knockout

mice exhibit abnormal patterns of coupling between θ and γ oscillations in the CA1 area of the hippocampus, enhanced mGluR5 signalling resulting in altered neocortical rhythmic activity, and abnormally weak changes during tasks involving cognitive challenge (Hays et al. 2011, Radwan et al. 2016). FOXP1 interacts with FOXP2 to form functional heterodimers and *FOXP1* haplo-insufficiency has been associated with epileptiform discharges, speech delay, and motor delay (Carr et al. 2010). In mice, the knockout of *Pdgfrb*, which is involved in central nervous system development, gives rise to reduced auditory phase-locked γ, which correlates with distinctive features in schizophrenia, such as deficient spatial memory and impaired social behaviour (Nguyen et al. 2011, Nakamura et al. 2015). Finally, the knockdown of *Shank3*, which encodes a postsynaptic scaffolding protein important for the balance between neuronal excitation and inhibition, affects long-term potentiation and synaptic activity (Lanz et al. 2013), whereas mutation of the gene impacts neuronal excitation and inhibition (Lee et al. 2015, Lu et al. 2016), along with social interaction and grooming behaviour (Peça et al. 2011).

Although in this paper we are interested in changes that are specific to AMHs and that allegedly contributed to the emergence of our species-specific pattern of brain oscillations, we wish to note that two of our candidate genes exhibit a differential methylation pattern in Neanderthals compared to other hominins: *EGR1*, discussed above, and *KANSL1* (Table S2). The latter encodes a component of the NSL1 complex, which functions as a regulator of gene transcription and chromatin organization, and is a candidate for Koolen-de Vries syndrome, which presents with epilepsy, developmental delay, and moderate intellectual disability, that mostly affects expressive language abilities (Koolen et al. 2016).

3. The language network in Neanderthals: Paleoneurology and oscillomics

Comparative studies of Neanderthal and AMH cognition are vital if we are to hone our understanding of cognitive features unique to our species. As Mounier et al. (2016: 22) note, 'the use of palaeoneurology to infer phylogenies of our genus is rare'. An emerging consensus indicates that while the skull size and shape of AMHs and Neanderthals are comparable, their internal neural organisation was in a number of respects distinct (Pearce et al. 2013). For instance, Neanderthals lived at much higher latitudes than AMHs and would have likely had larger eyes and, consequently, larger visual cortices (Pearce and Dunbar 2012, Balzeau et al. 2012). Over the past couple of decades, digital anatomy and computed morphometrics have led to a seismic shift in functional craniology, exposing the generally plesiomorph structure of the Neanderthal braincase (Weber and Bookstein 2011, Bruner 2014, Williams and Cofran 2016). Figure 1 summarises the major findings of this research. Naturally, not every neuroanatomical difference between Neanderthals and AMHs will be relevant to evolutionary linguistics, but we will conservatively review those differences which strike us as most significant. Notably, Neanderthals display upper parietal bulging which approximates the AMH shape but crucially lacks what Bruner (2014: 123) describes as 'the overall bulging of the parietal surface characterizing the modern human brain globularity, in which the parietal proportions also enlarge in the longitudinal and vertical directions'. As we have already noted (Murphy and Benítez-Burraco 2016), the globularity of the AMH braincase appears to have had direct consequences for the efficiency of oscillatory couplings across the cortex and subcortex which we have, in turn, used to derive the cross-modular nature of linguistic representations. Bruner (2014: 123) concludes that the type of parietal changes exhibited by the AMH brain likely resulted in modifications to visuo-spatial integration and 'mental experiment capabilities', which we interpret as recursively generated symbolic manipulations or what Tattersall (2017: 65) regards as 'imagination'. Likewise, due to the expanded occipital lobe and visual cortices in Neanderthals, and due to the role of this region in α-inhibition vital

to the coordination of cross-cortical feature binding (Murphy 2016a), it is a possibility that Neanderthals were capable of executing a degree of control over representation integration not seen in other primates close to AMHs. Further support for this hypothesis comes from Table 2: *COMT* exhibits hypermethylation in AMHs compared to Neanderthals, and since low-voltage α has been associated with low activity levels in the gene (Enoch et al. 2003), with *COMT* additionally being associated with language performance and verbal fluency (Krug et al. 2009, Soeiro-De-Souza et al. 2013), it is reasonable to assume that this hypermethylation contributes to the type of α-inhibition required to control inter-modular featural integration.

Figure 1. Paleoneurological characters of Neanderthals from a replica of the skull and endocasts of Saccopastore 1. White arrows indicate derived traits while black arrows indicate plesiomorph traits, which impose constraints as a result of large brain size (reproduced from Bruner 2014: 123).

It was shown in Murphy (2016b) that a range of monkeys (putty-nosed monkeys, Campbell's monkeys, King Colobus monkeys, and New World monkeys) seem capable only of single-instance concatenation and not phrasal construction. Drawing on much neuroethological data, the likely oscillatory basis of this capacity was explored. It was proposed that phase-amplitude coupling between parahippocampal θ and cross-cortical γ was responsible for constructing individual call representations, and that primarily basal ganglia β increases signalled the maintenance of such calls in memory (Murphy 2016b).

At this point, we need to distinguish between the possibility that Neanderthals were capable of speech from the possibility that they were capable of internally generating hierarchically organised sets of mental representations. Neanderthals may have been capable of speech similar to AMHs due to the hard portion of the hyoid apparatus typically being preserved and their aural ability to be sensitive to frequencies (up to 5Hz) relevant for speech (Martinez et al. 2004). But what about the generation of hierarchically organised representations, the core of natural language? We believe that since current work is revealing an oscillatory code for language, and since the molecular basis of this code is in turn becoming amenable to reconstruction (Murphy and Benítez-Burraco 2017), the most reliable existing method of examining whether Neanderthals had anything resembling a proto-language is to draw

inferences from emerging genetic databases. For instance, while Neanderthals may have been sensitive to language-relevant frequencies, the question we are interested in is whether neural assemblies internally oscillating at the same frequencies were used in the service of constructing hierarchically organised representations. Relatedly, Mithen (2014) reviews the evidence for cognitive similarities and differences in AMHs and Neanderthals and concludes, for example, that while pigment use likely indicates a complex social and emotional life, it does not provide evidence for symbolic thought.

It is likely that the human-specific oscillatory profile documented in Murphy (2016a), differing from the Neanderthal oscillome proposed here, is responsible for Mithen's (1996, 2014) seminal distinction between the domain-specific Neanderthal mind (in which distinct core knowledge systems are largely isolated from each other) and the 'cognitive fluidity' of the AMH mind. Without this fluidity, integrating representations from distinct domains, Neanderthals 'could never develop such [complex social] relationships and initiate a process of domestication' (Mithen 2014: 12; concerning domestication, see also Benítez-Burraco et al. 2016). More specifically, we would like to suggest that the distinct θ-γ code we predicted for the Neanderthal brain can also explain its limited working memory (Wynn and Coolidge 2012), given the crucial role that coupling between these rhythms appears to have in working memory operations across a number of modalities (Murphy 2016a, Schomburg et al. 2014, Vosskuhl et al. 2015). A widening in the anterior fossa in Neanderthals (Figure 1) is associated with an expansion in Broca's area, argued in Murphy (2016a) to be implicated in language-relevant memory buffers. However, taken alongside the oscillogenomic evidence presented above – that Neanderthals may have exhibited reduced cross-frequency coupling between θ and γ due to differences in *CACNA1C* and *ELP4* expression – we should also conclude that the Neanderthal brain was likely incapable of exploiting this expanded Broca's area, at least in the service of maintaining sets of symbolic representations. As with the partly globular parietal modifications discussed above, this appears to be another major feature of the Neanderthal brain which begins to approximate neuroanatomical characteristics that we hypothesize are required for core aspects of AMH syntax, but which ultimately displays other features which are likely incompatible with it.

Returning to Table 1, *CACNA1C*, which is differentially methylated in AMHs, contributes to β and γ generation, and has been correlated with semantic verbal fluency, may set AMHs apart from Neanderthals by contributing to speech-related fluency through β (implicated in storing ongoing speech representations in memory) and γ (involved in the initial construction of discourse representations, as discussed above). Relatedly, the abnormal dendritic spine morphology associated with *FMR1*, along with the decreased β power (Table 2), strengthens this notion. Finally, the hypomethylation of *PDGFRB* in AMHs may have contributed to enhancing phase-locked γ in our species, with this distinct difference from the Neanderthal genome also likely playing a role in reducing the efficacy of any attempted cross-cortical representation integration in Neanderthals, confining them to domain-specific search spaces as opposed to the cognitive fluidity exhibited by AMH concatenation processes.

4. Conclusion

Overall, it is likely that the oscillatory differences we have documented between AMHs and Neanderthals are connected to the major cultural and technological novelties which occurred towards the end of the Middle Pleistocene and which distinguish AMHs from our closest relatives. Expanding our molecular understanding of the Neanderthal brain can contribute to expanding the line of inquiry into what we have here called paleo-oscillomics. While it is hard

to imagine anything in the fossil record shedding light on the *representational* basis of the Neanderthal brain (Berwick et al. 2013), reconstructing the Neanderthal oscillome can at least reveal aspects of its *computational* capacity.

Acknowledgments: This work was supported by an Economic and Social Research Council scholarship (1474910) to EM, and by funds from the Spanish Ministry of Economy and Competitiveness (grant numbers FFI2014-61888-EXP and FFI-2013-43823-P) to ABB. The authors wish express their gratitude to David Gokhman for sharing his data on methylation maps of Neanderthals and humans, and for kindly reading an earlier draft of the paper.

References

Balzeau, A., Holloway, R. L. and Grimaud-Hervé, D. (2012) Variations and asymmetries in regional brain surface in the genus *Homo*. *J. Hum. Evol.* 62: 696-706.

Benítez-Burraco, A. and Boeckx, C. (2015) Possible functional links among brain- and skull-related genes selected in modern humans. *Front, Psychol.* 6: 794.

Benítez-Burraco, A. and Murphy, E. (2016) The oscillopatic nature of language deficits in autism: from genes to language evolution. *Front. Hum. Neurosci.* 10: 120.

Benítez-Burraco, A., Lattanzi, W. and Murphy, E. (2016) Language impairments in ASD resulting from a failed domestication of the human brain. *Front. Neurosci.* 10: 373.

Berwick, R. C., Hauser, M. D. and Tattersall, I. (2013) Neanderthal language? Just-so stories take center stage. *Front. Psychol.* 4: 671.

Boeckx, C. and Benítez-Burraco, A. (2014a) The shape of the human language-ready brain. *Front. Psychol.* 5: 282

Boeckx, C. and Benítez-Burraco, A. (2014b) Globularity and language-readiness: Generating new predictions by expanding the set of genes of interest. *Front. Psychol.* 5: 1324.

Bruner, E. (2004) Geometric morphometrics and paleoneurology: brain shape evolution in the genus *Homo*. *J. Hum. Evol.* 47: 279-303.

Bruner, E. (2014) Functional craniology, human evolution, and anatomical constraints in the Neanderthal braincase. In *Dynamics of Learning in Neanderthals and Modern Humans*, vol. II: *Cognitive and Physical Perspectives. Replacement of Neanderthals by Modern Humans Series* (Akazawa, T., Ogihara, N., Tanabe, H.C. and Terashima, H., eds). Tokyo: Springer Japan. pp. 121-129.

Buzsáki, G. (2006) *Rhythms of the Brain.* New York: Oxford University Press.

Carr, C. W., Moreno-De-Luca, D., Parker, C., Zimmerman, H. H., Ledbetter, N., Martin, C. L., et al. (2010) Chiari I malformation, delayed gross motor skills, severe speech delay, and epileptiform discharges in a child with FOXP1 haploinsufficiency. *Eur. J. Hum. Genet.* 18: 1216-20.

Damaj, L., Lupien-Meilleur, A., Lortie, A., Riou,É.,Ospina, L. H., Gagnon, L., et al. (2015) CACNA1A haploinsufficiency causes cognitive impairment, autism and epileptic encephalopathy with mild cerebellar symptoms. *Eur. J. Hum. Genet.* 23: 1505-12.

Egan, M. F., Goldberg, T. E., Kolachana, B. S., Callicott, J. H., Mazzanti, C. M., Straub, R. E., et al. (2001) Effect of COMT Val108/158 Met genotype on frontal lobe function and risk for schizophrenia. *Proc. Natl. Acad. Sci. U.S.A.* 98: 6917-6922.

Enoch, M. A., Xu, K., Ferro, E., Harris, C. R. and Goldman, D. (2003) Genetic origins of anxiety in women: a role for a functional catechol-O-methyltransferase polymorphism. *Psychiatr. Genet.*13: 33–41.

Gokhman, D., Lavi, E., Prüfer, K., Fraga, M. F., Riancho, J. A., Kelso, J., et al. (2014) Reconstructing the DNA methylation maps of the Neandertal and the Denisovan. *Science* 344: 523-7.

Gokhman, D., Agranat-Tamir, L., Housman, G., Nissim-Rafinia, M., Nieves-Colón, M., Gu, H., et al. (2017) Recent regulatory changes shaped human facial and vocal anatomy. bioRxiv 106955; doi: https://doi.org/10.1101/106955

Green, R. E., Krause, J., Briggs, A. W., Maricic, T., Stenzel, U., Kircher, M., et al. (2010) A draft sequence of the Neandertal genome. *Science* 328: 710-22.

Hays, S. A., Huber, K. M. and Gibson, J. R. (2011) Altered neocortical rhythmic activity states in Fmr1 KO mice are due to enhanced mGluR5 signaling and involve changes in excitatory circuitry. *J. Neurosci.* 31: 14223-34.

Hori, K. and Hoshino, M. (2017) Neuronal migration and AUTS2 syndrome. *Brain Sci.* doi: 10.3390/brainsci7050054.

Johansson, S. (2015) Language abilities in Neanderthals. *Ann. Rev. Ling.* 1: 311-332.

Jones, P. A. (2012) Functions of DNA methylation: islands, start sites, gene bodies and beyond. *Nat. Rev. Genet.* 13: 484-92.

Kaufmann, W. E., Cortell, R., Kau, A. S., Bukelis, I., Tierney, E., Gray, R. M., et al. (2004) Autism spectrum disorder in fragile X syndrome: Communication, social interaction, and specific behaviors. *Am J Med Genet A* 129: 225–234

Konopka, G., Bomar, J. M., Winden, K., Coppola, G., Jonsson, Z. O., Gao, F., et al. (2009) Human-specific transcriptional regulation of CNS development genes by FOXP2. *Nature* 462: 213-7.

Koolen, D. A., Pfundt, R., Linda, K., Beunders, G., Veenstra-Knol, H. E., Conta, J. H., et al. (2016) The Koolen-de Vries syndrome: a phenotypic comparison of patients with a 17q21.31 microdeletion versus a KANSL1 sequence variant. *Eur. J. Hum. Genet.* 24: 652–659

Krug, A., Markov, V., Sheldrick, A., Krach, S., Jansen, A., Zerres, K., et al. (2009) The effect of the COMT val(158)met polymorphism on neural correlates of semantic verbal fluency. *Eur. Arch. Psychiatry Clin. Neurosci.* 259: 459-65.

Krug, A., Nieratschker, V., Markov, V., Krach, S., Jansen, A., Zerres, K., et al. (2010) Effect of CACNA1C rs1006737 on neural correlates of verbal fluency in healthy individuals. *Neuroimage* 49, 1831-6.

Kumar, A., Sundaram, S. K., Sivaswamy, L., Behen, M. E., Makki, M. I., Ager, J., et al. (2010). Alterations in frontal lobe tracts and corpus callosum in young children with autism spectrum disorder. *Cereb. Cortex* 20, 2103–13.

Kumar, D., Dedic, N., Flachskamm, C., Voulé, S., Deussing, J. M., and Kimura, M. (2015) Cacna1c (Cav1.2) modulates electroencephalographic rhythm and rapid eye movement sleep recovery. *Sleep* 38: 1371-1380.

Landi, N., Frost, S. J., Mencl, W. E., Preston, J. L., Jacobsen, L. K., et al. (2013) The COMT Val/Met polymorphism is associated with reading-related skills and consistent patterns of functional neural activation. *Dev. Sci.* 16: 13-23.

Lanz, T. A., Guilmette, E., Gosink, M.M., Fischer, J.E., Fitzgerald, L.W., Stephenson, D.T., et al. (2013) Transcriptomic analysis of genetically defined autism candidate genes reveals common mechanisms of action. *Mol. Autism* 4: 45.

Lee, J., Chung, C., Ha, S., Lee, D., Kim, D. Y., Kim, H. et al. (2015) Shank3-mutant mice lacking exon 9 show altered excitation/inhibition balance, enhanced rearing, and spatial memory deficit. *Front. Cell. Neurosci.* 9: 94.

Lev Maor, G., Yearim, A. and Ast, G. (2015) The alternative role of DNA methylation in splicing regulation. *Trends Genet.* 31: 274-80.

Lu, C., Chen, Q., Zhou, T., Bozic, D., Fu, Z., Pan, J. Q., et al. (2016) Micro-electrode array recordings reveal reductions in both excitation and inhibition in cultured cortical neuron networks lacking Shank3. *Mol. Psychiatry* 21:159-68.

Luciano, M., Hansell, N. K., Lahti, J., Davies, G., Medland, S. E., Räikkönen, K., et al. (2011) Whole genome association scan for genetic polymorphisms influencing information processing speed. *Biol. Psychol.* 86, 193–202

Malhotra, A. K., Kestler, L. J., Mazzanti, C., Bates, J. A., Goldberg, T. and Goldman, D. (2002) A functional polymorphism in the COMT gene and performance on a test of prefrontal cognition. *Am. J. Psychiatry* 159: 652- 654

Martinez, I., Rosa, M., Arsuaga, J.-L., Jarabo, P., Quam, R., Lorenzo, C., et al. (2004). Auditory capacities in Middle Pleistocene humans from the Sierra de Atapuerca in Spain. *PNAS* 101: 9976-9981.

Matsunoshita, Y., Ijiri, K., Ishidou, Y., Nagano, S., Yamamoto, T., Nagao, H., et al. (2011) Suppression of osteosarcoma cell invasion by chemotherapy is mediated by urokinase plasminogen activator activity via up-regulation of EGR1. *PLoS One* 6: e16234.

Mefford, H. C., Muhle H., Ostertag P., von Spiczak S., Buysse K., Baker, C., et al. (2010) Genome-wide copy number variation in epilepsy: novel susceptibility loci in idiopathic generalized and focal epilepsies. *PLoS Genet.* 6, e1000962.

Miller, J. L. and Grant, P. A. (2013) The role of DNA methylation and histone modifications in transcriptional regulation in humans. *Subcell Biochem.* 61:289-317.

Mithen, S. (1996). *The Prehistory of the Mind: A Search for the Origins of Art, Science and Religion*. London: Thames & Hudson.

Mithen, S. (2014) The cognition of *homo neanderthalensis* and *H. sapiens*: does the use of pigment necessarily imply symbolic thought? In *Dynamics of Learning in Neanderthals and Modern Humans, vol. II: Cognitive and Physical Perspectives* (Akazawa, T., Ogihara, N., Tanabe, H.C. and Terashima, H., eds). Replacement of Neanderthals by Modern Humans Series . Tokyo: Springer Japan. pp. 7-16.

Mounier, A., Balzeau, A., Caparros, M. and Grimaud-Hervé, D. (2016) Brain, calvarium, cladistics: A new approach to an old question, who are modern humans and Neandertals? *J. Hum. Evol.* 92: 22-36.

Murphy, E. (2015a) Labels, cognomes and cyclic computation: an ethological perspective. *Front. Psychol.* 6: 715.

Murphy, E. (2015b) The brain dynamics of linguistic computation. *Front. Psychol.* 6: 1515.

Murphy, E. (2016a) A theta-gamma neural code for feature set composition with phase-entrained delta nestings. *UCL Working Papers in Linguistics* 28: 1-23.

Murphy, E. (2016b). Evolutionary monkey oscillomics: generating linking hypotheses from preserved brain rhythms. *Theor. Ling.* 42: 117-137.

Murphy, E. and Benítez-Burraco, A. (2016) Bridging the gap between genes and language deficits in schizophrenia: an oscillopathic approach. *Front. Hum. Neurosci.* 10: 422.

Murphy, E., and Benítez-Burraco, A. (2017) The language oscillogenome. BioRxiv doi: https://doi.org/10.1101/114033

Nakamura, T., Matsumoto, J., Takamura, Y., Ishii, Y., Sasahara, M., Ono, T, et al. (2015) Relationships among parvalbumin-immunoreactive neuron density, phase-locked gamma oscillations, and autistic/schizophrenic symptoms in PDGFR-β knock-out and control mice. *PLoS One* 10: e0119258.

Nguyen, P. T., Nakamura, T., Hori, E., Urakawa, S., Uwano, T., Zhao, J., et al. (2011) Cognitive and socio-emotional deficits in platelet-derived growth factor receptor-β gene knockout mice. *PLoS One* 6: e18004.

Oksenberg, N., Haliburton, G. D., Eckalbar, W. L., Oren, I., Nishizaki, S., Murphy, K. et al. (2014) Genome-wide distribution of Auts2 binding localizes with active neurodevelopmental genes. *Transl. Psychiatry* 4, e431.

Pearce, E. and Dunbar, R.I.M. (2012) Latitudinal variation in light levels drives human visual system size. *Biol. Lett.* 8: 90-93.

Pearce, E., Stringer, C. and Dunbar, R. I. M. (2013) New insights into differences in brain organization between Neanderthals and anatomically modern humans. *Proc. Biol. Sci.* 280: 20130168.

Peça, J., Feliciano, C., Ting, J. T., Wang, W., Wells, M. F., et al. (2011) Shank3 mutant mice display autistic-like behaviours and striatal dysfunction. *Nature* 472:437-42.

Ponce de León, M. S., Bienvenu, T., Akazawa, T. and Zollikofer, C. P. (2016) Brain development is similar in Neanderthals and modern humans. *Curr. Biol.* 26: R665-6.

Prüfer, K., Racimo, F., Patterson, N., Jay, F., Sankararaman, S., Sawyer, S., et al. (2014) The complete genome sequence of a Neanderthal from the Altai Mountains. *Nature* 505: 43-9.

Radwan, B., Dvorak, D. and Fenton, A. A. (2016) Impaired cognitive discrimination and discoordination of coupled theta-gamma oscillations in Fmr1 knockout mice. *Neurobiol. Dis.* 88: 125-38.

Rakhade, S. N., Shah, A. K., Agarwal, R., Yao, B., Asano, E., Loeb, J. A. (2007) Activity-dependent gene expression correlates with interictal spiking in human neocortical epilepsy. *Epilepsia* 48: 86-95.

Roll, P., Vernes, S. C., Bruneau, N., Cillario, J., Ponsole-Lenfant, M., Massacrier, A., et al. (2010) Molecular networks implicated in speech-related disorders: FOXP2 regulates the SRPX2/uPAR complex. *Hum. Mol. Genet.* 19: 4848-60.

Schomburg, E. W., Fernandez-Ruiz, A., Mizuseki, K., Berenyi, A., Anastassiou, C. A., Koch, C., et al. (2014) Theta phase segregation of input-specific gamma patterns in entorhinal-hippocampal networks. *Neuron* 84: 470-485.

Smith, L. E., Barker, E. T., Seltzer, M. M., Abbeduto, L. and Greenberg, J. S. (2012) Behavioral phenotype of fragile X syndrome in adolescence and adulthood. *Am. J. Intellect. Dev. Disabil.* 117: 1-17.

Soeiro-De-Souza, M. G., Bio, D. S., David, D. P., Missio, G., Lima, B., Fernandes, F., et al. (2013) Gender effects of the COMT Val 158 Met genotype on verbal fluency in healthy adults. *Mol. Med. Rep.* 8: 837-44

Sugiura, L., Toyota, T., Matsuba-Kurita, H., Iwayama, Y., Mazuka, R., Yoshikawa, T., et al. (2017) Age-dependent effects of Catechol-O-Methyltransferase (COMT) gene Val158Met polymorphism on language function in developing children. *Cereb Cortex* 27: 104-116.

Tattersall, I. (2017). How can we detect when language emerged? *Psychon. Bull. Rev.* 24: 64-67.

Veyrac, A., Besnard, A., Caboche, J., Davis, S. and Laroche, S. (2014) The transcription factor Zif268/Egr1, brain plasticity, and memory. *Prog. Mol. Biol. Transl. Sci.* 122: 89-129.

Vosskuhl, J., Huster, R. J. and Herrmann, C. S. (2015) Increase in short-term memory capacity induced by down-regulating individual theta frequency via transcranial alternating current stimulation. *Front. Hum. Neurosci.* 9: 257.

Weber, G. W. and Bookstein, F. L. (2011) *Virtual Anthropology: A Guide to a New Interdisciplinary Field*. Springer: Wien.

Williams, F. L. and Cofran, Z. (2016) Postnatal craniofacial ontogeny in neandertals and modern humans. *Am. J. Phys. Anthropol.* 159: 394-409.

Wynn, T. and Coolidge, F. L. (2012). *How to Think Like a Neandertal*. Oxford: Oxford University Press

Zollikofer, C. P. and Ponce de León, M. S. (2013) Pandora's growing box: Inferring the evolution and development of hominin brains from endocasts. *Evol. Anthropol.* 22:20-33.