The human auditory brainstem response to running speech reveals a subcortical mechanism for selective attention

Antonio Elia Forte, Octave Etard and Tobias Reichenbach*

Department of Bioengineering and Centre for Neurotechnology, Imperial College London, South Kensington Campus, SW7 2AZ, London, U.K.

*To whom correspondence should be addressed (email: reichenbach@imperial.ac.uk)

Abstract

Humans excel at selectively listening to a target speaker in background noise such as competing voices. While the encoding of speech in the auditory cortex is modulated by selective attention, it remains debated whether such modulation occurs already in subcortical auditory structures. Investigating the contribution of the human brainstem to attention has, in particular, been hindered by the tiny amplitude of the brainstem response. Its measurement normally requires a large number of repetitions of the same short sound stimuli, which may lead to a loss of attention and to neural adaptation. Here we develop a mathematical method to measure the auditory brainstem response to running speech, an acoustic stimulus that does not repeat and that has a high ecological validity. We employ this method to assess the brainstem's activity when a subject listens to one of two competing speakers, and show that the brainstem response is consistently modulated by attention.

Introduction

It is well known that selective attention to one of several competing acoustic signals affects the encoding of sound in the auditory cortex (Shinn-Cunningham 2008; Hackley et al. 1990; Choi et al. 2013; Fritz et al. 2007b; Hillyard et al. 1973; Womelsdorf & Fries 2007; Fritz et al. 2007a; Näätänen et al. 2001). Because extensive auditory centrifugal pathways carry information from central to more peripheral levels of the auditory system (Winer 2006; Pickels 1988; Song et al. 2008; Bajo et al. 2010), neural activity in the subcortical structures may contribute to attention as well. Previous attempts to determine an attentional modulation from recording the auditory brainstem response through scalp electrodes have, however, yielded highly inconclusive results.

In particular, one investigation found that selective attention alters the brainstem's response to the fundamental frequency of a speech signal (Galbraith et al. 1998), while another study concluded that this response is modulated in an unsystematic but subject-specific manner (Lehmann & Schönwiesner 2014) and a third recent experiment did not find a significant attentional effect (Varghese et al. 2015). Results on the effects of attention on the auditory-brainstem response to short clicks or pure tones are similarly inconclusive (Brix 1984; Gregory et al. 1989; Hoormann et al. 2000; Galbraith et al. 2003). These inconsistencies may result from a main experimental limitation in these studies: because the brainstem response is tiny, its measurement requires hundred- to thousandfold repetition of the same sound. The large number of repetitions may lead to difficulties for subjects in sustaining selective attention, to adaptation in the nervous system, and to a reduction in efferent feedback (Lasky 1997; Neupane et al. 2014).

To overcome this limitation, we develop here a method to measure the auditory brainstem's response to natural running speech that does not repeat. We then use this method to assess the modulation of the auditory brainstem response to one of two competing speakers by selective attention.

Results

Assessing the brainstem's response to continuous non-repetitive speech does not allow to average over many repeated presentations of the same sound. Instead, we sought to quantify the brainstem's response to the fundamental frequency of speech. Neuronal activity in the brainstem, and in particular in the inferior colliculus, can indeed phase lock to the periodicity of voiced speech (Skoe & Kraus 2010). The fundamental frequency of running speech varies over time, however, compounding a direct read-out of the evoked brainstem response.

To overcome this difficulty, we employed empirical mode decomposition (EMD) of the speech stimuli to identify an empirical mode that, at each time instance, oscillates at the fundamental frequency of the speech signal (Huang & Pan 2006) (Methods). This mode is a nonlinear and

nonstationary oscillation with a temporally-varying amplitude and frequency that we refer to as the

'fundamental waveform' of the speech stimulus (Figure 1a).

We then recorded the brainstem response to running non-repetitive speech stimuli of several

minutes in duration from human volunteers through scalp electrodes. We cross-correlated the obtained

recording with the fundamental waveform of the speech signal (Figure 1b). Because the brainstem

response may occur at a phase that is different from that of the fundamental waveform, we also

correlated the neural signal to the Hilbert transform of the fundamental waveform that has a phase

delay of 90˚. The two correlations can be viewed as the real and imaginary part of a complex

correlation function that can trace the brainstem response at any phase delay. The amplitude of the

complex correlation informs then on the strength of the brainstem response.

We found that the amplitude of the complex correlation peaked at a mean latency of 9.3 ± 0.7

ms, and our statistical analysis showed that this peak was significantly different from the noise in

fourteen out of sixteen subjects ($p < 0.05$, Methods). The average value of the correlation at the peak

was 0.015 ± 0.003. Moreover, the latency agrees with that found previously regarding the brainstem's

response to short repeated speech stimuli (Skoe & Kraus 2010). We checked that the response does

not contain a stimulus artifact or a contribution from the cochlear microphonic, and that the latency of

the response is not affected by the processing of the speech signal or of the neural response (Methods;

Figure 1–figure supplement 1). This demonstrates that the brainstem's response to continuous speech

can be reliably extracted through the developed method, and the response can be characterized

through the latency and amplitude of the correlation's peak.

Armed with the ability to quantify the brainstem's response to running non-repetitive speech,

we sought to investigate if this neural activity is affected by selective attention. Employing a well-

established paradigm of attention to one of two speakers (Ding & Simon 2012), we presented

volunteers diotically with two concurrent speech streams of equal intensity, one by a male and another

by a female voice. For parts of the speech presentation subjects attended the male voice and ignored

the female voice, and *vice versa* for the remaining parts.

We quantified the brainstem's response to both the male and the female voice by extracting

the fundamental waveforms of both speech signals and correlating the neural recording separately to

both. We found that the latency of the response was unaffected by attention: the response to the

unattended speaker occurred 0.8 ± 0.5 ms later than that to the attended speaker, which was not

statistically significant ($p = 0.2$; average over the responses to the male and the female voice as well

as all subjects).

In contrast, all subjects showed a larger response of the auditory brainstem to the male voice

when attending rather than ignoring it (Figure 2a). The difference in the responses was statistically

significant in nine of the fourteen subjects ($p < 0.05$). The brainstem's response to the attended female

speaker similarly exceeded that to the unattended female voice in all but one subject, with eight

subjects showing a statistically-significant difference ($p < 0.05$; Figure 2b). The ratio of the

brainstem's response to attended and to ignored speech, averaged over all subjects, was 1.5 ± 0.1 and

1.6 ± 0.2 for the male and for the female speaker, respectively. Both ratios were significantly different

from unity ($p < 0.001$, male voice; $p < 0.01$, female voice). The male and the female voice elicited a

comparable attentional modulation: the difference between the corresponding ratios was insignificant

($p = 0.7$). The magnitude of the brainstem's response was hence significantly enhanced through

attention, and consistently so across subjects and speakers.

Discussion

Our results show that the human auditory brainstem response to continuous speech is larger when

attending than when ignoring a speech signal, and consistently so across different subjects and

speakers. In particular, the strength of the phase locking of the neural activity to the pitch structure of

speech is larger for an attended than for an unattended speech stream. In contrast, we did not observe

a difference in the latency of this activity.

The fundamental waveform of speech that we have obtained from EMD has a temporally

varying frequency and amplitude and is therefore not a simple component of Fourier analysis. While

it may be obtained from short-time Fourier transform or wavelet analysis, both methods suffer from

an inherently limited time-frequency resolution that makes them inferior to the EMD analysis (Huang

& Pan 2006).

Because we have employed a diotic stimulus presentation in which the same acoustical

stimulus was presented to each ear, the attentional modulation cannot result from a general

modulation of the brainstem's activity to acoustic stimuli between the two hemispheres. Moreover,

although the fundamental frequencies of the two competing speakers differ at most time points, their

spectra largely overlap. The attentional modulation can therefore not result from a broad-band

modulation of the neural activity either. Instead, the attentional effect must result from a modulation

of the brainstem's response to the specific pitch structure of a speech stimulus.

The brainstem response to the pitch of continuous speech that we have measured can reflect a

response both to the fundamental frequency of speech as well as to higher harmonics. Indeed,

previous studies have found that the brainstem responds at the fundamental frequency of a speech

stimulus even when that frequency itself is removed from the acoustic signal (Galbraith & Doan

1995), or when it cancels out due to presentation of stimuli with opposite polarities and averaging of

the obtained responses (Aiken & Picton 2008). The attentional modulation of the brainstem response

can thus reflect a modulation of the response to the fundamental frequency itself or to higher

harmonics. Moreover, attentional modulation of higher harmonics may depend on frequency as shown

recently in recordings of otoacoustic emissions from the inner ear (Maison et al. 2001).

The attentional modulation of the brainstem's response to the pitch of a speaker may result

from an enhancement of the neural response to an attended speech signal, from the suppression of the

response to an ignored speech stimulus, or from both. Further investigation into this issue may

compare brainstem responses to speech when attending to the acoustical signal and when attending to

a visual stimulus (Woods et al. 1992; Karns & Knight 2009; Saupe et al. 2009).

The response at the fundamental frequency of speech can result from multiple sites in the

brainstem (Chandrasekaran & Kraus 2010). However, we observed a single peak with a width of a

few ms in the correlation of the neural signal to the fundamental waveform of speech. The brainstem

response to running speech that we have measured here can therefore only reflect neural sources

whose latencies vary by a few ms or less from the peak latency. . The neural delay of about 9 ms as

well as the similarity of the speech-evoked brainstem response to the frequency-following response

suggest that the main neural source may be in the inferior colliculus (Sohmer et al. 1977). The

attentional effect that we have observed may then result from the multiple feedback loops between the

inferior colliculus, the medial geniculate body and the auditory cortex (Huffman & Henson 1990).

Our study provides the mathematical tools to analyse the brainstem response to complex, real

world stimuli such as speech. Since our method does not require artificial and repeated stimuli, it

fosters sustained attention and avoids potential neural adaptation. This method can therefore pave the

way to further explore how the brainstem contributes to the processing of complex real-world acoustic

environments. It may also be relevant for better understanding and diagnosing the recently discovered

cochlear neuropathy or 'hidden hearing loss' (Kujawa & Liberman 2009). Because the latter alters the

brainstem's activity (Schaette & McAlpine 2011; Mehraei et al. 2016), assessing the auditory

brainstem response to speech as well as its modulation by attention may further clarify the origin,

prevalence and consequences of such poorly understood supra-threshold hearing loss.

Methods

Participants. 16 healthy adult volunteers aged 18 to 32, eight of which were female, participated in

the study. All subjects were native English speakers and had no history of hearing or neurological

impairments. All participants had pure-tone hearing thresholds better than 20 dB hearing loss in both

ears at octave frequencies between 250 Hz and 8 kHz. Each subject provided written informed

consent. All experimental procedures were approved by the Imperial College Research Ethics

Committee.

Auditory brainstem recordings to running speech. Samples of continuous speech from a male and
a female speaker were obtained from publicly available audiobooks (https://librivox.org). All samples
had a duration of at least two minutes and ten seconds; some were slightly longer to end upon
completion of a sentence. To construct speech samples with two competing speakers, samples from
the male and from the female speaker were normalized to the same root-mean-square amplitude and
then superimposed.

Participants were placed in a comfortable chair in an acoustically and electrically insulated
room (IAC Acoustics, U.K.). A personal computer outside the room controlled audio presentation and
data acquisition. Speech stimuli were presented at a sampling frequency of 44.1 kHz through a high-
performance sound card (Xonar Essence STX, Asus, U.S.A.). Stimuli were delivered diotically
through insert earphones (ER-3C, Etymotic, U.S.A.) at a level of 78 dB(C) SPL (C-weighted
frequency response). Sound intensity was calibrated with an ear simulator (Type 4157, Brüel & Kjaer,
Denmark). All subjects reported that the stimulus level was comfortable.

The response from the auditory brainstem was measured through five passive Ag/AgCl
electrodes (Multitrode, BrainProducts, Germany). Two electrodes were positioned at the cranial
vertex (Cz), two further electrodes were placed on the left and right mastoid processes, and the
remaining electrode was positioned on the forehead to measure the ground. The impedance between
each electrode and the skin was reduced to below 5 kΩ using abrasive electrolyte-gel (Abralyt HiCl,
Easycap, Germany). The electrode on the left mastoid, at the cranial vertex and the ground electrode
were connected to a bipolar amplifier with low-level noise and a gain of 50 (EP-PreAmp,
BrainProducts, Germany). The remaining two electrodes were connected to a second identical bipolar
amplifier. The output from both bipolar amplifiers was fed into an integrated amplifier (actiCHamp,
BrainProducts, Germany) where it was low-pass filtered through a hardware anti-aliasing filter with a
corner frequency of 4.9 kHz and sampled at 25 kHz. The audio signals were measured by the
integrated amplifier as well through an acoustic adapter (Acoustical Stimulator Adapter and StimTrak,
BrainProducts, Germany). The electrophysiological data were acquired through PyCorder
(BrainProducts, Germany). The simultaneous measurement of the audio signal and the brainstem
response from the integrated amplifier was employed to temporally align both signals to a precision of
less than 40 μs, the inverse of the sampling rate (25 kHz).

Experimental design. In the first part of the experiment, each volunteer listened to four speech
samples of the female speaker only. Comprehension questions were asked at the end of each part in
order to verify the subject's attention to the story.

The second part of the experiment employed eight samples of speech that contained both a
male and a female voice. During the presentation of the first four samples, subjects were asked to
attend either the male or the female speaker. Volunteers were then presented with the next four speech

samples and asked to attend to the speaker that they had ignored earlier. Whether the subject was

asked to attend first to the male or to the female voice was determined randomly for every subject.

Comprehension questions were asked after each sample.

Computation of the fundamental waveform of speech. The fundamental waveform of each speech

sample with a single speaker was computed through a custom-written Matlab program (code available

on Github; Forte 2017). The fundamental waveform of a speech sample with two speakers followed

from the two corresponding samples with a single speaker only.

First, each speech signal was downsampled to 8,820 Hz, low-pass filtered at 1,500 Hz (FIR,

transition band 1,500 – 1,650 Hz, stopband attenuation -80 dB, passband ripple 1 dB, order 296) and

time-shifted to compensate for the filter delay. Silent parts between words were identified by

computing the envelope of the speech signal. Each part where the envelope was less than 10% of the

maximal value found in the speech was considered silent, and the speech signal there was set to zero.

Second, the instantaneous fundamental frequency of the voiced parts of the speech signal was

detected through the autocorrelation method, employing rectangular windows of 50 ms duration with

a successive overlap of 49 ms. Speech segments that yielded a fundamental frequency outside the

range of 60 Hz to 400 Hz, or in which the fundamental frequency varied by more than 10 Hz between

two successive windows were considered voiceless. The speech segments that corresponded to voiced

speech, as well as their fundamental frequency, were thus obtained. The fundamental frequency of

each segment was interpolated through a cubic spline, and varied between 100 and 300 Hz in each

segment. Note that this method yields the fundamental frequency but not by itself the fundamental

wavemode.

Third, the voiced speech segments where analysed through the Hilbert-Huang transform. The

latter is an adaptive signal processing based on empirical basis functions and can thus be better suited

for analysing nonlinear and nonstationary signals such as speech than Fourier analysis (Huang & Pan

2006). The transform consists of two parts. First, empirical mode decomposition extracts intrinsic

mode functions (IMFs) that satisfy two properties: (i) the numbers of extrema and zero crossings are

either equal or differ by one; (ii) the mean of the upper and lower envelope vanishes. The signal

follows as the linear superposition of the IMFs. Second, the Hilbert spectrum of each IMF is

determined, which yields, in particular, the mode's instantaneous frequency. This analysis was

performed for each short segment of voiced speech, that is, for each part of voiced speech that was

preceded and followed by a pause or voiceless speech.

Fourth, the fundamental frequency of each short speech segment was compared to the

instantaneous frequencies of the segment's IMFs at each individual time point. All IMFs with an

instantaneous frequency that differed by less than 20% from the segment's fundamental frequency

were determined, and the IMF with the largest amplitude was therefrom selected as the fundamental

wavemode of that segment and at that time point (Huang & Pan 2006). If no IMF had an
instantaneous frequency within 20% of the fundamental frequency, or if a speech segment was
unvoiced, that time point was assigned a fundamental waveform of zero. The fundamental waveforms
obtained at the different time points were combined through cosine crossfading functions with a
window width of 10 ms to obtain the fundamental waveform of the speech signal. The Hilbert
transform of that fundamental waveform was computed as well.

To control for latency changes in the acoustic signal induced by the subsequent processing
steps, and in particular by the involved frequency filtering, the cross-correlation between the original
speech signal and the fundamental waveform as well as with its Hilbert transform was computed
(Figure 1–figure supplement 1**a**). The cross-correlations show that the fundamental waveform has no
latency change and no phase difference with respect to the original speech stimulus.

Analysis of the auditory-brainstem response. The brainstem responses from the two measurement
channels were averaged. A frequency-domain regression technique (CleanLine, EEGLAB) was used
to attenuate noise from the power line in the brainstem recording. Moreover, because a voltage
amplitude above 20 mV cannot result from the brainstem but represents artefacts such as spurious
muscle activity, the signal was set to zero during episodes of such high voltage. The
electrophysiological recording was then filtered between $100 - 300$ Hz since the fundamental
frequency of the speech was in that range (high-pass filter: FIR, transition band from $90 - 100$ Hz,
stopband attenuation -80 dB, passband ripple 1 dB, order 6862; low-pass filter: FIR, transition band
$300 - 360$ Hz, stopband attenuation -80 dB, passband ripple 1 dB, order 1054). In particular, the high-
pass filter eliminated neural signals from the cerebral cortex that occur predominantly below 100 Hz.
To avoid transient activity at the beginning of each speech sample, the first ten seconds of each
brainstem recording in response to a speech sample were discarded. The following two minutes of
data were divided into 40 epochs of a duration of 3 s each, and the remaining data were discarded, if
any.

The processing of the neural signal did not induce a latency. This was confirmed by
computing the cross-correlation between the processed neural response and the original signal,
demonstrating a maximum correlation at zero temporal delay (Figure 1–figure supplement 1**b**).

As set out above, the first part of the experiment measured the brainstem response to running
speech without background noise. For each subject and each epoch, the cross-correlation of the
brainstem response with the corresponding segment of the fundamental waveform as well as with its
Hilbert transform were computed. A delay of 1 ms of the acoustic signal produced by the earphones
was taken into account. The two cross-correlation functions were interpreted as the real and the
imaginary part of a complex correlation function. For each individual subject, the average of the

complex cross-correlation over all epochs was then computed, and the latency at which the amplitude

peaked was determined.

The obtained latencies of about 9 ms affirmed that the signal resulted from the auditory

brainstem and not from the cerebral cortex, whose latencies exceed 20 ms. The latency also evidenced

that the signal resulted neither from stimulus artifacts nor from the cochlear microphonic, which

would occur at or near zero delay (Skoe & Kraus 2010). As an additional control, the brainstem

response was recorded when the earphones were near the ear, but not inserted into the ear canal, so

the subject could not hear the speech signals. The recording did then not yield a measurable brainstem

response (Figure 1–figure supplement 1**c**). Two presentations of the same speech stimulus, but with

opposite polarities, were employed as well, and the neural response to both presentations was

averaged before computing the correlation to the fundamental waveform. The correlation was

identical to that obtained by a single stimulus presentation, demonstrating the absence of a stimulus

artifact and of the cochlear microphonic (Figure 1–figure supplement 1**d**).

To determine whether the peak in the cross-correlation obtained from a given subject was

significant, the values of the complex cross-correlation from the individual epochs, and at the peak

latency, were analysed. Because each correlation value is an average of many measurements, it

follows from the Central Limit Theorem that the complex correlations from the different epochs

exhibit a two-dimensional normal distribution with a mean of zero if the measurements are randomly

distributed. A one-sample Hotelling's T-squared test was therefore used to assess the significance of

the complex correlation at the peak latency. Two subjects who did not show a significant correlation

($p > 0.05$) were not included in the further analysis.

The population mean and standard error of the mean of the latency were computed from the

latencies of the individual subjects.

The brainstem responses to competing speakers were then analysed for each individual

subject. For each epoch, the complex cross-correlation between the brainstem response and the

fundamental waveform was computed, both for the fundamental waveform of the attended and for

that of the unattended speaker. The corresponding complex correlation functions were averaged

across epochs, and the amplitudes as well as latencies of the peaks were determined.

Statistical significance of the difference in latency of the brainstem responses to the attended

and the unattended speaker, obtained from the eight samples, was tested by computing population

mean as well as standard error of the mean for the differences in latencies obtained from individual

subjects. A two-tailed Student's t-test was employed to test if the difference was significantly different

from zero.

To control for differences in the voice of the male and the female speaker, differences in
amplitude of the brainstem response to the attended and ignored male speaker were determined
separately from differences in the amplitude of the brainstem response to the attended and ignored
female speaker. The amplitudes of the complex cross-correlations, at the peak latencies, were
computed for all epochs. A two-sample Student's t-test was then employed to test for a significant
difference between the amplitude in response to the attended and the ignored speaker.

The amplitude of the brainstem response to speech can vary widely between subjects (Figure 2), due
to variations such as in anatomy and scalp conductivity. The ratios of the amplitudes of the brainstem
responses to attended and ignored speech, rather than the differences, were thus computed for each
individual. The population mean and standard error of the mean were therefrom obtained. A one-
tailed Student's t-test assessed whether the population average of the ratio was significantly larger
than unity. A two-tailed two-sample Student's t-test was employed to assess whether the ratios
obtained from the responses to the male and to the female speaker were significantly different.

Acknowledgement

We thank Steve Bell, Karolina Kluk-de Kort, Patrick Naylor, David Simpson and Malcolm Slaney for
discussion as well as for comments on the manuscript. This research was supported by EPSRC grant
EP/M026728/1 to T.R. as well as in part by the National Science Foundation under Grant No. NSF
PHY-1125915.

Competing financial interests

The authors declare no competing financial interests.

References

Aiken, S.J. & Picton, T.W., 2008. Envelope and spectral frequency-following responses to vowel
sounds. *Hear. Res.*, 245(1–2), pp.35–47.

Bajo, V.M., Nodal, F.R., Moore, D.R. & King, A.J., 2010. The descending corticocollicular pathway
mediates learning-induced auditory plasticity. *Nat. Neurosci.*, 13(2), pp.253–260.

Brix, R., 1984. The influence of attention on the auditory brain stem evoked responses preliminary
report. *Acta Otolaryngol.*, 98(1–2), pp.89–92.

Chandrasekaran, B. & Kraus, N., 2010. The scalp-recorded brainstem response to speech: neural

origins and plasticity. *Psychophysiology*, 47(2), pp.236–246.

Choi, I., Rajaram, S., Varghese, L.A. & Shinn-Cunningham, B.G., 2013. Quantifying attentional
modulation of auditory-evoked cortical responses from single-trial electroencephalography.
Front. Hum. Neurosci., 7, p.115.

Ding, N. & Simon, J.Z., 2012. Emergence of neural encoding of auditory objects while listening to
competing speakers. *Proc. Natl. Acad. Sci. U. S. A.*, 109(29), pp.11854–9.

Forte, A.E., 2017. Fundamental_waveforms_extraction, GitHub,
https://github.com/antn85/fundamental_waveforms_extraction, version 1.

Fritz, J.B., Elhilali, M., David, S. V. & Shamma, S.A., 2007a. Auditory attention - focusing the
searchlight on sound. *Curr. Opin. Neurobiol.*, 17(4), pp.437–455.

Fritz, J.B., Elhilali, M., David, S. V. & Shamma, S.A., 2007b. Does attention play a role in dynamic
receptive field adaptation to changing acoustic salience in a1? *Hear. Res.*, 229(1–2), pp.186–
203.

Galbraith, G.C., Bhuta, S.M., Choate, A.K., Kitahara, J.M. & Mullen, T.A., 1998. Brain stem
frequency-following response to dichotic vowels during attention. *Neuroreport*, 9(8), pp.1889–
1893.

Galbraith, G.C. & Doan, B.Q., 1995. Brainstem frequency-following and behavioral responses during
selective attention to pure tone and missing fundamental stimuli. *Int. J. Psychophysiol.*, 19(3),
pp.203–214.

Galbraith, G.C., Olfman, D.M. & Huffman, T.M., 2003. Selective attention affects human brain stem
frequency-following response. *Neuroreport*, 14(5), pp.735–8.

Gregory, S.D., Heath, J.A. & Rosenberg, M.E., 1989. Does selective attention influence the brain-
stem auditory evoked potential? *Electroencephalogr. Clin. Neurophysiol.*, 73(6), pp.557–60.

Hackley, S.A., Woldorff, M. & Hillyard, S.A., 1990. Cross-modal selective attention effects on
retinal, myogenic, brainstem, and cerebral evoked potentials. *Psychophysiology*, 27(2), pp.195–
208.

Hillyard, S.A., Hink, R.F., Schwent, V.L. & Picton, T.W., 1973. Electrical signs of selective attention
in the human brain. *Science*, 182(October), pp.177–180.

Hoormann, J., Falkenstein, M. & Hohnsbein, J., 2000. Early attention effects in human auditory-
evoked potentials. *Psychophysiology*, 37(1), pp.29–42.

Huang, H. & Pan, J., 2006. Speech pitch determination based on hilbert-huang transform. *Signal*

Process., 86(4), pp.792–803.

Huffman, R.F. & Henson, O.W., 1990. The descending auditory pathway and acousticomotor
systems: connections with the inferior colliculus. *Brain Res. Rev.*, 15(3), pp.295–323.

Karns, C.M. & Knight, R.T., 2009. Intermodal auditory, visual, and tactile attention modulates early
stages of neural processing. *J. Cogn. Neurosci.*, 21(4), pp.669–83.

Kujawa, S.G. & Liberman, M.C., 2009. Adding insult to injurt: cochlear nerve degeneration after
"temporary" noise- induced hearing loss. *J. Neurosci.*, 29(45), p.14077-.

Lasky, R.E., 1997. Rate and adaptation effects on the auditory evoked brainstem response in human
newborns and adults. *Hear. Res.*, 111(1–2), pp.165–176.

Lehmann, A. & Schönwiesner, M., 2014. Selective attention modulates human auditory brainstem
responses: relative contributions of frequency and spatial cues. *PLoS One*, 9(1), pp.1–10.

Maison, S., Micheyl, C. & Collet, L., 2001. Influence of focused auditory attention on cochlear
activity in humans. *Psychophysiology*, 38(1), pp.35–40.

Mehraei, G., Hickox, A.E., Bharadwaj, H.M., Goldberg, H., Verhulst, S., Liberman, M.C., Barbara,
X. & Shinn-Cunningham, G., 2016. Auditory brainstem response latency in noise as a marker of
cochlear synaptopathy. *J. Neurosci.*, 36(13), pp.3755–3764.

Näätänen, R., Tervaniemi, M., Sussman, E., Paavilainen, P. & Winkler, I., 2001. "Primitive
intelligence" in the auditory cortex. *Trends Neurosci.*, 24(5), pp.283–288.

Neupane, A.K., Gururaj, K., Mehta, G. & Sinha, S.K., 2014. Effect of repetition rate on speech
evoked auditory brainstem response in younger and middle aged individuals. *Audiol. Res.*, 4(1),
pp.21–27.

Pickels, J.O., 1988. *An introduction to the physiology of hearing*, Emerald.

Saupe, K., Widmann, A., Bendixen, A., Müller, M.M. & Schröger, E., 2009. Effects of intermodal
attention on the auditory steady-state response and the event-related potential.
Psychophysiology, 46(2), pp.321–327.

Schaette, R. & McAlpine, D., 2011. Tinnitus with a normal audiogram: physiological evidence for
hidden hearing loss and computational model. *J. Neurosci.*, 31(38), pp.13452–13457.

Shinn-Cunningham, B.G., 2008. Object-based auditory and visual attention. *Trends Cogn. Sci.*,
12(April), pp.182–186.

Skoe, E. & Kraus, N., 2010. Auditory brain stem response to complex sounds: a tutorial. *Ear Hear.*,
31(3), pp.302–324.

Sohmer, H., Pratt, H. & Kinarti, R., 1977. Sources of frequency following responses (ffr) in man.
Electroencephalogr. Clin. Neurophysiol., 42(5), pp.656–664.

Song, J.H., Skoe, E., Wong, P.C.M. & Kraus, N., 2008. Plasticity in the adult human auditory
brainstem following short-term linguistic training. *J. Cogn. Neurosci.*, 20(10), pp.1892–902.

Varghese, L., Bharadwaj, H.M. & Shinn-Cunningham, B.G., 2015. Evidence against attentional state
modulating scalp-recorded auditory brainstem steady-state responses. *Brain Res.*, 1626, pp.146–
164.

Winer, J.A., 2006. Decoding the auditory corticofugal systems. *Hear. Res.*, 212(1–2), pp.1–8.

Womelsdorf, T. & Fries, P., 2007. The role of neuronal synchronization in selective attention. *Curr.*
Opin. Neurobiol., 17(2), pp.154–160.

Woods, D.L., Alho, K. & Algazi, A., 1992. Intermodal selective attention. i. effects on event-related
potentials to lateralized auditory and visual stimuli. *Electroencephalogr. Clin. Neurophysiol.*,
82(5), pp.341–355.

Figures

Figure 1 The brainstem response to running speech. (**a**) Speech (black) contains voiced parts with
irregular oscillations at a time-varying fundamental frequency and higher harmonics. We extract a
fundamental waveform (red) that oscillates nonlinearly at the fundamental frequency. (**b**) The
autocorrelation of the fundamental waveform (red) peaks when the delay vanishes and oscillates at the
average fundamental frequency. The cross-correlation of the fundamental waveform with its Hilbert
transform (blue) can be seen as an imaginary part of the autocorrelation. The amplitude of the
resulting complex cross-correlation (black) shows a life-time of a few ms. (**c**) The correlation of the
speech-evoked brainstem response, recorded from one subject, to the fundamental waveform of the
speech signal (red) as well as to its Hilbert transform (blue) can serve as real and imaginary parts of a
complex correlation function. Its amplitude (black) peaks at a latency of 9 ms. The latency of the
correlation is not altered by the processing of the speech signal or of the neural recording, and
contains neither a stimulus artifact nor the cochlear microphonic (**Figure 1–figure supplement 1**).

Figure 1–figure supplement 1. Controls for latencies induced by signal processing as well as for the
source of the measured brainstem response to running speech. (**a**) The cross-correlation between the
original speech signal with the fundamental waveform (red) as well as with its Hilbert transform
(blue) and the resulting amplitude (black) show a peak at 0 ms and no phase delay. The processing of
the acoustic signal does accordingly not change the latency or phase of that signal. (**b**) The
computation of the cross-correlation of the fundamental waveform to the neural recording involved
processing of the neural signal such as through filtering. However, the cross-correlation between the
recorded neural signal and the filtered version shows a peak at vanishing latency. The processing of
the neural signal did therefore not alter the latency. (**c**) When the earphones are placed close to the
ears, but not inside the ear canal, preventing a subject from hearing the speech signal, the cross-
correlation between the recorded neural signal and the fundamental waveform of speech (red) as well
as its Hilbert transform (blue) do not yield a measurable peak. The amplitude of the resulting complex
correlation function (black) does not peak either, demonstrating the absence of a stimulus artifact. (**d**)

When a subject listened to a speech signal and then to the same signal with reversed polarity, and
when the average over the neural recordings to both stimulus presentations was employed for the
analysis, the complex cross-correlation showed the same structure as when it was computed using the
neural response to one stimulus only. This shows the absence of a stimulus artifact as well as the
absence of the cochlear microphonic in the measured response. To enable comparison, all recordings
were obtained from the same subject for whom we report the exemplary recording in Figure 1 (**c**).

Figure 2 Modulation of the brainstem response to speech by selective attention. (**a**) The brainstem's
response to the male speaker is larger for each subject when attending the speaker (dark blue) than
when ignoring it (light blue). The average ratio of the brainstem responses to the attended and to the
ignored male speaker is significantly larger than 1 (black, mean and standard error of the mean). (**b**)
With the exception of subject 13, the neural response to the female voice is also larger when subjects
attend to it (dark red) instead of ignoring it (light red). The average ratio of the brainstem responses to
the attended and to the ignored female speaker is significantly larger than 1 as well (black, mean and
standard error of the mean).

453