1 **Ancient genomic variation underlies recent and repeated ecological adaptation**
2
3

4 Thomas C. Nelson[1] and William A. Cresko[1]*
5
6 [1]Institute of Ecology and Evolution, University of Oregon, Eugene, OR
7 *To whom correspondence should be addressed: wcresko@uoregon.edu
8

Adaptation in the wild often involves the use of standing genetic variation (SGV), allowing rapid responses to selection on ecological timescales. Despite increasing documentation of evolutionarily important SGV in natural populations, we still know little about how the genetic and genomic structure and molecular evolutionary history of SGV relate to adaptation. Here, we address this knowledge gap using the threespine stickleback fish (*Gasterosteus aculeatus*) as a model. We demonstrate that adaptive genetic variation is structured genome-wide into distinct marine and freshwater haplogroups. This divergent variation averages six million years old, nearly twice the genome-wide average, but has been evolving over the 15-million-year history of the species. Divergent marine and freshwater genomes maintain regions of ancient ancestry that include multiple chromosomal inversions and extensive linked variation. These discoveries about ancient SGV demonstrate the intertwined nature of selection on ecological timescales and genome evolution over geological timescales.

The mode and tempo of adaptive evolution depend on the sources of genetic variation affecting fitness[1,2]. While new mutation is ultimately the source of all genetic variation, recent studies of adaptation in the wild document adaptive genetic variation that was either segregating in the ancestral population as standing genetic variation (SGV)[3-5] or introgressed from a separate population or species[6,7]. The use of SGV appears particularly important when dramatic responses to selection occur on ecological timescales, in dozens of generations or fewer[3]. When environments change rapidly, SGV can propel rapid evolution in ecologically relevant traits even in populations of long-lived organisms like Darwin's finches[8], monkeyflowers[9], and threespine stickleback fish[10].

Existing genetic variants have evolutionary histories that are often unknown but that may have significant impacts on subsequent adaptation[9,11]. The abundance, genomic distribution, and fitness effects[10-14] of SGV are themselves products of evolution, and their unknown history raises fascinating questions for the genetics of adaptation in the wild. When did adaptive variants originally arise? How are they structured, across both geography and the genome? Which evolutionary forces shaped

their distribution? And how does this evolutionary history of SGV potentially channel

future evolutionary change?

Answers to these questions are critical for our understanding of the importance of

SGV in nature and our ability to predict the paths available to adaptation on ecological

timescales[9]. Biologists are beginning to probe evolutionary histories of SGV using

genome-wide sequence variation across multiple individuals in numerous populations[15],

but this level of inference has been unavailable for most natural systems because of

methodological limitations that remove phase information (e.g. pool-seq[16]) or produce

very short reads (e.g. RAD-seq[17]). Here, we investigate the structure and evolutionary

history of divergent SGV by implementing a novel haplotyping method based on

restriction site-associated DNA sequencing (RAD-seq). This approach creates nearly

1kb haplotypes at thousands of densely sampled loci, allowing us to accurately

measure sequence variation and estimate divergence times across the genome.

SGV is likely critical to adaptation in this species. Marine stickleback have

repeatedly colonized freshwater lakes and streams[18,19], and adaptive divergence in

isolated freshwater habitats is highly parallel at the phenotypic[20,21] and genomic

levels[19,22] (but see Stuart et al.[23]). In addition, analyses of haplotype variation at the

genes *eda*[10,24] and *atp1a1*[24] present two clear results: separate freshwater populations

share common 'freshwater' haplotypes that are identical-by-descent (IBD), and

sequence divergence between the major marine and freshwater haplogroups suggests

their ancient origins, perhaps over two million years ago in the case of *eda*[10]. While

intriguing, it is not clear whether the deep evolutionary histories of these loci are outliers

or representative of more widespread ancient history across the genome. To address

this fundamental question we utilize the new haplotype RAD-seq approach to assay

genome-wide variation associated with adaptive divergence in two young freshwater

ponds, which formed during the end- the end-Pleistocene glacial retreat (c. 12,000

years ago[21,25]; Fig. 1). Our results demonstrate a suite of adaptive variation structured

into distinct marine and freshwater haplotypes that evolved over millions of years.

Figure 1. Stickleback sampling and RAD sequencing to measure haplotype variation. A) Threespine stickleback sampling locations in this study. Colors represent habitat type: red: marine; blue: freshwater. B-D: We modified the original RAD-seq protocol to generate local haplotypes. Colored bars represent polymorphic sites. For a detailed description of haplotype construction, see *Methods*. B) Overlapping paired-end reads are anchored to *PstI* restriction sites. C) Paired reads mapping to each halfsite are merged into contigs. Contigs mapping to the same restriction site are identified by alignment to the reference genome. D) Sequences from each half of a restriction site are phased to generate a single RAD locus.

RESULTS AND DISCUSSION

Parallel adaptation to freshwater environments has been a major theme of

stickleback evolutionary history[26]. Stereotypical morphological changes (e.g. bony

armor[20] and craniofacial structures[27]) presumably reflect adaptation to similar selective

regimes[28,29] and are accompanied by parallel genomic divergence[19,22], which involves

large regions spanning many megabases[24,30], including multiple chromosomal

inversions[19]. The leading hypothesis for the genetics of parallel divergence in

stickleback posits that distinct freshwater-adaptive haplotypes that are identical-by-

descent (IBD) are shared among fresh water populations due to historical gene flow

between marine and freshwater populations[30]. To test for the presence of these

haplotypes directly, we characterized the genomic architecture and evolutionary history

of SGV by modifying the RAD-seq protocol[31] to generate phased haplotypes similar in

length to Sanger sequencing reads, each anchored to tens of thousands of *PstI*

restriction sites spread across the genome (Fig. 1.B-D). We sampled five fish (10

haploid genomes) each from Boot Lake (BL) and Rabbit Slough (RS), and four fish (8

genomes) from Bear Paw Lake (BP). After stringent data filtering (see Methods), this

resulted in a dataset of 57,992 RAD loci (locus = two tags representing one cut site)

with 694 potential variable sites per locus and a median of seven segregating sites per

locus (range: 2-155, Suppl. Fig. 1, Suppl. Table 1). We then used these phased

haplotypes to estimate genealogies at each RAD locus. By including haplotypes from all

three populations in these genealogical analyses, we were able to jointly calculate

population genetic statistics (F_{ST}, π, d_{XY}), estimate the degree of lineage sorting within

populations, and identify patterns of IBD among populations.

We find that indeed, parallel population genomic divergence in each freshwater

site consistently involved haplotypes that were IBD among both freshwater populations

(Fig. 2). Background F_{ST} between populations ranged from 0.139-0.226, with

divergence between the freshwater populations BL and BP being highest ($F_{ST(RS-BL)}$ =

0.139, $F_{ST(RS-BP)}$ = 0.194, $F_{ST(BL-BP)}$ = 0.226; two-sided Mann-Whitney test for all pairwise

comparisons: $p \leq 1\times10^{-10}$). The degree and genomic distribution of pairwise F_{ST} between

the BL, BP, and RS populations were similar to those previously reported[22], including

marine-freshwater F_{ST} outlier regions on chromosome 4 over a broad span in which the

eda gene is embedded (orange triangle in Fig. 2A) and three regions now known to be

associated with chromosomal inversions on chromosomes 1, 11, and 21 (yellow bars in

Fig. 2; hereafter referred to as *inv1*, *inv11*, and *inv21*). The gene *atp1a1* (green triangle

in Fig. 2A) is contained within *inv1*. As expected, we found distinct haplogroups

associated with marine and freshwater habitats at both *eda* and *atp1a1* (Fig. 3, insets).

Strikingly, this finding of habitat specific haplogroups was not at all unique to

these well studied genes or chromosomal inversions. The two isolated freshwater

populations shared IBD haplotypes within all common marine-freshwater F_{ST} peaks

even though IBD was rare elsewhere (Fig. 2B). Furthermore, we observed a separate

clade of haplotypes representing the marine RS population at the majority (1129 of

Figure 2. The genealogical structure of parallel genomic divergence. A) Genome-wide F_{ST} for both marine-freshwater comparisons was kernel-smoothed using a normally distributed kernel with a window size of 500 kb. Inverted triangles indicate the locations of two genes known to show extensive marine-freshwater haplotype divergence, *Eda* and *Atp1a1*. Three chromosomal inversions are highlighted in yellow. B) Lineage sorting patterns were identified from maximum clade credibility trees for each RAD locus. Blue bars: haplotypes from both freshwater populations form a single monophyletic group; red: haplotypes from the marine population form a monophyletic group; black: A RAD locus is structured into reciprocally monophyletic marine and freshwater haplogroups.

2172, 52%) of RAD loci showing freshwater IBD. The result was a genome-wide pattern

of reciprocal monophyly between marine and freshwater haplotypes. Notably, this is the

same genealogical structure previously reported at *eda* [10,24] and *atp1a1* [24],

demonstrating that these loci are but a small part of a genome-wide suite of genetic

variation sharing similar habitat-specific evolutionary histories, and the previous

documentation of their genealogies was a harbinger of a much more extensive pattern

across the genome revealed here. Hereafter, we refer collectively to this class of RAD

loci as 'divergent loci'.

Because the genealogical structure of divergence across the genome mirrors

that at *eda* and *atp1a1*, we asked whether levels of sequence variation and divergence

also showed consistent genomic patterns. At all RAD loci we therefore calculated π

within each population, as well as in the combined freshwater populations, and d_{XY}

between marine and freshwater habitat types. Genome-wide diversity was similar

across populations and habitat types (m ean π_{RS} = 0.0032, π_{BL} = 0.0034, π_{BP} = 0.0026,

π_{FW} = 0.0038) and comparable to previous estimates[22]. Likewise, genome-wide d_{XY}

among habitat types was modest (0.0049)

when compared to π across all

populations (π = 0.0042, two-sided Mann-

Whitney test: $p \le 1\times10^{-10}$). Among

divergent loci, however, we observed

reductions in diversity in both habitats

(mean $\pi_{RS-divergent}$ = 0.0012, $\pi_{RS-divergent}$ =

0.0016, two-sided permutation test: $p \le$

1×10^{-4}, Fig. 3, Suppl. Fig. 2), indicating

natural selection in both habitats.

Sequence divergence associated with

reciprocal monophyly was striking,

however, averaging nearly three times the

genome-wide mean (mean $d_{XY-divergent}$ =

0.0124). This divergence ranged more

than an order of magnitude (0.0013–

0.0442), from substantially lower than the

genome-wide average to ten times greater

than the average. These findings indicate

that much of the genetic variation

underlying adaptive divergence was not

just standing and structured by habitat,

but has been segregating and

accumulating for millennia.

These data clearly support the

hypothesis of Schluter and Conte[30] of

ancient haplotypes 'transported' among

Figure 3. Extensive sequence divergence between marine and freshwater haplogroups accompanies reciprocal monophyly. For each reciprocally monophyletic RAD locus, we calculated sequence variation (π) within and sequence divergence between habitat types (d_{XY}). Each RAD locus is shown as a pair of lines connecting estimates of π and d_{XY}. Boxplots show distributions across all reciprocally monophyletic RAD loci: Boxes are upper and lower quartiles, including the median; whiskers extend to 1.5x interquartile range. Dashed lines are the genome-wide medians. Single RAD loci from within the transcribed regions of *Eda* and *Atp1a1* are shown as gold and green lines, respectively, and presented as haplotype networks. Dots represent mutational steps. Circle sizes indicate the number of haplotypes and colors indicate population of origin as in Figure 1. Each network = 29 haplotypes.

freshwater populations. Much of the divergence we observed was ancient in origin, with

levels of sequence divergence at some RAD loci exceeding that observed at *eda* (Fig.

3, gold line) and suggestive of divergence times of at least two million years ago[10]. Our

observation that sequence variation was consistently reduced in both habitat types

emphasizes that alternative haplotypes at these loci are likely selected for in the marine

population as well as the freshwater. These alternative fitness optima — driven by

different ecologies — provide a favorable landscape for the maintenance of

variation[32,33], but also lead to a more potent barrier to gene flow among freshwater

populations if there are fitness consequences in the marine habitat for stickleback

carrying freshwater-adaptive variation. Conditional fitness effects through genetic

interactions (e.g., dominance[34] or epistasis[35]) and genotype-by-habitat interactions[36]

could potentially extend the residence time of freshwater haplotypes in the marine

habitat. Future work should consider the phenotypic effects of divergently adaptive

variation in different external environments[36,37].

A steady accumulation of divergently adaptive variation between marine and

freshwater stickleback genomes may also have been critical to the rapid divergence in

the young pond populations we study here. We found reciprocal monophyly associated

with a spectrum of sequence divergence, including a substantial fraction of divergent

loci (11.0%, 124/1129) with d_{XY} below the genome-wide average. Thus, ongoing

marine-freshwater ecological divergence has yielded continuing marine-freshwater

genomic divergence. Moreover, while this younger variation is shared between the

freshwater populations in this study, and localizes to genomic regions of divergence

shared globally[19], some adaptive variants may be distributed only locally (e.g. to

southern Alaska or the eastern Pacific basin). In addition to the globally distributed suite

of variation, there may also exist a substantial amount of regional variation contributing

to stickleback genomic and phenotypic diversity.

Sequence divergence provides an important relative evolutionary timescale.

However, to more directly compare the timescales of ecological adaptation and genomic

evolution, we translated patterns of sequence variation into the time to the most recent

common ancestor (T_{MRCA}) of allelic variation, in years. To do so, we performed a *de*

novo genome assembly of the ninespine stickleback (*Pungitius pungitius*), a member of

the Gasterosteidae that diverged from the threespine stickleback lineage approximately

15 million years ago[38] (Fig. 4A, Suppl. Table 2). We then aligned our RAD dataset to

Figure 4. Marine-freshwater divergence has evolved over millions of years, affecting large genomic regions. We performed Bayesian estimation of the time to the most recent common ancestor (T_{MRCA}) of alleles at threespine stickleback RAD loci. We calibrated coalescence times within threespine stickleback by including a *de novo* genome assembly from the ninespine stickleback (*Pungitius pungitius*) and setting threespine-ninespine divergence at 15 million years ago. A) Maximum clade credibility RAD gene tree representative of the genome-wide average T_{MRCA}. Branches within threespine are colored by population of origin. B) Kernel-smoothed densities of T_{MRCA} distributions for all RAD loci containing a monophyletic group of threespine stickleback alleles (light gray) and those structured into reciprocally monophyletic marine and freshwater haplogroups. C) The genomic distribution of reciprocally monophyletic RAD loci (black, as in Figure 2) is associated with increased T_{MRCA} at a genomic scale. T_{MRCA} outlier windows (those exceeding 99.9% of permuted genomic windows) are shown as gray bars. Genome-wide T_{MRCA} was kernel-smoothed using a normally distributed kernel with a window size of 500 kb. Inverted triangles indicate the locations of *Eda* and *Atp1a1*. Three chromosomal inversions are highlighted in yellow.

this assembly and estimated gene trees for each alignment with BEAST[39], setting

divergence to the ninespine stickleback at 15 MYA (see Methods).

We find that the divergence of key marine and freshwater haplotypes has been

ongoing for millions of years and extends back to the split with the ninespine stickleback

lineage (Fig. 4B). Genome-wide variation averaged 4.1 MY old, and T_{MRCA} for the vast

majority of RAD loci was under 5 MY old. In contrast, divergent loci averaged 6.4 MY

old and, amazingly, the most ancient 10% (118 of 1129 loci) are estimated at over 10

MY old. This deep genomic divergence not only underscores that the marine-freshwater

transition has been occurring throughout the history of the threespine stickleback

lineage, for which there is evidence in the fossil record going back 10 million years[40],

but it also demonstrates that at least some of the variation fueling those ancient events

has persisted until the present day. In some genomic regions, then, marine and

freshwater threespine stickleback are as divergent as threespine and ninespine

stickleback, which are classified into separate genera.

Adaptive divergence has impacted the history of the stickleback genome as a

whole (Fig. 4C). We identified 32.6 Mb, or 7.5%, of the genome as having elevated

T_{MRCA} (gray boxes in Fig. 4C; two-sided permutation test, $p \leq 0.001$). Outside of the

non-recombining portion of the sex chromosome (chr. 19), the oldest regions of the

stickleback genome were those enriched for divergent loci. Patterns of ancient ancestry

closely mirrored recent divergence in allele frequencies (Fig. 2A) and it appears that

historical and contemporary marine-freshwater divergence has impacted ancestry

across much of the length of some chromosomes. Chromosome 4, for example,

contains at least three broad peaks in T_{MRCA} and a total of 5.9 Mb identified as genome-

wide outliers (two-sided permutation test, $p \leq 0.001$). This chromosome has been of

particular interest because of its association with a number of phenotypes [20,41], including

fitness[42]. We found the major-effect armor plate locus *eda* comprised a local peak

(mean T_{MRCA} = 6.4 MYA) nested within a large region of deep ancestry spanning 8.1

Mb. Moreover, at least two other peaks distal to *eda,* centered at 21.4 Mb and 26.6 Mb,

were also several million years older than the genomic average at 6.8 MYA and 7.0

MYA, respectively.

Intriguingly, genomic regions of elevated T_{MRCA} remained outliers even after

removing marine-freshwater relative divergence outliers as measured by F_{ST} (Suppl.

Fig. 3). We estimated that 7.5% of the genome had increased T_{MRCA} even though only

1.9% of RAD loci (1129 of 57,992) were classified as divergent. When we removed

these loci along with loci with extreme values of marine-freshwater F_{ST} ($F_{ST} > 0.5$),

many of the regions in which they resided were still T_{MRCA} outliers. It is possible that the

remainder of this old variation is neutral with respect to fitness. However, we identified

divergence outliers based on only a single axis of divergence: the marine-freshwater

axis. Throughout the entire species range, populations are locally experiencing multiple

axes of divergence, including lake-stream and benthic-limnetic axes[43], that often shares

a common genomic architecture[44,45]. Our data may indicate underlying similarities in

selection regimes. Alternatively, this co-localized ancient variation may represent the

accumulation of adaptive divergence along multiple axes in the same genomic regions,

whether or not the underlying adaptive variants are the same. Aspects of the genomic

architecture, such as gene density or local recombination rates, may in part govern

where in the genome adaptive divergence can occur[46-48]. Multiple axes of divergence

may therefore act synergistically to maintain genomic variation across the stickleback

metapopulation.

Nevertheless, much of the ancient variation we observe may in fact itself be

neutral, having been maintained by close linkage to loci under divergent selection

between the marine and freshwater habitats[32]. Indeed, the broadest peaks of T_{MRCA} we

observe occur in genomic regions with low rates of recombination[47,49] in other

stickleback populations, which would extend the size of the linked region affected by

divergent selection. On ecological timescales, low recombination rates in stickleback

are thought to promote divergence by making locally adapted genomic regions resistant

to gene flow[47]. Our results potentially extend the inferred impact of recombination rate

variation on genomic variation to timescales that are 1000-fold longer, maintaining both

multimillion-year-old adaptive variation and large stores of linked genetic variation.

Future modeling efforts will be needed to explore the range of population genetic

parameter values (e.g. selection coefficients, migration rates, and recombination rates)

required to produce the extent of divergence we see here.

Lastly, our findings demonstrate that known chromosomal inversions maintain

globally distributed, multilocus haplotypes. The three chromosomal inversions (*inv1*,

inv11, and *inv21*; yellow bars in Fig. 4C) all showed sharp spikes in T_{MRCA}. Genomic

signatures of these inversions are distributed throughout the species range, including

coastal marine-freshwater population pairs in the Pacific and Atlantic basins[19] and

inland lake-stream pairs in Switzerland. [44]. Despite our limited geographic sampling, our

finding that all three of these inversions are over six million years old is further evidence

of single, ancient origins of each, followed by their spread across the species range.

Each inversion contained a high density of divergent RAD loci (*inv1*: 64% of loci

divergent; *inv11*: 60%; *inv21*: 71%) but we also identified regions within these inversions

in which haplotypes from marine or freshwater habitats, or both, were not monophyletic.

inv1 and *inv11* both contained two regions separated by loci in which neither habitat

type was monophyletic; *inv21*, the largest of the three, contained ten such regions.

Additionally, T_{MRCA} and F_{ST} decreased sharply to background levels outside of the

inversions, demonstrating the potential for gene flow and recombination to homogenize

variation in these regions. We interpret this as evidence that these inversions help

maintain linkage disequilibrium among multiple divergently adaptive variants in regions

susceptible to homogenization[11,50]. The presence of these inversions, therefore, further

supports the hypothesis that the recombinational landscape can influence where in the

genome adaptive divergence can occur and emphasizes the degree to which gene flow

among divergently adapted stickleback populations has impacted global genomic

diversity.

CONCLUSIONS

Selection operating on two very different timescales — the ecological and the

geological — has shaped genomic patterns of SGV in the threespine stickleback.

Selection on ecological timescales drives phenotypic divergence in decades or millennia

by sorting SGV across geography and throughout the genome[22,44,51,52]. Our findings

show that the persistence of this ecological diversity and local adaptation of stickleback

has set the stage for long-term divergent selection and for the continual accumulation

and maintenance of adaptive variation over millions of years. A number of genetic

variants fueling contemporary, rapid adaptation may even have been present - and

under selection - since before the threespine-ninespine stickleback lineages split. The

extent to which ecological adaptation in a single population drew on haplotypes that

have evolved over millions of years and persisted in multiple populations, many of which

are now extinct, underscores the need to understand macroevolutionary patterns when

studying microevolutionary processes, and vice versa.

METHODS

Sample collection and library preparation

Wild threespine stickleback were collected from Rabbit Slough (N 61.5595, W

149.2583), Boot Lake (N 61.7167, W 149.1167), and Bear Paw Lake (N 61.6139, W

149.7539). Rabbit Slough is an offshoot of the Knik Arm of Cook Inlet and is known to

be populated by anadromous populations of stickleback that are stereotypically oceanic

in phenotype and genotype [22,53]. Boot Lake and Bear Paw Lake are both shallow lakes

formed during the end-Pleistocene glacial retreat. Fish were collected in the summers of

2009 (Rabbit Slough), 2010 (Bear Paw Lake), and 2014 (Boot Lake) using wire minnow

traps and euthanized *in situ* with Tricaine solution. Euthanized fish were immediately

fixed in 95% ethanol and shipped to the Cresko Laboratory at the University of Oregon

(Eugene, OR, USA). DNA was extracted from fin clips preserved in 95% ethanol using

either Qiagen DNeasy spin column extraction kits or Ampure magnetic beads (Beckman

Coulter, Inc) following manufacturer's instructions. Yields averaged 1-2 µg DNA per

extraction (~30 mg tissue). Treatment of animals followed protocols approved the

University of Oregon Institutional Animal Care and Use Committee (IACUC).

We designed our library preparation strategy to identify sufficient sequence

variation for gene tree reconstruction and to simplify downstream sequence processing

and analysis by taking advantage of the phase information captured by paired-end

sequencing. We generated RAD libraries from these samples using the single-digest

sheared RAD protocol from Baird et al. with the following specifications and

adjustments: 1 µg of genomic DNA per fish was digested with the restriction enzyme

PstI-HF (New England Biolabs), followed by ligation to P1 Illumina adaptors with 6 bp

inline barcodes. Ligated samples were multiplexed and sheared by sonication in a

Bioruptor (Diagenode). To ensure that most of our paired-end reads would overlap

unambiguously and produce longer contiguous sequences, we selected a narrow

fragment size range of 425-475 bp. The remainder of the protocol was per Baird et al.

[31]. All fish were sequenced on an Illumina HiSeq 2500 using paired-end 250 bp

sequencing reads at the University of Oregon's Genomics and Cell Characterization

Core Facility (GC3F).

Sequence preparation

Raw Illumina sequence reads were demultiplexed, cleaned, and processed

primarily using the Stacks pipeline[54]. Paired-end reads were demultiplexed with

process_shortreads and cleaned using **process_radtags** using default criteria

(throughout this document, names of scripts, programs, functions, and command-line

arguments will appear in **fixed-width font**). Overlapping read pairs were then

merged with **fastq-join** [55] (Fig. S1). Pairs that failed to merge were removed from

further analysis. In order to retain the majority of the sequence data for analysis in

Stacks and still maintain adequate contig lengths, merged contigs were trimmed to 350

bp and all contigs shorter than 350 bp were discarded. We aligned these contigs to the

stickleback reference genome[19,49] using **bbmap** with the most sensitive alignment

settings ('**vslow=t**'; http://jgi.doe.gov/data-and-tools/bbtools/) and used the **pstacks**,

cstacks, and **sstacks** components of the Stacks pipeline to create stacks and call

SNPs and haplotypes, create a catalog of RAD tags across individuals, and match tags

across individuals. All data were then passed through the Stacks error correction

module **rxstacks** to prune unlikely haplotypes. We ran the Stacks component program

populations on the final dataset to filter loci genotyped in fewer than four individuals in

each population and to create output files for sequence analysis. We use the naming

conventions of Baird et al. [56]: A "RAD tag" refers to sequence generated from a single

end of a restriction site and the pair of RAD tags sequenced at a restriction site

comprises a "RAD locus" (Figure 2.1).

We used the program **phase** [57] to phase pairs of RAD tags originating from the

same restriction site. We coded haplotypes present at each RAD tag, which often

contain multiple SNPs, into multiallelic genotypes. This both simplified and reduced

computing time for the phasing process. Custom Python scripts automated this process

and are included as supplementary files. We required that each individual had at least

one sequenced haplotype at each tag for phasing to be attempted. If a sample had

called genotypes at only one tag in the pair, the sample was removed from further

analysis of that locus. The resultant phased haplotypes were used to generate

sequence alignments for import into BEAST.

We recovered a total of 236,787 RAD tags after filtering, mapping to 151,813 *PstI*

restriction sites. At 84,974 restriction sites, we recovered and successfully phased

adjacent RAD tags (169,948 RAD tags) into single RAD loci. We retained these 84,974

RAD loci for our analysis.

Ninespine stickleback genome assembly

In order to estimate the T_{MRCA} of threespine stickleback RAD alleles, we used the

ninespine stickleback (*Pungitius pungitius*) as an outgroup (Figure 3.1, see Figure 1.2).

RAD sequence analysis, however, relies on the presence of homologous restriction

sites among sampled individuals and results in null alleles when mutations occur within

a restriction site[58]. Because this probability increases with greater evolutionary distance

among sampled sequences, we elected to use RAD-seq to only estimate sequence

variation within the threespine stickleback. We then generated a contig-level *de novo*

ninespine stickleback genome assembly from a single ninespine stickleback individual

from St. Lawrence Island, Alaska (collected by J. Postlethwait) using DISCOVAR *de*

novo (https://software.broadinstitute.org/software/discovar). We used this single

ninespine stickleback haplotype to estimate threespine-ninespine sequence divergence

and time calibrate coalescence times within the threespine stickleback. DISCOVAR de

novo requires a single shotgun library of paired-end 250-bp sequence reads from short-

insert-length DNA fragments. High molecular weight genomic DNA was extracted from

an ethanol-preserved fin clip by proteinase K digestion followed by DNA extraction with

Ampure magnetic beads. Purified genomic DNA was mechanically sheared by

sonication and size selected to a range of 200-800 bp by gel electrophoresis and

extraction. We selected this fragment range to agree with the recommendations for *de*

novo assembly using the DISCOVAR *de novo*

(https://software.broadinstitute.org/software/discovar/blog). This library was sequenced

on a single lane of an Illumina HiSeq2500 at the University of Oregon's Genomics and

Cell Characterization Core Facility (GC3F: https://gc3f.uoregon.edu/). We assembled

the draft ninespine stickleback genome using DISCOVAR de novo. Raw sequence read

pairs were first quality filtered and adaptor sequence contamination removed using the

program **process_shortreads**, which is included in the Stacks analysis pipeline [59]. We

ran the genome assembly on the University of Oregon's Applied Computational

Instrument for Scientific Synthesis (ACISS: http://aciss-computing.uoregon.edu).

Alignment of RAD tags to the ninespine assembly

We included the single ninespine stickleback haplotype into our sequence

analyses by aligning a single phased threespine stickleback RAD haplotype from each

locus to the ninespine genome assembly. For those that aligned uniquely (59,254 RAD

loci), we used a custom Python script to parse the output BAM file [60] and reconstruct the

ninespine haplotype from the query sequence and alignment fields. The final dataset

consists of 57,992 RAD loci that mapped to the 21 threespine stickleback chromosomes

and aligned uniquely to the ninespine assembly.

Lineage sorting and time to the most recent common ancestor

Allelic divergence can occur by multiple modes of lineage sorting during

adaptation. To identify patterns of lineage sorting associated with freshwater

colonization, we analyzed gene tree topologies at all RAD loci using BEAST v. 1.7[39,61].

We used blanket parameters and priors for BEAST analyses across all RAD loci.

Markov chain Monte Carlo (MCMC) runs of 1,000,000 states were specified, and trees

logged every 100 states. We used a coalescent tree prior and the GTR+Γ substitution

model with four rate categories and uniform priors for all substitution rates. We identified

evidence of lineage sorting by using the program **treeannotator** to select the

maximum clade credibility (MCC) tree for each RAD locus and the

`is.monophyletic()` function included in the R package 'ape' [62]. We determined for

each MCC tree whether tips originating from marine (RS) or freshwater (BL+BP) formed

monophyletic clades.

To convert node ages estimated in BEAST into divergence times, in years, we

assumed a 15 million-year divergence time between threespine and ninespine

stickleback at each RAD locus [38]. The T_{MRCA} of all alleles in each gene tree was set at

15 Mya at each node age of interest was converted into years relative to the total height

of the tree. Additionally, to use the ninespine stickleback as an outgroup, we required

that threespine stickleback haplotypes at a RAD locus were monophyletic to the

exclusion of the ninespine haplotype. Doing so reduced our analysis to 49,672 RAD loci

for analyses included in Fig. 4 of the main text. RAD loci not showing this pattern of

lineage sorting did not show evidence of a genome-wide correlation with marine-

freshwater divergence and thus do not impact the assertions in the main text. We used

medians of the posterior distributions as point estimates of T_{MRCA} for each RAD locus.

Because of the somewhat limited information from any single RAD locus, and because

the facts of the genealogical process mean that the true T_{MRCA} at any locus likely differs

from the 15 My estimate [63-65], we do not rely heavily on T_{MRCA} estimates at individual

RAD loci. Rather, we use these estimates to understand patterns of broad patterns of

ancestry throughout the threespine stickleback genome — spatially along chromosomes

and genome-wide patterns.

We determined T_{MRCA} outlier genomic regions by permuting and kernel

smoothing the genomic distribution of T_{MRCA} estimates using the same window sizes as

we present in the main text. Windows where the actual T_{MRCA} value exceeded 99.9% of

permuted windows were considered outliers. This method controls for the local density

of RAD loci (poorly sampled regions will have larger confidence bands) and the size of

the windows used.

Sequence diversity and haplotype networks

We quantified sequence diversity within and among populations and sequence

divergence between populations using R (R Core Team [66]). We used the R package

'ape' [62] to compute pairwise distance matrices for all alleles at each RAD locus and

used these matrices to calculate the average pairwise nucleotide distances, π, within

and among populations along with d_{XY}, the average pairwise distance between two

sequences using only across-population comparisons [67]. We also calculated the

haplotype-based F_{ST} from Hudson et al. [68] implemented in the R package 'PopGenome'

[69]. We used permutation tests written in R to identify differences in variation within- and

between-habitat type at divergent RAD loci versus the genome-wide distributions.

Mann-Whitney-Wilcoxon tests implemented in R were used to identify variation in

genome-wide diversity among populations and habitat types.

We constructed haplotype networks of the RAD loci at *eda* and *atp1a1* using the

infinite sites model with the function **haploNet()** in the R package 'pegas'[70]. The

atp1a1 network was constructed from from a RAD locus spanning exon 15 of *atp1a1*

and including portions of introns 14 and 15 at (chr1:21,726,729-21,727,381 [BROAD

S1, v89]; chr1: 26,258,117-26,257,465 [re-scaffolding from Glazer, *et al* [49]]). The *eda*

network spans exon 2 and portions of introns 1 and 3 of *eda* (chr4: 12,808,396-

12,809,030).

Code availability

Scripts used to phase RAD-tags, summarize gene trees, calculate population genetic

statistics, and produce figures and statistics presented in paper are available at

https://github.com/thomnelson/ancient-divergence. Scripts for processing raw sequence

data are available from the authors upon request.

DATA AVAILABILITY

Raw sequence data supporting these findings are available on the Sequence Read

Archive at PRJNAXXXXXX. The final datasets needed to reproduce the figures and

statistics presented in the paper are available at https://github.com/thomnelson/ancient-

divergence.

ACKNOWLEDGEMENTS

We thank P. Phillips, M. Streisfeld, J. Postlethwait, K. Sterner for valuable input and

lively discussion throughout this project. We also thank K. Alligood, E. Beck, S.

Bassham, M. Chase, M. Currey, M. Hahn, L. Fishman, C. Small, S. Stankowski, J.

Willis, two anonymous reviewers, and members of the Cresko Lab and the Institute of

Ecology and Evolution for advice and comments on previous versions of this

manuscript. J. Postlethwait graciously donated ninespine stickleback tissue, collected

under award XXXXXXXX. We acknowledge National Science Foundation awards NSF

DEB 1501423 (WAC and TCN), NSF DEB 0949053 (WAC), and National Institutes of

Health award NIH T32GM007413 (TCN).

AUTHOR CONTRIBUTIONS

TCN and WAC conceived of the project and designed sampling, sequencing, and

analysis. TCN prepared sequencing libraries, wrote software, and performed data

analysis. TCN and WAC wrote the paper.

COMPETING FINANCIAL INTERESTS

The authors declare no competing financial interests.

LITERATURE CITED

1   Wright, S. The roles of mutation, inbreeding, crossbreeding and selection in
evolution. *Proceedings of the Sixth International Congress on Genetics* **1**, 356-
366 (1932).
2   Orr, H. A. The genetic theory of adaptation: a brief history. *Nature Reviews*
Genetics **6**, 119-127, doi:10.1038/nrg1523 (2005).
3   Barrett, R. D. H. & Schluter, D. Adaptation from standing genetic variation.
Trends in Ecology and Evolution **23**, 38-44, doi:10.1016/j.tree.2007.09.008
(2008).
4   Domingues, V. S. *et al.* Evidence of adaptation from ancestral variation in young
populations of beach mice. *Evolution* **66**, 3209-3223, doi:10.1111/j.1558-
5646.2012.01669.x (2012).

5   Schrider, D. R. & Kern, A. D. Soft sweeps are the dominant mode of adaptation
in the human genome. *Molecular Biology and Evolution*,
doi:10.1093/molbev/msx154 (2017).
6   Huerta-Sánchez, E. *et al.* Altitude adaptation in Tibetans caused by introgression
of Denisovan-like DNA. *Nature* **512**, 194-197, doi:10.1038/nature13408 (2014).
7   Fontaine, M. C. *et al.* Mosquito genomics. Extensive introgression in a malaria
vector species complex revealed by phylogenomics. *Science* **347**, 1258524,
doi:10.1126/science.1258524 (2015).
8   Grant, P. R. & Grant, B. R. Unpredictable evolution in a 30-year study of Darwin's
finches. *Science* **296**, 707-711, doi:DOI 10.1126/science.1070315 (2002).
9   Wright, K. M., *et al.* Indirect Evolution of Hybrid Lethality Due to Linkage with
Selected Locus in Mimulus guttatus. *PLoS Biology* **11**,
doi:10.1371/journal.pbio.1001497 (2013).
10   Colosimo, P. F. *et al.* Widespread parallel evolution in sticklebacks by repeated
fixation of Ectodysplasin alleles. *Science* **307**, 1928-1933,
doi:10.1126/science.1107239 (2005).
11   Kirkpatrick, M. & Barton, N. Chromosome inversions, local adaptation and
speciation. *Genetics* **173**, 419-434, doi:10.1534/genetics.105.047985 (2006).
12   Charlesworth, B., Morgan, M. T. & Charlesworth, D. The effect of deleterious
mutations on neutral molecular variation. *Genetics* **134**, 1289-1303 (1993).
13   Linnen, C. R., *et al.* On the origin and spread of an adaptive allele in deer mice.
Science **325**, 1095-1098, doi:10.1126/science.1175826 (2009).
14   Stankowski, S. & Streisfeld, M. A. Introgressive hybridization facilitates adaptive
divergence in a recent radiation of monkeyflowers. *Proceedings of the Royal
Society B: Biological Sciences* **282**, 20151666, doi:10.1098/rspb.2015.1666
(2015).
15   Pease, J. B., *et al.* Phylogenomics reveals three sources of adaptive variation
during a rapid radiation. *PLoS Biology* **14**, e1002379,
doi:10.1371/journal.pbio.1002379 (2016).
16   Schlotterer, C., *et al.* Sequencing pools of individuals - mining genome-wide
polymorphism data without big funding. *Nature Reviews Genetics* **15**, 749-763,
doi:10.1038/nrg3803 (2014).
17   Davey, J. W. *et al.* Genome-wide genetic marker discovery and genotyping using
next-generation sequencing. *Nature Reviews Genetics* **12**, 499-510,
doi:10.1038/nrg3012 (2011).
18   Bell, M. A. & Foster, S. A. in *The Evolutionary Biology of the Threespine
Stickleback*   (eds M. A. Bell & S. A. Foster) Ch. 1, 1-27 (Oxford University Press,
1994).
19   Jones, F. C. *et al.* The genomic basis of adaptive evolution in threespine
sticklebacks. *Nature* **484**, 55-61, doi:10.1038/nature10944 (2012).
20   Colosimo, P. F. *et al.* The genetic architecture of parallel armor plate reduction in
threespine sticklebacks. *PLoS Biology* **2**, 635-641,
doi:10.1371/journal.pbio.0020109 (2004).

21   Cresko, W. A. *et al.* Parallel genetic basis for repeated evolution of armor loss in
532   Alaskan threespine stickleback populations. *Proc Natl Acad Sci U S A* **101**, 6050-
533   6055, doi:10.1073/pnas.0308479101 (2004).

534   22   Hohenlohe, P. A. *et al.* Population genomics of parallel adaptation in threespine
535   stickleback using sequenced RAD tags. *Plos Genet* **6**, e1000862,
536   doi:10.1371/journal.pgen.1000862 (2010).

537   23   Stuart, Y. E. *et al.* Contrasting effects of environment and genetics generate a
538   continuum of parallel evolution. *Nature Ecology & Evolution* **1**, 0158,
539   doi:10.1038/s41559-017-0158 (2017).

540   24   Roesti, M., *et al.* The genomic signature of parallel adaptation from shared
541   genetic variation. *Mol Ecol* **23**, 3944-3956, doi:10.1111/mec.12720 (2014).

542   25   Francis, R. C., *et al.* Historical and ecological sources of variation among lake
543   populations of threespine sticklebacks, *Gasterosteus aculeatus*, near Cook Inlet,
544   Alaska. *Canadian Journal of Zoology* **64**, 2257-2265 (1986).

545   26   Bell, M. A. & Foster, S. A. in *The Evolutionary Biology of the Threespine
546   Stickleback* (eds M. A. Bell & S. A. Foster) Ch. 16, 472-486 (Oxford University
547   Press, 1994).

548   27   Kimmel, C. B. *et al.* Evolution and development of facial bone morphology in
549   threespine sticklebacks. *Proc Natl Acad Sci U S A* **102**, 5791-5796,
550   doi:10.1073/pnas.0408533102 (2005).

551   28   Reimchen, T. E. in *The Evolutionary Biology of the Threespine Stickleback* (eds
552   M. A. Bell & S. A. Foster) Ch. 9, 240-276 (Oxford University Press, 1994).

553   29   Arnegard, M. E. *et al.* Genetics of ecological divergence during speciation.
554   *Nature* **511**, 307-311, doi:10.1038/nature13301 (2014).

555   30   Schluter, D. & Conte, G. L. Genetics and ecological speciation. *Proc Natl Acad
556   Sci U S A* **106 Suppl 1**, 9955-9962, doi:10.1073/pnas.0901264106 (2009).

557   31   Baird, N. A. *et al.* Rapid SNP discovery and genetic mapping using sequenced
558   RAD markers. *Plos One* **3**, e3376, doi:10.1371/journal.pone.0003376 (2008).

559   32   Charlesworth, B., Nordborg, M. & Charlesworth, D. The effects of local selection,
560   balanced polymorphism and background selection on equilibrium patterns of
561   genetic diversity in subdivided populations. *Genetics Research* **70**, 155-174,
562   doi:Doi 10.1017/S0016672397002954 (1997).

563   33   Lenormand, T. Gene flow and the limits to natural selection. *Trends Ecol Evol* **17**,
564   183-189, doi:Doi 10.1016/S0169-5347(02)02497-7 (2002).

565   34   Otto, S. P. & Bourguet, D. Balanced polymorphisms and the evolution of
566   dominance. *American Naturalist* **153**, 561-574, doi:Doi 10.1086/303204 (1999).

567   35   Phillips, P. C. Epistasis--the essential role of gene interactions in the structure
568   and evolution of genetic systems. *Nature Reviews Genetics* **9**, 855-867,
569   doi:10.1038/nrg2452 (2008).

570   36   McGuigan, K., *et al.* Cryptic genetic variation and body size evolution in
571   threespine stickleback. *Evolution* **65**, 1203-1211, doi:10.1111/j.1558-
572   5646.2010.01195.x (2011).

573   37   McCairns, R. J. S. & Bernatchez, L. Plasticity and heritability of morphological
574   variation within and between parapatric stickleback demes. *Journal of*

575   *Evolutionary Biology* **25**, 1097-1112, doi:10.1111/j.1420-9101.2012.02496.x
576   (2012).
577 38 Aldenhoven, J. T., *et al.* Phylogeography of ninespine sticklebacks (*Pungitius*
578   *pungitius*) in North America: glacial refugia and the origins of adaptive traits. *Mol*
579   *Ecol* **19**, 4061-4076, doi:10.1111/j.1365-294X.2010.04801.x (2010).
580 39 Drummond, A. J., Suchard, M. A., Xie, D. & Rambaut, A. Bayesian phylogenetics
581   with BEAUti and the BEAST 1.7. *Mol Biol Evol* **29**, 1969-1973,
582   doi:10.1093/molbev/mss075 (2012).
583 40 Bell, M. A., Baumgartner, J. V. & Olson, E. C. Patterns of temporal change in
584   single morphological characters of a Miocene stickleback fish. *Paleobiology* **11**,
585   258-271 (1985).
586 41 Miller, C. T. *et al.* Modular skeletal evolution in sticklebacks is controlled by
587   additive and clustered quantitative trait loci. *Genetics* **197**, 405-420,
588   doi:10.1534/genetics.114.162420 (2014).
589 42 Barrett, R. D., Rogers, S. M. & Schluter, D. Natural selection on a major armor
590   gene in threespine stickleback. *Science* **322**, 255-257,
591   doi:10.1126/science.1159978 (2008).
592 43 McKinnon, J. S. & Rundle, H. D. Speciation in nature:the threespine stickleback
593   model systems. *Trends in Ecology and Evolution* **17**, 480-488 (2002).
594 44 Roesti, M., Kueng, B., Moser, D. & Berner, D. The genomics of ecological
595   vicariance in threespine stickleback fish. *Nature Communications* **6**, 8767,
596   doi:10.1038/ncomms9767 (2015).
597 45 Deagle, B. E. *et al.* Population genomics of parallel phenotypic evolution in
598   stickleback across stream-lake ecological transitions. *Proceedings of the Royal*
599   *Society B: Biological Sciences* **279**, 1277-1286, doi:10.1098/rspb.2011.1552
600   (2012).
601 46 Samuk, K. *et al.* Gene flow and selection interact to promote adaptive divergence
602   in regions of low recombination. *Molecular Ecology*, doi:10.1111/mec.14226
603   (2017).
604 47 Roesti, M., Moser, D. & Berner, D. Recombination in the threespine stickleback
605   genome - Patterns and consequences. *Molecular Ecology* **22**, 3014-3027,
606   doi:10.1111/mec.12322 (2013).
607 48 Aeschbacher, S., *et al.* Population-genomic inference of the strength and timing
608   of selection against gene flow. *Proceedings of the National Academy of Sciences*
609   *USA* **114**, 7061-7066, doi:10.1073/pnas.1616755114 (2017).
610 49 Glazer, A. M., *et al.* Genome assembly improvement and mapping of
611   convergently evolved skeletal traits in sticklebacks with genotyping-by-
612   sequencing. *G3-Genes Genomes Genetics* **5**, 1463-1472,
613   doi:10.1534/g3.115.017905 (2015).
614 50 Guerrero, R. F., Rousset, F. & Kirkpatrick, M. Coalescent patterns for
615   chromosomal inversions in divergent populations. *Philosophical Transactions of*
616   *the Royal Society B: Biological Sciences* **367**, 430-438,
617   doi:10.1098/rstb.2011.0246 (2012).

618 51   Hendry, A. P., Taylor, E. B. & McPhail, J. D. Adaptive divergence and the
619   balance between selection and gene flow: lake and stream stickleback in the
620   Misty system. *Evolution* **56**, 1199-1216, doi:10.1554/0014-
621   3820(2002)056[1199:ADATBB]2.0.CO;2 (2002).
622 52   Lescak, E. A. *et al.* Evolution of stickleback in 50 years on earthquake-uplifted
623   islands. *Proceedings of the National Academy of Sciences USA* **112**, E7204-
624   7212, doi:10.1073/pnas.1512020112 (2015).
625 53   Cresko, W. A. *et al.* Parallel genetic basis for repeated evolution of armor loss in
626   Alaskan threespine stickleback populations. *Proceedings of the National*
627   *Academy of Sciences USA* **101**, 6050-6055, doi:10.1073/pnas.0308479101
628   (2004).
629 54   Catchen, J. M., *et al.* Stacks: building and genotyping Loci de novo from short-
630   read sequences. *G3 - Genes Genomes Genetics* **1**, 171-182,
631   doi:10.1534/g3.111.000240 (2011).
632 55   Aronesty, E. ea-utils: Command-line tools for processing biological sequencing
633   data. (2011).
634 56   Baird, N. A. *et al.* Rapid SNP discovery and genetic mapping using sequenced
635   RAD markers. *Plos One* **3**, 1-7, doi:10.1371/journal.pone.0003376 (2008).
636 57   Stephens, M., Smith, N. J. & Donnelly, P. A new statistical method for haplotype
637   reconstruction from population data. *American Journal of Human Genetics* **68**,
638   978-989, doi:Doi 10.1086/319501 (2001).
639 58   Arnold, B., *et al.* RADseq underestimates diversity and introduces genealogical
640   biases due to nonrandom haplotype sampling. *Mol Ecol* **22**, 3179-3190,
641   doi:10.1111/mec.12276 (2013).
642 59   Catchen, J., *et al.* Stacks: An analysis tool set for population genomics.
643   *Molecular Ecology* **22**, 3124-3140, doi:10.1111/mec.12354 (2013).
644 60   Li, H. *et al.* The Sequence alignment/map (SAM) format and SAMtools.
645   *Bioinformatics* **25**, 2078-2079 (2009).
646 61   Drummond, A. J. & Rambaut, A. BEAST: Bayesian evolutionary analysis by
647   sampling trees. *Bmc Evol Biol* **7**, 214, doi:10.1186/1471-2148-7-214 (2007).
648 62   Paradis, E., Claude, J. & Strimmer, K. APE: Analyses of phylogenetics and
649   evolution in R language. *Bioinformatics* **20**, 289-290,
650   doi:10.1093/bioinformatics/btg412 (2004).
651 63   Kingman, J. F. C. The coalescent. *Stochastic Processes and their Applications*
652   **13**, 235-248, doi:10.1016/0304-4149(82)90011-4 (1982).
653 64   Kingman, J. F. C. On the genealogy of large populations. *Journal of Applied*
654   *Probability* **19**, 27-43 (1982).
655 65   Tajima, F. Evolutionary relationship of DNA-sequences in finite populations.
656   *Genetics* **105**, 437-460 (1983).
657 66   R Core Team. R Foundation for Statistical Computing, Vienna, Austria (2016).
658 67   Nei, M. *Molecular Evolutionary Genetics*. (Columbia university press, 1987).
659 68   Hudson, R. R., Slatkin, M. & Maddison, W. P. Estimation of levels of gene flow
660   from DNA-sequence data. *Genetics* **132**, 583-589 (1992).

69   Pfeifer, B., *et al*. PopGenome: An efficient swiss army knife for population
genomic analyses in R. *Molecular Biology and Evolution* **31**, 1929-1936,
doi:10.1093/molbev/msu136 (2014).
70   Paradis, E. pegas: an R package for population genetics with an integrated-
modular approach. *Bioinformatics* **26**, 419-420,
doi:10.1093/bioinformatics/btp696 (2010).
667

Supplementary Figure S1. RAD-seq effectively samples genome-wide sequence diversity. Histograms of the distance between adjacent RAD loci (left: calculated as the distance between the centers of each restriction site) and the number of variable sites per locus show that most RAD loci were within 4 kb of their nearest neighbor and contained ≥ 7 variable sites. Means for each metric are shown as dashed vertical lines. Medians are solid lines. Each histogram is truncated to highlight the bulk of the distribution. Maximum values: distance = 455 kb; variable sites = 155.

Supplementary Figure S2. Relative (F_{ST}) and absolute (d_{XY}) sequence divergence are positively correlated genome-wide in two instances of marine-freshwater divergence. Points are 250 kb non-overlapping genomic windows. Left panel compares the marine Rabbit Slough population (RS) to the freshwater Boot Lake population (BL) (type-II linear model: $r^2 = 0.314$, permuted p-value [reduced major axis] = 0.01). Right panel compares RS to the freshwater Bear Paw Lake population (BL) (type-II linear model: $r^2 = 0.311$, permuted p-value [reduced major axis] = 0.01).

Supplementary Figure S3. T_{MRCA} outlier regions remain outliers after removing highly differentiated RAD loci. Panel A is taken from Fig. 2 and shows the genomic distribution of reciprocally monophyletic ("divergent"; black bars) RAD loci. Panel B shows the distributions of T_{MRCA} outlier regions (increased T_{MRCA}) including all RAD loci (magenta boxes, "Y"). Below are the T_{MRCA} outlier regions after removing divergent loci and any RAD locus with a marine-freshwater (RS vs. [BL+BP]) $F_{ST} > 0.5$, which is approximately the top 7% of the F_{ST} distribution. Panel C: Genome scans of T_{MRCA} using all RAD loci (magenta) and excluding marine-freshwater outliers (gray).

Supplementary table 1. Sequencing summary for threespine stickleback samples

Sample	population	raw reads	filtered reads	merged pairs	mean coverage per locus
1827.05	Rabbit Slough	10167407	10031967	7269377	12X
1827.06	Rabbit Slough	10265078	10172621	7591801	13X
1827.07	Rabbit Slough	9175983	9040625	6771332	11X
1827.08	Rabbit Slough	7896938	7814081	5879351	10X
1827.09	Rabbit Slough	8773502	8668261	6405777	11X
2827.01	Boot Lake	8917575	8810382	6373001	11X
2827.07	Boot Lake	10064876	9917732	7255989	13X
2827.13	Boot Lake	9099831	9002717	6528704	12X
2827.19	Boot Lake	11021084	10792092	7911026	14X
2827.25	Boot Lake	9920574	9814758	7287485	13X
1902.02	Bear Paw Lake	4780489	4365505	2942926	5X
1902.03	Bear Paw Lake	5073434	4643909	3192582	5X
1902.04	Bear Paw Lake	4902931	4600877	3138791	6X
1902.06	Bear Paw Lake	4501906	4339253	2983345	5X

Supplementary table 2. Genome assembly statistics for *Pungitius pungitius*.

	contig (scaffold)
n	393,037 (391,396)
Max length (bp)	165,088 (182,644)
N50 (bp)	9,202 (9,886)
Average length (bp)	1,314 (1,320)
Gaps (%)	0.03
Total assembly length (bp)	516,674,741