Title

A GAL80 collection to inhibit GAL4 transgenes

in *Drosophila* olfactory sensory neurons

Authors

Eliason, Jessica*[†][1]; Afify, Ali[‡]; Potter, Christopher[‡]; Matsumura, Ichiro[†]

Affiliations

[*]*Janelia Research Campus, Howard Hughes Medical Institute; Ashburn, VA, 20147;

United States

[†] Department of Biochemistry; Emory University School of Medicine; Atlanta, GA,

30322; United States

[‡] Solomon H. Snyder Department of Neuroscience; Johns Hopkins University School of

Medicine; Baltimore, MD, 21205; United States

---

[1] 19700 Helix Drive; Ashburn, VA; 20147

Running Title

GAL80 in *Drosophila* olfactory neurons

Key Words

Olfaction; GAL80; Drosophila; Odorant Receptor; Sensory Neurons

Corresponding authors' contact information

Jessica Eliason

19700 Helix Drive

2E.120.2

Ashburn, VA 20147

(209) 712-9343

eliasonj@janelia.hhmi.org;

&

Ichiro Matsumura

1510 Clifton Rd NE

4119 Rollins Research Center

Atlanta, GA 30322

(404) 727-5625

 imatsum@emory.edu

ABSTRACT

Fruit flies recognize hundreds of ecologically relevant odors and respond appropriately to them. The complexity, redundancy and interconnectedness of the olfactory machinery complicate efforts to pinpoint the functional contributions of any component neuron or receptor to behavior. Some contributions can only be elucidated in flies that carry multiple mutations and transgenes, but the production of such flies is currently labor-intensive and time-consuming. Here, we describe a set of transgenic flies that express the *Saccharomyces cerevisiae* GAL80 in specific olfactory sensory neurons (*OrX-GAL80s*). The GAL80s effectively and specifically subtract the activities of GAL4-driven transgenes that impart anatomical and physiological phenotypes. *OrX-GAL80s* can allow researchers to efficiently activate only one or a few types of functional neurons in an otherwise nonfunctional olfactory background. Such experiments will improve our understanding of the mechanistic connections between odorant inputs and behavioral outputs at the resolution of only a few functional neurons.

INTRODUCTION

The olfactory system of *Drosophila melanogaster* is often the subject in studies of memory, evolution, gene choice, development and odorant-induced behavior. It is a good model system because of its relatively stereotyped neuronal circuitry, complex behaviors and convenient genetic tools.

In *Drosophila*, most olfactory sensory neurons (OSNs) typically expresses a

single odorant receptor (OR) from a genomic repertoire of 60 genes (Vosshall et al.

1999; Robertson, Warr, and Carlson 2003; Vosshall, Wong, and Axel 2000; Clyne et al.

1999; Goldman et al. 2005). The promoter of an OR gene can be employed to label

specific subsets of OSNs with a particular transgene (Fishilevich and Vosshall 2005;

Couto, Alenius, and Dickson 2005). ORs, which vary in sensitivity and specificity to a

wide range of different odorants, determine the firing kinetics and odor-response

dynamics of each OSN (Hallem and Carlson 2006; Hallem, Ho, and Carlson 2004;

Couto, Alenius, and Dickson 2005; Fishilevich and Vosshall 2005; de Bruyne, Clyne,

and Carlson 1999; de Bruyne, Foster, and Carlson 2001; Dobritsa et al. 2003; Elmore et

al. 2003; Kreher et al. 2008).

Most OSNs express Odorant Receptor Co-Receptor (Orco), a highly conserved

member of the olfactory receptor family (Krieger et al. 2003; Vosshall and Hansson

2011), in addition to a single selected OR. Though Orco usually does not contribute to

the structure of the odorant binding site (Jung, borst, and Haag 2011; Nakagawa and

Vosshall 2009; Nichols and Luetje 2010; P. L. Jones, Pask, and Rinker 2011), it is

essential for odorant-invoked signaling in flies. Without Orco, the co-expressed OR

cannot localize to the dendritic membrane or relay an odor-evoked signal (Larsson et al.

2004; Benton et al. 2006). Orco null flies are largely anosmic, though some

chemosensation remains due to the presence of ionotropic receptors and gustatory

receptors, which do not require Orco to function (Silbering et al. 2011; Benton et al.

2009; Kwon et al. 2007; W. D. Jones et al. 2007). The Orco promoter is consequently a

convenient device for the expression of transgenes in most OSNs.

The olfactory organs, the antenna and maxillary palp, contain OSNs dendrites

within structures called sensilla. ORs and Orco are embedded in the dendritic

membrane. OSN axons project to the antennal lobes in the brain of the animal. Each

antennal lobe consists of ~50 globular synaptic sites called glomeruli. All OSNs on the

periphery that expresses the same OR converge onto their own unique glomerulus. For

example, all OSNs expressing Or22a will send axons to the DM2 glomerulus in the

antennal lobe while all OSNs expressing Or82a will send axons to the VA6 glomerulus

(**Figure 1**). The stereotyped organization of OSNs and their projections is known as the

olfactory sensory map (Vosshall, Wong, and Axel 2000; Fishilevich and Vosshall 2005;

Couto, Alenius, and Dickson 2005; Stocker et al. 1990). The regularity of this map is a

key feature that makes *Drosophila* olfaction such a useful model, as any aberration to

the typical pattern will be apparent. The apparent simplicity of the map (**Figure 1**),

however, obscures mechanistic complexities that are yet to be discovered, in part

because necessary tools remain unavailable.

Drosophila geneticists have traditionally relied on genetic mutations or deletions

to understand how complex biological systems normally work. Most alleles are

recessive, so homozygotes must be bred over multiple generations. Achieving

homozygosity of a mutation while also adding transgenes to the system often requires

the creation of recombinant chromosomes produced after multiple generations of

crossing and PCR screening. Classical genetic strategies thus limit the number and

complexity of combinatorial genotypes that one can achieve. More challenging

experimental questions demand more facile and versatile genetic tools.

The GAL4/UAS gene regulation system has become a *defacto* standard in

studies of *Drosophila*. GAL4 is a yeast transcription activator that binds to the Upstream

Activating Sequence (UAS) and induces expression of downstream genes (Giniger,

Varnum, and Ptashne 1985). By driving GAL4 expression from an OR promoter,

specific expression of a *UAS-transgene* can be obtained for any OSN subtype. An *OrX-*

GAL4 line exists for almost every OR. This collection of GAL4 lines is a powerful toolbox

since different *UAS-transgenes* can be introduced into a line via conventional mating.

For example, human α-synuclein has been expressed in OSNs to model human

Parkinson's disease (A. Y. Chen et al. 2014). Alternatively, protein expression levels

can be knocked down using any specified *UAS-RNAi* transgene.

A variety of existing compatible effectors can be used study different aspects of

neuronal communication. The *UAS-Kir2.1* effector is used as an example in

experiments described below. This inward rectifier potassium channel electrically

inactivates the neurons that express it (Hodge 2009; Baines et al. 2001; Johns et al.

1999). Similarly, *shibire^{ts}* or tetanus toxin can be used to silence synaptic

communications (van der Bliek and Meyerowitz 1991; kitamoto 2002; Kitamoto 2001; M.

S. Chen et al. 1991; Sweeney et al. 1995; Baines et al. 1999), *reaper/grim/hid* genes

can be used to physically kill neurons by inducing their own apoptotic pathways (Song

and Steller 1999; Abrams 1999), or ricin toxin can be expressed ectopically to kill

neurons. Conversely, neurons can be selectively activated with *trp1a* or a variety of

other channelrhodopsin transgenes (Boyden 2011; Pulver et al. 2009).

If GAL4 is a standard on-switch for nearly any desired transgene, GAL80 is the

logical off-switch. GAL80 binds the GAL4 transcriptional activation domain, thereby

preventing recruitment of RNA polymerase (Ma and Ptashne 1987). GAL80 crosses are

much more convenient than classical breeding approaches (**Figure 2**). In order to have

a single functional OSN in an otherwise silent olfactory system, the traditional method

uses an Orco null mutation (Larsson et al. 2004). In this genetic setup, Orco mutant flies

are mostly anosmic, but function is restored to one OSN subset with *Or-GAL4, UAS-*

Orco transgenes (Olsen, Bhandawat, and Wilson 2007; DasGupta and Waddell 2008;

Hoare, McCrohan, and Cobb 2008; Hoare et al. 2011; Benton et al. 2006; Fishilevich et

al. 2005) (**Figure 2a**). An *Orco-GAL4, UAS-effector, Or-GAL80* method can be used

instead (**Figure 2b**). Kir2.1 is used as an example of an effector (Hodge 2009; Baines et

al. 2001; Johns et al. 1999). Classical breeding strategies (**Figure 2a**) may look less

complicated on paper than GAL80 crosses (**Figure 2b**) but are actually more time-

consuming and limited. The Orco mutation must be homozygous. Since most

Drosophila transgenes are embedded into the same two chromosomes (2 or 3)

recombination and PCR screening may be required to achieve this homozygosity.

Neurons seldom operate autonomously, but rather groups of neurons coordinate

within a circuit to provide an organism with perception and behavior. An investigation of

the behavioral impact provided by a limited number of different functional neuronal

types would require additional genes. The elaboration of genotypes (**Figure 2**) to

restore pairs or groups of functional OSNs in a nonfunctional background normally

requires generations of crosses (followed by PCR screens for desired recombinants). A

GAL80 strategy can shorten this process by achieving similar results in only one or two

generations with no necessary recombinant creation. Furthermore, a GAL80 strategy

takes advantage of the interchangeable variety of existing *UAS-transgene* lines. Here

we describe a new collection of *OrX-GAL80* lines designed to complement existing *OrX-*

GAL4 lines, and demonstrate their potential utility for neuroanatomical studies of the

Drosophila olfaction model.

170   **MATERIALS AND METHODS**

Fly Stocks

Flies were reared on standard cornmeal/molasses food and kept at 25C with a

16 hours on/8hours off light cycle. All lines were obtained from the Indiana University

Bloomington Stock Center and the Janelia Research Campus. Any recombinants made

were validated with PCR.

Stock List:

Or7a-GAL4 #23907

Or7a-GAL4 #23908

Or10a-GAL4 #9944

Or13a-GAL4 #9946

Or13a-GAL4 #23886

182	Or19a-Gal4 #24617
183	Or22a-GAL4 #9951
184	Or22a-GAL4 #9952
185	Or22b-GAL4 #23891
186	Or33c-GAL4 #23893
187	Or35a-GAL4 #9967
188	Or42a-GAL4 #9970
189	Or42b-GAL4 #9971
190	Or43b-Gal4 #23894
191	Or46a-GAL4 #23291
192	Or47a-GAL4 #9981
193	Or56a-GAL4 #9988
194	Or59b-GAL4 #23897
195	Or59c-GAL4 #23899
196	Or67a-GAL4 #23904
197	Or67d-GAL4 #9998
198	Or71a-GAL4 #23121
199	Or82a-GAL4 #23125
200	Orco-GAL4 #23292
201	Orco-GAL4 #26818
202	Or85a-GAL4 #23133
203	Or85b-GAL4 #23911
204	Or85c-GAL4 #23913

Gr21a-GAL4 #24147

Or22a-mcd8::GFP #52620

Gr21a-mcd8::GFP #52619

Orco[2] #23130

UAS-Orco #23145

UAS-mcd8::GFP #5130

UAS-mcd8::GFP #5137

UAS-Kir2.1 Janelia stock #3015545

UAS-Kir2.1 Janelia stock #3015298

UAS-Kir2.1::eGFP Janelia stock #BS00312

pJFRC19-13xLexAop2-IVS-myr::GFP-p10 (attP8) Janelia stock #1171146

pJFRC59-13xLexAop2-IVS-myr::GFP-p10 (attP40) Janelia stock #3015445

GAL80 Creation

Primers were designed to capture the entire promoters described by (Couto,

Alenius, and Dickson 2005) (see **Table S1**). Promoters were amplified from genomic

DNA using Q5 High Fidelity PCR (NEB #M0491S) and added to entry vectors using the

pENTR/D-TOPO system (Invitrogen 2012b). Recombination with the pBP-GAL80Uw-6

(Addgene #26236) destination vector was done using the LR Clonase II system

(Invitrogen 2012a). To ensure no mutations, no gaps, and correct orientation, the

complete promoters were sequenced in the destination vector using the sequencing

primers shown in **Table S2**. PhiC31 site-directed transgenesis was performed by

Genetivision Inc. All GAL80 transgenes were inserted at the attP2 site. A single Or-

LexA line was also created using the Or22a-promoter entry vector and

pBPnlsLexA::p65Uw (Addgene #26230).

Immunohistochemistry

Female adult brains were dissected one day after eclosion in cold S2 Schneider's

Insect Medium (Sigma Aldrich #S0146) and fixed while nutating for 55 minutes at room

temperature in 2mL 2%PFA (Electron Microscopy Sciences #15713) in protein loBind

Tubes (Eppendorf #022431102). Brains were washed 4x, 15min per wash while

nutating with 2mL PBT buffer (1xPBS, Cellgro #21-040, with 0.5% TritonX-100, Sigma

Aldrich #X100). Brains were then blocked with 200µL 5% Goat serum (ThermoFischer.

#16210064) in PBT for 90 minutes while nutating, upright. Block was removed and 200

µL primary antibodies in PBT were added for 4 hours at room temperature and then

transferred to 4C for 36-48 hours while nutating, upright. Primary antibodies: mouse α-

bruchpilot (Developmental Studies Hybridoma Bank. #nc82-s) at 1:30, rabbit α-GFP at

1:1000 (Thermo Fischer #A11122), or rabbit α-Tom at 1:500 (clontech #632393).

Monoclonal antibody nc82 identifies Bruchpilot. Bruchpilot can serve as a general

neuropil marker because it is required in synaptic zones (Wagh et al. 2006). Larval

brains were collected from third instar larvae and fixed in 4% PFA. Primary antibodies:

mouse α-neuroglian (Developmental Studies Hybridoma Bank. #BP104) at 1:50 and

rabbit α-GFP at 1:500. Brains were washed 4x, 15min per wash while nutating with 2mL

PBT. 200µL secondary antibodies in PBT were then added for 4 hours at room

temperature and then 3 overnights at 4C while nutating upright. Secondary antibodies:

AF568 goat α-mouse (Life Technologies #A11031) at 1:400 and AF488 goat α-rabbit

(ThermoFischer #A11034) at 1:800. Tubes were protected from light at all times after

secondary antibodies had been added. Brains were washed again 4x, 15min per wash

while nutating with 2mL PBT. Then washed with 1xPBS and mounted using Vectashield

mounting media (Vector Labs #H-1000). Confocal images were taken with Leica800

microscope.

Single Sensillum Recordings

SSRs were performed as described in Lin et al 2015 (Lin and Potter 2015). GFP

labeled ab1 and ab3 sensilla were identified using a Zeiss AxioExaminer D1 compound

microscope with eGFP filter cube (FL Filter Set 38 HE GFP shift free). A glass recording

electrode filled with ringers solution (7.5g of NaCl+0.35g of KCl+0.279g of $CaCl_2$-$2H_2O$

in 1L of H_2O) was inserted into the base of the sensillum. To test ab1 (Gr21a) response,

CO_2 was delivered through a tube ending with a Pasteur pipette that was inserted for 1

second into a hole in a plastic pipette directed at the antenna. This plastic pipette

(Denville Scientific Inc, 10ml pipette) carried a purified continuous air stream (8.3 ml/s)

that used a stimulus controller (Syntech) at the time of CO_2 delivery to correct for the

increased air flow. To test ab3 (Or22a) response, 20 µl of E2-Hexenal or Isoamyl

acetate (diluted to 1% in mineral oil) was pipetted on a piece of filter paper (1X2 cm) in

a Pasteur pipette. The Pasteur pipette was then inserted into the hole of the plastic

pipette that carried continuous air stream to the antenna. For odorant delivery, the

stimulus controller (Syntech) was used to divert a 1 s pulse of charcoal-filtered air (5

ml/s) into the Pasteur pipette containing the odorant.

Signals were acquired and analyzed using AUTOSPIKE software (USB-IDAC System;

Syntech). Spikes were counted in a 500 ms window from 500 ms after CO_2 delivery and

multiplied by 2 to calculate spikes/second. Then, the spikes in 1000ms before CO_2

delivery were subtracted to calculate the increase in spike rate in response to CO_2

(Δspikes/second). For each genotype, 6 flies (4-8 days old) were tested, with 1-3

sensilla tested in each fly.

RESULTS

Design of GAL80 Constructs

The following criteria were used to choose OR promoters for the collection. i) The

ORs should be relevant to current research as shown by the number of studies that

used it. ii) The ORs should represent a variety of expression patterns (larval or adult,

antennae or maxillary palps, sensillary class etc.). iii) Finally, the ORs should reflect a

variety of different odorant response profiles. The promoter regions were defined based

largely on the work of Couto *et al*, 2005.

Equimolar expression of GAL4 and GAL80 is not always sufficient to effectively

eliminate GAL4 activity so the pBP-GAL80uW-6 vector was used. This vector contains a

modified GAL80 sequence, designed to increase the stability and expression of its gene

product (Pfeiffer et al. 2010). A few *OrX-GAL80*s were already made with this vector

and used effectively. (Gao, Clandinin, and Luo 2015) pioneered the technique by

creating a limited number of OrX-GAL80s. This work is a logical extension and makes

many additional *OrX-GAL80s* available for general use.

Testing GAL80 Efficacy and Specificity

GAL80 lines were created for the following odorant receptor promoters: Or7a,

Or10a, Or13a, Or19a, Or22a, Or22b, Or33c, Or35a, Or42a, Or42b, Or43b, Or47a,

Or56a, Or59b, Or59c, Or67a, Or67d, Or71a, Or82a, Orco, Or85a, Or85b, Or85c, and

Gr21a. To examine GAL4 subtraction *in vivo*, *OrX-GAL80* flies were crossed to flies with

the genotype *OrX-GAL4, UAS-GFP*. OSNs expressing the same OR can be identified

from their specific glomerulus in the antennal lobe (**Figure 1**). *OrX-GAL4, UAS-GFP*

flies show robust expression of the GFP reporter gene in their respective glomeruli.

However, when *OrX-GAL80* is added to the genotype, GFP expression is entirely

absent, indicating a robust antagonism of GAL4 activity (**Figure 3**). The efficacy of

Or22b-GAL80 could not be determined because the *OR22b-GAL4, UAS-GFP* control

did not show robust or reliable GFP signaling in the first place. The created *Or7a-*

GAL80 line was not effective at subtracting GFP signal. Though these lines are not

included in Figure 3, they will still be available in the Bloomington Stock Center. Several

of the GAL80 lines also have expression in larvae. GAL4 subtraction was examined in

larval brains using the *UAS-GFP* reporter gene. In larvae, GAL80 reduced but did not

eliminate GAL4 activity (**Figure S1a**).

The *OrX-GAL80* lines were checked to ensure they would not have aberrant

expression in untargeted OSN subtypes. The pBP-GAL80uW-6 vector contains a

Drosophila Synthetic Core Promoter (DSCP). DSCP is an effective means of using

enhancer elements to drive strong expression (Pfeiffer et al. 2008), but it could also

cause the GAL80s to have nonspecific or leaky expression. Therefore, a version of

pBP-GAL80uW-6 was cloned with the DSCP removed. However, when the DSCP was

absent, GAL80 expression was insufficient to subtract GAL4 activity (**Figure S1b**). A

few lines were tested to see if DSCP causes nonspecific GAL80 expression. For these

lines, an *OrY-GAL80* did not impede GAL4 activity of an *OrX-GAL4* neuron (**Figure**

S1c). Due to the uneven expression in an *Orco-GAL4, UAS-GFP* line, it could not be

determined if each OrX-GAL80 subtracts GAL4 from only one glomerulus in an

otherwise fully-labeled brain, but results shown in Figure S1c give reasonable

confidence that the GAL80s do not have widespread nonspecific expression. It can also

be noted that the GAL80 subtraction does not interfere with reporter gene expression in

a genetic system that does not use GAL4. When *Or22a-GAL80* is used in conjunction

with *Or22a-GFP*, containing no GAL4/UAS intermediary, the GFP is still expressed

(**Figure S1d**). These images, showing subtraction of reporter gene expression, confirm

that GAL4 activity is suppressed anatomically by the GAL80 lines.

To confirm GAL4 was suppressed physiologically by the GAL80s, Single

Sensillum Recordings (SSRs) were used to assay electrical activity of OSNs. *Gr21a-*

GFP was used to identify sensilla of interest without interfering with the

GAL4/UAS/GAL80 system. Gr21a neurons are housed in ab1 sensilla. Carbon Dioxide

exposure causes a robust response in Gr21a ab1C neurons (Hallem and Carlson 2006;

Hallem, Ho, and Carlson 2004; Fishilevich and Vosshall 2005; deBruyne:2001bs W. D.

Jones et al. 2007; Kwon et al. 2007; de Bruyne, Foster, and Carlson 2001). When

Gr21a-GFP flies were exposed to CO_2, their ab1C sensillar neurons showed robust

responses (mean Δspikes/s=88, N=8 sensilla). Kir2.1-containing neurons are expected

to show little to no spontaneous firing (Olsen, Bhandawat, and Wilson 2007; Hoare,

McCrohan, and Cobb 2008). Adding Kir2.1 to Gr21a neurons (genotype *Gr21a-GFP,*

Gr21a-GAL4, UAS-Kir2.1) greatly reduced spiking responses to CO_2 (mean

Δspikes/s=14, N=12 sensilla, *p*=0.01). When Gr21a-GAL80 was added (genotype

Gr21a-GFP, Gr21a-GAL4, UAS-Kir2.1, Gr21a-GAL80), responses to CO_2 were restored

(mean Δspikes/s=94, N=6 sensilla, *p*< 0.001. No significant difference from genotype

Gr21a-GFP, p= .26) (**Figure 4a**).

To make sure the system also worked for neurons expressing an OR protein (in

additional to a GR), SSR was also done with ab3 sensilla. Ab3 houses Or22a-

expressing neurons. This receptor is known to be activated by a diverse set of odorants,

including Isoamyl acetate and E2-hexenal (Hallem and Carlson 2006; Hallem, Ho, and

Carlson 2004; Fishilevich and Vosshall 2005; deBruyne:2001bs W. D. Jones et al.

2007; Kwon et al. 2007; de Bruyne, Foster, and Carlson 2001). A transgene was

necessary to visualize the neurons without interfering with the GAL4/UAS/GAL80

system, but *Or22a-GFP* was insufficiently bright to identify sensilla for SSR. A new

Or22a-LexA transgenic animal was therefore created using the same promoter that was

used to create the *Or22a-GAL80* gene (this line is also available through Bloomington).

When crossed to a *LexAop2-GFP* line, the ab3 sensilla showed bright fluorescence.

When *Or22a-LexA, LexAop2-GFP* flies were exposed to Isoamyl acetate or to E2-

hexenal, their sensillar neurons showed robust responses (mean Δspikes/s=67.86 and

27.71, N=7 and 7 sensilla, respectively). Unlike the Gr21a neurons, Kir2.1 expression in

the Or22a neurons effectively eliminated both spontaneous and odor-evoked activity.

(mean Δspikes/s=0, N=7 sensilla, p<.001 for both odorants). Both activities could be

restored with the addition of the *Or22a-GAL80* gene (Isoamyl acetate: mean Δspikes/s

=74.29, N=7 sensilla, p<.001; E2-Hexenal: mean Δspikes/s =25, N=7 sensilla, p<.001).

Neurons showed some low-level responses to mineral oil alone, the solvent used for the

odorants (**Figure 4b**). Only genotype 3 *Or22a-LexA, LexAop2-GFP, Or22a-GAL4, UAS-*

Kir2.1, Or22a-GAL80 showed significantly higher responses to mineral oil than

genotypes *Or22a-LexA, LexAop2-GFP* (p=.01) and *Or22a-LexA, LexAop2-GFP, Or22a-*

GAL4, UAS-Kir2.1 (p=.005), but the latter two genotypes showed no significant

response to mineral oil alone. The results in **Figure 4** confirm that GAL80 functions

effectively to prevent GAL4-induced activity in OSNs.

379   **DISCUSSION**

The collection of GAL80 lines subtracts GAL4 activity efficiently and specifically

in OSNs. In anatomical studies, reporter gene expression from the GAL4/UAS system is

suppressed. Neurons silenced with Kir2.1 expression have normal firing capacity

restored when GAL4 is antagonized using the GAL80 lines.

In behavioral assays, using a GAL80 transgene will be more flexible than mutant

lines and less cumbersome than crafting the required recombinants as the complexity of

the genotype increases. Though in some special circumstances, olfactory sensory

neurons have been shown to produce behaviors autonomously, this is not a widely

applicable principle, and further investigation upon this principle requires better tools.

For example, Fishilevish et al (2005) used larvae in their study to restore aversion with a

single functional OSN subtype, but the larval olfactory system may be fundamentally

different from adults in this respect., (Bhandawat et al. 2010) also showed that single

glomerular activity is sufficient to invoke a behavioral response, but that study was done

using an intact and fully functional olfactory background, so some neuronal cooperation

may still have occurred. (DasGupta and Waddell 2008) provided evidence that a single

functional OSN subtype is sufficient to learn odor discrimination, and Gao et al (2015)

gave convincing evidence of aversive restoration in adults with only one functional OSN.

However, the extent to which the restoration of single-OSN behavior depends on

the odorant and receptor used is still unknown. Only a small subset of receptors have

been tested. The current models of odor coding by the olfactory system predict that a

coordinated effort of many OSNs is usually required to produce a behavioral output.

Paired neurons in sensilla can affect the firing dynamics of their neighbors in the

periphery (Dobritsa et al. 2003; Su et al. 2012; Kazama and Wilson 2009), and

downstream neurons such as interneurons and projection neurons may rely on

synchronized input from multiple OSN types (Chou et al. 2010; Wilson 2011; Yaksi and

Wilson 2010; Hong and Wilson 2015; Kazama, Yaksi, and Wilson 2011; Olsen,

Bhandawat, and Wilson 2007; Ng et al. 2002; Acebes et al. 2011). GAL80 tools open

more possibilities to combinatorially activate subsets of neurons. The hope is that

additional researchers will use the reagents and validate them in their own assays.

Researchers encounter a significant technical obstacle to the understanding of

olfactory function if they need to create genotypes with small groups of interacting

neurons in isolation. The tools presented here facilitate the activation or deactivation of

combinations of particular neurons, thereby overcoming this obstacle. The lines are

available to order through Bloomington Stock Center.

AUTHOR CONTRIBUTIONS

Conceptualization, JE and IM; Methodology, JE; Formal Analysis, JE and AA;

Investigation, JE and AA; Writing—Original Draft, JE; Writing—Review and Editing, AA,

CP, IM; Visualization, JE and AA; Supervision, IM; Funding Acquisition, IM

ACKNOWLEDGEMENTS

Funding for this project comes from the National Science Foundation (MCB- 1413062).

The authors report no conflicts of interest. We thank Heather Dionne for cloning advice,

Yoshi Aso for the use of his behavioral setup and Janelia FlyLight for consulting on

imaging and immunostaining.

FIGURE LEGENDS

Figure 1: Olfactory Sensory Map. Each neuron in the olfactory system expresses one

type of odorant receptor (OR). Or22a (teal) and Or82a (gold) are used here as

examples. Neurons usually exist in pairs or groups in sensilla within the olfactory

organs—antenna or maxillary palp. Neurons expressing the same OR are distributed

throughout the periphery, but project their axons onto the same glomerulus in the

antennal lobe of the brain. For example, all Or22a-expressing neurons synapse onto the

DM2 glomerulus while all Or82a-expressing neurons synapse onto the VA6 glomerulus.

Figure 2: Advantages of using a GAL80 approach over a null mutation. a) **Current**

method with available reagents. In order to examine a single type of Olfactory

Sensory Neuron (OSN) without interference from other OSNs, one can use an Orco null

mutant. Without Orco, ORs cannot reach the cell membrane or function properly. Orco

mutants are mostly anosmic (unable to smell.) A single OR can then be restored using

two transgenes, *OrX-GAL4* and *UAS-Orco*. *Or22a-GAL4* is shown here as an example.

This fly may require the making and validating of one or more recombinant

chromosomes, since the Orco mutation must be homozygous. In more complicated

systems, e.g. restoring more than one OSN, multiple recombinants would need to be

made and validated at a cost of several months of crossing. b) **Using a GAL80**. GAL80

is a potent GAL4 inhibitor. All olfactory neurons could be silenced using any number of

transgenes in an *Orco-GAL4, UAS-effector* (such as *UAS-Kir2.1*) genotype. A single

OSN subtype can then be restored using an *OrX-GAL80* (such as *Or22a-GAL80*). This

system requires no recombinant creation, and is amenable to the use of various

effectors or additional transgenes without requiring recombinant construction. (Receptor

appearance, orientation, and heterodimerization is based on previous designs by

Neuhaus et al. (2005), Benton et al. (2006), Smart et al. (2008), and (Smart et al. 2008;

Neuhaus et al. 2005; Benton et al. 2006; Benton 2009)

Figure 3: OR-GAL80 reagents eliminate GAL4 activity. All antennal lobes are stained

with anti-nc82 (a general neuropil marker, grey) and anti-GFP (green). The orientation

of each image is dorsal-up, ventral-down, lateral-right, medial-left. Scale bars indicate

20μm. Each of the brains shown has the genotype *OrX-GAL4, UAS-GFP*. The specific

receptor promoter is given above each column. The top row in each set shows GFP

expression in these lines without GAL80. Notice how each neuron's target in the

antennal lobe glomeruli is expressing GFP. Each bottom row shows the brains

containing an additional *Or-GAL80* gene. Note how GAL80 effectively inhibits GAL4

activity, as seen by the elimination of GFP expression. The images are representative of

the 5-20 brains examined per genotype. GAL4 inactivation was 100% penetrant in one

day old female flies.

Figure 4: Olfactory neuron responses towards odors in Single Sensillum

Recordings (SSR). In box plots on the left, each circle shows response in an individual

sensillum, and filled squares indicate the means. On the right of each plot, example

SSR traces are shown for each genotype. (* indicates $.01>p>.005$, ** indicates $p<.001$)

a) **Ab1C SSR responses.** Ab1C neurons are visualized using the *Gr21a-GFP* gene.

Top: Sensilla respond strongly to CO_2, and adding *Kir2.1* reduces response to CO_2.

Response is restored when *Gr21a-GAL80* is added. Bottom: Air was used as a control

for CO_2 experiments. Air does not cause an odor-evoked neuronal response, and

adding the *Kir2.1* or *GAL80* genes does not affect the spontaneous signaling

responses. b) **Ab3 SSR responses.** Ab3 neurons are visualized using the *Or22a-LexA*

and LexAop2-GFP genes*.* Sensilla respond to Isoamyl Acetate and to E2-Hexenal.

Adding *Kir2.1* eliminates both odor-evoked and spontaneous activity in these neurons.

Spontaneous and odor-evoked activity is restored when *Or22a-GAL80* is added.

Odorants were diluted in mineral oil and neurons from the GAL80 restorative genotype

did show low-level responses to mineral oil alone (bottom).

486   **REFERENCES**

Abrams, J M. 1999. "An Emerging Blueprint for Apoptosis in Drosophila.." *Trends in Cell*

Biology 9 (11): 435–40.

Acebes, Angel, Alfonso Martín-Peña, Valérie Chevalier, and Alberto Ferrús. 2011.

"Synapse Loss in Olfactory Local Interneurons Modifies Perception.." *The Journal of*

Neuroscience : the Official Journal of the Society for Neuroscience 31 (8): 2734–45.

doi:10.1523/JNEUROSCI.5046-10.2011.

Baines, R A, S G Robinson, M Fujioka, J B Jaynes, and M Bate. 1999. "Postsynaptic

Expression of Tetanus Toxin Light Chain Blocks Synaptogenesis in Drosophila.."

Current Biology 9 (21): 1267–70.

Baines, Richard A, Jay P Uhler, Annemarie Thompson, Sean T Sweeney, and Michael

Bate. 2001. "Altered Electrical Properties in DrosophilaNeurons Developing Without

Synaptic Transmission." *Journal of Neuroscience* 21 (5). Society for Neuroscience:

1523–31. doi:10.1016/S0960-9822(99)80510-7.

Benton, R. 2009. "Evolution and Revolution in Odor Detection." *Science* 326 (5951):

382–83. doi:10.1126/science.1181998.

Benton, Richard, Kirsten S Vannice, Carolina Gomez-Diaz, and Leslie B Vosshall. 2009.

"Variant Ionotropic Glutamate Receptors as Chemosensory Receptors in

Drosophila." *Cell* 136 (1): 149–62. doi:10.1016/j.cell.2008.12.001.

Benton, Richard, Silke Sachse, Stephen W Michnick, and Leslie B Vosshall. 2006.

"Atypical Membrane Topology and Heteromeric Function of Drosophila Odorant

Receptors in Vivo." *PLoS Biology* 4 (2): e20–18. doi:10.1371/journal.pbio.0040020.

Bhandawat, V, G Maimon, M H Dickinson, and R I Wilson. 2010. "Olfactory Modulation

of Flight in Drosophila Is Sensitive, Selective and Rapid." *Journal of Experimental*

Biology 213 (21): 3625–35. doi:10.1242/jeb.040402.

Boyden, Edward S. 2011. "A History of Optogenetics: the Development of Tools for

Controlling Brain Circuits with Light.." *F1000 Biology Reports* 3 (11): 11.

doi:10.3410/B3-11.

Chen, Alex Y, Shouzhen Xia, Paul Wilburn, and Tim Tully. 2014. "Olfactory Deficits in

an Alpha-Synuclein Fly Model of Parkinson's Disease.." Edited by Mel B Feany.

PLoS ONE 9 (5): e97758. doi:10.1371/journal.pone.0097758.

Chen, M S, R A Obar, C C Schroeder, T W Austin, C A Poodry, S C Wadsworth, and R

B Vallee. 1991. "Multiple Forms of Dynamin Are Encoded by Shibire, a Drosophila

Gene Involved in Endocytosis.." *Nature* 351 (6327): 583–86. doi:10.1038/351583a0.

Chou, Ya-Hui, Maria L Spletter, Emre Yaksi, Jonathan C S Leong, Rachel I Wilson, and

Liqun Luo. 2010. "Diversity and Wiring Variability of Olfactory Local Interneurons in

the Drosophila Antennal Lobe." *Nature Neuroscience*, February. Nature Publishing

Group, 1–13. doi:10.1038/nn.2489.

Clyne, P J, C G Warr, M R Freeman, D Lessing, J Kim, and J R Carlson. 1999. "A Novel

Family of Divergent Seven-Transmembrane Proteins: Candidate Odorant Receptors

in Drosophila.." *Neuron* 22 (2): 327–38.

Couto, Africa, Mattias Alenius, and Barry J Dickson. 2005. "Molecular, Anatomical, and

Functional Organization of the Drosophila Olfactory System." *Current Biology* 15

(17): 1535–47. doi:10.1016/j.cub.2005.07.034.

DasGupta, Shamik, and Scott Waddell. 2008. "Learned Odor Discrimination in

Drosophila Without Combinatorial Odor Maps in the Antennal Lobe." *Current*

Biology 18 (21): 1668–74. doi:10.1016/j.cub.2008.08.071.

de Bruyne, M, P J Clyne, and J R Carlson. 1999. "Odor Coding in a Model Olfactory

Organ: the Drosophila Maxillary Palp.." *The Journal of Neuroscience : the Official*

Journal of the Society for Neuroscience 19 (11). Society for Neuroscience: 4520–32.

de Bruyne, Marien, Kara Foster, and John R Carlson. 2001. "Odor Coding in the

Drosophila Antenna." *Neuron* 30 (2): 537–52. doi:10.1016/S0896-6273(01)00289-6.

Dobritsa, Anna A, Wynand van der Goes van Naters, Coral G Warr, R Alexander

Steinbrecht, and John R Carlson. 2003. "Integrating the Molecular and Cellular

Basis of Odor Coding in the Drosophila Antenna.." *Neuron* 37 (5): 827–41.

Elmore, Tamara, R Ignell, John R Carlson, and Dean P Smith. 2003. "Targeted

Mutation of a Drosophila Odor Receptor Defines Receptor Requirement in a Novel

Class of Sensillum | Journal of Neuroscience." *The Journal of Neuroscience : the*

Official Journal of the Society for Neuroscience 23 (30). Society for Neuroscience:

9906–12. doi:10.1038/81774.

Fishilevich, Elane, Ana I Domingos, Kenta Asahina, Félix Naef, Leslie B Vosshall, and

Matthieu Louis. 2005. "Chemotaxis Behavior Mediated by Single Larval Olfactory

Neurons in Drosophila.." *Current Biology* 15 (23): 2086–96.

doi:10.1016/j.cub.2005.11.016.

Fishilevich, Elane, and Leslie B Vosshall. 2005. "Genetic and Functional Subdivision of

the Drosophila Antennal Lobe." *Current Biology* 15 (17): 1548–53.

doi:10.1016/j.cub.2005.07.066.

Gao, Xiaojing J, Thomas R Clandinin, and Liqun Luo. 2015. "Extremely Sparse

Olfactory Inputs Are Sufficient to Mediate Innate Aversion in Drosophila.." Edited by

Matthieu Louis. *PLoS ONE* 10 (4). Public Library of Science: e0125986.

doi:10.1371/journal.pone.0125986.

Giniger, E, S M Varnum, and M Ptashne. 1985. "Specific DNA Binding of GAL4, a

Positive Regulatory Protein of Yeast.." *Cell* 40 (4): 767–74.

Goldman, Aaron L, Wynand van der Goes van Naters, Derek Lessing, Coral G Warr,

and John R Carlson. 2005. "Coexpression of Two Functional Odor Receptors in

One Neuron." *Neuron* 45 (5): 661–66. doi:10.1016/j.neuron.2005.01.025.

Hallem, Elissa A, and John R Carlson. 2006. "Coding of Odors by a Receptor

Repertoire." *Cell* 125 (1): 143–60. doi:10.1016/j.cell.2006.01.050.

Hallem, Elissa A, Michael G Ho, and John R Carlson. 2004. "The Molecular Basis of

Odor Coding in the Drosophila Antenna." *Cell* 117 (7): 965–79.

doi:10.1016/j.cell.2004.05.012.

Hoare, D J, C R McCrohan, and M Cobb. 2008. "Precise and Fuzzy Coding by Olfactory

Sensory Neurons." *Journal of Neuroscience* 28 (39): 9710–22.

doi:10.1523/JNEUROSCI.1955-08.2008.

Hoare, Derek J, James Humble, Ding Jin, Niall Gilding, Rasmus Petersen, Matthew

Cobb, and Catherine McCrohan. 2011. "Modeling Peripheral Olfactory Coding in

Drosophila Larvae." Edited by Bradley Steven Launikonis. *PLoS ONE* 6 (8):

e22996–11. doi:10.1371/journal.pone.0022996.

Hodge, James J L. 2009. "Ion Channels to Inactivate Neurons in Drosophila." *Frontiers*

in Molecular Neuroscience 2: 1–10. doi:10.3389/neuro.02.013.2009.

Hong, Elizabeth J, and Rachel I Wilson. 2015. "Simultaneous Encoding of Odors by

Channels with Diverse Sensitivity to Inhibition." *Neuron* 85 (3). Elsevier Inc.: 573–

89. doi:10.1016/j.neuron.2014.12.040.

Invitrogen. 2012a. "pBAD/Thio His TOPO Manual," March, 1–74.

Invitrogen. 2012b. "pENTR™ Directional TOPO® Cloning Kits," March, 1–52.

Johns, D C, R Marx, R E Mains, B O'Rourke, and E Marbán. 1999. "Inducible Genetic

Suppression of Neuronal Excitability.." *Journal of Neuroscience* 19 (5): 1691–97.

Jones, P L, G M Pask, and D C Rinker. 2011. "Functional Agonism of Insect Odorant

Receptor Ion Channels." In. doi:10.1073/pnas.1102425108/-

/DCSupplemental/pnas.201102425SI.pdf.

Jones, Walton D, Pelin Cayirlioglu, Ilona Grunwald Kadow, and Leslie B Vosshall. 2007.

"Two Chemosensory Receptors Together Mediate Carbon Dioxide Detection in

Drosophila." *Nature* 445 (7123). Nature Publishing Group: 86–90.

doi:10.1038/nature05466.

Jung, Sarah Nicola, alexander borst, and Juergen Haag. 2011. "Flight Activity Alters

Velocity Tuning of Fly Motion-Sensitive Neurons.." *The Journal of Neuroscience :*

the Official Journal of the Society for Neuroscience 31 (25). Society for

Neuroscience: 9231–37. doi:10.1523/JNEUROSCI.1138-11.2011.

Kazama, H, E Yaksi, and R I Wilson. 2011. "Cell Death Triggers Olfactory Circuit

Plasticity via Glial Signaling in Drosophila." *Journal of Neuroscience* 31 (21): 7619–

30. doi:10.1523/JNEUROSCI.5984-10.2011.

Kazama, Hokto, and Rachel I Wilson. 2009. "Origins of Correlated Activity in an

Olfactory Circuit." *Nature Neuroscience* 12 (9): 1136–44. doi:10.1038/nn.2376.

Kitamoto, T. 2001. "Conditional Modification of Behavior in Drosophila by Targeted

Expression of a Temperature-Sensitive Shibire Allele in Defined Neurons.." *Journal*

of Neurobiology 47 (2): 81–92.

kitamoto, Toshihiro. 2002. "Conditional Disruption of Synaptic Transmission Induces

Male-Male Courtship Behavior in Drosophila.." *Proceedings of the National*

Academy of Sciences 99 (20): 13232–37. doi:10.1073/pnas.202489099.

Kreher, Scott A, Dennis Mathew, Junhyong Kim, and John R Carlson. 2008.

"Translation of Sensory Input Into Behavioral Output via an Olfactory System.."

Neuron 59 (1): 110–24. doi:10.1016/j.neuron.2008.06.010.

Krieger, J, O Klink, C Mohl, K Raming, and H Breer. 2003. "A Candidate Olfactory

Receptor Subtype Highly Conserved Across Different Insect Orders." *Journal of*

Comparative Physiology A 189 (7): 519–26. doi:10.1007/s00359-003-0427-x.

Kwon, Jae Young, Anupama Dahanukar, Linnea A Weiss, and John R Carlson. 2007.

"The Molecular Basis of CO_2 Reception in Drosophila.." *Proceedings of the National*

Academy of Sciences 104 (9): 3574–78. doi:10.1073/pnas.0700079104.

Larsson, Mattias C, Ana I Domingos, Walton D Jones, M Eugenia Chiappe, Hubert

Amrein, and Leslie B Vosshall. 2004. "Or83b Encodes a Broadly Expressed Odorant

Receptor Essential for Drosophila Olfaction." *Neuron* 43 (5): 703–14.

doi:10.1016/j.neuron.2004.08.019.

Lin, Chun-Chieh, and Christopher J Potter. 2015. "Re-Classification of Drosophila

Melanogaster Trichoid and Intermediate Sensilla Using Fluorescence-Guided Single

Sensillum Recording.." Edited by Matthieu Louis. *PLoS ONE* 10 (10). Public Library

of Science: e0139675. doi:10.1371/journal.pone.0139675.

Ma, Jun, and Mark Ptashne. 1987. "The Carboxy-Terminal 30 Amino Acids of GAL4 Are

Recognized by GAL80." *Cell* 50 (1): 137–42. doi:10.1016/0092-8674(87)90670-2.

Nakagawa, Takao, and Leslie B Vosshall. 2009. "Controversy and Consensus:

Noncanonical Signaling Mechanisms in the Insect Olfactory System." *Current*

Opinion in Neurobiology 19 (3): 284–92. doi:10.1016/j.conb.2009.07.015.

Neuhaus, Eva M, Günter Gisselmann, Weiyi Zhang, Ruth Dooley, Klemens Störtkuhl,

and Hanns Hatt. 2005. "Odorant Receptor Heterodimerization in the Olfactory

System of Drosophila Melanogaster." *Nature Neuroscience* 8 (1): 15–17.

doi:10.1038/nn1371.

Ng, Minna, Robert D Roorda, Susana Q Lima, Boris V Zemelman, Patrick Morcillo, and

Gero Miesenböck. 2002. "Transmission of Olfactory Information Between Three

Populations of Neurons in the Antennal Lobe of the Fly.." *Neuron* 36 (3): 463–74.

Nichols, Andrew S, and Charles W Luetje. 2010. "Transmembrane Segment 3 of

Drosophila Melanogaster Odorant Receptor Subunit 85b Contributes to Ligand-

Receptor Interactions.." *The Journal of Biological Chemistry* 285 (16): 11854–62.

doi:10.1074/jbc.M109.058321.

Olsen, Shawn R, Vikas Bhandawat, and Rachel I Wilson. 2007. "Excitatory Interactions

639 Between Olfactory Processing Channels in the Drosophila Antennal Lobe." *Neuron*

640 54 (1): 89–103. doi:10.1016/j.neuron.2007.03.010.

641 Pfeiffer, B D, A Jenett, A S Hammonds, T T B Ngo, S Misra, C Murphy, A Scully, et al.

642 2008. "Tools for Neuroanatomy and Neurogenetics in Drosophila." *Proceedings of*

643 *the National Academy of Sciences* 105 (28): 9715–20.

644 doi:10.1073/pnas.0803697105.

645 Pfeiffer, B D, T T B Ngo, K L Hibbard, C Murphy, A Jenett, J W Truman, and G M

646 Rubin. 2010. "Refinement of Tools for Targeted Gene Expression in Drosophila."

647 *Genetics* 186 (2): 735–55. doi:10.1534/genetics.110.119917.

648 Pulver, Stefan R, Stanislav L Pashkovski, Nicholas J Hornstein, Paul A Garrity, and

649 Leslie C Griffith. 2009. "Temporal Dynamics of Neuronal Activation by

650 Channelrhodopsin-2 and TRPA1 Determine Behavioral Output in Drosophila

651 Larvae.." *Journal of Neurophysiology* 101 (6). American Physiological Society:

652 3075–88. doi:10.1152/jn.00071.2009.

653 Robertson, Hugh M, Coral G Warr, and John R Carlson. 2003. "Molecular Evolution of

654 the Insect Chemoreceptor Gene Superfamily in Drosophila Melanogaster.."

655 *Proceedings of the National Academy of Sciences* 100 Suppl 2 (Supplement 2):

656 14537–42. doi:10.1073/pnas.2335847100.

657 Silbering, A F, R Rytz, Y Grosjean, L Abuin, P Ramdya, G S X E Jefferis, and R Benton.

658 2011. "Complementary Function and Integrated Wiring of the Evolutionarily Distinct

659 Drosophila Olfactory Subsystems." *Journal of Neuroscience* 31 (38): 13357–75.

660 doi:10.1523/JNEUROSCI.2360-11.2011.

661 Smart, Renee, Aidan Kiely, Morgan Beale, Ernesto Vargas, Colm Carraher, Andrew V

Kralicek, David L Christie, Chen Chen, Richard D Newcomb, and Coral G Warr.

2008. "Drosophila Odorant Receptors Are Novel Seven Transmembrane Domain

Proteins That Can Signal Independently of Heterotrimeric G Proteins." *Insect*

Biochemistry and Molecular Biology 38 (8): 770–80.

doi:10.1016/j.ibmb.2008.05.002.

Song, Z, and H Steller. 1999. "Death by Design: Mechanism and Control of Apoptosis.."

Trends in Cell Biology 9 (12): M49–M52.

Stocker, R F, M C Lienhard, A Borst, and K F Fischbach. 1990. "Neuronal Architecture

of the Antennal Lobe in Drosophila Melanogaster.." *Cell and Tissue Research* 262

(1). Springer-Verlag: 9–34. doi:10.1007/BF00327741.

Su, Chih-Ying, Karen Menuz, Johannes Reisert, and John R Carlson. 2012. "Non-

Synaptic Inhibition Between Grouped Neurons in an Olfactory Circuit." *Nature* 492

(7427). Nature Publishing Group: 66–71. doi:10.1038/nature11712.

Sweeney, Sean T, Kendal Broadie, John Keane, Heiner Niemann, and Cahir J O'Kane.

1995. "Targeted Expression of Tetanus Toxin Light Chain in Drosophila Specifically

Eliminates Synaptic Transmission and Causes Behavioral Defects." *Neuron* 14 (2):

341–51. doi:10.1016/0896-6273(95)90290-2.

van der Bliek, A M, and E M Meyerowitz. 1991. "Dynamin-Like Protein Encoded by the

Drosophila Shibire Gene Associated with Vesicular Traffic.." *Nature* 351 (6325):

411–14. doi:10.1038/351411a0.

Vosshall, L B, A M Wong, and R Axel. 2000. "An Olfactory Sensory Map in the Fly

Brain.." *Cell* 102 (2): 147–59.

Vosshall, L B, and B S Hansson. 2011. "A Unified Nomenclature System for the Insect

Olfactory Coreceptor." *Chemical Senses* 36 (6): 497–98.

doi:10.1093/chemse/bjr022.

Vosshall, Leslie B, Hubert Amrein, Pavel S Morozov, Andrey Rzhetsky, and Richard

Axel. 1999. "A Spatial Map of Olfactory Receptor Expression in the Drosophila

Antenna." *Cell* 96 (5): 725–36. doi:10.1016/S0092-8674(00)80582-6.

Wagh, Dhananjay A, Tobias M Rasse, Esther Asan, Alois Hofbauer, Isabell

Schwenkert, Heike Dürrbeck, Sigrid Buchner, et al. 2006. "Bruchpilot, a Protein with

Homology to ELKS/CAST, Is Required for Structural Integrity and Function of

Synaptic Active Zones in Drosophila.." *Neuron* 49 (6): 833–44.

doi:10.1016/j.neuron.2006.02.008.

Wilson, Rachel I. 2011. "Understanding the Functional Consequences of Synaptic

Specialization: Insight From the Drosophila Antennal Lobe." *Current Opinion in*

Neurobiology 21 (2). Elsevier Ltd: 254–60. doi:10.1016/j.conb.2011.03.002.

Yaksi, Emre, and Rachel I Wilson. 2010. "Electrical Coupling Between Olfactory

Glomeruli." *Neuron* 67 (6). Elsevier Inc.: 1034–47.

doi:10.1016/j.neuron.2010.08.041.

Figure 1

724　　Figure 2

728   Figure 3

729   Figure 4