Cytokinin signaling in *Mycobacterium tuberculosis*

Marie I. Samanovic[1,8], Hao-Chi Hsu[2,8], Marcus B. Jones[3,4], Victoria Jones[5], Michael R.

McNeil[5], Samuel H. Becker[1], Ashley T. Jordan[1], Miroslav Strnad[6], Changcheng Xu[7],

Mary Jackson[5], Huilin Li[2]* and K. Heran Darwin[1]*

[1]Department of Microbiology, New York University School of Medicine, New York, NY

10016, USA

[2]The Van Andel Research Institute, Grand Rapids, MI 49503, USA

[3]Human Longevity, Inc., San Diego, CA 92121, USA

[4]Present address: Regeneron Pharmaceuticals, Inc, Tarrytown, NY, USA

[5]Mycobacteria Research Laboratories, Department of Microbiology, Immunology and

Pathology, Colorado State University, Fort Collins, CO 80523-1682, USA

[6]Laboratory of Growth Regulators, Centre of the Region Haná for Biotechnological and

Agricultural Research, Institute of Experimental Botany ASCR and Palacký University,

78371 Olomouc, Czech Republic

[7]Biology Department, Brookhaven National Laboratory, Upton, New York 11973, USA

[8]These authors contributed equally to this manuscript.

Corresponding authors: heran.darwin@med.nyu.edu

21   Huilin.Li@vai.org

22 **Keywords**: *Mycobacterium tuberculosis*, cytokinins, cell envelope, proteasome, Pup,

23 antibiotic resistance

Summary

It was reported the human-exclusive pathogen *Mycobacterium (M.) tuberculosis* secretes cytokinins, which previously had only been known as plant hormones. Cytokinins are adenine-based signaling molecules in plants that have never been shown to participate in signal transduction in other kingdoms of life. Here, we show that cytokinins induce the strong expression of the *M. tuberculosis* gene, Rv0077c. We found that a TetR-like transcriptional regulator, Rv0078, directly repressed expression of the Rv0077c gene. Strikingly, cytokinin-induced expression of Rv0077c resulted in a loss of acid-fast staining of *M. tuberculosis*. Although acid-fast staining is thought to be associated with changes in the bacterial cell envelope and virulence, Rv0077c-induced loss of acid-fastness did not affect antibiotic susceptibility or attenuate bacterial growth in mice. Collectively, these findings show cytokinins signal transcriptional changes that affect the *M. tuberculosis* cell envelope, and that cytokinin signaling is no longer limited to the kingdom plantae.

Introduction

M. tuberculosis is the causative agent of tuberculosis, one of the world's leading causes

of mortality(WHO, 2017). For this reason, researchers are eager to identify pathways

that could be targeted for the development of new therapeutics to treat this devastating

disease. Among the current prioritized targets is the mycobacterial proteasome. *M.*

tuberculosis strains with defects in proteasome-dependent degradation are highly

attenuated in mice, partly because they are sensitive to nitric oxide (NO) (Cerda-Maira

et al., 2010, Darwin *et al.*, 2003, Gandotra *et al.*, 2010, Gandotra *et al.*, 2007,

Lamichhane *et al.*, 2006, Lin *et al.*, 2009). The NO-sensitive phenotype of mutants

defective for proteasomal degradation has been attributed to a failure to degrade an

enzyme called Log (<u>Lo</u>nely <u>g</u>uy), a homologue of a plant enzyme involved in the

synthesis of a family of N^6-substituted adenine-based molecules called cytokinins

(Samanovic *et al.*, 2015). The accumulation of Log in *M. tuberculosis* results in a

buildup of cytokinins, a break down product of which includes aldehydes that effectively

sensitize mycobacteria to NO (Samanovic *et al.*, 2015).

While we determined that a lack of proteasome-dependent degradation results in

cytokinin accumulation, we were left with more questions, namely, what is the function

of cytokinin production by *M. tuberculosis*? In plants, cytokinins are hormones that

regulate growth and development (Sakakibara, 2006). In addition, bacterial plant

pathogens and symbionts use cytokinins to facilitate the parasitism of their plant hosts

(Frebort *et al.*, 2011). Outside of the laboratory, *M. tuberculosis* exclusively infects

humans and is not known to have an environmental reservoir; therefore, it is unlikely *M.*

tuberculosis secretes cytokinins to modulate plant development. Instead, we

hypothesized that *M. tuberculosis*, like plants, uses cytokinins to signal intra-species

transcriptional changes to its benefit. Here, we show that cytokinins induce the

transcription of a gene of unknown function. Moreover, we identified and characterized

a TetR-like regulator that represses the expression of this gene. While we have not yet

identified an *in vivo* phenotype associated with this cytokinin-inducible gene, we found

that its expression altered the cell envelope of *M. tuberculosis*, changing its staining

properties. Collectively, these studies provide a foundation to characterize cytokinin

signaling in *M. tuberculosis* and other cytokinin-producing bacterial species.

Results

Cytokinins induce the specific and high expression of Rv0077c in *M. tuberculosis*

To test if a cytokinin (CK) could induce gene expression in *M. tuberculosis*, we grew

wild type (WT) *M. tuberculosis* H37Rv to mid-logarithmic phase and incubated the

bacteria for five hours with N^6-(Δ^2-isopentenyl)adenine (iP), one of the most abundantly

produced cytokinins in *M. tuberculosis* that is also commercially available(Samanovic *et*

al., 2015) (**see Materials and Methods--table supplement 1**). Using RNA sequencing

(RNA-Seq), we discovered the expression of four genes, Rv0076c, Rv0077c, Rv0078,

and *mmpL6*, was significantly induced upon iP treatment compared to treatment with

the vehicle control (dimethylsulfoxide, DMSO) (**Figure 1A--table supplement 2**).

Rv0077c is conserved among many mycobacterial species, while Rv0076c and Rv0078

are present only in several mycobacterial genomes (**Figure 1B**) (Lechat *et al.*, 2008).

Notably*, M. smegmatis*, a distant, non-pathogenic relative of *M. tuberculosis*, has a

weak homologue of Rv0077c and no conspicuous Rv0078 homologue (**Figure 1B**).

mmpL6 is one of 13 *mmpL* (mycobacterial membrane protein large) genes in *M.*

tuberculosis. In strain H37Rv, *mmpL6* is predicted to encode a 42 kD protein with five

trans-membrane-domains and is truncated compared to the same gene in ancestral

tuberculosis strains (Brosch *et al.*, 2002). Thus, it is unclear if *mmpL6* encodes a

functional protein in strain H37Rv.

Rv0077c was by far the most strongly induced gene in *M. tuberculosis* upon iP

treatment therefore we chose it for follow-up studies. We raised polyclonal antibodies to

recombinant Rv0077c protein and showed that protein levels were increased in *M.*

tuberculosis treated with iP for 24 hours (**Figure 1C,** first two lanes). Rv0077c was

barely detectable in cell lysates of bacteria that had not been incubated with iP and was

undetectable in a strain with a transposon insertion mutation in Rv0077c (**Figure 1C,**

center two lanes). Rv0077c protein was restored to WT levels in the mutant upon

complementation with an integrative plasmid encoding Rv0077c expressed from its

native promoter (**Figure 1C,** last two lanes). We also found a dose-dependent induction

of Rv0077c production using iP concentrations from 1 nM to 100 µM (**Figure 1D**).

We next synthesized and tested if the most abundantly produced cytokinin in *M.*

tuberculosis, 2-methyl-thio-iP (2MeSiP) (Samanovic *et al.*, 2015), could also induce

Rv0077c production. 2MeSiP strongly induced Rv0077c production (**Figure 1E,** lane 4).

Importantly, we did not observe induction of Rv0077c when we incubated the bacteria

with the appropriate cytokinin riboside (R) precursors iPR or 2MeSiPR (**Figure 1E,**

lanes 3 and 5). Similarly, adenosine monophosphate (AMP) or the closely related

molecule adenine could not induce Rv0077c expression (**Figure 1E,** lanes 6 and 7). We

hypothesized while adenine could not induce Rv0077c expression, at high enough

concentrations it could possibly inhibit Rv0077c induction by competing with cytokinin

for access to a transporter or receptor. Indeed, adenine reduced the induction of

Rv0077c by iP in a dose-dependent manner (**Figure 1F**).

Identification of an operator for the TetR-like transcriptional repressor Rv0078

Rv0077c is divergently expressed from Rv0078; the proposed translational start codons

for these genes are separated by 61 base pairs therefore we hypothesized that Rv0078

encodes a repressor of Rv0077c expression. We identified the promoter regions for

each gene by performing rapid amplification of 5' complementary DNA ends (5'RACE)

analysis for Rv0077c and Rv0078 and determined the likely start of transcription for one

gene was within the 5' untranslated region of the other gene (**Figure 2A**). We sought to

identify an operator sequence by using an electrophoretic mobility shift assay (EMSA,

Figure 2B). We narrowed down a putative Rv0078 binding site to a region overlapping

the proposed starts of transcription (+1) of both Rv0077c and Rv0078 (**Figure 2A**, red

box). TetR-like regulators generally bind to inverted repeat sequences; we identified a

21-base pair (bp) sequence within -14/+19 of Rv0077c containing two 10-bp inverted

repeat sequences. Mutagenesis of the probe in the repeat sequences disrupted Rv0078

binding to the DNA (**Figure 2B**). Importantly, the binding sequence overlaps the

putative transcriptional start sites of both genes, suggesting that expression of both

genes is repressed by Rv0078. Notably, the addition of iP did not result in the release of

Rv0078 from the DNA probe (**Figure 2B--figure supplement 1**). Therefore, cytokinins

do not appear to directly bind to Rv0078 to induce gene expression.

Based on the 5'RACE analysis we were able to delete and replace most of the

Rv0078 gene with the hygromycin-resistance gene (*hyg*) without disrupting the

promoter of Rv0077c in *M. tuberculosis* H37Rv. A ΔRv0078::*hyg* strain displayed

constitutively high expression of Rv0077c irrespective of the presence of cytokinin,

supporting a model where Rv0078 directly represses Rv0077c expression (**Figure 2C**).

A single copy of Rv0078 expressed from its native promoter restored iP-regulated

control of Rv0077c in this strain (**Figure 2C,** lanes 5 and 6). Interestingly, in the process

of making the complementation plasmid, we acquired a random mutation (likely

generated during PCR) in Rv0078 that changed a tryptophan to arginine (W100R); this

allele was unable to complement the ΔRv0078::*hyg* strain (**Figure 2C,** lanes 7 and 8).

Two Rv0078 dimers bind to one operator

To gain an understanding of how Rv0078 represses gene expression, we solved the

crystal structure of Rv0078 to 1.85 Å by single-wavelength anomalous dispersion

method. As previously reported, Rv0078 forms dimers resembling other TetR-like

proteins (Orth *et al.*, 2000, Wohlkonig *et al.*, 2017). Unlike the canonical TetR binding

site, *tetO*, which is 15 bp long, the Rv0078 binding site is 21 bp long suggesting Rv0078

binds to DNA differently than TetR. We performed an isothermal titration calorimetry

(ITC) experiment and determined that the apparent K_D of Rv0078 for operator DNA was

357 nM (**Figure 3A**). The estimated stoichiometry of DNA duplex to Rv0078 dimer was

0.66, suggesting that two Rv0078 dimers bind to one operator sequence.

We next co-crystallized Rv0078 with a 23-bp DNA fragment that was extended

one base pair at each end of the 21-bp operator sequence identified in Figure 2 (-8/+13

of Rv0077c) and solved the structure to 3.0 Å resolution (**Table supplement 3**). As

predicted by the ITC experiments, we observed two Rv0078 dimers bound to a DNA

duplex where each dimer bound on opposite sides of the DNA (**Figure 3B**). The top or

"central" dimer bound the DNA palindrome symmetrically while the bottom or

"peripheral" dimer bound to DNA off-center, staggered from the central dimer binding

site by seven base pairs. Thus, the longer-than-canonical (i.e., *tetO*) binding site is

required for accommodating two Rv0078 dimers, unlike TetR, which binds to *tetO* as a

single dimer.

Rv0078 has an N-terminal DNA-binding domain (DBD) and a C-terminal ligand-

binding domain (LBD) (**Figure 3B-D**). The DBD is comprised of three α-helices (α1 -

α3), and within this domain, α2 and α3 form a helix-turn-helix (HTH) motif that directly

contacts DNA. The LBD is formed by six α-helices (α4 - α9). The four Rv0078 protein

structures within a complex were nearly the same; the root mean square deviation

(rmsd) ranged from 0.07 Å to 0.17 Å in a pair-wise superimposition. The Rv0078 dimer

structure is held together by α8 and α9 of each monomer, forming a four-helix bundle.

The two dimer structures were highly similar with an rmsd of 0.14 Å. DNA binding

induced significant conformational changes across the Rv0078 dimer structure, with an

rmsd of 2.13 Å when compared to the DNA-free dimer. In particular, the two α3 helices

move towards each other by ~6 Å, in order to reduce their distance to 36.4 Å and fit in

the DNA major grooves (**Figure 3C**). In the LBD, the ligand entry between helices α4

and α5 is open and the ligand-binding pocket is empty. It appears that changes initiated

in the DBD regions upon binding to the DNA are transmitted to the LBD via the DBD-

LBD interface that involves extensive interactions, including two salt bridges and six H-

bonds (**Figure 3D**).

The ligand-binding pocket of Rv0078 is largely hydrophobic. Within this pocket,

we observed an elongated density resembling a long aliphatic chain of a fatty acid

(**Figure supplement 2**). Gas chromatograph-mass spectrometry of the compounds

extracted from Rv0078 purified from *E. coli* revealed fatty acids commonly found in this

organism, and a palmitate molecule fit well with the electron density (**Figure**

supplement 2A-E). A fatty acid carboxylate formed a hydrogen bond with Rv0078 Glu-

70 (**Figure 3E--figure supplement 2C**), and the long alkyl chain had numerous

hydrophobic interactions within the extended ligand pocket. When we purified Rv0078

under denaturing conditions and refolded the protein to remove the lipid, we observed

the same EMSA results as we observed with protein purified under native conditions

(data not shown), suggesting this fatty acid is unlikely to be the native Rv0078 ligand;

however, these data may suggest that the native ligand is fatty acid-like. Interestingly,

we found that Trp100 faces the ligand-binding pocket (**Figure 3E**), which suggests

Trp100 interacts with the natural ligand. This hypothesis is supported by our data

showing an Rv0078$_{W100R}$ mutant was unable to complement the Rv0078-deletion mutant

strain (**Figure 2C, lane 8**).

In addition to characterizing the LBD, we identified nine amino acids that were

important for interacting with DNA (**Fig. 4A-E**). Specifically, the hydroxyl groups of

Thr37, Thr47, and Tyr52 interacted strongly with DNA phosphates at a distance of 2.6

Å. Arg48 is the only residue that interacted with DNA by recognizing a guanine base at

a distance of 2.6 Å. To further examine the importance of these residues, we introduced

single amino acid substitutions (T37V, T47V, R48M, and Y52F) or double mutations

(T47V, R48M) into Rv0078 and performed EMSA assays. While all of the mutant

proteins were soluble and behaved like the WT protein in solution, Rv0078$_{R48M}$ did not

bind to DNA, and the other mutant proteins bound to DNA with reduced affinities

(**Figure 4F**).

To examine the DNA sequence specificity of Rv0078, we synthesized three

EMSA probes by changing the Rv0078 Arg48-interacting guanines to adenines (G-to-A)

in the central dimer binding region (probe S1), the peripheral dimer-binding region

(probe S2), or both dimer-binding regions (probe S3). These G-to-A substitutions either

abolished or compromised Rv0078 binding to DNA, affirming the critical role of

guanines in the binding site (**Figure 4G**). Notably, substitutions in probe S1 entirely

abolished Rv0078 binding, while substitutions in S2 retained partial gel shift at a high

concentration (white arrowhead in **Figure 4G**). This observation suggests that the

binding of the two Rv0078 dimers is cooperative, with the central dimer likely the first

one to bind to DNA.

Constitutive expression of Rv0077c does not affect antibiotic susceptibility or

virulence in mice

During our ongoing studies, a report was published on the identification of a small

molecule of the spiroisoxazoline family, SMARt-420, which strongly induces the

expression of the Rv0077c orthologue *bcg_0108c* in *M. bovis* Bacille Calmette-Guerin

(Blondiaux *et al.*, 2017). Using x-ray crystallography and surface plasmon resonance

techniques, the authors of this study found SMARt-420 binds to Rv0078 to derepress

binding from the Rv0077c promoter (Blondiaux *et al.*, 2017, Wohlkonig *et al.*, 2017).

SMARt-420 was identified in a search for compounds that boost the efficacy of the

second-line tuberculosis drug ethionamide (ETH). ETH is a pro-drug that is activated by

the mono-oxygenase EthA, which transforms ETH into highly reactive intermediates.

Activated ETH and nicotinamide adenine dinucleotide form a stable adduct, which binds

to and inhibits InhA, an essential enzyme needed for mycolic acid synthesis in

mycobacteria (DeBarber *et al.*, 2000, Vannelli *et al.*, 2002). While spontaneous

inactivating mutations in *ethA* result in resistance to ETH, it was proposed that the

induction of Rv0077c expression could bypass the need for EthA and transform ETH

into its toxic form (Blondiaux *et al.*, 2017). Based on this study, we predicted that an

Rv0078 mutant of *M. tuberculosis*, which expresses high levels of Rv0077c, should be

hypersensitive to ETH compared to the parental strain H37Rv. However, we observed

either little to no significant change in the 50% minimum inhibitory concentration (MIC_{50})

of ETH between the WT and ΔRv0078::*hyg* strains (**Table supplement 4**). We also

tested if the constitutive expression of Rv0077c changed the susceptibility of *M.*

tuberculosis to other antibiotics, including two cell wall synthesis inhibitors. We

observed no differences in the MIC_{50} of these antibiotics between the WT and

ΔRv0078::*hyg* strains (**Table supplement 4**).

We also tested if either the Rv0077c or Rv0078 mutant had growth defects *in*

vivo compared to the WT H37Rv strain. We infected C57BL/6J mice by a low-dose

aerosol route with the parental, mutant, and complemented mutant strains, as well as

with the Rv0078 mutant transformed with the $Rv0078_{W100R}$ allele. Interestingly, none of

the strains revealed a difference in growth or survival compared to WT *M. tuberculosis*

in mice as determined by the recovery of colony forming units (CFU) from the lungs and

spleens (**Figure 5--figure supplement 3**).

Expression of Rv0077c alters acid-fast staining of *M. tuberculosis*

To gain insight into the function of Rv0077c, we performed metabolomics analysis of

strains expressing Rv0077c to potentially determine how its presence alters bacterial

physiology. We prepared total cell lysates of WT and Rv0077c mutant strains treated

with or without 100 µM iP for 24 hours (see Materials and Methods). From a total of 337

detectable metabolites, we observed a significant change in 24 molecules after the

addition of iP to WT *M. tuberculosis*. Seventeen metabolites showed a consistent

difference between samples in which iP was added and these changes disappeared in

an Rv0077c-disrupted strain, suggesting the changes were specifically due to the

presence of Rv0077c (**Table supplement 4**, highlighted in yellow). We observed an

increased abundance of several phospholipids and a decrease in a major precursor of

peptidoglycan, N-acetyl-glucosamine-1-phosphate (GlcNAc1P). We therefore

hypothesized that Rv0077c modified one or more components of the cell envelope.

Microscopic examination of Ziehl-Neelsen (ZN) stained WT *M. tuberculosis* treated with

iP showed a loss of acid-fast staining (**Figure 6**, panel a compared to b), and this

phenotype depended on the presence of Rv0077c (**Figure 6**, panel c compared to d).

Complementation of the mutation with Rv0077c alone restored the iP-induced loss of

acid-fast staining (**Figure 6**, panels e and f). Deletion of Rv0078 resulted in a

constitutive loss of staining, irrespective of the presence of iP (**Figure 6**, panels g and

h). Complementation of the Rv0078 deletion with the WT gene restored iP-control of

loss of acid-fast staining (**Figure 6**, panels i and j) but complementation with

Rv0078$_{W100R}$, which could not fully de-repress Rv0077c expression in the presence of iP

(**Figure. 2C**), could not restore iP-induced loss of acid-fast staining (**Figure 6**, panels k

and l). Mixing and simultaneous staining of the Rv0077c and Rv0078 mutants further

showed the staining differences were not a result of a technical artifact (**Figure 6,** panel

m).

Acid-fast staining is primarily thought to be associated with mycolic acids;

alterations in mycolic acid synthesis result in negative effects on cell growth *in vitro* and

in vivo(Barkan *et al.*, 2009, Bhatt *et al.*, 2007). We used thin layer and liquid or gas

chromatography coupled to mass spectrometry to analyze the fatty acid, mycolic acid

and lipid contents of the WT, Rv0077c and Rv0078 strains and observed no significant

qualitative or quantitative differences between strains (**Figure supplement 4, table**

supplement 6). In particular, the expression of Rv0077c did not conspicuously modify

the chain length of mycolic acids, their cyclopropanation or the relative abundance of

keto- to alpha- mycolates, and did not alter the wax ester and triglyceride content of the

strains that have been previously been linked to acid-fast staining (Bhatt *et al.*, 2007,

Deb *et al.*, 2009).

Discussion

Our studies are the first to demonstrate CKs induce robust and specific transcriptional

and physiologic changes in a bacterial species. *M. tuberculosis* treated with CK

expressed high levels of Rv0077c, which altered its metabolome and staining

properties. Despite these changes, we did not observe an effect on the susceptibility of

these bacteria to antibiotics of different classes or on survival in mice. In addition we

found the transcriptional regulator Rv0078 represses Rv0077c expression in the

absence of CK, and we defined the operator to which Rv0078 binds. We determined

that two dimers of Rv0078 bind to inverted repeats in the intergenic region between

Rv0077c and Rv0078, which likely represses the expression of both genes.

In plants, CKs are sensed by membrane receptors related to the sensors of

bacterial two-component systems (TCS) (To & Kieber, 2008, Inoue *et al.*, 2001, Steklov

et al., 2013). In these systems, CKs interact with membrane receptors with a CHASE

(cyclases/histidine kinases associated sensing extracellular) domain (Mougel & Zhulin,

2001, Anantharaman & Aravind, 2001). As in bacteria, the ligand-receptor interaction

stimulates a phospho-relay that ultimately leads to the phosphorylation of a response

regulator protein, which either represses or activates gene expression. *M. tuberculosis*

has at least 12 known TCS, and none has a predicted CHASE domain (L. Aravind,

February 2012; and I. Jouline, August 2017, personal communications). Thus, it

remains to be determined if a TCS sensor protein is involved in CK signal transduction

in *M. tuberculosis*.

It also remains to be determined what the natural ligand is for Rv0078; while

SMARt-420 binds robustly to Rv0078 (Wohlkonig *et al.*, 2017, Blondiaux *et al.*, 2017)

the cytokinin iP did not in our studies. This result may not be surprising when

considering SMARt-420 and cytokinins bear no resemblance to each other.

Furthermore, we could not co-crystalize iP with Rv0078 (data not shown). We

hypothesize that the interaction of CK with an unknown receptor or enzyme leads to the

synthesis of a small molecule (e.g., a lipid) that binds to Rv0078, resulting in the

derepression of Rv0077c. This possibility may be supported by our observation that

there is an increase in several phospholipids in iP-treated *M. tuberculosis*. We are

currently working to identify factors required for CK-mediated gene induction.

Rv0077c is predicted to have an α/β hydrolase fold (Soding *et al.*, 2005), which

can have a variety of substrate types (Nardini & Dijkstra, 1999). Based on our

metabolomics and microscopy studies, we predict Rv0077c targets one or more

components of the cell envelope. The identification of the natural target of Rv0077c may

help provide new insight into the elusive molecular basis of acid-fastness of this human-

exclusive pathogen. Our results may also indicate that acid fast staining can be affected

by different cell envelope chemistries in addition to changes in mycolic acid structure or

content.

While we did not observe any differences in bacterial burden in mice infected

with strains that either were disrupted for or constitutively expressed Rv0077c, it is

possible that Rv0077c function might only be needed during very late or specific stages

of infection. This idea may be supported by a previous observation that the cytokinin

synthase *log* gene in *M. marinum* is specifically expressed in late granulomas of

infected frogs (Ramakrishnan *et al.*, 2000). Furthermore, mice might not provide an

optimal model to observe a role for this pathway in tuberculosis.

A previous report suggested the expression of Rv0077c increases the sensitivity

of *M. tuberculosis* to the antibiotic ETH. ETH must be activated by a mono-oxygenase,

EthA, in order for it to be toxic to *M. tuberculosis* (DeBarber *et al.*, 2000, Vannelli *et al.*,

2002). We found the expression of Rv0077c did not confer increased susceptibility to

ETH or any other antibiotic we tested, an observation consistent with the likelihood that

Rv0077c is not a mono-oxygenase. It is possible that the effects of SMARt-402 is

growth-condition dependent, that this molecule affects another pathway to either

increase the susceptibility of *M. tuberculosis* to ETH, or that another SMARt-420-

induced enzyme in addition to Rv0077c synergize together to activate ETH (A. Baulard,

personal communication). Irrespective of these possibilities, it is unlikely that Rv0077c

has a considerable role in the activation of ETH. While a previous report named

Rv0077c and Rv0078, EthA2 and EthR2, respectively, we propose to rename them

"LoaA" and "LoaR" for "loss of acid-fast staining A and Repressor" due to the lack of

association of these proteins with ETH susceptibility.

Finally, our studies have opened the door to the possibility that numerous

commensal and pathogenic microbes (including fungi) use cytokinins for intra- or inter-

species communication in complex systems such as the gut microbiome. The

identification of one or more CK receptors and the signal transduction pathway that

leads to the induction of Rv0077c expression will likely lay the foundation for

understanding CK signaling in hundreds of bacterial species.

Author Contributions M.I.S. and K.H.D. performed *in vitro* and *in vivo M. tuberculosis*

work. H.C.H. and H.L. determined the structure of Rv0078 and performed the

mutagenesis and EMSA assays. M.B.J. performed the RNA-Seq analysis. S.H.B. and

A.T.J. performed antibiotic susceptibility assays. M.S. provided cytokinins and their

ribosides. V.J., M.R.M. and M.J. performed the lipid analysis. C.X. performed mass

spectrometry of fatty acids extracted from purified Rv0078 protein. M.S. synthesized

cytokinins and their precursors. M.I.S., H.C.H., H.L. and K.H.D. wrote the manuscript.

Acknowledgments We thank A. Darwin and V. Torres for critical review of a draft

version of this manuscript. We thank S. Zhang for assistant in preparing reagents. This

work was supported by NIH grant R01 HL092774 awarded to K.H.D. K.H.D. holds an

Investigators in the Pathogenesis of Infectious Disease Award from the Burroughs

Wellcome Fund. M.I.S. and S.H.B were supported by the Jan T. Vilcek Endowed

Fellowship fund. S.H.B. is also supported by NIH grant T32 AT007180. H.C.H. and H.L.

were supported by R01 AI070285. M. Strnad was supported by LO1204 from the

National Program of Sustainability I. We thank A. Liang and Y. Deng of the Microscopy

Laboratory at New York University Langone Medical Center for assistance with

microscopy and imaging. Diffraction data for this study were collected at the Lilly

Research Laboratories Collaborative Access Team (LRL-CAT) beamline and the Life

Sciences Collaborative Access Team (LS-CAT) beamline at the Advanced Photon

Source (APS), Argonne National Laboratory. APS was supported by the U.S.

Department of Energy, Office of Science, Office of Basic Energy Sciences, under

Contract DE-AC02-06CH11357. Use of the LRL-CAT beamline at Sector 31 of the APS

was provided by Eli Lilly Co., which operates the facility.

Materials and Methods

Bacterial strains, plasmids, primers, chemicals, and culture conditions. Bacterial

strains, plasmids, and primer sequences used in this study are listed in Table

supplement 1. All primers for cloning and sequencing were from Invitrogen, Inc. *M.*

tuberculosis strains were grown in Middlebrook 7H9 broth (Difco) supplemented with

0.2% glycerol, 0.05% Tween-80, 0.5% fraction V bovine serum albumin, 0.2% dextrose

and 0.085% sodium chloride ("7H9c"). *M. tuberculosis* cultures were grown without

shaking in 25 or 75 cm^2 vented flasks (Corning) at 37 ºC. 7H11 agar (Difco)

supplemented with 0.5% glycerol and BBL TM Middlebrook OADC enrichment (BD) was

used for growth on solid medium ("7H11"). *M. tuberculosis* was transformed as

described(Hatfull & Jacobs, 2000). *E. coli* strains used for cloning and expression were

grown in LB-Miller broth (Difco) at 37 °C with aeration on a shaker or on LB agar. *E. coli*

strains were chemically transformed as previously described(Sambrook *et al.*, 1989).

The final concentrations of antibiotics used for *M. tuberculosis* growth: kanamycin, 50

µg/ml; hygromycin, 50 µg/ml; streptomycin, 25 µg/ml; and for *E. coli*: hygromycin, 150

µg/ml; kanamycin, 100 µg/ml; and streptomycin 50 µg/ml. AMP, adenine and iP were

purchased from Sigma. iPR, 2MeSiP and 2MeSiPR were synthesized as previously

described(Sugiyama & Hashizume, 1978). The purity of the synthesized cytokinin and

derivatives was >98% for each as determined by high performance liquid

chromatography and mass spectrometry.

The Rv0077c::MycoMarT7 mutant was isolated from a library of ordered

transposon insertion mutants as previously described(Darwin *et al.*, 2003, Festa *et al.*,

2007). The ΔRv0078::*hyg* mutant was made by deletion-disruption mutagenesis as

described in detail elsewhere using pYUB854(Festa *et al.*, 2011, Bardarov *et al.*, 2002).

Protein purification and immunoblotting. DNA sequence encompassing the full-

length Rv0078 or Rv0077c gene was cloned into pET24b(+) vector using primers listed

in Table supplement 1. Recombinant proteins were produced in *E. coli* ER2566 and

purified under native conditions for Rv0078 and denaturing conditions for Rv0077c

according to the manufacturer's specifications (Qiagen). Polyclonal rabbit antibodies

were raised by Covance (Denver, PA). For all immunoblots, cell lysates or purified

proteins were separated sodium dodecyl sulfate polyacrylamide gel electrophoresis

(SDS-PAGE); transferred to nitrocellulose and incubated with rabbit polyclonal

antibodies to the protein of interest at 1:1000 dilution in 3% bovine serum albumin in

TBST (25 mM Tris-HCl, pH 7.4, 125 mM NaCl, 0.05% Tween 20, pH 7.4). Equal loading

was determined by stripping the nitrocellulose membranes with 0.2 N NaOH for 5 min,

rinsing, blocking and incubating the nitrocellulose with polyclonal rabbit antibodies to

dihydrolipoamide acyltransferase (DlaT)(Tian *et al.*, 2005). Horseradish peroxidase

conjugated anti-rabbit antibody (GE-Amersham Biosciences) was used for

chemiluminescent detection (SuperSignal West Pico; ThermoScientific).

For crystallography studies of Rv0078-His$_6$, bacteria were grown at 37°C to an

OD_{600} = 0.5-0.6 before being induced with 0.5 mM IPTG and incubated at 16°C

overnight. After harvesting by centrifugation, cells were lysed by passing through a

microfluidizer cell disruptor in 10 mM potassium phosphate, pH 8.0, 10 mM imidazole,

and 500 mM NaCl. The homogenate was clarified by spinning at 27,000 *g* and the

supernatant was applied to a HiTrap-Ni column (GE Healthcare) pre-equilibrated with

the lysis buffer. Histidine-tagged protein was eluted with a 10–300 mM imidazole

gradient in 10 mM potassium phosphate, pH 8.0, containing 300 mM M NaCl. The

Rv0078 fractions were further purified by a Superdex 75 column (16 x 1000 mm, GE

Healthcare) pre-equilibrated with 20 mM potassium phosphate, pH 8.0, and 300 mM

NaCl. The purified Rv0078 was concentrated to 40 mg/ml for crystallization screen.

RNA-Seq and 5'RACE. Three biological replicate cultures of WT *M. tuberculosis* were

grown to an OD_{580} ~1 and incubated in 100 μM iP or DMSO (control) for five hours.

Cells were harvested and RNA was purified as described previously (Festa *et al.*, 2011).

Briefly, an equal volume of 4 M guanidinium isothiocyanate, 0.5% sodium N-lauryl

sarcosine, 25 mM trisodium citrate solution was added to cultures to arrest transcription.

RNA was isolated with Trizol Reagent (Invitrogen) and further purified using RNeasy

Miniprep kits and DNase I (Qiagen). Transcriptome profiling by RNA-seq was performed

and analyzed as follows: RNA from *M. tuberculosis* cultures were extracted for library

construction. Libraries were constructed and barcoded with the Epicentre ScriptSeq

Complete Gold low input (Illumina, Inc) and sequenced on Illumina HiSeq 2000

sequencer using version 3 reagents. Unique sequence reads were mapped to the

corresponding reference genome and RPKM values were calculated in CLC (CLC

version 7.0.4). Genes with significantly different RPKM values were identified using the

Significant Analysis for Microarray (SAM) statistical analysis component of MeV(Saeed

et al., 2003).

5'RACE was performed as described by the manufacturer (Invitrogen). Briefly, 1

μg of RNA was used as template for cDNA production using a reverse primer 150–300

bp downstream of annotated translational start sites. A 3' poly-C tail was added to

cDNA by recombinant Tdt. The cDNA was then amplified using a nested reverse primer

and a primer that anneals to the poly-C tail. Products were cloned and sequenced.

Likely transcriptional start sites were selected based on clones that had the most

nucleotide sequence upstream of the start codon.

EMSA. A series of double stranded DNA probes consisting of sequences in the

intergenic region between Rv0077c and Rv0078 were generated by annealing two

complementary oligonucleotides and a 5'-end IRDye700-labeled 14-nucleotide

oligomers (5'dye-GTGCCCTGGTCTGG-3') (Integrated DNA Technologies). Binding

assays were performed by incubating 100 nM of probes and various concentrations of

Rv0078 at room temperature for 30 min in 20 mM HEPES, pH 7.5, 3 mM DTT, 0.1 mM

EDTA, 100 mM KCl, 5% glycerol, 5 mg/ml BSA, 10 mM MgCl$_2$, and 0.25% Tween 20,

and were subsequently resolved in 6% polyacrylamide gels in 0.5 × TBE buffer. Mobility

shifts of protein-DNA complex were visualized in LI-COR Odyssey imager.

Crystallization and structure determination. DNA-free Rv0078 crystals were

obtained by screening at 20 °C using the sitting-drop vapor diffusion method. The C2

space group crystals were grown in 0.1 M sodium cacodylate, pH 6.4, and 1.3 M

Lithium sulfate. SeMet substituted Rv0078 crystals with C2 space group were grown in

in 0.1 M sodium cacodylate, pH 6.6, 1.3 M Lithium sulfate, 0.2 M magnesium sulfate,

and 2% PEG400. Diffraction data to a resolution of 1.85 Å were collected at the Lilly

Research Laboratories Collaborative Access Team (LRL-CAT) beamline of Advanced

Photon Source (APS), Argonne National Laboratory, and were processed with Mosflm

software(Winn *et al.*, 2011). The program Hybrid-Substructure-Search in the Phenix

package was used to locate the Se sites and the initial phasing was carried out using

the program Autosol of Phenix. The 2.7 Å map phased by SAD method allowed us to

build Rv0078 model unambiguously. The native Rv0078 structure was subsequently

determined by the program PHASER using SeMet substituted Rv0078 as the initial

search model. To obtain the Rv0078-DNA crystals, purified Rv0078 was co-crystallized

with a 23-mer DNA duplex (5'- TTTACAAGCAGACTGCCGGTAAC-3') at a molar ratio

of 2:1 (protein dimer:DNA) in the presence of 150 mM $MgCl_2$. The DNA-bound Rv0078

crystals were grown in the buffer containing only 0.2 M Magnesium formate. Diffraction

data to 3.0 Å were collected at the Life Sciences Collaborative Access Team (LS-CAT)

beamline of APS and were processed with Mosflm. The Rv0078-DNA structure was

determined by PHASER using DNA-free structure as the search model. All the

refinements were performed using Phenix-refine (Adams *et al.*, 2010). The statistics

were provided in **Supplemental Table 3**.

Mass spectrometry of fatty acids co-purified with Rv0078. Fatty acids were

extracted from 5 mg Rv0078 that was purified from *E. coli*, and analyzed as described

previously (Fan *et al.*, 2013). Separation and identification of the FA methyl esters were

performed on an HP5975 gas chromatograph-mass spectrometer (Hewlett-Packard)

fitted with a 60 m × 250 µm SP-2340 capillary column (Supelco) with helium as the

carrier gas

Isothermal calorimetry. ITC experiment was performed in a Microcal PEAQ-ITC at 25

°C. The stirring speed was 750 rpm and the interval between each titration was 150 sec.

The concentration of Rv0078 in the reaction cell was 25 µM and the concentration of the

titration DNA ligand was 500 µM. The recorded thermal data was analyzed using

Microcal PEAQ-ITC analysis software.

MIC$_{50}$ Determination. To determine the MIC$_{50}$ of each antibiotic, *M. tuberculosis* strains were grown to an OD$_{580}$ ~0.7 and diluted into fresh media to an OD$_{580}$ of 0.02. Diluted cultures were transferred to a 96-well microtiter plate containing triplicate, 10-fold serial dilutions of antibiotic. Cell wall-active antibiotics (vancomycin, meropenem) were supplemented at all concentrations with potassium clavalunate to inhibit the intrinsic β-lactamase activity of *M. tuberculosis*. After five days of incubation at 37ºC, growth of each strain was measured by OD$_{580}$. MIC$_{50}$ values were interpolated from a non-linear least squares fit of log$_2$-transformed OD$_{580}$ measurements. Data are representative of two independent experiments. Antibiotics were purchased from Sigma-Aldrich (clavulanate, ethionamide, meropenem, rifampicin, vancomycin) or Thermo-Fisher Scientific (ciprofloxacin, ethambutol, isoniazid, norfloxacin, streptomycin).

Mouse Infections. Mouse infections were performed essentially as described previously(Darwin *et al.*, 2003). 7-9-week-old female C57BL6/J mice (The Jackson Laboratory) were infected by aerosol to deliver ~200 bacilli per mouse, using a Glas-Col Inhalation Exposure System (Terre Haute, IN). Strains used were WT (MHD761) or (MHD794), Rv0077c (MHD1086), Rv0077c complemented (MHD1077), Rv0078 (MHD1315), Rv0078 complemented with WT Rv0078 (MHD1318) and Rv0078$_{W100R}$ (MHD1316). This study was performed in strict accordance with the recommendations in the Guide for the Care and Use of Laboratory Animals of the National Institutes for Health. Mice were humanely euthanized according to an approved Institutional Animal Care and Use Committee protocol at New York University School of Medicine. Lungs

and spleens were harvested and homogenized PBS/0.05% Tween-80 at indicated time

points to determine bacterial CFU.

Metabolomic analysis of *M. tuberculosis* cell lysates. Four independent cultures of

each analyzed strain were grown in 7H9 to an OD_{580} ~0.7 and treated with iP in DMSO

at a final concentration of 100 µM or an equal volume of DMSO for 24 hours. Bacteria

were harvested the next day at OD_{580} ~1. Sixty-five OD-equivalents per replicate were

processed by chloroform:methanol extraction(Layre *et al.*, 2011, Samanovic *et al.*,

2015). Metabolomic profiling was performed by Metabolon, Inc.

Staining and microscopy. *M. tuberculosis* strains were grown to mid-logarithmic

phase (OD_{580} ~0.5-0.7). 5 µl of culture was spotted onto glass slides and heat-killed

over a flame or on a heat block (15 min, 80˚C). Staining was performed according the

method of Ziehl-Neelson as per the manufacturer's instructions (BD Stain Kit ZN).

Images were acquired on a Zeiss Axio Observer with a Plan-Aprochromat 63x/1.4 oil

lens. Images were taken with an Axiocam503 camera at the NYULMC Microscopy

Laboratory.

Analysis of total lipids, mycolic acids and shorter chain fatty acids. For lipid

analysis 400 ml cultures were grown up to OD_{580} ~0.7 and treated with iP in DMSO at a

final concentration of 100 µM or an equal volume of DMSO-only for 24 hours. 400 ml of

Rv0077c and Rv0078 cultures were treated with DMSO only. Cells were washed three

times in DPBS and heated at 100°C for 45 min for sterilization before freezing at -20 °C.

Total lipids extraction from bacterial cells and preparation of fatty acid and mycolic acid

methyl esters from extractable lipids and delipidated cells followed earlier procedures

(Stadthagen *et al.*, 2005). Total lipids and fatty acid/mycolic acid methyl esters were

analyzed by one and two-dimensional thin-layer chromatography (TLC) in a variety of

solvent systems on aluminum-backed silica gel 60-precoated plates F_{254} (E. Merck).

TLC plates were revealed by spraying with cupric sulfate (10% in a 8% phosphoric acid

solution) and heating. Alternatively, total lipids were run in both positive and negative

mode and the released fatty acids/mycolic acids in negative mode only, on a high

resolution Agilent 6220 TOF mass spectrometer interfaced to a LC as described(Sartain

et al., 2011, Bhamidi *et al.*, 2012). Data files were analyzed with Agilent's Mass hunter

work station software and most compounds were identified using a database of *M.*

tuberculosis lipids developed in-house (Sartain *et al.*, 2011). Fatty acids methyl esters

from extractable lipids were treated with 3 M HCl in CH_3OH (Supelco) overnight at 80˚C,

dried and dissolved in *n*-hexane(s) prior to GC/MS analysis. GC/MS analyses of fatty

acid methyl esters were carried out using a TRACE 1310 gas chromatograph (Thermo

Fisher) equipped with a TSQ 8000 Evo Triple Quadrupole in the electron impact mode

and scanning from *m/z* 70 to *m/z* 1000 over 0.8 s. Helium was used as the carrier gas

with a flow rate of 1 ml per min. The samples were run on a ZB-5HT column (15 m x

0.25 mm i.d.) (Zebron). The injector (splitless mode) was set for 300°C (350°C for

mycolic acid methyl esters). The oven temperature was held at 60°C for 2 min,

programmed at 20°C per min to 375°C, followed by a 10 min hold. The data analyses

were carried out on Chromeleon data station.

References

Adams, P.D., P.V. Afonine, G. Bunkoczi, V.B. Chen, I.W. Davis, N. Echols, J.J. Headd, L.W. Hung, G.J. Kapral, R.W. Grosse-Kunstleve, A.J. McCoy, N.W. Moriarty, R. Oeffner, R.J. Read, D.C. Richardson, J.S. Richardson, T.C. Terwilliger & P.H. Zwart, (2010) PHENIX: a comprehensive Python-based system for macromolecular structure solution. *Acta Crystallogr D Biol Crystallogr* **66**: 213-221.

Anantharaman, V. & L. Aravind, (2001) The CHASE domain: a predicted ligand-binding module in plant cytokinin receptors and other eukaryotic and bacterial receptors. *Trends Biochem Sci* **26**: 579-582.

Bardarov, S., S. Bardarov Jr, Jr., M.S. Pavelka Jr, Jr., V. Sambandamurthy, M. Larsen, J. Tufariello, J. Chan, G. Hatfull & W.R. Jacobs Jr, Jr., (2002) Specialized transduction: an efficient method for generating marked and unmarked targeted gene disruptions in Mycobacterium tuberculosis, M. bovis BCG and M. smegmatis. *Microbiology* **148**: 3007-3017.

Barkan, D., Z. Liu, J.C. Sacchettini & M.S. Glickman, (2009) Mycolic acid cyclopropanation is essential for viability, drug resistance, and cell wall integrity of Mycobacterium tuberculosis. *Chem Biol* **16**: 499-509.

Bhamidi, S., L. Shi, D. Chatterjee, J.T. Belisle, D.C. Crick & M.R. McNeil, (2012) A bioanalytical method to determine the cell wall composition of Mycobacterium tuberculosis grown in vivo. *Anal Biochem* **421**: 240-249.

Bhatt, A., N. Fujiwara, K. Bhatt, S.S. Gurcha, L. Kremer, B. Chen, J. Chan, S.A. Porcelli, K. Kobayashi, G.S. Besra & W.R. Jacobs, Jr., (2007) Deletion of kasB in Mycobacterium tuberculosis causes loss of acid-fastness and subclinical latent tuberculosis in immunocompetent mice. *Proc Natl Acad Sci U S A* **104**: 5157-5162.

Blondiaux, N., M. Moune, M. Desroses, R. Frita, M. Flipo, V. Mathys, K. Soetaert, M. Kiass, V. Delorme, K. Djaout, V. Trebosc, C. Kemmer, R. Wintjens, A. Wohlkonig, R. Antoine, L. Huot, D. Hot, M. Coscolla, J. Feldmann, S. Gagneux, C. Locht, P. Brodin, M. Gitzinger, B. Deprez, N. Willand & A.R. Baulard, (2017) Reversion of antibiotic resistance in Mycobacterium tuberculosis by spiroisoxazoline SMARt-420. *Science* **355**: 1206-1211.

Brosch, R., S.V. Gordon, M. Marmiesse, P. Brodin, C. Buchrieser, K. Eiglmeier, T. Garnier, C. Gutierrez, G. Hewinson, K. Kremer, L.M. Parsons, A.S. Pym, S. Samper, D. van Soolingen & S.T. Cole, (2002) A new evolutionary scenario for the Mycobacterium tuberculosis complex. *Proc Natl Acad Sci U S A* **99**: 3684-3689.

Cerda-Maira, F.A., M.J. Pearce, M. Fuortes, W.R. Bishai, S.R. Hubbard & K.H. Darwin, (2010) Molecular analysis of the prokaryotic ubiquitin-like protein (Pup) conjugation pathway in Mycobacterium tuberculosis. *Mol Microbiol* **77**: 1123-1135.

Darwin, K.H., S. Ehrt, J.C. Gutierrez-Ramos, N. Weich & C.F. Nathan, (2003) The proteasome of Mycobacterium tuberculosis is required for resistance to nitric oxide. *Science* **302**: 1963-1966.

Deb, C., C.M. Lee, V.S. Dubey, J. Daniel, B. Abomoelak, T.D. Sirakova, S. Pawar, L. Rogers & P.E. Kolattukudy, (2009) A novel in vitro multiple-stress dormancy model for Mycobacterium tuberculosis generates a lipid-loaded, drug-tolerant, dormant pathogen. *PLoS One* **4**: e6077.

DeBarber, A.E., K. Mdluli, M. Bosman, L.G. Bekker & C.E. Barry, 3rd, (2000) Ethionamide activation and sensitivity in multidrug-resistant Mycobacterium tuberculosis. *Proc Natl Acad Sci U S A* **97**: 9677-9682.

Fan, J., C. Yan & C. Xu, (2013) Phospholipid:diacylglycerol acyltransferase-mediated triacylglycerol biosynthesis is crucial for protection against fatty acid-induced cell death in growing tissues of Arabidopsis. *Plant J* **76**: 930-942.

Festa, R.A., M.B. Jones, S. Butler-Wu, D. Sinsimer, R. Gerads, W.R. Bishai, S.N. Peterson & K.H. Darwin, (2011) A novel copper-responsive regulon in Mycobacterium tuberculosis. *Mol Microbiol* **79**: 133-148.

Festa, R.A., M.J. Pearce & K.H. Darwin, (2007) Characterization of the proteasome accessory factor (paf) operon in Mycobacterium tuberculosis. *J Bacteriol* **189**: 3044-3050.

Frebort, I., M. Kowalska, T. Hluska, J. Frebortova & P. Galuszka, (2011) Evolution of cytokinin biosynthesis and degradation. *J Exp Bot* **62**: 2431-2452.

Gandotra, S., M.B. Lebron & S. Ehrt, (2010) The Mycobacterium tuberculosis proteasome active site threonine is essential for persistence yet dispensable for replication and resistance to nitric oxide. *PLoS Pathog* **6**.

Gandotra, S., D. Schnappinger, M. Monteleone, W. Hillen & S. Ehrt, (2007) In vivo gene silencing identifies the Mycobacterium tuberculosis proteasome as essential for the bacteria to persist in mice. *Nat Med* **13**: 1515-1520.

Hatfull, G.F. & W.R.J. Jacobs, (2000) *Molecular Genetics of Mycobacteria*. ASM Press, Washington, DC.

Inoue, T., M. Higuchi, Y. Hashimoto, M. Seki, M. Kobayashi, T. Kato, S. Tabata, K. Shinozaki & T. Kakimoto, (2001) Identification of CRE1 as a cytokinin receptor from Arabidopsis. *Nature* **409**: 1060-1063.

Lamichhane, G., T.R. Raghunand, N.E. Morrison, S.C. Woolwine, S. Tyagi, K. Kandavelou & W.R. Bishai, (2006) Deletion of a Mycobacterium tuberculosis proteasomal ATPase homologue gene produces a slow-growing strain that persists in host tissues. *J Infect Dis* **194**: 1233-1240.

Layre, E., L. Sweet, S. Hong, C.A. Madigan, D. Desjardins, D.C. Young, T.Y. Cheng, J.W. Annand, K. Kim, I.C. Shamputa, M.J. McConnell, C.A. Debono, S.M. Behar, A.J. Minnaard, M. Murray, C.E. Barry, 3rd, I. Matsunaga & D.B. Moody, (2011) A comparative lipidomics platform for chemotaxonomic analysis of Mycobacterium tuberculosis. *Chem Biol* **18**: 1537-1549.

Lechat, P., L. Hummel, S. Rousseau & I. Moszer, (2008) GenoList: an integrated environment for comparative analysis of microbial genomes. *Nucleic Acids Res* **36**: D469-474.

Lin, G., D. Li, L.P. de Carvalho, H. Deng, H. Tao, G. Vogt, K. Wu, J. Schneider, T. Chidawanyika, J.D. Warren, H. Li & C. Nathan, (2009) Inhibitors selective for mycobacterial versus human proteasomes. *Nature* **461**: 621-626.

Mougel, C. & I.B. Zhulin, (2001) CHASE: an extracellular sensing domain common to
transmembrane receptors from prokaryotes, lower eukaryotes and plants. *Trends*
Biochem Sci **26**: 582-584.
Nardini, M. & B.W. Dijkstra, (1999) Alpha/beta hydrolase fold enzymes: the family keeps
growing. *Curr Opin Struct Biol* **9**: 732-737.
Orth, P., D. Schnappinger, W. Hillen, W. Saenger & W. Hinrichs, (2000) Structural basis
of gene regulation by the tetracycline inducible Tet repressor-operator system.
Nat Struct Biol **7**: 215-219.
Ramakrishnan, L., N.A. Federspiel & S. Falkow, (2000) Granuloma-specific expression
of Mycobacterium virulence proteins from the glycine-rich PE-PGRS family.
Science **288**: 1436-1439.
Saeed, A.I., V. Sharov, J. White, J. Li, W. Liang, N. Bhagabati, J. Braisted, M. Klapa, T.
Currier, M. Thiagarajan, A. Sturn, M. Snuffin, A. Rezantsev, D. Popov, A.
Ryltsov, E. Kostukovich, I. Borisovsky, Z. Liu, A. Vinsavich, V. Trush & J.
Quackenbush, (2003) TM4: a free, open-source system for microarray data
management and analysis. *Biotechniques* **34**: 374-378.
Sakakibara, H., (2006) Cytokinins: activity, biosynthesis, and translocation. *Annu Rev*
Plant Biol **57**: 431-449.
Samanovic, M.I., S. Tu, O. Novak, L.M. Iyer, F.E. McAllister, L. Aravind, S.P. Gygi, S.R.
Hubbard, M. Strnad & K.H. Darwin, (2015) Proteasomal control of cytokinin
synthesis protects Mycobacterium tuberculosis against nitric oxide. *Mol Cell* **57**:
984-994.
Sambrook, J., T. Maniatis & E. Fritsch, (1989) *Molecular Cloning: A Laboratory Manual.*
. Cold Spring Harbor: Cold Spring Harbor Laboratory Press.
Sartain, M.J., D.L. Dick, C.D. Rithner, D.C. Crick & J.T. Belisle, (2011) Lipidomic
analyses of Mycobacterium tuberculosis based on accurate mass measurements
and the novel "Mtb LipidDB". *J Lipid Res* **52**: 861-872.
Soding, J., A. Biegert & A.N. Lupas, (2005) The HHpred interactive server for protein
homology detection and structure prediction. *Nucleic Acids Res* **33**: W244-248.
Stadthagen, G., J. Kordulakova, R. Griffin, P. Constant, I. Bottova, N. Barilone, B.
Gicquel, M. Daffe & M. Jackson, (2005) p-Hydroxybenzoic acid synthesis in
Mycobacterium tuberculosis. *J Biol Chem* **280**: 40699-40706.
Steklov, M.Y., S.N. Lomin, D.I. Osolodkin & G.A. Romanov, (2013) Structural basis for
cytokinin receptor signaling: an evolutionary approach. *Plant Cell Rep* **32**: 781-
793.
Sugiyama, T. & T. Hashizume, (1978) Ribosylation of 6-Chloropurine and Its 2-
Methylthio Derivative by a Fusion Procedure Using
Iodin. *Agric. Biol. Chem.* **42**: 1791-1792.
Tian, J., R. Bryk, S. Shi, H. Erdjument-Bromage, P. Tempst & C. Nathan, (2005)
Mycobacterium tuberculosis appears to lack alpha-ketoglutarate dehydrogenase
and encodes pyruvate dehydrogenase in widely separated genes. *Mol Microbiol*
57: 859-868.
To, J.P. & J.J. Kieber, (2008) Cytokinin signaling: two-components and more. *Trends*
Plant Sci **13**: 85-92.

Vannelli, T.A., A. Dykman & P.R. Ortiz de Montellano, (2002) The antituberculosis drug
707   ethionamide is activated by a flavoprotein monooxygenase. *J Biol Chem* **277**:
708   12824-12829.
709   WHO, (2017) http://www.who.int/mediacentre/factsheets/fs104/en/.
710   Winn, M.D., C.C. Ballard, K.D. Cowtan, E.J. Dodson, P. Emsley, P.R. Evans, R.M.
711   Keegan, E.B. Krissinel, A.G. Leslie, A. McCoy, S.J. McNicholas, G.N.
712   Murshudov, N.S. Pannu, E.A. Potterton, H.R. Powell, R.J. Read, A. Vagin & K.S.
713   Wilson, (2011) Overview of the CCP4 suite and current developments. *Acta*
714   *Crystallogr D Biol Crystallogr* **67**: 235-242.
715   Wohlkonig, A., H. Remaut, M. Moune, A. Tanina, F. Meyer, M. Desroses, J. Steyaert, N.
716   Willand, A.R. Baulard & R. Wintjens, (2017) Structural analysis of the interaction
717   between spiroisoxazoline SMARt-420 and the Mycobacterium tuberculosis
718   repressor EthR2. *Biochem Biophys Res Commun* **487**: 403-408.
719

A

Gene	Putative function	Fold increase
Rv0076c	Hypothetical protein	3
Rv0077c	Oxidoreductase	81
Rv0078	TetR-like regulator	2
Rv1557	*mmpL6*; possible lipid transporter	8

B

Gene	*Mma*	*Mav*	*Mul*	*Mle*	*Mgi*	*Msm*
Rv0076c	76	68	0	0	0	58
Rv0077c	87	88	86	0	53	52
Rv0078	94	90	92	67	0	0
mmpL6*	88	88	76	63	80	83

Figure 1. Cytokinins induce the expression of Rv0077c in *M. tuberculosis*. (A) Genes significantly regulated by the presence of 100 µM of iP for five hours as analyzed by RNA-Seq. **(B)** Percent identity between *M. tuberculosis* H37Rv proteins and proteins of targeted mycobacterial genomes including *M. marinum* (*Mma*), *M. avium* (*Mav*), *M. ulcerans* (*Mul*), *M. gilvum* (*Mgi*), and *M. smegmatis* (*Msm*). Asterisk (*) indicates *mmpL6* encodes a truncated protein in H37Rv, unlike in the other mycobacterial species in the table. **(C)** Immunoblot for Rv0077c in total cell lysates of WT *M. tuberculosis*. "compl." = complemented. iP was used at a final concentration of 100 µM when added. **(D)** Dose-dependent production of Rv0077c protein. Bacteria were incubated with cytokinin at the indicated concentrations for 24 hours. **(E)** Only cytokinins, and not closely related molecules, induce the production of Rv0077c. Each compound was added to a final concentration of 100 µM. "R" indicates the riboside form of the preceding indicated cytokinin. **(F)** Adenine inhibits the induction of Rv0077c by 100 µM iP. For all panels, we added an equal volume of DMSO to samples where iP was not added. For all immunoblots (IB), we stripped the membranes and incubated them with antibodies to dihydrolipoamide acyltransferase (DlaT) to confirm equal loading of total lysates. Molecular weight standards are indicated to the left of the blots and are in kilodaltons (kD). Ade, adenine.

Figure 2. Rv0078 represses the expression of Rv0077c. (A) The putative transcriptional start sites (+1) of Rv0077c and Rv0078 as determined by 5'RACE and represented as bent arrows. The predicted start codons are in capital bold letters. **(B)** EMSA analysis identifies a putative repressor binding site. Probe sequences. +19/-14 refers to positions relative to the Rv0077c +1. In bold is the presumed binding site. Mutated residues are in red. Not shown at the end of each probe is a sequence for annealing to a fluorescent tag (Table supplement 1). Rv0078 was purified under native conditions from *E. coli*. **(C)** Deletion and disruption of Rv0078 results in the constitutive expression of Rv0077c. Total cell lysates were prepared and separated on a 10% SDS-PAGE gel. IB = immunoblot. The membrane was stripped and incubated them with antibodies to DlaT to confirm equal loading of samples. Molecular weight standards are indicated to the left of the blots and are in kilodaltons (kD).

Figure 3. Crystal structure of Rv0078 in complex with DNA. (A) ITC of Rv0078 binding to the +13/-8 DNA probe. The binding stoichiometry, ΔH, and K_D are marked. **(B)** Overall structure of Rv0078-DNA complex in cartoon view. Two Rv0078 dimers ("central" and "peripheral") bind to one DNA molecule. **(C)** The distance between two DNA-binding domains decreases by ~6 Å when bound to DNA. The DNA-free Rv0078 is in green; the DNA-bound Rv0078 is in magenta. **(D)** Interactions between the LBD (magenta) and DBD (green) in a monomer. **(E)** The ligand binding pocket (magenta mesh) of Rv0078 is enclosed by a four-helix bundle (helices α5 to α8).

Figure 4. Rv0078-DNA interactions. The hydroxyl group of **(A)** Thr37, **(B)** Thr47, and **(C)** Tyr52 interact with the backbone phosphate with a distance of 2.6 Å. **(D)** Arg48 interacts with guanidine with a distance of 2.6 Å. The 2Fo-Fc maps are contoured at 1σ level. **(E)** A schematic representation of Rv0078-DNA contacts. Residues of the central dimer are labeled in red, and residues of the peripheral dimer are in purple. **(F)** EMSA using WT DNA probe and Rv0078 with mutations in Thr37, Thr47, Arg48, or Tyr52. **(G)** Left panel: EMSA of Rv0078 with DNA probes with G-to-A substitutions in the central dimer binding region (S1), peripheral dimer binding region (S2), or in both DNA regions (S3). The white arrowhead marks the partial shift with the S2 probe at high protein concentration. Right panel: sequences of the four DNA probes used in EMSA experiments. The nucleotides contacting the central dimer are boxed in red and those contacting the peripheral dimer are boxed in purple.

Figure 5. Loss of Rv0077c or Rv0078 does not attenuate bacterial survival in mice.
(A) Bacterial colony forming units (CFU) after infection of C57BL/6J mice with WT, Rv0077c and complemented strains. **(B)** Bacterial CFU after infection of C57BL/6J mice with WT, Rv0078 and complemented strains. For both panels, data in each are from a single experiment that is representative of two independent experiments. Error bars indicate the standard error of the mean.

Figure 6. Induction of Rv0077c expression results in loss of acid-fast staining. *M. tuberculosis* strains were examined by Ziehl-Neelsen staining. Magnification is 63-fold. Scale bars are applicable to all images. See text for details.