1

**Determining the specificity of Cascade binding, interference, and primed adaptation *in vivo***

**in the *Escherichia coli* type I-E CRISPR-Cas system**

Lauren A. Cooper[1], Anne M. Stringer [2] and Joseph T. Wade[1,2*]

[1]Department of Biomedical Sciences, School of Public Health, University at Albany, Albany,

New York, USA.

[2]Wadsworth Center, New York State Department of Health, Albany, New York, USA

*Corresponding author:

joseph.wade@health.ny.gov

**ABSTRACT**

In CRISPR-Cas immunity systems, short CRISPR RNAs (crRNAs) are bound by CRISPR-

associated (Cas) proteins, and these complexes target invading nucleic acid molecules for

degradation in a process known as interference. In type I CRISPR-Cas systems, the Cas protein

complex that binds DNA is known as Cascade. Association of Cascade with target DNA can also

lead to acquisition of new immunity elements in a process known as primed adaptation. Here, we

assess the specificity determinants for Cascade-DNA interaction, interference, and primed

adaptation *in vivo*, for the type I-E system of *Escherichia coli*. Remarkably, as few as 5 bp of

crRNA-DNA are sufficient for association of Cascade with a DNA target. Consequently, a single

crRNA promotes Cascade association with numerous off-target sites, and the endogenous *E. coli*

crRNAs direct Cascade binding to >100 chromosomal sites. In contrast to the low specificity of

Cascade-DNA interactions, >18 bp are required for both interference and primed adaptation.

Hence, Cascade binding to sub-optimal, off-target sites is inert. Our data support a model in

which initial Cascade association with DNA targets requires only limited sequence

complementarity at the crRNA 5□ end, whereas recruitment and/or activation of the Cas3

nuclease, a prerequisite for interference and primed adaptation, requires extensive base-pairing.

**IMPORTANCE**

Many bacterial and archaeal species encode CRISPR-Cas immunity systems that protect against

invasion by foreign DNA. In the *Escherichia coli* CRISPR-Cas system, a protein complex,

Cascade, binds 61 nt CRISPR RNAs (crRNAs). The Cascade complex is directed to invading

DNA molecules through base-pairing between the crRNA and target DNA. This leads to

recruitment of the Cas3 nuclease that destroys the invading DNA molecule and promotes

acquisition of new immunity elements. We make the first *in vivo* measurements of Cascade

binding to DNA targets. Thus, we show that Cascade binding to DNA is highly promiscuous;

endogenous *E. coli* crRNAs can direct Cascade binding to >100 chromosomal locations. By

contrast, we show that target degradation and acquisition of new immunity elements requires

highly specific association of Cascade with DNA, limiting CRISPR-Cas function to the

appropriate targets.

**INTRODUCTION**

Clustered Regularly Interspaced Short Palindromic Repeats (CRISPR)-Cas (CRISPR-associated) systems are adaptive immune systems found in approximately 40% of bacteria and 90% of archaea [1]. CRISPR-Cas systems are characterized by the presence of CRISPR arrays and Cas proteins. CRISPR arrays are genomic loci that consist of short repetitive sequences ("repeats"), interspaced with short sequences of viral or plasmid origin ("spacers") [2–5]. Spacers are acquired during a process known as "adaptation", in which a complex of Cas1 and Cas2 integrates invading DNA into a CRISPR array, effectively immunizing the organism from future assault by the invader [6]. In the archetypal type I-E CRISPR system of *Escherichia coli*, immunity occurs via two processes known as "biogenesis" and "interference". During biogenesis, a CRISPR array is transcribed, and Cas6e processes the transcript into individual 61 nt CRISPR RNAs (crRNAs) that each include a single 32 nt spacer sequence flanked by partial repeat sequences [7,8]. Individual crRNAs are then incorporated into Cascade, a protein complex composed of five different Cas proteins (Cse1 (Cas8e)-$Cse2_2$-$Cas7_6$-Cas5-Cas6e) [7,9]. During interference, Cascade complexes bind to target DNA sequences known as "protospacers" that are complementary to the crRNA spacer, and are immediately adjacent to a short DNA sequence known as a "Protospacer-Associated Motif" (PAM) [10] that is bound by Cse1 [11,12]. The crRNA bound by Cascade forms an R-loop with the target DNA, which in turn leads to recruitment of the Cas3 nuclease, DNA cleavage, and elimination of the invader [13–19].

For type I CRISPR-Cas systems, adaptation can occur by two mechanisms: "naïve" and "primed". Typically, naïve adaptation requires only Cas1 and Cas2 [6,20]. Primed adaptation, by

contrast, requires all Cas proteins and an existing crRNA [21]. The molecular details of primed

adaptation are poorly understood. Spacers acquired by primed adaptation correspond to locations

on the same DNA molecule as the protospacer [21–24]. Some type I CRISPR-Cas systems

acquire spacers preferentially from one strand [21,23,25], whereas others acquire spacers from

both strands [24,26,27]. Primed adaptation has been proposed to involve translocation of Cas3

away from the Cascade-bound protospacer [21,28].

There are conflicting reports on the relationship between interference and primed adaptation.

Initially, it was proposed that primed adaptation occurs only when Cascade-protospacer

interactions are sub-optimal and cannot lead to interference, e.g. with a sub-optimal PAM, or

with mismatches in the PAM-proximal region of the protospacer known as the "seed" [21,23,29–

31]. However, more recent studies have shown that at least some protospacers can lead to both

interference and primed adaptation, indicating that the requirements for interference and primed

adaptation overlap [24,32,33].

Prior to interference or primed adaptation, Cascade must bind to the target protospacer. This

requires an interaction between Cse1 and the PAM, and base-pairing between the crRNA and

protospacer DNA [11,12,14]. PAM recognition is required for both Cascade binding and later

recruitment and activation of Cas3 [11,13,16,19]. Changes to the optimal PAM weaken Cascade

binding to a protospacer [34]. Nonetheless, some sub-optimal PAMs are sufficient for

interference, albeit with lower efficiency than the optimal PAM [31]. Sequences within the

crRNA spacer are also required for initial binding of Cascade to a protospacer; mutations in

positions 1-5 and 7-8 adjacent to the PAM of the protospacer (the seed sequence) reduce the

affinity of Cascade for the protospacer [29].

Reports of the sequence determinants for Cascade binding, interference and primed adaptation

are conflicting [16,18,23,31,35]. In particular, the impact of extensive mismatches in the

crRNA:DNA hybrid on Cascade binding and primed adaptation is unclear [16,35]. Importantly,

association of Cascade with protospacer DNA has not previously been studied in an *in vivo*

context. Here, we use ChIP-seq to perform the first *in vivo* assessment of Cascade binding to its

DNA targets. Our data show that base-pairing between the crRNA and protospacer with as few

as 5 nt in the seed region, coupled with an optimal PAM, is often sufficient for Cascade binding.

Hence, crRNAs, including those transcribed from the native *E. coli* CRISPR loci, drive off-target

binding at over a hundred chromosomal sites. If Cascade binding to DNA were sufficient for

interference or primed adaptation, these off-target binding events would likely be catastrophic

for the bacterium [36,37]. However, we show that extensive base-pairing between the crRNA

and protospacer from the PAM-proximal end is required for efficient interference and primed

adaptation. Thus, under native conditions, Cascade samples potential DNA target sites, but limits

nuclease activity to protospacers that meet a higher specificity threshold that would only be

expected of on-target sites.

**RESULTS**

**An AAG PAM and seed matches are sufficient for Cascade binding to DNA target sites *in***

***vivo***

Previous studies of Cascade association with protospacer DNA have been performed *in vitro*

using purified Cascade and crRNA. To determine the *in vivo* target specificity of *E. coli* Cascade,

we used ChIP-seq to map the association of Cse1-FLAG$_3$ and FLAG$_3$-Cas5 (FLAG-tagged

strains retain CRISPR function; Supplementary Figure 1) across the *E. coli* chromosome in

Δ*cas3* (interference-deficient) cells constitutively expressing all other *cas* genes and each of two

crRNAs that target either the *lacZ* promoter or the *araB* promoter (both targets are chromosomal;

Figure S2A-B). ChIP-seq data for Cse1 and Cas5 were highly correlated ($R^2$ values of 0.93-0.99

for *lacZ*-targeting cells, and 0.99 for *araB*-targeting cells), consistent with Cse1 and Cas5

binding DNA together in the context of Cascade. We detected association of Cascade with many

genomic loci for each of the two spacers tested (Figure 1A-B; Table S1). In all cases, the

genomic region with strongest Cascade association was the on-target site at *lacZ* or *araB*. Off-

target binding events occurred with <20% of the ChIP signal of on-target binding. To determine

the sequence requirements for off-target Cascade binding with each of the two crRNAs used, we

searched for enriched sequence motifs in the Cascade-bound regions, excluding the on-target site

(Table S2). For both the *lacZ* and *araB* spacers, the most enriched sequence motif we identified

was a close match to the canonical PAM, AAG, on the non-target strand, followed by 5 nt of

sequence complementarity at the start of the crRNA seed region (Figure 1C-D; c.f. Figure S2A-

B). In some cases, we observed Cascade binding events associated with non-AAG PAMs;

however, these sites were more weakly bound, and/or had matches in seed region beyond

position 5. For example, Cascade binding events targeted by the *araB* spacer were significantly

more likely to have matches at positions 7-9 in cases where the PAM was not AAG (an average

of 2.2 out of 3 matches per target sequence if the PAM was not AAG [n = 19], and 1.2 out of 3

matches if the PAM was AAG [n=41]; Fisher's Exact Test $p = 0.00005$). We conclude that as

few as 5 bp in the seed region, together with an AAG PAM, are sufficient for Cascade binding,

with additional base-pairing in/near the seed region increasing binding and/or overcoming the

need for an AAG PAM.

**Extensive off-target Cascade binding driven by endogenous spacers**

We identified several sites of Cascade binding that were shared between cells targeting *lacZ* and

cells targeting *araB*. These bound regions were not associated with sequences matching the seed

regions of either crRNA. We reasoned that these off-target binding events may be due to

Cascade association with the endogenous *E. coli* crRNAs. To test this hypothesis, we performed

ChIP-seq of Cse1-FLAG$_3$, as described above, for cells expressing only the endogenous CRISPR

RNAs from their native loci. Thus, we identified 188 binding sites for Cascade (Figure 2A;

Table S1). These sites were associated with four enriched sequence motifs, with each motif

corresponding to a canonical AAG PAM and 5-10 nt matching the seed region of a crRNA from

the CRISPR-I array (spacers #1, #3, #4, and #8; Figure 2B; Figure S2C; Table S2). The strongest

binding events were associated with spacer #8 of CRISPR-I ("sp1.8"; Figure 2B; Figure S2C).

To confirm that Cascade binding events were due to association with endogenous crRNAs, we

repeated the ChIP-seq experiment in cells lacking the CRISPR-I array and in cells lacking the

CRISPR-II array. Deletion of CRISPR-II had little effect on the profile of Cascade binding

(Figure 2C; Table S1). In contrast, deletion of CRISPR-I resulted in loss of Cascade binding to

almost all sites bound in wild-type cells (Figure 2D; Table S1). Instead, low-level binding of

Cascade was observed at a small number of sites that were associated with a weakly enriched

sequence motif corresponding to a perfect PAM and 8 nt matching the seed region of spacer #2

of CRISPR-II (Figure S2D and Figure S3; Table S2).

**CRISPR-I spacer #8 is the major determinant of off-target Cascade binding in cells**

**expressing endogenous crRNAs.**

Our data suggested that the majority of Cascade binding associated with endogenous crRNAs is

due to CRISPR-I, and that the dominant spacer from CRISPR-I is sp1.8. To confirm this, we

measured Cascade binding by ChIP-seq in cells lacking CRISPR-I but expressing a plasmid-

encoded sp1.8 crRNA. Note that the plasmid-expressed sp1.8 crRNA differs from sp1.8 at the

last two nucleotides of the spacer. However, these mismatches are not expected to affect Cascade

binding [23,38]. Most of the Cascade binding sites we observed were identical to those seen in

cells expressing both CRISPR arrays, or cells expressing only CRISPR-I (Figure 3A; Table S1),

and corresponded to regions containing strong matches to sp1.8 (orange dots in Figure 3A

correspond to regions containing a match to the sp1.8 motif shown in Figure 2B). As expected,

and unlike for cells expressing CRISPR-I, we detected only a single strongly enriched sequence

motif (Figure S4A; Table S2). This motif, as expected, corresponds to an AAG PAM and 9 nt

matching the seed region of sp1.8 (Figure S2C). We also detected a weakly enriched sequence

motif (Figure S4B; Table S2) that corresponds to an AAG PAM and the 11 nt immediately

downstream of the second repeat on the plasmid encoding the sp1.8 crRNA. This is likely due to

formation of a non-canonical crRNA that consists of the sequence between the second repeat and

the transcription terminator (Figure S2E). A transcription terminator hairpin has previously been

shown to function analogously to repeat sequence in the *E. coli* crRNAs [39].

The most enriched Cascade target region in cells with CRISPR-I, and cells expressing sp1.8

crRNA, was inside the *yggX* gene. We identified a sequence in this region with an AAG PAM

and matches to positions 1-5 and 7-10 of sp1.8 (Figure 3B). We used targeted ChIP-qPCR to

measure Cascade binding to this site in cells lacking CRISPR-I but expressing a plasmid-

encoded sp1.8 crRNA (mismatches to sp1.8 at the last two nucleotide positions, as described

above). We compared binding of Cascade to *yggX* in wild-type cells, and cells where the putative

protospacer was mutated in the region predicted to bind the sp1.8 crRNA seed. As expected, we

observed greatly reduced Cascade binding at the mutated site relative to the wild-type site.

Similarly, we observed greatly reduced Cascade binding at the wild-type site when we expressed

a mutant sp1.8 with changes in the seed region (Figure 3C). However, when we combined the

mutant spacer with the mutant protospacer, base-pairing potential was restored, and we observed

wild-type levels of Cascade binding (Figure 3C). We conclude that sp1.8 is the major

determinant for off-target Cascade binding in cells expressing endogenous crRNAs.

**Off-target Cascade binding events do not affect local gene expression**

Cascade binding events can lead to transcription repression by preventing initiating RNA

polymerase binding to a promoter, or acting as a roadblock to elongating RNA polymerase

within a transcription unit [38,40]. To determine if off-target events driven by endogenous

spacers affect local gene expression, we measured global RNA levels using RNA-seq in Δ*cas3*

cells with other *cas* genes constitutively expressed, with either intact CRISPR arrays or a

ΔCRISPR-I deletion. We detected few differences in RNA levels between the two strains (Table

S3), and none of the differences correspond to genes within 1 kb of a Cascade binding site

identified by ChIP-seq. We conclude that off-target binding by a Cas3-deficient complex does

not impact local gene expression.

**No evidence for RNA-targeting by *E. coli* Cascade**

A recent report suggested that Cascade binding to RNAs in *Pseudomonas aeruginosa,* which has

a type I-F system, leads to Cas3-mediated degradation of the target RNA [41]. Moreover, this

study suggested that only 8 nt of sequence complementarity between the crRNA and target RNA,

and a flanking 5□-GGA-3□ sequence, are required to recruit Cas3. This is similar to the

sequence requirement for off-target binding to DNA sites (Figures 1 and 2), suggesting that

Cascade could target many endogenous RNAs [42]. To determine whether the *E. coli* type I-E

CRISPR-Cas system targets RNA in a similar way, we measured global RNA levels using RNA-

seq in cells expressing *cas3* from a plasmid and all other *cas* genes from their chromosomal loci,

with either intact CRISPR arrays or a ΔCRISPR-I deletion. We compared these data to the data

described above for Δ*cas3* cells with either intact CRISPR arrays or a ΔCRISPR-I deletion. We

reasoned that targeted RNAs would be less abundant in cells expressing both Cas3 and CRISPR-

I. However, we detected only two genes, *ykgE* and *glpD*, for which RNA levels were

significantly lower in the *cas3*[+] CRISPR-I[+] strain than strains lacking either or both of *cas3* and

CRISPR-I (Table S3). Only one of these genes (*glpD*) contains an 8 nt sequence complementary

to the 3□ end of a spacer in CRISPR-I (spacer #4; we included the predicted 5' UTR in the

search for both RNAs). Given the length of the two genes, finding an 8 nt match by chance is not

unlikely. Moreover, three other genes contain the same 8 nt match to spacer #4, with the same 3

nt flanking sequence, but these genes did not have the RNA profile expected for a Cascade

target. Thus, our data strongly suggest that the type I-E CRISPR-Cas system in *E. coli* does not

target RNA using a mechanism similar to that described for the type I-F system in *P. aeruginosa.*

**Off-target Cascade binding is not associated with interference**

Previous studies have suggested that extensive mismatches at the PAM-proximal end of the

spacer/protospacer prevent interference [16,35]. To determine whether off-target Cascade

binding events lead to interference, we constructed a Δ*yggX* Δ*cas3* strain expressing all other *cas*

genes, with both CRISPR arrays intact. We introduced a plasmid expressing *cas3*, or an

equivalent empty vector. We then transformed these strains with a plasmid containing the off-

target protospacer from *yggX* that is an imperfect match to sp1.8, an equivalent plasmid with a

protospacer that is a perfect match to sp1.8, a plasmid with a protospacer that is a perfect match

to CRISPR-I spacer 2 ("sp1.2"), or an empty vector. We reasoned that the number of viable

transformants for plasmids with interference-proficient protospacers would be low for cells

expressing Cas3, since interference would cause loss of the protospacer-containing plasmid,

leading to killing by the antibiotic selection. In contrast, the number of viable transformants for

plasmids with interference-deficient protospacers, or cells not expressing Cas3, should be high.

We measured the transformation efficiency for plasmids containing each of the two protospacers

in cells with a Cas3-expressing plasmid, or an equivalent empty vector. The efficiency of

interference was calculated using the ratio of transformation efficiency for cells with/without

Cas3. As expected, the protospacer that perfectly matches sp1.8 resulted in highly efficient

interference. Similarly, the protospacer that perfectly matches sp1.2 resulted in highly efficient

interference. We conclude that sp1.2 is efficiently assembled into Cascade, despite the lack of

chromosomal off-target binding events detected by ChIP-seq. By contrast, the protospacer with

the native *yggX* sequence (i.e. imperfect match to sp1.8) resulted in no detectable interference

(Figure 4A). We conclude that off-target Cascade binding events do not cause interference.

**Off-target Cascade binding is not associated with primed adaptation**

Protospacers with multiple mismatches to a crRNA can still cause primed adaptation [23], and a

recent study concluded Cascade can bind to a protospacer with extensive mismatches, including

in the seed region or at the PAM-distal end, and that these binding events cause primed

adaptation [35]. To test whether off-target Cascade binding is sufficient for primed adaptation,

we used the strains described above that contained a plasmid with a protospacer that is either an

imperfect or a perfect match to sp1.8. We then introduced a plasmid with an inducible copy of

*cas3*, under non-inducing conditions, to avoid interference. Following induction of *cas3*

expression, we harvested cells and PCR-amplified the 5□ end of the CRISPR-II array to

determine whether new spacers had been acquired because of primed adaptation. We observed

robust primed adaptation for the protospacer with a perfect match to sp1.8, but no detectable

adaptation for the off-target protospacer with an imperfect match to sp1.8 (Figure 4B). We

conclude that off-target Cascade binding events do not lead to primed adaptation.

**Strong Cascade binding to protospacers with extensive mismatches at the crRNA PAM-**

**distal end**

To further delineate the protospacer sequence requirements for Cascade binding, interference and

primed adaptation, we constructed 13 variants of a protospacer that matches sp1.8. We selected

sp1.8 because it elicits robust Cascade binding, interference, and primed adaptation (Figures 3

and 4). Protospacer variants (Figure 5A) included those with (variant i; the "optimal"

protospacer) full sequence complementarity, and an optimal, AAG PAM; (variants ii - iii) non-

optimal PAMs: CCG, which is expected to completely abolish Cascade binding [34], and ATT, a

sub-optimal sequence previously shown to cause primed adaptation but not detectable

interference [31]; (variants iv - viii) two or three mismatches in the first three positions of the

seed; and (variants ix – xiii) stretches of $\geq 6$ nt mismatches at various positions within the

protospacer.

We pooled cells containing each of the protospacer variants. We used ChIP of Cse1-FLAG$_3$ in

$\Delta cas3$ cells to measure association of Cascade with all protospacers within the pool (see

Methods). As expected, the protospacer with a CCG PAM (variant ii) had far less Cascade

association than did the optimal protospacer (variant i) (Figure 5A). We presume that the level of

ChIP signal for the protospacer with the CCG PAM (variant ii) represents the background of this

experiment. The protospacer with a sub-optimal, ATT PAM (iii), showed reduced Cascade

binding relative to the optimal protospacer (variant i), but was well above the experimental

background (Figure 5A). Similarly, mismatches in the seed region (variants iv - viii) resulted in a

reduction in Cascade association (Figure 5A). Our data for PAM and seed mutants are consistent

with earlier studies showing that these sequences are important for Cascade binding

[17,29,30,34].

Mismatches in the protospacer from positions 1-6 (variants xi and xii) or 7-20 (variant xiii)

abolished Cascade binding (Figure 5A). This is consistent with the observation from our ChIP-

seq data that sequence matches in positions 1-8 appear to be required for Cascade binding to off-

target sites using sp1.8 (Figure 2B; Figure S4A). Strikingly, mismatches across positions 25-32

(variant ix) or positions 19-32 (variant x) did not reduce Cascade association relative to the

optimal protospacer (variant i) (Figure 5A). Thus, our data confirm that PAM-proximal sequence

is necessary for Cascade binding, while PAM-distal sequence is insufficient for Cascade binding.

**Extensive crRNA-protospacer base-pairing is required for interference and primed**

**adaptation**

We next determined which of the protospacer variants lead to interference. Using a modification

of a previously described assay (see Methods) [23,31], we measured the level of interference

with a plasmid target for each of the 13 protospacers, using $\Delta cas1$ cells that cannot acquire new

spacers; primed adaptation cannot contribute to the level of interference in these cells. As

expected, the optimal protospacer (variant i) was associated with robust levels of interference,

whereas protospacer variants that do not bind Cascade (variants ii, xi, xii, and xiii; Figure 5A)

were not associated with detectable interference (Figure 5B). Protospacers with PAM and seed

variants that showed reduced but not abolished Cascade binding (variants iii, vi, vii, and viii;

Figure 5A) were associated with a range of interference levels that correlate well with the level

of Cascade binding. Seed mutants with stronger defects in binding exhibited on detectable

interference. However, the ability of protospacers to cause interference did not always correlate

with the level of Cascade association. Specifically, we detected no interference for either of the

protospacer variants with mismatches only at the PAM distal end (variants ix and x; Figure 5B),

even though these protospacers bind Cascade at least as well as the optimal protospacer (Figure

5A).

Previous studies have proposed that some protospacers with sub-optimal PAMs or mismatches in

the seed region are not subject to detectable interference, but do cause primed adaptation

[21,23,28,31,35]. We determined whether the 13 protospacer variants caused primed adaptation

in a plasmid context. We used a highly sensitive assay for adaptation that relies on expression of

a *yfp* reporter gene that is encoded immediately upstream of a CRISPR array. Translation is

terminated upstream of *yfp* in cells without newly acquired spacers, whereas acquisition of one

spacer/repeat puts *yfp* back in frame [43], causing cells to fluoresce. We introduced an inducible

copy of *cas3* into cells with an intact CRISPR-I array, and containing each of the protospacers on

a high-copy plasmid, and the *yfp* reporter construct. We then induced expression of *cas3* and

measured the level of primed adaptation using flow cytometry. In this experiment we expect the

sp1.8 crRNA from the native CRISPR-I array to cause primed adaptation. We detected primed

adaptation for all protospacers associated with detectable interference (variants i, iii, vii, and viii;

Figure 5C), although the level of adaptation was lower for two of the constructs with the highest

levels of interference (variants i and viii). This is likely to be due to high levels of interference

reducing the available substrate for adaptation [33]. By contrast, we observed no adaptation for

protospacers that do not bind Cascade (variants ii, xi, xii, and xiii; Figure 5C). Strikingly, we

observed primed adaptation for four protospacers that were not associated with detectable

interference (Figure 5C). Three of these protospacers have seed mismatches, and exhibited the

lowest levels of Cascade binding (Figure 5A; variants iv, v and vi). The other protospacer has

mismatches across positions 25-32 (variant ix). Thus, for these protospacers, we detected robust

Cascade binding and primed adaptation but we were unable to detect interference. For the

protospacer with mismatches across positions 19-32 (variant x), we detected no primed

adaptation. Thus, for this protospacer, we detected robust Cascade binding, but no primed

adaptation or interference. Overall, our data suggest that extensive crRNA-protospacer base

pairing from the PAM-proximal end is required for both interference and primed adaptation, and

that primed adaptation is a more sensitive assay of CRISPR-Cas function than interference.

**DISCUSSION**

**Base-pairing in the seed region together with an AAG PAM is sufficient for Cascade to**

**bind DNA**

No previous studies have measured Cascade binding to protospacer DNA *in vivo*. Our ChIP data

indicate that an AAG PAM and as little as 5 nucleotides of base-pairing at the start of the seed

region are sufficient for *E. coli* Cascade to bind DNA targets. The sequence requirements for

protospacer binding in type II systems are similarly relaxed [44–46]. The affinity of Cascade for

a protospacer increases as the extent of base-pairing increases, but maximal affinity occurs with

no more than an 18 bp match at the PAM-proximal end (Figure 5A). Analysis of Cascade

interactions with DNA *in vitro* suggests that Cascade associates for brief periods with PAM-

containing sequences, and for longer periods if there is partial base-pairing in the seed region

[28]. Our data support these observations, although we did not detect ChIP signal at PAM

sequences that lack seed matches, suggesting that seed base-pairing contributes more to Cascade

association *in vivo*. Consistent with this suggestion, the difference in ChIP signal for off-target

and on-target sites is considerably less than the difference between dwell times *in vitro* [28],

although the use of crosslinking in ChIP may also contribute to this difference since crosslinking

"locks" Cascade on the DNA.

**AAG is the optimal PAM in** *E. coli*

Three previous studies proposed that AAG, GAG, TAG, AGG, and ATG are optimal PAMs in *E.*

*coli* [23,31,47], while another study suggested that AAG, ATG and GAG PAMs were associated

with moderately higher affinity Cascade binding than an AGG PAM [34]. Our data clearly

indicate that AAG is the optimal PAM for off-target sites, with most off-target Cascade binding

events being associated with an AAG PAM. Specifically, 65% of Cascade binding sites

associated with a detectable motif have an AAG PAM for the crRNAs targeting *lacZ* and *araB*,

and the plasmid-encoded sp1.8 crRNA. Moreover, off-target Cascade binding events with higher

enrichment scores, suggestive of higher Cascade affinity, were more likely to be associated with

an AAG PAM than Cascade binding events with lower enrichment scores (76% vs 61% for the

top 20% and bottom 80% of bound regions, respectively, after sorting by Cse1 enrichment level).

We hypothesize that the dependence on the PAM for Cascade binding is increased in situations

where base-pairing only occurs in the seed region. According to this model, complete or near-

complete base-pairing between the crRNA and protospacer would weaken the requirement for an

optimal PAM, obscuring differences in PAM affinity. This would explain why previous studies

suggested that there are at least three optimal PAMs [23,31,34,47].

**Defining the crRNA seed**

The seed region of a crRNA has been previously defined as positions 1-5 and 7-8, with position

1 being immediately adjacent to the PAM [29]. However, our data suggest that the length of the

seed varies between crRNAs, since we observed off-target binding with some crRNAs that

requires base-pairing in positions 1-5, whereas off-target binding for other crRNAs requires

base-pairing up to position 9 (Figures 1 and 2; Figure S3 and S4). We propose that the crRNA

sequence determines the length of the seed, and that this reflects the initial binding mode, prior to

extended base-pair formation. Every $6^{th}$ position of the crRNA is flipped out in the Cascade-

protospacer complex, and hence does not contribute to base-pairing [15,48,49]. Consistent with

this, the importance of position 6 for off-target binding is substantially less than that of positions

1-5 (Figures 1 and 2; Figure S3 and S4). Nonetheless, off-target protospacers had a sequence

match to the crRNA at position 6 far more frequently than expected by chance (45% for the

crRNAs targeting *lacZ* and *araB*, and the plasmid-encoded sp1.8 crRNA; Binomial Test *p*-value

$= 2.4e^{-10}$). We hypothesize that the initial binding of Cascade to a protospacer includes base-

pairing interactions at position 6, but that the complex rapidly transitions to a conformation in

which the 6[th] position is flipped out of the helix. Our data are consistent with an *in vitro* study of

another type I-E system, where position 6 was also shown to contribute to off-target Cascade

binding [50]. The apparent requirement for a sequence match at position 6 is not consistent

across all crRNAs we tested, suggesting that the pathway towards stable seed base-pairing differs

in a sequence-dependent manner.

**Interference and primed adaptation require extended R-loop formation**

Although binding of Cascade to a DNA target requires relatively little sequence identity, our data

indicate that robust interference and primed adaptation require at least 18-25 bp, beginning in the

seed region. This is consistent with *in vitro* data showing that near-complete R-loop formation is

required to license Cas3 activity [16]. Thus, although Cascade binds DNA promiscuously,

functional binding occurs with high specificity. Our data support a previously proposed model in

which extended R-loop formation triggers a conformational change in Cascade at the PAM-distal

end of the spacer, which is then transmitted, presumably through Cse2, to PAM-associated Cse1

[16,51]. This change in Cse1 conformation then recruits Cas3, and/or activates the nuclease

activity of Cas3, as suggested by a recent structural study [51].

**Evidence that interference and primed adaptation are obligately coupled processes**

Primed adaptation was initially proposed to be an alternative pathway to interference, with

optimal PAM/seed sequences leading to interference, and sub-optimal sequences leading to

primed adaptation [21,23,28,31,35,52]. However, primed adaptation has been observed in

situations where interference occurs (Figure 5, variants i, iii, vii, and viii) [22,24,32,33],

suggesting that primed adaptation and interference can be coupled processes, and supporting the

idea that primed adaptation is a positive feedback loop [22]. While these data show that primed

adaptation and interference can occur at the same time at a population level, they do not

necessarily indicate that individual primed adaptation and interference events are coupled.

Moreover, while it has been proposed that interference and primed adaptation are obligately

coupled [53], this has not been tested. There are many examples where primed adaptation has

been observed in the absence of detectable interference [21,23,31,32,35,52]. However, this can

be explained by the fact that primed adaptation is likely to be a more sensitive assay of CRISPR-

Cas function than interference, as there would be detectable primed adaptation but not detectable

interference in cells where target DNA replication outpaces interference [53]. Our data are

consistent with a model in which primed adaptation and interference are coupled processes: seed

mismatches reduce Cascade binding, and we observe a corresponding effect on interference and

primed adaptation, with primed adaptation being a more sensitive assay for CRISPR-Cas

function (Figure 5). The only exception to this trend is the seed mismatch with the highest level

of binding (Figure 5, variant viii), which has relatively low levels of primed adaptation.

However, very efficient interference with this variant likely depletes substrate for primed

adaptation [33]. Unexpectedly, we observed primed adaptation in the absence of detectable

interference for a protospacer with mismatches across positions 25-32 (Figure 5, variant ix). We

propose that this degree of mismatch at the 3☐ end of the crRNA greatly reduces, but does not

abolish, the isomerization of Cascade into the "active" state that recruits/activates Cas3.

**Extensive, inert, off-target binding of Cascade**

Cascade has many off-target binding sites due to its ability to bind DNA with low sequence-

specificity. Consequently, the endogenous crRNAs transcribed from the bacterial genome result

in extensive off-target binding, even in the absence of an on-target site. Since off-target binding

does not involve extended R-loop formation, it has no deleterious effects on genome integrity.

We also observed no impact on transcription associated with any of the off-target binding events,

despite that fact that targeted Cascade binding is known to repress transcription by occluding

promoters or acting as a roadblock for elongating RNA polymerase [38,40]. Transcription

repression by Cascade is considerably weaker when targeting within a transcribed region (i.e.

acting as a roadblock) [38]. Given that the location of off-target Cascade binding sites is

essentially random with respect to genome organization, and that genes make up ~90% of the *E.*

*coli* genome, off-target Cascade binding is expected to be primarily intragenic. This may partly

explain the lack of impact on transcription. Moreover, a recent study showed that the level of

repression by Cascade occlusion of a promoter is greatly reduced with as few as 6 bases

mismatched at the PAM-distal end of the spacer/protospacer [54], suggesting that even

intergenic off-target Cascade binding sites would be transcriptionally inert. We propose that

incomplete R-loop formation results in an unstable Cascade-DNA complex with a relatively high

rate of dissociation, such that it cannot compete effectively with initiating or elongating RNA

polymerase. Consistent with this model, stable association of Cascade with DNA *in vitro* has

been shown to require near-complete R-loop formation [18]. We conclude that type I CRISPR-

Cas systems have evolved to tolerate off-target binding driven by the endogenous crRNAs, and

are only functional at on-target sites. Given the length of crRNA spacers in type I systems, there

is no expectation of complete or near-complete spacer-protospacer base-pairing by chance. It is

important to note that self-targeting by type I CRISPR-Cas systems has been described

previously, but these would be considered "on-target" events, likely caused by acquisition of

spacers from the chromosome. As expected for spacers with perfect sequence complementarity,

these self-targeting crRNAs are typically functional in gene regulation and interference

[36,37,55].

**Not all crRNAs are created equal**

The *E. coli* genome encodes at least 19 crRNAs, yet our data suggest that only four crRNAs

contribute to off-target binding of Cascade. All four of these crRNAs are encoded in the

CRISPR-I array, and the majority of off-target binding is driven by just one, sp1.8. The lack of

off-target binding driven by CRISPR-II crRNAs is likely due to weak transcription of this array,

which is repressed by H-NS [56]. In contrast, the CRISPR-I array is likely co-transcribed with

the upstream *cas* genes, which are strongly transcribed in the strain used in this study. The

preference for specific spacers within CRISPR-I cannot be explained by differences in

expression levels, since the crRNAs are transcribed as a single RNA. Rather, biases in spacer

usage are more likely due to differential assembly of specific crRNAs into Cascade. Consistent

with this, a previous study surveyed crRNAs associated with Cascade. Spacers #2, #4 and #8

from CRISPR-I represented 68% of the Cascade-associated crRNAs [7]. The cause of this bias is

unclear, but may in part be due to differences in RNA secondary structure between spacers,

which could impact the efficiency of RNA processing by Cas6e. Consistent with this, RNA

secondary structure of repeat sequences, and associated processing by Cas6, has been shown to

be impacted by spacer sequences in the type I-D system of *Synechocystis* sp. PCC 6803 [57].

Nonetheless, it is likely that other factors influence the level of off-target binding, since the

relative association of crRNAs for spacers #2, #4 and #8 with Cascade is likely to be similar [7],

and sp1.2 causes efficient interference (Figure 4A), but sp1.8 drives a disproportionately high

level of off-target binding relative to sp1.2. Strikingly, there are many more chromosomal

sequence matches to the seed sequence of sp1.8 coupled with an AAG PAM than for any other

spacer (Table S4). This is likely due to the fact that the sequence from position -1 (i.e. last base

of the PAM) to +8 of sp1.8 differs from the canonical Chi site sequence (5□-GCTGGTGG-3□)

[58] by a single nucleotide; Chi sites are strongly enriched in the *E. coli* K-12 genome [59].

Moreover, positions 3-7 of sp1.8 (5□-GCTGG-3□) are a perfect match to a sequence that is

strongly enriched in the *E. coli* K-12 genome [59]. We conclude that extensive off-target binding

driven by sp1.8 is likely due to a combination of high association with Cascade, and a relatively

high abundance of potential binding sites in the genome.

**METHODS**

**Strains and plasmids**

All strains, plasmids, oligonucleotides and purchased, chemically synthesized dsDNA fragments

are listed in Table S5. All strains are derivatives of MG1655 [59]. CB386 has been previously

described [38]. CB386 contains a chloramphenicol resistance cassette in place of *cas3*. We

removed this cassette using Flp recombinase, expressed from plasmid pCP20 [60], to generate

strain AMD536. Epitope-tagged strains AMD543 and AMD554 (Cse1-FLAG$_3$ and FLAG$_3$-Cas5,

respectively) are derivatives of CB386, and were generated using the previously described

FRUIT method of recombineering [61]. Cse1 was C-terminally tagged in AMD543 by inserting

a FLAG$_3$ tag immediately upstream of codon 495 using oligonucleotides JW6364 and JW6365.

Tagging of Cse1 resulted in an 8 amino acid C-terminal truncation. We predicted based on

phylogenetic comparisons and on structural data [49] that this truncation would not impact the

function of Cse1. Cas5 was N-terminally tagged in AMD554 by inserting FLAG$_3$ using

oligonucleotides JW6272 and JW6273. LC060 is a derivative of AMD536 and was generated

using (i) FRUIT [61] with oligonucleotides JW7537-JW7540 to delete the CRISPR-II locus, (ii)

P1 transduction of the CB386 (Δ*cas3* P*cse1*)::(cat::P$_{J23199}$) region, (iii) FRUIT [61] to C-

terminally tag Cse1 with FLAG$_3$ (as described above for AMD543), and (iv) pCP20-expressed

Flp recombinase [60] to remove the *cat* cassette. LC074 is a derivative of AMD536 in which the

CRISPR-I array was deleted using FRUIT [61] with oligonucleotides JW7529 and JW7530 and a

synthesized dsDNA fragment (gBlock 14148263; Integrated DNA technologies). LC077 is a

derivative of LC074 in which Cse1 was C-terminally tagged with FLAG$_3$ (as described above for

AMD543). AMD566 is a derivative of AMD536 in which Cse1 was C-terminally tagged with

FLAG$_3$ (as described above for AMD543). LC099 is a derivative of AMD566 in which the off-

target binding site for Cascade in *yggX* was mutated using FRUIT [61] with oligonucleotides

JW7635-8. LC103 is a derivative of AMD536 in which the the *yggX* gene was replaced with a

kanamycin resistance cassette using P1 transduction from the Keio Collection Δ*yggX*::*kan*$^R$ strain

[62]. LC106 is a derivative of LC103 with an unmarked, scar-free deletion of *cas1* made using

FRUIT with oligonucleotides JW7898-JW7901. AMD688 is a strain that contains a previously

reported *yfp* reporter construct that can be used to measure adaptation levels [43]. AMD688 was

constructed by P1-transduction of the Δ*cas3*::*cat* cassette from CB386 into MLS1003 (provided

by the Lundgren laboratory). The *cat* gene was removed using Flp recombinase, expressed from

plasmid pCP20 [60]. AMD688 has an intact copy of the CRISPR-I array (co-transduced with the

Δ*cas3*::*cat* cassette from CB386) but lacks the CRISPR-II array.

Plasmids that express crRNAs targeting the *lacZ* promoter (pCB380) and *araB* promoters

(pCB381) have been described previously [38]. All other crRNA-expressing plasmids are

derivatives of pAMD179. pAMD179 was constructed by amplifying a DNA fragment from

plasmid pAMD172 (Integrated DNA Technologies) using with oligonucleotides JW6421 and

JW6513. This DNA fragment was cloned into pBAD24 [63] cut with *Nhe*I and *Hin*dIII (NEB)

using the In-Fusion method (Clontech). The inserted fragment contains two repeats from the

CRISPR-I array, separated by a stuffer fragment containing *Xho*I and *Sac*II restriction sites, and

an intrinsic transcription terminator downstream of the second repeat. To clone individual

spacers, pairs of oligonucleotides were annealed, extended, and inserted using In-Fusion

(Clontech) into the *Xho*I and *Sac*II sites of pAMD179 to generate pLC008 (with oligonucleotides

JW6518 and JW7911), pLC010 (with oligonucleotides JW6518 and JW7912), and pAMD189

(with oligonucleotides JW7598 and JW7693). Note that the derivatives of sp1.8 expressed from

pLC008 and pLC010 differ from sp1.8 at the last two nucleotide positions to facilitate cloning.

These mismatches are not expected to affect crRNA function [23,38].

pLC021, pLC022, and pLC057 are derivatives of pBAD24 [63] that contain a protospacer

matching the off-target Cascade binding site in *yggX* (pLC021), a protospacer with a perfect

match to sp1.8 (pLC022), or a protospacer with a perfect match to sp1.2 (pLC057). These

plasmids were constructed by annealing and extending pairs of oligonucleotides (JW7913 and

JW7914 for pLC021, JW7924 and JW7925 for pLC022, and JW9131 and JW9132 for pLC057),

and cloning the resultant DNA fragments into the *Eco*RV and *Sph*I sites of pBAD24. pAMD191

is a derivative of pBAD33 [63] that expresses *cas3* under arabinose control. To construct

pAMD191, *cas3* was amplified by colony PCR using oligonucleotides JW7736 and JW7738.

The PCR product was cloned into the *Sac*I and *Hin*dIII sites of pBAD33 using In-Fusion

(Clontech). All protospacers described in Figure 5 are cloned into plasmid pLC020, the "pre-

protospacer plasmid", which is a derivative of pBAD24 [63]. pLC020 was generated by cloning

the ~500 bp region upstream of *E. coli thyA* (amplified by colony PCR using oligonucleotides

JW8040 and JW8128) and the ~500 bp region downstream of *E. coli thyA* (amplified by colony

PCR using oligonucleotides JW8042 and JW8043) into the *Eco*RI site of pBAD24 using In-

Fusion (Clontech), simultaneously generating a new *Eco*RI site between the upstream and

downstream regions of *thyA*. The *thyA* gene was then amplified by colony PCR using a universal

forward primer (oligonucleotide JW8129) and each of 13 reverse primers (oligonucleotides

JW8130, JW8139, JW8145, JW8169, JW8499-JW8502, and JW8675-JW8679) containing the

13 protospacer variants described in Figure 5. The resulting PCR products were cloned into the

*Eco*RI site of the pBAD24 derivative using In-Fusion (Clontech) to generate plasmids pLC023-

pLC035 (see Table S5 for details). Note that plasmids pLC024 and pLC025 differ from pLC023

and pLC026-pLC035 at the nucleotide position immediately adjacent to the protospacer, on the

PAM-distal side. Differences at this nucleotide position are not expected to affect Cascade

binding, interference or primed adaptation.

**ChIP-qPCR**

Cells were grown overnight in LB, subcultured in LB supplemented with 0.2% arabinose and

100 µg/mL ampicillin at 37 °C with aeration to an $OD_{600}$ of ~0.6. AMD566 and LC099 with

either pLC008 or pLC010 were used for ChIP-qPCR. ChIP-qPCR was performed as described

previously [64], except that 2 µL anti-FLAG M2 monoclonal antibody (Sigma) and 1 µL anti-$\sigma^{54}$

monoclonal antibody (NeoClone) were included simultaneously in the immunoprecipitation step.

qPCR was performed using oligonucleotides JW7490-1 (amplifies the off-target site in *yggX*)

and JW7922-3 (amplifies the region upstream of *hypA*). Since $\sigma^{54}$ is known not to bind within

*yggX* [65], we were able to normalize binding of Cse1 within *yggX* to the binding of $\sigma^{54}$

upstream of *hypA*.

**ChIP-seq**

Strains AMD543, LC060, LC077, AMD543/AMD554 with pCB380/pCB381, and strain LC077

with pLC008, were used for ChIP-seq of Cse1-FLAG$_3$ and FLAG$_3$-Cas5, except that ampicillin

was only included for experiments involving a crRNA-expressing plasmid, and arabinose was

only included for experiments using pLC008. Cells were grown and processed as described for

ChIP-qPCR. ChIP-seq was performed in duplicate, following a previously described protocol

[66] using 2 µL anti-FLAG M2 monoclonal antibody (Sigma). Sequencing was performed on an

Illumina High-Seq 2000 Instrument (Next-Generation Sequencing and Expression Analysis

Core, State University of New York at Buffalo) or an Illumina Next-Seq Instrument (Wadsworth

Center Applied Genomic Technologies Core). ChIP-seq data analysis was performed as

previously described [67], with reads mapped to the updated MG1655 *E. coli* genome (accession

code U00096.3). Relative sequence coverage values were calculated by dividing the sequence

read coverage at a given genomic location by (total number of sequence reads in the

run/100,000). Values plotted in Figures 1A-B, 2A and 2D are the maximum values in 1 kbp

regions across the genome. $R^2$ values comparing ChIP-seq datasets were calculated by

comparing read coverage at peak centers for all peaks identified for the analyzed datasets. Read

coverage at peak centers was determined using a custom Python script. Sequence motifs were

identified using MEME (version 4.12.0) [68] with default parameters.

**RNA-seq**

RNA-seq was performed in duplicate with strains AMD536 and LC074, with and without

pAMD191. Cells were grown overnight in LB, subcultured in LB (supplemented with 0.2%

arabinose and 100 µg/mL ampicillin for experiments involving pAMD191) at 37 °C with

aeration to an $OD_{600}$ of ~0.6. RNA was purified using a modified hot phenol method, as

previously described [69]. Purified RNA was treated with 2 µL DNase (TURBO DNA-free kit;

Life Technologies) for 45 minutes at 37 °C, followed by phenol extraction and ethanol

precipitation. The RiboZero kit (Epicentre) was used to remove rRNA, and strand-specific

cDNA libraries were created using the ScriptSeq 2.0 kit (Epicentre). Sequencing was performed

using an Illumina Next-Seq Instrument (Wadsworth Center Applied Genomic Technologies

Core). Differential RNA expression analysis was performed using Rockhopper (version 2.03)

with default parameters [70]. Differences in RNA levels were considered statistically significant

for genes with $q$-values $\leq 0.01$.

**Plasmid transformation efficiency assay**

LC103 was transformed with either empty pBAD33 or pAMD191 (expresses *cas3*) and these

strains were then transformed with either pBAD24 (no protospacer), pLC021 (protospacer with a

perfect match to sp1.8), pLC022 (protospacer with an imperfect match to sp1.8, corresponding to

the off-target site in *yggX*) or pLC057 (protospacer with a perfect match to sp1.2). Cells were

plated on M9 medium supplemented with 0.2% glycerol, 0.2% arabinose, 100 μg/mL ampicillin

and 30 μg/mL chloramphenicol at 37 °C. After overnight growth, colonies were counted, and the

relative transformation efficiency was calculated as the ratio of transformants for pAMD191-

containing cells to transformants for pBAD33-containing cells for each transformed protospacer-

containing plasmid.

**PCR to assess primed adaptation**

Primed adaptation was assessed for AMD536 with pAMD191 and either pLC021 or pLC022

(Figure 4B), and MG1655/AMD536/AMD543/AMD544 with pAMD191 and pAMD189

(expresses a self-targeting crRNA; Figure S1). Cells were grown overnight in LB supplemented

with 100 μg/mL ampicillin and 30 μg/mL chloramphenicol at 37°C with aeration, and sub-

cultured the next day in LB supplemented with chloramphenicol and 0.2% arabinose at 37°C

with aeration for six hours. Cells were pelleted from 1 mL of culture by centrifugation, and cell

pellets were frozen at -20°C. PCRs were then performed on the cell pellets, amplifying the

CRISPR-II array using oligonucleotides JW7818 and JW7819. PCR products were visualized on

acrylamide gels.

**Sequence analysis of protospacers from a pooled ChIP library**

LC099 with each of the 13 protospacer variant plasmids (pLC23-pLC035), was grown overnight

in LB supplemented with 100 μg/mL ampicillin. 10 mL subcultures were grown in LB

supplemented with 100 μg/mL ampicillin and 0.2% arabinose at 37°C with aeration to an $OD_{600}$

of ~0.6. 3 mL from each culture were combined. ChIP was performed on mixed cultures using 2

μL M2 anti-FLAG monoclonal antibody (Sigma), as previously described [64]. A Zymo "PCR

Clean and Concentrate" kit was used to purify ChIP and input DNA. A 50 μL FailSafe

(Epicentre) PCR reaction using FailSafe PCR 2X PreMix "C" and 5.48 ng of ChIP DNA was

performed following the manufacturer's instructions, using oligonucleotide JW8567 and each of

oligonucleotides JW8537, JW8556, JW8557, JW8558, JW8559, JW8561, JW8562, JW8563,

JW8564, and JW8565 (these incorporate different Illumina indices). PCR products were purified

and concentrated using 0.8X Ampure Beads (Beckman Coulter Life Sciences) and sequenced on

an Illumina Mi-Seq Instrument (Wadsworth Center Applied Genomic Technologies Core).

Sequence reads were mapped to each of the 13 protospacer variants using a custom Python

script. Relative protospacer abundance in input and ChIP samples for each protospacer were

normalized to the total sequence reads. Values for normalized protospacer abundance were

further normalized to values from the input sample. Protospacer abundance values are reported

relative to those for the optimal protospacer (variant i in Figure 5).

**Measuring interference for a pooled protospacer library**

Overnight cultures of LC106 strains with each of the 13 protospacer plasmids (pLC023-pLC035)

were grown in LB with 100 μg/mL ampicillin and 30 μg/mL kanamycin. All 13 cultures were

combined to make a single subculture (7.7 μL of each overnight culture into a single 10 mL

culture). Electrocompetent cells were made and transformed with either empty pBAD33 or

pAMD191 (pBAD33-*cas3*). Transformants were plated onto M9 agar supplemented with 0.2%

glycerol, 0.2% arabinose, and 30 μg/mL chloramphenicol, and grown overnight at 37°C. Cells

were scraped off plates, washed in LB, and protospacers were PCR amplified from cell pellets

with oligonucleotide JW8567 and each of oligonucleotides JW8537, JW8558, JW8559, JW8562,

JW8563, and JW8566 (these incorporate different Illumina indices). PCR products were purified

and concentrated with 0.8X AMPure Beads (Beckman Coulter Life Sciences), and sequenced

using a Illumina MiSeq Instrument (Wadsworth Center Applied Genomic Technologies Core).

Sequence reads were mapped to each of the 13 protospacer variants using a custom Python

script. "Relative Interference Efficiency" was calculated for each protospacer variant by dividing

the number of sequence reads from cells transformed with empty pBAD33 by the number of

sequence reads from cells transformed with pAMD191 (pBAD33-*cas3*), and normalizing to the

value for the protospacer with a CCG PAM (variant ii in Figure 5).

**Measuring primed adaptation using a YFP fluorescent reporter**

MLS1003 was transformed with each of plasmids LC023-LC035, and each of these strains was

transformed with the *cas3*-expressing plasmid pAMD191. Cells were grown overnight at 37 °C

with shaking in LB supplemented with 100 μg/ml ampicillin and 30 μg/ml chloramphenicol.

Cells were subcultured 1:100 for six hours in LB supplemented with 0.2% L-arabinose and 20

μg/ml chloramphenicol at 37 °C with shaking. Cells were pelleted by centrifugation and

resuspended in M9 minimal medium in twice the original volume ($OD_{600}$ values of ~1). Cells

were transferred to 5 ml polystyrene round-bottom tubes and were analyzed by flow cytometry

for single-cell detection of *yfp* expression using the BD FACSAria iiu cell sorter. 100,000 events

were recorded for each sample. Experiments were performed for between three and ten

independent biological replicates.

**ACKNOWLEDGEMENTS**

We thank Chase Beisel for sharing strains and plasmids. We thank Lina Amlinger and Magnus

Lundgren for the YFP-expressing adaptation reporter strain. We thank the Wadsworth Center

Applied Genomic Technologies Core Facility and the University at Buffalo Genomics and

Bioinformatics Core Facility for Sanger and MiSeq sequencing. We thank the Wadsworth Center

Media and Tissue Culture and Glassware Core Facilities. We thank Todd Gray, Keith

Derbyshire, and Shailab Shrestha for helpful discussions. This study was supported by NIH

Grant AI126416 (to J.T.W.) and a University at Albany, SUNY, RNA Fellowship (to L.A.C.).

**REFERENCES**

1.  Grissa I, Vergnaud G, Pourcel C. The CRISPRdb database and tools to display CRISPRs

and to generate dictionaries of spacers and repeats. BMC Bioinformatics. 2007;8: 1–10.

doi:10.1186/1471-2105-8-172

2.  Bolotin A, Quinquis B, Sorokin A, Ehrlich SD. Clustered regularly interspaced short

palindrome repeats (CRISPRs) have spacers of extrachromosomal origin. Microbiol Read

Engl. 2005;151: 2551–2561. doi:10.1099/mic.0.28048-0

3.  Makarova KS, Grishin NV, Shabalina SA, Wolf YI, Koonin EV. A putative RNA-

interference-based immune system in prokaryotes: computational analysis of the predicted

enzymatic machinery, functional analogies with eukaryotic RNAi, and hypothetical

mechanisms of action. Biol Direct. 2006;1: 7. doi:10.1186/1745-6150-1-7

4.  Mojica FJM, Díez-Villaseñor C, García-Martínez J, Soria E. Intervening sequences of

regularly spaced prokaryotic repeats derive from foreign genetic elements. J Mol Evol.

2005;60: 174–182. doi:10.1007/s00239-004-0046-3

5.  Pourcel C, Salvignol G, Vergnaud G. CRISPR elements in Yersinia pestis acquire new

repeats by preferential uptake of bacteriophage DNA, and provide additional tools for

evolutionary studies. Microbiology. 2005;151: 653–663. doi:10.1099/mic.0.27437-0

6.  Nuñez JK, Kranzusch PJ, Noeske J, Wright AV, Davies CW, Doudna JA. Cas1–Cas2

complex formation mediates spacer acquisition during CRISPR–Cas adaptive immunity.

Nat Struct Mol Biol. 2014;21: 528–534. doi:10.1038/nsmb.2820

7.    Brouns SJJ, Jore MM, Lundgren M, Westra ER, Slijkhuis RJH, Snijders APL, et al. Small

CRISPR RNAs guide antiviral defense in prokaryotes. Science. 2008;321: 960–964.

doi:10.1126/science.1159689

8.    Carte J, Wang R, Li H, Terns RM, Terns MP. Cas6 is an endoribonuclease that generates

guide RNAs for invader defense in prokaryotes. Genes Dev. 2008;22: 3489–3496.

doi:10.1101/gad.1742908

9.    Jore MM, Lundgren M, van Duijn E, Bultema JB, Westra ER, Waghmare SP, et al.

Structural basis for CRISPR RNA-guided DNA recognition by Cascade. Nat Struct Mol

Biol. 2011;18: 529–536. doi:10.1038/nsmb.2019

10.   Mojica FJM, Díez-Villaseñor C, García-Martínez J, Almendros C. Short motif sequences

determine the targets of the prokaryotic CRISPR defence system. Microbiol Read Engl.

2009;155: 733–740. doi:10.1099/mic.0.023960-0

11.   Sashital DG, Wiedenheft B, Doudna JA. Mechanism of foreign DNA selection in a

bacterial adaptive immune system. Mol Cell. 2012;46: 606–615.

doi:10.1016/j.molcel.2012.03.020

12.   Hayes RP, Xiao Y, Ding F, van Erp PB, Rajashankar K, Bailey S, et al. Structural basis for

promiscuous PAM recognition in type I-E Cascade from E. coli. Nature. 2016;530: 499–

503.

13.   Hochstrasser ML, Taylor DW, Bhat P, Guegler CK, Sternberg SH, Nogales E, et al. CasA

mediates Cas3-catalyzed target degradation during CRISPR RNA-guided interference. Proc

Natl Acad Sci U S A. 2014;111: 6618–6623. doi:10.1073/pnas.1405079111

14.  Mulepati S, Bailey S. In vitro reconstitution of an Escherichia coli RNA-guided immune

system reveals unidirectional, ATP-dependent degradation of DNA target. J Biol Chem.

2013;288: 22184–22192. doi:10.1074/jbc.M113.472233

15.  Mulepati S, Héroux A, Bailey S, 19;345(6203):1479-84. S 2014 S. Crystal structure of a

CRISPR RNA-guided surveillance complex bound to a ssDNA target. Science. 2014;345:

1479–1484.

16.  Rutkauskas M, Sinkunas T, Songailiene I, Tikhomirova MS, Siksnys V, Seidel R.

Directional R-Loop Formation by the CRISPR-Cas Surveillance Complex Cascade

Provides Efficient Off-Target Site Rejection. Cell Rep. 2015;

doi:10.1016/j.celrep.2015.01.067

17.  Westra ER, van Erp PB, Künne T, Wong SP, Staals RH, Seegers CL, et al. CRISPR

immunity relies on the consecutive binding and degradation of negatively supercoiled

invader DNA by Cascade and Cas3. Mol Cell. 2012;46: 595–605.

18.  Szczelkun MD, Tikhomirova MS, Sinkunas T, Gasiunas G, Karvelis T, Pschera P, et al.

Direct observation of R-loop formation by single RNA-guided Cas9 and Cascade effector

complexes. Proc Natl Acad Sci U S A. 2014;111: 9798–9803.

doi:10.1073/pnas.1402597111

19.  Sinkunas T, Gasiunas G, Waghmare SP, Dickman MJ, Barrangou R, Horvath P, et al. In

vitro reconstitution of Cascade-mediated CRISPR immunity in Streptococcus thermophilus.

EMBO J. 2013;32: 385–394. doi:10.1038/emboj.2012.352

20.  Yosef I, Goren MG, Qimron U. Proteins and DNA elements essential for the CRISPR

adaptation process in Escherichia coli. Nucleic Acids Res. 2012;40: 5569–5576.

doi:10.1093/nar/gks216

21.  Datsenko KA, Pougach K, Tikhonov A, Wanner BL, Severinov K, Semenova E. Molecular

memory of prior infections activates the CRISPR/Cas adaptive bacterial immunity system.

Nat Commun. 2012;3: 945. doi:10.1038/ncomms1937

22.  Swarts DC, Mosterd C, van Passel MWJ, Brouns SJJ. CRISPR Interference Directs Strand

Specific Spacer Acquisition. PLoS ONE. 2012;7: e35888.

doi:10.1371/journal.pone.0035888

23.  Fineran PC, Gerritzen MJH, Suárez-Diez M, Künne T, Boekhorst J, Hijum SAFT van, et al.

Degenerate target sites mediate rapid primed CRISPR adaptation. Proc Natl Acad Sci.

2014;111: E1629–E1638. doi:10.1073/pnas.1400071111

24.  Staals RHJ, Jackson SA, Biswas A, Brouns SJJ, Brown CM, Fineran PC. Interference-

driven spacer acquisition is dominant over naive and primed adaptation in a native

CRISPR–Cas system. Nat Commun. 2016;7: 12853. doi:10.1038/ncomms12853

25.  Savitskaya E, Semenova E, Dedkov V, Metlitskaya A, Severinov K. High-throughput

analysis of type I-E CRISPR/Cas spacer acquisition in E. coli. RNA Biol. 2013;10: 716–

725. doi:10.4161/rna.24325

26.  Li M, Wang R, Zhao D, Xiang H. Adaptation of the Haloarcula hispanica CRISPR-Cas

system to a purified virus strictly requires a priming process. Nucleic Acids Res. 2014;42:

2483–2492. doi:10.1093/nar/gkt1154

27.    Richter C, Dy RL, McKenzie RE, Watson BNJ, Taylor C, Chang JT, et al. Priming in the

783            Type I-F CRISPR-Cas system triggers strand-independent spacer acquisition, bi-

784            directionally from the primed protospacer. Nucleic Acids Res. 2014;42: 8516–8526.

785            doi:10.1093/nar/gku527

786    28.    Redding S, Sternberg SH, Marshall M, Gibb B, Bhat P, Guegler CK, et al. Surveillance and

787            Processing of Foreign DNA by the Escherichia coli CRISPR-Cas System. Cell. 2015;163:

788            854–865. doi:10.1016/j.cell.2015.10.003

789    29.    Semenova E, Jore MM, Datsenko KA, Semenova A, Westra ER, Wanner B, et al.

790            Interference by clustered regularly interspaced short palindromic repeat (CRISPR) RNA is

791            governed by a seed sequence. Proc Natl Acad Sci USA. 2011;108: 10098–10103.

792    30.    Wiedenheft B, van Duijn E, Bultema JB, Bultema J, Waghmare SP, Waghmare S, et al.

793            RNA-guided complex from a bacterial immune system enhances target recognition through

794            seed sequence interactions. Proc Natl Acad Sci U S A. 2011;108: 10092–10097.

795            doi:10.1073/pnas.1102716108

796    31.    Xue C, Seetharam AS, Musharova O, Severinov K, Brouns SJ, Severin AJ, et al. CRISPR

797            interference and priming varies with individual spacer sequences. Nucleic Acids Res.

798            2015;43: 10831–10847.

799    32.    Künne T, Kieper SN, Bannenberg JW, Vogel AIM, Miellet WR, Klein M, et al. Cas3-

800            Derived Target DNA Degradation Fragments Fuel Primed CRISPR Adaptation. Mol Cell.

801            2016;63: 852–864. doi:10.1016/j.molcel.2016.07.011

33.   Semenova E, Savitskaya E, Musharova O, Strotskaya A, Vorontsova D, Datsenko KA, et

al. Highly efficient primed spacer acquisition from targets destroyed by the Escherichia coli

type I-E CRISPR-Cas interfering complex. Proc Natl Acad Sci. 2016; 201602639.

doi:10.1073/pnas.1602639113

34.   Westra ER, Semenova E, Datsenko KA, Jackson RN, Wiedenheft B, Severinov K, et al.

Type I-E CRISPR-Cas Systems Discriminate Target from Non-Target DNA through Base

Pairing-Independent PAM Recognition. PLoS Genet. 2013;9.

doi:10.1371/journal.pgen.1003742

35.   Blosser TR, Loeff L, Westra ER, Vlot M, Künne T, Sobota M, et al. Two distinct DNA

binding modes guide dual roles of a CRISPR-Cas protein complex. Mol Cell. 2015;58: 60–

70. doi:10.1016/j.molcel.2015.01.028

36.   Stern A, Keren L, Wurtzel O, Amitai G, Sorek R. Self-targeting by CRISPR: gene

regulation or autoimmunity? Trends Genet TIG. 2010;26: 335–340.

doi:10.1016/j.tig.2010.05.008

37.   Vercoe RB, Chang JT, Dy RL, Taylor C, Gristwood T, Clulow JS, et al. Cytotoxic

Chromosomal Targeting by CRISPR/Cas Systems Can Reshape Bacterial Genomes and

Expel or Remodel Pathogenicity Islands. PLoS Genet. 2013;9: e1003454.

doi:10.1371/journal.pgen.1003454

38.   Luo ML, Mullis AS, Leenay RT, Beisel CL. Repurposing endogenous type I CRISPR-Cas

systems for programmable gene repression. Nucleic Acids Res. 2014;43: 674–681.

39.    Semenova E, Kuznedelov K, Datsenko KA, Boudry PM, Savitskaya EE, Medvedeva S, et

823           al. The Cas6e ribonuclease is not required for interference and adaptation by the E. coli

824           type I-E CRISPR-Cas system. Nucleic Acids Res. 2015;43: 6049–6061.

825           doi:10.1093/nar/gkv546

826    40.    Rath D, Amlinger L, Hoekzema M, Devulapally PR, Lundgren M. Efficient programmable

827           gene silencing by Cascade. Nucleic Acids Res. 2015;43: 237–246.

828    41.    Li R, Fang L, Tan S, Yu M, Li X, He S, et al. Type I CRISPR-Cas targets endogenous

829           genes and regulates virulence to evade mammalian host immunity. Cell Res. 2016;26:

830           1273–1287. doi:10.1038/cr.2016.135

831    42.    Müller-Esparza H, Randau L. Commentary: Type I CRISPR-Cas targets endogenous genes

832           and regulates virulence to evade mammalian host immunity. Front Microbiol. 2017;8: 319.

833           doi:10.3389/fmicb.2017.00319

834    43.    Amlinger L, Hoekzema M, Wagner EGH, Koskiniemi S, Lundgren M. Fluorescent

835           CRISPR Adaptation Reporter for rapid quantification of spacer acquisition. Sci Rep.

836           2017;7: 10392. doi:10.1038/s41598-017-10876-z

837    44.    Wu X, Scott DA, Kriz AJ, Chiu AC, Hsu PD, Dadon DB, et al. Genome-wide binding of

838           the CRISPR endonuclease Cas9 in mammalian cells. Nat Biotechnol. 2014;32: 670–676.

839           doi:10.1038/nbt.2889

840    45.    Kuscu C, Arslan S, Singh R, Thorpe J, Adli M. Genome-wide analysis reveals

841           characteristics of off-target sites bound by the Cas9 endonuclease. Nat Biotechnol. 2014;32:

842           677–683. doi:10.1038/nbt.2916

843    46.    Duan J, Lu G, Xie Z, Lou M, Luo J, Guo L, et al. Genome-wide identification of

844            CRISPR/Cas9 off-targets in human genome. Cell Res. 2014;24: 1009–1012.

845            doi:10.1038/cr.2014.87

846    47.    Leenay RT, Maksimchuk KR, Slotkowski RA, Agrawal RN, Gomaa AA, Briner AE, et al.

847            Identifying and Visualizing Functional PAM Diversity across CRISPR-Cas Systems. Mol

848            Cell. 2016;62: 137–147. doi:10.1016/j.molcel.2016.02.031

849    48.    Jackson RN, Golden SM, van Erp PBG, Carter J, Westra ER, Brouns SJJ, et al. Crystal

850            structure of the CRISPR RNA-guided surveillance complex from Escherichia coli. Science.

851            2014;345: 1473–1479. doi:10.1126/science.1256328

852    49.    Zhao H, Sheng G, Wang J, Wang M, Bunkoczi G, Gong W, et al. Crystal structure of the

853            RNA-guided immune surveillance Cascade complex in Escherichia coli. Nature. 2014;515:

854            147–150. doi:10.1038/nature13733

855    50.    Jung C, Hawkins JA, Jones SK, Xiao Y, Rybarski JR, Dillard KE, et al. Massively Parallel

856            Biophysical Analysis of CRISPR-Cas Complexes on Next Generation Sequencing Chips.

857            Cell. 2017;170: 35–47.e13. doi:10.1016/j.cell.2017.05.044

858    51.    Xiao Y, Luo M, Hayes RP, Kim J, Ng S, Ding F, et al. Structure Basis for Directional R-

859            loop Formation and Substrate Handover Mechanisms in Type I CRISPR-Cas System. Cell.

860            2017;170: 48–60.e11. doi:10.1016/j.cell.2017.06.012

861    52.    Xue C, Whitis NR, Sashital DG. Conformational Control of Cascade Interference and

862            Priming Activities in CRISPR Immunity. Mol Cell. 2016;64: 826–834.

863            doi:10.1016/j.molcel.2016.09.033

864    53.    Severinov K, Ispolatov I, Semenova E. The Influence of Copy-Number of Targeted

865           Extrachromosomal Genetic Elements on the Outcome of CRISPR-Cas Defense. Front Mol

866           Biosci. 2016;3: 45. doi:10.3389/fmolb.2016.00045

867    54.    Luo ML, Jackson RN, Denny SR, Tokmina-Lukaszewska M, Maksimchuk KR, Lin W, et

868           al. The CRISPR RNA-guided surveillance complex in Escherichia coli accommodates

869           extended RNA spacers. Nucleic Acids Res. 2016;44: 7385–7394. doi:10.1093/nar/gkw421

870    55.    Heussler GE, O'Toole GA. Friendly Fire: Biological Functions and Consequences of

871           Chromosomal Targeting by CRISPR-Cas Systems. J Bacteriol. 2016;198: 1481–1486.

872           doi:10.1128/JB.00086-16

873    56.    Pul Ü, Wurm R, Arslan Z, Geißen R, Hofmann N, Wagner R. Identification and

874           characterization of E. coli CRISPR-cas promoters and their silencing by H-NS. Mol

875           Microbiol. 2010;75: 1495–1512. doi:10.1111/j.1365-2958.2010.07073.x

876    57.    Reimann V, Alkhnbashi OS, Saunders SJ, Scholz I, Hein S, Backofen R, et al. Structural

877           constraints and enzymatic promiscuity in the Cas6-dependent generation of crRNAs.

878           Nucleic Acids Res. 2017;45: 915–925.

879    58.    Smith GR, Kunes SM, Schultz DW, Taylor A, Triman KL. Structure of chi hotspots of

880           generalized recombination. Cell. 1981;24: 429–436.

881    59.    Blattner FR, Plunkett G, Bloch CA, Perna NT, Burland V, Riley M, et al. The complete

882           genome sequence of Escherichia coli K-12. Science. 1997;277: 1453–1462.

883    60.    Cherepanov PP, Wackernagel W. Gene disruption in Escherichia coli: TcR and KmR

884           cassettes with the option of Flp-catalyzed excision of the antibiotic-resistance determinant.

885           Gene. 1995;158: 9–14.

886    61.    Stringer AM, Singh N, Yermakova A, Petrone BL, Amarasinghe JJ, Reyes-Diaz L, et al.

887           FRUIT, a scar-free system for targeted chromosomal mutagenesis, epitope tagging, and

888           promoter replacement in Escherichia coli and Salmonella enterica. PLoS One. 2012;7:

889           e44841.

890    62.    Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, et al. Construction of

891           Escherichia coli K-12 in-frame, single-gene knockout mutants: the Keio collection. Mol

892           Syst Biol. 2006;2: 2006.0008. doi:10.1038/msb4100050

893    63.    Guzman LM, Belin D, Carson MJ, Beckwith J. Tight regulation, modulation, and high-level

894           expression by vectors containing the arabinose PBAD promoter. J Bacteriol. 1995;177:

895           4121–4130.

896    64.    Bonocora RP, Fitzgerald DM, Stringer AM, Wade JT. Non-canonical protein-DNA

897           interactions identified by ChIP are not artifacts. BMC Genomics. 2013;14: 254.

898           doi:10.1186/1471-2164-14-254

899    65.    Bonocora RP, Smith C, Lapierre P, Wade JT. Genome-Scale Mapping of Escherichia coli

900           σ54 Reveals Widespread, Conserved Intragenic Binding. PLoS Genet. 2015;11: e1005552.

901           doi:10.1371/journal.pgen.1005552

66.  Singh SS, Singh N, Bonocora RP, Fitzgerald DM, Wade JT, Grainger DC. Widespread

suppression of intragenic transcription initiation by H-NS. Genes Dev. 2014;

doi:10.1101/gad.234336.113

67.  Fitzgerald DM, Bonocora RP, Wade JT. Comprehensive Mapping of the Escherichia coli

Flagellar Regulatory Network. PLOS Genet. 2014;10: e1004649.

doi:10.1371/journal.pgen.1004649

68.  Bailey TL, Elkan C. Fitting a mixture model by expectation maximization to discover

motifs in biopolymers. Proc Int Conf Intell Syst Mol Biol. 1994;2: 28–36.

69.  Stringer AM, Currenti SA, Bonocora RP, Petrone BL, Palumbo MJ, Reilly AE, et al.

Genome-Scale Analyses of Escherichia coli and Salmonella enterica AraC Reveal Non-

Canonical Targets and an Expanded Core Regulon. J Bacteriol. 2014;196: 660–671.

70.  McClure R, Balasubramanian D, Sun Y, Bobrovskyy M, Sumby P, Genco CA, et al.

Computational analysis of bacterial RNA-Seq data. Nucleic Acids Res. 2013;41: e140.

 **FIGURE LEGENDS**

**Figure 1. Extensive off-target Cascade binding in *E. coli*. (A)** Binding profile of Cse1 across

the *E. coli* genome, as determined by ChIP-seq, for Cse1-FLAG$_3$ cells (AMD543) expressing a

plasmid-encoded crRNA targeting the *lacZ* promoter region (pCB380). The graph indicates the

relative sequence read coverage (see Methods for details) across the genome in a Cse1 ChIP-

enriched sample. A scale-bar is shown for 1 Mbp. The location of the on-target binding site at

the *lacZ* promoter is shown by the red arrow. **(B)** Binding profile of Cas5 across the *E. coli*

genome, as determined by ChIP-seq, for FLAG$_3$-Cas5 cells (AMD554) expressing a plasmid-

encoded crRNA targeting the *araB* promoter region (pCB381). The location of the on-target

binding site at the *araC* promoter is shown by the red arrow. **(C)** Enriched sequence motif

associated with off-target Cascade binding sites when targeting the *lacZ* promoter, as determined

by MEME. The likely PAM sequence is indicated. The number of identified motifs and the

MEME E-value are shown. **(D)** Enriched sequence motif associated with off-target Cascade

binding sites when targeting the *araB* promoter, as determined by MEME. The likely PAM

sequence is indicated.

**Figure 2. Endogenous crRNAs drive Cascade association with over a hundred**

**chromosomal sites. (A)** Binding profile of Cse1 across the *E. coli* genome, as determined by

ChIP-seq, for Cse1-FLAG$_3$ cells (AMD543) expressing only endogenous crRNAs. The location

of the binding site within *yggX* is shown by the red arrow. **(B)** Enriched sequence motifs

associated with Cascade binding sites in cells expressing only endogenous crRNAs. The four

motifs are associated with four of the CRISPR-I spacers, as indicated. The likely PAM sequence

is also indicated. The number of identified motifs and the MEME E-value are shown. **(C)**

Comparison of Cascade binding events in Cse1-FLAG$_3$ cells with both CRISPR arrays intact

(AMD543) and or CRISPR-II deleted (LC060). Sequence read coverage is shown for CRISPR-

1$^+$ CRISPR-2$^+$ (AMD543) and CRISPR-I$^+$ ΔCRISPR-II cells (LC060), for all ChIP-seq peaks

identified for either strain. **(D)** Binding profile of Cse1 across the *E. coli* genome, as determined

by ChIP-seq, for Cse1-FLAG$_3$ cells expressing only endogenous crRNAs, but with CRISPR-I

deleted (LC077).

**Figure 3. CRISPR-I Spacer #8 is responsible for the majority of Cascade binding in cells**

**expressing only endogenous crRNAs. (A)** Comparison of Cse1-FLAG$_3$ binding events in cells

with both CRISPR arrays intact (AMD543), and cells with CRISPR-I deleted (LC077) that

express CRISPR-I spacer #8 from a plasmid (pLC008). Sequence read coverage is shown for all

ChIP-seq peaks identified for either strain. ChIP-seq peaks associated with the CRISPR-I spacer

#8 motif (first motif listed in Figure 2B) are shown in orange. **(B)** Predicted base-pairing

interaction between CRISPR-I spacer #8 and a protospacer within *yggX*. The PAM is underlined.

**(C)** ChIP-qPCR measurement of Cse1 binding at wild-type (i and iii; AMD566) and mutant (ii

and iv; LC099) protospacers in *yggX* for cells expressing wild-type (i and ii; pLC008) or mutant

(iii and iv; pLC010) CRISPR-I spacer #8 from a plasmid. The mutations in spacer #8 restore

base-pairing potential with the mutant protospacer, as indicated. Values represent the average of

three independent replicate experiments. Error bars show one standard deviation from the mean.

**Figure 4. Off-target Cascade binding events are not associated with interference or primed**

**adaptation. (A)** Relative efficiency of transformation of a *cas3*-expressing plasmid (pAMD191)

into LC103 cells expressing spacer #8 from a native CRISPR-I array, and containing (i) empty

pBAD24 ("No Protospacer"), (ii) a plasmid with a protospacer that base-pairs perfectly with

spacer #8 (pLC022), (iii) a plasmid with a protospacer that has only partial base-pairing with

CRISPR-I spacer #8 (pLC021; protospacer sequence matches the off-target Cascade binding site

in *yggX*), or (iv) a plasmid with a protospacer that base-pairs perfectly with CRISPR-I spacer #2

(pLC057). Note that crRNAs were expressed from the chromosome since both CRISPR arrays

are intact in these strains. Transformation efficiency was calculated relative to that of empty

pBAD33, as described in the Methods. Values represent the average of three independent

replicate experiments. Error bars show one standard deviation from the mean. The calculated

transformation efficiency for protospacers (ii) and (iv) was 0, but the limit of detection in this

assay was $3e^{-5}$. **(B)** PCR-amplification of the start of the CRISPR-II array to detect primed

adaptation in cells expressing CRISPR-I spacer #8 (AMD536), *cas3* (pAMD191), and with (i) a

protospacer that base-pairs perfectly with CRISPR-I spacer #8 (pLC022), (ii) the protospacer

from *yggX* that has only partial base-pairing with CRISPR-I spacer #8 (pLC021), or (iii) empty

vector (pBAD24). L = molecular weight ladder, with marker sizes (bp) indicated. The expected

PCR product sizes are indicated. Note that there is a non-specific PCR product of ~380 bp for

both samples i and ii (indicated by the asterisks).

**Figure 5. Assessment of Cascade binding, interference and primed adaptation for a panel**

**of protospacer variants. (A)** Relative Cse1 association (in strain LC099, which expresses

CRISPR-I spacer #8 crRNA from the native CRISPR-I array) for each of 13 protospacer

variants. The protospacer variants are: (i) optimal protospacer that has a perfect match to

CRISPR-I spacer #8 and an AAG PAM (pLC023); (ii) CCG PAM (pLC027); (iii) ATT PAM

(pLC029); (iv) mismatches at positions 1-3 (pLC031; wild-type is CTG); (v) mismatches at

positions 1 and 3 (pLC033); (vi) mismatches at positions 1 and 3 (pLC035); (vii) mismatches at

positions 2 and 3 (pLC032); (viii) mismatches at positions 2 and 3 (pLC034); (ix) mismatches

across positions 25-32 (pLC024); (x) mismatches across positions 19-32 (pLC025); (xi)

mismatches across positions 1-6 (pLC026); (xii) mismatches across positions 1-6 and 25-32

(pLC028); (xiii) mismatches across positions 7-24 (pLC030). Values represent the average of

five independent replicate experiments. Error bars show one standard deviation from the mean.

**(B)** Relative efficiency of interference (see Methods for details) for each of the indicated

protospacer variants (pLC023-pLC035). Values represent the average of three independent

replicate experiments. Error bars show one standard deviation from the mean. **(C)** Levels of

primed adaptation in AMD688 cells expressing CRISPR-I spacer #8 from a native CRISPR-I

array , *cas3* (pAMD191), and with each of the 13 indicated protospacer variants (pLC023-

pLC035). Adaptation was measured by monitoring the conversion of a *yfp* reporter construct into

the actively expressed state (see methods for details). Values shown are the percentage of the

population that converted to $YFP^+$. Background levels of adaptation were detected for variants

(ii), (x), (xi), (xii) and (xiii).

1002

1003 **SUPPORTING INFORMATION**

1004

**Figure S1. FLAG$_3$-tagged Cse1 and Cas5 are fully functional for primed adaptation.** PCR-

amplification of the start of each of the CRISPR-I and CRISPR-II arrays to detect primed

adaptation in cells expressing *cas3* (pAMD191) and a plasmid-encoded crRNA that perfectly

targets a sequence on the same plasmid (pAMD189). Adaptation was assessed for (i) MG1655

(no *cas* gene expression, except for plasmid-encoded *cas3* (pAMD191)), (ii) MG1655 with

constitutive expression of *cas* genes (AMD536), (iii) MG1655 *cse1*-FLAG$_3$ with constitutive

expression of *cas* genes (AMD543), and (iv) MG1655 FLAG$_3$-Cas5 with constitutive expression

of *cas* genes (AMD554).

1013

**Figure S2. crRNA spacers used in this study. (A)** Sequence of the crRNA spacer targeting the

*lacZ* promoter (pCB380). **(B)** Sequence of the crRNA spacer targeting the *araB* promoter

(pCD381). **(C)** Sequence of the CRISPR-I array. Spacers #1, #3, #4 and #8 are underlined. **(D)**

Sequence of the CRISPR-II array. Spacer #2 is underlined. **(E)** Sequence of a portion of the

CRISPR-I spacer #8 crRNA-expressing plasmid (pLC008). Note that the sequence downstream

of the second repeat (underlined) can be used as a spacer.

1020

1021 **Figure S3. Spacer #2 of CRISPR-II directs Cascade binding in cells lacking CRISPR-I.**

1022 Enriched sequence motif associated with Cascade binding sites in cells expressing only

1023 endogenous crRNAs, where CRISPR-I is deleted (LC077). The motif is associated with

1024 CRISPR-II spacer #2, as indicated. The likely PAM sequence is also indicated. The number of

1025 identified motifs and the MEME E-value are shown.

1026

**Figure S4. Sequence motifs associated with Cascade binding in cells expressing CRISPR-I**

**spacer #8 from a plasmid. (A)** Sequence of the most strongly enriched motif, as identified by

MEME, in ΔCRISPR-I cells (LC077) expressing CRISPR-I spacer #8 from a plasmid (pLC008).

The motif is associated with CRISPR-I spacer #8, as indicated. The likely PAM sequence is also

indicated. The number of identified motifs and the MEME E-value are shown. **(B)** The second

enriched sequence motif, as identified by MEME, in ΔCRISPR-I cells (LC077) expressing

CRISPR-I spacer #8 from a plasmid (pLC008). The motif is associated with the sequence

immediately downstream of the second repeat on the crRNA plasmid, as indicated.

1035

1036

1037     **Table S1. Lists of ChIP-seq peak coordinates.**

1038

1039     **Table S2. Lists of regions used to search for enriched sequence motifs.**

1040

1041     **Table S3. Analysis of RNA-seq data.**

1042

1043     **Table S4. Numbers of potential off-target chromosomal binding sites for spacers in the**

1044     **CRISPR-I array.**

1045

1046     **Table S5. Strains, Plasmids, Oligonucleotides, and Chemically Synthesized dsDNA**

1047     **fragments used in this study.**

# Figure 1

# Figure 2

**A**

Relative Sequence Coverage

CRISPR-I⁺
CRISPR-II⁺

1 Mbp

Genome Position

**B**

n = 74
$E = 1.1e^{-113}$

PAM

sp1.8    CTGCTGGAGCT...

n = 45
$E = 2.1e^{-29}$

PAM

sp1.1    CTTTCGCA...

n = 25
$E = 1.7e^{-6}$

PAM

sp1.3    GGCTCCCTG...

n = 20
$E = 8.0e^{-6}$

PAM

sp1.4    TTTGGATC...

**C**

Relative Sequence Coverage (ΔCRISPR-II)

Relative Sequence Coverage (Wild-Type)

**D**

Relative Sequence Coverage

ΔCRISPR-I
CRISPR-II⁺

1 Mbp

Genome Position

# Figure 3

# Figure 4

## A

Relative Transformation Efficiency

| i | No Protospacer |
| ii | sp1.8 |
| iii | sp1.8 |
| iv | sp1.2 |

## B

```
300
200
        L      i      ii     iii
```

← R S R S R  296 bp

← R S R  235 bp

# Figure 5