1 **Double-digest RAD sequencing outperforms microsatellite loci at assigning**
2 **paternity and estimating relatedness: a proof of concept in a highly promiscuous**
3 **bird**
4

5 Derrick J. Thrasher[3,4], Bronwyn G. Butcher[1], Leonardo Campagna[1,2], Michael S.
6 Webster[3,4], Irby J. Lovette[1,2]
7
8
[1]Fuller Evolutionary Biology Program, Cornell Laboratory of Ornithology, Ithaca, NY
14850, USA
[2]Department of Ecology and Evolutionary Biology, Cornell University, E145 Corson Hall,
Ithaca, NY 14853, USA
[3]Macaulay Library, Cornell Lab of Ornithology, 159 Sapsucker Woods Rd, Ithaca, NY
14850, USA
[4]Department of Neurobiology and Behavior, Cornell University, W361 Mudd Hall, 215
Tower Rd, Ithaca, NY 14853, USA

Keywords: double-digest restriction site-associated DNA sequencing (ddRAD-seq),
microsatellite, single nucleotide polymorphism (SNP), parentage, relatedness,
cooperative breeding

*Corresponding author: Derrick J. Thrasher, Macaulay Library, Cornell Lab of
Ornithology, 159 Sapsucker Woods Rd, Ithaca, NY 14850, USA. Fax: (607) 254-2439.
Email: djt224@cornell.edu

25
26
27 Running title: SNPs outperform microsatellite loci

Abstract

Information on genetic relationships among individuals is essential to many

studies of the behavior and ecology of wild organisms. Parentage and relatedness

assays based on large numbers of SNP loci hold substantial advantages over the

microsatellite markers traditionally used for these purposes. We present a double-digest

restriction site-associated DNA sequencing (ddRAD-seq) analysis pipeline that, as such,

simultaneously achieves the SNP discovery and genotyping steps and which is

optimized to return a statistically powerful set of SNP markers (typically 150-600 after

stringent filtering) from large numbers of individuals (up to 240 per run). We explore the

tradeoffs inherent in this approach through a set of experiments in a species with a

complex social system, the variegated fairy-wren (*Malurus lamberti*), and further validate

it in a phylogenetically broad set of other bird species. Through direct comparisons with

a parallel dataset from a robust panel of highly variable microsatellite markers, we show

that this ddRAD-seq approach results in substantially improved power to discriminate

among potential relatives and substantially more precise estimates of relatedness

coefficients. The pipeline is designed to be universally applicable to all bird species (and

with minor modifications to many other taxa), to be cost- and time-efficient, and to be

replicable across independent runs such that genotype data from different study periods

can be combined and analyzed as field samples are accumulated.

Introduction

Advances in molecular techniques over the past several decades have substantially improved our ability to test questions about animal social behavior by providing reliable information on the genetic relationships among individuals (Westneat *et al.* 1990; Hughes 1998; Avise *et al.* 2002; Griffith *et al.* 2002; Solomon *et al.* 2004; Myers & Zamudio 2004). Microsatellites have been the molecular 'tool-of-choice' for this application since the 1990s, as microsatellite loci are often highly polymorphic, with up to dozens of co-segregating alleles at a single locus (Queller *et al.* 1993; Li *et al.* 2002; Selkoe & Toonen 2006; Guichoux *et al.* 2011). Accordingly, a small number of highly variable microsatellite loci can provide considerable power for discerning genetic relationships among individuals (Queller *et al.* 1993; Blouin 2003; Webster & Reichart 2005). However, microsatellite assays also have some practical drawbacks. Microsatellite laboratory protocols developed for one species are often not suitable for use in other species, especially in more distantly related taxa (Galbusera 2000; Decroocq *et al.* 2003; Hedgecock *et al.* 2004; Primmer *et al.* 2005). Traditional PCR-based microsatellite assays incur substantial financial and lab-bench time investments. The manual scoring of microsatellite alleles also requires substantial researcher time, and can involve various forms of error arising from alleles that have more than one clearly defined peak, allelic drop-out and null allele issues, and the various sources of human error that are inherent in any complicated workflow (Pemberton *et al.* 1995; Hedgecock *et al.* 2004; Hoffman & Amos 2005; Kalinowski *et al.* 2007).

Many of these limitations are less severe in assays based on single-nucleotide

polymorphisms (SNPs), which require fewer steps and have greater automation (Gut

2001; Syvänen 2001; Seeb *et al.* 2011; Davey *et al.* 2011). SNPs are appropriate

alternatives for studies of parentage and relatedness data because they are abundant in

the genome, have low mutation rates (Brumfield *et al.* 2003; Morin *et al.* 2004), and can

be scored semi-automatically (Garvin *et al.* 2010; Guichoux *et al.* 2011). In comparison

to microsatellite-based relationship tests, the primary limitation of SNPs is that SNPs are

typically biallelic, whereas microsatellite loci are often multiallelic, and hence the

statistical power of SNP loci for discriminating parentage and relatedness is far lower on

a per-locus basis (Ball *et al.* 2010). Compared to highly variable microsatellite loci, a

substantially higher number of SNP markers is therefore required to achieve appropriate

power in parentage and relatedness studies (Glaubitz *et al.* 2003; Morin *et al.* 2004;

Coates *et al.* 2009).

Recently, the application of SNPs for use in analyses of parentage, relatedness,

and overall population structure has received greater attention (Glaubitz *et al.* 2003;

Anderson & Garza 2006; Coates *et al.* 2009). Studies in birds (Cramer *et al.* 2011;

Weinman *et al.* 2014; Kaiser *et al.* 2016), fish (Hauser *et al.* 2011), and several

domesticated taxa (Tokarska *et al.* 2009; Fernández *et al.* 2013) have developed SNP

panels with a sufficient number of SNPs to attain a comparable, if not better, level of

resolving power as highly polymorphic microsatellite panels. While each of these studies

manage to identify powerful SNP panels, the SNP genotyping methods used are often

labor intensive, requiring a significant amount of preparatory work at the discovery stage

prior to genotyping of large numbers of individuals. Many of these methods also rely on

reference genomes (Anderson & Garza 2006; Heylar *et al.* 2011), or other genomic

resources (Fernández *et al.* 2013; Weinman *et al.* 2014; Kaiser *et al.* 2016) (e.g.

transcriptome, SNP microarray), for SNP identification. Ultimately, this has afforded

several beneficial examples of the utility of SNPs for parentage and relatedness

analyses, but without an efficient, universal method of SNP discovery and identification.

Restriction site-associated DNA sequencing (RAD-seq) is widely used in

molecular genetic studies (Davey & Blaxter 2010; Etter *et al.* 2012; Puritz *et al.* 2014),

particularly for linkage and quantitative trait locus (QTL) mapping (Baird *et al.* 2008),

genome wide association studies (Davey *et al.* 2011), and phylogeography (Andrews *et*

al. 2016). RAD-seq uses a restriction enzyme to fragment and sample a fraction of a

genome; as it identifies SNPs with no prior knowledge of the genome, it provides a more

universal method of SNP discovery (Willing *et al.* 2011). Double-digest restriction site-

associated DNA sequencing (ddRAD-seq) allows for selection of an even smaller

fraction of the genome through the combined use of two restriction enzymes, affording

the ability to target a smaller total number of SNPs in a greater number of individuals

(Peterson *et al.* 2012; Puritz *et al.* 2014; Kess *et al.* 2016). This ability, in concert with

the fact that no prior knowledge of the genome is needed, makes ddRAD-seq an

attractive method of simultaneous SNP discovery and screening for use in discerning

genetic relationships among individuals.

Here, we describe a ddRAD-based approach to the simultaneous discovery and

screening of high numbers of SNP loci with high power for testing questions about

parentage and relatedness. These protocols are optimized to generate an appropriately

robust set of SNP markers for 240 individuals per run, to be repeatable across runs to

allow the combination of SNP datasets generated at different times, and to be

universally applicable to birds (and with small modifications, to other organisms) without

requiring a species-specific marker discovery step. We validate these methods by

conducting a SNP-based parentage and relatedness study in the highly promiscuous,

and socially complex, variegated fairy-wren (*Malurus lamberti*). We compare the results

with previously generated paternity assignments and relatedness information, based on

microsatellite screens of the same fairy-wren individuals and social groups. To illustrate

the broad utility of this method we report summary statistics for equivalent studies of

parentage in a variety of other species that collectively span much of the phylogenetic

diversity of living birds.

Methods & Materials

Study Population

The variegated fairy-wren, endemic to Australia, is a cooperatively breeding bird

that lives in social groups composed of kin and non-kin (Schodde 1982; Rowley &

Russell 1997). Male dispersal is limited, and rates of extra-pair fertilizations (EPFs) are

high (~68% of all young, assessed with a panel of 12 species-specific microsatellites; DJ

Thrasher, unpublished data). We intensively monitored a color-banded population of the

nominate subspecies, *M. l. lamberti*, on Lake Samsonvale (27^O16' S, 152^O 41' E), 30 km

northwest of Brisbane, Queensland, Australia, from 2012 – 2016. The population

ranges from about 250-300 adults depending on year-to-year conditions. The study site

is bounded on most sides by Lake Samsonvale, and on its westernmost side by a major

highway, which increases our confidence in sampling most, if not all, of the adults in the

population. We also monitored all nesting attempts to measure, mark, and collect blood

samples from nestlings 6 days after hatching. Blood samples were immediately stored in

lysis buffer (White & Densmore 1992), and genomic DNA was later extracted using

Qiagen DNeasy Blood and Tissue kits. DNA concentration was determined using the

Qubit dsDNA BR Assay Kit and the Qubit® Fluorometer (Life Technologies) following

the manufacturers protocol.

Microsatellite development and genotyping

We previously developed twelve polymorphic microsatellite loci for the variegated

fairy-wren (Table S1, Supporting information) following methods described previously in

Nali *et al.* (2014). Briefly, we extracted genomic DNA from blood in lysis buffer from eight

adults in our study population, enriched the mix of DNA with repetitive sequences to

develop an enriched microsatellite library, and conducted an Illumina MiSeq sequencing

run. From this pool of sequences, we optimized twelve loci that amplified well using

polymerase chain reaction (PCR), were polymorphic, and exhibited clearly defined

peaks for genotyping. We designed three multiplexed PCRs for genotyping, and each

amplification reaction contained 1ul of genomic DNA of varying concentrations (1 ng-40

ng per 1 ul). PCR products were combined with the GeneScan 500 base pair LIZ

internal size standard for size-sorting using a 3730 DNA Analyzer. We used Geneious

version 8.0 (Kearse *et al.* 2012) to score alleles. The program automatically identifies

alleles at each locus, and we manually inspected allele calls to minimize genotyping

error. In total, we genotyped 287 adults and 482 nestlings from 226 nests sampled

during the 2012-2016 breeding seasons.

ddRAD sequencing

We selected a subset of the individuals genotyped for microsatellite loci (120

adults and 40 nestlings) for use in our ddRAD-seq experiment and subsequent

analyses. To assess the reliability of our SNP panel for parentage and relatedness

analysis, we chose representative nestlings from all the years of our study. Typically, we

selected one nestling from any individual nest. In a few cases, we selected two nestlings

that prior microsatellite analysis had assigned to the same mother but different fathers.

For each nestling, we included the mother, the social father, and the genetic father as

assigned by previous microsatellite analysis. Our pool of candidate parents included 24

mothers and 78 putative fathers, and 18 randomly selected individuals of both sexes, for

a total of 120 adults.

Our ddRAD-seq protocol is adapted from Peterson et al. (2012) (see Supporting

information for a detailed protocol). Briefly, for each individual, 100ng - 500ng of DNA

(20ul of DNA between concentrations of 5ng/ul - 25ng/ul) were digested with either SbfI

and MspI, or SbfI and EcoRI (NEB), and ligated with one of 20 P1 adapters (each

containing a unique inline barcode) and a P2 adapter (P2-MspI or P2-EcoRI). After

digestion and ligation these samples were pooled in groups of 20 (each with a unqiue P1

adpater) and purified using 1.5X volumes of homemade MagNA made with Sera-Mag

Magnetic Speed-beads (FisherSci) as described by Rohland & Reich (2012). Fragments

of between 450 bp and 600 bp were selected using BluePippin (Sage Science) by the

Cornell University Biotechnology Resource Center (BRC). Following size selection,

index groups and Illumina sequencing adapters were added by performing 11 PCR

cycles with Phusion DNA Polymerase (NEB). These reactions were cleaned up with 0.7x

volumes of MagNA and pooled in equimolar ratios to create a single library for

sequencing on one lane of Illumina HiSeq 2500 (100bp single end, performed by BRC).

By replicating 20 samples (run as a separate index group) with a wider

BluePippin size selection range of 400-700 bp, we explored the inherent trade-off

between the number of samples sequenced on a lane at a given coverage threshold and

the number of SNPs recovered per sample. We similarly explored the effects of using a

less frequent restriction enzyme digest (by substituting the 6 bp cutter EcoRI in place of

the 4 bp cutter MspI) to generate a smaller total number of fragments in our size range,

which in turn should increase the sequencing coverage of the loci screened.

To assess repeatability across index groups, we replicated two index groups,

each comprised of 20 samples: one index group from the standard protocol, and the

index group generated with the rare-cutter EcoRI enzyme (Table 1). We multiplexed the

resulting 240 samples in 12 index groups (with 20 individuals each) which were pooled

into a final library that was sequenced on one lane of an Illumina HiSeq 2500, producing

220,300,739 100 bp single end reads (see Table 1).

SNP data analysis

<u>Quality filtering and demultiplexing</u>

222   After the quality of the reads was assessed using FASTQC version 0.11.5

(www.bioinformatics.babraham.ac.uk/projects/fastqc), we trimmed all sequences to 97bp

using fastX_trimmer (FASTX-Toolkit) to exclude low quality calls near the 3' of the reads.

We subsequently removed reads containing a single base with a Phred quality score of

less than 10 (using fastq_quality_filter). We additionally removed sequences if more

than 5% of the bases had a Phred quality score of less than 20. Using process_radtags

module from the Stacks version 1.37 pipeline (Catchen *et al.* 2013), we demultiplexed

the reads to obtain files with sequences that were specific to each individual.

<u>De novo assembly of RAD loci</u>

232   Because we do not have a sequenced genome for the variegated fairy wren or a

close relative – which is likely to be the case for many non-model organisms involved in

parentage studies – we assembled the sequences de novo using the Stacks pipeline

(Catchen *et al.* 2013). First, we used denovo_map.pl to assemble the reads into a

catalog allowing a minimum stack depth of 5 (m parameter), up to 5 mismatches per

locus within an individual (M paramenter), and 5 mismatches between loci of different

individuals when building the catalog (n parameter). We ran the rxstacks module to filter

loci with a log likelihood of less than -50 (lnl_lim -50) or that were confounded in at least

25% of the population (conf_lim 0.25). We then built a new catalog by rerunning cstacks

and obtained individual genotype calls with sstacks.

<u>SNP filtering</u>

SNPs were exported using the populations module of the Stacks pipeline. All of

our samples were grouped in one population and a locus was exported if it was present

in 95% of the individuals in this population (r parameter) at a stack depth of at least 10

(m parameter). The data were restricted to the first SNP per locus (--write_single_snp),

and a minor allele frequency of at least 0.25 was required to process a nucleotide site (--

min_maf).

Using vcftools version 0.1.14 (Danecek *et al.* 2011), we removed loci that were

not in Hardy-Weinberg equilibrium (---hwe). We obtained a variant call format file that

was converted to structure format in PGD Spider version 2.0.5.0 (Lischer & Excoffier

2012) and further modified to a format compatible with CERVUS version 3.0.7

(Kalinowski *et al.* 2007) using a custom perl script.

Parentage Analysis

We used CERVUS version 3.0.7 (Kalinowski *et al.* 2007) to assign paternity for all

nestlings using our microsatellite and SNP datasets separately. CERVUS uses a two-

step, likelihood-based approach to assign parentage. First, CERVUS compares each

offspring's genotype to that of a candidate parent and a random individual in the

population to calculate a likelihood ratio. This relationship is presented as an LOD score,

which is simply the natural logarithm of the calculated likelihood ratio. Positive LOD

scores indicate that a candidate parent is much more likely to be the true parent,

whereas negative LOD scores indicate that the candidate parent is highly unlikely to be

a true parent. Second, CERVUS conducts a simulation of parentage analysis based on

population allele frequencies and the proportion of potential parents included in the

analysis. The simulation accounts for the possibility of unsampled parents, missing data,

and genotyping errors. Considering these parameters, the simulation calculates critical

LOD scores by comparing the LOD distributions of the most likely parent and all other

candidate parents. The critical LOD score is used to determine the confidence (95% or

80%) of each parentage assignment.

CERVUS allows for different types of parentage analysis, including parent-pair

(sexes known or unknown), maternity (known father, but not mother), and paternity

(known mother, but not father). Variegated fairy-wrens at Lake Samsonvale are relatively

easy to observe, and we were able to assign known mothers behaviorally. We

subsequently confirmed this with microsatellite analyses: females that built and attended

a nest throughout incubation were always the mothers of the nestlings in that nest. In

many systems, a comparable level of demographic knowledge may not be available, so

a marker set must be powerful enough to assign parentage with minimal social

information. To investigate the broader utility of our ddRAD-seq method, we conducted

analyses that relied on the inclusion of the known mother, in addition to analyses

independent of the known mother, which were based only on the father-offspring

relationship. We simulated paternity assignments for 10,000 offspring to determine

critical LOD scores, using slightly different input parameters for each panel

(microsatellites and ddRAD sequencing derived SNPs). Simulations for both used the

following parameters: 78 candidate males, 95% of candidate males sampled, estimated

error rate of 0.01 for mistyped loci and likelihood scores. The proportion of loci typed

across all individuals was different for both panels: 0.997 for the microsatellite

simulation, and 0.961 for the SNP simulation.

For both paternity analyses, we used the trio LOD score and the father-offspring

LOD score from CERVUS to make assignments. The trio LOD score was calculated by

comparing the genotypes of the candidate male and offspring, relative to that of the

known mother. The father-offspring LOD score only accounts for the relationship of the

candidate male and the offspring, independent of the known mother. CERVUS ranked

candidate males by LOD scores in each category, and the highest-ranking males were

assigned as fathers. These rankings should be in agreement, but ambiguous

assignments (different top-ranking males assigned in each category) may occur when

multiple candidate male genotypes closely match an offspring's genotype.

We assessed each CERVUS assignment to determine whether it was plausible,

and whether the assigned male was the social father or an extra-pair sire. Our criteria for

accepting assignments differed slightly for microsatellites and SNPs. For microsatellites,

we automatically accepted the CERVUS assignment if the highest-ranking male was in

agreement for both the trio LOD and the father-offspring LOD, and if the number of

mismatches between the assigned male and the offspring was ≤ 1 (8% of 12 loci). For

SNPs, we also accepted the assignment if the highest-ranking males by LOD score type

were in agreement, and an allowable number of mismatches were not exceeded.

However, for SNPs, our allowable number of mismatches was based on the observed

maximum number of mismatches between a known mother and her known offspring

(max. = 7, mean = 3.4, 2% of 411 loci). For both panels, we accepted the social father

as the genetic sire if he met these respective criteria. If the social father mismatched the

offspring at higher numbers, or had negative LOD scores, the offspring was considered

sired by an extra-pair father. We accepted assignments of extra-pair fathers using the

same criteria outlined above. We did not observe cases in which an offspring could not

be assigned to either its social father, or an extra-pair sire.

Relatedness Analysis

We used the package, "related" (Pew *et al.* 2015), in R version 3.2.5 (R Core

Team 2016) to estimate pairwise relatedness (r) between all pairs of individuals in this

study. This package accounts for genotyping errors, missing data, and can estimate

relatedness using any of seven different estimators (4 non-likelihood-based, and 3

likelihood-based). "related" includes the function, **compareestimators**, which tests the

performance of different estimators on simulated data that share the same

characteristics as the real data. The program uses an allele frequency file to generate

simulated pairs of individuals of known relatedness, and automatically estimates

relatedness using four of the most commonly used estimators (all non-likelihood-based).

The function calculates a correlation coefficient between observed and expected values,

to evaluate which estimator performs best with the data set. Using **compareestimators**

to generate 200 simulated pairs of individuals for each degree of relatedness (i.e., half-

sib, full-sib, parent-offspring, unrelated), we determined that the Wang (2002) estimator

performed best for both our microsatellite and SNP datasets. We obtained point

estimates of relatedness using the Wang (2002) estimator, and evaluated all parent-

offspring relationships that were previously determined in our parentage analysis.

"Related" also evaluates how well different marker sets resolve degrees of

relatedness, given simulated genotypes based on allele frequency files. For both panels,

we used the **familysim** function to generate 200 pairs of individuals for each degree of

relatedness. We then used the **coancestry** function to analyze all pairwise relatedness

values with the Wang (2002) estimator. We created density plots representing

histograms of the relatedness values. These plots show the overlap in relatedness

values for degree of relatedness, and we used them to infer how well each panel

performed at discerning different relationships.

Results

SNP development and analysis

After trimming, filtering and demultiplexing the data, we retained a total of

109,524,874 reads across all index groups, with an average of approximately 9,000,000

reads per index group. Two individuals failed (i.e., had less than 66,000 reads each; one

individual from each of two index groups) and were excluded from further analysis. The

number of reads per sample for the remaining individuals ranged from 213,544 to

810,966 (mean = 459,726 ± 118,771 std. dev.).

Further analysis using the population program from stacks identified loci with at

least 10X coverage, present in 95% of the individuals, and a minimum allele frequency

greater than 25%. We further retained only those loci that were in Hardy-Weinberg

equilibrium. When performing analyses on all 160 individuals from the primary

comparison runs (Table 1), we identified 411 loci that fulfilled these criteria and were

used for downstream analyses.

We varied two aspects of the ddRAD-seq protocol to assess the number of loci

recovered and the reproducibility of the method. As expected, both a reduction in the

range of fragment sizes selected during the construction of the library (compare index 1

(150 bp size selection) and index 10 (300 bp size selection)) or the use of the less

frequent cutter, EcoRI (index 11 and 12), resulted in fewer loci recovered (Table 3).

Those from the EcoRI digest were non-overlapping with those from the MspI index

groups. More importantly, between the replicated index groups in our standard protocol

(index 1 and index 9) we recovered similar numbers of loci (797 and 645, respectively)

with 549 (85.1%) loci found in both datasets (under our stringent filtering criteria).

Paternity assignments

Both panels produced highly concordant results when assigning paternity, but the

SNP panel showed substantially higher power overall. Generally, the microsatellite loci

were more polymorphic, resulting in greater mean polymorphic information content (PIC)

for any given locus. Despite this, the SNP panel performed better because of the large

number of loci obtained through RAD sequencing. This greatly improved the non-

exclusion probabilities across different parentage assignment contexts (Table 2), and

reduced uncertainty in our assignments. Given the known mother, the microsatellite and

SNP panels assigned the same fathers to all 40 offspring with 95% confidence. When

paternity assignments were made without the known mother (no known candidate

parents), both panels again assigned fathers for all 40 offspring with 95% confidence.

However, 5 of these assignments were not in agreement between the two panels. For

these 5 cases, two candidate males had very similar LOD scores under the

microsatellite panel, and the assigned males did not match the males that were

assigned given the known mother. These (and all other) cases were resolved

unambiguously when the SNP panel was used, and the paternity assignments with and

without the known mother were in complete agreement for all offspring (Fig. 1).

Overall, both panels assigned 23 out of 40 nestlings (57.5%) to males that were

not their social father. Due to the nature of our non-random sampling of individuals for

this experiment, and the overall smaller sample size, this value is slightly lower than the

overall rate of 67.6% extra-pair young observed for all years of the study (unpublished

data).

As a measure of certainty for our assignments, we calculated the difference

between LOD scores for the two top-ranked males assigned to each nestling, under

each panel (Fig. 2). Typically, this difference was 8 – 10x higher for the SNP panel

(n=40, mean = 165.0) than for the microsatellite panel (n=40, mean 19.1), a reflection of

the much higher discriminatory power of the SNP dataset. For the SNP panel, many of

the second-ranked males had a strongly negative LOD score, making them extremely

unlikely to be the true father. This was less often true for the microsatellite panel, as the

second-ranked males often had positive, or just slightly negative, LOD scores. Overall

this result illustrates the increased discrimination power achieved by the SNP panel

compared to the microsatellites, which allowed us to assign paternity in cases in which

the microsatellite assignments remained ambiguous or (albeit rarely) misleading.

Relatedness analysis

The SNP panel produced simulated data that closely matched the observed allele

and genotype frequencies (Pearson's correlation coefficient = 0.975). The microsatellite

panel also matched well, but was not as reliable as the SNP panel (Pearson's

correlation coefficient = 0.877). This resulted in better estimates of pairwise relatedness

for parent-offspring using the SNP panel (Fig. 3). Overall, the SNP panel produced

better simulated estimates for each degree of relatedness (Fig. 4), greatly reducing the

variance around expected relatedness values (unrelated = 0, half-sib = 0.25, full-sib =

0.5, and parent-offspring = 0.5). This bolsters the confidence with which actual

relationships can be discerned when calculating pairwise relatedness of a population for

which there is little prior knowledge of social relationships.

Discussion

Several recent studies have rigorously investigated the use of SNPs in population

genetic studies for several non-model organisms (Morin *et al.* 2004; Slate *et al.* 2010;

Garvin *et al.* 2010; Heylar *et al.* 2011; Seeb *et al.* 2011), with growing support for the use

of SNPs in studies of parentage (Anderson & Garza 2006; e.g. Hauser *et al.* 2011; e.g.

Kaiser *et al.* 2016; e.g. Kess *et al.* 2016) and relatedness (e.g. Glaubitz *et al.* 2003;

Wang 2007). SNPs have proven to perform as well, if not better, than microsatellites in

these types of studies. To our knowledge this is the first study to describe a universal

ddRAD-seq method for use in parentage and relatedness analyses of wild populations of

birds. Our study is also the first to compare the efficiency of microsatellites versus SNPs

for determining genetic relationships in a species that is both socially complex, and

highly promiscuous. We show that SNPs developed from our modified ddRAD-seq

method are substantially more powerful than a moderate number of species-specific

microsatellite loci at assigning paternity and estimating relatedness among individuals.

Our method is highly attractive as an alternative to traditional microsatellite genotyping,

especially for systems where no microsatellites have been developed. This is largely

due to the combination of its cost and researcher time efficiency, the ease of this non-

species-specific method that combines the SNP discovery and screening steps, and the

large number of SNPs reliably recovered.

The total approximate materials cost for our ddRAD-seq analyses, including DNA

extraction, normalization of the DNA concentrations, library preparation, sequencing and

computational time was US$3,270.00 for 240 samples, or approximately $13.63 per

sample. The use of a homemade MagNA in place of commercial SPRI beads provides

significant savings. This cost is similar to that for genotyping 240 individuals at 12

microsatellite loci (in 3 multiplexed PCR mixes), in a situation where the labelled primers

have already been designed, purchased, and tested. However, a substantial additional

benefit of this ddRAD-seq method is that it does not require any locus discovery or

development before starting. The time required for library preparation, once DNA has

been extracted, is modest, and once the sequence data have been obtained, SNP

calling for the entire dataset can be performed in less than a day through a largely

automated bioinformatics pipeline. Unlike manually scoring peaks as in a traditional

microsatellite genotyping analysis, the identification of SNPs is less subjective and takes

far fewer hours of hands-on analysis (as most is performed computationally). The tools

for analyzing these ddRAD data are freely available and widely used (e.g., Stacks,

VCFtools).

For this study, our conditions and protocol allowed us to recover 411 high quality

SNP loci for 240 individual samples. However, we show that through simple variations in

the size selection window or the specificity of the restriction enzyme, more or fewer loci

can be obtained. For some applications, it could be advantageous to multiplex a greater

number of individuals and achieve similar coverage by aiming to recover fewer loci (e.g.,

using EcoRI rather than MspI). Alternatively, for applications where more loci are

required, the size selection window could be widened and concordantly the number of

individuals would have to be lowered.

The number of SNPs needed to perform robust parentage and relatedness

analyses depends on characteristics of the study population. Populations with reduced

genetic diversity will likely require a greater number of loci than those that are more

genetically diverse (Saunders *et al.* 2007; Strucken *et al.* 2016; Tortereau *et al.* 2017).

Obtaining more loci from the outset would aid in overcoming any issues relating to

population genetic diversity. Additionally, when studying species with complex social

systems, including for example both variable levels of genetic relatedness among

individuals and high rates of extra-pair fertilizations, it is imperative to obtain a sufficient

number of markers to discern genetic relationships robustly (Hughes 1998; Ross 2001;

Weinman *et al.* 2014). Our case study, using the variegated fairy-wren, shows that our

modified ddRAD-seq method recovers more than enough SNP loci to confidently discern

relationships in a species with a complex social system. Most parentage and

relatedness analysis programs are well equipped to handle large numbers of loci, so a

greater number of loci would not hinder analyses. Once an appropriate number of SNPs

are identified for performing robust analyses, conditions can be varied to maximize the

number of individuals to be genotyped.

For both paternity and relatedness analyses, our SNP panel far outperformed our

microsatellite panel by providing much more power and improving the overall confidence

for assignments. Variegated fairy-wrens are relatively easy to observe, and every nest

found can be assigned to a known mother by watching the female that builds the nest

and/or incubates the eggs. This level of knowledge may not be the norm for most study

systems, so we also investigated the CERVUS output for male-offspring relationships,

independent of known mothers. In doing so, the reliability of the SNP panel became

even more evident. In CERVUS, the higher the LOD score, the more likely that a given

male is the true father. Using SNPs, CERVUS typically output only a single male with a

positive LOD score, and the difference in LOD scores between the top-two ranked males

was dramatically different for SNP assignments (Fig. 2). When social information about

the known mother was excluded from the paternity analysis, the microsatellite panel

sometimes produced assignments that were ambiguous (two males had similar LOD

scores), and occasionally the wrong male was assigned paternity of the offspring. Under

the SNP panel, ambiguous assignments were nonexistent, and these cases were clearly

resolved (Fig. 1).

It is sometimes difficult to obtain appropriate demographic data to use in a formal

parentage analysis, and for many studies, this level of detail may not be necessary.

Population allele frequencies can be used to estimate pairwise relatedness for

individuals, and to reconstruct pedigrees using maximum likelihood-based methods.

Variance in estimates of pairwise relatedness (r) for known parent-offspring pairs was

dramatically reduced when using SNPs (Fig. 3). For our simulations, SNPs greatly

improved the differentiation between distributions for individuals of known degrees of

relatedness (Fig. 4). This is particularly important for systems with minimal demographic

and observational data, where these distributions can be used to determine familial

relationships between individuals, in conjunction with actual estimated r-values.

In summary, our ddRAD-seq method provides a cost effective and robust way to

identify SNPs for use in studies utilizing parentage and relatedness analyses. Our

experiment shows that a majority of the same SNPs can be obtained across groups,

using the same size selection windows and restriction enzymes. Future individuals can

be genotyped and incorporated to the analysis by re-running the Stacks pipeline. Using

a bird exhibiting great social complexity, and high promiscuity, we have shown that

SNPs identified by ddRAD-seq are more effective at assigning paternity and estimating

relatedness than highly polymorphic, species-specific microsatellite loci. This protocol

was designed to be universally applicable across bird species, and we have successfully

applied it in a range of other avian study systems (Table 4). While different numbers of

individuals were used in each study and therefore different numbers of loci were

recovered, in all cases paternity was confidently assigned to nestlings using CERVUS

(unpublished results). Applying this general protocol to many non-avian taxa may simply

require ensuring that the specific restriction enzymes and fragment size windows are

chosen appropriately.

Acknowledgements

We thank D. Baldassarre, K. Gielow, J. Welklin, and field technicians at Lake

Samsonvale who assisted with field efforts for this study. We are grateful to L. Stenzler

and S. Bogdanowicz for help with microsatellite discovery, development, and

genotyping. D. Baldassarre, E. Greig, & A. Dalziel provided valuable discussion on the

study design. This research was supported by the US National Science Foundation

(IOS-1353681 and DEB-1721662) .

References

Anderson EC, Garza JC (2006) The power of single-nucleotide polymorphisms for large-scale parentage inference. *Genetics*, **172**, 2567–2582.

Andrews KR, Good JM, Miller MR, Luikart G, Hohenlohe PA (2016) Harnessing the power of RADseq for ecological and evolutionary genomics. *Nature Reviews Genetics*, **17**, 81–92.

Avise JC, Jones AG, Walker D, Dewoody JA (2002) Genetic mating systems and reproductive natural histories of fishes: lessons for ecology and evolution. *Annual Review of Genetics*, **36**, 19–45.

Baird NA, Etter PD, Atwood TS *et al.* (2008) Rapid SNP discovery and genetic mapping using sequenced RAD markers. *PLoS ONE*, **3**, e3376.

Ball AD, Stapley J, Dawson DA *et al.* (2010) A comparison of SNPs and microsatellites as linkage mapping markers: lessons from the zebra finch (Taeniopygia guttata). *BMC Genomics*, **11**, 218.

Blouin MS (2003) DNA-based methods for pedigree reconstruction and kinship analysis in natural populations. *Trends in Ecology & Evolution*, **18**, 503–511.

Brumfield RT, Beerli P, Nickerson DA, Edwards SV (2003) The utility of single nucleotide polymorphisms in inferences of population history. *Trends in Ecology & Evolution*, **18**, 249–256.

Catchen J, Hohenlohe PA, Bassham S, Amores A, Cresko WA (2013) Stacks: an analysis tool set for population genomics. *Molecular Ecology*, **22**, 3124–3140.

Coates BS, Sumerford DV, Miller NJ *et al.* (2009) Comparative Performance of Single Nucleotide Polymorphism and Microsatellite Markers for Population Genetic Analysis. *Journal of Heredity*, **100**, 556–564.

Cramer ERA, Hall ML, De Kort SR, Lovette IJ, Vehrencamp SL (2011) Infrequent Extra-Pair Paternity in the Banded Wren, a Synchronously Breeding Tropical Passerine. *The Condor*, **113**, 637–645.

Danecek P, Auton A, Abecasis G *et al.* (2011) The variant call format and VCFtools. *Bioinformatics*, **27**, 2156–2158.

Davey JW, Blaxter ML (2010) RADSeq: next-generation population genetics. *Briefings in Functional Genomics*, **9**, 416–423.

Davey JW, Hohenlohe PA, Etter PD *et al.* (2011) Genome-wide genetic marker discovery and genotyping using next-generation sequencing. *Nature Reviews Genetics*, **12**, 499–510.

Decroocq V, Fave MG, Hagen L, Bordenave L, Decroocq S (2003) Development and transferability of apricot and grape EST microsatellite markers across taxa. *Theoretical and Applied Genetics*, **106**, 912–922.

Etter PD, Bassham S, Hohenlohe PA, Johnson EA, Cresko WA (2012) SNP discovery and genotyping for evolutionary genetics using RAD sequencing. In: *Molecular Methods for Evolutionary Genetics* Methods in Molecular Biology. pp. 157–178. Humana Press, Totowa, NJ.

Fernández ME, Goszczynski DE, Lirón JP *et al.* (2013) Comparison of the effectiveness

of microsatellites and SNP panels for genetic identification, traceability and assessment of parentage in an inbred Angus herd. *Genetics and molecular biology*, **36**, 185–191.

Galbusera P (2000) Cross-species amplification of microsatellite primers in passerine birds. *Conservation Genetics*, **1**, 163–168.

Garvin MR, Saitoh K, Gharrett AJ (2010) Application of single nucleotide polymorphisms to non-model species: a technical review. *Molecular Ecology Resources*, **10**, 915–934.

Glaubitz JC, Rhodes OE, Dewoody JA (2003) Prospects for inferring pairwise relationships with single nucleotide polymorphisms. *Molecular Ecology*, **12**, 1039–1047.

Griffith SC, Owens IPF, Thuman KA (2002) Extra pair paternity in birds: a review of interspecific variation and adaptive function. *Molecular Ecology*, **11**, 2195–2212.

Guichoux E, Lagache L, Wagner S *et al.* (2011) Current trends in microsatellite genotyping. *Molecular Ecology Resources*, **11**, 591–611.

Gut IG (2001) Automation in genotyping of single nucleotide polymorphisms. *Human Mutation*, **17**, 475–492.

Hauser L, Baird M, Hilborn R, Seeb LW, Seeb JE (2011) An empirical comparison of SNPs and microsatellites for parentage and kinship assignment in a wild sockeye salmon (Oncorhynchus nerka) population. *Molecular Ecology Resources*, **11**, 150–161.

Hedgecock D, Li G, Hubert S, Bucklin K (2004) Widespread null alleles and poor cross-species amplification of microsatellite DNA loci cloned from the Pacific oyster, Crassostrea gigas. *Journal of Shellfish Research*, **23**, 379–385.

Heylar SJ, Hemmer Hansen J, Bekkevold D *et al.* (2011) Application of SNPs for population genetics of nonmodel organisms: new opportunities and challenges. *Molecular Ecology Resources*, **11**, 123–136.

Hoffman JI, Amos W (2005) Microsatellite genotyping errors: detection approaches, common sources and consequences for paternal exclusion. *Molecular Ecology*, **14**, 599–612.

Hughes C (1998) Integrating molecular techniques with field methods in studies of social behavior: a revolution results. *Ecology*, **79**, 383–399.

Kaiser SA, Taylor SA, Chen N *et al.* (2016) A comparative assessment of SNP and microsatellite markers for assigning parentage in a socially monogamous bird. *Molecular Ecology Resources*, **17**, 183–193.

Kalinowski ST, Taper ML, Marshall TC (2007) Revising how the computer program CERVUS accommodates genotyping error increases success in paternity assignment. *Molecular Ecology*, **16**, 1099–1106.

Kearse M, Moir R, Wilson A *et al.* (2012) Geneious Basic: An integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics*, **28**, 1647–1649.

Kess T, Gross J, Harper, F, Boulding EG (2016) Low-cost ddRAD method of SNP discovery and genotyping applied to the periwinkle *Littorina saxatilis*. *Journal of*

Molluscan Studies, **82**, 104–109.

Li YC, Korol AB, Fahima T, Beiles A, Nevo E (2002) Microsatellites: genomic
distribution, putative functions and mutational mechanisms: a review. *Molecular*
Ecology, **11**, 2453–2465.

Lischer HEL, Excoffier L (2012) PGDSpider: an automated data conversion tool for
connecting population genetics and genomics programs. *Bioinformatics*, **28**, 298–
299.

Morin PA, Luikart G, Wayne RK (2004) SNPs in ecology, evolution and conservation.
Trends in Ecology & Evolution, **19**, 208–216.

Myers EM, Zamudio KR (2004) Multiple paternity in an aggregate breeding amphibian:
the effect of reproductive skew on estimates of male reproductive success.
Molecular Ecology, **13**, 1951–1963.

Nali RC, Zamudio KR, Prado CPA (2014) Microsatellite markers for Bokermannohyla
species (Anura, Hylidae) from the Brazilian Cerrado and Atlantic Forest domains.
Amphibia-Reptilia, **35**, 355–360.

Pemberton JM, Slate J, Bancroft DR, Barrett JA (1995) Nonamplifying alleles at
microsatellite loci: a caution for parentage and population studies. *Molecular*
Ecology, **4**, 249–252.

Peterson BK, Weber JN, Kay EH, Fisher HS, Hoekstra HE (2012) Double digest
RADseq: An inexpensive method for de novo SNP discovery and genotyping in
model and non-model species (L Orlando, Ed,). *PLoS ONE*, **7**, e37135.

Pew J, Muir PH, Wang J, Frasier TR (2015) related: an R package for analysing pairwise
relatedness from codominant molecular markers. *Molecular Ecology Resources*, **15**,
557–561.

Primmer CR, N Painter J, T Koskinen M, U Palo J, Merilä J (2005) Factors affecting
avian cross-species microsatellite amplification. *Journal of Avian Biology*, **36**, 348–
360.

Puritz JB, Matz MV, Toonen RJ *et al.* (2014) Demystifying the RAD fad. *Molecular*
Ecology, **23**, 5937–5942.

Queller DC, Strassmann JE, Hughes CR (1993) Microsatellites and kinship. *Trends in*
Ecology & Evolution, **8**, 285–288.

R Core Team (2016) R: A Language and Environment for Statistical Computing. R
Foundation for Statistical Computing, Vienna, Austria. http:// www.R-project.org/.

Rohland N, Reich D (2012) Cost-effective, high-throughput DNA sequencing libraries for
multiplexed target capture. *Genome Research*, **22**, 939–946.

Ross KG (2001) Molecular ecology of social behaviour: analyses of breeding systems
and genetic structure. *Molecular Ecology*, **10**, 265–284.

Rowley I, Russell EM (1997) *Fairy-wrens and Grasswrens: Maluridae*. Oxford University
Press.

Saunders IW, Brohede J, Hannan GN (2007) Estimating genotyping error rates from
Mendelian errors in SNP array genotypes and their impact on inference. *Genomics*,
90, 291–296.

Schodde, R (1982). *The fairy-wrens* (ed. Bass T). The Craftsman Press. Victoria,

Australia.

Seeb JE, Carvalho G, Hauser L *et al.* (2011) Single-nucleotide polymorphism (SNP)
discovery and applications of SNP genotyping in nonmodel organisms. *Molecular*
Ecology Resources, **11**, 1–8.

Selkoe KA, Toonen RJ (2006) Microsatellites for ecologists: a practical guide to using
and evaluating microsatellite markers. *Ecology letters*, **9**, 615–629.

Slate J, Gratten J, Beraldi D *et al.* (2010) Gene mapping in the wild with SNPs:
guidelines and future directions. *Genetica*, **136**, 97–107.

Solomon NG, Keane B, Knoch LR (2004) Multiple paternity in socially monogamous
prairie voles (Microtus ochrogaster). *Canadian Journal of Zoology*, **82**, 1667–1671.

Strucken EM, Lee SH, Lee HK *et al.* (2016) How many markers are enough? Factors
influencing parentage testing in different livestock populations. *Journal of Animal*
Breeding and Genetics, **133**, 13–23.

Syvänen A-C (2001) Accessing genetic variation: genotyping single nucleotide
polymorphisms. *Nature Reviews Genetics*, **2**, 930–942.

Tokarska M, Marshall T, Kowalczyk R *et al.* (2009) Effectiveness of microsatellite and
SNP markers for parentage and identity analysis in species with low genetic
diversity: the case of European bison. *Heredity*, **103**, 326–332.

Tortereau F, Moreno CR, Tosser-Klopp G, Servin B, Raoul J (2017) Development of a
SNP panel dedicated to parentage assignment in French sheep populations. *BMC*
Genetics, **18**, 50.

Wang J (2002) An estimator for pairwise relatedness using molecular markers. *Genetics*,
160, 1203–1215.

Wang J (2007) Triadic IBD coefficients and applications to estimating pairwise
relatedness. *Genetical research*, **89**, 135–153.

Webster MS, Reichart L (2005) Use of microsatellites for parentage and kinship
analyses in animals. *Methods in enzymology*, **395**, 222–238.

Weinman LR, Solomon JW, Rubenstein DR (2014) A comparison of single nucleotide
polymorphism and microsatellite markers for analysis of parentage and kinship in a
cooperatively breeding bird. *Molecular Ecology Resources*, **15**, 502–511.

Westneat DF, Sherman PW, Morton ML (1990) The ecology and evolution of extra-pair
copulations in birds. *Current Ornithology*, **7**, 331–370.

White PS, Densmore LD (1992) Mitochondrial DNA isolation. In: *Molecular Genetic*
Analysis of Populations: A Practical Approach (ed. Hoelzel AR), pp. 50–51. Oxford
University Press, New York, NY.

Willing E-M, Hoffmann M, Klein JD, Weigel D, Dreyer C (2011) Paired-end RAD-seq for
de novo assembly and marker design without available reference. *Bioinformatics*,
27, 2187–2193.

Author Contributions

D.J.T designed the study, collected field data, performed microsatellite development and analysis, conducted parentage and relatedness analyses, and drafted the manuscript with help from all co-authors. B.G.B and L.C. designed the study, and performed SNP discovery and analysis. M.S.W and I.J.L. helped design the study, and secured funding.

741 **Tables**

742 Table 1: Experimental design.

# of samples (index groups)	enzymes	size selection interval
160 samples (8 index groups, index 1-8)	SbfI - MspI	450 - 600 bp
20 samples (Index 9)	SbfI - MspI	450 - 600 bp (replicate of above)
20 samples (index 10)	SbfI - MspI	400 - 700 bp (wide size selection)
20 samples (index 11)	SbfI - EcoRI	450 - 600 bp (infrequent 3' cutter)
20 samples (index 12)	SbfI - EcoRI	450 - 600 bp (replicate of above)

748 Table 2. Marker characteristics

Marker Panel	Number of loci	Mean proportion loci typed	Mean alleles per locus	Mean H_e	Mean H_o	Mean PIC	Nonexclusion probability (first parent)	Nonexclusion probability (second parent)	Nonexclusion probability (parent pair)
Microsatellites	12	0.99	14.17	0.77	0.76	0.74	1.9×10^{-4}	1.9×10^{-6}	1.5×10^{-10}
SNPs	411	0.98	2.00*	0.45	0.45	0.35	5.2×10^{-20}	6.9×10^{-35}	1.0×10^{-55}

749 *Only biallelic SNPs were retained. If a locus had 3 alleles across the population, it was filtered from the dataset.

Table 3. Overlap in the RAD loci that were obtained while varying different steps of the
protocol (size selection and restriction enzymes). The diagonal indicates the total
number of loci recovered for each treatment. Values above the diagonal represent the
percent overlapping loci between groups (relative to the group with the smallest number
of loci), while values below the diagonal list the number of loci that were overlapping
between groups.

	All	A	B	C	D	E
All	411	85.4	78.1	79.1	1.2	1.0
A	351	797	85.1	75.8	0.8	0.8
B	321	549	645	83.4	2.0	2.3
C	325	604	538	1440	4.2	3.9
D	5	15	12	25	596	68.4
E	4	13	12	20	353	516

All: 160 samples; Sbf1/Msp1; 450-600 bp.
A: 20 samples; Sbf1/Msp1; 450-600 bp.
B: 20 samples; Sbf1/Msp1; 450-600 bp.
C: 20 samples; Sbf1/Msp1; 400-700 bp.
D: 20 samples; SbfI/EcoRI; 450-600 bp.
E: 20 samples; SbfI/EcoRI; 450-600 bp.

776
777 Table 4: Summary information for ddRAD-seq studies performed to investigate
778 parentage in other bird species.

Species	Scientific name	Number of individuals	Number of loci	Number of loci in HWE
Variegated fairy-wren	*Malurus lamberti*	160	552	411
Hispaniolan woodpecker*	*Melanerpes striatus*	288	179	135
Northern Red-billed hornbill	*Tockus erythrorhynchus*	40	475	414
Von der Decken's hornbill	*Tockus deckeni*	112	490	410
Sapayoa	*Sapayoa aenigma*	6	672	671
Red-backed fairy-wren**	*Malurus melanocephalus*	240	483	233
		240	291	174

779 * Two independent ddRAD-seq experiments were performed - one with 240 samples
780 and the other with 48. After quality filtering and demultiplexing the data from 288
781 samples was combined for denovo assembly and SNP identification.
782 ** Two independent ddRAD-seq experiments were run on each set of 240 samples.
783 There is 74% overlap in the loci identified in each experiment (of the loci in HWE).
784
785

Figures
Figure 1. Resolved paternity assignments for 5 nestlings with ambiguous assignments
under the microsatellite panel, but not with the SNP panel. Each panel in the graph
represents an individual offspring, and the two top-ranked males are depicted as a
triangle and a circle, respectively. Lines connecting like shapes show the change in LOD
score for each male, using each marker type (microsatellites versus SNPs). Note that y-
axis scale varies among panels in the graph.

794   Figure 2. Difference in CERVUS LOD scores (delta LOD) between the most likely father
795   of a nestling and the second possible father in the population, for both marker panels.

813   Figure 3. Box plot of pairwise relatedness values for all parent-offspring (40 mother-
814   offspring and 40 father-offspring) relationships, using population allele frequencies from
815   each marker panel.
816

Combined pairwise r-values for all parent-offspring pairs

Figure 4. Density plots of relatedness values for simulated pairs of known relatedness
(unrelated, half-sibling, full-sibling, and parent-offspring) using population allele
frequencies from each marker panel (a. MSAT; b. SNP). Overlap in distributions
indicates the overlap between relatedness value estimators for pairs of individuals of
different relationships. The spread of each distribution indicates the reliability of
observed relatedness values based on their deviation from expected relatedness values
(Unrelated = 0, Half-sib = 0.25, Full-sib = 0.5, and Parent-offspring (P-O) = 0.5, denoted
by vertical dashed lines).
840

841