1   **Insects and incest: sib-mating tolerance in natural populations of a parasitoid wasp**

Marie Collet[1*], Isabelle Amat, Sandrine Sauzet[1], Alexandra Auguste[2], Xavier Fauvergue[2], Laurence

Mouton[1], Emmanuel Desouhant[1]

[1] Univ Lyon, Université Lyon 1, CNRS, Laboratoire de Biométrie et Biologie Evolutive UMR5558, F-69622

Villeurbanne, France

2   [2] INRA, CNRS, Université Côte d'Azur, ISA, France

3   *Corresponding author: marie.collet@univ-lyon1.fr

Abstract

1. Sib-mating avoidance is a pervasive behaviour, which likely evolves in species that are

subject to inbreeding depression. Laboratory studies have provided elegant

demonstrations, but small-scale bioassays often minimize the costs associated with mate

finding and mate-choice and may for this reason produce spurious findings.

2. We inferred the mating behaviour of the parasitoid wasp *Venturia canescens* from lab

experiment and genetic analyses of natural populations. We used *V. canescens* as a model

organism because in this species laboratory experiments have shown that sib-mating

yields a 25% decrease in fertile offspring, and as a consequence, sib-mating is partially

avoided.

3. Our study consisted of a mate choice experiment in laboratory cages and a field study

based on the genotyping of 86 wild-caught males, 155 wild-caught females and their 226

daughters at eighteen microsatellite loci. With these field data, we reconstructed the

genotype of each female's mate and estimated the relatedness of each mating pair.

4. The mate choice experiment confirmed that kin discrimination occurs in this species.

Time to mating depended on the frequency of female encounters with related and

unrelated males. However, contrary to previously published results, no sib-mating

avoidance was detected during these experiments. In the field, we found that the effective

rate of sib-mating did not differ from the probability that sibs encounter at random, which

corroborates the absence of sib-mating avoidance. We also detected a weak but

significant male bias in dispersal, a pattern which could reduce encounters between sibs.

5. These results suggest that despite kin discrimination, *Venturia canescens* tolerates sib-

mating in the field. The weak male-biased dispersal cannot explain entirely this pattern.

The observed sib-mating tolerance raises the question as to why kin discrimination is

maintained in this species. It also calls into question common beliefs on inbreeding

depression in species with single-locus complementary sex determination.

Key-words: Field study**,** Inbreeding tolerance, Kin recognition, Parasitic wasp, single-locus

Complementary Sex Determination, sl-CSD.

Introduction

When inbreeding lowers fitness-related components such as survival and fertility, natural

selection should favour behaviours preventing the reproduction of genetically-related individuals, or

mitigating harmful consequences referred to as inbreeding depression (Pusey and Wolf, 1996; Keller

and Waller, 2002; Angeloni et al., 2011). Inbreeding depression is caused by the expression of

deleterious recessive alleles at the homozygous state or by a loss of heterosis in inbred crosses

(Charlesworth and Charlesworth, 1987; Pusey and Wolf, 1996). Selection on behaviours underlying

inbreeding avoidance should thus depend on inbreeding load, which balances the advantages of

inbreeding avoidance with the costs of implementing adapted physiological or behavioural responses.

Selection may also depend on the benefits of inbreeding in terms of inclusive fitness, as inbreeding

results in an increased representation of genes identical by descent in the offspring (Kokko and Ots,

2006; Puurtinen, 2011; Duthie and Reid, 2016).

Responses to inbreeding risk reflect the diversity of benefits and costs associated with

inbreeding and inbreeding avoidance. Observations range from systematic inbreeding avoidance to

inbreeding tolerance, or even, inbreeding preference (Szulkin et al., 2013). For example, inbreeding

preference is observed in the cichlid fish *Pelvicachromis taeniatus* ; in this species, inbreeding does not

lower the survival and growth rate of the youngs, and related pairs invest more in parental care and

cooperate better (Thünken et al., 2007). In rodents such as *Marmota flavipentris*, inbreeding

preference was shown despite the lower survival of inbred progeny. This preference can be explained

by the large variation of male reproductive success and the composition of social groups, where males

mating with related females do not suffer decreased annual reproductive success compared to the

majority of males, that achieve little or no reproductive success (Olson et al., 2012). In contrast, ring-

tailed lemurs, or mole rats living in socially structured groups, avoid inbreeding. In mole rats, avoidance

leads to a high reproductive skew in males, with a subset of males that never access to reproduction

(Cooney and Bennett, 2000). In ring-tailed lemurs, inbreeding avoidance allows females to save energy

(pregnancy and lactation) that would be lost for the production of inbred offspring suffering from

depressed immune functions and a reduced life expectancy (Boulet et al., 2009; Charpentier et al.,

2008). Hence, quantifying the costs and benefits of inbreeding avoidance and the balance between

them is necessary to decipher the selective pressures acting on inbreeding avoidance, including the

consequences of inbreeding, and in turn, to understand the evolution of this behaviour.

Additionally, the balance between benefits and costs associated with inbreeding avoidance

depends on environmental factors, such as population density, spatial or social population structure.

A low population density constrains mate availability and consequent encounter rate, and may

therefore influence mate choice (Kokko and Rankin, 2006; Duthie and Reid, 2016). In the lizard *Zootoca*

vivipara, female choosiness is reduced when mate encounter rate decreases (Breedveld and Fitze,

2015). Similarly, genetically-related partners are less avoided in small and isolated populations of

moose, with an amplified effect when the sex-ratio is female-biased (Herfindal et al., 2014). In the

marsupial carnivore *Antechinus agilis*, habitat selectivity following natal dispersal is negatively

correlated with the abundance and relatedness of females occupying habitats, which suggests a

pervasive effect of inbreeding risk on dispersal (Banks and Lindenmayer, 2014). In cooperative

breeding species, inbreeding avoidance strategies can lead to the suppression of reproduction outside

dominant pairs due to the lack of opportunity of outbred matings (as in the Damaraland mole-rats

Fukomys damarensis; (Cooney and Bennett, 2000). Therefore, the reproductively supressed

subordinates that are related to the dominant pair should increase their involvement in help to

increase their own inclusive fitness and maintain the social structure of the group (Nichols, 2017). This

interplay between environment and inbreeding avoidance patterns requires field data to be able to

quantify cost and benefits of each strategy.

Various behavioural and physiological strategies to avoid inbreeding risks have been selected

for in animals, including sib-mating avoidance, dispersal (Greenwood et al., 1978; Szulkin and Sheldon,

2008), polyandry, extra-pair paternity, divorce (Hatchwell et al., 2000; Cornell and Tregenza, 2007;

Cohas et al., 2008; Lardy et al., 2011; Reid et al., 2015; Duthie and Reid, 2016), and postcopulatory

mechanisms, such as the preferential use of sperm from unrelated males by inseminated females

(Tregenza and Wedell, 2002; Bretman et al., 2004). Mate choice is probably the most pervasive

behaviour for sib-mating avoidance (Pusey and Wolf, 1996) and relies mostly on sib recognition

mediated by chemical communication (Howard and Blomquist, 2005; Johansson and Jones, 2007;

Charpentier et al., 2008a; Bonadonna and Sanz-Aguilar, 2012). It is nonetheless important to note that

kin recognition mechanisms can arise from selective forces other than inbreeding avoidance, such as

nest-mate recognition, parental care, territory defence, dominance hierarchies to name a few (Mateo,

2004).

Assessing inbreeding avoidance patterns, that is, the occurrence of inbreeding avoidance and

the behavioural strategies implied, is a difficult task that requires the estimation of relatedness

coefficients between actual and potential mates (Szulkin et al., 2013). This may explain why most field

studies have been conducted on large species of mammals and birds, for which monitoring is much

easier compared to small invertebrates (Cohas et al., 2008; Herfindal et al., 2014; Arct et al., 2015;

Hardouin et al., 2015). It is therefore not surprising that inbreeding avoidance patterns have been

rarely documented in insects in the wild (but see Robinson et al., 2012 for a field assessment of optimal

inbreeding in *Drosophila melanogaster*). Laboratory studies nonetheless suggest various strategies of

inbreeding avoidance in different insect genus, from active choices such as precopulatory avoidance

of related males, which represent low quality partners with a high genetic load, in the butterfly *Bicyclus*

anynana (Fischer et al., 2015) or postcopulatory choice with the preferential use of the sperm stored

from unrelated males in *Gryllus bimaculatus* (Bretman et al., 2004, 2009). In a social species of termite,

Neotermes chilensis, the dispersal of individuals founding new colonies is the main mechanism to avoid

inbreeding risks (Aguilera-Olivares et al., 2015). Absence of inbreeding avoidance has also been

evidenced in insects such as parasitoid wasps (Bourdais and Hance, 2009) or *Drosophila melanogaster*

(Tan et al., 2012). The diversity of strategies unveiled in the laboratory renders more striking our lack

of knowledge on insect inbreeding avoidance patterns *in natura*, and calls for field studies.

Here, we take up this challenge by studying, both in the laboratory and in the field, a parasitoid

wasp with a simple form of inbreeding depression. Many species of the order Hymenoptera have

single-locus Complementary Sex Determination (sl-CSD) whereby sex is determined by both ploidy and

heterozygosity at a unique sex determination locus (Cook, 1993; van Wilgenburg et al., 2006; Heimpel

and de Boer, 2008; Asplen et al., 2009; Schmieder et al., 2012). In haplodiploid species, males develop

from unfertilized haploid eggs whereas females develop from fertilized diploid eggs. Inbreeding

depression arises for diploids that are homozygous at the sex locus: instead of developing normally

into females, they turn into abnormal diploid males that are generally unviable or sterile (Cook, 1993;

van Wilgenburg et al., 2006; Fauvergue et al., 2015). Whatever the diversity of sex alleles in the

population, full sibs have 50% chance of sharing a common sex allele and, if they mate, half of their

diploid offspring will develop into unfit males (Cook, 1993). Hence, assuming females fertilize half of

their eggs, sib-mating results in 12.5% fewer offspring, on average. sl-CSD is thus a form of inbreeding

depression underlined by overdominance, with no deleterious alleles at the sex locus but a strong

heterozygous advantage (e.g., Charlesworth and Willis, 2009). The null fitness of diploid males should

favour sib-mating avoidance.

The parasitoid wasp *Venturia canescens* Gravenhost (Hymenoptera: Ichneumonidae) has a

single locus complementary sex determination (first demonstrated by Beukeboom, 2001) and

inbreeding reduces the fitness of both males and females via the production of sterile diploid offspring

and a negative impact on egg load and hatching rate (Vayssade et al., 2014; Chuine et al., 2015).

Consistently, in no-choice bioassays, mating success decreases with increasing genetic relatedness

between mates (Metzger et al., 2010a; Chuine, 2014). Circumstantial evidence suggests that females

are the choosy sex, which makes sense in a species with a monoandrous/polygenous mating system

(Metzger et al., 2010a). However, when confronting females to choice between a brother and an

unrelated male, inbreeding avoidance vanishes, likely because the mixture of volatiles in small test

tubes confounds females (Metzger et al., 2010a).

In this study, we address the question of inbreeding avoidance by combining field and

laboratory approaches. We first tested the effects of male density and average relatedness on

inbreeding avoidance with more realistic conditions than before (Metzger et al., 2010a). We expected

sib-mating avoidance to decrease with decreasing density and increasing relatedness. We further

implemented a population genetic approach based on microsatellite genotyping in order to assess

inbreeding avoidance and underlying behaviours in field populations of *V. canescens*. For this, we

sampled two natural populations of *V. canescens* and compared the observed rate of sib-mating to the

probability of random encounter among genetically-related individuals. Under the hypothesis of sib-

mating avoidance, we expected the rate of sib-mating to be smaller than the probability that relatives

encounter at random. Besides, we used the field data to test whether sex-biased dispersal limits

encounters among sibs in the field. We show that sib-mating tolerance occurs in the wild, and propose

an evolutionary scenario to explain why this result contrasts with earlier findings of inbreeding

avoidance in this species as well as prior expectations about inbreeding depression in species with sl-

CSD (van Wilgenburg et al., 2006).

Materials and Methods

Biological model

Venturia canescens is a monoandrous solitary endoparasitoid found in most circum-

Mediterranean areas (Salt, 1976). Females lay eggs in a large range of hosts found on various fruits

(Salt, 1976; Driessen and Bernstein, 1999). Knowledge of the mating system of *V. canescens* under

natural conditions is limited but can partly be inferred from lab results. Females do not discriminate

between host patches with more than four host larvae, suggesting that, in the field, they rarely

encounter aggregated host (Driessen and Bernstein, 1999). Therefore, it is unlikely that this species

experiences local mate competition. Males search for females via a synergistic combination of volatile

sex pheromones emitted by females and kairomones from hosts (Metzger et al., 2010b). Both

immature development and adult life last around three weeks (at 25°C), and females lay eggs during

most of their adult life (Metzger et al., 2008). This leads to overlapping generations. Depending on

studies, sex-ratio is considered balanced (Beukeboom, 2001) or weakly biased toward females

(Metzger et al., 2008). Balanced sex ratio is congruent with the absence of aggregation (Driessen and

Bernstein, 1999), the absence of local mate competition, and mate-finding via volatile cues. The

proportion of diploid males corresponds to that expected under the hypothesis of a balanced sex ratio

(Fauvergue et al., 2015). Field and laboratory studies showed that adult females are good dispersers,

with a flight velocity estimated at 0.2 $m.s^{-1}$ (Schneider et al., 2002; Desouhant et al 2003; Amat et al.,

2012).

Insect sampling in the field

Adults *V. canescens* were captured from June to September 2014, across 10 non-consecutive

days. Two field sites located 300 km apart were sampled: Valence (N 44°58'21" E 4°55'39", composed

of fruit trees in an orchard) and Nice (N 43°41'23" E 7°18'6", composed of carob trees in a semi-natural

park). We captured only females in Valence and wasps of both sexes in Nice.

Two different types of traps were used: kairomone-baited traps for the capture of live females,

and yellow sticky traps for males (which were generally found dead). Female traps constituted of an

open cup containing host larvae, *Ephestia kuehniella* Zeller (Lepidoptera: Pyralidae), along with used

rearing medium (semolina). Saliva secreted by host larvae when feeding serves as a kairomone for

female *V. canescens* (Corbet, 1971)*.* Traps were hung in trees (Metzger et al., 2008) and visited every

20 minutes to capture attracted females. For males, 125 mm × 200 mm yellow sticky cards were baited

with extracts from females and hosts, a method that has already proved successful (Collet et al., 2016).

For each trap, a vial containing the extracts was attached at the centre of the card. As for females,

traps were hanged within host trees, and all the *V. canescens* males that were found stuck on the traps

were collected and conserved individually in 96% ethanol to preserve DNA. We captured 77 females

and 86 males in Nice and 78 females in Valence (see Results and Table 1).

Insect rearing

The wild females trapped were brought back to the laboratory where they were handled in a

climatic chamber under constant conditions (24±1°C, 60±10% RH, LD 16:8) and fed with 50% water-

diluted honey. Each female was allowed to lay eggs individually in *E. kuehniella* hosts for three days.

As we could not control for the age and previous egg-laying experience of captured females, the

number of offspring was highly variable (1-10 daughters and 1-7 sons). As sexual (arrhenotokous) and

asexual (thelytokous) strains coexist in *V. canescens* (Beukeboom et al., 1999; Schneider et al., 2002),

we used the presence of males among offspring as evidence for arrhenotoky, and thelytokous

individuals were discarded. Mothers and their offspring were then preserved individually in 96%

ethanol.

For laboratory experiments, we used a standardized laboratory rearing. The wasp culture had

been established with about 60 females collected in the field near Valence (same location as the field

capture, N 44°58'21" E 4°55'39", INRA, Gotheron), southern France, on several occasions during the

summer 2015. *Ephestia kuehniella* was reared in the laboratory on organic wheat semolina medium

for three weeks before being exposed to parasitoids (eggs were obtained from the Biotop rearing

facility located in Livron sur Drôme, France). Parasitoid development took place in a controlled

environment (24±1°C; 60±5% RH; LD regime 12 h:12 h).

Experiments on female mate choice required related and unrelated males. We thus initiated

families by collecting females randomly from the mass rearing (from different rearing boxes), and

maintaining them individually in a little circular box (∅: 8 cm, height: 3 cm) with meshed lid with two

males and about 30 three-weeks host larvae on 1 cm of organic wheat semolina. The wasps were fed

ad libitum with water-diluted honey (50-50) and allowed to lay eggs during their whole life. About

three weeks after, the parents were removed from the box if still alive, and their offspring were

collected at emergence, before reproduction. Females (the daughters) were kept individually in plastic

vials with a drop of water-diluted honey, and males were kept with their brothers and fed similarly. All

individuals used in the experiments were two-days old.

Effect of male density and sib frequency on sib-mating avoidance in the laboratory

We aimed at untangling the effects of male density and the frequency of encounters with

related males on female mate choice. We thus designed a factorial experiment with two densities of

males (high density, noted *D*, with nine males, and low density, noted *d*, with three males) and two

frequencies of related males (high frequency, noted *F*, with 2/3 of brothers, and low frequency, noted

f, with 1/3 of brothers). Males were marked according to kinship (brother *vs* unrelated) with a drop of

paint applied on their thorax 24h before being tested. The two colours used do not modify behaviours

(data not shown, and Desouhant et al., 2003) but were nonetheless alternated randomly across trials.

Each trial took place as follows: a virgin female was released in a mesh cage ($30 \times 30 \times 30$cm) containing

the males at 25°C, 65% RH and with a constant luminosity. Courtship and mating behaviours were

monitored for 30 min or until a mating had occur between 10 am and 4 pm. We recorded mating

latency, copulation duration, relatedness (colour) of the successful male, and courtship effort (*i.e.* the

time and the duration of each male courtship behaviour) (van Santen and Schneider 2002). For each

courtship, we identified the relatedness (colour) of the candidate. Therefore, all courtships were

counted independently of mating success. Twenty trials were performed per factor level, excepted for

d-F (18 replicates).

We analysed the proportion of sib-mating with a GEE-GLM (Generalized Estimating Equation –

Generalized Linear Model; Liang and Zeger, 1986, GEEPACK package in R software) implemented with a

binomial distribution of error, a logit link function, an exchangeable correlation structure, and a fixed

scale. The model was fitted with mate relatedness as a response variable (binomial: 0 if the female

mated with her brother, 1 if she mated with an unrelated male), density and frequency of brothers as

main effects and in interaction as explanatory variables, and families as a random factor (*i.e.* clusters,

Højsgaard et al., 2006). We added an offset term to take into account the theoretical mating

proportion with related males (*i.e.* 1/3 in low frequency *f* and 2/3 in high frequency *F*).

We also tested the effects of density and proportion of brothers on male courtship effort and latency

before copulation. For courtship effort, we implemented a GLM based on a binomial distribution and

a logit link function, with the percentage of courtships done by brothers as a response variable and the

density and proportion of brothers as explanatory variables. For latency before copulation, we

performed a Cox model with the same explanatory variables as fixed effects and family as a random

effect (COXME package, Therneau, 2015 in R software). We also included the number and timing of

courtships (from a related or unrelated male) as a time-dependent variable. If females assess their

relatedness with males during courtship, an increasing number of encounters with related males could

lead the female to decrease its selectivity and, as a consequence, the latency before mating. In the

other hand, an increasing number of encounters with unrelated males should not affect its selectivity

and therefore its latency to mate.

Genotyping

Genotyping was performed at 18 microsatellite loci developed by Collet et al. (2016) and

Mateo Leach et al. (2012) (See SEM-A for details on the genotyping and SEM-D for details on the

number of microsatellites used). We genotyped all field-captured males and females (hereafter

referred to as wild females) and up to three daughters per female (see SEM-B for a justification of the

number of offspring genotyped). When fewer than three daughters emerged (ca. 34%, in which 40%

produced only one daughter), all were genotyped.

We estimated polymorphism and linkage disequilibrium at each microsatellite locus for the

wild females of Nice and Valence with the GENEPOP software version 4.3 (Raymond and Rousset, 1995).

The frequency of null alleles was estimated with FREE NA software (Chapuis and Estoup, 2007).

Population differentiation was estimated based on Wright's F estimator F_{st} with the FSTAT software

version 2.9.3 (Goudet, 1995). Effective population sizes were estimated using the full likelihood

method, assuming random mating, developed by Wang (2009), and implemented in the software

COLONY (Jones and Wang, 2010). As effective population sizes were small (< 80), we did not check for

Hardy-Weinberg equilibrium. We also sought for diploid males as a proxy of consanguinity and of the

number of alleles at *csd* locus. A male was considered diploid if at least one microsatellite locus was

heterozygous, haploid otherwise (Collet et al., 2016).

Inference of relatedness among mates

• Male genotypes

We inferred the genotype of wild females' mates based on the genotypes of wild females and

that of their daughters. For this, we used the software COLONY, which reconstructs families based on

haplodiploid genotypes (Jones and Wang, 2010). When alternative genotypes were proposed for a

given male, we selected the one that was compatible with the genotype of the daughter and the

mother. When several alternatives were possible (e.g. if mother and daughters were heterozygous

with the same alleles at a given locus), it was impossible to assign the right allele to the father with

certitude. In this case, we assigned the allele with the highest probability of occurrence (the probability

is given by COLONY). Doing so, we successfully inferred the genotype of 51 mates for the 54 wild

females captured (see Results and Table 1).

272   • Relatedness between mates

273   To assess if *V. canescens* females avoid mating with their relatives in the field, we compared

the observed number of wild females mated with a related male to the number expected under the

assumption of random mating. For this, we determined a "relatedness threshold" above which

individuals were considered genetically related. To determine this threshold, we simulated

populations with similar allele frequencies at microsatellite loci to that of natural populations, and a

balanced sex ratio (see SEM-C for details on simulations). Simulations allowed to keep track of

pedigrees, yielding a perfect knowledge of kinship for all potential mates, which we classified in three

categories: full-sibs, half-sibs, or unrelated. Using microsatellite genotypes from the same simulated

dataset, we estimated the relatedness coefficient (r) for all female-male pairs (TrioML coefficient,

software COANCESTRY; Wang, 2011). We then used these estimated relatedness coefficients with

different relatedness thresholds to assign each dyad to related vs unrelated categories (SEM-C). By

comparing these posterior assignations to the pedigrees known a priori, we estimated the relatedness

threshold that minimized the number of wrong assignations (*i.e.* unrelated pairs assigned related

based on the threshold, or vice-versa) under random encounter hypothesis. We found r_{crit} = 0.16 (SEM-

D, Fig. S2). Logically, this threshold is lower than the theoretical relatedness between a brother and a

sister ($r_{\text{sister-brother}}$ = 0.25; $r_{\text{brother-sister}}$ = 0.5) and in the interval of relatedness for 1st degree cousins (r =

0.125 or r = 0.1875 depending on the cross). With this threshold, we expect to wrongly assign 11.4%

of the relationships.

We applied this r_{crit} = 0.16 threshold to field data. First, we estimated the number of wild

females that had been inseminated by a related male. Second, we generated a theoretical distribution

of the number of couples with a r ≥ 0.16 under the null hypothesis of random mating. For this, each of

the 51 wild females from Nice, for which a mate genotype had been successfully inferred, was paired

randomly with one of the 79 genotyped males, and we counted the number of pairs with r ≥ 0.16. This

calculation was repeated 1000 times to produce the null distribution. We finally compared the

observed number of related pairs with the null distribution via a permutation test implemented in the

function AS.RANDTEST from the ADE4 package in R software version 3.2.3 (Chessel et al., 2004; R Core

Team, 2015). We also estimated the probability that a female encountered a related male in the field

the day of her capture. Indeed, partners present in the same day in the same area have a high chance

to encounter each other and mate due to the high dispersal capacity of this species.

Sex-biased dispersal in the field

Sex-biased dispersal can decrease the burden of inbreeding depression by reducing encounters

and subsequent mating among sibs. We thus assessed the dispersal ability of males and females from

our genotyping data in the population of Nice, where two trapping locations were distinguishable

(Figure 1). The first patch contained a group of 21 trees and the second patch 15 trees. Trees were

located a few meters from one another within patches, and the two patches were about 200 m away.

Six sampled trees further apart were not included in this scheme (black dots on Figure 1).

We estimated the relatedness between individuals of each sex within and between patches

using COANCESTRY software (Wang, 2011). Sex-biased dispersal should result in higher coefficients of

relatedness within patches than between patches for the sex dispersing at smaller distances (Prugnolle

and de Meeus, 2002). Coefficients of relatedness inter-patch and intra-patch were compared via a

Kruskal-Wallis test (with patch as an explanatory factor) for each sex separately. When significant,

post-hoc (with Bonferroni corrections) Mann-Whitney's *U*-tests between the three modalities (*i.e.*

relatedness within patch 1, relatedness within patch 2, and relatedness between patches) were

performed.

Results

Effect of male density and sib frequency on sib-mating avoidance in the laboratory

Mating success was higher at high male density (*D;* 81.6%, 40 matings/49 trials) than at low

density (*d*; 44.1%; 38/86) (Table 3; proportion test: χ^2 = 16.44, *df* = 1, *p* = 5.02 .10^{-5}). The observed

proportion of sib-mating did not depart from that expected under random encounters (GEE-GLM

model, *p*-value of the intercept = 0.3). This was confirmed by a positive relationship between the

proportion of related males and the probability of sib-mating (Wald statistic table, χ^2 = 15.36, *df* = 1,

p = 8.9 .10^{-5}). Male density had no effect on mate choice, neither as a main effect (χ^2 = 0.92, *df* = 1,

p = 0.34) nor in interaction with the proportion of related males (χ^2 = 0, *df* = 1, *p* = 1). Males did not

differ in their courtship effort, as the proportion of courtships observed for unrelated males and

brothers did not depart from the expected values (GLM model, *p*-value of the Intercept = 0.16): for

example, in the *D-F* modality with 6 brothers and 3 unrelated males, we expected that 2/3 of the

courtships should be done by brothers (i.e. 13.3 on 20 matings) and observed 12 sib-mating. Time

before copulation decreased with increases in male density and increases in the number of sib

rejections (Table 4), but it was not affected by the proportion of related males or by the number of

rejections of unrelated males (Table 4).

Genotyping

A total of 241 wasps were captured in Valence and in Nice. In Valence, 78 wild females were

captured (Table 1) among which 35 produced a total of 86 daughters (Table 1; 5.1 ± 0.4 offspring

including 2.4 ± 0.2 daughters per female). In Nice, 86 males and 77 females were captured, among

which 54 produced 140 daughters (Table 1; 4.8 ± 0.3 offspring including 2.6 ± 0.1 daughters per

female). We genotyped 467 of these individuals at 18 microsatellite markers. We obtained the

genotype at all 18 microsatellite loci for 402 individuals, at 15-17 loci for 50 individuals, 1-10 loci for

11 individuals, and for four individuals, we obtained no amplification (Table 1). Only individuals with

successful amplification at >14 loci were included in the dataset, which represented 452 individuals

(96.8%).

No linkage disequilibrium was detected and null allele frequency was always lower than 6.5%

in the two populations. We found no genetic differentiation between the two populations (F_{st} = 0.01).

We nonetheless treated the two populations separately because the probability that a male captured

in one population had inseminated a female captured 300 km apart is negligible. Effective population

sizes were estimated to be 79 for Nice (IC 95% [44; 82]) and 51 for Valence (IC 95% [26; 56]). These

values are low and not significantly different.

Sib-mating avoidance in the field? Relatedness of actual versus potential mates

In Nice, the mean coefficient of relatedness between wild males and females was

r = 0.047 ± 0.001. We estimated that 11% ± 0.79% (IC 95%) of the relatedness coefficients between

these individuals were higher than the threshold r_{crit} = 0.16 (671 comparisons over 6083, from 79 males

× 77 females); the corresponding pairs were thus considered genetically related. The average

probability that a female encountered a related male in the field the day of its capture was 12.0% ± 4.9

(Fig. 2).

Thirty-five and 54 wild females from Valence and Nice, respectively, produced at least one

daughter which allowed to infer the genotype of their mate (Table 1). The mean relatedness between

these inferred pairs was 0.050 ± 0.009 in Nice and 0.044 ± 0.013 in Valence. Assuming r_{crit} = 0.16, we

found six and four genetically related pairs in Nice and Valence, respectively, that is, 11.1% ± 8.4% and

11.4% ± 10.5% of all pairs. For the population of Nice, this observed number of related pairs did not

differ from the number expected under the assumption of random mating (1000 permutations; mean

number of related pairs expected from random mating = 5.264; number observed = 6, p = 0.822; Fig.

3).

Proportion of diploid males and number of CSD alleles

Estimating inbreeding depression is required to understand the balance between costs and

benefits of inbreeding avoidance in a given species. Here, we assessed the cost of inbreeding by

estimating the number of diploid males in the population of Nice (no male was captured in Valence).

In species with sl-CSD, diploid males result from matched matings (parents sharing a common allele at

the *csd* locus), and the probability of matched mating in a panmictic population is $2/k$, where k is the

effective number of *csd* alleles. The number of alleles at the *csd* locus k could be estimated from the

proportion of diploid males among the males m and the proportion fertilization s (see SEM-E for the

calculation). As every evidence points to a balanced sex-ratio, we set s=0.5. We estimated the number

of *csd* alleles in the population of Nice, where random matings are suggested by the result on the

number of related pairs in the wild (see previous section). Six males were found diploids among the 79

males captured (m=7.6% ± 5.8%, IC 95%) yielding an estimation of 8.8 alleles at the *csd* and a

probability $2/k$ = 0.23 of matched matings.

Sex-biased dispersal in the field

During the field experiment, 50 females and 47 males were genotyped in patch 1, 18 females

and 22 males in patch 2. Genetic relatedness was estimated among females and among males within

patches (referred to as 1-1 for intra Patch 1 and 2-2 for intra Patch 2) as well as between patches

(referred to as 1-2) (Table 2). Mean relatedness differed within and between patches for females

(Kruskal-Wallis test: χ^2 = 14.88, *df* = 2, *p* < 1.10^{-4}) but not for males (Kruskal-Wallis test: χ^2 = 1.72, *df* =

2, *p* = 0.424). For females, relatedness was similar within each of two patches (1-1 and 2-2; Mann-

Whitney test: *U* = 80550, *p* = 0.520; Table 2) and higher than between patches (1-1 / 1-2 comparison,

Mann-Whitney test: *U* = 56724, *p* < 4.10^{-4}; 2-2 / 1-2 comparison, Mann-Whitney test: *U* = 65356, *p* =

0.012). Therefore, sex-biased dispersal occurs in *V. canescens*, with males dispersing more than

females.

Discussion

Sib-mating avoidance in *Venturia canescens* had been observed in no-choice behavioural

experiments conducted in laboratory conditions (Metzger et al., 2010a, Chuine, 2014). We show here

that when females can choose between related and unrelated males, they do mate with their sibs. The

density of males and the proportion of related individuals in the surroundings do not change the

pattern of inbreeding avoidance, suggesting that relatedness is not a cue for mate choice. Females

nonetheless appear to discriminate sibs from non-sibs, as shown by decreasing latency before

copulation after rejection of a sib. The sib-mating tolerance that we observed in the laboratory is

confirmed by our genetic approach of field populations. We showed that the observed frequency of

sib-mating corresponds to the probability of sib-mating expected from random encounters. Besides,

our field data suggest a slightly sex-biased dispersal, although without yielding passive sib-mating

avoidance. In the remainder of the discussion, we propose an evolutionary scenario to explain the

maintenance of kin recognition system and its apparent absence of use in the mate choice context, in

the lab or in the field.

In laboratory cages, the proportion of sib-mating perfectly matched the proportion of sibs

present. Consistently, mating rates between relatives, be they inferred from genetic analyses or from

a simulation model assuming random encounters, were both equal to 11%. These two results are

congruent with an absence of sib-mating avoidance in *V. canescens*, and questions past evidence of

kin recognition and avoidance in the contexts of both host parasitization (Bernstein and Driessen,

1996; Marris et al., 1996) and mate choice (Metzger et al., 2010a; Chuine et al., 2015). In particular,

no-choice bioassays had shown that females *V. canescens* mated twice more often with unrelated

males than with their brothers, and were therefore capable of sib-mating avoidance (Metzger et al.

2010a). In the presence of volatiles from brothers, females also discriminated against unrelated males,

suggesting that mate-choice could be underlined by volatile cues (Metzger et al., 2010a; Chuine et al.,

2015). Our laboratory cage experiment also suggests the ability of females to recognize kin and modify

a component of their mating behaviour (latency before copulation), but do not yield sib-mating

avoidance. We further discuss eight hypotheses to explain the differences between our findings and

those of Metzger et al., (2010a).

First, the observed lack of evidence for kin discrimination in our field study could result from a

misclassification of cousins or more distant individuals as full sibs. The reason is that when genetic

relatedness decreases to that of cousins (r ≤ 0.1875) sib-mating avoidance in *V. canescens* declines

steeply (Chuine, 2014). Hence, misclassifying cousins as brothers would yield an absence of sib-mating

avoidance, with no surprise. This is nonetheless unlikely. We did classify pairs as sibs versus non-sibs

based on a threshold r_{crit} = 0.16 that proved to minimize the probability of misclassification (about 10%

of errors). Besides, increasing this threshold to further restrict the definition of kinship did not change

our conclusions, that is, the observed number of related pairs was always similar to that expected

under random matings (r_{crit} = 0.2: 1000 permutations, 2 observed related pairs, 1.88 expected related

pairs, *p* = 1; r_{crit} = 0.25: 1000 permutations, 0 observed related pairs, 0.999 expected related pairs, *p* =

0.637). Hence, the hypothesis that poor kinship assignation has yielded spurious results falls short of

argument, and we conclude that the lack of evidence for sib-mating avoidance truly reflects a lack of

sib-mating avoidance, called sib-mating tolerance.

The second hypothesis is that we could face the ghost of past selection for sib-mating

avoidance. If the cost of inbreeding was higher in the evolutionary history of *V. canescens*, behaviours

allowing kin recognition and reducing the risk of inbreeding could have been selected for.

A third hypothesis is that sex-biased dispersal shapes the pattern of inbreeding avoidance in

the field (Gandon, 1999; Roze and Rousset, 2005). In *V. canescens* we found a slight male-biased

dispersal at a local scale (Table 2). However, in the population of Nice, the relatedness values between

and within patches (between which dispersal occurs) in both sexes are low (between 0.042 and 0.065,

Table2). Therefore, if this male-biased dispersal affects the structure of this population and encounter

rates between relatives, it should be a weakly effect.

The fourth hypothesis is that female choosiness in *V. canescens* is density-dependent, as

shown in other species (Kokko and Rankin, 2006; Duthie and Reid, 2016). Sib-mating avoidance would

arise in dense or aggregated populations with high mate encounter rates, but not in scarce populations

where the cost of discrimination – remaining virgin – is higher. This was shown in a small isolated

population of moose, where females accepted mates with higher relatedness during years where

males were less available (Herfindal et al., 2014). A similar reasoning may also hold true for highly

inbred populations, where sib-mating avoidance would result in a high probability of not mating at all,

a penalty that could exceed inbreeding depression (Kokko and Ots, 2006). In *V. canescens*, population

density affects mating success but not sib-mating probability. Besides, the proportions of related males

that we tested in the laboratory were respectively twice and four times higher than those found in the

field, and still, no sib-mating avoidance was observed. In the field, the distribution of the sib-

encounters probability is skewed towards zero, leading to more than half of the *V. canescens* females

having no chance to encounter a related male on a particular day, possibly because of the low number

of individuals in the field (Fig. 2). The average probability for a female to encounter a relative is

nonetheless in the same range as actual sib-mating (12%; Fig. 2). Kin recognition mechanisms could

then have persisted, but not sib-mating avoidance. We therefore conclude that *V. canescens* does not

avoid sib-mating, whatever the density of males and the level of inbreeding.

A fifth hypothesis to explain the absence of sib-mating avoidance despite kin recognition ability

is that female discrimination against related males decays over time. Encountering a succession of low-

quality males (*i.e.* genetically-related in our study) could decrease female choosiness, as demonstrated

in a monoandrous wolf spider (Stoffer and Uetz, 2016). We have shown in the lab experiment that

even if females did not avoid sib-mating, their latency before copulation decreased significantly after

having rejected a brother. This result confirms that *V. canescens* is capable of discriminating relatives

in a context of mating (see also Metzger et al., 2010a; Chuine, 2014; Chuine et al., 2015) and suggests

that the presence of relatives could affect mating by decreasing the latency before copulation, as

expected in this hypothesis. We cannot therefore rule out this hypothesis but more experiments are

needed to firmly test the evolution of female choosiness over time.

The sixth hypothesis is that tolerance to sib-mating, as observed in *Venturia canescens*, is often

explained by low inbreeding depression or high cost of outbreeding (Kokko and Ots, 2006; Puurtinen,

2011). We found a low effective size in both sample sites (Ne = 51 in Valence, Ne = 79 in Nice), in the

same range as threatened bumblebees (Ellis et al., 2006) or Hine's emerald dragonflies (Monroe and

Britten, 2015). *V. canescens* could be prone to loss of genetic diversity, possibly amplified by inbreeding

tolerance, which would in turn increase inbreeding load. However, alleles at the *csd* locus, the major

mechanism of inbreeding depression in *V. canescens*, are under strong balancing selection, and are

thus expected to overcome genetic impoverishment, as observed in populations of Asiatic honeybee

(Gloag et al., 2016). Moreover, in *V. canescens*, inbreeding has only a moderate negative impact on

egg load and hatching rate, and no effect on other fitness components such as mating probability or

body size (Vayssade et al., 2014).

The seventh hypothesis is that despite common beliefs, the genetic load resulting from sl-CSD

might not be dramatic in *Venturia canescens*. In the population of Nice, we estimated 8.8 alleles at the

csd locus, which is in the range of what was previously found in other species of the order

Hymenoptera (8-86 alleles, mostly between 9 to 20; Cook and Crozier, 1995). Under random matings,

this yields a probability of matched-matings of 23% (2/8.8). In case of brother-sister mating, the

probability of matched matings increases to 50%. Assuming that females fertilize half of their eggs so

that matched matings results in 25% diploid males, and that diploid males are sterile (Fauvergue et al.

2015), the expected reduction in the number of fertile offspring under random mating would be 5.8%

(0.23 x 0.25). If sib-mating was perfectly avoided, the expected reduction in the number of fertile

offspring would be 4.8% (0.19 x 0.25, see SEM-E for the calculation). Sib-mating avoidance would thus

result in about 1% more fertile offspring in the populations we studied, a fitness gain that may appear

insignificant. Hence, the relatively low probability of matched matings and the consequent benign

inbreeding load may be insufficient to select for sib-mating avoidance. *V. canescens* could so satisfy

the hypothesis of inbreeding tolerance (Waser et al., 1986; Kokko and Ots, 2006).

At last, the eight hypothesis lean on the possibility that the discrepancies between sib-mating

avoidance evidenced in previous research and our finding of inbreeding tolerance could highlight the

evolution of kin-recognition mechanisms for other purposes than sib-mating avoidance. This would

have produced artefactual behaviours when observed in small arenas (Metzger et al., 2010a) but

vanishing in larger cages and in the field. Evidence for active kin recognition is pervasive in various

ecological contexts such as territory defence, reciprocal altruism, or dominance hierarchies (Mateo,

2004). In *V. canescens*, the evolution of kin recognition for the avoidance of superparasitism could

explain the maintenance of kin recognition without sib-mating avoidance. Indeed, females are able to

discriminate the self, their relatives and their conspecifics when laying eggs, and avoid already

parasitized hosts. As a single offspring can develop in a host, superparasitism has a high cost for female

fitness. Consistently, the intensity of superparasitism avoidance increases with genetic relatedness to

the first laying female (Hubbard et al., 1987; Marris et al., 1996; Amat et al., 2009). If kin recognition

evolved in this context, kin discrimination in a situation of mate choice could be a by-product of this

primary selection on kin recognition. We could therefore expect a relatively general molecular

pathway, which would have been expressed in a small-scale mating bioassay: kin recognition during

mating is probably mediated by semiochemicals from the insect cuticle (cuticular hydrocarbons,

Metzger et al., 2010a; Chuine, 2014) and discrimination against superparasitized larvae is driven by

weakly volatile hydrocarbons left by females during oviposition (Fisher, 1961; Harrison et al., 1985).

These mechanistic similarities point towards a unique pathway for kin recognition. Superparasitism

avoidance could act as a strong selective pressure for the maintenance of kin recognition, because the

cost of superparasitism is positively correlated with the relatedness among competing females.

Further experiments are needed to test the hypothesis that sib-mating avoidance observed in

laboratory conditions is a by-product of cognitive abilities which have evolved in a context of

competition.

Bibliography

Aguilera-Olivares, D., Flores-Prado, L., Véliz, D., Niemeyer, H.M., 2015. Mechanisms of inbreeding
avoidance in the one-piece drywood termite Neotermes chilensis. Insectes Sociaux 62, 237–
245. doi:10.1007/s00040-015-0399-1
Amat, I., Besnard, S., Foray, V., Pelosse, P., Bernstein, C., Desouhant, E., 2012. Fuelling flight in a
parasitic wasp: which energetic substrate to use? Ecol. Entomol. 37, 480–489.
doi:10.1111/j.1365-2311.2012.01388.x
Amat, I., Desouhant, E., Bernstein, C., 2009. Differential use of conspecific-derived information by
sexual and asexual parasitic wasps exploiting partially depleted host patches. Behav. Ecol.
Sociobiol. 63, 563–572. doi:10.1007/s00265-008-0691-4
Angeloni, F., Ouborg, N.J., Leimu, R., 2011. Meta-analysis on the association of population size and
life history with inbreeding depression in plants. Biol. Conserv. 144, 35–43.
doi:10.1016/j.biocon.2010.08.016
Bernstein, C., Driessen, G., 1996. Patch-Marking and Optimal Search Patterns in the Parasitoid
Venturia canescens. J. Anim. Ecol. 65, 211. doi:10.2307/5724
Beukeboom, L., Driessen, G., Luckerhoff, L., Bernstein, C., Lapchin, L., Van Alphen, J., 1999.
Distribution and relatedness of sexual and asexual Venturia canescens (Hymenoptera). Proc
Exp Appl Entomol Neth NEV Amst. 10, 23–28.
Beukeboom, L.W., 2001. Single-locus complementary sex determination in the ichneumonid Venturia
canescens (Gravenhorst) (Hymenoptera). Neth. J. Zool. 51, 1–15.
Billing, A.M., Lee, A.M., Skjelseth, S., Borg, Å.A., Hale, M.C., Slate, J., Pärn, H., Ringsby, T.H., Sæther,
B.-E., Jensen, H., 2012. Evidence of inbreeding depression but not inbreeding avoidance in a
natural house sparrow population. Mol. Ecol. 21, 1487–1499. doi:10.1111/j.1365-
294X.2012.05490.x
Biosa, D., Grignolio, S., Sica, N., Pagon, N., Scandura, M., Apollonio, M., 2015. Do relatives like to stay
closer? Spatial organization and genetic relatedness in a mountain roe deer population. J.
Zool. 296, 30–37. doi:10.1111/jzo.12214
Bonadonna, F., Sanz-Aguilar, A., 2012. Kin recognition and inbreeding avoidance in wild birds: the
first evidence for individual kin-related odour recognition. Anim. Behav. 84, 509–513.
doi:10.1016/j.anbehav.2012.06.014
Bourdais, D., Hance, T., 2009. Lack of behavioural evidence for kin avoidance in mate choice in a
hymenopteran parasitoid (Hymenoptera: Braconidae). Behav. Processes 81, 92–94.
doi:10.1016/j.beproc.2009.02.015
Breedveld, M.C., Fitze, P.S., 2015. A matter of time: delayed mate encounter postpones mating
window initiation and reduces the strength of female choosiness. Behav. Ecol. Sociobiol. 69,
533–541. doi:10.1007/s00265-014-1864-y
Bretman, A., Newcombe, D., Tregenza, T., 2009. Promiscuous females avoid inbreeding by controlling
sperm storage. Mol. Ecol. 18, 3340–3345. doi:10.1111/j.1365-294X.2009.04301.x
Bretman, A., Wedell, N., Tregenza, T., 2004. Molecular evidence of post-copulatory inbreeding
avoidance in the field cricket Gryllus bimaculatus. Proc. R. Soc. B-Biol. Sci. 271, 159–164.
doi:10.1098/rspb.2003.2563
Chapman, Partridge, 1998. Sex-specific selection on time to remate in Drosophila melanogaster.
Anim. Behav. 56, 1267–1278. doi:10.1006/anbe.1998.0900
Chapuis, M.-P., Estoup, A., 2007. Microsatellite Null Alleles and Estimation of Population
Differentiation. Mol. Biol. Evol. 24, 621–631. doi:10.1093/molbev/msl191
Charlesworth, D., Charlesworth, B., 1987. Inbreeding Depression and its Evolutionary Consequences.
Annu. Rev. Ecol. Syst. 18, 237–268.
Charpentier, M.J.E., Boulet, M., Drea, C.M., 2008a. Smelling right: the scent of male lemurs advertises
genetic quality and relatedness. Mol. Ecol. 17, 3225–3233. doi:10.1111/j.1365-
294X.2008.03831.x

Charpentier, M.J.E., Williams, C.V., Drea, C.M., 2008b. Inbreeding depression in ring-tailed lemurs
(Lemur catta): genetic diversity predicts parasitism, immunocompetence, and survivorship.
Conserv. Genet. 9, 1605–1615. doi:10.1007/s10592-007-9499-4
Chessel, D., Dufour, A.-B., Thioulouse, J., 2004. The ade4 package-I- One-table methods. R News 4, 5–
10.
Chuine, A., 2014. Des mécanismes aux conséquences adaptatives du choix du partenaire sexuel pour
la compatibilité génétique.   Exemple d'un hyménoptère parasitoïde soumis à la dépression
de consanguinité. (PhD thesis). Université Claude Bernard Lyon 1.
Chuine, A., Sauzet, S., Debias, F., Desouhant, E., 2015. Consequences of genetic incompatibility on
fitness and mate choice: the male point of view. Biol. J. Linn. Soc. 114, 279–286.
doi:10.1111/bij.12421
Cohas, A., Yoccoz, N.G., Bonenfant, C., Goossens, B., Genton, C., Galan, M., Kempenaers, B., Allainé,
D., 2008. The genetic similarity between pair members influences the frequency of extrapair
paternity in alpine marmots. Anim. Behav. 76, 87–95. doi:10.1016/j.anbehav.2008.01.012
Collet, M., Vayssade, C., Auguste, A., Mouton, L., Desouhant, E., Malausa, T., Fauvergue, X., 2016.
Diploid male production correlates with genetic diversity in the parasitoid wasp Venturia
canescens: a genetic approach with new microsatellite markers. Ecol. Evol. 6, 6721–6734.
doi:10.1002/ece3.2370
Cook, J.M., 1993. Sex determination in the Hymenoptera: a review of models and evidence. Heredity
71, 421–435.
Cook, J.M., Crozier, R.H., 1995. Sex determination and population biology in the hymenoptera.
Trends Ecol. Evol. 10, 281–286. doi:10.1016/0169-5347(95)90011-X
Cooney, R., Bennett, N.C., 2000. Inbreeding avoidance and reproductive skew in a cooperative
mammal. Proc. R. Soc. Lond. B Biol. Sci. 267, 801–806. doi:10.1098/rspb.2000.1074
Corbet, S.A., 1971. Mandibular gland secretion of larvae of the flour moth, Anagasta kuehniella,
contains an epideictic pheromone and elicts oviposition movements in a Hymenopteran
parasite. Nature 232, 481–484.
Cornell, S.J., Tregenza, T., 2007. A new theory for the evolution of polyandry as a means of
inbreeding avoidance. Proc. R. Soc. Lond. B Biol. Sci. 274, 2873–2879.
doi:10.1098/rspb.2007.0926
Desouhant, E., Driessen, G., Lapchin, L., Wielaard, S., Bernstein, C., 2003. Dispersal between host
populations in field conditions: navigation rules in the parasitoid Venturia canescens. Ecol.
Entomol. 28, 257–267. doi:10.1046/j.1365-2311.2003.00511.x
Dowling, D.K., Friberg, U., Arnqvist, G., 2007. A comparison of nuclear and cytoplasmic genetic effects
on sperm competitiveness and female remating in a seed beetle. J. Evol. Biol. 20, 2113–2125.
doi:10.1111/j.1420-9101.2007.01433.x
Driessen, G., Bernstein, C., 1999. Patch departure mechanisms and optimal host exploitation in an
insect parasitoid. J. Anim. Ecol. 68, 445–459.
Duthie, A.B., Reid, J.M., 2016. Evolution of Inbreeding Avoidance and Inbreeding Preference through
Mate Choice among Interacting Relatives. Am. Nat. 188, 651–667. doi:10.1086/688919
Ellis, J.S., Knight, M.E., Darvill, B., Goulson, D., 2006. Extremely low effective population sizes, genetic
structuring and reduced genetic diversity in a threatened bumblebee species, Bombus
sylvarum (Hymenoptera: Apidae): GENETIC STRUCTURING IN DECLINING BUMBLEBEES. Mol.
Ecol. 15, 4375–4386. doi:10.1111/j.1365-294X.2006.03121.x
Fauvergue, X., Chuine, A., Vayssade, C., Auguste, A., Desouhant, E., 2015. Sterile males in a parasitoid
wasp with complementary sex determination: from fitness costs to population extinction.
BMC Ecol. 15. doi:10.1186/s12898-014-0032-6
Fischer, K., Karl, I., Heuskin, S., Janowitz, S., Dötterl, S., 2015. Kin Recognition and Inbreeding
Avoidance in a Butterfly. Ethology 121, 977–984. doi:10.1111/eth.12410
Fisher, R., 1961. Study in Insect Multiparasitism .1. Host Selection and Oviposition. J. Exp. Biol. 38,
267-.

Gandon, S., 1999. Kin Competition, the Cost of Inbreeding and the Evolution of Dispersal. J. Theor.
616   Biol. 200, 345–364. doi:10.1006/jtbi.1999.0994
617   Gloag, R., Ding, G., Christie, J.R., Buchmann, G., Beekman, M., Oldroyd, B.P., 2016. An invasive social
618   insect overcomes genetic load at the sex locus. Nat. Ecol. Evol. 1, s41559-016-0011-016.
619   doi:10.1038/s41559-016-0011
620   Goudet, J., 1995. FSTAT (Version 1.2): A Computer Program to Calculate F-Statistics. J. Hered. 86,
621   485–486.
622   Greenwood, P.J., Harvey, P.H., Perrins, C.M., 1978. Inbreeding and dispersal in the great tit. Nature
623   271, 52–54. doi:10.1038/271052a0
624   Harrison, E.G., Fisher, R.C., Ross, K.M., 1985. The temporal effects of Dufour's gland secretion in host
625   discrimination by Nemeritis canescens. Entomol. Exp. Appl. 38, 215–220. doi:10.1111/j.1570-
626   7458.1985.tb03521.x
627   Hatchwell, B.J., Russell, A.F., Ross, D.J., Fowlie, M.K., 2000. Divorce in cooperatively breeding long-
628   tailed tits: a consequence of inbreeding avoidance? Proc. R. Soc. B Biol. Sci. 267, 813–819.
629   Herfindal, I., Haanes, H., Roed, K.H., Solberg, E.J., Markussen, S.S., Heim, M., Saether, B.-E., 2014.
630   Population properties affect inbreeding avoidance in moose. Biol. Lett. 10, 20140786–
631   20140786. doi:10.1098/rsbl.2014.0786
632   Højsgaard, S., Halekoh, U., Yan, J., 2006. The R Package geepack for Generalized Estimating
633   Equations. J. Stat. Softw. 15. doi:10.18637/jss.v015.i02
634   Howard, R.W., Blomquist, G.J., 2005. Ecological, Behavioral, and Biochemical Aspects of Insect
635   Hydrocarbons. Annu. Rev. Entomol. 50, 371–393.
636   doi:10.1146/annurev.ento.50.071803.130359
637   Hubbard, S.F., Marris, G., Reynolds, A., Rowe, G.W., 1987. Adaptive Patterns in the Avoidance of
638   Superparasitism by Solitary Parasitic Wasps. J. Anim. Ecol. 56, 387. doi:10.2307/5055
639   Johansson, B.G., Jones, T.M., 2007. The role of chemical communication in mate choice. Biol. Rev. 82,
640   265–289. doi:10.1111/j.1469-185X.2007.00009.x
641   Jones, O.R., Wang, J., 2010. COLONY: a program for parentage and sibship inference from multilocus
642   genotype data. Mol. Ecol. Resour. 10, 551–555. doi:10.1111/j.1755-0998.2009.02787.x
643   Keller, L.F., Waller, D.M., 2002. Inbreeding effects in wild populations. Trends Ecol. Evol. 17, 230–241.
644   doi:10.1016/S0169-5347(02)02489-8
645   Kokko, H., Ots, I., 2006. When Not to Avoid Inbreeding. Evolution 60, 467–475. doi:10.1111/j.0014-
646   3820.2006.tb01128.x
647   Kokko, H., Rankin, D.J., 2006. Lonely hearts or sex in the city? Density-dependent effects in mating
648   systems. Philos. Trans. R. Soc. B Biol. Sci. 361, 319–334. doi:10.1098/rstb.2005.1784
649   Laloi, D., Eizaguirre, C., Fédérici, P., Massot, M., 2011. Female choice for heterozygous mates changes
650   along successive matings in a lizard. Behav. Processes 88, 149–154.
651   doi:10.1016/j.beproc.2011.08.011
652   Lardy, S., Cohas, A., Figueroa, I., Allaine, D., 2011. Mate change in a socially monogamous mammal:
653   evidences support the "forced divorce" hypothesis. Behav. Ecol. 22, 120–125.
654   doi:10.1093/beheco/arq168
655   Liang, K.-Y., Zeger, S.L., 1986. Longitudinal Data Analysis Using Generalized Linear Models. Biometrika
656   73, 13–22. doi:10.2307/2336267
657   Marris, G.C., Hubbard, S.F., Scrimgeour, C., 1996. The perception of genetic similarity by the solitary
658   parthenogenetic parasitoid Venturia canescens, and its effects on the occurrence of
659   superparasitism. Entomol. Exp. Appl. 78, 167–174. doi:10.1111/j.1570-7458.1996.tb00778.x
660   Mateo, J.M., 2004. Recognition systems and biological organization: The perception component of
661   social recognition. Ann. Zool. Fenn. 41, 729–745.
662   Mateo Leach, I., Ferber, S., van de Zande, L., Beukeboom, L.W., 2012. Genetic variability of
663   arrhenotokous and thelytokous Venturiacanescens (Hymenoptera). Genetica 140, 53–63.
664   doi:10.1007/s10709-012-9657-6

665   Metzger, M., Bernstein, C., Desouhant, E., 2008. Does constrained oviposition influence offspring sex
666   ratio in the solitary parasitoid wasp Venturia canescens? Ecol. Entomol. 33, 167–174.
667   doi:10.1111/j.1365-2311.2007.00953.x
668   Metzger, M., Bernstein, C., Hoffmeister, T.S., Desouhant, E., 2010a. Does Kin Recognition and Sib-
669   Mating Avoidance Limit the Risk of Genetic Incompatibility in a Parasitic Wasp? PLoS ONE 5,
670   e13505. doi:10.1371/journal.pone.0013505
671   Metzger, M., Fischbein, D., Auguste, A., Fauvergue, X., Bernstein, C., Desouhant, E., 2010b. Synergy in
672   information use for mate finding: demonstration in a parasitoid wasp. Anim. Behav. 79,
673   1307–1315. doi:10.1016/j.anbehav.2010.03.003
674   Monroe, E.M., Britten, H.B., 2015. Single-sample estimation of effective population size in several
675   populations of the endangered Hine's emerald dragonfly. Freshw. Sci. 34, 1058–1064.
676   doi:10.1086/682073
677   Nichols, H.J., 2017. The causes and consequences of inbreeding avoidance and tolerance in
678   cooperatively breeding vertebrates. J. Zool. 303, 1–14. doi:10.1111/jzo.12466
679   Nichols, H.J., Cant, M.A., Hoffman, J.I., Sanderson, J.L., 2014. Evidence for frequent incest in a
680   cooperatively breeding mammal. Biol. Lett. 10, 20140898–20140898.
681   doi:10.1098/rsbl.2014.0898
682   Olson, L.E., Blumstein, D.T., Pollinger, J.R., Wayne, R.K., 2012. No evidence of inbreeding avoidance
683   despite demonstrated survival costs in a polygynous rodent: NO INBREEDING AVOIDANCE IN
684   A POLYGYNOUS RODENT. Mol. Ecol. 21, 562–571. doi:10.1111/j.1365-294X.2011.05389.x
685   Prugnolle, F., de Meeus, T., 2002. Inferring sex-biased dispersal from population genetic tools: a
686   review. Heredity 88, 161–165. doi:10.1038/sj.hdy.6800060
687   Pusey, A., Wolf, M., 1996. Inbreeding avoidance in animals. Trends Ecol. Evol. 11, 201–206.
688   Puurtinen, M., 2011. Mate Choice for Optimal (k)inbreeding. Evolution 65, 1501–1505.
689   doi:10.1111/j.1558-5646.2010.01217.x
690   R Core Team, 2015. R: A language and environment for statistical   computing. R Found. Stat.
691   Comput. Vienna Austria.
692   Raymond, M., Rousset, F., 1995. An Exact Test for Population Differentiation. Evolution 49, 1280–
693   1283. doi:10.2307/2410454
694   Reid, J.M., Arcese, P., Keller, L.F., Germain, R.R., Duthie, A.B., Losdat, S., Wolak, M.E., Nietlisbach, P.,
695   2015. Quantifying inbreeding avoidance through extra-pair reproduction. Evol. Int. J. Org.
696   Evol. 69, 59–74. doi:10.1111/evo.12557
697   Reynolds, S.M., Uy, J.A.C., Patricelli, G.L., Coleman, S.W., Braun, M.J., Borgia, G., 2014. Tests of the
698   kin selection model of mate choice and inbreeding avoidance in satin bowerbirds. Behav.
699   Ecol. 25, 1005–1014. doi:10.1093/beheco/aru065
700   Roze, D., Rousset, F., 2005. Inbreeding Depression and the Evolution of Dispersal Rates: A Multilocus
701   Model. Am. Nat. 166, 708–721. doi:10.1086/497543
702   Salt, G., 1976. The hosts of Nemeritis canescens: a problem in the host specificity of insect
703   parasitoids. Ecol. Entomol. 1, 63–67.
704   Schneider, M.V., Beukeboom, L.W., Driessen, G., Lapchin, L., Bernstein, C., Van Alphen, J.J.M., 2002.
705   Geographical distribution and genetic relatedness of sympatrical thelytokous and
706   arrhenotokous populations of the parasitoid Venturia canescens (Hymenoptera). J. Evol. Biol.
707   15, 191–200. doi:10.1046/j.1420-9101.2002.00394.x
708   Stoffer, B., Uetz, G.W., 2016. Tuft size matters: the effects of adult visual social experience on female
709   mate preferences in a wolf spider. Behav. Ecol. Sociobiol. 70, 2211–2221.
710   doi:10.1007/s00265-016-2225-9
711   Szulkin, M., Sheldon, B.C., 2008. Dispersal as a means of inbreeding avoidance in a wild bird
712   population. Proc. R. Soc. B Biol. Sci. 275, 703–711. doi:10.1098/rspb.2007.0989
713   Szulkin, M., Zelazowski, P., Nicholson, G., Sheldon, B.C., 2009. Inbreeding avoidance under different
714   null models of random mating in the great tit. J. Anim. Ecol. 78, 778–788. doi:10.1111/j.1365-
715   2656.2009.01544.x

Tan, C.K.W., Løvlie, H., Pizzari, T., Wigby, S., 2012. No evidence for precopulatory inbreeding
avoidance in Drosophila melanogaster. Anim. Behav. 83, 1433–1441.
doi:10.1016/j.anbehav.2012.03.015
Therneau, T.M., 2015. coxme: Mixed Effects Cox Models.
Thünken, T., Bakker, T.C.M., Baldauf, S.A., Kullmann, H., 2007. Active Inbreeding in a Cichlid Fish and
Its Adaptive Significance. Curr. Biol. 17, 225–229. doi:10.1016/j.cub.2006.11.053
Tregenza, T., Wedell, N., 2002. Polyandrous females avoid costs of inbreeding. Nature 415, 71–73.
doi:10.1038/415071a
van Wilgenburg, E., Driessen, G., Beukeboom, L.W., 2006. Single locus complementary sex
determination in Hymenoptera: an "unintelligent" design? Front. Zool. 3, 1.
doi:10.1186/1742-9994-3-1
Vayssade, C., de Fazio, C., Quaglietti, B., Auguste, A., Ris, N., Fauvergue, X., 2014. Inbreeding
Depression in a Parasitoid Wasp with Single-Locus Complementary Sex Determination. PLoS
ONE 9, e97733. doi:10.1371/journal.pone.0097733
Wang, J., 2011. coancestry: a program for simulating, estimating and analysing relatedness and
inbreeding coefficients. Mol. Ecol. Resour. 11, 141–145. doi:10.1111/j.1755-
0998.2010.02885.x
Wang, J., 2009. A new method for estimating effective population sizes from a single sample of
multilocus genotypes. Mol. Ecol. 18, 2148–2164. doi:10.1111/j.1365-294X.2009.04175.x
Waser, P.M., Austad, S.N., Keane, B., 1986. When Should Animals Tolerate Inbreeding? Am. Nat. 128,
529–537. doi:10.1086/284585
737

Authors' contributions

M.C, I.A., X.F., L.M. and E.D. designed the experiment and M.C., S.S and A.A collected the data. M.C.,

I.A., L.M. and E.D. designed and interpreted the analyses. M.C. conducted the analyses and assembled

a first draft. M.C., I.A., X.F., L.M. and E.D. all contributed to revisions.

Acknowledgments

We want to thank François Debias for his help in the field capture of insects, rearing and technical

assistance, and Aurélie Blin for technical advices on genotyping. We thank Angèle, Anatole and Auguste

Desouhant for their help in the field sampling. This work was partly funded thanks to Agence Nationale

de la Recherche ("Sextinction" project, ANR-2010-BLAN-1717). We declare no conflict of interest.

Table 1 – Number of adults *V. canescens* captured in the wild populations of Nice and Valence.

Pop = population; Number of mothers = number of wild females captured in the field and which have

produced at least one daughter; F = female, M = male.

Pop	Sex	Wild individuals	Wild individuals successfully genotyped	Number of mothers	Total number of daughters	Daughters successfully genotyped
Nice	F	77	77 (100%)	54	140	134 (96%)
	M	86	79 (92%)	-	-	-
Valence	F	78	77 (99%)	35	86	85 (99%)

754 **Table 2 –** Mean relatedness *r* (± SE) among females and among males of *V. canescens*, either within

755 patches (1-1 or 2-2) or between patches (1-2) of host plants (carob trees) in the population of Nice.

756 F = female, M = male.

Sex	Patch	Mean *r* (± SE)
F	1-1	0.065 ± 0.002
	2-2	0.060 ± 0.006
	1-2	0.050 ± 0.002
M	1-1	0.042 ± 0.002
	2-2	0.048 ± 0.006
	1-2	0.043 ± 0.003

758   **Table 3 -** Effect of male density and sib frequency on sib-mating avoidance in *V. canescens*. Two

759   densities of males (*D*: high density = 9 males, and *d*: low density = 3 males) and two frequencies of

760   brothers (*F*: high frequency = 2/3 brothers and *f*: low frequency = 1/3 brothers) were manipulated in a

761   cage experiment. Here, we present the number of successful matings for each factor level (20, except

762   for the *d-F* modality where only 18 were reached).

		D	d
F	Brother	12	12
	Unrelated	8	6
	Total	20/23 trials	18/44 trials
f	Brother	6	6
	Unrelated	14	14
	Total	20/26 trials	20/42 trials

766   **Table 4 –** Effect of male density, sib frequency (and their interaction) and the number of male

767   rejections on mating latency in *V. canescens*. We analysed the deviance table of the mixed-effect Cox

768   model with the female family as a random effect.

	Log likelihood	χ^2	*df*	*p*-value
NULL	-351.51			
Male density	-338.44	26.14	1	$3.17 \cdot 10^{-7}$
Sib frequency	-337.57	1.74	1	0.19
Rejection of sib	-332.45	10.25	1	$1.36 \cdot 10^{-3}$
Rejection of unrelated	-331.46	1.97	1	0.16
Density $\times$ sib frequency	-331.46	0.01	1	0.92

Figure 1 – Location of carob trees from the Nice field site where *V. canescens* were captured during the summer 2014. Each dot represents a tree. Axes are GPS coordinates in decimal degrees. Blue dots represent trees of "Patch 1" and orange dots represent trees of "Patch 2" (see also Table 2). Six trees (black dots) were not included because they did not fit our definition of patches.

Figure 2 – Distribution of the probability that a female *V. canescens* encountered a genetically related male during the day of capture in the Nice population. For each female caught, we computed its relatedness with all males captured the same day. We then counted the number of relationships with $r \geq$ 0.16 and we divided this number by the total number of relatedness computed (*i.e.* the number of males captured the same day).

Figure 3 – Theoretical distribution of the number of crosses between genetically related *V. canescens* male and female under the assumption of random mating. The theoretical distribution was built using random sampling of a mate for each of the 51 females from the Nice population, estimating *r* for each pair and counting those with *r* > 0.16. Sampling was reiterated 1000 times. The red line represents the observed number of pairs with a *r* > 0.16.