Neuronal origin of the temporal dynamics of spontaneous BOLD activity correlation

Teppei Matsui[1,2*], Tomonari Murakami[1,2] and Kenichi Ohki[1,2,3*]

1. Department of Physiology, The University of Tokyo School of Medicine, Tokyo, Japan
2. Department of Molecular Physiology, Graduate School of Medical Sciences, Kyushu University, Fukuoka, Japan
3. International Research Center for Neurointelligence (WPI-IRCN), The University of Tokyo, Tokyo, Japan

Send Correspondence To:
Teppei Matsui, Ph.D. (Email: tematsui@m.u-tokyo.ac.jp)
or
Kenichi Ohki, M.D, Ph.D. (Email: kohki@m.u-tokyo.ac.jp)
Department of Physiology
The University of Tokyo School of Medicine
Tokyo 113-0033, Japan
Phone: (+81)-3-5841-3459

Abstract (196/200 words)

Resting-state functional connectivity (FC) has become a major fMRI method to study network organization of human brains. There is recent interest in the temporal fluctuations of FC calculated using short time-windows ("dynamic FC") because it could provide information inaccessible with conventional "static" FC that is typically calculated using the entire scan lasting several tens of minutes. Although multiple studies have revealed considerable temporal fluctuations in FC, it is still unclear whether the fluctuations of FC measured in hemodynamics reflect the dynamics of underlying neural activity. We addressed this question using simultaneous imaging of neuronal calcium and hemodynamic signals in mice, and found coordinated temporal dynamics of calcium FC and hemodynamic FC measured in the same short time windows. Moreover, we found that variation in transient neuronal coactivation patterns (CAPs) was significantly related to temporal fluctuations of sliding window FC in hemodynamics. Finally, we show that observed dynamics of FC cannot be fully accounted for by simulated data assuming stationary FC. These results provide evidence for the neuronal origin of dynamic FC and further suggest that information relevant to FC is condensed in temporally sparse events that can be extracted using a small number of time points.

Keywords: calcium imaging, BOLD, resting state, functional connectivity, stationarity

Introduction

Resting state functional connectivity (FC) uses temporal correlation of spontaneous neuronal activity to assess network organization of brain regions in a non-invasive manner (Fox and Raichle 2007). Traditionally, FC has been calculated using all time points in a scan that typically lasts between several minutes to tens of minutes (Biswal et al. 1995; Fox et al. 2005; Van Dijk et al. 2010). Such "static" FC has been shown to largely reflect anatomical connectivity (Adachi et al. 2012; Honey et al. 2009; Matsui et al. 2012; Matsui et al. 2011; Vincent et al. 2007). Recently, in contrast to traditional analysis of "static" FC, the temporal fluctuation of FC across short time windows is increasingly recognized as a useful aspect of FC (Allen et al. 2014; Hutchison et al. 2013; Zalesky et al. 2014). Such "dynamic" FC (dFC) calculated using short time-windows could provide information that is inaccessible with static FC about the functional network organizations of healthy and diseased brains (Calhoun et al. 2014; Preti et al. 2016) [We note that the term "dynamics" refers to the non-stationarity of FC obtained with the sliding window analyses and does not refer to a process that is not invariant against to temporal reordering of the samples (Liégeois et al. 2017)]. The presence of temporal fluctuations in FC has also informed theoreticians to constrain realistic models of brain networks (Deco et al. 2013; Hansen et al. 2015; Messé et al. 2014).

However, despite growing interest, the neurophysiological basis of dFC is still weak. Previous attempts to investigate neural origin of dFC by simultaneous measurement of electrophysiological and functional magnetic resonance imaging (fMRI) are limited in several ways (Lu et al. 2007; Pan et al. 2011; Tagliazucchi et al. 2012b; Thompson et al. 2013). In some studies, electrophysiological recording was limited to a small number of recording sites due to technical difficulty (Lu et al. 2007; Pan et al. 2011; Thompson et al. 2013); hence, information on the global pattern of neuronal activity was lacking. In another study, electrophysiological signals were obtained with an electroencephalogram, which records global neuronal activity but lacks precise spatial information (Tagliazucchi et al. 2012b). Thus, the link between temporal fluctuations of FC in hemodynamics and that of large-scale neuronal activity has not been adequately proven.

Several studies have also questioned whether the apparent "dynamics" of FC calculated using the sliding window method is related to temporal instability of

spontaneous brain network (Hindriks et al. 2016; Laumann et al. 2016). While many studies have attributed temporal fluctuations of sliding window FC to non-stationarity of spontaneous neuronal activity correlation (Allen et al. 2014; Zalesky et al. 2014), some recent studies have demonstrated that the temporal fluctuations of FC observed in the real and the simulated data, which is stationary by construction, are statistically indistinguishable (Hindriks et al. 2016; Laumann et al. 2016). Furthermore, Laumann and colleagues have shown that, in the human resting-state BOLD time series are largely stationary, discounting head-motion and fluctuating arousal (Laumann et al. 2016). Therefore, not only the neuronal basis of dFC, but also the existence of statistical non-stationarity of FC, or at least the capability of sliding window methods to detect the non-stationarity, is called into question.

In the present study, we addressed these questions using simultaneous imaging of neuronal calcium and blood oxygen level dependent (BOLD) hemodynamic signals in the entire neocortex of transgenic mice expressing a genetically encoded calcium indicator (Matsui et al. 2016; Vanni and Murphy 2014; White et al. 2011). In the present experimental setup, wide-field calcium signal provided access to neuronal activity at higher temporal resolution and signal-to-noise ratio compared to that of hemodynamic signal (Matsui et al. 2016; Murakami et al. 2017; Murakami et al. 2015; Tohmi et al. 2014; Vanni and Murphy 2014). Moreover, unlike human fMRI data, in the present dataset, mice were tightly head-fixed and lightly anesthetized; thus, excluding head motions from contaminating FC. Main findings of the present study are as follows. First, we found consistency between the "dynamics" of FC calculated using calcium and hemodynamic signals, suggesting the neuronal origin of the temporal fluctuations of hemodynamic FC. Second, we found that temporal fluctuations of the spatial pattern of transient neuronal coactivations as measured in calcium signal were significantly correlated with temporal fluctuations of hemodynamic FC. Finally, we found that statistical properties of sliding window FC were significantly different between the real and the simulated data suggesting non-stationarity of resting-state FC.

Materials and Methods

Animals

Emx1-IRES-cre and Ai38 (Zariwala et al. 2012) mice were obtained from the Jackson Laboratory (Sacramento, CA). These mice were crossed and all cortical excitatory neurons expressed GCaMP3. Mice (P60–P90) were prepared for *in vivo* wide-field simultaneous imaging. Anesthesia was induced with isoflurane (3 %) and maintained with isoflurane (1 – 2 % in surgery, 0.5 – 0.8 % during imaging) and chlorprothixene (0.3 – 0.8 mg/kg, intramuscular injection). For simultaneous imaging of calcium and hemodynamic signals, a custom-made metal head plate was attached to the skull using dental cement (Sun Medical Company, Ltd, Shiga, Japan) and a large craniotomy was made over the whole cortex. The craniotomy was sealed with 1 % agarose and a glass coverslip. During the imaging, body temperature was maintained by a heat pad. All experiments were carried out in accordance with the NIH Guide for the Care and Use of Laboratory Animals, the institutional animal welfare guidelines set forth by the Animal Care and Use Committee of Kyushu University, and the study was approved by the Ethical Committee of Kyushu University.

Simultaneous Calcium and Intrinsic Signal Imaging

The data for simultaneous imaging of calcium and hemodynamic signals was taken from a published report (Matsui et al. 2016). Briefly, simultaneous imaging of calcium and intrinsic signals *in vivo* was performed using a macro zoom fluorescence microscope (MVX-10, Olympus, Tokyo, Japan) or an upright fluorescence microscope (ECLIPSE Ni-U, Nikon, Tokyo, Japan), equipped with a 1x objective. A 625 nm LED light source was used to obtain intrinsic signals, which we referred to as the hemodynamic signal (Vanni and Murphy 2014). At this wavelength, the optical intrinsic signal primarily reflects deoxyhemoglobin signal (HbR) (Ma et al. 2016). GCaMP was excited by a 100 W mercury lamp through a GFP mirror unit (Olympus). Intrinsic signal data was collected at a frame rate of 5 Hz using a CCD camera (1,000m; Adimec, Boston, MA, U.S.A.) and calcium signal data was collected at a frame rate of 10 Hz using a CCD camera (DS-Qi1 Mc; Nikon). The emission filters were 625 nm long pass (SC-60, Fuji film, Tokyo, Japan) for intrinsic signals, and 505-535 nm band pass (FF01-520/35-25, Semrock, Lake Forest, Illinois) for calcium signals. Data were acquired for 30-60 min per animal (5 min per scan).

Data Preprocessing

All data analyses were conducted in Matlab (MathWorks, Natick, MA) using a method

described previously (Matsui et al. 2016). Briefly, all the image frames were corrected

for possible within-scan motion by rigid-body transformation. Calcium and

hemodynamic images were then coregistered by rigid-body transformation using

manually selected anatomical landmarks that were visible in both images (e.g.,

branching points of blood vessels). All of the images were then spatially down-sampled

by a factor of two. Pixels within the cortex (at this point including large blood vessels

including the sinus) were extracted manually. For both calcium and hemodynamics,

slow drift in each pixel's time course was removed using a high-pass filter (> 0.01 Hz,

second order Butterworth. No low-pass filter was used). After filtering, each pixel's

time course was normalized by subtracting the mean across time and then dividing by

the standard deviation across time. Global signal regression was conducted by

regressing out the time course of average signal within the brain from each pixel's time

course. Finally, hemodynamic signal was multiplied by -1 to set the polarity of the

activity change equal to that in the calcium signal.

In some analyses, the calcium signal was further preprocessed. To obtain the high

frequency calcium signal, an additional high-pass filter (> 0.1 Hz) followed by a median

filter was applied. The median filter was applied as follows: For each frame, we defined

a time-window (width = 200 frames) whose center was positioned at the frame. Then,

the signal of the frame was replaced by the median of the time-window [B(t) =

median(A(t-100), …, A(t+100)); where A(k) denotes the original signal at frame k and

B(k) denotes median filtered signal at frame k]. To obtain the low frequency calcium

signal, an additional low-pass filter (< 0.1 Hz) was applied.

Extraction of Region-of-Interest (ROI) Time Courses

Selection of ROI and time courses are conducted as described previously (Matsui et al.

2016). Briefly, 38 cortical regions (19 for each hemisphere) were selected as ROIs

based on a previous mouse functional connectivity study (White et al. 2011)

(Supplementary Fig. 1). Each ROI was a 6×6 pixel square (0.5 mm $\times$ 0.5 mm)

centered at a selected coordinate. The time course for each ROI was calculated by

averaging the time courses of pixels within the ROI that corresponded to gray matter.

ROIs located outside of the FOV were discarded.

Analysis of FC

For both calcium and hemodynamic signals, FC was calculated using a standard
seed-based correlation method (Matsui et al. 2016). First, the correlation coefficient
between the time course of a selected ROI ("seed time course") and the time course of
every pixel within the brain was calculated. Second, FC values were averaged across
scans to obtain FC values for each pixel. The spatial correlation between FC maps of
calcium and hemodynamic signals was calculated by taking the pixel-by-pixel
correlation coefficient between the two maps using all the gray matter pixels. FC with
short time window was obtained by taking correlation coefficient using all the frames
within a 30-sec window. Steps of 3 sec and 30 sec were used for the sliding window and
non-overlapping window, respectively. Scan-shifted control was calculated by shifting
the scan number of hemodynamics data relative to simultaneously obtained calcium
data.

Analysis of Co-Activation Patterns (CAPs)

CAP analysis was adopted from previous fMRI studies (Liu and Duyn 2013). Briefly,
calcium time course from each ROI was z-normalized. CAPs were calculated for each
ROIs. Frames corresponding to large peaks (> 2 s.d.) of the time course of a given ROI
were considered CAPs. We examined if CAPs calculated using calcium signal could
predict sliding-window FC calculated using hemodynamic signal. For each ROI and
each time window, we quantified the similarity between the sliding window FC and
CAPs by calculating spatial correlation between the FC map and CAPs. To quantify
coordinated temporal variations in CAPs and FCs, we also calculated temporal
deviation of CAP and FC from the mean patterns as described in the followings. First,
we calculated average patterns of CAPs and FCs in a given scan. For a given scan and a
ROI_i, average of all CAPs was calculated using the entire scan$(CAP_{scan})_i$. Similarly, a
FC map was calculated for the same scan and the same ROI [$(FC_{scan})_i$]. We repeated this
procedure to obtain ROI-by-ROI matrices of CAP_{scan} and FC_{scan}. Next, we calculated
ROI-by-ROI matrices of CAPs and FCs in a short time window (CAP_{window} and
FC_{window}) using the same procedure used to calculate CAP_{scan} and FC_{scan} but for each 30
sec time window. Then difference between the CAP_{window} and CAP_{scan} was taken to

quantify the deviation of CAP in a time window from the mean CAP pattern in the entire scan [$\Delta CAP = CAP_{window} - CAP_{scan}$]. Similarly, ΔFC was obtained by subtracting FC_{scan} from FC_{window}. Finally, correlation coefficient between non-diagonal elements of the ROI-by-ROI matrices of ΔCAP and ΔFC were calculated. When CAPs were absent for a particular ROI in a time window, that ROI was omitted from the calculation for the time window.

Cluster Analysis and Kurtosis Analysis

For the state analysis of sliding window FC, we adopted the k-means clustering algorithm used in the previous studies (Allen et al. 2014; Laumann et al. 2016). Correlation distance (1-r) was used to compute the separation between each window's FC-matrix (using all 38 ROIs) and the k-means clustering was iterated 100 times with random centroid positions to avoid local minima. The windowed FC-matrices were mean-centered by scan to eliminate scan-level and subject-level features from contributing the clustering result. K-means clustering was applied in the same manner to the simulated data that was matched in size to the real data. The cluster validity index was used to evaluate the quality of clustering for the range of cluster numbers (k = 2-10). The cluster validity index was computed as the average ratio of within-cluster distance to between-cluster distance.

Non-stationarity of spontaneous neuronal signal correlation was assessed by calculating multivariate kurtosis using the same procedure as described by Laumann and colleagues (Laumann et al. 2016). One value of kurtosis was calculated for each FC-matrix (using all 38 ROIs) obtained each scan. The same procedure was applied to the simulated data that was matched in size to the real data. Significant difference between the kurtosis measure of the real and the simulated data indicates either non-stationarity of FC or non-gaussianity of the signal or both.

Time Course Simulation

To obtain a null dataset to evaluate the non-stationarity of the real data, we constructed simulated time courses using a method developed by Laumann and colleagues (Laumann et al. 2016). Briefly, random normal deviates having the same dimensionality as a real dataset are sampled. These time courses are multiplied in the spectral domain by the average power spectrum of the (bandpass filtered) real data. These time courses

are then projected onto the eigenvectors derived from the covariance matrix of the real

data. This procedure produces simulated data that are stationary by construction but

matched to real data in the covariance structure and mean spectral content.

Assessment of motion-related and physiological artifacts

To estimate the strength of motion-related and physiological artifacts from functional

images, we calculated DVARS using the hemodynamic signal with a procedure

described in fMRI literatures (Power et al. 2012; Power et al. 2014). Specifically, for

each scan, after the preprocessing, root mean squares of the temporal derivatives of the

hemodynamic time courses were calculated and averaged across ROIs to obtain one

time course of DVARS. To remove the data potentially contaminated by the artifact, we

conducted two types of analyses. First, for each scan, the histogram of DVARS was

calculated to exclude scans with strong skewness (Supplementary Fig. 3)

(scan-censoring). To define the DVARS histogram with strong skewness, we calculated

DVARS histograms for 1000 sets of simulation described in the preceding section. If the

skewness of the data was larger than the 99^{th} percentiles of the simulation, the scan was

considered to be strongly skewed. Second, frame-censoring was conducted at multiple

DVARS thresholds as described in the fMRI literatures (Power et al. 2012; Power et al.

2014). For a given threshold, frames with DVARS larger than the threshold were simply

discarded from the subsequent analyses.

Assessment of the normality of the signal

To check for the normality of the hemodynamic and the calcium signals, we computed

the 4^{th} moment of the signal distribution for each scan. For each scan, signals in all the

ROIs were concatenated before calculating the 4^{th} moment. Similar results were

obtained when the 4^{th} moment was calculated separately for individual ROIs and then

the mean of the 4^{th} moment was obtained for each scan (data not shown). The calculated

4^{th} moment of the real data was then compared with that of the 1000 simulated time

courses obtained by the procedure described above. If the 4^{th} moment of the real data

was larger than the 99^{th} percentile of the simulation, the signal in the scan was

considered to be non-normal. In the case of calcium signal, we applied median filter to

remove high frequency noise and enforce normality of the signal. Calcium time courses

often contained some high frequency noise on top of the slower calcium activity (Fig.

1A), which was most likely due to the moderate level of excitation light adjusted to

avoid bleaching of GCaMP fluorophore. We consider that the median filter removed

these high frequency noise while retaining slower calcium signal that was

approximately normally distributed.

Results

Consistent FC dynamics in calcium and hemodynamic signals

Transgenic mice expressing GCaMP in neocortical neurons were used to simultaneously measure neuronal calcium signal and hemodynamics in a large portion of bilateral neocortex (Fig. 1A) (Matsui et al. 2016). Mice were lightly anesthetized and head-fixed with metal plates so that head-motion could not contaminate the signals. For both calcium and hemodynamic signals, power spectra of the signals exhibited approximately linear trend on log-log plots (Supplementary Fig. 2) suggesting that the non-neuronal artifact was small. We used sliding window correlation (30 sec window at 3 sec steps) to examine if calcium and hemodynamic FC in mice exhibited "dynamic" changes. Consistent with previous reports in humans (Allen et al. 2014; Chang and Glover 2010; Zalesky et al. 2014) and other animals (Hutchison et al. 2014; Majeed et al. 2009), FC between pairs of ROIs calculated with sliding windows showed considerable variability over different time points both in calcium signal and hemodynamics (Fig. 1B-C). Consistent with the idea that variability in hemodynamic FC arises from underlying neuronal activity, we found close matches between dFC of calcium and hemodynamics (correlation coefficients, 0.631 and 0.675 for Figs. 1B and 1C, respectively). Correlation between the time courses of calcium FC and hemodynamic FC was significantly larger for the data than that of the scan-shifted control ($P < 10^{-20}$, Kolmogorov-Smirnov test; Fig. 1D).

To further examine the consistency between dFC in calcium and hemodynamics in the entire neocortex, we calculated calcium and hemodynamic FC among all pairs of ROIs and compared them across time windows (Fig. 2A-B). The ROI-based FC-matrices in calcium and hemodynamics both showed variability across time windows. On the other hand, FC matrices in calcium and hemodynamics within each time window were similar. If dFC in calcium and hemodynamics were matched, the similarity between calcium and hemodynamic FC in the same time window should be higher than that calculated using different time windows (e.g., similarity between Ca-FC$_{window\#1}$ and Hemo-FC$_{window\#1}$ would be higher than the similarity between Ca-FC$_{window\#1}$ and Hemo-FC$_{window\#2}$). Otherwise, the similarity between FC-matrices in calcium and hemodynamics merely reflects the overall similarity of FC in calcium and hemodynamics but not the coordinated "dynamics" of calcium and hemodynamic FC. Across all the data, we found that the distribution of the correlation coefficient between

the FC-matrices in calcium and hemodynamics was shifted toward positive values
compared with that calculated with the scan-shifted data (P < 10^{-14},
Kolmogorov-Smirnov test; Fig. 2C). The difference between the real data and the
trial-shifted data was also consistently positive across animals (p < 0.0156, n = 7 mice,
sign-rank test; Fig. 2D) and was seen across various window sizes ranging from 1 sec to
60 sec (Fig. 2E). Together these results suggest that temporal variability in
hemodynamic FC, as measured with sliding window, arises from neural activity rather
than from movement-related artifacts (Laumann et al. 2016) or non-neuronal
physiological artifacts such as heartbeat and respiration (Bianciardi et al. 2009; Shmueli
et al. 2007).

Variations in transient neuronal coactivations explained variations in FC
What are the potential neuronal events that create dFC? Recent fMRI studies proposed
that variability in the neuronal coactivation pattern (CAP) of brain areas is directly
reflected in the "dynamic" change of FC observed with the sliding window correlation
(Liu and Duyn 2013). To address this possibility, for each scan, we compared sliding
window FC calculated using hemodynamics with the CAPs calculated using calcium
signal. The use of calcium signal for extracting CAPs allowed us to capture faster
spatiotemporal dynamics than the hemodynamics. More importantly, the use of two
different signals also allowed us to avoid comparing sliding window FC and CAPs that
were derived from the same signals and could lead to circular logic.

For each anatomical ROI, we first detected peaks in the calcium signal within a
given time-window and then defined CAPs as the frames in the calcium signal
corresponding to the detected peak locations (Fig. 3A) (Liu and Duyn 2013). Similar to
the previous reports in fMRI (Liu et al. 2013; Liu and Duyn 2013), we found variations
in the spatial patterns of CAPs extracted from the same ROI (Fig. 3A, panels above time
courses). We used CAPs to examine if variations of the spatial pattern of CAPs in
different time windows could explain the spatial variation of sliding window FC. For
each ROI in each 30 sec window, we extracted CAPs and FC using calcium and
hemodynamic signals, respectively. In the example 30 sec windows shown in Figure 3A,
time courses of the chosen ROI showed transient activations that resulted in 11 and 3
frames of CAPs (corresponding to 1.1 and 0.3 sec of data, respectively). Despite the
small number of frames corresponding to CAPs, the average spatial pattern of CAPs in

the time window closely matched the spatial pattern of hemodynamic FC calculated in

the same time window (compare mean CAP and mean FC in Fig. 3A).

To further compare CAPs with sliding window hemodynamic FC across ROIs, we

calculated CAPs for all pairs of ROIs and compared them with FC of the same

ROI-pairs in the same time window (Fig. 3B). Across all the data, CAP-matrices and

FC matrices showed high positive correlation (Fig. 3C-D; mean R = 0.525 across

animals) suggesting that CAP and FC calculated using the same sliding window were

similar.

The similarity between calcium CAP and hemodynamic FC in short time window

does not necessarily indicate coordinated temporal variation between CAP and FC, but

could result entirely from similarity between the time-average patterns of CAP and FC.

Therefore, to further examine if coordinated temporal variations in CAPs and FCs exist,

we calculated ΔCAP and ΔFC by subtracting from each CAP and FC in each time

window the average pattern of CAP and FC, respectively, that were calculated using the

entire scan (Fig. 4A). Coordinated change in ΔCAP and ΔFC indicates similar temporal

fluctuation of CAP and sliding-window FC that cannot be accounted for by the

similarity in the mean pattern of CAP and FC calculated using the entire scan. We found

that the distribution of the correlation between ΔCAP and ΔFC for the real data was

shifted toward positive values whereas the same distribution calculated using

trial-shifted data was centered near zero ($P < 10^{-30}$, Kolmogorov-Smirnov test; Fig. 4B).

Furthermore, the correlation between ΔCAP and ΔFC was consistently positive across

all animals ($P < 0.156$, n = 7 mice, sign rank test; Fig. 4C) and was seen across various

sizes of time-windows ranging from 1 to 60 sec (Fig. 4D). Excluding scans with

potential artifacts using DVARS (Power et al. 2012; Power et al. 2014) still yielded a

significant difference between the real and the trial-shifted data (Supplementary Fig. 9).

Taken together, these results suggest temporal fluctuations of the spatial pattern of

CAPs at least partly explain temporal fluctuations of hemodynamic FC calculated using

sliding windows.

"Dynamics" of FC arise from non-stationarity of resting-state activity

Because FC is estimated by using finite number of time-points, temporal fluctuations of

FC observed in short time-windows could arise from mere sampling error even when

underlying FC is stationary (Laumann et al. 2016). We next addressed whether the

sampling error could explain the dFC observed in the present data. We compared two

indices used in a previous study, namely cluster validity index and kurtosis, for real data

and simulated data that are matched in spectral and covariance properties (Fig. 5A)

(Laumann et al. 2016). The cluster validity index measures degree of clustering of

multiple sliding window FC calculated within the scan. Note that smaller cluster

validity index indicates more clustering (see Methods for details). For both calcium and

hemodynamic signals, we found cluster validity index of real data to be significantly

smaller than that of simulated data (Fig. 5B), suggesting that the real data had cluster

structure that could not be fully accounted for by sampling error. Similarly, we

calculated kurtosis of the covariance matrices of real and simulated data. If the kurtosis

of real data were larger than that of simulated data assuming a stationary Gaussian

process, non-stationarity is implied for the real data, if the real data is normally

distributed (Laumann et al. 2016). For the hemodynamic signal, the real data was

approximately normally distributed (Supplementary Fig. 8A). Thus, we calculated the

kurtosis using the hemodynamic signal without further preprocessing. For the calcium

signal, because the original signal was not normally distributed, we applied further

preprocessing before calculating the kurtosis. First, we applied an additional high pass

temporal filter to remove low frequencies (< 0.1 Hz). Then we used a median filter to

enforce a normality of the signal (Supplementary Fig. 6). For both hemodynamic and

calcium signals, we found that the kurtosis of the real data was significantly higher than

that of the simulated data ($P < 10^{-10}$ for calcium, sign rank test, n = 50 scans that showed

approximately normal distribution of the signal (see Methods for the details of the

assessment of normality); $P < 10^{-11}$ for hemodynamics, sign rank test, n = 64 scans; Fig.

5C, Supplementary Fig. 8B). Calcium signal at low frequency (< 0.1 Hz) was

approximately normal without additional preprocessing (Supplementary Fig. 7A). The

kurtosis of the real and the simulated data was also significantly different at this

frequency range, though the magnitude of the difference was smaller (Supplementary

Fig. 7B). Together, these results suggest that dFC arise from non-stationarity of

spontaneous neuronal activity, and analyses based on sliding window correlation have

the potential to detect non-stationarity.

To assess potential contribution of motion-related and physiological artifacts on the

analysis of the non-stationarity using kurtosis, we used DVARS calculated using the

hemodynamic signal to exclude time points for which the artifacts might have been

problematic (Power et al. 2012; Power et al. 2014). In some scans with large peaks in
the time courses of DVARS, indicating potential body movements, the histogram of
DVARS was strongly right-skewed (Supplementary Fig. 3). Analysis of the kurtosis
using a subset of scans for which DVARS distribution was not right-skewed compared
to the stationary simulation also revealed a significant difference between the real and
the simulated data ($P < 10^{-14}$, sign rank test, n = 25; Supplementary Fig. 4; see Materials
and Methods for details). Furthermore, frame-censoring at several thresholds of DVARS
revealed that the difference of kurtosis between the real and the simulated data did not
depend on the levels of DVARS thresholds (Supplementary Fig. 5). These results
suggest that the motion-related and physiological artifacts, as detected by DVARS, did
not significantly affect the present results.

Discussion

In the present study, we used simultaneous imaging of calcium and hemodynamic signals to show that temporal fluctuations in hemodynamic FC calculated in a short time window closely follow that of calcium FC, suggesting the neuronal origin of dFC. We have further shown that the spatial pattern of hemodynamic FC in a short time window is predicted by averaging transient coactivations in the calcium signal (CAPs) contained within the same time-window suggesting that temporally interspersed transient neuronal events underlie resting-state FC. Finally, we have shown that in both calcium and hemodynamic signals, statistical properties of FC calculated in a short time window was significantly different from that obtained with simulated signals that were stationary by construction. These results advocate for the analysis of the "dynamic" aspect of FC obtained in human fMRI experiments. Insights of the neuronal events underlying dFC provided by the present study would also be informative for developing appropriate analysis methods for dFC.

Relationship to previous investigations of the neuronal origin of dFC

To provide direct evidence linking neuronal activity and dFC, several groups have conducted simultaneous recording of fMRI and local field potential (LFP) (Lu et al. 2007; Pan et al. 2011; Thompson et al. 2013) or EEG (Chang et al. 2013; Tagliazucchi et al. 2012b). However, these previous studies were limited in several ways. Since LFP recordings were limited from a small number of recording sites whereas EEG recording did not have enough spatial resolution, evidence directly linking global spatial pattern of neuronal activity with hemodynamic FC has been lacking. Using simultaneous imaging of calcium and hemodynamic signals, the present study provides evidence suggesting that temporal variability of hemodynamic FC and its time-to-time spatial patterns reflect spatial patterns of large-scale neuronal activity. Moreover, since the present study used anesthetized and head-fixed mice, the results are unlikely to be attributable to head motion.

Recent human fMRI studies have proposed that neuronal activity important for FC is condensed into transient large scale neuronal coactivations (i.e. CAPs) (Liu and Duyn 2013; Tagliazucchi et al. 2012a; Tagliazucchi et al. 2011). Consistent with this idea, imaging studies in mice revealed transient neuronal coactivations across brain areas (Matsui et al. 2016; Mohajerani et al. 2013; Vanni and Murphy 2014). In our previous

study, we searched for neuronal coactivations that resembled the spatial patterns of
(static) FC and showed that such neuronal coactivations were converted into spatially
similar hemodynamic signals (Matsui et al. 2016). In the present study, we took a
different approach that was similar to single frame analysis methods employed in recent
human fMRI studies (Karahanoğlu and Van De Ville 2015; Liu et al. 2013; Liu and
Duyn 2013; Tagliazucchi et al. 2011). We were especially interested whether CAPs
represent the potential neuronal events underlying the temporal variability of sliding
window FC. Because derivation of CAPs and sliding-window FC using identical BOLD
signals could potentially lead to circular logic, in the present study, we examined the
link between CAPs and sliding-window FC derived from different signals (calcium and
hemodynamics, respectively). Instead of specifically looking at neuronal coactivations
that resembled "static" FC, we took all the individual CAPs into our analysis and
showed that variation of the spatial pattern of individual CAPs across time windows
was significantly related to variations of hemodynamic FC across time windows. Thus,
the present findings suggest importance of the development of analysis that specifically
focuses on CAPs (Karahanoğlu and Van De Ville 2015; Liu et al. 2013). However, the
modest correlation found between CAP and sliding window FC implies that the
fluctuations of calcium and hemodynamic signals from the average pattern maybe small.
It should also be noted that, although statistically significant, the correlation between
ΔCAP and ΔFC was relatively weak. Part of the reason for this could be non-neuronal
physiological noise that contributed to hemodynamics (Matsui et al. 2016). In the
present study, because of the use of anesthesia and head-fixation, head motion is
unlikely to be the primary source of the non-neuronal noise. However, other
physiological activities, e.g. respiration and heartbeat, are known to affect
hemodynamics (Chang et al. 2009; Chang and Glover 2009) and, thus, likely to affect
temporal fluctuation of hemodynamic FC as well. Although scan-wise data exclusion
using DVARS suggested that the motion-contamination was not the major cause of the
observed temporal correlation between ΔCAP and ΔFC (Supplementary Fig. 9), the
common artifacts on hemodynamic and calcium signals could have contributed to the
observed temporal correlation. Our results (i.e. relatively low correlation between
ΔCAP and ΔFC) indicate that correction for such non-neuronal physiological noise
(Glover et al. 2000) is likely to be essential for the analysis of dFC.
482

Non-stationarity of spontaneous brain activity correlation

It has been of a matter of debate to what extent temporal fluctuations of FC is

attributed to the dynamics of underlying neuronal activity but not to non-neuronal

sources of noise (e.g., head motion, sampling variability; reviewed in (Hutchison et al.

2013)). Laumann and colleagues have reported that most of the temporal fluctuations of

single subject FC is explained by head motion (Laumann et al. 2016). After controlling

for the head motion, Laumann and colleagues have concluded that statistical properties

of resting-state FC in human fMRI is indistinguishable from those obtained with

simulated signals that are stationary by construction. A similar study by Hindriks and

colleagues has also indicated the apparent dFC calculated with the sliding window

method does not necessarily indicate non-stationarity of resting brain network (Hindriks

et al. 2016). However, in terms of spontaneous neuronal activity itself, there is

substantial evidences showing that spontaneous neuronal activity is non-stationary

(Foster and Wilson 2006; Ji and Wilson 2007; Logothetis et al. 2012). In particular,

under both awake and anesthetized states, transient neuronal events such as

sharp-wave-ripples have been shown to produce coordinated activity across the entire

brain (Logothetis et al. 2012). The present results are consistent with these previous

studies supporting the non-stationarity of neuronal activity, and further showed that FC

calculated using such non-stationary neuronal activity also showed non-stationarity, as

expected.

Potential contribution of arousal

Fluctuation in the level of arousal has been shown to contribute to apparent "dynamics"

and non-stationarity of FC (Laumann et al. 2016; Tagliazucchi and Laufs 2014). The

present study observed larger temporal variability of hemodynamic FC in anesthetized

mice than in awake, eyes open-fixated humans (Laumann et al. 2016). Interestingly, a

recent study reports large within-subject FC variability in subjects instructed to close

their eyes (Pannunzi et al. 2017), which predisposes to sleep (Tagliazucchi and Laufs

2014). In the present study, it could be possible that fluctuating level of anesthesia,

instead of the subject's sleep state, could have contributed to the greater variability of

FC. Alternatively, rapid fluctuation between awake and sleep states in mice (Adler et al.

2014), could have contributed to the greater variability of FC compared to humans.

According to the results of the analysis using power spectra (Supplementary Fig. 2) and

DVARS (Supplementary Figs. 3-5), we consider that the level of arousal rarely reached

a point at which the animal started to struggle. Nevertheless, the present data alone was

not sufficient to exclude the possibility that the fluctuation of the level of arousal

accounted for a large fraction of the FC "dynamics" and non-stationarity observed here.

Future experiments conducting simultaneous recordings of functional images and

physiological signals (e.g. electroencephalogram, respiration-rates) in mice would be

able to assess the exact amount of non-stationarity in FC under a defined state of

arousal.

Other limitations of the study

It should be clearly stated that the present results do not guarantee that sliding window

methods are always capable of detecting non-stationarity in human resting-state fMRI

data. The present study used tightly head-restrained animals and high signal to

noise-ration (SNR) imaging at a high frame rate (5 and 10 Hz for hemodynamics and

calcium signal, respectively). Compared to the present experimental conditions, overall

SNR in typical human resting-state fMRI is likely to be substantially compromised.

Under such low SNR conditions, it is not clear whether simple sliding window

correlation methods can detect the non-stationarity of FC (Hindriks et al. 2016;

Laumann et al. 2016). With respect to SNR, we expect that the recent development of

high-speed fMRI (Feinberg et al. 2010) will significantly improve the detectability of

non-stationarity. Nevertheless, the present results suggest that, rather than the sliding

window based method, an alternative analysis strategy that directly extracts CAPs from

hemodynamic signals (Karahanoğlu and Van De Ville 2015; Liu et al. 2013; Liu and

Duyn 2013; Tagliazucchi et al. 2012a) may be more appropriate for extracting relevant

information related to dFC. Care should be taken, however, since the smaller brain and

the low dimensionality of FC in the mouse (compared to the human) could also have

made CAPs from just a few frames look similar to those of FC.

It should also be noted that the present results do not claim that dFC has significant

behavioral or cognitive consequences. Instead of examining the potential relationship

between dFC and cognitive dynamics or behavioral variability [see for recent review

(Preti et al. 2016)], here we focused on validating the neuronal origin of dFC.

Experiments under anesthesia greatly reduced potential confounding factors, such as

head motion and arousal state (Hutchison et al. 2014; Laumann et al. 2016).

Nevertheless, the present wide-field imaging setup can be naturally extended to awake

imaging with task-performing mice (Ferezou et al. 2007; Wekselblatt et al. 2016). Such

experiments would reveal the potential consequences of the dFC on its behavioral

outcome.

Acknowledgements

We thank A. Honda and Y. Sono for animal care and genotyping; the Research Support Center, Graduate School of Medical Sciences, Kyushu University, for technical support: and K. Jimura and Y. Noro for helpful discussion. Support for this work was provided by grants from Brain Mapping by Integrated Neurotechnologies for Disease Studies (Brain/MINDS) – Japan Agency for Medical Research and Development (AMED), Core Research for Evolutionary Science and Technology (CREST) – AMED and Strategic International Research Cooperative Program (SCIP) – AMED, and Japan Society for Promotions of Sciences (JSPS) KAKENHI Grants 25221001 and 25117004 (to K.O.), World Premium Institute (WPI), JSPS (to K.O.); JSPS KAKENHI Grant 17K14931 (to T. Matsui) and JSPS Research Fellowship 20153597 (to T. Murakami).

References

Adachi Y, Osada T, Sporns O, Watanabe T, Matsui T, Miyamoto K, Miyashita Y. 2012. Functional connectivity between anatomically unconnected areas is shaped by collective network-level effects in the macaque cortex. Cereb Cortex. 22(7):1586-1592.

Adler DA, Ammanuel S, Lei J, Dada T, Borbiev T, Johnston MV, Kadam SD, Burd I. 2014. Circadian cycle-dependent eeg biomarkers of pathogenicity in adult mice following prenatal exposure to in utero inflammation. Neuroscience. 275:305-313.

Allen EA, Damaraju E, Plis SM, Erhardt EB, Eichele T, Calhoun VD. 2014. Tracking whole-brain connectivity dynamics in the resting state. Cereb Cortex. 24(3):663-676.

Bianciardi M, Fukunaga M, van Gelderen P, Horovitz SG, de Zwart JA, Shmueli K, Duyn JH. 2009. Sources of functional magnetic resonance imaging signal fluctuations in the human brain at rest: A 7 t study. Magn Reson Imaging. 27(8):1019-1029.

Biswal B, Yetkin FZ, Haughton VM, Hyde JS. 1995. Functional connectivity in the motor cortex of resting human brain using echo-planar mri. Magn Reson Med. 34(4):537-541.

Calhoun VD, Miller R, Pearlson G, Adalı T. 2014. The chronnectome: Time-varying connectivity networks as the next frontier in fmri data discovery. Neuron. 84(2):262-274.

Chang C, Cunningham JP, Glover GH. 2009. Influence of heart rate on the bold signal: The cardiac response function. Neuroimage. 44(3):857-869.

Chang C, Glover GH. 2009. Relationship between respiration, end-tidal co2, and bold signals in resting-state fmri. Neuroimage. 47(4):1381-1393.

Chang C, Glover GH. 2010. Time-frequency dynamics of resting-state brain connectivity measured with fmri. Neuroimage. 50(1):81-98.

Chang C, Liu Z, Chen MC, Liu X, Duyn JH. 2013. Eeg correlates of time-varying bold functional connectivity. Neuroimage. 72:227-236.

Deco G, Ponce-Alvarez A, Mantini D, Romani GL, Hagmann P, Corbetta M. 2013. Resting-state functional connectivity emerges from structurally

and dynamically shaped slow linear fluctuations. J Neurosci.
33(27):11239-11252.
Feinberg DA, Moeller S, Smith SM, Auerbach E, Ramanna S, Gunther M,
Glasser MF, Miller KL, Ugurbil K, Yacoub E. 2010. Multiplexed echo
planar imaging for sub-second whole brain fmri and fast diffusion
imaging. PLoS One. 5(12):e15710.
Ferezou I, Haiss F, Gentet LJ, Aronoff R, Weber B, Petersen CC. 2007.
Spatiotemporal dynamics of cortical sensorimotor integration in
behaving mice. Neuron. 56(5):907-923.
Foster DJ, Wilson MA. 2006. Reverse replay of behavioural sequences in
hippocampal place cells during the awake state. Nature.
440(7084):680-683.
Fox MD, Raichle ME. 2007. Spontaneous fluctuations in brain activity
observed with functional magnetic resonance imaging. Nat Rev
Neurosci. 8(9):700-711.
Fox MD, Snyder AZ, Vincent JL, Corbetta M, Van Essen DC, Raichle ME.
2005. The human brain is intrinsically organized into dynamic,
anticorrelated functional networks. Proc Natl Acad Sci U S A.
102(27):9673-9678.
Glover GH, Li TQ, Ress D. 2000. Image-based method for retrospective
correction of physiological motion effects in fmri: Retroicor. Magn
Reson Med. 44(1):162-167.
Hansen EC, Battaglia D, Spiegler A, Deco G, Jirsa VK. 2015. Functional
connectivity dynamics: Modeling the switching behavior of the resting
state. Neuroimage. 105:525-535.
Hindriks R, Adhikari MH, Murayama Y, Ganzetti M, Mantini D, Logothetis
NK, Deco G. 2016. Can sliding-window correlations reveal dynamic
functional connectivity in resting-state fmri? Neuroimage.
127:242-256.
Honey CJ, Sporns O, Cammoun L, Gigandet X, Thiran JP, Meuli R,
Hagmann P. 2009. Predicting human resting-state functional
connectivity from structural connectivity. Proc Natl Acad Sci U S A.
106(6):2035-2040.

Hutchison RM, Hutchison M, Manning KY, Menon RS, Everling S. 2014.
Isoflurane induces dose-dependent alterations in the cortical
connectivity profiles and dynamic properties of the brain's functional
architecture. Hum Brain Mapp. 35(12):5754-5775.
Hutchison RM, Womelsdorf T, Allen EA, Bandettini PA, Calhoun VD,
Corbetta M, Della Penna S, Duyn JH, Glover GH, Gonzalez-Castillo J
et al. 2013. Dynamic functional connectivity: Promise, issues, and
interpretations. Neuroimage. 80:360-378.
Ji D, Wilson MA. 2007. Coordinated memory replay in the visual cortex and
hippocampus during sleep. Nat Neurosci. 10(1):100-107.
Karahanoğlu FI, Van De Ville D. 2015. Transient brain activity disentangles
fmri resting-state dynamics in terms of spatially and temporally
overlapping networks. Nat Commun. 6:7751.
Laumann TO, Snyder AZ, Mitra A, Gordon EM, Gratton C, Adeyemo B,
Gilmore AW, Nelson SM, Berg JJ, Greene DJ et al. 2016. On the
stability of bold fmri correlations. Cereb Cortex.
Liu X, Chang C, Duyn JH. 2013. Decomposition of spontaneous brain
activity into distinct fmri co-activation patterns. Front Syst Neurosci.
7:101.
Liu X, Duyn JH. 2013. Time-varying functional network information
extracted from brief instances of spontaneous brain activity. Proc Natl
Acad Sci U S A. 110(11):4392-4397.
Liégeois R, Laumann TO, Snyder AZ, Zhou J, Yeo BTT. 2017. Interpreting
temporal fluctuations in resting-state functional connectivity mri.
Neuroimage. 163:437-455.
Logothetis NK, Eschenko O, Murayama Y, Augath M, Steudel T, Evrard HC,
Besserve M, Oeltermann A. 2012. Hippocampal-cortical interaction
during periods of subcortical silence. Nature. 491(7425):547-553.
Lu H, Zuo Y, Gu H, Waltz JA, Zhan W, Scholl CA, Rea W, Yang Y, Stein EA.
2007. Synchronized delta oscillations correlate with the resting-state
functional mri signal. Proc Natl Acad Sci U S A. 104(46):18265-18269.
Ma Y, Shaik MA, Kim SH, Kozberg MG, Thibodeaux DN, Zhao HT, Yu H,
Hillman EM. 2016. Wide-field optical mapping of neural activity and

brain haemodynamics: Considerations and novel approaches. Philos
Trans R Soc Lond B Biol Sci. 371(1705).
Majeed W, Magnuson M, Keilholz SD. 2009. Spatiotemporal dynamics of low
frequency fluctuations in bold fmri of the rat. J Magn Reson Imaging.
30(2):384-393.
Matsui T, Koyano KW, Tamura K, Osada T, Adachi Y, Miyamoto K, Chikazoe
J, Kamigaki T, Miyashita Y. 2012. Fmri activity in the macaque
cerebellum evoked by intracortical microstimulation of the primary
somatosensory cortex: Evidence for polysynaptic propagation. PLoS
One. 7(10):e47515.
Matsui T, Murakami T, Ohki K. 2016. Transient neuronal coactivations
embedded in globally propagating waves underlie resting-state
functional connectivity. Proc Natl Acad Sci U S A. 113(23):6556-6561.
Matsui T, Tamura K, Koyano KW, Takeuchi D, Adachi Y, Osada T, Miyashita
Y. 2011. Direct comparison of spontaneous functional connectivity and
effective connectivity measured by intracortical microstimulation: An
fmri study in macaque monkeys. Cereb Cortex. 21(10):2348-2356.
Messé A, Rudrauf D, Benali H, Marrelec G. 2014. Relating structure and
function in the human brain: Relative contributions of anatomy,
stationary dynamics, and non-stationarities. PLoS Comput Biol.
10(3):e1003530.
Mohajerani MH, Chan AW, Mohsenvand M, LeDue J, Liu R, McVea DA,
Boyd JD, Wang YT, Reimers M, Murphy TH. 2013. Spontaneous
cortical activity alternates between motifs defined by regional axonal
projections. Nat Neurosci. 16(10):1426-1435.
Murakami T, Matsui T, Ohki K. 2017. Functional segregation and
development of mouse higher visual areas. J Neurosci.
37(39):9424-9437.
Murakami T, Yoshida T, Matsui T, Ohki K. 2015. Wide-field ca(2+) imaging
reveals visually evoked activity in the retrosplenial area. Front Mol
Neurosci. 8:20.
Pan WJ, Thompson G, Magnuson M, Majeed W, Jaeger D, Keilholz S. 2011.
Broadband local field potentials correlate with spontaneous

fluctuations in functional magnetic resonance imaging signals in the
rat somatosensory cortex under isoflurane anesthesia. Brain Connect.
1(2):119-131.
Pannunzi M, Hindriks R, Bettinardi RG, Wenger E, Lisofsky N, Martensson
J, Butler O, Filevich E, Becker M, Lochstet M et al. 2017.
Resting-state fmri correlations: From link-wise unreliability to whole
brain stability. Neuroimage. 157:250-262.
Power JD, Barnes KA, Snyder AZ, Schlaggar BL, Petersen SE. 2012.
Spurious but systematic correlations in functional connectivity mri
networks arise from subject motion. Neuroimage. 59(3):2142-2154.
Power JD, Mitra A, Laumann TO, Snyder AZ, Schlaggar BL, Petersen SE.
2014. Methods to detect, characterize, and remove motion artifact in
resting state fmri. Neuroimage. 84:320-341.
Preti MG, Bolton TA, Van De Ville D. 2016. The dynamic functional
connectome: State-of-the-art and perspectives. Neuroimage.
Shmueli K, van Gelderen P, de Zwart JA, Horovitz SG, Fukunaga M, Jansma
JM, Duyn JH. 2007. Low-frequency fluctuations in the cardiac rate as
a source of variance in the resting-state fmri bold signal. Neuroimage.
38(2):306-320.
Tagliazucchi E, Balenzuela P, Fraiman D, Chialvo DR. 2012a. Criticality in
large-scale brain fmri dynamics unveiled by a novel point process
analysis. Front Physiol. 3:15.
Tagliazucchi E, Balenzuela P, Fraiman D, Montoya P, Chialvo DR. 2011.
Spontaneous bold event triggered averages for estimating functional
connectivity at resting state. Neurosci Lett. 488(2):158-163.
Tagliazucchi E, Laufs H. 2014. Decoding wakefulness levels from typical
fmri resting-state data reveals reliable drifts between wakefulness
and sleep. Neuron. 82(3):695-708.
Tagliazucchi E, von Wegner F, Morzelewski A, Brodbeck V, Laufs H. 2012b.
Dynamic   bold   functional   connectivity   in   humans   and   its
electrophysiological correlates. Front Hum Neurosci. 6:339.
Thompson GJ, Merritt MD, Pan WJ, Magnuson ME, Grooms JK, Jaeger D,
Keilholz SD. 2013. Neural correlates of time-varying functional

connectivity in the rat. Neuroimage. 83:826-836.

Tohmi M, Meguro R, Tsukano H, Hishida R, Shibuki K. 2014. The
extrageniculate visual pathway generates distinct response properties
in the higher visual areas of mice. Curr Biol. 24(6):587-597.

Van Dijk KR, Hedden T, Venkataraman A, Evans KC, Lazar SW, Buckner RL.
2010. Intrinsic functional connectivity as a tool for human
connectomics: Theory, properties, and optimization. J Neurophysiol.
103(1):297-321.

Vanni MP, Murphy TH. 2014. Mesoscale transcranial spontaneous activity
mapping in gcamp3 transgenic mice reveals extensive reciprocal
connections between areas of somatomotor cortex. J Neurosci.
34(48):15931-15946.

Vincent JL, Patel GH, Fox MD, Snyder AZ, Baker JT, Van Essen DC, Zempel
JM, Snyder LH, Corbetta M, Raichle ME. 2007. Intrinsic functional
architecture in the anaesthetized monkey brain. Nature.
447(7140):83-86.

Wekselblatt JB, Flister ED, Piscopo DM, Niell CM. 2016. Large-scale
imaging of cortical dynamics during sensory perception and behavior.
J Neurophysiol. 115(6):2852-2866.

White BR, Bauer AQ, Snyder AZ, Schlaggar BL, Lee JM, Culver JP. 2011.
Imaging of functional connectivity in the mouse brain. PLoS One.
6(1):e16322.

Zalesky A, Fornito A, Cocchi L, Gollo LL, Breakspear M. 2014. Time-resolved
resting-state brain networks. Proc Natl Acad Sci U S A.
111(28):10341-10346.

Zariwala HA, Borghuis BG, Hoogland TM, Madisen L, Tian L, De Zeeuw CI,
Zeng H, Looger LL, Svoboda K, Chen TW. 2012. A cre-dependent
gcamp3 reporter mouse for neuronal imaging in vivo. J Neurosci.
32(9):3131-3141.

Figure Captions

Figure 1. Representative dynamics of simultaneously observed calcium and hemodynamic FC. (A) Experimental setup. The left most panel shows the setup for simultaneous imaging. Right side shows example calcium time courses for two ROIs (green and cyan traces indicate M1 and V1 ROIs, respectively. Positions of the ROIs indicated in the example field of view. See Supplementary Figure 1 for abbreviations). FC with short time window uses subset of frames contained in short (30 sec) windows (red dotted squares). Sliding FC for hemodynamic signal was carried out similarly. **(B)-(C)** Examples dynamics of calcium and hemodynamic FC. (B) FC between right V1 and right AC. (C) FC between left M1 and left pPar. See Supplementary Figure 1 for ROI positions and abbreviations. **(D)** Histogram of correlation between time courses of Ca-FC and Hemo-FC for the data (solid line) and the scan-shifted control (dotted line). Data from all pairs of ROIs for all scans obtained in all mice were used.

Figure 2. Significant relationship between calcium and hemodynamic FC calculated in short time windows. (A)-(B) Example ROI-by-ROI FC matrices for calcium and hemodynamics for different (non-overlapping) 30 sec windows. FC matrices were similar for calcium and hemodynamics in the same time window, but not across different time windows. (A) and (B) are two different examples from different animals. **(C)** Cumulative histogram of correlation between FC matrices for calcium and hemodynamics. Dotted line indicates trial-shifted control. **(D)** Correlation between FC matrices for calcium and hemodynamics was larger for the data than for the trial-shifted control significantly across animals. **(E)** Correlation between FC matrices of calcium and hemodynamics was larger for the data than the trial-shifted control across different window-sizes (1, 2, 3, 5, 6, 10, 12, 15, 20, 30 and 60 sec). Error bars indicate s.e.m. across animals (n = 7).

Figure 3. Comparison of calcium CAPs and hemodynamic FC across time-windows. (A) Procedure for detection of CAPs in calcium signal. For a given ROI, a calcium time course was extracted and z normalized (green time courses). Then, peaks exceeding 2 s.d. (red dots) were extracted. The frames corresponding to the peaks were considered CAPs (panels above the time courses). For each window, CAPs in calcium signal were averaged to obtain mean calcium CAP. Hemodynamic CAPs were

calculated similarly (see Methods). Maps of Ca-FC and Hemo-FC were also calculated

using the same time window. **(B)** Schematics to show the procedure of comparing

calcium CAP and Hemo-FC across all ROI pairs in each time window. In each 30 sec

time window, mean calcium CAPs and Hemo-FC maps were calculated for all ROIs as

seeds (left). Then, for each seed-ROI j, calcium CAP and Hemo-FC values in ROI i

were extracted to obtain a pair of CAP-FC values for the ROI-pairs (i, j) (middle).

Finally, for each time-window, CAP-FC values were compared across all pairs of ROIs

(right). **(C)** Histograms of correlation between CAP and Hemo-FC for all time windows

across all animals. Vertical line indicates mean across time windows. **(D)** Mean

correlation between CAP and Hemo-FC across animals. Note that (C) shows entire

distribution of the data whereas (D) shows reproducibility across mice. Error bar

indicates s.e.m. across animals (n = 7).

Figure 4. Temporal fluctuations in calcium CAPs and Hemo-FC was significantly

related. (A) Schematics of the analysis. In each 30 sec time-window, mean calcium

CAP and Hemo-FC were calculated (indicated as window CAP and window FC,

respectively). From window CAP and window FC, average calcium CAP and average

Hemo-FC that were calculated using the entire scan, in which the 30-sec window

belongs to, were subtracted to obtain maps of ΔCAP and ΔFC, respectively. Finally,

values of ΔCAP and ΔFC were compared across ROI pairs similarly as in Figure 3B.

(B) Histograms of correlation between ΔCAP and ΔFC for all time windows across all

animals. Vertical lines indicate mean across time windows. Solid and dotted lines

indicate real and trial-shifted data, respectively. **(C)** Correlation between ΔCAP and

ΔFC was significantly larger for the data than for trial-shifted control across animals.

(D) Same as (C) but with different window-sizes (1, 2, 3, 5, 6, 10, 12, 15, 20, 30 and 60

sec). Error bars indicate s.e.m. across animals (n = 7).

Figure 5. Comparison with simulated data indicated non-stationarity of the real

data. (A) Examples of real and simulated time courses. Simulated time course (black)

was matched to real data (green; calcium) in mean spectral content (middle panels) and

ROI-by-ROI covariance matrix (right panels). The same procedure was applied to

create simulated hemodynamic data (not shown). **(B)** Cluster validity index calculated

for different number of clusters (k = 2-10). In both calcium (left) and hemodynamics

(right), the cluster validity index was smaller for the real (solid lines) than the simulated

data (dotted lines) indicating that the real data tended to be more clustered. **(C)** Kurtosis

of real and simulated covariance matrices. For both calcium (left) and hemodynamics

(right), multivariate kurtosis was larger for the real than for the simulated data. Error

bars indicate s.e.m. across animals (n = 7).

Supplementary Figure 1. Anatomical locations of ROIs. Anatomical locations of 19

ROIs are shown for the right hemisphere. Anatomical nomenclatures of ROIs are shown

on the right. ROIs in the left hemisphere are taken at mirror symmetric positions to

yield a total of 38 ROIs.

Supplementary Figure 2. Power spectrum of calcium and hemodynamic signals.

(A) Example power spectrum of hemodynamic signal for one mouse. Hemodynamic

signals were not temporally filtered. Black, individual scan. Red, average across scans.

(B) Average power spectrum of hemodynamic signal across mice (Red). Black,

individual mouse (averaged across scans). **(C)** Example power spectrum of calcium

signal for one mouse. Calcium signals were not temporally filtered. Black, individual

scan. Red, average across scans. **(D)** Average power spectrum of calcium signal across

mice (red). Black, individual mouse (average across scans).

Supplementary Figure 3. Example time courses and histograms of DVARS

calculated using hemodynamic signal. (A) Example time course of DVARS for one

mouse. **(B)** Histogram of DVARS for the data shown in (A). Inset, same histogram but

for different y-axis range. The histogram was not markedly right skewed (skewness =

1.60). **(C)** Another example time course of DVARS obtained in the same mouse as in

(A). In this scan, large peaks probably representing motion and/or physiological

artifacts were observed. **(D)** Histogram of DVARS calculated using data shown in (C).

Inset, same histogram but for different y-axis range. The histogram was strongly

right-skewed (skewness = 6.54).

Supplementary Figure 4. Difference between the kurtosis of real and simulated

data in Hemodynamics. Kurtosis was calculated for scans selected based on the

distribution of DVARS (n = 25). See Methods for details of the selection of the scans. *,

P < 0.0001 (sign rank test). Error bars indicate s.e.m.

Supplementary Figure 5. Difference between the kurtosis of real and simulated
data in hemodynamics as a function of DVARS threshold for frame-censoring. The
difference of the kurtosis between the real and the simulated data (blue and red,
respectively) was largely insensitive to the level of the DVARS threshold. Note that the
smaller DVARS threshold indicates stricter threshold. Error bars indicate s.e.m.

Supplementary Figure 6. Kurtosis analysis using calcium signal processed with
additional high-pass and media filters. (A) Example histogram of high frequency
calcium signal (> 0.1 Hz) in one ROI of one mouse before the application of median
filter. Red, Gaussian fit. **(B)** Same as (A) but for the signal after the application of
median filter. Red, Gaussian fit. Note better fitting with Gaussian after median filter
application, suggesting enforced normality of the signal.

Supplementary Figure 7. Kurtosis analysis using low frequency calcium signal. (A)
Example histogram of low frequency calcium signal (0.01 Hz < f < 0.1 Hz) in one ROI
of one mouse. Red, Gaussian fit. (B) Difference between the kurtosis of real and
simulated data in low frequency calcium signal for scans selected based on the
normality of the signal (55 scans). *, $P < 10^{-9}$, sign rank test. Error bars indicate s.e.m.

Supplementary Figure 8. Hemodynamic signal was approximately normally
distributed. (A) Example histogram of hemodynamic signal in one ROI of one mouse.
Red, Gaussian fit. Note that no additional preprocessing was performed. **(B)** Difference
between the kurtosis of real and simulated data in hemodynamic signal for scans
selected based on the normality of the signal (45 scans). *, $P < 10^{-8}$, sign rank test. Error
bars indicate s.e.m.

Supplementary Figure 9. Significant temporal correlation between CAP and
windowed-FC remained after removing scans with potential artifacts. (A) Same
convention as in Figure 4B but for the data using scans selected for small movement
and/or physiological artifacts based on DVARS (25 scans). *, $P < 10^{-17}$,
Kolmogorov-Smirnov test. **(B)** Same convention as in Figure 4C but for the selected

scans. Positive difference between the real and the shifted data was observed in 5 out of

7 mice.

A

CCD

625nm
HP

BP
520nm
CCD

BP
470nm

Dichroic
mirror

Mercury lamp

LED

625nm

m131003

FC with the entire scan (5 min)

FC with sliding time-windows (30 sec)

2 s.d.

50 sec

B

R = 0.631

m131027s3r7r10

Ca
Hemo

Sliding WIndow FC (R)

Time (sec)

C

R = 0.675

m131003s6r20r23

Sliding WIndow FC (R)

Time (sec)

D

data
shift

Number of ROI-pairs

Corr. b/w FC Time Courses (R)

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

1. Primary Motor Area (M1)
2. Secondary Motor Area (M2)
3. Parietal Medial Area (mPar)
4. Parietal Lateral Area (lPar)
5. Parietal Posterior Area (pPar)
6. Restrosplenial Area (Rsp)
7. Primary Visual Area (V1)
8. Medial Secondary Visual Area (mV2)
9. Lateral Secondary VIsual Area (lV2)
10. Anterior Cingulate Area (AC)
11. Frontal Association Area (FAsc)
12. Trunk Region of Primary Somatosensory Area (Tr-S1)
13. Hindlimb Region of Primary Somatosensory Area (HL-S1)
14. Shoulder Region of Primary Somatosensory Area (Sh-S1)
15. Forelimb Region of Primary Somatosensory Area (FL-S1)
16. Barrel Region of Primary Somatosensory Area (B-S1)
17. Head Region of Primary Somatosensory Area (H-S1)
18. Auditory Area (Aud)
19. Temporal Association Area (TA)

Supplementary Figure 1

Supplementary Figure 2

Supplementary Figure 3

Supplementary Figure 4

Supplementary Figure 5

A

B

Calcium Signal (z)

Supplementary Figure 6

Supplementary Figure 7

Supplementary Figure 8

Supplementary Figure 9