Neuronal origin of the temporal dynamics of spontaneous BOLD activity correlation

Teppei Matsui[1,2*], Tomonari Murakami[1,2] and Kenichi Ohki[1,2,3]

1. Department of Physiology, The University of Tokyo School of Medicine, Tokyo, Japan
2. Department of Molecular Physiology, Graduate School of Medical Sciences, Kyushu University, Fukuoka, Japan
3. International Research Center for Neurointelligence (WPI-IRCN), The University of Tokyo, Tokyo, Japan

Send Correspondence To:

Teppei Matsui, Ph.D. (Email: tematsui@m.u-tokyo.ac.jp)

Department of Physiology

The University of Tokyo School of Medicine

Tokyo 113-0033, Japan

Phone: (+81)-3-5841-3459

Abstract (199/200 words)

Resting-state functional connectivity (FC) has become a major fMRI method to study network organization of human brains. There has been recent interest in the temporal fluctuations of FC calculated using short time windows ("dynamic FC") because this method could provide information inaccessible with conventional "static" FC, which is typically calculated using the entire scan lasting several tens of minutes. Although multiple studies have revealed considerable temporal fluctuations in FC, it is still unclear whether the fluctuations of FC measured in hemodynamics reflect the dynamics of underlying neural activity. We addressed this question using simultaneous imaging of neuronal calcium and hemodynamic signals in mice and found coordinated temporal dynamics of calcium FC and hemodynamic FC measured in the same short time windows. Moreover, we found that variation in transient neuronal coactivation patterns (CAPs) was significantly related to temporal fluctuations of sliding window FC in hemodynamics. Finally, we show that the observed dynamics of FC cannot be fully accounted for by simulated data assuming stationary FC. These results provide evidence for the neuronal origin of dynamic FC and further suggest that information relevant to FC is condensed in temporally sparse events that can be extracted using a small number of time points.

Keywords: calcium imaging, BOLD, resting state, functional connectivity, stationarity

Introduction

Resting-state functional connectivity (FC) uses temporal correlation of spontaneous neuronal activity to assess network organization of brain regions in a non-invasive manner (Fox and Raichle 2007). Traditionally, FC has been calculated using all time points in a scan that typically lasts between several minutes to tens of minutes (Biswal et al. 1995; Fox et al. 2005; Van Dijk et al. 2010). Such "static" FC has been shown to largely reflect anatomical connectivity (Adachi et al. 2012; Honey et al. 2009; Matsui et al. 2012; Matsui et al. 2011; Vincent et al. 2007). Recently, in contrast to the traditional analysis of "static" FC, the temporal fluctuation of FC across short time windows is increasingly recognized as a useful aspect of FC (Allen et al. 2014; Hutchison et al. 2013; Zalesky et al. 2014). Such "dynamic" FC (dFC) calculated using short time windows could provide information that is inaccessible with static FC about the functional network organizations of healthy and diseased brains (Calhoun et al. 2014; Preti et al. 2016) [We note that the term "dynamics" refers to the non-stationarity of FC obtained with the sliding window analyses and does not refer to a process that is not invariant to temporal reordering of the samples (Liégeois et al. 2017)]. The presence of temporal fluctuations in FC has also informed theoreticians on how to constrain realistic models of brain networks (Deco et al. 2013; Hansen et al. 2015; Messé et al. 2014).

However, despite growing interest, the neurophysiological basis of dFC is still weak. Previous attempts to investigate the neural origin of dFC by simultaneous measurement of electrophysiological and functional magnetic resonance imaging (fMRI) have been limited in several ways (Lu et al. 2007; Pan et al. 2011; Tagliazucchi et al. 2012b; Thompson et al. 2013). In some studies, electrophysiological recording was limited to a small number of recording sites due to technical difficulty (Lu et al. 2007; Pan et al. 2011; Thompson et al. 2013); hence, information on the global pattern of neuronal activity was lacking. In another study, electrophysiological signals were obtained with an electroencephalogram, which records global neuronal activity but lacks precise spatial information (Tagliazucchi et al. 2012b). Thus, the link between temporal fluctuations of FC in hemodynamics and that of large-scale neuronal activity has not been adequately proven.

Several studies have also questioned whether the apparent "dynamics" of FC calculated using the sliding window method is related to temporal instability of spontaneous brain network (Hindriks et al. 2016; Laumann et al. 2016). Whereas many

studies have attributed temporal fluctuations of sliding window FC to the non-stationarity
of spontaneous neuronal activity correlation (Allen et al. 2014; Zalesky et al. 2014), some
recent studies have demonstrated that the temporal fluctuations of FC observed in the real
and the simulated data, which is stationary by construction, are statistically
indistinguishable (Hindriks et al. 2016; Laumann et al. 2016). Furthermore, Laumann and
colleagues have shown that, in the human resting state, blood oxygen level dependent
(BOLD) time series are largely stationary, discounting head-motion and fluctuating
arousal (Laumann et al. 2016). Therefore, not only the neuronal basis of dFC, but also the
existence of the statistical non-stationarity of FC, or at least the capability of sliding
window methods to detect the non-stationarity, is called into question.

In the present study, we addressed these questions using simultaneous imaging of
neuronal calcium and BOLD hemodynamic signals in the entire neocortex of transgenic
mice expressing a genetically encoded calcium indicator (Matsui et al. 2016; Vanni and
Murphy 2014; White et al. 2011). In the present experimental setup, the wide-field
calcium signal provided access to neuronal activity at higher temporal resolution and
signal-to-noise ratio compared to that of the hemodynamic signal (Matsui et al. 2016;
Murakami et al. 2017; Murakami et al. 2015; Tohmi et al. 2014; Vanni and Murphy 2014).
Moreover, unlike human fMRI data, in the present dataset, mice were tightly head-fixed
and lightly anesthetized; thus, excluding head motions from contaminating FC. The main
findings of the present study were as follows. First, we found consistency between the
"dynamics" of FC calculated using calcium and hemodynamic signals, suggesting a
neuronal origin of the temporal fluctuations of hemodynamic FC. Second, we found that
temporal fluctuations of the spatial pattern of transient neuronal coactivations as
measured in the calcium signal were significantly correlated with temporal fluctuations
of hemodynamic FC. Finally, we found that the statistical properties of sliding window
FC were significantly different between the real and the simulated data suggesting
non-stationarity of resting-state FC.

Materials and Methods

Animals

Emx1-IRES-cre and Ai38 (Zariwala et al. 2012) mice were obtained from the Jackson Laboratory (Sacramento, CA). These mice were crossed and all cortical excitatory neurons expressed GCaMP3. Mice (P60–P90) were prepared for *in vivo* wide-field simultaneous imaging. Anesthesia was induced with isoflurane (3 %) and maintained with isoflurane (1 – 2 % in surgery, 0.5 – 0.8 % during imaging) and chlorprothixene (0.3 – 0.8 mg/kg, intramuscular injection). For simultaneous imaging of calcium and hemodynamic signals, a custom-made metal head plate was attached to the skull using dental cement (Sun Medical Company, Ltd, Shiga, Japan) and a large craniotomy was made over the whole cortex. The craniotomy was sealed with 1 % agarose and a glass coverslip. During the imaging, body temperature was maintained by a heat pad. All experiments were carried out in accordance with the NIH Guide for the Care and Use of Laboratory Animals, the institutional animal welfare guidelines set forth by the Animal Care and Use Committee of Kyushu University, and the study was approved by the Ethical Committee of Kyushu University.

Simultaneous Calcium and Intrinsic Signal Imaging

The data for simultaneous imaging of calcium and hemodynamic signals was taken from a published report (Matsui et al. 2016). Briefly, simultaneous imaging of calcium and intrinsic signals *in vivo* was performed using a macro zoom fluorescence microscope (MVX-10, Olympus, Tokyo, Japan) or an upright fluorescence microscope (ECLIPSE Ni-U, Nikon, Tokyo, Japan), equipped with a 1× objective. A 625 nm LED light source was used to obtain intrinsic signals, which we referred to as the hemodynamic signal (Vanni and Murphy 2014). At this wavelength, the optical intrinsic signal primarily reflects deoxyhemoglobin signal (HbR) (Ma et al. 2016). GCaMP was excited by a 100 W mercury lamp through a GFP mirror unit (Olympus, Tokyo, Japan). Intrinsic signal data was collected at a frame rate of 5 Hz using a CCD camera (1,000m; Adimec, Boston, MA, U.S.A.) and calcium signal data was collected at a frame rate of 10 Hz using a CCD camera (DS-Qi1 Mc; Nikon, Tokyo, Japan). The emission filters were 625 nm long pass (SC-60, Fuji film, Tokyo, Japan) for intrinsic signals, and 505-535 nm band pass (FF01-520/35-25, Semrock, Lake Forest, Illinois) for calcium signals. Data were acquired for 30-60 min per animal (5 min per scan).

Data Preprocessing

All data analyses were conducted in Matlab (MathWorks, Natick, MA) using a method

described previously (Matsui et al. 2016). Briefly, all the image frames were corrected for

possible within-scan motion by rigid-body transformation. Calcium and hemodynamic

images were then coregistered by rigid-body transformation using manually selected

anatomical landmarks that were visible in both images (e.g., branching points of blood

vessels). All of the images were then spatially down-sampled by a factor of two. Pixels

within the cortex (at this point including large blood vessels including the sinus) were

extracted manually. For both calcium and hemodynamics, slow drift in each pixel's time

course was removed using a high-pass filter (> 0.01 Hz, second order Butterworth. No

low-pass filter was used). After filtering, each pixel's time course was normalized by

subtracting the mean across time and then dividing by the standard deviation across time.

Global signal regression was conducted by regressing out the time course of average

signal within the brain from each pixel's time course. Finally, hemodynamic signal was

multiplied by -1 to set the polarity of the activity change equal to that in the calcium

signal.

    In some analyses, the calcium signal was further preprocessed. To obtain the high

frequency calcium signal, an additional high-pass filter (> 0.1 Hz) followed by a median

filter was applied. The median filter was applied as follows: For each frame, we defined a

time-window (width = 200 frames) whose center was positioned on the frame. Then, the

signal of the frame was replaced by the median of the time-window [B(t) =

median(A(t-100), …, A(t+100)); where A(k) denotes the original signal at frame k and

B(k) denotes the median filtered signal at frame k]. To obtain the low frequency calcium

signal, an additional low-pass filter (< 0.1 Hz) was applied.

Extraction of Region-of-Interest (ROI) Time Courses

Selection of ROI and time courses are conducted as described previously (Matsui et al.

2016). Briefly, 38 cortical regions (19 for each hemisphere) were selected as ROIs based

on a previous mouse functional connectivity study (White et al. 2011) (Supplementary

Fig. 1). Each ROI was a 6×6 pixel square (0.5 mm $\times$ 0.5 mm) centered at a selected

coordinate. The time course for each ROI was calculated by averaging the time courses of

pixels within the ROI that corresponded to gray matter. ROIs located outside of the FOV

were discarded.

Analysis of FC

For both calcium and hemodynamic signals, FC was calculated using a standard

seed-based correlation method (Matsui et al. 2016). First, the correlation coefficient

between the time course of a selected ROI ("seed time course") and the time course of

every pixel within the brain was calculated. Second, FC values were averaged across

scans to obtain FC values for each pixel. The spatial correlation between FC maps of

calcium and hemodynamic signals was calculated by taking the pixel-by-pixel correlation

coefficient between the two maps using all the gray matter pixels. FC with short time

window was obtained by taking correlation coefficient using all the frames within a 30

sec window. Steps of 3 sec and 30 sec were used for the sliding window and

non-overlapping window, respectively. Scan-shifted control was calculated by shifting

the scan number of hemodynamics data relative to simultaneously obtained calcium data.

Analysis of Co-Activation Patterns (CAPs)

CAP analysis was adopted from previous fMRI studies (Liu and Duyn 2013). Briefly, the

calcium time course from each ROI was z-normalized. CAPs were calculated for each

ROIs. Frames corresponding to large peaks (> 2 s.d.) of the time course of a given ROI

were considered CAPs. We examined whether CAPs calculated using the calcium signal

could predict the sliding window FC calculated using the hemodynamic signal. For each

ROI and each time window, we quantified the similarity between the sliding window FC

and CAPs by calculating the spatial correlation between the FC map and CAPs. To

quantify coordinated temporal variations in CAPs and FC, we also calculated the

temporal deviation of CAP and FC from the mean patterns as described in the following.

First, we calculated average patterns of CAPs and FC in a given scan. For a given scan

and each ROI_i, the average of all CAPs was calculated using the entire scan [$(CAP_{scan})_i$].

Similarly, an FC map was calculated for the same scan and the same ROI [$(FC_{scan})_i$]. We

repeated this procedure to obtain ROI-by-ROI matrices of CAP_{scan} and FC_{scan}. Next, we

calculated ROI-by-ROI matrices of CAPs and FC in a short time window (CAP_{window} and

FC_{window}) using the same procedure used to calculate CAP_{scan} and FC_{scan} but for each 30

sec time window. Then, the difference between CAP_{window} and CAP_{scan} was taken to

quantify the deviation of CAP in a time window from the mean CAP pattern in the entire

scan [$\Delta CAP = CAP_{window} - CAP_{scan}$]. Similarly, ΔFC was obtained by subtracting FC_{scan}

from FC_{window}. Finally, the correlation coefficient between non-diagonal elements of the

ROI-by-ROI matrices of ΔCAP and ΔFC were calculated. When CAPs were absent for a

particular ROI in a time window, that ROI was omitted from the calculation for the time

window.

Cluster Analysis and Kurtosis Analysis

For the state analysis of sliding window FC, we adopted the k-means clustering algorithm

used in the previous studies (Allen et al. 2014; Laumann et al. 2016). Correlation distance

(1-r) was used to compute the separation between each window's FC-matrix (using all 38

ROIs) and the k-means clustering was iterated 100 times with random centroid positions

to avoid local minima. The windowed FC-matrices were mean-centered by scan to

eliminate scan-level and subject-level features from contributing the clustering result.

The k-means clustering was applied in the same manner to the simulated data that was

matched in size to the real data. The cluster validity index was used to evaluate the quality

of clustering for the range of cluster numbers (k = 2–10). The cluster validity index was

computed as the average ratio of within-cluster distance to between-cluster distance.

The non-stationarity of spontaneous neuronal signal correlation was assessed by

calculating the multivariate kurtosis using the same procedure as described by Laumann

and colleagues (Laumann et al. 2016). One value of kurtosis was calculated for each FC

matrix (using all 38 ROIs) obtained at each scan. The same procedure was applied to the

simulated data that was matched in size to the real data. A significant difference between

the kurtosis measure of the real and the simulated data indicates either non-stationarity of

FC or non-Gaussianity of the signal or both.

Time Course Simulation

To obtain a null dataset to evaluate the non-stationarity of the real data, we constructed

simulated time courses using a method developed by Laumann and colleagues (Laumann

et al. 2016). Briefly, random normal deviates having the same dimensionality as a real

dataset are sampled. These time courses are multiplied in the spectral domain by the

average power spectrum of the (bandpass filtered) real data. These time courses are then

projected onto the eigenvectors derived from the covariance matrix of the real data. This

procedure produces simulated data that are stationary by construction but matched to real

data in the covariance structure and mean spectral content.

Assessment of motion-related and physiological artifacts

To estimate the strength of motion-related and physiological artifacts from functional

images, we calculated DVARS using the hemodynamic signal with a procedure described

in fMRI literatures (Power et al. 2012; Power et al. 2014). Specifically, for each scan,

after the preprocessing, root mean squares of the temporal derivatives of the

hemodynamic time courses were calculated and averaged across ROIs to obtain one time

course of DVARS. To remove the data potentially contaminated by the artifacts, we

conducted two types of analyses. First, for each scan, the histogram of DVARS was

calculated to exclude scans with strong skewness (Supplementary Fig. 3)

(scan-censoring). To define a DVARS histogram with strong skewness, we calculated

DVARS histograms for 1000 sets of the simulation described in the preceding section. If

the skewness of the data was larger than the 99[th] percentiles of the simulation, the scan

was considered to be strongly skewed. Second, frame-censoring was conducted at

multiple DVARS thresholds as described in the fMRI literatures (Power et al. 2012;

Power et al. 2014). For a given threshold, frames with DVARS larger than the threshold

were simply discarded from the subsequent analyses.

Assessment of the normality of the signal

To check for the normality of the hemodynamic and the calcium signals, we computed the

4[th] moment of the signal distribution for each scan. For each scan, signals in all the ROIs

were concatenated before calculating the 4[th] moment. Similar results were obtained when

the 4[th] moment was calculated separately for individual ROIs and then the mean of the 4[th]

moment was obtained for each scan (data not shown). The calculated 4[th] moment of the

real data was then compared with that of the 1000 simulated time courses obtained by the

procedure described above. If the 4[th] moment of the real data was larger than the 99[th]

percentile of the simulation, the signal in the scan was considered to be non-normal. In

the case of the calcium signal, we applied a median filter to remove high frequency noise

and enforce the normality of the signal. Calcium time courses often contained some high

frequency noise on top of the slower calcium activity (Fig. 1A), which was most likely

due to the moderate level of excitation light adjusted to avoid bleaching of GCaMP

fluorophore. We consider that the median filter removed this high frequency noise while

retaining the slower calcium signal that was approximately normally distributed.

Results

Consistent FC dynamics in calcium and hemodynamic signals

Transgenic mice expressing GCaMP in neocortical neurons were used to simultaneously

measure the neuronal calcium signal and hemodynamics in a large portion of the bilateral

neocortex (Fig. 1A) (Matsui et al. 2016). Mice were lightly anesthetized and head-fixed

with metal plates so that head-motion could not contaminate the signals. For both calcium

and hemodynamic signals, power spectra of the signals exhibited an approximately linear

trend on log-log plots (Supplementary Fig. 2) suggesting that the non-neuronal artifact

was small. We used sliding window correlation (30 sec window at 3 sec steps) to examine

whether the calcium and hemodynamic FC in mice exhibited "dynamic" changes.

Consistent with previous reports in humans (Allen et al. 2014; Chang and Glover 2010;

Zalesky et al. 2014) and other animals (Hutchison et al. 2014; Majeed et al. 2009), FC

between pairs of ROIs calculated with sliding windows showed considerable variability

over different time points both in calcium signal and hemodynamics (Fig. 1B–C).

Consistent with the idea that variability in hemodynamic FC arises from underlying

neuronal activity, we found close matches between dFC of calcium and hemodynamics

(correlation coefficients, 0.631 and 0.675 for Figs. 1B and 1C, respectively). Correlation

between the time courses of calcium FC and hemodynamic FC was significantly larger

for the data than for the scan-shifted control ($P < 10^{-20}$, Kolmogorov-Smirnov test; Fig.

1D).

To further examine the consistency between dFC in the calcium signal and

hemodynamics in the entire neocortex, we calculated the calcium and hemodynamic FC

among all pairs of ROIs and compared them across time windows (Fig. 2A–B). The

ROI-based FC-matrices in calcium and hemodynamics both showed variability across

time windows. On the other hand, FC matrices in calcium and hemodynamics within each

time window were similar. If dFC in calcium and hemodynamics are matched, the

similarity between calcium and hemodynamic FC in the same time window should be

higher than that calculated using different time windows (e.g., similarity between

Ca-FC$_{window\#1}$ and Hemo-FC$_{window\#1}$ would be higher than the similarity between

Ca-FC$_{window\#1}$ and Hemo-FC$_{window\#2}$). Otherwise, the similarity between FC-matrices in

calcium and hemodynamics merely reflects the overall similarity of FC in calcium and

hemodynamics but not the coordinated "dynamics" of calcium and hemodynamic FC.

Across all the data, we found that the distribution of the correlation coefficient between

the FC-matrices in calcium and hemodynamics was shifted toward positive values

compared with that calculated with the scan-shifted data ($P < 10^{-14}$,

Kolmogorov–Smirnov test; Fig. 2C). The difference between the real data and the

trial-shifted data was also consistently positive across animals ($P < 0.0156$, n = 7 mice,

sign rank test; Fig. 2D) and was seen across various window sizes ranging from 1 sec to

60 sec (Fig. 2E). Together these results suggest that temporal variability in hemodynamic

FC, as measured with a sliding window, arises from neural activity rather than from

movement-related artifacts (Laumann et al. 2016) or non-neuronal physiological artifacts

such as heartbeat and respiration (Bianciardi et al. 2009; Shmueli et al. 2007).

Variations in transient neuronal coactivations explained variations in FC

What are the potential neuronal events that create dFC? Recent fMRI studies have

proposed that variability in the neuronal coactivation pattern (CAP) of brain areas is

directly reflected in the "dynamic" change of FC observed with the sliding window

correlation (Liu and Duyn 2013). To address this possibility, for each scan, we compared

the sliding window FC calculated using hemodynamics with the CAPs calculated using

the calcium signal. The use of the calcium signal for extracting CAPs allowed us to

capture faster spatiotemporal dynamics than the hemodynamics. More importantly, the

use of two different signals also allowed us to avoid comparing sliding window FC and

CAPs that were derived from the same signals and could lead to circular logic.

For each anatomical ROI, we first detected peaks in the calcium signal within a

given time-window and then defined CAPs as the frames in the calcium signal

corresponding to the detected peak locations (Fig. 3A) (Liu and Duyn 2013). Similar to

the previous reports in fMRI (Liu et al. 2013; Liu and Duyn 2013), we found variations in

the spatial patterns of CAPs extracted from the same ROI (Fig. 3A, panels above time

courses). We used CAPs to examine whether variations of the spatial pattern of CAPs in

different time windows could explain the spatial variation of sliding window FC. For

each ROI in each 30 sec window, we extracted CAPs and FC using the calcium and

hemodynamic signals, respectively. In the example 30 sec windows shown in Figure 3A,

time courses of the chosen ROI showed transient activations that resulted in 11 and 3

frames of CAPs (corresponding to 1.1 and 0.3 sec of data, respectively). Despite the small

number of frames corresponding to CAPs, the average spatial pattern of CAPs in the time

window closely matched the spatial pattern of hemodynamic FC calculated in the same

time window (compare the mean CAP and mean FC in Fig. 3A).

To further compare CAPs with sliding window hemodynamic FC across ROIs, we

calculated CAPs for all pairs of ROIs and compared them with the FC of the same

ROI-pairs in the same time window (Fig. 3B). Across all the data, CAP-matrices and FC

matrices showed a high positive correlation (Fig. 3C–D; mean R = 0.525 across animals)

suggesting that CAP and FC calculated using the same sliding window were similar.

The similarity between calcium CAP and hemodynamic FC in a short time window

does not necessarily indicate coordinated temporal variation between CAP and FC, but

could result entirely from similarity between the time-average patterns of CAP and FC.

Therefore, to further examine whether coordinated temporal variations in CAPs and

sliding-window FC exist, we calculated ΔCAP and ΔFC by subtracting from each CAP

and FC in each time window the average pattern of CAP and FC, respectively, that were

calculated using the entire scan (Fig. 4A). A coordinated change in ΔCAP and ΔFC

indicates a similar temporal fluctuation of CAP and sliding-window FC that cannot be

accounted for by the similarity in the mean pattern of CAP and FC calculated using the

entire scan. We found that the distribution of the correlation between ΔCAP and ΔFC for

the real data was shifted toward positive values whereas the same distribution calculated

using trial-shifted data was centered near zero ($P < 10^{-30}$, Kolmogorov–Smirnov test; Fig.

4B). Furthermore, the correlation between ΔCAP and ΔFC was consistently positive

across all animals ($P < 0.156$, n = 7 mice, sign rank test; Fig. 4C) and was seen across

various sizes of time windows ranging from 1 to 60 sec (Fig. 4D). Excluding scans with

potential artifacts using DVARS (Power et al. 2012; Power et al. 2014) still yielded a

significant difference between the real and the trial-shifted data (Supplementary Fig. 9).

Taken together, these results suggest that temporal fluctuations of the spatial pattern of

CAPs at least partly explain temporal fluctuations of hemodynamic FC calculated using

sliding windows.

"Dynamics" of FC arise from non-stationarity of resting-state activity

Because FC is estimated by using a finite number of time-points, temporal fluctuations of

FC observed in short time windows could arise from mere sampling error even when the

underlying FC is stationary (Laumann et al. 2016). We next addressed whether the

sampling error could explain the dFC observed in the present data. We compared two

indices used in a previous study, namely cluster validity index and kurtosis, for real data

and simulated data that were matched in spectral and covariance properties (Fig. 5A)

(Laumann et al. 2016). The cluster validity index measures the degree of clustering of

multiple sliding window FC calculated within the scan. Note that a smaller cluster

validity index indicates more clustering (see Materials and Methods for details). For both

calcium and hemodynamic signals, we found the cluster validity index of the real data to

be significantly smaller than that of the simulated data (Fig. 5B), suggesting that the real

data had a cluster structure that could not be fully accounted for by sampling error.

Similarly, we calculated the kurtosis of the covariance matrices of real and simulated data.

If the kurtosis of real data is larger than that of simulated data assuming a stationary

Gaussian process, non-stationarity is implied for the real data, if the real data is normally

distributed (Laumann et al. 2016). For the hemodynamic signal, the real data was

approximately normally distributed (Supplementary Fig. 8A). Thus, we calculated the

kurtosis using the hemodynamic signal without further preprocessing. For the calcium

signal, because the original signal was not normally distributed, we applied further

preprocessing before calculating the kurtosis. First, we applied an additional high pass

temporal filter to remove low frequencies (< 0.1 Hz). Then we used a median filter to

enforce the normality of the signal (Supplementary Fig. 6). For both hemodynamic and

calcium signals, we found that the kurtosis of the real data was significantly higher than

that of the simulated data ($P < 10^{-10}$ for calcium, sign rank test, n = 50 scans that showed

the approximately normal distribution of the signal (see Materials and Methods for the

details of the assessment of normality); $P < 10^{-11}$ for hemodynamics, sign rank test, n = 64

scans; Fig. 5C, Supplementary Fig. 8B). The calcium signal at low frequency (< 0.1 Hz)

was approximately normal without additional preprocessing (Supplementary Fig. 7A).

The kurtosis of the real and the simulated data was also significantly different at this

frequency range, though the magnitude of the difference was smaller (Supplementary Fig.

7B). Together, these results suggest that dFC arises from the non-stationarity of

spontaneous neuronal activity, and analyses based on sliding window correlation have the

potential to detect non-stationarity.

To assess the potential contribution of motion-related and physiological artifacts on

the analysis of the non-stationarity using kurtosis, we used DVARS calculated using the

hemodynamic signal to exclude time points for which the artifacts might have been

problematic (Power et al. 2012; Power et al. 2014). In some scans with large peaks in the

time courses of DVARS, indicating potential body movements, the histogram of DVARS

was strongly right-skewed (Supplementary Fig. 3). Analysis of the kurtosis using a subset

of scans for which the DVARS distribution was not right-skewed compared to the

stationary simulation also revealed a significant difference between the real and the

simulated data ($P < 10^{-14}$, sign rank test, n = 25; Supplementary Fig. 4; see Materials and

Methods for details). Furthermore, frame-censoring at several thresholds of DVARS

revealed that the difference of kurtosis between the real and the simulated data did not

depend on the levels of DVARS thresholds (Supplementary Fig. 5). These results suggest

that the motion-related and physiological artifacts, as detected by DVARS, did not

significantly affect the present results.

Discussion

In the present study, we used simultaneous imaging of calcium and hemodynamic signals to show that temporal fluctuations in hemodynamic FC calculated in a short time window closely follow that of calcium FC, suggesting the neuronal origin of dFC. We have further shown that the spatial pattern of hemodynamic FC in a short time window is predicted by averaging the transient coactivations in the calcium signal (CAPs) contained within the same time window suggesting that temporally interspersed transient neuronal events underlie resting-state FC. Finally, we have shown that in both calcium and hemodynamic signals, statistical properties of FC calculated in a short time window were significantly different from those obtained with simulated signals that were stationary by construction. These results advocate for the analysis of the "dynamic" aspect of FC obtained in human fMRI experiments. Insights of the neuronal events underlying dFC provided by the present study would also be informative for developing appropriate analysis methods for dFC.

Relationship to previous investigations of the neuronal origin of dFC

To provide direct evidence linking neuronal activity and dFC, several groups have conducted simultaneous recording of fMRI and local field potential (LFP) (Lu et al. 2007; Pan et al. 2011; Thompson et al. 2013) or EEG (Chang et al. 2013; Tagliazucchi et al. 2012b). However, these previous studies were limited in several ways. Because LFP recordings were limited by a small number of recording sites and EEG recordings did not have enough spatial resolution, evidence directly linking the global spatial pattern of neuronal activity with hemodynamic FC has been lacking. Using simultaneous imaging of calcium and hemodynamic signals, the present study provides evidence suggesting that temporal variability of hemodynamic FC and its time-to-time spatial patterns reflect spatial patterns of large-scale neuronal activity. Moreover, because the present study used anesthetized and head-fixed mice, the results are unlikely to be attributable to head motion.

Recent human fMRI studies have proposed that neuronal activity important for FC is condensed into transient large scale neuronal coactivations (i.e. CAPs) (Liu and Duyn 2013; Tagliazucchi et al. 2012a; Tagliazucchi et al. 2011). Consistent with this idea, imaging studies in mice have revealed transient neuronal coactivations across brain areas (Matsui et al. 2016; Mohajerani et al. 2013; Vanni and Murphy 2014). In our previous

study, we searched for neuronal coactivations that resembled the spatial patterns of
(static) FC and showed that such neuronal coactivations were converted into spatially
similar hemodynamic signals (Matsui et al. 2016). In the present study, we took a
different approach that was similar to single frame analysis methods employed in recent
human fMRI studies (Karahanoğlu and Van De Ville 2015; Liu et al. 2013; Liu and
2013; Tagliazucchi et al. 2011). We were especially interested whether CAPs represent
the potential neuronal events underlying the temporal variability of sliding window FC.
Because derivation of CAPs and sliding-window FC using identical BOLD signals
potentially lead to circular logic, in the present study, we examined the link between
CAPs and sliding window FC derived from different signals (calcium and
respectively). Instead of specifically looking at neuronal coactivations that resembled
"static" FC, we included all the individual CAPs into our analysis and showed that
variation of the spatial pattern of individual CAPs across time windows was
related to variations of hemodynamic FC across time windows. Thus, the present
suggest the importance of the development of an analysis that specifically focuses on
CAPs   (Karahanoğlu and Van De Ville 2015; Liu et al. 2013). However, the modest
correlation found between CAP and sliding window FC implies that the fluctuations of
calcium and hemodynamic signals from the average pattern maybe small. It should also
be noted that, although statistically significant, the correlation between ΔCAP and ΔFC
was relatively weak. Part of the reason for this could be non-neuronal physiological
noise that contributed to the hemodynamics (Matsui et al. 2016). In the present study,
because of the use of anesthesia and head-fixation, head motion was unlikely to be the
primary source of the non-neuronal noise. However, other physiological activities (e.g.
respiration and heartbeat) are known to affect hemodynamics (Chang et al. 2009; Chang
and Glover 2009) and, thus, are likely to affect temporal fluctuation of hemodynamic
FC as well. Although scan-wise data exclusion using DVARS suggested that the
motion-contamination was not the major cause of the observed temporal correlation
between ΔCAP and ΔFC (Supplementary Fig. 9), the common artifacts on
hemodynamic and calcium signals could have contributed to the observed temporal
correlation. Our results (i.e. relatively low correlation between ΔCAP and ΔFC) indicate
that correction for such non-neuronal physiological noise (Glover et al. 2000) is likely
to be essential for the analysis of dFC.
473

Non-stationarity of spontaneous brain activity correlation

It has been a matter of debate as to what extent temporal fluctuations of FC are

attributed to the dynamics of underlying neuronal activity and not to non-neuronal

sources of noise (e.g., head motion, sampling variability; reviewed in (Hutchison et al.

2013)). Laumann and colleagues have reported that most of the temporal fluctuations of

single subject FC is explained by head motion (Laumann et al. 2016). After controlling

for the head motion, Laumann and colleagues concluded that the statistical properties of

resting-state FC in human fMRI are indistinguishable from those obtained with

simulated signals that are stationary by construction. A similar study by Hindriks and

colleagues has also indicated that the apparent dFC calculated with the sliding window

method does not necessarily indicate non-stationarity of the resting brain network

(Hindriks et al. 2016). However, in terms of spontaneous neuronal activity itself, there

is substantial evidences showing that spontaneous neuronal activity is non-stationary

(Foster and Wilson 2006; Ji and Wilson 2007; Logothetis et al. 2012). In particular,

under both awake and anesthetized states, transient neuronal events such as sharp wave

ripples have been shown to produce coordinated activity across the entire brain

(Logothetis et al. 2012). The present results are consistent with these previous studies

supporting the non-stationarity of neuronal activity and further showed that FC

calculated using such non-stationary neuronal activity also showed non-stationarity, as

expected.

Potential contribution of arousal

Fluctuation in the level of arousal has been shown to contribute to the apparent

"dynamics" and non-stationarity of FC (Laumann et al. 2016; Tagliazucchi and Laufs

2014). The present study observed larger temporal variability of hemodynamic FC in

anesthetized mice than in awake, eyes open-fixated humans (Laumann et al. 2016).

Interestingly, a recent study reported large within-subject FC variability in subjects

instructed to close their eyes (Pannunzi et al. 2017), which predisposes them to sleep

(Tagliazucchi and Laufs 2014). In the present study, it could be possible that a

fluctuating level of anesthesia, instead of the subject's sleep state, could have

contributed to the greater variability of FC. Alternatively, rapid fluctuation between

awake and sleep states in mice (Adler et al. 2014) could have contributed to the greater

variability of FC compared to humans. According to the results of the analysis using

power spectra (Supplementary Fig. 2) and DVARS (Supplementary Figs. 3–5), we

consider that the level of arousal rarely reached a point at which the animal started to

struggle. Nevertheless, the present data alone were not sufficient to exclude the

possibility that the fluctuation of the level of arousal accounted for a large fraction of

the FC "dynamics" and non-stationarity observed here. Future experiments conducting

simultaneous recordings of functional images and physiological signals (e.g.

electroencephalogram, respiration-rates) in mice would be able to assess the exact

amount of non-stationarity in FC under a defined state of arousal.

Other limitations of the study

It should be clearly stated that the present results do not guarantee that sliding window

methods are always capable of detecting non-stationarity in human resting-state fMRI

data.   The   present   study   used   tightly   head-restrained   animals   and   high

signal-to-noise-ratio (SNR) imaging at a high frame rate (5 and 10 Hz for the

hemodynamics and calcium signal, respectively). Compared to the present experimental

conditions, overall SNR in typical human resting-state fMRI is likely to be substantially

compromised. Under such low SNR conditions, it is not clear whether simple sliding

window correlation methods can detect the non-stationarity of FC (Hindriks et al. 2016;

Laumann et al. 2016). With respect to SNR, we expect that the recent development of

high-speed fMRI (Feinberg et al. 2010) will significantly improve the detectability of

non-stationarity. Nevertheless, the present results suggest that, rather than the sliding

window based method, an alternative analysis strategy that directly extracts CAPs from

hemodynamic signals (Karahanoğlu and Van De Ville 2015; Liu et al. 2013; Liu and

Duyn 2013; Tagliazucchi et al. 2012a) may be more appropriate for extracting relevant

information related to dFC. Care should be taken, however, because the smaller brain

and the low dimensionality of FC in the mouse (compared to the human subjects) could

also have made CAPs from just a few frames look similar to those of FC.

It should also be noted that the present results do not claim that dFC has significant

behavioral or cognitive consequences. Instead of examining the potential relationship

between dFC and cognitive dynamics or behavioral variability [see (Preti et al. 2016)

for a recent review], here we focused on validating the neuronal origin of dFC.

Experiments under anesthesia greatly reduced potential confounding factors, such as

head   motion   and   arousal   state   (Hutchison   et   al.   2014;   Laumann   et   al.   2016).

Nevertheless, the present wide-field imaging setup can be naturally extended to awake

imaging with task-performing mice (Ferezou et al. 2007; Wekselblatt et al. 2016). Such

experiments would reveal the potential consequences of dFC on the behavior.

Acknowledgements

We thank A. Honda and Y. Sono for animal care and genotyping; the Research Support Center, Graduate School of Medical Sciences, Kyushu University, for technical support: and K. Jimura and Y. Noro for helpful discussion. Support for this work was provided by grants from Brain Mapping by Integrated Neurotechnologies for Disease Studies (Brain/MINDS) – Japan Agency for Medical Research and Development (AMED), Core Research for Evolutionary Science and Technology (CREST) – AMED and Strategic International Research Cooperative Program (SCIP) – AMED, and Japan Society for Promotions of Sciences (JSPS) KAKENHI Grants 25221001 and 25117004 (to K.O.), World Premium Institute (WPI), JSPS (to K.O.); JSPS KAKENHI Grant 17K14931 (to T. Matsui) and JSPS Research Fellowship 20153597 (to T. Murakami).

References

Adachi Y, Osada T, Sporns O, Watanabe T, Matsui T, Miyamoto K, Miyashita Y. 2012. Functional connectivity between anatomically unconnected areas is shaped by collective network-level effects in the macaque cortex. Cereb Cortex. 22(7):1586-1592.

Adler DA, Ammanuel S, Lei J, Dada T, Borbiev T, Johnston MV, Kadam SD, Burd I. 2014. Circadian cycle-dependent eeg biomarkers of pathogenicity in adult mice following prenatal exposure to in utero inflammation. Neuroscience. 275:305-313.

Allen EA, Damaraju E, Plis SM, Erhardt EB, Eichele T, Calhoun VD. 2014. Tracking whole-brain connectivity dynamics in the resting state. Cereb Cortex. 24(3):663-676.

Bianciardi M, Fukunaga M, van Gelderen P, Horovitz SG, de Zwart JA, Shmueli K, Duyn JH. 2009. Sources of functional magnetic resonance imaging signal fluctuations in the human brain at rest: A 7 t study. Magn Reson Imaging. 27(8):1019-1029.

Biswal B, Yetkin FZ, Haughton VM, Hyde JS. 1995. Functional connectivity in the motor cortex of resting human brain using echo-planar mri. Magn Reson Med. 34(4):537-541.

Calhoun VD, Miller R, Pearlson G, Adalı T. 2014. The chronnectome: Time-varying connectivity networks as the next frontier in fmri data discovery. Neuron. 84(2):262-274.

Chang C, Cunningham JP, Glover GH. 2009. Influence of heart rate on the bold signal: The cardiac response function. Neuroimage. 44(3):857-869.

Chang C, Glover GH. 2009. Relationship between respiration, end-tidal co2, and bold signals in resting-state fmri. Neuroimage. 47(4):1381-1393.

Chang C, Glover GH. 2010. Time-frequency dynamics of resting-state brain connectivity measured with fmri. Neuroimage. 50(1):81-98.

Chang C, Liu Z, Chen MC, Liu X, Duyn JH. 2013. Eeg correlates of time-varying bold functional connectivity. Neuroimage. 72:227-236.

Deco G, Ponce-Alvarez A, Mantini D, Romani GL, Hagmann P, Corbetta M. 2013. Resting-state functional connectivity emerges from structurally and dynamically shaped slow linear fluctuations. J Neurosci.

33(27):11239-11252.

Feinberg DA, Moeller S, Smith SM, Auerbach E, Ramanna S, Gunther M,
Glasser MF, Miller KL, Ugurbil K, Yacoub E. 2010. Multiplexed echo
planar imaging for sub-second whole brain fmri and fast diffusion
imaging. PLoS One. 5(12):e15710.

Ferezou I, Haiss F, Gentet LJ, Aronoff R, Weber B, Petersen CC. 2007.
Spatiotemporal dynamics of cortical sensorimotor integration in
behaving mice. Neuron. 56(5):907-923.

Foster DJ, Wilson MA. 2006. Reverse replay of behavioural sequences in
hippocampal place cells during the awake state. Nature.
440(7084):680-683.

Fox MD, Raichle ME. 2007. Spontaneous fluctuations in brain activity
observed with functional magnetic resonance imaging. Nat Rev
Neurosci. 8(9):700-711.

Fox MD, Snyder AZ, Vincent JL, Corbetta M, Van Essen DC, Raichle ME.
2005. The human brain is intrinsically organized into dynamic,
anticorrelated functional networks. Proc Natl Acad Sci U S A.
102(27):9673-9678.

Glover GH, Li TQ, Ress D. 2000. Image-based method for retrospective
correction of physiological motion effects in fmri: Retroicor. Magn
Reson Med. 44(1):162-167.

Hansen EC, Battaglia D, Spiegler A, Deco G, Jirsa VK. 2015. Functional
connectivity dynamics: Modeling the switching behavior of the resting
state. Neuroimage. 105:525-535.

Hindriks R, Adhikari MH, Murayama Y, Ganzetti M, Mantini D, Logothetis
NK, Deco G. 2016. Can sliding-window correlations reveal dynamic
functional connectivity in resting-state fmri? Neuroimage.
127:242-256.

Honey CJ, Sporns O, Cammoun L, Gigandet X, Thiran JP, Meuli R,
Hagmann P. 2009. Predicting human resting-state functional
connectivity from structural connectivity. Proc Natl Acad Sci U S A.
106(6):2035-2040.

Hutchison RM, Hutchison M, Manning KY, Menon RS, Everling S. 2014.

Isoflurane induces dose-dependent alterations in the cortical
connectivity profiles and dynamic properties of the brain's functional
architecture. Hum Brain Mapp. 35(12):5754-5775.

Hutchison RM, Womelsdorf T, Allen EA, Bandettini PA, Calhoun VD,
Corbetta M, Della Penna S, Duyn JH, Glover GH, Gonzalez-Castillo J
et al. 2013. Dynamic functional connectivity: Promise, issues, and
interpretations. Neuroimage. 80:360-378.

Ji D, Wilson MA. 2007. Coordinated memory replay in the visual cortex and
hippocampus during sleep. Nat Neurosci. 10(1):100-107.

Karahanoğlu FI, Van De Ville D. 2015. Transient brain activity disentangles
fmri resting-state dynamics in terms of spatially and temporally
overlapping networks. Nat Commun. 6:7751.

Laumann TO, Snyder AZ, Mitra A, Gordon EM, Gratton C, Adeyemo B,
Gilmore AW, Nelson SM, Berg JJ, Greene DJ et al. 2016. On the
stability of bold fmri correlations. Cereb Cortex.

Liu X, Chang C, Duyn JH. 2013. Decomposition of spontaneous brain activity
into distinct fmri co-activation patterns. Front Syst Neurosci. 7:101.

Liu X, Duyn JH. 2013. Time-varying functional network information
extracted from brief instances of spontaneous brain activity. Proc Natl
Acad Sci U S A. 110(11):4392-4397.

Liégeois R, Laumann TO, Snyder AZ, Zhou J, Yeo BTT. 2017. Interpreting
temporal fluctuations in resting-state functional connectivity mri.
Neuroimage. 163:437-455.

Logothetis NK, Eschenko O, Murayama Y, Augath M, Steudel T, Evrard HC,
Besserve M, Oeltermann A. 2012. Hippocampal-cortical interaction
during periods of subcortical silence. Nature. 491(7425):547-553.

Lu H, Zuo Y, Gu H, Waltz JA, Zhan W, Scholl CA, Rea W, Yang Y, Stein EA.
2007. Synchronized delta oscillations correlate with the resting-state
functional mri signal. Proc Natl Acad Sci U S A. 104(46):18265-18269.

Ma Y, Shaik MA, Kim SH, Kozberg MG, Thibodeaux DN, Zhao HT, Yu H,
Hillman EM. 2016. Wide-field optical mapping of neural activity and
brain haemodynamics: Considerations and novel approaches. Philos
Trans R Soc Lond B Biol Sci. 371(1705).

Majeed W, Magnuson M, Keilholz SD. 2009. Spatiotemporal dynamics of low
frequency fluctuations in bold fmri of the rat. J Magn Reson Imaging.
30(2):384-393.
Matsui T, Koyano KW, Tamura K, Osada T, Adachi Y, Miyamoto K, Chikazoe
J, Kamigaki T, Miyashita Y. 2012. Fmri activity in the macaque
cerebellum evoked by intracortical microstimulation of the primary
somatosensory cortex: Evidence for polysynaptic propagation. PLoS
One. 7(10):e47515.
Matsui T, Murakami T, Ohki K. 2016. Transient neuronal coactivations
embedded in globally propagating waves underlie resting-state
functional connectivity. Proc Natl Acad Sci U S A. 113(23):6556-6561.
Matsui T, Tamura K, Koyano KW, Takeuchi D, Adachi Y, Osada T, Miyashita
Y. 2011. Direct comparison of spontaneous functional connectivity and
effective connectivity measured by intracortical microstimulation: An
fmri study in macaque monkeys. Cereb Cortex. 21(10):2348-2356.
Messé A, Rudrauf D, Benali H, Marrelec G. 2014. Relating structure and
function in the human brain: Relative contributions of anatomy,
stationary dynamics, and non-stationarities. PLoS Comput Biol.
10(3):e1003530.
Mohajerani MH, Chan AW, Mohsenvand M, LeDue J, Liu R, McVea DA, Boyd
JD, Wang YT, Reimers M, Murphy TH. 2013. Spontaneous cortical
activity alternates between motifs defined by regional axonal
projections. Nat Neurosci. 16(10):1426-1435.
Murakami T, Matsui T, Ohki K. 2017. Functional segregation and
development of mouse higher visual areas. J Neurosci.
37(39):9424-9437.
Murakami T, Yoshida T, Matsui T, Ohki K. 2015. Wide-field ca(2+) imaging
reveals visually evoked activity in the retrosplenial area. Front Mol
Neurosci. 8:20.
Pan WJ, Thompson G, Magnuson M, Majeed W, Jaeger D, Keilholz S. 2011.
Broadband local field potentials correlate with spontaneous
fluctuations in functional magnetic resonance imaging signals in the
rat somatosensory cortex under isoflurane anesthesia. Brain Connect.

1(2):119-131.

Pannunzi M, Hindriks R, Bettinardi RG, Wenger E, Lisofsky N, Martensson
J, Butler O, Filevich E, Becker M, Lochstet M et al. 2017. Resting-state
fmri correlations: From link-wise unreliability to whole brain stability.
Neuroimage. 157:250-262.

Power JD, Barnes KA, Snyder AZ, Schlaggar BL, Petersen SE. 2012.
Spurious but systematic correlations in functional connectivity mri
networks arise from subject motion. Neuroimage. 59(3):2142-2154.

Power JD, Mitra A, Laumann TO, Snyder AZ, Schlaggar BL, Petersen SE.
2014. Methods to detect, characterize, and remove motion artifact in
resting state fmri. Neuroimage. 84:320-341.

Preti MG, Bolton TA, Van De Ville D. 2016. The dynamic functional
connectome: State-of-the-art and perspectives. Neuroimage.

Shmueli K, van Gelderen P, de Zwart JA, Horovitz SG, Fukunaga M, Jansma
JM, Duyn JH. 2007. Low-frequency fluctuations in the cardiac rate as
a source of variance in the resting-state fmri bold signal. Neuroimage.
38(2):306-320.

Tagliazucchi E, Balenzuela P, Fraiman D, Chialvo DR. 2012a. Criticality in
large-scale brain fmri dynamics unveiled by a novel point process
analysis. Front Physiol. 3:15.

Tagliazucchi E, Balenzuela P, Fraiman D, Montoya P, Chialvo DR. 2011.
Spontaneous bold event triggered averages for estimating functional
connectivity at resting state. Neurosci Lett. 488(2):158-163.

Tagliazucchi E, Laufs H. 2014. Decoding wakefulness levels from typical fmri
resting-state data reveals reliable drifts between wakefulness and
sleep. Neuron. 82(3):695-708.

Tagliazucchi E, von Wegner F, Morzelewski A, Brodbeck V, Laufs H. 2012b.
Dynamic bold functional connectivity in humans and its
electrophysiological correlates. Front Hum Neurosci. 6:339.

Thompson GJ, Merritt MD, Pan WJ, Magnuson ME, Grooms JK, Jaeger D,
Keilholz SD. 2013. Neural correlates of time-varying functional
connectivity in the rat. Neuroimage. 83:826-836.

Tohmi M, Meguro R, Tsukano H, Hishida R, Shibuki K. 2014. The

extrageniculate visual pathway generates distinct response properties
in the higher visual areas of mice. Curr Biol. 24(6):587-597.
Van Dijk KR, Hedden T, Venkataraman A, Evans KC, Lazar SW, Buckner RL.
2010. Intrinsic functional connectivity as a tool for human
connectomics: Theory, properties, and optimization. J Neurophysiol.
103(1):297-321.
Vanni MP, Murphy TH. 2014. Mesoscale transcranial spontaneous activity
mapping in gcamp3 transgenic mice reveals extensive reciprocal
connections between areas of somatomotor cortex. J Neurosci.
34(48):15931-15946.
Vincent JL, Patel GH, Fox MD, Snyder AZ, Baker JT, Van Essen DC, Zempel
JM, Snyder LH, Corbetta M, Raichle ME. 2007. Intrinsic functional
architecture in the anaesthetized monkey brain. Nature.
447(7140):83-86.
Wekselblatt JB, Flister ED, Piscopo DM, Niell CM. 2016. Large-scale
imaging of cortical dynamics during sensory perception and behavior.
J Neurophysiol. 115(6):2852-2866.
White BR, Bauer AQ, Snyder AZ, Schlaggar BL, Lee JM, Culver JP. 2011.
Imaging of functional connectivity in the mouse brain. PLoS One.
6(1):e16322.
Zalesky A, Fornito A, Cocchi L, Gollo LL, Breakspear M. 2014. Time-resolved
resting-state brain networks. Proc Natl Acad Sci U S A.
111(28):10341-10346.
Zariwala HA, Borghuis BG, Hoogland TM, Madisen L, Tian L, De Zeeuw CI,
Zeng H, Looger LL, Svoboda K, Chen TW. 2012. A cre-dependent
gcamp3 reporter mouse for neuronal imaging in vivo. J Neurosci.
32(9):3131-3141.
746
747

Figure Captions

Figure 1. Representative dynamics of simultaneously observed calcium and

hemodynamic FC. (A) Experimental setup. The left most panel shows the setup for

simultaneous imaging. Right side shows example calcium time courses for two ROIs

(green and cyan traces indicate M1 and V1 ROIs, respectively. Positions of the ROIs

indicated in the example field of view. See Supplementary Figure 1 for abbreviations).

FC with short time window uses subset of frames contained in short (30 sec) windows

(red dotted squares). Sliding FC for hemodynamic signal was carried out similarly.

(B)-(C) Examples dynamics of calcium and hemodynamic FC. (B) FC between right V1

and right AC. (C) FC between left M1 and left pPar. See Supplementary Figure 1 for

ROI positions and abbreviations. **(D)** Histogram of correlation between time courses of

Ca-FC and Hemo-FC for the data (solid line) and the scan-shifted control (dotted line).

Data from all pairs of ROIs for all scans obtained in all mice were used.

Figure 2. Significant relationship between calcium and hemodynamic FC calculated

in short time windows. (A)-(B) Example ROI-by-ROI FC matrices for calcium and

hemodynamics for different (non-overlapping) 30 sec windows. FC matrices were

similar for calcium and hemodynamics in the same time window, but not across

different time windows. (A) and (B) are two different examples from different animals.

(C) Cumulative histogram of correlation between FC matrices for calcium and

hemodynamics. Dotted line indicates trial-shifted control. **(D)** Correlation between FC

matrices for calcium and hemodynamics was larger for the data than for the trial-shifted

control significantly across animals. **(E)** Correlation between FC matrices of calcium

and hemodynamics was larger for the data than the trial-shifted control across different

window-sizes (1, 2, 3, 5, 6, 10, 12, 15, 20, 30 and 60 sec). Error bars indicate s.e.m.

across animals (n = 7).

Figure 3. Comparison of calcium CAPs and hemodynamic FC across time-windows.

(A) Procedure for detection of CAPs in calcium signal. For a given ROI, a calcium time

course was extracted and z normalized (green time courses). Then, peaks exceeding 2

s.d. (red dots) were extracted. The frames corresponding to the peaks were considered

CAPs (panels above the time courses). For each window, CAPs in calcium signal were

averaged to obtain mean calcium CAP. Hemodynamic CAPs were calculated similarly

(see Methods). Maps of Ca-FC and Hemo-FC were also calculated using the same time

window. **(B)** Schematics to show the procedure of comparing calcium CAP and

Hemo-FC across all ROI pairs in each time window. In each 30 sec time window, mean

calcium CAPs and Hemo-FC maps were calculated for all ROIs as seeds (left). Then,

for each seed-ROI j, calcium CAP and Hemo-FC values in ROI i were extracted to

obtain a pair of CAP-FC values for the ROI-pairs (i, j) (middle). Finally, for each

time-window, CAP-FC values were compared across all pairs of ROIs (right). **(C)**

Histograms of correlation between CAP and Hemo-FC for all time windows across all

animals. Vertical line indicates mean across time windows. **(D)** Mean correlation

between CAP and Hemo-FC across animals. Note that (C) shows entire distribution of

the data whereas (D) shows reproducibility across mice. Error bar indicates s.e.m.

across animals (n = 7).

Figure 4. Temporal fluctuations in calcium CAPs and Hemo-FC was significantly

related. (A) Schematics of the analysis. In each 30 sec time-window, mean calcium

CAP and Hemo-FC were calculated (indicated as window CAP and window FC,

respectively). From window CAP and window FC, average calcium CAP and average

Hemo-FC that were calculated using the entire scan, in which the 30-sec window

belongs to, were subtracted to obtain maps of ΔCAP and ΔFC, respectively. Finally,

values of ΔCAP and ΔFC were compared across ROI pairs similarly as in Figure 3B.

(B) Histograms of correlation between ΔCAP and ΔFC for all time windows across all

animals. Vertical lines indicate mean across time windows. Solid and dotted lines

indicate real and trial-shifted data, respectively. **(C)** Correlation between ΔCAP and

ΔFC was significantly larger for the data than for trial-shifted control across animals.

(D) Same as (C) but with different window-sizes (1, 2, 3, 5, 6, 10, 12, 15, 20, 30 and 60

sec). Error bars indicate s.e.m. across animals (n = 7).

Figure 5. Comparison with simulated data indicated non-stationarity of the real

data. (A) Examples of real and simulated time courses. Simulated time course (black)

was matched to real data (green; calcium) in mean spectral content (middle panels) and

ROI-by-ROI covariance matrix (right panels). The same procedure was applied to

create simulated hemodynamic data (not shown). **(B)** Cluster validity index calculated

for different number of clusters (k = 2-10). In both calcium (left) and hemodynamics

(right), the cluster validity index was smaller for the real (solid lines) than the simulated

data (dotted lines) indicating that the real data tended to be more clustered. **(C)** Kurtosis

of real and simulated covariance matrices. For both calcium (left) and hemodynamics

(right), multivariate kurtosis was larger for the real than for the simulated data. Error

bars indicate s.e.m. across animals (n = 7).

Supplementary Figure 1. Anatomical locations of ROIs. Anatomical locations of 19

ROIs are shown for the right hemisphere. Anatomical nomenclatures of ROIs are shown

on the right. ROIs in the left hemisphere are taken at mirror symmetric positions to

yield a total of 38 ROIs.

Supplementary Figure 2. Power spectrum of calcium and hemodynamic signals. (A)

Example power spectrum of hemodynamic signal for one mouse. Hemodynamic signals

were not temporally filtered. Black, individual scan. Red, average across scans. **(B)**

Average power spectrum of hemodynamic signal across mice (Red). Black, individual

mouse (averaged across scans). **(C)** Example power spectrum of calcium signal for one

mouse. Calcium signals were not temporally filtered. Black, individual scan. Red,

average across scans. **(D)** Average power spectrum of calcium signal across mice (red).

Black, individual mouse (average across scans).

Supplementary Figure 3. Example time courses and histograms of DVARS

calculated using hemodynamic signal. (A) Example time course of DVARS for one

mouse. **(B)** Histogram of DVARS for the data shown in (A). Inset, same histogram but

for different y-axis range. The histogram was not markedly right skewed (skewness =

1.60). **(C)** Another example time course of DVARS obtained in the same mouse as in

(A). In this scan, large peaks probably representing motion and/or physiological

artifacts were observed. **(D)** Histogram of DVARS calculated using data shown in (C).

Inset, same histogram but for different y-axis range. The histogram was strongly

right-skewed (skewness = 6.54).

Supplementary Figure 4. Difference between the kurtosis of real and simulated data

in Hemodynamics. Kurtosis was calculated for scans selected based on the distribution

of DVARS (n = 25). See Methods for details of the selection of the scans. *, $P < 0.0001$

(sign rank test). Error bars indicate s.e.m.

Supplementary Figure 5. Difference between the kurtosis of real and simulated data

in hemodynamics as a function of DVARS threshold for frame-censoring. The

difference of the kurtosis between the real and the simulated data (blue and red,

respectively) was largely insensitive to the level of the DVARS threshold. Note that the

smaller DVARS threshold indicates stricter threshold. Error bars indicate s.e.m.

Supplementary Figure 6. Kurtosis analysis using calcium signal processed with

additional high-pass and media filters. (A) Example histogram of high frequency

calcium signal (> 0.1 Hz) in one ROI of one mouse before the application of median

filter. Red, Gaussian fit. **(B)** Same as (A) but for the signal after the application of

median filter. Red, Gaussian fit. Note better fitting with Gaussian after median filter

application, suggesting enforced normality of the signal.

Supplementary Figure 7. Kurtosis analysis using low frequency calcium signal. (A)

Example histogram of low frequency calcium signal (0.01 Hz < f < 0.1 Hz) in one ROI

of one mouse. Red, Gaussian fit. (B) Difference between the kurtosis of real and

simulated data in low frequency calcium signal for scans selected based on the

normality of the signal (55 scans). *, $P < 10^{-9}$, sign rank test. Error bars indicate s.e.m.

Supplementary Figure 8. Hemodynamic signal was approximately normally

distributed. (A) Example histogram of hemodynamic signal in one ROI of one mouse.

Red, Gaussian fit. Note that no additional preprocessing was performed. **(B)** Difference

between the kurtosis of real and simulated data in hemodynamic signal for scans

selected based on the normality of the signal (45 scans). *, $P < 10^{-8}$, sign rank test. Error

bars indicate s.e.m.

Supplementary Figure 9. Significant temporal correlation between CAP and

windowed-FC remained after removing scans with potential artifacts. (A) Same

convention as in Figure 4B but for the data using scans selected for small movement

and/or physiological artifacts based on DVARS (25 scans). *, $P < 10^{-17}$,

Kolmogorov-Smirnov test. **(B)** Same convention as in Figure 4C but for the selected

scans. Positive difference between the real and the shifted data was observed in 5 out of

7 mice.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

1. Primary Motor Area (M1)
2. Secondary Motor Area (M2)
3. Parietal Medial Area (mPar)
4. Parietal Lateral Area (lPar)
5. Parietal Posterior Area (pPar)
6. Restrosplenial Area (Rsp)
7. Primary Visual Area (V1)
8. Medial Secondary Visual Area (mV2)
9. Lateral Secondary VIsual Area (lV2)
10. Anterior Cingulate Area (AC)
11. Frontal Association Area (FAsc)
12. Trunk Region of Primary Somatosensory Area (Tr-S1)
13. Hindlimb Region of Primary Somatosensory Area (HL-S1)
14. Shoulder Region of Primary Somatosensory Area (Sh-S1)
15. Forelimb Region of Primary Somatosensory Area (FL-S1)
16. Barrel Region of Primary Somatosensory Area (B-S1)
17. Head Region of Primary Somatosensory Area (H-S1)
18. Auditory Area (Aud)
19. Temporal Association Area (TA)

Supplementary Figure 1

Supplementary Figure 2

Supplementary Figure 3

Supplementary Figure 4

Supplementary Figure 5

Supplementary Figure 6

Supplementary Figure 7

Supplementary Figure 8

Supplementary Figure 9