1   **The nucleoid as a scaffold for the assembly of bacterial signaling complexes**
2
3

4   Audrey Moine[1], Leon Espinosa[1], Eugenie Martineau[1], Mutum Yaikhomba [2], P J Jazleena[2],
5   Deborah Byrne[3], Emanuele G. Biondi[1], Eugenio Notomista[4], Matteo Brilli[5], Virginie Molle[6],
6   Pananghat Gayathri[2], Tâm Mignot[1]*and Emilia M.F. Mauriello[1]*
7
[1]Laboratoire de Chimie Bactérienne, CNRS-Université Aix-Marseille, Marseille, France;
[2]Biology Division, Indian Institute of Science Education and Research, Pune, India;
[3]Protein Purification Platform, Institut de Microbiologie de la Méditerranée, CNRS,
Marseille, France; [4]Dipartimento di Biologia, Università degli Studi di Napoli "Federico II",
Naples, Italy; [5]Centre for Research and Innovation, Fondazione Edmund MachTrento, Italy;
[6]Laboratoire de Dynamique des Interactions Membranaires Normales et Pathologiques,
CNRS-Universités de Montpellier II et I, Montpellier, France.
15
16
*Correspondence to:
Emilia M.F. Mauriello
Laboratoire de Chimie Bactérienne,
CNRS-Université Aix-Marseille
31 Chemin Joseph Aiguier, Marseille 13402, France
Tel. +33491164321
Email : emauriello@imm.cnrs.fr
24
Tâm Mignot
Laboratoire de Chimie Bactérienne,
CNRS-Université Aix-Marseille
31 Chemin Joseph Aiguier, Marseille 13402, France
Tel. +33491164348
Email: tmignot@imm.cnrs.fr

31
Running Title: The *Myxococcus xanthus* FrzCD cytoplasmic chemoreceptor binds DNA
to enhance the signaling activity of Che-like clusters

ABSTRACT

The FrzCD chemoreceptor from the gliding bacterium *Myxococcus xanthus* forms

cytoplasmic clusters that occupy a large central region of the cell body also occupied by the

nucleoid. In this work, we show that FrzCD directly binds to the nucleoid with its N-terminal

positively charged tail and recruits active signaling complexes at this location. The FrzCD

binding to the DNA leads to the formation of multiple distributed clusters that explore

constrained areas. This supra-molecular organization is required for cooperative interactions

between clustered receptors, in turn important for the modulation of bacterial social

behaviors.

1 INTRODUCTION

The bacterial cytoplasm is not a homogeneous solution of macromolecules, but rather a

highly organized and compartmentalized space where the clustering and segregation of

macromolecular complexes at certain cell regions confers functional efficiency (Typas and

Sourjik, 2015). Bacterial chemoreceptors represent a versatile model system to study the

subcellular localization of macromolecules, as they are universally present in prokaryotes

where they form highly ordered arrays that occupy different positions in cells.

Chemoreceptors, also called Methyl-accepting Chemotaxis Proteins (MCP), are capable of

transducing external signals to downstream signaling pathways where phospho-cascades,

initiating at the level of histine kinases CheAs and culminating at the level of output response

regulators CheYs, translate the initial signal into cell behaviors such as regulation of motility,

cell development or aggregation (Sourjik and Berg, 2002; Berleman and Bauer, 2005; Bible *et*

al., 2012). A common feature of the MCPs is their ability to form highly ordered hexagonal

structures, which, by cryoelectron tomography, look like lattices with each unit composed of

an MCP trimer of dimers, two CheW docking proteins and one CheA dimer (Studdert and

Parkinson, 2005; Li *et al.*, 2011; Briegel *et al.*, 2012a; Liu *et al.*, 2012). Receptor clustering is

not strictly required for signal transduction, as one functional unit is enough to generate

phosphorylated CheY (Francis *et al.*, 2002; Li *et al.*, 2011; Li and Hazelbauer, 2011; Piñas *et*

al., 2016). However, MCP clustering is essential to ensure the amplification of the initial

signal, which is a direct consequence of the cooperative interactions between clustered

chemoreceptors (Sourjik and Berg, 2004; Ames and Parkinson, 2006; Li and Hazelbauer,

2014; Piñas *et al.*, 2016; Frank *et al.*, 2016).

While the MCP lattices have been described as universal among prokaryotes (Briegel *et al.*,

2015) their subcellular localization and distribution can vary among different bacterial

species, often reflecting life style complexity, behaviors and functions. For example,

Escherichia coli MCPs localize in one or two polar clusters and more lateral clusters appear as cells become longer (Thiem and Sourjik, 2008). Differently, the TlpT cytoplasmic chemoreceptor from *Rhodobacter sphaeroides* forms a cluster positioned at the center of cells or two clusters positioned at the two and three cell quarters (Thompson *et al.*, 2006). The determinants of these different localization patterns also vary. In *E. coli*, membrane-anchored MCPs form clusters stochastically and through a self-assembly mechanism (Thiem and Sourjik, 2008). The TlpA *Bacillus subtilis* polar chemoreceptor recognizes and associates with strongly curved membrane regions generated during cell septation. These regions become the new poles after cell division, which explains the TlpA polar localization (Strahl *et al.*, 2015). While in *E. coli* and *B. subtilis* the polar targeting of bacterial chemoreceptors is due to intrinsic properties of these proteins, in *Vibrio* species the Che proteins are recruited to the cell poles by a set of specialized proteins responsible of the general maturation of these cell regions (Ringgaard *et al.*, 2011; Ringgaard *et al.*, 2014). The presence of CheWs and CheAs also seem to be universally important in chemoreceptor cluster formation (Sourjik and Berg, 2000; Martin *et al.*, 2001).

Myxococcus xanthus is a gliding bacterium that uses the Frz chemosensory system to modulate the frequency at which cells periodically reverse the direction of their movement on solid surfaces to reorient in the environment, analogously to controlled tumbles in *E. coli* (Blackhart and Zusman, 1985). The Frz regulation of directionality in *M. xanthus* is essential to achieve fruiting body formation, a behavior that bacteria initiate when they are exposed to unfavorable growth conditions. In the Frz pathway, the FrzCD chemoreceptor activates the autophosphorylation of a CheA-CheY fusion, FrzE, which in turn phosphorylates the response regulator FrzZ (Guzzo *et al.*, 2015). The system also possesses two CheW homologues (FrzA and FrzB), a methyltransferase (FrzF) and methylesterase (FrzG). The chemoreceptor of the Frz pathway, FrzCD, lacks the transmembrane and periplasmic

domains, which are replaced by a N-terminal domain of unknown function (Bustamante *et al.*,

2004). When FrzCD was first localized in cells, it appeared organized in multiple dynamic

cytoplasm clusters that aligned when cells made side-to-side contacts, which has been

proposed to be part of a signaling process that synchronizes cell reversals (Mauriello, Astling,

et al., 2009). However the determinants of FrzCD localization and its exact link with the

regulation of the cell reversal are still unclear.

In this work we show that FrzCD forms cytoplasmic signaling clusters by directly binding to

the nucleoid. Analogously to membrane chemotaxis clusters in *E. coli*, nucleoid binding and

cluster formation confer to the Frz system important regulatory properties such as signal

amplification. We propose that the nucleoid functions as a scaffold for the formation of

bacterial chemosensory complexes like the membrane for enteric chemosensory systems.

RESULTS

The Frz chemosensory system co-localizes with the nucleoid of different bacterial species

Fluorescence microscopy has shown that a FrzCD-GFP fusion appears as multiple distributed

clusters co-localizing with the nucleoid in *M. xanthus* cells (Figure 1A, 1B and Figure 1-

figure supplement 1) (Mauriello, Astling, *et al.*, 2009; Moine *et al.*, 2014; Kaimer and

Zusman, 2016). It has been also shown that an inducible FrzE-YFP fusion colocalizes with

FrzCD at the nucleoid (Kaimer and Zusman, 2016). To further analyze the co-localization of

FrzCD and FrzE with the nucleoid, we first constructed a strain expressing a *frzE-mcherry*

fusion that replaced the *frzE* locus and is, thus, expressed under its endogenous promoter. The

chimeric protein was functional (Figure 1-figure supplement 2) and formed clusters very

similar to that of FrzCD and also co-localizing with the *M. xanthus* nucleoid (Figure 1A, 1B

and Figure 1-figure supplement 1).

To directly show that the nucleoid supports Frz protein localization, we constructed a *M.*

xanthus conditional mutant that lacked ParB, a protein important for nucleoid segregation

whose absence causes the presence of anucleated cells, cells with abnormal nucleoid

condensation and cells where the division septum is improperly positioned over the nucleoid

("guillotines") (Harms *et al.*, 2013; Iniesta, 2014). When *frzCD-gfp* and *frzE-mCherry* were

expressed in the *parB* mutant, we observed that both FrzCD and FrzE clusters always co-

localized with the nucleoid and cells lacking the nucleoid also lacked Frz clusters. Similarly,

in cells with "guillotines", septa formed in regions occupied by both the nucleoid and Frz

clusters instead than in DNA-free regions (Figure 1E).

As a control, we looked at the cellular localization of another chemoreceptor fusion, DifA-

GFP, in the absence of *parB*. DifA has been recently shown to form membrane bound and

uniformly distributed clusters (Moine *et al.*, 2014). In the absence of *parB*, even cells without

nucleoid still carried DifA-GFP clusters and these clusters localized similar to the wild type

(Moine *et al.*, 2014) (Figure 1E). These results confirmed that nucleoid-mediated cluster

formation is a specific feature of Frz proteins.

To test if FrzCD and FrzE were capable of associating with the nucleoid independently of

each other, we expressed *frzE-mcherry* in a strain lacking *frzCD* and *frzCD-gfp* in a strain

lacking *frzE* (Mauriello et al., 2009 and this study). As previously shown, in the absence of

FrzCD, FrzE-mCherry was homogeneously dispersed in the cytoplasm, and notably also in

the polar regions (Figure 1A, 1B and Figure 1-figure supplement 1) (Mauriello, Astling, *et al.*,

2009; Kaimer and Zusman, 2016). Here we additionally showed that in the absence of *frzE*,

FrzCD was no longer capable to form clusters, but the fluorescent signal was still retained

towards the center of the cell body and strictly co-localized with the nucleoid (Figure 1A-D

and Figure 1-figure supplement 1). The aberrant localization patterns observed in the deletion

mutants were not due to a change in protein levels (Figure 1-figure supplement 3). Thus, this

result confirms that both FrzCD and FrzE are important for cluster formation, whereas FrzCD

is responsible for the recruitment of FrzE to the nucleoid.

To check whether the association between FrzCD and the nucleoid was *M. xanthus*-specific,

we constructed an *E. coli* strain expressing *frzCD-gfp* from a plasmid and under the control of

an IPTG inducible promoter. Under these conditions, FrzCD also co-localized with the

nucleoid and this co-localization was particularly evident in elongated undivided cells

containing multiple segregated nucleoids (Figure 2A-D).

These observations suggest that FrzCD can associate with the chromosomes of different

bacterial species, either directly or by the aid of a docking factor common to *M. xanthus* and

E. coli.

FrzCD directly binds to the DNA

The possibility that FrzCD interacted with the nucleoid was puzzling especially considering

that the direct binding between a chemoreceptor and the DNA has not been reported prior to

this study. To explore this possibility, we generated a 6His-tagged FrzCD version, purified it

from *E. coli* and tested its ability to form complexes with DNA. FrzCD binds directly to DNA

because its presence altered DNA mobility on agarose gels (Figure 3A). Binding did not

require extended DNA fragments because it was also observed with oligomers of DNA as

small as 69 bp (Figure 3-figure supplement 1). Moreover, FrzCD DNA-binding did not seem

to depend on the DNA sequence nor on its GC content (Figure 3-figure supplement 1) as

anticipated by the *in vivo* results showing that FrzCD is distributed all over the nucleoid in the

absence of FrzE (Figure 1A and B).

The shift pattern depended on the FrzCD concentration (Figure 3A). More specifically, the

shift of DNA fragments gradually increased as the FrzCD concentration was increased

(Figure 3A and Figure 3-figure supplement 1). Such migration profiles have been previously

described for proteins that can nucleate on DNA molecules in a non -specific manner, i.e.

some Type Ib ParA-like proteins (Hester and Lutkenhaus, 2007; Castaing *et al.*, 2008).

However, to exclude that such profiles were due to the formation of FrzCD unfolded

aggregates, we checked the oligomerization state of our purified 6His-FrzCD. As expected for

an MCP, FrzCD forms a homogenous dimer in solution, corresponding to a molecular weight

of ~ 90 kDa (Figure 3-figure supplement 1). Last, 6His-FrzCD activated the

autophosphorylation of the FrzE kinase *in vitro*, proving that it is functional (Figure 3-figure

supplement 1).

While FrzCD does not appear to bind specific DNA motifs in vitro, it could bind to specific

sites *in vivo* (perhaps with the help of additional factors), explaining the formation of clusters.

To test this possibility, we performed chromatin-immunoprecipitation (ChIP) experiments

using the *frzCD-gfp* strain and polyclonal GFP antibodies. As expected, FrzCD-GFP but not a

FrzCD variant that cannot bind DNA (see below) was able to co-immunopreipicate

significant amounts of DNA. Deep-sequence (ChIP-Seq) (Fioravanti *et al.*, 2013) of the

immunoprecipitated DNA revealed no enrichment in the pool of DNA fragments obtained

with the ChIP meaning that FrzCD-GFP can bind any DNA sequence from the *M. xanthus*

genome (Figure 3-figure supplement 2; the ChipSeq results have also been deposited on GEO

(https://www.ncbi.nlm.nih.gov/geo/) and an accession number is being created). As a positive

control we used a *parB-yfp* strain (Harms *et al.*, 2013). In this case, as expected, the nucleoid

region corresponding to positions 9,109 to 9,110 Kb and containing *parS* (Harms *et al.*, 2013)

was highly represented in the DNA pool obtained with ParB-YFP (Figure 3-figure

supplement 2). We conclude that FrzCD binds DNA in a non sequence-specific manner to

recruit FrzE and thus form clusters containing Frz signaling complexes.

The FrzCD N-terminal region is required for the FrzCD DNA binding

Beside a very conserved C-terminal methylation domain, FrzCD contains a unique 137

residue N-terminal region (Figure 3B). We then asked whether this region corresponded to the

FrzCD nucleoid-binding domain and tested its ability to form complexes with the DNA in our

gel shift assays. Indeed, the FrzCD N-terminal domain (FrzCD$^{\Delta131\text{-}417}$) also provoked a

mobility shift of DNA fragments of different length, albeit at a lower efficiency (compare

Figure 3A and C). On the other hand, the FrzCD C-terminal methyl-accepting domain alone

(FrzCD$^{\Delta1\text{-}130}$) did not associate with any DNA fragment (Figure 3D) showing that the

methylation domain does not bind DNA.

To confirm a direct interaction between FrzCD and DNA and also better compare the DNA-

binding properties of FrzCD and FrzCD$^{\Delta131\text{-}417}$, we performed DNA-protein interaction

experiments using Bio-Layer Interferometry (BLI), a technique previously used to study

protein-protein interactions (Arlet *et al.*, 2014). FrzCD and DNA interaction was also detected

in this assay (Figure 3E). Consistent with the gel-shift experiments, binding appeared

complex and could not be fitted to a 1:1 interaction model, precluding precise determination

of a K_D. Nevertheless, when we compared the DNA binding curves of FrzCD$^{\Delta131-417}$ and

FrzCD, the results confirmed that FrzCD$^{\Delta131-417}$ binds the DNA at a lower efficiency than

FrzCD (showing slower association and faster dissociation, Figure 3E).

By further analyzing the FrzCD N-terminal domain, we realized that it contains a positively

charged peptide of approximately 30 amino acids followed by a more negative region

predicted to contain alpha helices (Figure 3B and F, Figure 3-figure supplement 3). By

searching for homologs in the UniProtKB/SwissProt Data Base, we found that such FrzCD N-

terminal basic peptide was similar to the basic tail present at the N terminus of eukaryotic

histones (Figure 3-figure supplement 3) whose deletion has been shown to substantially affect

histone-DNA interactions and decrease nucleosome stability (Parra *et al.*, 2006; Iwasaki *et al.*,

2013). To test whether this sequence had a histone-tail-like function in the binding of FrzCD

to DNA *in vitro*, we generated a 6His-tagged FrzCD version only lacking the basic amino

acid sequence from residue 7 to 27 (FrzCD$^{\Delta7-27}$), purified it from *E. coli* and tested its ability

to form complexes with the DNA in our gel shift assays. Remarkably, FrzCD$^{\Delta7-27}$ did not shift

the migration of DNA fragments on agarose gels, similarly to FrzCD$^{\Delta1-130}$ missing the entire

N-terminal domain (Figure 3D and 3G). In the BLI assay, binding of FrzCD$^{\Delta7-27}$ to DNA was

still detectable, however it was severely impaired (Figure 3E). This result suggests that the

positively charged motif is required for efficient DNA binding but it may not be the sole

determinant.

The different DNA binding efficiencies of the four FrzCD, FrzCD$^{\Delta1-130}$, FrzCD$^{\Delta131-417}$ and

FrzCD$^{\Delta7-27}$ alleles were not due to altered protein stability and folding because all

recombinant proteins, except as expected FrzCD$^{\Delta 131\text{-}417}$ (the signaling domain) were able to

activate the autophosphorylation of the FrzE kinase *in vitro* (Figure 3-figure supplement 1).

The binding of FrzCD to the nucleoid is required for FrzCD cluster formation *in vivo*

To check whether the absence of the N terminus or the basic tail also affected the binding of

FrzCD to DNA *in vivo*, we used *M. xanthus* strains expressing *frzCD$^{\Delta 6\text{-}130}$-gfp* or *frzCD$^{\Delta 7\text{-}27}$-*

gfp at the *frzCD* locus (Mauriello *et al.*, 2009 and this study). The FrzCD$^{\Delta 6\text{-}130}$-GFP and

FrzCD$^{\Delta 7\text{-}27}$-GFP fluorescence appeared mostly diffused, also occupying the polar regions

(Figure 1A-D). The two protein fusions could only rarely form short-lived clusters that

localized anywhere in the cells (not only at the central region). In all cases, the aberrant

localization patterns where not due to protein stability (Figure 1-figure supplement 3). In

addition, when FrzCD$^{\Delta 6\text{-}130}$-GFP and FrzCD$^{\Delta 7\text{-}27}$-GFP were produced in *E. coli* they also lost

their co-localization with the nucleoid. However, instead of looking dispersed in the

cytoplasm as in *M. xanthus*, FrzCD$^{\Delta 6\text{-}130}$-GFP and FrzCD$^{\Delta 7\text{-}27}$-GFP were confined at one cell

pole in *E. coli*. It is likely that FrzCD$^{\Delta 6\text{-}130}$-GFP and FrzCD$^{\Delta 7\text{-}27}$-GFP formed aggregates

targeted to the poles due to the absence of the nucleoid anchor (Figure 2).

FrzCD cluster dynamics are confined to small nucleoid areas

To understand how FrzCD clusters are formed along the nucleoid, we analyzed the cluster

dynamics at high temporal resolution. Contrarily to previous assumptions based on lower

resolution analysis (Mauriello, Astling, *et al.*, 2009), this analysis showed that FrzCD clusters

are quite fixed and only featured by Brownian-like motions in highly constrained areas of the

nucleoid (Figure 4A-C). This mobility decreased with the increase of cluster intensity,

suggesting that clusters containing more molecules might be more tightly anchored to the

chromosome and, hence, more fixed (Figure 4D). To test whether the signaling state of

FrzCD also affects the cluster mobility, we tested whether clusters were also constrained in

strains carrying point mutations either generating FrzCD loss of function or, oppositely,

FrzCD gain of function (Astling *et al.*, 2006; Mauriello, Astling, *et al.*, 2009; Guzzo *et al.*,

2015). There were no notable differences between the tested conditions, suggesting that

signaling does not affect the nucleoid dynamics of Frz signaling complexes along the

nucleoid (Figure 4E).

The nucleoid-dependent assembly could promote cooperative interactions between

FrzCD receptors

Consistent with transmembrane chemosensory clusters (Francis *et al.*, 2002; Li *et al.*, 2011;

Li and Hazelbauer, 2011; Piñas *et al.*, 2016), the formation of Frz nucleoid-associated clusters

is not strictly required for signaling. In fact, it has been previously shown that a *frzCD$^{\Delta6-130}$*

strain, where FrzCD molecules can no longer bind DNA and are, thus, diffused (Figure 1),

can still produce reversals (Bustamante et al., 2004). This is not surprising because in *E. coli*

chemosensory cluster formation is also not critical for the signal transduction, but it confers

properties such as the amplification of signal, a direct consequence of the cooperative

interactions between clustered chemoreceptors (Sourjik and Berg, 2004; Ames and Parkinson,

2006; Li and Hazelbauer, 2014; Piñas *et al.*, 2016; Frank *et al.*, 2016). Thus, we decided to

check if the formation of Frz clusters also led to cooperativity in the signaling activity of the

Frz chemosensory system. For this, we took advantage of a newly developed microfluidic

single cell assay where the frequency of reversals can be measured as a function of increasing

concentrations of an artificial Frz-signal activator, the isoamyl alcohol (IAA) (Guzzo *et al.*,

2015). Consistent with previous observations (Guzzo *et al.*, 2015), in cells where FrzCD

formed nucleoid bound clusters, IAA induced a dose-dependent response with a sigmoidal

shaped curve that could be fitted by the Hill equation with a coefficient $n = 3.017$ +/- 0.2 (P =

0.0007), which is significantly higher than one and reveals the presence of cooperativity in

the FrzCD activity (Figure 5A). Such response is FrzCD-dependent because a $\Delta frzCD$ strain

does not reverse at any IAA dose (Figure 5A). In the $frzCD^{\Delta 6\text{-}130}$ mutant, the dose-dependent

response curve to IAA resulted to a Hill coefficient $n = 1.15$ +/- 0.01 (P = 0.008) when it was

fitted by the Hill equation, revealing that cooperation is lost in this mutant. These results

suggest that while the IAA response in the $frzCD^{\Delta 6\text{-}130}$ mutant only depends on the number of

the $FrzCD^{\Delta 6\text{-}130}$ molecules in the cytoplasm, in cells where FrzCD formed nucleoid bound

clusters, such response also depends on the supra-molecular organization of the FrzCD

proteins, presumably because of cooperative protein interactions in the clusters (Lai *et al.*,

2005; Li and Hazelbauer, 2014; Piñas *et al.*, 2016). This signal amplification is advantageous

to *M. xanthus* social behaviors because swarming and predation are defective in the $frzCD^{\Delta 6\text{-}}$

^{130} mutant compared to the wild type (Figure 5B and C).

DISCUSSION

In this work, we show that analogously to how transmembrane chemoreceptors use the

bacterial inner membrane as a platform to form the well described arrays of trimers of dimers

(Briegel *et al.*, 2012b), the *M. xanthus* Frz system forms signaling clusters on the bacterial

chromosome. Cluster assembly is directed by the chemoreceptor FrzCD, which binds to the

DNA by a N-terminal domain carrying a positively charged tail similar to that found in

eukaryotic histones (Parra *et al.*, 2006; Iwasaki *et al.*, 2013). While the binding of FrzCD to

DNA is essential to target the Frz chemosensory system to the nucleoid, it is not sufficient for

Frz cluster formation, as it requires downstream interactions with the FrzE kinase. Because

FrzCD appears to bind DNA in a non-sequence specific manner, DNA-bound clusters do not

occupy fixed localization sites and move across small areas on the nucleoid surface.

Analogous to trans-membrane proteins diffusing in the bacterial membrane, the FrzCD cluster

dynamic behavior may be affected by the size of the complex (and thus the number of

interactions with DNA), explaining why bright clusters show Brownian-like motions that only

explore constrained nucleoid areas.

Several lines of evidence suggest that the Frz cluster formation on the nucleoid occurs in a

stochastic manner similarly to the assembly of the *E. coli* Che lattices in the membrane. First,

the initial binding of FrzCD to DNA might take place anywhere on the nucleoid as such

binding is not DNA-sequence specific. Once recruited to the nucleoid, small FrzCD foci

diffuse, non-directionally, across confined small areas until they might nucleate large fixed

clusters by attracting more FrzCD molecules. "Newborn" FrzCD foci might also, at one point,

be incorporated by existing neighboring clusters. Thus, the areas explored by FrzCD clusters

might represent the minimal critical distance from other clusters at which foci can exist. The

existence of such minimal critical distance is supported by the fact that the number of Frz

clusters increases linearly with the nucleoid size (Figure 1-figure supplement 4), suggesting

that more clusters can form when more surface becomes available. Thus, like for transmembrane chemoreceptors, FrzCD molecules might either nucleate new dynamic foci if they are far enough from existing clusters, or encounter and join neighboring clusters.

Transmembrane chemoreceptors are arrayed in a monolayer tightly associated with the inner membrane (Briegel *et al.*, 2009). However, in the absence of a scaffold *Vibrio sp.* and *R. sphaeroides* cytoplasmic chemosensory arrays, are instead organized in two sandwiched arrays (Briegel *et al.*, 2014; Briegel *et al.*, 2015; Briegel *et al.*, 2016). Our data suggest that the *M. xanthus* chromosome functions as a platform to form signaling clusters, much like the inner membrane supports transmembrane chemosensory complexes. In the future, it will be interesting to test if the Frz proteins, indeed, form single layered arrays at the surface of the chromosome by Cryo-EM tomography.

But, what is the outcome of this nucleoid driven cluster assembly? FrzCD cluster formation has been previously been associated to regulation of the cell reversal frequency in response to cell-cell contact. In fact, FrzCD clusters align in adjacent *M. xanthus* cells, a behavior that also seemed to induce cell reversals (Mauriello, Astling, *et al.*, 2009). The authors' argument in favor of a retro-regulated FrzCD cluster rearrangement in response to cell contact was that adjacent cells of a *frzE* strain, which also seemed to form FrzCD-GFP clusters even if more diffused, did not produce cluster alignment. In light of our new results, what seemed to be more diffused FrzCD clusters in the *frzE* strain, are, in fact, FrzCD molecules dispersed on the nucleoid. Moreover, we now show that the Frz cluster organization is independent on the Frz signaling activity. Thus, the observed FrzCD cluster alignment might be more likely determined by similar dynamic rearrangements of the nucleoid of adjacent cells rather than by an active regulated mechanism.

Therefore, a simpler outcome of the nucleoid-driven Frz cluster assembly could be to confer to a cytoplasmic receptor the universal properties of transmembrane receptors but here, in

response to intracellular signals. One of these properties is the signal amplification, which, in enteric bacteria, strictly requires the MCP clustering to the membrane. Remarkably, this work as well as previous studies shows that wild type *M. xanthus* cells respond to increasing concentrations of the Frz activator IAA with a dose-dependent response curve typical of a cooperative response that strongly suggests the presence of signal amplification. Analogously to enteric bacteria, signal amplification must be due to the supra-molecular organization of FrzCD receptors on the nucleoid, because such property is lost in FrzCD$^{\Delta 6-130}$ lacking the nucleoid binding domain. In cells of this strain, the reversal frequencies increase linearly with the IAA doses, suggesting that, in this case, signaling is only function of the number of activated receptor-signaling complexes dispersed in the cytoplasm.

Why does FrzCD need to form many and not just one single cluster? One possibility could be to prevent the diffusion of CheY-P to be a limiting factor in the control of the polar and lateral motility motors due to the length of *M. xanthus* cells (5-10 μm in average). However, a more attractive explanation could be that the nucleoid-dependent formation of multiple distributed clusters represents a simple mechanism to segregate clusters during cell division without the need for a faithful partitioning system (Figure 1-figure supplement 5). Such partitioning system would be, in fact, likely required in the presence of a single cluster like in *R. sphaeroides* where PpfA ensures the faithful segregation of Tlp clusters (Thompson *et al.*, 2006; Ringgaard *et al.*, 2011).

Finally, the analysis of the FrzCD sequences from some related species of δ-proteobacteria shows that while the FrzCD C-terminal region is very conserved, its N-terminus largely varies. Nevertheless, the FrzCD N-terminus always shows a positively charged sequence (Figure 3-figure supplement 4) suggesting that the non sequence-specific recruitment of Frz

proteins to the nucleoid essentially requires the presence of a positively charged protein

domain rather than a specific amino acid sequence.

This type of cellular organization may be common to other bacterial macromolecular

complexes to provide important regulatory functions. In this sense, the Frz example provides

new perspectives to the role of the bacterial nucleoid as a scaffold for the spatial control of

cellular functions.

MATERIALS AND METHODS

Bacterial strains, plasmids and growth

Strains and plasmids are listed in Table S1 and S2. *M. xanthus* strains were grown at 32°C in

CYE rich media as previously described. *Pcuo::parB-ΔparB* cells were grown at 32°C in

CTT supplied with 300 μM CuSO₄.

Plasmids were introduced into *M. xanthus* cells by electroporation. Deletions and GFP fusions

were inserted in frame to avoid polar effects on the downstream gene expression. These

strains were obtained by homologous recombination based on a previously reported method

using the pBJ113 or pBJ114 vectors (Bustamante *et al.*, 2004; Moine *et al.*, 2014). To

generate strains expressing GFP or mCherry fusion proteins, we constructed DNA cassettes

including the last approximately 800 bp of each gene, with the exception of the stop codon;

the gene encoding the *egfp* or *mcherry* genes from pEGFP-N1 (Invitrogen) or pEM147

(Mauriello, Nan, *et al.*, 2009) excluding the start codon and including the stop codon; the

intergenic region between the gene of interest and its immediately downstream gene, if any;

the first 800 bp of the downstream gene.

To construct the *parB* conditional mutant, we transferred the previously described *parB*

conditional depletion (Harms *et al.*, 2013) in our wild type strain, DZ2 (Harms *et al.*, 2013;

Iniesta, 2014).

To construct strains carrying a FrzS-YFP fusion, we used the pEFrzSY plasmid (Guzzo *et al.*,

2015)

Escherichia coli cells were grown under standard laboratory conditions in Luria-Bertani broth

supplemented with antibiotics if necessary.

For swarming assays, cells (5 μl) at a concentration of 5×10^9 cfu ml^{-1} were spotted on CYE

agar plates and incubated at 32°C and photographed after respectively 48h with an Olympus

SZ61 binocular stereoscope. For predation assays, *E. coli* (3μl at a concentration of 5×10^9 cfu

ml^{-1}) and *M. xanthus* cells (3µl at a concentration of 5×10^9 cfu ml^{-1}) were spotted at 0.7 mm

distance from each other on CF agar plates, incubated at 32°C and photographed after 72 h.

Protein purification

BL21(DE3) [F$^-$ ompT hsdSB(rB$^-$ mB$^-$) gal dcm (DE3)] cells were grown in Luria-Bertani

broth supplemented 100 µg/ml ampicillin to mid-exponential phase at 37°C. For the

experiments shown on Figure 3 and phosphorylation assays on Figure 3-figure supplement 1,

overexpression was induced by adding 0,1 mM IPTG for cells containing plasmid pEM414 or

0,5mM for pEM415 and pEM433. Cells were then grown at 16°C over night. Cells were

washed and resuspended in lysis buffer (50 mM TrisHCl, pH 8; 300 mM NaCl; 100 µg/ml

PMSF; 30 U/mL Benzonase) and lysed at the French press. The cell lysates were centrifuged

at 4°C for 30 min at 13000× rpm. Soluble tagged His$_6$-proteins were purified on 1ml

HisTrap^{TM}FF columns (GE Healthcare) and desalted with PD-10 columns (GE Healthcare).

Ultimately, purified proteins were eluted in 50 mM TrisHCl, pH 8 and 300 mM NaCl.

For the experiments shown on Figure 3-figure supplement 1A-E, the C-terminal-tagged

FrzCD-(His)$_6$ was expressed in *E. coli* BL21-AI (Invitrogen) cells. The cells were grown in

Luria-Bertani broth containing 100 µg/ml ampicillin at 37°C till OD$_{600}$ of 0.8 − 1.0, and

incubated at 30°C for 5 hours post induction. The harvested cells were resuspended in Buffer

A (50 mM Tris, 200 mM NaCl, pH 8.0) containing 10 % glycerol and lysed by sonication

(Sonics VibraCell, 5 minutes, 60% amplitude, 1" ON, 3" OFF cycle). The cell lysate was

spun at 39, 191 g for an hour and the supernatant loaded to a 5-ml HisTrap^{TM} FF (GE

Healthcare) equilibrated with Buffer A. Following wash and elution with increasing

concentrations (5%, 10%, 20%, 50% and 100%) of Buffer B (50 mM Tris, 200 mM NaCl,

500 mM imidazole, pH 8.0), pure fractions containing FrzCD-H$_6$ were pooled and dialysed

into buffer containing 50 mM Tris, 25 mM NaCl, 1 mM EDTA, pH 8.0 to enable further

purification through ion exchange chromatography using a MonoQ-PE (GE Healthcare)

column. The fractions containing the protein were pooled and concentrated using centricons.

The protein was finally in buffer containing 50 mM NaCl, 1 mM EDTA and 50 mM Tris pH

8.0 either through dialysis or by buffer exchange during the concentration step. The

concentrated protein was flash frozen and stored in small aliquots in -80°C till further use.

Electrophoretic mobility shift assays (EMSA)

EMSAs on Figure 3 were carried out by incubating different concentrations of purified

proteins with 10 nM PCR-amplification products of different sizes (Table S4), in buffer (10

mM of TrisHCl at pH 8; 60 mM of NaCl; 10% glycerol). Reactions were incubated for 40

min at 4°C before being loaded on 1% agarose gels. Gel migration was performed in 1X TBE

at 4°C for 55 min. Gels were, then, stained with ethidium bromide and revealed at the UV

light.

EMSAs for shorter oligos (Figure 3-figure supplement 1 and Table S4) were carried out

in buffer containing 50 mM Tris, 50 mM NaCl, 1 mM dithiothreitol and 10 % glycerol, pH

7.4. All the oligonucleotide fragments were PCR-amplification products, except the 69 bp

DNA that was generated by annealing the custom-synthesized oligonucleotides (Sigma)

corresponding to the 5' to 3' sequence and its complementary strand (Table S4). The samples

were incubated at 25°C for 20 min and loaded onto the appropriate % of agarose gel (1% for

1.3 kbp, and electrophoresed for 60 – 90 minutes in 1x TAE (40 mM Tris-Acetate and 1 mM

EDTA).

Oligomerization study using Size Exclusion Chromatography coupled with Multi-Angle

light Scattering (SEC-MALS)

The expected elongated structure of an MCP precludes estimation of the oligomeric status by

size exclusion chromatography alone, and hence we carried out SEC coupled with multi-angle

light scattering to estimate the molecular mass in solution. The mass was determined using a

Wyatt Dawn Heleos II equipped with light scattering detectors at 18 angles and an Optilab

TrEX differential refractive index detector. Protein sample (100 µl of 2 mg/ml solution or 45

µM) was injected into the size exclusion column (BioRad EnRich650) equilibrated with the

buffer 50 mM Tris, 50 mM NaCl, pH 8.0, and the run carried out at a flow rate of 0.4 ml/min.

The observed molecular masses at various points along the peak in the elution curve were

calculated using the protein concentration estimated from the differential refractive index

(dRI), and measured scattered intensities. The Debye model in the ASTRA software provided

with the equipment was used for fitting the data.

Biolayer interferometry

Protein-DNA interaction experiments were conducted at 25°C with the BLItz instrument from

ForteBio (Menlo Park, CA, USA). The BLI consists in a real time optical biosensing

technique exploits the interference pattern of white light reflected from two surfaces to

measure biomolecular interactions (Concepcion *et al.*, 2009). Purified 6His-FrzCD, 6His-

FrzCD$^{\Delta131\text{-}417}$, 6His-FrzCD$^{\Delta6\text{-}130}$ and 6His-FrzCD$^{\Delta7\text{-}27}$ protein ligands were immobilized onto

two different Ni-NTA biosensors (ForteBio) in duplicate at 1µM concentrations. A PCR

amplified DNA fragment (474bp) with primers AGACCCCCGCACCCACGGAG and

TCACGCGGGCTCCGGCTC (Eurogentec) was used as the analyte throughout the study at

the 38nM. The assay was conducted in PBS pH 7.5, 0.001% tween-20. The binding reactions

were performed with an initial baseline during 30 seconds, an association step at 120 seconds

and a dissociation step of 120 seconds with lateral shaking at 2200rpm. A double reference

subtraction (sensor reference and 6His-FrzCD$^{\Delta 1\text{-}130}$) was applied to account for non-specific

binding, background, and signal drift to minimize sensor variability.

Chromatin Immunoprecipitation-deep sequencing (ChIp-seq)

ChIp-seq was performed as previously described (Fioravanti *et al.*, 2013). In particular, mid-

log phase cells (80 ml, OD_{600} of 0.6) were cross-linked in 10 mM sodium phosphate (pH 7.6)

and 1% formaldehyde at room temperature for 10 min and on ice for 30 min thereafter,

washed thrice in phosphate buffered saline (PBS) and lysed with lysozyme 2.2 mg ml^{-1} in

TES (Tris-HCl 10 mM pH 7.5, EDTA 1 mM, NaCl 100 mM). Lysates (Final volume 1ml)

were sonicated (Branson Digital Sonicator 450, Branson Sonic Power. Co.,

www.bransonic.com/) on ice using 10 bursts of 30 sec (50% duty) at 30% amplitude to shear

DNA fragments to an average length of 0.3–0.5 kbp and cleared by centrifugation at 14,000

rpm for 2 min at 4°C. Lysates were normalized by protein content by measuring the

absorbance at 280 nm; ca. 7.5 mg of protein was diluted in 1 mL of ChIP buffer (0.01% SDS,

1.1% Triton X-100, 1.2 mM EDTA, 16.7 mM Tris-HCl [pH 8.1], 167 mM NaCl plus protease

inhibitors (Euromedex, https://www.euromedex.com/) and pre-cleared with 80 μL of protein-

A agarose (Sigma-Aldrich, www.sigmaaldrich.com) and 100 μg BSA. Polyclonal GFP

antibodies were added to the remains of the supernatant (1:1,000 dilution), incubated

overnight at 4°C with 80 μL of protein-A agarose beads pre-saturated with BSA, washed once

with low salt buffer (0.1% SDS, 1% Triton X-100, 2 mM EDTA, 20 mM Tris-HCl (pH 8.1),

150 mM NaCl), high salt buffer (0.1% SDS, 1% Triton X-100, 2 mM EDTA, 20 mM Tris-

HCl (pH 8.1), 500 mM NaCl) and LiCl buffer (0.25 M LiCl, 1% NP-40, 1% sodium

deoxycholate, 1 mM EDTA, 10 mM Tris-HCl (pH 8.1) and twice with TE buffer (10 mM

Tris-HCl (pH 8.1) and 1 mM EDTA). The protein-DNA complexes were eluted in 500 μL

freshly prepared elution buffer (1% SDS, 0.1 M NaHCO3), supplemented with NaCl to a final

concentration of 300 mM and incubated overnight at 65°C to reverse the crosslinks. The

samples were treated with 2 µg of Proteinase K for 2 h at 45°C in 40 mM EDTA and 40 mM

Tris-HCl (pH 6.5). DNA was extracted using QIAgen minelute kit and resuspended in 30 µl

of Elution Buffer. ChIp DNA sequencing was performed using Illumina MiSeq and analyzed

using Galaxy Web Portal (usegalaxy.org/). Reads were analyzed by MatLab.

Protein sequence analyses

In order to search for homologs of the FrzCD N-terminal domain, the first 130 aminoacids of

the FrzCD sequence were BLAST into the UniProtKB/SwissProt Data Base

(http://blast.ncbi.nlm.nih.gov). Predictions of secondary structures and protein sequence

alignments were obtained with Jpred (Cole *et al.*, 2008) and Clustal Omega (Sievers and

Higgins, 2002), respectively. To analyze the FrzCD N-terminal region protein charge,

"Sliding window" analyses were performed with Microsoft Excel.

***In vitro* autophosphorylation assay**

In vitro phosphorylation assays were performed with *E. coli* purified recombinant proteins. 4

µg of FrzE^{kinase} were incubated with 1µg of FrzA and increasing concentrations (3µg) of

different FrzCD proteins (entire FrzCD, FrzCD^c, FrzCDΔ6-130, FrzCDΔ7-27 or FrzCDΔ131-

417) in 25 µl of buffer P (50 mM Tris-HCl, pH 7.5; 1 mM DTT; 5 mM MgCl$_2$; 50mM KCl; 5

mM EDTA; 50µM ATP, 10% glycerol) supplemented with 200 µCi ml^{-1} (65 nM) of [γ-

33P]ATP (PerkinElmer, 3000 Ci mmol^{-1}) for 10 minutes at room temperature in order to

obtain the optimal FrzE^{kinase} autophosphorylation activity. Each reaction mixture was stopped

by addition of 5 × Laemmli and quickly loaded onto SDS-PAGE gel. After electrophoresis,

proteins were revealed using Coomassie Brilliant Blue before gel drying. Radioactive proteins

were visualized by autoradiography using direct exposure to film (Carestream).

Fluorescence microscopy and image analysis

For fluorescence microscopy analyses, 5 μl of cells from 4 x 10^8 cfu ml^{-1} vegetative CYE

cultures were spotted on a thin fresh TPM agar pad at the top a slide (Mignot *et al.*, 2005). A

cover slip was added immediately on the top of the pad, and the obtained slide was analyzed

by microscopy using a Nikon Eclipse TE2000 E PFS inverted epifluorescence microscope

(100 x oil objective NA 1.3 Phase Contrast) (Ducret *et al.*, 2009).

To study the colocalization with the DNA, the TPM agar pads were supplied with 1μg/ml

DAPI stain and 1 mM IPTG. Prior to imaging, *E. coli* cells were grown in 1 mM IPTG for

one hour then spotted on agar pads containing or not 10μg/ml cephalexin and incubated for 1

hour. Cell fluorescence profiles were obtained with the "plot profile" function of FIJI

(Schindelin *et al.*, 2012). FrzCD clusters numbers and distances, nucleoid areas and cell areas

were automatically determined and verified manually with the "MicrobeJ" Fiji/ImageJ plugin

created by A. Ducret, Brun Lab (http://www.indiana.edu/~microbej/index.html). All data

plots and statistical tests were obtained with the R software (https://www.r-project.org/).

To study the dynamic of FrzCD-GFP clusters we automatically tracked clusters (imaged

every second) by MicrobeJ and recorded parameters such as the mean square displacement

(MSD), the confinement radius and the fluorescence intensity along the time.

For manual image analysis we chose a sample size that allowed an error on the mean (sem)

lower than 10%. For results generated by automated analyses, we analyzed all the available

samples. We chose all cells where clusters and nucleoids were tractable by our image analysis

tools (Fiji and MicrobeJ). For the cluster dynamic determinations we analyzed all clusters that

were tractable for at least 5 consecutive frames.

Reversal frequencies

These assays were performed as previously described (Guzzo *et al.*, 2015) by using

homemade PDMS glass microfluidic chambers (Ducret *et al.*, 2013) treated with 0.015%

carboxymethylcellulose after extensive washing of the glass slide with water. For each

experiment, 1mL of a CYE grown culture of OD = 0.5 was injected directly into the chamber

and the cells were allowed to settle for 5 min. Motility was assayed after the chamber was

washed with TPM 1mM CaCl buffer. For IAA injections, IAA solutions made in TPM 1mM

CaCl buffer at appropriate concentrations were injected directly into the channels and motility

was assayed directly under the microscope. Time-lapse movies of strains carrying a FrzS-YFP

fusion were shot for 20 minutes with frames captured every 15 seconds.

To discriminate bona fide reversal events from stick-slip motions (Guzzo *et al.*, 2015), the

fluorescence intensity of FrzS-YFP was measured at cell poles over time. In fact, this protein

has been shown to switch from the leading cell pole to the lagging pole when *M. xanthus* cells

reverse their movement direction (Mignot *et al.*, 2005). For each cell that was tracked, the

fluorescence intensity and reversal profiles were correlated to distinguish bona fide reversals

from stick-slip events. About one hundred cells for the wild type and *frzCD$^{Δ1-130}$* strains and

fifty for the *ΔfrzCD* strain were analyzed (refer to Supplementary Table 3 for the exact

number of cells analyzed for each strain and IAA dose). The number of reversals was plotted

against time. The best fits for the reversal frequencies values at the different IAA doses were

obtained with the following Hill equation:

$$\frac{N_{rev}}{\Delta T} = \frac{[IAA]^n}{K_d + [IAA]^n} B + A$$

where the $N_{rev}/\Delta T$ is the number of reversal events per hour; K_d is the apparent affinity

constant; *[IAA]* is the IAA dose; *B* is the plateau; *A* is the basal reversal frequency and *n* is the

Hill coefficient describing cooperativity. Reversal frequency values for each IAA dose and

each strain are the results of two independent biological triplicates.

We chose a sample size that allowed an error on the mean (sem) lower than 10%. We used all

cells that moved, remained isolated for at least 20 consecutive frames and where FrzS-YFP

foci were detectable by FIJI.

Acknowledgments

We would like to thank Dr. Romain Mercier and Dr. Thierry Doan for their critical reading of the manuscript and discussions; Dr. Mireille Ansaldi for her advise on the EMSA assays; Hanna Bismuth for helping with the ChIP experiments; Pravin Dewangan and Dr. Radha Chauhan for use of the SEC-MALS facility at National Centre for Cell Science, Pune, India.

Research on chemotaxis in our laboratory is funded by the Agence National de la Recherche Jeune Chercheur-Jeune Chercheuse (ANR-14-CE11-0023-01) to E.M.M and the "Fondation Amidex" award to EM and TM. YM, PJJ and PG acknowledge fellowships from INSPIRE, Department of Science and Technology (DST), Govt. of India and the work in the lab at IISER Pune is supported by Innovative Young Biotechnologist Award (IYBA from Department of Biotechnology, Govt. of India), Indian National Science Academy (INSA), and Science and Engineering Research Board (SERB), DST.

BIBLIOGRAPHY

Ames, P., and Parkinson, J.S. (2006) Conformational suppression of inter-receptor signaling
defects. *Proc Natl Acad Sci U S A* **103**: 9292–9297.
Arlet, J.-B., Ribeil, J.-A., Guillem, F., Negre, O., Hazoume, A., Marcion, G., *et al.* (2014)
HSP70 sequestration by free α-globin promotes ineffective erythropoiesis in β-thalassaemia.
Nature **514**: 242–246.
Astling, D.P., Lee, J.Y., and Zusman, D.R. (2006) Differential effects of chemoreceptor
methylation-domain mutations on swarming and development in the social bacterium
Myxococcus xanthus. *Mol Microbiol* **59**: 45–55.
Berleman, J.E., and Bauer, C.E. (2005) Involvement of a Che-like signal transduction cascade
in regulating cyst cell development in Rhodospirillum centenum. *Mol Microbiol* **56**: 1457–
1466.
Bible, A., Russell, M.H., and Alexandre, G. (2012) The Azospirillum brasilense Che1
chemotaxis pathway controls swimming velocity, which affects transient cell-to-cell
clumping. *J Bacteriol* **194**: 3343–3355.
Blackhart, B.D., and Zusman, D.R. (1985) "Frizzy" genes of Myxococcus xanthus are
involved in control of frequency of reversal of gliding motility. *Proc Natl Acad Sci U S A* **82**:
8767–8770.
Briegel, A., Ladinsky, M.S., Oikonomou, C., Jones, C.W., Harris, M.J., Fowler, D.J., *et al.*
(2014) Structure of bacterial cytoplasmic chemoreceptor arrays and implications for
chemotactic signaling. *eLife* **3**: e02151.
Briegel, A., Li, X., Bilwes, A.M., Hughes, K.T., Jensen, G.J., and Crane, B.R. (2012a)
Bacterial chemoreceptor arrays are hexagonally packed trimers of receptor dimers networked
by rings of kinase and coupling proteins. *Proc Natl Acad Sci U S A* **109**: 3766–3771.
Briegel, A., Li, X., Bilwes, A.M., Hughes, K.T., Jensen, G.J., and Crane, B.R. (2012b)
Bacterial chemoreceptor arrays are hexagonally packed trimers of receptor dimers networked
by rings of kinase and coupling proteins. *Proc Natl Acad Sci U S A* **109**: 3766–3771.
Briegel, A., Ortega, D.R., Huang, A.N., Oikonomou, C.M., Gunsalus, R.P., and Jensen, G.J.
(2015) Structural conservation of chemotaxis machinery across Archaea and Bacteria.
Environ Microbiol Rep **7**: 414–419.
Briegel, A., Ortega, D.R., Mann, P., Kjær, A., Ringgaard, S., and Jensen, G.J. (2016)
Chemotaxis cluster 1 proteins form cytoplasmic arrays in Vibrio cholerae and are stabilized
by a double signaling domain receptor DosM. *Proc Natl Acad Sci U S A* **113**: 10412–10417.
Briegel, A., Ortega, D.R., Tocheva, E.I., Wuichet, K., Li, Z., Chen, S., *et al.* (2009) Universal
architecture of bacterial chemoreceptor arrays. *Proc Natl Acad Sci U S A* **106**: 17181–17186.
Bustamante, V.H., Martinez-Flores, I., Vlamakis, H.C., and Zusman, D.R. (2004) Analysis of
the Frz signal transduction system of Myxococcus xanthus shows the importance of the
conserved C-terminal region of the cytoplasmic chemoreceptor FrzCD in sensing signals. *Mol*
Microbiol **53**: 1501–13.
Castaing, J.-P., Bouet, J.-Y., and Lane, D. (2008) F plasmid partition depends on interaction
of SopA with non-specific DNA. *Mol Microbiol* **70**: 1000–1011.
Cole, C., Barber, J.D., and Barton, G.J. (2008) The Jpred 3 secondary structure prediction
server. *Nucleic Acids Res* **36**: W197–W201.
Concepcion, J., Witte, K., Wartchow, C., Choo, S., Yao, D., Persson, H., *et al.* (2009) Label-
free detection of biomolecular interactions using BioLayer interferometry for kinetic
characterization. *Comb Chem High Throughput Screen* **12**: 791–800.
Ducret, A., Maisonneuve, E., Notareschi, P., Grossi, A., Mignot, T., and Dukan, S. (2009) A
microscope automated fluidic system to study bacterial processes in real time. *PloS One* **4**:
e7282.
Ducret, A., Théodoly, O., and Mignot, T. (2013) Single cell microfluidic studies of bacterial

motility. *Methods Mol Biol Clifton NJ* **966**: 97–107.
Fioravanti, A., Fumeaux, C., Mohapatra, S.S., Bompard, C., Brilli, M., Frandi, A., *et al.*
(2013) DNA Binding of the Cell Cycle Transcriptional Regulator GcrA Depends on N6-
Adenosine Methylation in Caulobacter crescentus and Other Alphaproteobacteria. *PLOS*
Genet **9**: e1003541.
Francis, N.R., Levit, M.N., Shaikh, T.R., Melanson, L.A., Stock, J.B., and DeRosier, D.J.
(2002) Subunit organization in a soluble complex of tar, CheW, and CheA by electron
microscopy. *J Biol Chem* **277**: 36755–36759.
Frank, V., Piñas, G.E., Cohen, H., Parkinson, J.S., and Vaknin, A. (2016) Networked
Chemoreceptors Benefit Bacterial Chemotaxis Performance. *mBio* **7**.
Guzzo, M., Agrebi, R., Espinosa, L., Baronian, G., Molle, V., Mauriello, E.M.F., *et al.* (2015)
Evolution and Design Governing Signal Precision and Amplification in a Bacterial
Chemosensory Pathway. *PLoS Genet* **11**: e1005460.
Harms, A., Treuner-Lange, A., Schumacher, D., and Søgaard-Andersen, L. (2013) Tracking
of chromosome and replisome dynamics in Myxococcus xanthus reveals a novel chromosome
arrangement. *PLoS Genet* **9**: e1003802.
Hester, C.M., and Lutkenhaus, J. (2007) Soj (ParA) DNA binding is mediated by conserved
arginines and is essential for plasmid segregation. *Proc Natl Acad Sci U S A* **104**: 20326–
20331.
Iniesta, A.A. (2014) ParABS system in chromosome partitioning in the bacterium
Myxococcus xanthus. *PLoS One* **9**: e86897.
Iwasaki, W., Miya, Y., Horikoshi, N., Osakabe, A., Taguchi, H., Tachiwana, H., *et al.* (2013)
Contribution of histone N-terminal tails to the structure and stability of nucleosomes. *FEBS*
Open Bio **3**: 363–369.
Kaimer, C., and Zusman, D.R. (2016) Regulation of cell reversal frequency in Myxococcus
xanthus requires the balanced activity of CheY-like domains in FrzE and FrzZ. *Mol Microbiol*
100: 379–395.
Lai, R.-Z., Manson, J.M.B., Bormans, A.F., Draheim, R.R., Nguyen, N.T., and Manson, M.D.
(2005) Cooperative signaling among bacterial chemoreceptors. *Biochemistry (Mosc)* **44**:
14298–14307.
Li, M., and Hazelbauer, G.L. (2011) Core unit of chemotaxis signaling complexes. *Proc Natl*
Acad Sci U S A **108**: 9390–9395.
Li, M., and Hazelbauer, G.L. (2014) Selective allosteric coupling in core chemotaxis
signaling complexes. *Proc Natl Acad Sci U S A* **111**: 15940–15945.
Li, M., Khursigara, C.M., Subramaniam, S., and Hazelbauer, G.L. (2011) Chemotaxis kinase
CheA is activated by three neighbouring chemoreceptor dimers as effectively as by receptor
clusters. *Mol Microbiol* **79**: 677–685.
Liu, J., Hu, B., Morado, D.R., Jani, S., Manson, M.D., and Margolin, W. (2012) Molecular
architecture of chemoreceptor arrays revealed by cryoelectron tomography of Escherichia coli
minicells. *Proc Natl Acad Sci U S A* **109**: E1481-1488.
Martin, A.C., Wadhams, G.H., and Armitage, J.P. (2001) The roles of the multiple CheW and
CheA homologues in chemotaxis and in chemoreceptor localization in Rhodobacter
sphaeroides. *Mol Microbiol* **40**: 1261–1272.
Mauriello, E.M.F., Astling, D.P., Sliusarenko, O., and Zusman, D.R. (2009) Localization of a
bacterial cytoplasmic receptor is dynamic and changes with cell-cell contacts. *Proc Natl Acad*
Sci U S A **106**: 4852–4857.
Mauriello, E.M.F., Nan, B., and Zusman, D.R. (2009) AglZ regulates adventurous (A-)
motility in Myxococcus xanthus through its interaction with the cytoplasmic receptor, FrzCD.
Mol Microbiol **72**: 964–977.
Mignot, T., Merlie, J.P., and Zusman, D.R. (2005) Regulated pole-to-pole oscillations of a

bacterial gliding motility protein. *Science* **310**: 855–7.

Moine, A., Agrebi, R., Espinosa, L., Kirby, J.R., Zusman, D.R., Mignot, T., and Mauriello, E.M.F. (2014) Functional organization of a multimodular bacterial chemosensory apparatus. *PLoS Genet* **10**: e1004164.

Parra, M.A., Kerr, D., Fahy, D., Pouchnik, D.J., and Wyrick, J.J. (2006) Deciphering the Roles of the Histone H2B N-Terminal Domain in Genome-Wide Transcription. *Mol Cell Biol* **26**: 3842–3852.

Piñas, G.E., Frank, V., Vaknin, A., and Parkinson, J.S. (2016) The source of high signal cooperativity in bacterial chemosensory arrays. *Proc Natl Acad Sci U S A* **113**: 3335–3340.

Ringgaard, S., Schirner, K., Davis, B.M., and Waldor, M.K. (2011) A family of ParA-like ATPases promotes cell pole maturation by facilitating polar localization of chemotaxis proteins. *Genes Dev* **25**: 1544–1555.

Ringgaard, S., Zepeda-Rivera, M., Wu, X., Schirner, K., Davis, B.M., and Waldor, M.K. (2014) ParP prevents dissociation of CheA from chemotactic signaling arrays and tethers them to a polar anchor. *Proc Natl Acad Sci U S A* **111**: E255-264.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., *et al.* (2012) Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9**: 676–682.

Sievers, F., and Higgins, D.G. (2002) Clustal Omega. In *Current Protocols in Bioinformatics*. John Wiley & Sons, Inc., http://onlinelibrary.wiley.com.gate1.inist.fr/doi/10.1002/0471250953.bi0313s48/abstract. Accessed August 17, 2015.

Sourjik, V., and Berg, H.C. (2000) Localization of components of the chemotaxis machinery of Escherichia coli using fluorescent protein fusions. *Mol Microbiol* **37**: 740–751.

Sourjik, V., and Berg, H.C. (2002) Binding of the Escherichia coli response regulator CheY to its target measured in vivo by fluorescence resonance energy transfer. *Proc Natl Acad Sci U S A* **99**: 12669–12674.

Sourjik, V., and Berg, H.C. (2004) Functional interactions between receptors in bacterial chemotaxis. *Nature* **428**: 437–441.

Strahl, H., Ronneau, S., González, B.S., Klutsch, D., Schaffner-Barbero, C., and Hamoen, L.W. (2015) Transmembrane protein sorting driven by membrane curvature. *Nat Commun* **6**: 8728.

Studdert, C.A., and Parkinson, J.S. (2005) Insights into the organization and dynamics of bacterial chemoreceptor clusters through in vivo crosslinking studies. *Proc Natl Acad Sci U S A* **102**: 15623–15628.

Thiem, S., and Sourjik, V. (2008) Stochastic assembly of chemoreceptor clusters in Escherichia coli. *Mol Microbiol* **68**: 1228–1236.

Thompson, S.R., Wadhams, G.H., and Armitage, J.P. (2006) The positioning of cytoplasmic protein clusters in bacteria. *Proc Natl Acad Sci U S A* **103**: 8209–8214.

Typas, A., and Sourjik, V. (2015) Bacterial protein networks: properties and functions. *Nat Rev Microbiol* **13**: 559–572.

1 **Figure legends**
2

Figure 1. FrzCD-GFP colocalizes with the nucleoid in *M. xanthus*. (A) Micrographs of *M.*

xanthus cells carrying a GFP or a mCherry fusion and stained with the DNA DAPI stain. The

genetic backgrounds of the *M. xanthus* strains are indicated on the left. The white arrows

indicate the cells whose fluorescence profiles and correlation coefficients between the DAPI

and GFP localization are shown in (B) and (C), respectively. Cells surrounded by white boxes

are taken from separate original micrographs. Scale bars correspond to 1μm. **(B)** GFP or

mCherry (green or red) and DAPI fluorescence (blue) profiles with the fluorescence intensity

(arbitrary units) represented on the *y* axis and the cell length positions with -1 and +1

indicating the poles, on the *x* axis. **(C)** Correlation coefficients between the DAPI and GFP or

mCherry localization. R^2 values > 0,5 indicate significant correlations. **(D)** Box plots indicate

the medians of the correlation coefficients (R^2) from 10 cells (from one biological replicate)

of each of the indicated strains. **(E)** Micrographs of a *M. xanthus parB* conditional mutant

carrying *frzCD-gfp* or *frzE-mCherry* and DAPI stained. Micrographs were obtained upon 18h

depletion of ParB. The genetic backgrounds of the *M. xanthus* strains are indicated on the left.

Scale bars correspond to 1μm.

Figure 2. FrzCD-GFP colocalizes with the nucleoid in *E. coli*. (A) Micrographs of *E.coli*

cells carrying a GFP fusion on a plasmid and stained with the DNA DAPI stain. The genetic

fusions are indicated on the left. Cells carrying *frzCD-gfp* were also treated with 10μg/ml

cephalexin to visualize FrzCD-GFP colocalization with the multiple nucleoids of undivided

cells. The white arrows indicate the cells whose fluorescence profiles are shown in (B). Cells

surrounded by white boxes are taken from separate original micrographs. Scale bars

correspond to 1μm. **(B)** GFP (green) and DAPI fluorescence (blue) profiles with the

fluorescence intensity (arbitrary units) represented on the *y* axis and the cell length positions

with -1 and +1 indicating the poles, on the *x* axis. **(C)** Correlation coefficients between the

DAPI and GFP localization. R^2 values > 0,5 indicate significant correlations. **(D)** Box plots

indicate the medians of the correlation coefficients (R^2) from 10 cells (from one biological

replicate) of each of the indicated strains.

Figure 3. FrzCD directly interacts with the DNA *in vitro*. (A) Electrophoretic mobility

shift assays (EMSA) on 1% agarose gels stained with ethidium bromide and developed at the

UV light. The indicated concentrations of purified 6His-FrzCD were incubated with a 801 bp

DNA fragment. **(B)** Schematic representation of the FrzCD protein domains. **(C-D)** The

indicated increasing concentrations of 6His-FrzCD$^{\Delta131\text{-}417}$ (C) and 6His- FrzCD$^{\Delta1\text{-}130}$ (D) were

used in EMSA assays with a 801 bp DNA fragment. **(E)** Average binding curves and

duplicates in degraded colors of each immobilized FrzCD construct 6His- FrzCD, 6His-

FrzCD$^{\Delta131\text{-}417}$ or 6His- FrzCD$^{\Delta7\text{-}27}$, with a 474 bp DNA fragment at a concentration of 38nM**.**

(F) "Sliding window" representation indicating the protein charge of the first FrzCD N-

terminal region at the different positions and obtained with 10, 20 and 30 residue windows

(blu, green and red, respectively). **(G)** Increasing concentrations of 6His-FrzCD$^{\Delta7\text{-}27}$ were

used in EMSA assays. On the first lane of each gel, 500 ng of the 2-Log DNA ladder (0.1-10

kb, NEB) have been loaded. Data in panel (A, C, D, and G) are representative of three

independent experiments.

Figure 4. The organization of FrzCD clusters depends on cluster intensity and mobility.

(A) A representative fluorescence 1 second time-lapse (left panel) and the corresponding

kymograph (middle panel) of a *frzCD-gfp* cell (top panel). Big and small arrows indicate large

and small clusters, respectively. The right panel represents the trajectories of each cluster

(same color codes as on the top panel). Scale bars correspond to 1μm. **(B)** Cluster

displacement (r) from the mean position at each given time (t). L represents the cell length.

The color code corresponding to the logarithm of the ratio r/L indicates that the amplitude of

the cluster displacement never exceeds 5% of the cell length. **(C)** Box plots indicate the

distribution of the Mean Square Displacements at the different lag times; the mean of each lag

value is indicate by the black dots. **(D)** Box plots indicate a significant decrease of the median

confinement for clusters of low fluorescence intensity compared to high intensity clusters. For

panels B, C and D 1039 clusters from 297 cells (two biological replicates) were analyzed. **(E)**

The box plots and the violin plots show the measured confinements of *frzCD-gfp* strains

blocked in the ON (*frzCD$_{E168A-G169A}$::gfp*) and OFF (*frzCD$_{E202A-E203A}$::gfp*) states. 130 and 150

clusters were analyzed for the ON and OFF states, respectively.

Figure 5. Frz cluster formation generates signal sensitivity in turn important for social

behaviors. (A) The average reversal frequencies, calculated by scoring FrzS-YFP pole-to-

pole oscillations are shown as a function of the IAA concentration for wild type (black),

frzCD$^{\Delta6-130}$ (red) and *ΔfrzCD* (grey). Reversal frequencies values of wild type and *frzCD$^{\Delta6-130}$*

can be fitted by the Hill equation with an interval of confidence of 95% (dashed lines). Error

bars represent the standard errors of the means. Reversal frequency values for each IAA dose

and each strain are the results of two independent biological triplicates. About one hundred

cells for the wild type and *frzCD$^{\Delta1-130}$* strains and fifty for the *ΔfrzCD* strain were analyzed

(refer to Supplementary Table 3 for the exact number of analyzed cells for each strain and

IAA doses used in this experiment). **(B)** Colony expansion of wild type, *frzCD$^{\Delta6-130}$* and

ΔfrzCD cells. Error bars represent the standard deviations of the means from three biological

replicates. **(C)** The same strains were analyzed in *E. coli* predation assays.

Figure supplements

Figure 1-figure supplement 1. Representative micrographs of *M. xanthus* cells carrying

a GFP or a mCherry fusion and stained with the DNA DAPI stain. The genetic

backgrounds of the *M. xanthus* strains are indicated on the left. Scale bars correspond to 1μm.

Figure 1-figure supplement 2. *frzE-mCherry* cells can swarm and form fruiting bodies

like wild type. Scale bars correspond to 0,5 cm.

Figure 1-figure supplement 3. FrzCD and FrzE are stable in different *M. xanthus*

mutants. Western blot with anti-FrzCD **(A)** or anti-FrzE antibodies **(B)** on the cell extracts of

the indicated *M. xanthus* strains. Scale bars correspond to 1μm. Black lines are used to

indicate that two lanes from the same gel where separated by other lanes there were lanes

between separate lanes from different westerns blots.

Figure 1-figure supplement 4. Positive correlation between the number of FrzCD

clusters and the nucleoid length (pixels). Average numbers of clusters with standard

deviations (black dots and bars, respectively) for different nucleoid sizes are shown. Green

dots represent measurements for individual cells. Grey zones represent the variances. 2564

clusters from two biological replicates were analyzed.

Figure 1-figure supplement 5. FrzCD segregated in cells with segregated chromosomes.

(A) 909 cells were ordered according to their cell length (pixels, grey) and for each cells GFP

and DAPI fluorescence are represented as green and blue dots, respectively, at their

corresponding cell position. 0 is the cell center. **(B)** The three histograms show the average

GFP (green) and DAPI (blue) fluorescence density for each cell position for the indicated

cell-size ranges. 0 is the cell center.

Figure 3-figure supplement 1. FrzCD binds DNA fragments of different lengths, forms a

dimer and promotes FrzE phosphorylation. (A- D) Electrophoretic mobility shift assay

(EMSA) on agarose gel stained with ethidium bromide and developed at the UV light. DNA

fragments of different lengths and GC contents (Table S4) were incubated with increasing

concentrations of FrzCD-(His$_6$) as indicated, including **(A)** 4 nM of Oligo1300 (1% agarose

gel), **(B)** 40 nM of Oligo340-1 (38% GC content; 1.5% agarose gel), **(C)** 40 nM of Oligo340-

2 (61% GC content; 1.5% agarose gel), and **(D)** 300 nM of Oligo70 (2.5% agarose gel). **(E)**

SEC-MALS analysis showing the elution profile (refractive index; black; left y-axis) and the

estimated molar masses (green; right y-axis) of the FrzCD eluted protein. **(F)** SDS page of the

indicated proteins purified from *E. coli* and used for the different experiments shown in

Figure 3. **(G)** Kinetics of the FrzE kinase domain (FrzE^{CheA}) auto-phosphorylation were

tested *in vitro* by incubation of FrzE^{CheA} in the presence of FrzA , the indicated different form

of FrzCD and ATPγP^{33} as a phosphate donor.

Figure 3-figure supplement 2. FrzCD-GFP binds the nucleoid in a DNA-sequence

independent manner. A library of DNA fragments was obtained by ChIP experiments on

frzCD-gfp and *parB-yfp* strains, using GFP polyclonal antibodies. The figure shows the results

obtained by the deep sequencing of the DNA libraries. Only for *parB-yfp*, we observed an

enrichment corresponding a to the nucleoid region containing *parS* (rectangle in the middle

panel and last panel) (Harms *et al.*, 2013) (9,109 to 9,110 Kb). Note that while the number of

reads relative to ParB are represented with a logarithmic scale, for FrzCD we used a regular

scale.

Figure 3-figure supplement 3. The FrzCD N-terminal tail has the same properties of that

of eukaryotic histones. (A) Prediction of the FrzCD N-terminal secondary structures. The

nature of each amino acid is also indicated through color codes. **(B)** FrzCD first 50 amino

acid alignment with the N-terminal tail of Histones 2B. The alignment was obtained by

Clustal Omega. Dots indicate similarities and stars identities.

Figure 3-figure supplement 4. The FrzCD N-terminus from different species show

similar amounts of positively charged amino acids. The alignment of FrzCD sequences

from the indicated species was obtained by Clustal Omega. Stars indicate identities.

1 Table S1: Strains used in this study
2

Strains	Genotyp	Deletion	Source
DZ2	*wt*		Zusman et al., 1982
DZ4620	*frzCD-gfp*		Mauriello et al., 2009
DZ4480	Δ*frzCD*	Codons 6-393	Bustamante et al., 2004
DZ4485	*frzCD*$^{Δ6-130}$		Mauriello et al., 2009
DZ4743	*frzCD*$^{Δ6-130}$*-gfp*		Mauriello et al., 2009
EM231	*frzCD*$_{E202A-E203A}$*::gfp*		Mauriello et al., 2009
EM228	*frzCD*$_{E168A-G169A}$*::gfp*		Mauriello et al., 2009
EM434	*frzE-mCherry*		This study
EM506	*frzE-mCherry* Δ*frzCD*		This study
EM516	*frzCD-gfp frzE::kan*	Codons 171-438	This study
EM531	*difA-gfp* Δ*parB/P$_{cuoA}$-parB*		This study
EM532	*frzCD-gfp* Δ*parB/P$_{cuoA}$-parB*		This study
EM533	*frzE-mCherry* Δ*parB/P$_{cuoA}$-parB*		This study
EM543	*frzCD*$^{Δ7-27}$	Codons 7-27	This study
EM550	*frzCD*$^{Δ7-27}$*-gfp*	Codons 7-27	This study
TM26	*frzS-yfp*		Guzzo et al., 2015
EM622	*frzCD*$^{Δ6-130}$ *frzS-yfp*		This study
EM623	Δ*frzCD frzS-yfp*		This study

3
4
5 Table S2: Plasmids used in this study
6

Plasmid	Expression plasmid	Source
pAH57	*P$_{cuoA}$-parB,* copper-dependent expression of *parB*, Mx8 attB	Harms et al., 2013
pAH18	Construct for in frame deletion of *parB*	Harms et al., 2013
pETPhos_*frzCD*	pETPhos with *frzCD* tagged with 6-his inducible with IPTG	Guzzo et al., 2015
pETPhos_*frzE*^{kinase}	pETPhos with *frzE*^{CheA} tagged with 6-his inducible with IPTG	Guzzo et al., 2015
pETPhos_*frzCD*^{c}	pETPhos with *frzCD*^{c} tagged with 6-his inducible with IPTG	Guzzo et al., 2015
pGEX(M)_*frzA*	pETPhos with *frzA* tagged with 6-his inducible with IPTG	Guzzo et al., 2015
pEM365	pBJ113 with frzE-mCherry fusion	This study
pEM405	pBJ113 with an insertion cassette for *frzE::kan*	This study
pEM418	pBJ113 with a cassette for *frzCD*$^{Δ7-27}$	This study
pEM414	pETPhos with *frzCD*$^{Δ1-130}$ tagged with 6-his inducible with IPTG	This study
pEM415	pETPhos with *frzCD*$^{Δ131-417}$ tagged with 6-his inducible with IPTG	This study
pEM433	pETPhos with *frzCD*$^{Δ7-27}$ tagged with 6-his inducible with IPTG	This study
pEM409	pETDuet-1 with *frzCD-gfp* inducible with IPTG	This study
pEM417	pETDuet-1 with *frzCD*$^{Δ1-130}$*-gfp* inducible with IPTG	This study
pEM434	pETDuet-1 with *frzCD*$^{Δ7-27}$*-gfp* inducible with IPTG	This study
pEFrzSY	pEYFPN1 with a cassette to construct *frzS-yfp*	Guzzo et al., 2015
pFCD-H6	pHis17 with *frzCD* gene tagged with a C-terminal 6-his	This study

7
8

9 Table S3: Number of cells analyzed for Figure 5A

%IAA / Strain	0	0.005	0.01	0.03	0.05	0.1	0.3
frzS-yfp	101	113	77	114	64	47	170
frzCD$^{Δ6-130}$ *frzS-yfp*	98	103	75	128	112	139	109
Δ*frzCD frzS-yfp*	44	ND	ND	ND	48	ND	47

10
11
12
13

Table S4: DNA sequences used for EMSA in Figure 3 and Supplementary figure 4

Ds DNA	Sequence (5' → 3')	Length, GC content (%)
Oligo70	CTTGCAGTAGAGCTGACCATGATTACGCCATCAGCAGCTCCAGGTCGTAC CTCCAGCTACCAATCCCCG	69, 57
Oligo340-1	TAATACGACTCACTATAGGGAGACCACAACGGTTTCCCTCTAGAAATAAT TTTGTTTAACTTTAAGAAGGAGATATACATATGTCCCTGGACACCCCCAA CGAGAAGCCCGCTGGCAAGGCTCGCGCCCGGAAGGCCCCCGCCTCCAAGG CCGGCGCCACGAACGCGGCGTCGACCTCTTCCTCCACCAAGGCCATCACC GACACGCTGCTGACGGTGCTGTCCGGCAACCTGCAGGCCCGCGTGCCCAA GGAGCTGGTCGGTGAGTCCGGCGTGGAGCTGGCGCACCTGCTCAACCAGG TGCTGGACCAGTTCGCGGCCTCCGAGCACCGCAAGCATG	339, 61
Oligo340-2	GTTTAACTTTAAGAAGGAGATATACATATGAAAAAAGAAACGATTTTTTC CGAAGTAGAAACGGCTAACAGCAAGCAACTGGCTGTGTTGAAAGCTAATT TCCCACAGTGTTTTGATAAAAACGGAGCCTTCATTCAAGAAAAATTGCTT GAGATTATTAGGGCATCGGAAGTTGAACTCTCTAAAGAATCATACAGCTT GAACTGGCTGGGTAAATCTTATGCCCGTTTGTTGGCCAATCTACCACCGA AAACGTTGTTGGCAGAAGATAAAACTCATAACCAACAAGAAGAGAACAAG AACAGTCAACACCTGTTAATCAAAGGGGATAATCTCGAAGTATTG	345, 38
Oligo1300	*frzCD* gene amplified with the primers 5'-ATGTCCCTGGACACCCCCAACGAGAAGCCCGCTGG-3'and 5'-CTAGTCGGCCTTGAACCGCTTGATGAGCTCGGCCA-3' and the plasmid pFCD-H6 as template	1254, 70
Oligo801	Fragment amplified with the primers 5'-cgggatcctggctccgcccccgacgca-3' and 5'-cccaagcttttgatgaggcgcttggagat-3' and DZ2 chromosomal DNA as template	801, 68

	S-motility	**Fruiting body formation**

wt

frzE-mCherry

Δ*frzE*

A

M (Kda) wt ΔfrzCD frzCDgfp frzCDΔ6-130 frzCDΔ6-130-gfp frzCD Δ7-27 frzCD Δ7-27-gfp frzCDgfp frzCDgfp frzE::kan

B

M (Kda) wt ΔfrzE frzE-mCherry frzE-mCh ΔfrzCD

A

B

Cell length