Distress tolerance across self-report, behavioral and psychophysiological domains in women

with eating disorders and healthy controls

Authors: Angelina Yiu[1], M.A., Kara Christensen[2], M.A., Jean M. Arlt[1] MPhil., & Eunice Y.

Chen[1], Ph.D.

[1] Temple Eating Disorders program, Temple University, 1701 North 13th Street, Philadelphia,
PA 19122
[2] Cognition and Emotion Lab, Ohio State University, 1835 Neil Avenue, Columbus, OH 43210

Corresponding author: Eunice Y. Chen, Ph.D., Temple Eating Disorders program, Temple
University, 1701 North 13th Street, Philadelphia, PA 19122

Abstract word count: 250
Manuscript world count: 4,630

1   Abstract

Background and objectives: The tendency to engage in impulsive behaviors when distressed is

linked to engagement in disordered eating. The current study comprehensively examines

emotional responses to a distress tolerance task by utilizing self-report, psychophysiological

measures (respiratory sinus arrhythmia [RSA], skin conductance responses [SCRs] and tonic

skin conductance levels [SCLs]), and behavioral measures (i.e., termination of task, latency to

quit task). *Methods*: 26 healthy controls (HCs) and a sample of treatment-seeking women with

Bulimia Nervosa (BN), Binge Eating Disorder (BED) and Anorexia Nervosa (AN) (N=106)

completed the Paced Auditory Serial Addition Task- Computerized (PASAT-C).

Psychophysiological measurements were collected during baseline, PASAT-C, and recovery,

then averaged for each time-period. Self-reported emotions were collected at baseline, post-

PASAT-C and post-recovery. *Results*: Overall, we found an effect of Time, with all participants

reporting greater negative emotions, less happiness, lower RSA, more SCRs and higher tonic

SCLs after completion of the PASAT-C relative to baseline. There were no differences in

PASAT-C performance between groups. There was an effect of Group for negative emotions,

with women with BN, BED and AN reporting overall higher levels of negative emotions relative

to HCs. Furthermore, we found an effect of Group for greater urges to binge eat and lower RSA

values among BED, relative to individuals with BN, AN and HCs. *Limitations:* This study is

cross-sectional and lacked an overweight healthy control group. *Conclusion*: During the PASAT-

C, individuals with EDs compared to HCs report higher levels of negative emotions, despite

similar physiological and behavioral manifestations of distress.

Keywords: eating disorders; distress tolerance; psychophysiology; PASAT-C

Distress tolerance across self-report, behavioral and psychophysiological domains in women

2   with eating disorders and healthy controls

3   Eating disorders (EDs) affect up to 4.64% of adults (Le Grange, Swanson, Crow, &

Merikangas, 2012) and have significant medical and psychosocial outcomes (Baiano et al.,

2014). Disordered eating, in general, has been conceptualized as a maladaptive response to

alleviate distress, and thereby may negatively reinforce the disordered eating (Aldao, Nolen-

Hoeksema, & Schweizer, 2010; Anestis, Smith, Fink, & Joiner, 2009; Corstorphine, Mountford,

Tomlinson, Waller & Meyer, 2007; Fischer, Smith, & Cyders, 2008; Haynos & Fruzzetti, 2011;

Leyro, Zvolensky, & Bernstein, 2010).

10   A number of studies use multiple methodologies to examine individuals' with EDs

experience of and abilities to withstand negative emotions and/or aversive states, termed distress

tolerance. For example, it was found that self-report emotional distress tolerance, but not

behavioral and physical forms of distress tolerance, was negatively associated with symptoms of

bulimia nervosa (BN) (Anestis et al., 2012). Another study found that BN did not differ from

healthy controls (HCs) on heat and cold pain thresholds after a stress induction task (Schmahl et

al., 2010). While these studies expand our understanding of the utility of self-report and

behavioral measures of distress tolerance in EDs, our understanding of the course of affective

responding in EDs before, during, and after a stressor is still limited (Anestis et al., 2007;

Anestis, Smith, et al., 2009; Claes, Vandereycken, & Vertommen, 2005; Peterson & Fischer,

2012; Wenzel, Weinstock, Vander Wal, & Weaver, 2014; Wu et al., 2013).

21   To gain a comprehensive understanding of affective response when engaged in distress

tolerance in EDs, multiple methods of assessment are needed (Gross, 2013); however, there are

few studies that utilize this approach, therefore any relationship between the psychophysiological

components of emotional responding and the behavioral components in EDs remains unclear

(Gross, 2013; Lang, Greenwald, Bradley, & Hamm, 1993).

The psychophysiological component of emotional response is comprised of

parasympathetic and sympathetic nervous system activity. Parasympathetic nervous system

activity can be indexed by respiratory sinus arrhythmia (RSA), which consists of heart rate

variability (HRV) in conjunction with respiration. Neuroimaging studies show that there is a

positive relationship between RSA and increased activation in neural structures implicated in

emotion processing and physiological aspects of emotional responses (e.g., right pregenual

anterior cingulate, right subgenual anterior cingulate and right rostral medial prefrontal cortex

and the left sublenticular extended amygdala or ventral striatum) (Thayer, Åhs, Fredrikson,

Sollers, & Wager, 2012). Reduced resting RSA is theorized as a risk factor for psychopathology

(Crowell, Beauchaine, & Linehan, 2009), although systematic reviews and meta-analysis suggest

that higher resting HRV is associated with both BN (Peschel et al., 2016) and anorexia nervosa

(AN) (Mazurak, Enck, Muth, Teufel, & Zipfel, 2011) compared to controls. Low resting RSA,

decreased RSA when emotionally aroused, and a slow return to baseline after emotional arousal

are associated with greater symptoms of psychopathology (Beauchaine, 2015).

Experimental work examining RSA response to stressors among individuals with EDs

has produced mixed findings. Two separate studies found that women with BED and obesity

(Friederich et al., 2006) and women with BN (Messerli-Bürgy, Engesser, Lemmenmeier,

Steptoe, & Laederach-Hofmann, 2010) showed decreased RSA levels with a slow return to

baseline following a psychological stress induction task. However, there is also evidence to

suggest that RSA levels among women with BED and obesity did not change after psychological

stress was induced (Messerli-Bürgy et al., 2010) and conversely women without EDs who are

obese showed a return to baseline RSA levels after stress exposure (Messerli-Bürgy et al., 2010).

Additional research suggests that decreased RSA after psychological stress is associated with

engagement in clinically significant levels of binge eating, rather than weight status (Udo et al.,

2014).

The sympathetic nervous system response can be indexed by skin conductance response,

which measures the time it takes for a current to pass through the skin (Boucsein et al., 2012).

Higher skin conductance responses (SCRs) are associated with greater emotional arousal

(Kreibig, 2010; Lang et al., 1993). In contrast to the mixed relationship between RSA and EDs,

evidence suggests that skin conductance levels (SCL) are similar between women with BN and

BED in response to different stressors (Hilbert, Vögele, Tuschen-Caffier, & Hartmann, 2011)

and between women with BN, women with self-reported restrained eating or HCs (Tuschen-

Caffier & Vögele, 1999). Taken together, there is some evidence to suggest that there is a

relationship between EDs and decreased RSA when psychological stress is induced, and little

evidence pointing to a relationship between EDs and sympathetic responses as indexed through

skin conductance.

Thus far, the evidence base for the psychophysiological component of emotional

response while distressed across the range of EDs has been mixed, limited by small sample sizes,

and there is a lack of research utilizing multiple ED diagnoses. The current study extends past

research (Leehr et al., 2015; Naumann, Tuschen-Caffier, Voderholzer, Caffier, & Svaldi, 2015;

Svaldi, Griepenstroh, Tuschen-Caffier, & Ehring, 2012; Svaldi, Tuschen-Caffier, Lackner,

Zimmermann, & Naumann, 2012) through the multi-modal assessment of emotional responses

across three common ED diagnoses and HC participants to a commonly used behavioral distress

tolerance task. The Paced Auditory Serial Addition Task-Computerized (PASAT-C) (Lejuez,

Kahler, & Brown, 2003) was used as a behavioral measure of distress tolerance (Feldner, Leen-

Feldner, Zvolensky, & Lejuez, 2006; Gratz, Rosenthal, Tull, Lejuez, & Gunderson, 2006) and

self-reported emotions and psychophysiological measures of arousal were measured in response

to the PASAT-C. As individuals with EDs have been argued to have, "shared, but distinctive,

clinical features […] maintained by similar psychopathological processes," utilizing a

transdiagnostic approach that includes three ED groups may be most useful in identifying the

proposed "similar psychopathological processes" (Fairburn, Cooper, & Shafran, 2003)

We predicted a main effect of time, such that self-reported negative emotions would be

greater, happiness would be lower, and urges to binge eat would be greater after the PASAT-C,

in comparison to baseline and at recovery. Furthermore, we expected that diagnostic group

would moderate this effect. Specifically, we expected that HCs would demonstrate less

significant changes in negative emotions, happiness, and urges to binge eat during the PASAT-C,

compared to individuals with BN, BED or AN. As the PASAT-C is used as a demanding

behavioral distress tolerance measure (Gratz et al., 2006; Sauer & Baer, 2012), we expected that

individuals with BN, BED and AN would be more likely to prematurely terminate the PASAT-C

and exhibit shorter latency to terminate the PASAT-C than HCs. Given the proposed role of

negative affect in EDs during distress (Anestis, Peterson, et al., 2009; Anestis et al., 2007;

Peterson & Fischer, 2012), we expected that RSA values would be lower and SCRs and tonic

SCL values would be higher during the PASAT-C in ED groups compared to HC, relative to

baseline and recovery.

21   **Methods**

Participants

Participants were self-referred from the community using flyers, referred from local

eating disorder clinics and student health and counseling services, and recruited in a University-

based outpatient eating disorder program as part of several clinical trials. Participants were

invited to participate in the study if, after undergoing a clinical interview (described below) they

met for a diagnosis of BN, BED or AN (typical or atypical), without current drug or alcohol

dependence or symptoms of psychosis. HCs were self-referred from the community, student

health and counseling services, and the University Hospital, and were eligible to participate if,

after undergoing the same clinical interview as EDs, they did not meet diagnosis for current

psychopathology, and did not have a history of serious mental illness.

The sample consisted of 106 females with an ED and 26 female HCs with no history of

psychiatric disorders from the community. Of the participants with an ED, 34.90% of the sample

was diagnosed with BN (n=37), 51.89% with BED (n=55), and 13.21% with typical or atypical

AN (n=14). The sample identified predominantly as non-Hispanic (91.70%) and were 64.4%

Caucasian, 13.60% African-American, 4.50% Asian, and 9.1% of mixed race. The mean age of

participants was 35.06 years ($SD = 11.47$), and the mean body mass index (BMI) was 29.55

kg/m^2 ($SD = 9.50$).

Measures

Psychiatric symptoms. The Structured Clinical Interview for Diagnostic and Statistical

Manual of Mental Disorders – IV- Text Revision (DSM – IV – TR) Axis I Disorders (SCID – I)

(First, Spitzer, Gibbon, & Williams, 2002) was used to assess for psychiatric symptoms. The

SCID I was used to confirm ED diagnoses with the EDE-16.0 and to assess for exclusionary

psychiatric symptoms. The SCID I is considered the gold-standard measure for psychiatric

diagnoses (Lobbestael, Leurgans & Arntz, 2011).

Eating disorder behaviors. The Eating Disorder Examination 16.0 (EDE) (Fairburn,

Cooper, & O'Connor, 2008) is an investigator-based interview that assesses DSM– IV-TR

(American Psychiatric Association, 2000) ED symptoms. The EDE – 16.0 was used to diagnosis

AN, BN and BED and generate subscales that tap into ED behaviors and cognitions: Eating

Concern, Shape Concern, Weight Concern, Dietary Restraint, as well as a Global score. The

EDE – 16.0 was also used to assess for the number of binge eating and compensatory episodes in

the prior 3-month period. The EDE – 16.0 has good discriminant validity, such that individuals

with AN, BN and BED are distinguished from controls (Cooper, Cooper & Fairburn, 1989;

Wilson & Smith, 1989; Wilfley, Schwartz, Spurrell & Fairburn, 1999). In the current study, the

EDE total score demonstrated an internal consistency of α = .94.

Current subjective emotional state. A visual analogue scale (VAS) with an abbreviated

Positive and Negative Affect State (PANAS) (Watson, Clark, & Tellegen, 1988) measured

current subjective emotional state prior to the baseline, after the PASAT-C, and after recovery

from the PASAT-C (see Figure 1). Self-report negative and positive affective adjectives that

assessed anxiety, fear, frustration, happiness, sadness, and tension were scored on a 100-point

Likert scale. A single question to assess urges to binge eat was added due to its relevance in the

current ED sample. Higher scores indicated greater intensity of the response. In the present

study, we created a composite for negative emotions using an average of scores from anxiety,

fear, frustration, sadness and tension. Urge to binge eat and happiness were assessed separately.

The original PANAS shows good convergent validity, with correlations ranging from .51 to .74

with the Beck Depression Inventory, the State-Trait Anxiety Inventory State Anxiety Scale and

Hopkins Symptom Checklist (Watson, Clark & Tellegen, 1988). In the present study, internal

consistency of the negative emotion composite score at each time point ranged from α = .84 to .90.

Psychophysiological measures. Psychophysiological measures included respiratory

sinus arrhythmia (RSA) as a measure of parasympathetic activity (Grossman & Taylor, 2007;

Thayer et al., 2012) and skin conductance responses (SCRs) and tonic skin conductance levels

(tonic SCL) as measures of sympathetic activity (Boucsein, 2011). Psychophysiological

measures were assessed during the baseline, during the PASAT-C and during the recovery

period. Electrocardiogram (ECG) data were collected from two Biopac Ag-AgCl spot electrodes

placed on the bottom of the palm of the non-dominant hand, using a modified Lead II

configuration (Boucsein, 2011). We derived RSA by using a band pass filter on the ECG signal

and spectral analysis to extract the high-frequency component (> .15 hz) of heart rate variability.

Skin conductance was recorded from two electrodes attached to the palm of the non-dominant

hand. Increases in SCRs have been associated with negative affect following exposure to

emotional stimuli, even when cardiac measures did not change significantly (Boucsein et al.,

2012; Salters-Pedneault, Gentes, & Roemer, 2007). Tonic SCL referred to the level of skin

conductance per data collection period.

Diagnostic and Laboratory Procedures.

The present study was reviewed and approved by the university institutional review board

and took place over two separate testing sessions, with each session taking 2-4 hours. During the

first session, eligibility screening and informed consent were completed, followed by an

assessment of psychiatric and ED symptoms by Masters-level clinicians using the measures

described below. Individual participant diagnoses were agreed upon and finalized with the

licensed clinical psychologist supervisor at weekly best-estimate meetings (Klein, Ouimette,

Kelly, Ferro, & Riso, 1994; Kosten & Rounsaville, 1992), where all structured interview data

was presented. Height and weight were measured to calculate BMI. During the second session,

participants completed the laboratory procedures for the PASAT-C, described below. Diagnostic

and psychophysiological assessments occurred prior to outpatient treatment for individuals with

EDs.

Behavioral distress tolerance task. The Paced Auditory Serial Addition Task-

Computerized (PASAT-C) (Lejuez et al., 2003) is a behavioral distress tolerance task that has

been shown to induce negative affect (Daughters, Lejuez, Kahler, Strong, & Brown, 2005;

Feldner et al., 2006; Holdwick & Wingenfeld, 1999; Schloss & Haaga, 2011; Eichen, Chen,

Boutelle & McCloskey, 2017) and short-term anxiety, frustration, and irritability (Gratz et al.,

2006; Lejuez et al., 2003). The PASAT-C is a computer-based task that requires the participant

to add a visually presented digit to the previous visually presented digit. Explosion sounds

followed incorrect answers or when the participant failed to respond quickly enough. The

PASAT-C was presented for a maximum of twelve minutes, consisting of four levels that lasted

for three minutes each. With each successive level of the PASAT-C, the latency between trials

was decreased and negative feedback of "*go faster*," "*do better*," and "*go faster, do better*" was

added. By the fourth level (PASAT-C Level 4), there was a one second latency of trials with

negative feedback. Participants were provided written instructions at the beginning of the task

that they had the option to quit the task after completion of Level 2. If participants terminated the

task early, they went on to complete the recovery period, detailed below. Distress tolerance on

the PASAT-C was behaviorally operationalized dichotomously and continuously as termination

of the task when given the option and latency to quit, respectively.

Psychophysiology Capture Procedure. Psychophysiological measures were collected

during baseline, PASAT-C, and the recovery period. Average RSA, SCR and tonic SCL values

were computed from the 5-minute baseline, up to 12 minutes of the PASAT-C, and the 5-minute

recovery period, respectively. Please see Figure 1 for a schematic of the procedure. Once sensors

were attached, participants completed a baseline VAS and were asked to sit quietly without

moving for a 5-minute baseline period (Laborde, Mosley & Thayer, 2017). Following this,

participants were instructed to begin the PASAT-C. Upon completion of the PASAT-C,

participants completed a second VAS and were asked to sit quietly for an additional 5-minutes

without moving for a recovery period (Laborde et al., 2017). Participants completed a final VAS

rating after the recovery period.

Statistical Treatment

Chi-square tests were conducted to assess for group differences in all demographic

variables and medical and psychiatric comorbidities, with the exception of age, which was

assessed using an ANOVA. Variables related to presence of medication use (e.g., stimulant

medication, anxiolytics) and medical co-morbidities were included as covariates in the analyses

if there were significant differences between ED groups (BN, BED, AN) due to their effects on

psychophysiological responses (Grossman, Stemmler, & Meinhardt, 1990; Masi, Hawkley,

Rickett, & Cacioppo, 2007). As it was expected that HCs would exhibit significantly less

medical co-morbidities and medication use relative to the ED groups (Hudson, Hiripi, Pope, &

Kessler, 2007; Johnson, Spitzer, & Williams, 2001; Kessler et al., 2013; Padierna, Quintana,

Arostegui, Gonzalez, & Horcajo, 2000), medical co-morbidities and medication use were only

included as covariates if there were significant differences between BN, BED and AN.

For analyses involving self-reported emotions and psychophysiological measures,

independent variables were Time (baseline, PASAT-C, recovery) and Group (BN, BED, AN and

HC). A repeated measures ANOVA was chosen as it allows for individual differences in baseline

scores to be accounted (Field, 2013). Six repeated measures Time x Group ANOVAs were

conducted with the self-reported negative emotion composite score, happiness, urge to binge eat,

RSA, SCRs and Tonic SCL as dependent variables. Planned simple effects analyses were

conducted to probe significant interactions. To examine group differences in PASAT-C

completion vs. non-completion, a chi square test was conducted. To examine group differences

in PASAT-C latency to quit, a univariate ANOVA was conducted. A power analysis indicated

that we were powered to detect medium to large interaction effects ($f = .25$ to .40) with a power

of .95 (Faul, Erdfelder, Lang, & Buchner, 2007). Partial η^2 was used to report effect sizes for

repeated measures ANOVAs and ANOVAs, with the following cut-off conventions: small (.01),

medium (.06) and large (.14) (Cohen, 1988).

10   **Results**

Preliminary Analyses

As expected, there were significant group differences on presence of a lifetime mood

disorder ($p < .001$), anxiety disorder ($p < .001$), medical co-morbidities ($p = .006$) and

medication use ($p = .002$), which were driven by the lack of psychiatric and medical co-

morbidities among HCs. There were no significant differences between BN, BED and AN on

presence of a lifetime mood disorder, medical co-morbidities or medication use, although AN

were significantly more likely to have an anxiety disorder relative to BN ($p < .05$). Due to the

lack of significant systematic differences in psychiatric and medical co-morbidities between BN,

BED and AN, these variables were not included as covariates. Please see Table 1 for a

description of the severity of ED symptoms, demographic information and clinical characteristics

of the sample.

Main Analyses[1]

2   **Hypothesis 1: Self-reported emotions**. Overall, there was a main effect of Time, such

that participants reported higher negative emotions, $F_{(2, 107)}= 47.61$, $p < .001$, $\eta^2=.31$, and lower

happiness, $F_{(2, 107)}= 39.10$, $p < .001$, $\eta^2=.27$, after completion of the PASAT-C, in comparison to

baseline and recovery ($ps < .05$). There was a main effect of Group for negative emotions, $F_{(3,}$

$_{107)} = 12.09$, $p < .001$, $\eta^2=.25$, but not happiness ($p = .07$). Specifically, HC participants reported

lower negative emotions ($ps < .001$) in comparison to individuals with BN, BED or AN across

all measurement periods. There were no Time x Group interactions ($ps = .12 - .27$) for negative

emotions or happiness.

10   For urges to engage in binge eating, there was a main effect of Group, $F_{(3, 107)}= 16.51$, $p <$

$.001$, $\eta^2 = .32$, such that across all measurement periods individuals with BED demonstrated

significantly greater urges to binge eat compared to BN, AN and HCs ($ps < .04$), individuals with

BN demonstrated significantly greater urges to binge eat compared to HCs ($p < .001$), and

individuals with AN did not differ significantly from BN or HC ($ps > .06$). There was no effect

of Time ($p = .56$) or an interaction between Time x Group ($p = .93$), but individuals with AN

exhibited a trend for decreased urge to binge eat after completion of the PASAT-C. See Table 3

for a summary of findings.

18   **Hypothesis 2: PASAT-C termination and latency to quit.** A chi square test indicated

19   that there were no differences between groups on PASAT-C completion versus non-completion,

20   $\chi^2_{(3)}= 3.85$, $p = .28$. A univariate ANOVA suggested that there were no differences between

---

[1] Analyses were re-run with BMI as a covariate, with some differences in findings. Specifically, there was no longer an effect of Time for negative emotions ($p = .70$) or RSA ($p = .83$). All other findings remained the same.

1   groups on PASAT-C latency to quit, $F_{(3,129)} = 1.39$, $p = .25$, $\eta^2 = .03$. See Table 4 for means and

2   standard deviations.

3   **Hypothesis 3: Psychophysiological measures**. There was a significant effect of Time

for RSA values, $F_{(2,128)} = 11.87$, $p < .001$, $\eta^2 = .09$, such that all participants exhibited lowered

RSA during the PASAT-C, in comparison to baseline ($p = .05$) and recovery ($p < .001$). There

was a significant effect of Group, $F_{(3,128)} = 6.10$, $p = .001$, $\eta^2 = .13$, such that individuals with

BED exhibited lowered RSA levels in comparison to individuals with BN or HCs ($ps \leq .006$) but

not AN ($p = .325$) across all measurement periods. There was no Time x Group interaction for

RSA values.

10   There was a significant effect of Time for Tonic SCL values, $F_{(2,109)} = 50.57$, $p < .001$,

11    $\eta^2 = .32$, and SCR values, $F_{(2,107)} = 32.85$, $p < .001$, $\eta^2 = .24$, such that all participants exhibited

12   greater Tonic SCL and SCR values during the PASAT-C in comparison to baseline ($ps < .001$),

13   but not recovery ($ps = .45 - .75$). There was no effect of Group or Time x Group interaction for

14   Tonic SCL or SCR values. See Table 5 for means and standard deviations.

15   **Discussion**

16   The current study extends the current literature on affective response in EDs by

integrating self-report, behavioral and psychophysiological measures (RSA, SCRs and tonic

SCLs) of response to a distress tolerance task (Daughters et al., 2005; Feldner et al., 2006;

Holdwick & Wingenfeld, 1999; Schloss & Haaga, 2011). Despite similar psychophysiological

responding (RSA, SCRs and tonic SCL) to the PASAT-C and PASAT-C termination and latency

to quit across all groups, individuals with BN, BED and AN reported greater overall negative

emotions and lower overall happiness compared to HCs. The discrepancy between

psychophysiological responding and PASAT-C termination and latency to quit with self-report

of emotional experience suggests that individuals with EDs experience disproportionately greater

subjective distress relative to HCs when presented with the same stimuli. Overall, this may

suggest that while physiological and behavioral experience of distress manifests similarly across

groups, it is the interpretation of experiences during both baseline and when distressed that best

distinguish between groups with EDs and HCs.

Although HCs reported significantly lower negative emotions and greater happiness

relative to all other groups, individuals with BED were differentiated from other ED groups and

HCs by exhibiting significantly lower RSA values without concomitant differences from other

groups in Tonic SCL or SCR values. Research has demonstrated an association between BMI

and RSA and suggests that cardiovascular autonomic function changes as a function of weight

loss (Karason, Mølgaard, Wikstrand, & Sjöström, 1999; Rissanen, Franssila ‐ Kallunki, &

Rissanen, 2001). Consistent with past research (Dingemans & van Furth, 2012; Hudson et al.,

2007), HCs had significantly lower BMI relative to the BED group and significantly higher BMI

relative to the AN group, but did not differ significantly from the BN group; however, BMI was

not controlled for in the current study, as 92% of participants diagnosed with BED were

overweight or obese, likely due to the cumulative effects of objective binge episodes on weight

gain over time (Hudson et al., 2010). Therefore, participants with BED are "doubly diagnosed"

with BED and obesity, making it difficult to dissociate diagnosis and weight status. Future

studies may wish to dissociate diagnosis and weight status.

Contrary to expectations, we did not find a Time (baseline, PASAT-C and recovery) by

Group interaction in self-reported urge to binge eat. However, there were Group differences in

urge to binge eat, such that individuals with BED and BN reported overall greater urges to binge

eat across all time-points in comparison to individuals with AN or HC. Perhaps individuals who

engage in habitual binge eating experience consistent urges to engage in binge eating, but only

act on this urge when they experience negative affect (Hilbert & Tuschen‑Caffier, 2007) or are

under stress. As the current study only assessed urges to binge eat, future research may seek to

explore factors that influence whether one will or will not act upon urges to binge eat when

experiencing negative affect (Haedt-Matt & Keel, 2011; Hilbert & Tuschen‑Caffier, 2007).

The PASAT-C achieved the desired result of increasing distress as evidenced by

increased negative emotions, decreased happiness, decreased RSA values and increased SCRs

and tonic SCLs across the sample, without differences in termination or latency to quit. This

suggests that regardless of baseline level of negative emotions or diagnostic group, the PASAT-

C was distress inducing, which is consistent with past research utilizing samples with clinically

significant psychopathology (Gratz et al., 2006). The success of the PASAT-C to induce distress

may have produced a ceiling effect on performance, such that all participants were equally

sensitive to its effects regardless of diagnosis. However, the lack of significant group differences

on the PASAT-C is counter to another study that found significantly shorter latency to quit the

PASAT-C among undergraduate students who endorsed binge eating behaviors relative to

controls (Eichen et al., 2017). The difference in findings may be due to sampling differences

between the two studies. The current study recruited treatment-seeking patients with a range of

EDs; the other study recruited non-treatment seeking college students.

Future Directions

The inclusion of multiple ED diagnostic groups can be considered a strength and a

limitation of the current study. Comparisons were possible between ED diagnoses and HCs,

however the smaller group of individuals with AN may have hindered detecting differences

between that group and others. For example, individuals with AN exhibited a trend for decreased

urges to binge eat after completion of the PASAT-C in comparison to baseline, which is in

contrast to individuals with BED and BN who exhibited similar urges to binge eat at baseline and

after the PASAT-C. This may reflect an increased desire for control over eating when

experiencing negative affect among individuals with AN (Fairburn, Shafran, & Cooper, 1999). A

larger sample of individuals with AN may clarify the urge to binge eat in response to stress

among different ED diagnoses. Furthermore, as the PASAT-C generated similar levels of

negative affect across the entire sample, future research could utilize emotion inductions that

may be more salient to EDs, such as cues involving body shape/weight concerns or food. Such

disorder-related stimuli may produce differential results between groups with EDs and HCs and

offer increased insight into the negative affect states that are posited to maintenance disordered

eating behaviors. For example, participants could complete behavioral approach/avoidance tasks

that involve examining the body in a mirror or observing images of high-caloric food. In terms of

behavioral assessments, future research could utilize distress tolerance measures that more

closely map onto eating disorder concerns, such as sampling foods.

Finally, the current study's group of individuals with BED primarily consisted of

individuals who are also overweight. Although individuals with BED are at a higher risk for

obesity (e.g., de Zwaan, 2001), the medical condition of obesity is not synonymous with the

psychological condition of BED (e.g., Klatzkin, Gaffney, Cyrus, Bigus, & Brownley, 2015).

There are multiple physiological, cognitive, and psychological symptoms associated with obesity

and there is a risk of falsely conflating these symptoms with BED. Future studies may seek to

disentangle BED and obesity by examining whether similar patterns of findings are found for

individuals with BED who are of normal weight, or for individuals without EDs who are of

normal weight and overweight status.

The current study is one of the first to examine emotional responding across individuals

with EDs and HCs in response to a task that induces distress. It provides a step towards

enhancing our understanding of the similarities and differences in emotional responding across

ED diagnoses using multiple measures of emotional responding. As previously suggested, our

finding that individuals with BED exhibit overall reduced levels of RSA responding and

experience consistent urges to binge eat over time requires further study before treatment

implications can be made. The finding that EDs are associated with overall greater subjective

distress without deleterious effects on the PASAT-C termination or latency to quit relative to

HCs has treatment implications for the function of disordered eating to decrease distress

(Corstorphine et al., 2007). Overall, performance on the PASAT-C was comparable across

groups, suggesting there are not differences in the cognitive functioning required for the task,

thus clinical intervention for EDs may prioritize managing strong, negative emotion. Individuals

with EDs may benefit from treatments that focus on fostering greater acceptance of one's

emotional experiences, such as Dialectical Behavior Therapy (Chen et al., 2015; Safer,

Robinson, & Jo, 2010) or Acceptance and Commitment Therapy (Berman, Boutelle, & Crow,

2009; Juarascio et al., 2013; Juarascio, Forman, & Herbert, 2010). Our study supports the

proposed model that negative emotions may perpetuate and maintain EDs, with the caveat that

the experience of distress may manifest physiologically and behaviorally similarly to HCs.

1   References

2   Aldao, A., Nolen-Hoeksema, S., & Schweizer, S. (2010). Emotion-regulation strategies across

3   psychopathology: A meta-analytic review. *Clinical Psychology Review*, *30*, 217–237.

4   doi.org/10.1016/j.cpr.2009.11.004

5   American Psychiatric Association. (2000). *Diagnostic and statistical manual of mental disorders*

6   (4th ed., text revision). Washington, DC.

7   Anestis, M.D., Lavender, J.M., Marshall-Berenz, E.C., Gratz, K.L., Tull, M.T. & Joiner, T.E.

8   (2012). Evaluating distress tolerance measures: Interelations and associations with

9   impulsive behaviors. *Cognitive Therapy Research, 36*, 593-602.

https://doi.org/10.1007/s10608-011-9377-8

Anestis, M. D., Peterson, C. B., Bardone-Cone, A. M., Klein, M. H., Mitchell, J. E., Crosby, R.

D., … Joiner, T. E. (2009). Affective lability and impulsivity in a clinical sample of women

with bulimia nervosa: The role of affect in severely dysregulated behavior. *International*

Journal of Eating Disorders, *42*, 259–266. doi.org/10.1002/eat.20606

Anestis, M. D., Selby, E. A., Fink, E. L., & Joiner, T. E. (2007). The multifaceted role of distress

tolerance in dysregulated eating behaviors. *International Journal of Eating Disorders*, *40*,

718–726. https://doi.org/10.1002/eat.20471

Anestis, M. D., Smith, A. R., Fink, E. L., & Joiner, T. E. (2009). Dysregulated eating and

distress: Examining the specific role of negative urgency in a clinical sample. *Cognitive*

Therapy and Research, *33*, 390–397. doi.org/10.1007/s10608-008-9201-2

Baiano, M., Salvo, P., Righetti, P., Cereser, L., Baldissera, E., Camponogara, I., & Balestrieri,

M. (2014). Exploring health-related quality of life in eating disorders by a cross-sectional

study and a comprehensive review. *BMC Psychiatry*, *14*. doi.org/10.1186/1471-244x-14-

165

Beauchaine, T. P. (2015). Respiratory sinus arrhythmia: A transdiagnostic biomarker of emotion

dysregulation and psychopathology. *Current Opinion in Psychology*, *3*, 43–47.

doi.org/10.1016/j.copsyc.2015.01.017

Berman, M. I., Boutelle, K. N., & Crow, S. J. (2009). A case series investigating acceptance and

commitment therapy as a treatment for previously treated, unremitted patients with anorexia

nervosa. *European Eating Disorders Review*, *17*, 426–434. doi.org/10.1002/erv.962

Boucsein, W. (2011). *Electrodermal Activity* (2nd ed.). New York, NY, US: Springer.

Boucsein, W., Fowles, D., Grimnes, S., Ben-Shakhar, G., Roth, W., Dawson, M., & Filion, D.

(2012). Publication recommendations for electrodermal measurements. *Psychophysiology*,

49, 1017–1034. doi.org/0.1111/j.1469-8986.2012.01384.x

Chen, E. Y., Segal, K., Weissman, J., Zeffiro, T. A., Gallop, R., Linehan, M. M., … Lynch, T. R.

(2015). Adapting dialectical behavior therapy for outpatient adult anorexia nervosa—a pilot

study. *International Journal of Eating Disorders*, *48*, 123–132. doi.org/10.1002/erv.962

Claes, L., Vandereycken, W., & Vertommen, H. (2005). Impulsivity-related traits in eating

disorder patients. *Personality and Individual Differences*, *39*, 739–749.

doi.org/10.1016/j.paid.2005.02.022

Cohen, J. (1988). *Statistical power analysis for the behavioral sciences.* Hillsdale, NJ: Lawrence

Earlbaum Associates, 20-26.

Cooper, Z., Cooper, P.J., & Fairburn, C.G. (1989). The validity of the eating disorder

examination and its subscales. *British Journal of Psychiatry, 154*, 807-812.

https://doi.org/10.1192/bjp.154.6.807

Corstorphine, E., Mountford, V., Tomlinson, S., Waller, G., & Meyer, C. (2007). Distress

tolerance in the eating disorders. *Eating Behaviors*, *8*, 91–97.

doi.org/10.1016/j.eatbeh.2006.02.003

Crowell, S. E., Beauchaine, T. P., & Linehan, M. M. (2009). A biosocial developmental model

of borderline personality: Elaborating and extending Linehan's theory. *Psychological*

Bulletin, *135*, 495–510. doi.org/10.1037/a0015616

Daughters, S. B., Lejuez, C. W., Kahler, C. W., Strong, D. R., & Brown, R. A. (2005).

Psychological distress tolerance and duration of most recent abstinence attempt among

residential treatment-seeking substance abusers. *Psychology of Addictive Behaviors*, *19*,

208–211. doi.org/10.1037/0893-164X.19.2.208

de Zwaan, de Z. (2001). Binge eating disorder and obesity. *International Journal of Obesity and*

Related Metabolic Disorders : Journal of the International Association for the Study of

Obesity, *25 Suppl 1*, S51-5. doi.org/10.1038/sj.ijo.0801699

Dingemans, A. E., & van Furth, E. F. (2012). Binge eating disorder psychopathology in normal

weight and obese individuals. *International Journal of Eating Disorders*, *45*, 135–138.

doi.org/10.1002/eat.20905

Eichen, D. M., Chen, E., Boutelle, K. N., & McCloskey, M. S. (2017). Behavioral evidence of

emotion dysregulation in binge eaters. *Appetite*, *111*, 1–6.

doi.org/10.1016/j.appet.2016.12.021

Fairburn, C. G., Cooper, Z., & O'Connor, M. (2008). Eating Disorder Examination (Edition

16.0D). In *Cognitive behavior therapy and eating disorders* (pp. 309–314). New York, NY:

Guilford Press.

Fairburn, C. G., Cooper, Z., & Shafran, R. (2003). Cognitive behaviour therapy for eating

disorders: A "transdiagnostic" theory and treatment. *Behaviour research and therapy, 41*,

509-528. https://doi.org/10.1016/s0005-7967(02)00088-8

Fairburn, C. G., Shafran, R., & Cooper, Z. (1999). A cognitive behavioural theory of anorexia

nervosa. *Behaviour Research and Therapy*, *37*, 1–13. doi.org/10.1016/s0005-

7967(98)00102-8

Faul, F., Erdfelder, E., Lang, A. G., & Buchner, A. (2007). G* Power 3: A flexible statistical

power analysis program for the social, behavioral, and biomedical sciences. *Behavior*

research methods, 39, 175-191.

Feldner, M. T., Leen-Feldner, E. W., Zvolensky, M. J., & Lejuez, C. W. (2006). Examining the

association between rumination, negative affectivity, and negative affect induced by a paced

auditory serial addition task. *Journal of Behavior Therapy and Experimental Psychiatry*, *37*,

171–187. doi.org/10.1016/j.jbtep.2005.06.002

Field, A. (2013). *Discovering statistics using IBM SPSS statistics.* Sage.

First, M. B., Spitzer, R. L., Gibbon, M., & Williams, J. B. W. (2002). *Structured clinical*

interview for DSM-IV-TR axis I disorders, research version, patient edition. SCID-I/P. New

York, NY: New York State Psychiatric Institute.

Fischer, S., Smith, G. T., & Cyders, M. A. (2008). Another look at impulsivity: A meta-analytic

review comparing specific dispositions to rash action in their relationship to bulimic

symptoms. *Clinical Psychology Review*, *28*, 1413–1425. doi.org/10.1016/j.cpr.2008.09.001

Friederich, H.-C., Schild, S., Schellberg, D., Quenter, A., Bode, C., Herzog, W., & Zipfel, S.

(2006). Cardiac parasympathetic regulation in obese women with binge eating disorder.

International Journal of Obesity, *30*, 534–542. doi.org/10.1038/sj.ijo.0803181

Gratz, K. L., Rosenthal, M. Z., Tull, M. T., Lejuez, C. W., & Gunderson, J. G. (2006). An

experimental investigation of emotion dysregulation in borderline personality disorder.

Journal of Abnormal Psychology, *115*, 850–855. doi.org/10.1037/0021-843X.115.4.850

Gross, J. J. (2013). Emotion regulation: Taking stock and moving forward. *Emotion*, *13*, 359–

365. doi.org/10.1037/a0032135

Grossman, P., Stemmler, G., & Meinhardt, E. (1990). Paced respiratory sinus arrhythmia as an

index of cardiac parasympathetic tone during varying behavioral tasks. *Psychophysiology*,

27, 404–416. doi.org/10.1111/j.1469-8986.1990.tb02335.x

Grossman, P., & Taylor, E. W. (2007). Toward understanding respiratory sinus arrhythmia:

Relations to cardiac vagal tone, evolution and biobehavioral functions. *Biological*

Psychology, *74*, 263–285. doi.org/10.1016/j.biopsycho.2005.11.014

Haedt-Matt, A. A., & Keel, P. K. (2011). Revisiting the affect regulation model of binge eating:

A meta-analysis of studies using ecological momentary assessment. *Psychological Bulletin*,

137, 660–681. doi.org/10.1037/a0023660

Haynos, A. F., & Fruzzetti, A. E. (2011). Anorexia nervosa as a disorder of emotion

dysregulation: Evidence and treatment implications. *Clinical Psychology: Science and*

Practice, *18*, 183–202. doi.org/10.1111/j.1468-2850.2011.01250.x

Hilbert, A., & Tuschen-Caffier, B. (2007). Maintenance of binge eating through negative mood:

A naturalistic comparison of binge eating disorder and bulimia nervosa. *International*

Journal of Eating Disorders, *40*, 521–530. doi.org/10.1002/eat.20401

Hilbert, A., Vögele, C., Tuschen-Caffier, B., & Hartmann, A. S. (2011). Psychophysiological

responses to idiosyncratic stress in bulimia nervosa and binge eating disorder. *Physiology &*

Behavior, *104*, 770–777. doi.org/10.1016/j.physbeh.2011.07.013

Holdwick, D. J. J., & Wingenfeld, S. A. (1999). The subjective experience of PASAT testing:

Does the PASAT induce negative mood? *Archives of Clinical Neuropsychology*, *14*, 273–

284. doi.org/10.1093/arclin/14.3.273

Hudson, J. I., Hiripi, E., Pope, H. G. J., & Kessler, R. C. (2007). The prevalence and correlates of eating disorders in the National Comorbidity Survey Replication. *Biological Psychiatry*, *61*, 348–358. doi.org/10.1016/j.biopsych.2006.03.040

Hudson, J. I., Lalonde, J. K., Coit, C. E., Tsuang, M. T., McElroy, S. L., Crow, S. J., … Pope, H. G. J. (2010). Longitudinal study of the diagnosis of components of the metabolic syndrome in individuals with binge-eating disorder. *The American Journal of Clinical Nutrition*, *91*, 1568–1573. doi.org/10.3945/ajcn.2010.29203

Johnson, J. G., Spitzer, R. L., & Williams, J. B. W. (2001). Health problems, impairment and illnesses associated with bulimia nervosa and binge eating disorder among primary care and obstetric gynaecology patients. *Psychological Medicine*, *31*, 1455–1466. doi.org/10.1017/s0033291701004640

Juarascio, A. S., Forman, E. M., & Herbert, J. D. (2010). Acceptance and commitment therapy versus cognitive therapy for the treatment of comorbid eating pathology. *Behavior Modification*, *34*, 175–190. doi.org/10.1177/0145445510363472

Juarascio, A., Shaw, J., Forman, E., Timko, C. A., Herbert, J., Butryn, M., … Lowe, M. (2013). Acceptance and commitment therapy as a novel treatment for eating disorders: an initial test of efficacy and mediation. *Behavior Modification*, *37*, 459–489. doi.org/10.1177/0145445513478633

Karason, K., Mølgaard, H., Wikstrand, J., & Sjöström, L. (1999). Heart rate variability in obesity and the effect of weight loss. *The American Journal of Cardiology*, *83*, 1242–1247. doi.org/10.1016/s0002-9149(99)00066-1

Kessler, R. C., Berglund, P. A., Chiu, W. T., Deitz, A. C., Hudson, J. I., Shahly, V., … Xavier,

M. (2013). The prevalence and correlates of binge eating disorder in the World Health

Organization world mental health surveys. *Biological Psychiatry*, *73*, 904–914.

doi.org/10.1016/j.biopsych.2012.11.020

Klatzkin, R. R., Gaffney, S., Cyrus, K., Bigus, E., & Brownley, K. A. (2015). Binge eating

disorder and obesity: Preliminary evidence for distinct cardiovascular and psychological

phenotypes. *Physiology & Behavior*, *142*, 20–27. doi.org/10.1016/j.physbeh.2015.01.018

Klein, D. N., Ouimette, P. C., Kelly, H. S., Ferro, T., & Riso, L. P. (1994). Test-retest reliability

of team consensus best-estimate diagnoses of axis I and II disorders in a family study.

American Journal of Psychiatry, *151*, 1043–1047. doi.org/10.1176/ajp.151.7.1043

Kosten, T. A., & Rounsaville, B. J. (1992). Sensitivity of psychiatric diagnosis based on the best

estimate procedure. *The American Journal of Psychiatry*, *149*, 1225-1227.

doi.org/10.1176/ajp.149.9.1225

Kreibig, S. D. (2010). Autonomic nervous system activity in emotion: A review. *Biological

Psychology*, *84*, 394–421. doi.org/10.1016/j.biopsycho.2010.03.010

Laborde, S., Mosley, E., & Thayer, J. F. (2017). Heart Rate Variability and Cardiac Vagal Tone

in Psychophysiological Research–Recommendations for Experiment Planning, Data

Analysis, and Data Reporting. *Frontiers in psychology, 8*, 1-18.

https://doi.org/10.3389/fpsyg.2017.00213

Lang, P., Greenwald, M. K., Bradley, M. M., & Hamm, A. O. (1993). Looking at pictures:

Affective, facial, visceral, and behavioral reactions. *Psychophysiology*, *30*, 261–273.

doi.org/10.1111/j.1469-8986.1993.tb03352.x

Le Grange, D., Swanson, S. A., Crow, S. J., & Merikangas, K. R. (2012). Eating disorder not

otherwise specified presentation in the US population. *International Journal of Eating*

Disorders, *45*, 711–718. doi.org/10.1002/eat.22006

Leehr, E. J., Krohmer, K., Schag, K., Dresler, T., Zipfel, S., & Giel, K. E. (2015). Emotion

regulation model in binge eating disorder and obesity—A systematic review. *Neuroscience*

and Biobehavioral Reviews, *49*, 125–134. doi.org/10.1016/j.neubiorev.2014.12.008

Lejuez, C. W., Kahler, C. W., & Brown, R. A. (2003). A modified computer version of the Paced

Auditory Serial Addition Task (PASAT) as a laboratory-based stressor. *The Behavior*

Therapist, *26*, 290–293.

Leyro, T. M., Zvolensky, M. J., & Bernstein, A. (2010). Distress tolerance and

psychopathological symptoms and disorders: A review of the empirical literature among

adults. *Psychological Bulletin*, *136*, 576. doi.org/10.1037/a0019712

Lobbestael, J., Leurgans, M., & Arntz, A. (2011). Inter-rater reliability of the Structured Clinical

Interview for DSM-IV Axis I disorders (SCID I) and Axis II disorders (SCID II). *Clinical*

psychology & psychotherapy, 18, 75-79.

Masi, C. M., Hawkley, L. C., Rickett, E. M., & Cacioppo, J. T. (2007). Respiratory sinus

arrhythmia and diseases of aging: Obesity, diabetes mellitus, and hypertension. *Biological*

Psychology, *74*, 212–223. doi.org/10.1016/j.biopsycho.2006.07.006

Mazurak, N., Enck, P., Muth, E., Teufel, M., & Zipfel, S. (2011). Heart rate variability as a

measure of cardiac autonomic function in anorexia nervosa: A review of the literature.

European Eating Disorders Review, *19*, 87–99. doi.org/10.1002/erv.1081

Messerli-Bürgy, N., Engesser, C., Lemmenmeier, E., Steptoe, A., & Laederach-Hofmann, K.

(2010). Cardiovascular stress reactivity and recovery in bulimia nervosa and binge eating

disorder. *International Journal of Psychophysiology*, *78*, 163–168.

doi.org/10.1016/j.ijpsycho.2010.07.005

Naumann, E., Tuschen-Caffier, B., Voderholzer, U., Caffier, D., & Svaldi, J. (2015). Rumination

but not distraction increases eating-related symptoms in anorexia and bulimia nervosa.

Journal of Abnormal Psychology, *124*, 412-420. doi.org/10.1037/abn0000046

Padierna, A., Quintana, J. M., Arostegui, I., Gonzalez, N., & Horcajo, M. J. (2000). The health-

related quality of life in eating disorders. *Quality of Life Research*, *9*, 667–674.

doi.org/10.1007/springerreference_73510

Peschel, S. K. V., Feeling, N. R., Vögele, C., Kaess, M., Thayer, J. F., & Koenig, J. (2016). A

meta-analysis on resting state high-frequency heart rate variability in bulimia nervosa.

European Eating Disorders Review, *24*, 355–365. doi.org/10.1002/erv.2454

Peterson, C. M., & Fischer, S. (2012). A prospective study of the influence of the UPPS model of

impulsivity on the co-occurrence of bulimic symptoms and non-suicidal self-injury. *Eating

Behaviors*, *13*, 335–41. doi.org/10.1016/j.eatbeh.2012.05.007

Rissanen, P., Franssila-Kallunki, A., & Rissanen, A. (2001). Cardiac parasympathetic activity is

increased by weight loss in healthy obese women. *Obesity Research*, *9*, 637–643.

doi.org/10.1038/oby.2001.84

Safer, D. L., Robinson, A. H., & Jo, B. (2010). Outcome from a randomized controlled trial of

group therapy for binge eating disorder: Comparing dialectical behavior therapy adapted for

binge eating to an active comparison group therapy. *Behavior Therapy*, *41*, 106–120.

doi.org/10.1016/j.beth.2009.01.006

Salters-Pedneault, K., Gentes, E., & Roemer, L. (2007). The role of fear of emotion in distress,

arousal, and cognitive interference following an emotional stimulus. *Cognitive Behaviour

Therapy*, *36*, 12–22. doi.org/10.1080/16506070600874281

Sauer, S. E., & Baer, R. A. (2012). Ruminative and mindful self-focused attention in borderline

personality disorder. *Personality Disorders: Theory, Research, and Treatment*, *3*, 433–441.

doi.org/10.1037/a0025465

Schloss, H. M., & Haaga, D. A. F. (2011). Interrelating behavioral measures of distress tolerance

with self-reported experiential avoidance. *Journal of Rational-Emotive & Cognitive-*

Behavior Therapy, *29*, 53–63. doi.org/10.1007/s10942-011-0127-3

Schmahl, C., Meinzer, M., Zeuch, A., Fichter, M., Cebulla, M., Kleindienst, N….Bohus, M.

(2010). Pain sensitivity is reduced in borderline personality disorder, but not in post

traumatic stress disorder and bulimia nervosa. *The World Journal of Biological Psychiatry,*

11, 364-371. https://doi.org/10.1080/15622970701849952

Svaldi, J., Griepenstroh, J., Tuschen-Caffier, B., & Ehring, T. (2012). Emotion regulation deficits

in eating disorders: A marker of eating pathology or general psychopathology? *Psychiatry*

Research, *197*, 103–111. doi.org/10.1016/j.psychres.2011.11.009

Svaldi, J., Tuschen-Caffier, B., Lackner, H. K., Zimmermann, S., & Naumann, E. (2012). The

effects of emotion regulation on the desire to overeat in restrained eaters. *Appetite*, *59*, 256–

263. doi.org/10.1016/j.appet.2012.04.016

Thayer, J. F., Åhs, F., Fredrikson, M., Sollers, J. J. I. I. I., & Wager, T. D. (2012). A meta-

analysis of heart rate variability and neuroimaging studies: Implications for heart rate

variability as a marker of stress and health. *Neuroscience and Biobehavioral Reviews*, *36*,

747–756. doi.org/10.1016/j.neubiorev.2011.11.009

Tuschen-Caffier, B., & Vögele, C. (1999). Psychological and physiological reactivity to stress:

An experimental study on bulimic patients, restrained eaters and controls. *Psychotherapy*

and Psychosomatics, *68*, 333–340. doi.org/10.1159/000012352

Udo, T., Weinberger, A. H., Grilo, C. M., Brownell, K. D., DiLeone, R. J., Lampert, R., …

McKee, S. A. (2014). Heightened vagal activity during high-calorie food presentation in

obese compared with non-obese individuals - results of a pilot study. *Obesity Research &*

Clinical Practice, *8*, e201–e298. doi.org/10.1016/j.orcp.2013.05.006

Watson, D., Clark, L. A., & Tellegen, A. (1988). Development and validation of brief measures

of positive and negative affect: The PANAS scales. *Journal of Personality and Social*

Psychology, *54*, 1063–1070. doi.org/10.1037/0022-3514.54.6.1063

Wenzel, K. R., Weinstock, J., Vander Wal, J. S., & Weaver, T. L. (2014). Examining the role of

negative urgency in a predictive model of bulimic symptoms. *Eating Behaviors*, *15*, 343–

349. doi.org/10.1016/j.eatbeh.2014.04.014

Wilfley, D.E., Schwartz, M.B., Spurrell, E.B. & Fairburn, C.G. (1999). Using the eating disorder

examination to identify the specific psychopathology of binge eating disorder. *International*

Journal of Eating Disorders, 27, 259-269. https://doi.org/10.1002/(sici)1098-

108x(200004)27:3<259::aid-eat2>3.3.co;2-7

Wilson, G.T. & Smith, D. (1989). Assessment of bulimia nervosa: An evaluation of the Eating

Disorder Examination. *International Journal of Eating Disorders*, 8, 173-179.

https://doi.org/10.1002/1098-108x(198903)8:2<173::aid-eat2260080206>3.0.co;2-v

Wu, M., Giel, K. E., Skunde, M., Schag, K., Rudofsky, G., de Zwaan, M., … Friederich, H.

(2013). Inhibitory control and decision making under risk in bulimia nervosa and binge-

eating disorder. *International Journal of Eating Disorders*, *46*, 721–728.

doi.org/10.1002/eat.22143

Figure 1

Schematic of the procedure and time periods assessed.

Visual Analogue Scale: self-reported negative emotions, happiness, urges to binge eat

Psychophysiology: respiratory sinus arrhythmia, skin conductance response, tonic skin conductance levels

Table 1

Severity of eating disorder symptoms, demographic information and clinical characteristics of participants at baseline.

Variable	Anorexia Nervosa N = 14		Binge Eating Disorder N = 55		Bulimia Nervosa N = 37		Healthy Controls N = 26		Total Sample N = 132	
	M	SD	M	SD	M	SD	M	SD	M	SD
Body Mass Index[1]	18.82	3.67	36.42	9.47	25.45	5.88	26.68	4.61	29.56	9.50
EDE Global[2]	3.27	1.69	3.12	0.9	3.22	1.04	0.39	0.52	2.63	1.48
EDE Eating Concern[2]	3.27	1.94	2.82	1.42	2.76	1.15	0.04	0.08	2.30	1.69
EDE Dietary Restraint [3]	3.40	1.91	2.19	1.26	3.09	1.33	0.48	0.89	2.24	1.62
EDE Weight Concern[2]	2.90	1.83	3.59	1.07	3.42	1.41	0.48	0.67	2.85	1.69
EDE Shape Concern[2]	3.49	1.70	3.9	1.25	3.62	1.35	0.55	0.79	3.12	1.79
Objective Binge Episodes (3 months)[4]	1.21	2.51	4.58	4.11	6.29	6.04	0	0	3.80	4.82
Compensatory Behaviors Episodes (3 months)[5]	3.54	4.96	0.32	1.13	8.37	8.41	0	0	2.85	5.95
Age in years	34.64	11.47	41.65	9.49	29.81	9.77	28.81	10.59	35.06	11.47
	N	%	N	%	N	%	N	%	N	%
Ethnic minority	2	14.3	23	41.8	11	29.7	11	42.3	49	36.84
Single	9	64.3	24	43.6	22	59.5	15	62.5	70	52.63
Employed or Full-Time Student	13	92.9	51	92.7	33	89.2	25	96.2	122	91.73
Completed 4-year college	11	78.6	43	78.2	23	62.2	14	53.8	91	68.42
Low Income (< 25 000)	10	71.4	16	29.1	20	54.1	17	65.4	61	45.86
Mood Disorder	7	50	35	63.6	27	73	0	0	69	51.88
Anxiety Disorder	9	64.3	20	36.4	12	32.4	0	0	41	30.83
Medical Comorbidity	7	50	26	47.3	22	59.5	4	15.4	59	44.36
Medications	6	42.9	28	50.9	18	48.6	2	7.7	54	40.6

Notes:
[1] HCs had significantly lower BMI relative to the BED group ($p < .001$) and significantly higher BMI relative to the AN group ($p < .01$), but did not differ significantly from the BN group.
[2] Scores from the EDE Eating Concern subscale, Weight Concern subscale, Shape Concern subscale or Global scores did not differ significantly between the BN, BED or AN groups ($ps > .23$) but were significantly higher relative to HCs ($ps < .001$).
[3] AN and BN groups did not differ significantly on the EDE Dietary Restraint subscale ($p = .47$), but the BED group reported significantly lower dietary restraint relative to AN and BN and significantly greater dietary restraint relative to HCs ($ps < .001$).
[4] BED and BN reported significantly greater frequency of binge eating episodes than AN ($ps < .008$). BED and BN did not differ significantly on frequency of binge eating episodes ($p = .15$).
[5] BN and AN reported significantly greater frequency of compensatory behaviors ($ps < .001$) than BED, BN and AN did not differ significantly on frequency of compensatory behaviors ($p = .84$).

Table 2

Self-reported negative emotions, happiness and urges to binge eat at baseline, during the Paced

Auditory Serial Addition Task – Computerized (PASAT-C), and during recovery from PASAT-C

among women with Bulimia Nervosa, Binge Eating Disorder, Anorexia Nervosa and Healthy

Controls.

	Baseline M (SD)	PASAT-C M (SD)	Recovery M (SD)
Negative Emotions			
Anorexia Nervosa	48.99 (*22.51*)	60.80 (*26.58*)	47.53 (*21.66*)
Binge Eating Disorder	39.40 (*22.82*)	57.24 (*20.20*)	30.55 (*18.25*)
Bulimia Nervosa	38.90 (*22.83*)	58.38 (*20.22*)	35.90 (*20.33*)
Healthy Control	8.91 (*13.18*)	37.07 (*24.47*)	15.40 (*16.38*)
Happiness			
Anorexia Nervosa	37.56 (*25.16*)	22.47 (*26.08*)	17.68 (*24.51*)
Binge Eating Disorder	40.85 (*18.05*)	23.52 (*25.07*)	29.20 (*21.58*)
Bulimia Nervosa	43.98 (*18.74*)	21.03 (*21.58*)	29.50 (*23.21*)
Healthy Control	59.41 (*19.40*)	28.51 (*28.71*)	39.42 (*28.14*)
Urge to Binge			
Anorexia Nervosa	21.70 (*29.03*)	11.42 (*25.86*)	20.41 (*33.67*)
Binge Eating Disorder	46.87 (*27.10*)	42.41 (*36.48*)	44.21 (*32.50*)
Bulimia Nervosa	33.15 (*28.48*)	31.84 (*36.86*)	34.77 (*32.55*)
Healthy Control	.77 (*1.97*)	2.98 (*11.41*)	.34 (*.71*)

Table 3

Completion time and latency to quit on the Paced Auditory Serial Addition Task – Computerized

(PASAT- C) among women with Bulimia Nervosa, Binge Eating Disorder, Anorexia Nervosa and

Healthy Controls.

	PASAT-C Completion N (%)	PASAT-C Latency to Quit (seconds) M (*SD*)
Anorexia Nervosa	10 (71.4%)	678.64 (*103.62*)
Binge Eating Disorder	38 (69.1%)	683.84 (*89.32*)
Bulimia Nervosa	23 (62.2%)	643.84 (*128.05*)
Healthy Control	22 (84.6%)	690.85 (*110.35*)

Table 4

Respiratory sinus arrhythmia, skin conductance responses and tonic skin conductance levels for

Bulimia Nervosa, Binge Eating Disorder, Anorexia Nervosa and Healthy Controls at baseline,

during PASAT-C, and during recovery from PASAT-C.

	Baseline *M* (SD)	PASAT-C *M* (SD)	Recovery *M* (SD)
Respiratory sinus arrhythmia (in ms^2)			
Anorexia Nervosa	5.81 (1.20)	5.69 (1.16)	6.15 (1.17)
Binge Eating Disorder	5.25 (1.36)	5.16 (1.39)	5.40 (1.32)
Bulimia Nervosa	6.25 (1.16)	6.05 (1.06)	6.32 (1.09)
Healthy Control	6.21 (1.36)	6.06 (1.30)	6.38 (1.27)
Skin conductance response			
Anorexia Nervosa	1.80 (2.42)	3.96 (4.15)	3.91 (2.69)
Binge Eating Disorder	2.13 (1.90)	3.78 (3.02)	3.88 (2.75)
Bulimia Nervosa	1.24 (1.70)	3.54 (3.45)	2.77 (2.39)
Healthy Control	1.83 (1.72)	2.90 (2.54)	3.94 (2.94)
Tonic skin conductance levels (in µs)			
Anorexia Nervosa	2.19 (2.11)	6.38 (6.58)	6.72 (5.67)
Binge Eating Disorder	2.94 (2.27)	5.14 (4.06)	5.06 (4.05)
Bulimia Nervosa	2.35 (3.89)	5.28 (6.51)	4.98 (6.40)
Healthy Control	2.40 (3.11)	5.37 (6.04)	5.77 (6.76)