Live imaging of heart tube development in mouse reveals alternating phases of

cardiac differentiation and morphogenesis

Kenzo Ivanovitch[1,2], Susana Temiño[1] and Miguel Torres[1,*]

[1]Developmental Biology Program, Centro Nacional de Investigaciones

Cardiovasculares (CNIC), Madrid, Spain

[2]Current Address, The Francis Crick Institute, London, UK

Authors ORCID

Miguel Torres: orcid.org/0000-0003-0906-4767

Kenzo Ivanovitch: orcid.org/0000-0003-4706-7686

*Corresponding author: mtorres@cnic.es

ABSTRACT

During vertebrate heart development two progenitor populations, first and second

heart fields (FHF, SHF), sequentially contribute to longitudinal subdivisions of the

heart tube (HT), with the FHF contributing the left ventricle and part of the atria, and

the SHF the rest of the heart. Here we study the dynamics of cardiac differentiation

and morphogenesis by tracking individual cells in live analysis of mouse embryos.

We report that during an initial phase, FHF precursors differentiate rapidly to form a

cardiac crescent, while limited morphogenesis takes place. In a second phase, no

differentiation occurs while extensive morphogenesis, including splanchnic

mesoderm sliding over the endoderm, results in HT formation. In a third phase,

cardiac precursor differentiation resumes and contributes to SHF-derived regions and

the dorsal closure of the HT. These results reveal tissue-level coordination between

morphogenesis and differentiation during HT formation and provide a new

framework to understand heart development.

INTRODUCTION

The heart is the first organ to form and function during embryonic development. At

embryonic day (E) 7.5, cardiac precursors in the splanchnic mesoderm (mesoderm

apposed to the endoderm) differentiate into cardiomyocytes by assembling the

contractile sarcomere machinery (Tyser et al., 2016) and form a bilateral structure

known as the cardiac crescent (cc) in the mouse. Concomitant with foregut

invagination, the cc swings inwards to become placed underneath the developing head

folds. By a complex morphogenetic process, the cc subsequently transforms into an

early heart tube (HT) initially opened dorsally, which by E8.25 has transformed into a

closed and beating linear HT, also known as the primitive HT (Evans et al., 2010;

Kelly et al., 2014).

The cc and early HT mainly give rise to the left ventricle (Zaffran et al., 2004). The

right ventricle (RV), the outflow tract and most of the atria derive instead from

cardiac progenitors located dorso-medially to the cc in the splanchnic mesoderm, that

are progressively recruited at the poles of the HT at subsequent developmental stages

(Cai et al., 2003; Kelly et al., 2001; Mjaatvedt et al., 2001; Waldo et al., 2001; Galli et

al., 2008 135:1157-67). These findings led to the concept that cardiac mesodermal

progenitors contain two populations of cells: the first heart field (FHF) precursors,

recruited early in development to form the initial HT and mostly containing the LV

primordium, and the second heart field (SHF), recruited later and elongating the HT

(Buckingham et al., 2005).

Clonal analysis (Devine et al., 2014; Lescroart et al., 2014; Meilhac et al., 2004a)

supports the idea that FHF and SHF precursors are two independent developmental

fields with dedicated molecular pathways. Clonal analysis also showed that the SHF

shares a common origin with the skeletal muscles of the head and neck within the

pharyngeal mesoderm (Lescroart et al. 2010, Lescroart et al. 2015), further supporting

differences between FHF and SHF populations. However, the existence of a common

precursor between FHF and SHF was also reported in the early mouse embryo

(Meilhac et al., 2004a) and other views suggest that the heart would form by a

continuous differentiation process from a single population of cardiac precursors and

only timing of recruitment would distinguish cells of the FHF and SHF (Abu-Issa et

al., 2004; Moorman et al., 2007). In support of the latter, the typical marker of the

SHF, Islet1 (Cai et al., 2003), is also transiently expressed in FHF precursors and

must therefore be considered as widespread cardiac progenitor marker instead (Cai et

al., 2003; Prall et al., 2007; Yuan and Schoenwolf, 2000). Whether the recruitment of

cardiomyocytes from progenitors is a continuous process and how this coordinates

with morphogenesis, however, has not been directly studied. This is partly because

the spatial arrangement of progenitors and differentiated cardiomyocytes has so far

been analyzed on fixed embryos (Cai et al., 2003; Spater et al., 2013) and the

expression dynamics of genes reporting differentiation together with cell movements

during HT morphogenesis have not been captured so far (Abu-Issa, 2014).

Here, we report the live-imaging and 3D+t cell tracking of HT formation in whole

mouse embryos. Using this method, in conjunction with an Nkx2.5eGFP reporter line,

which provides high level of GFP in differentiated cardiomyocytes, we studied the

dynamics of cardiac field differentiation. During an initial phase, FHF cardiac

precursors differentiate rapidly to form a cardiac crescent, while limited

morphogenesis takes place. During a second phase, no differentiation events are

detected and extensive morphogenesis, including splanchnic mesoderm sliding over

the endoderm, results in HT formation. Finally, using an Isl1-Cre lineage tracing

assay combined with live-imaging, we show that during a third phase, cardiac

precursor differentiation resumes and contributes not only to the known SHF-derived

regions but also to the dorsal aspect of the HT. These results show essential properties

of FHF and SHF contribution to heart development and reveal tissue-level

coordination between alternating phases of differentiation and morphogenesis during

HT formation.

RESULTS

3D static analysis of mouse HT formation

To assess how the initial cardiogenic region transforms into a HT and differentiates,

we first analyzed $Nkx2.5^{cre/+}$; $Rosa26Rtdtomato^{+/-}$ embryos, in which both cardiac

precursors and cardiomyocytes are labeled (Stanley et al., 2002). Before cc

differentiation, at early head fold stage (EHF, ~E7.5), the cardiogenic region is

visualized as a flat horse shoe-shaped tdtomato+ mesodermal layer at the rostral

border of the embryo (Figure 1A, A', and Video 1). Figure 1-figure supplement-1

shows the criteria for embryo staging (Downs and Davies, 1993; Lawson, 2016). In

the $Nkx2.5^{cre/+}$; $Rosa26Rtdtomato^{+/-}$ embryos, tdtomato labeling is also observed in the

endocardium and endothelial cells (Stanley et al., 2002) but not in the endoderm

(Figure 1-figure supplement 2A, A'). We next studied the distribution of Cardiac

troponin T (cTnnT), one of the first evident sarcomeric proteins to appear in the

cardiac crescent (Tyser et al., 2016). At EHF stage (Figure 1B), while most embryos

are negative for cTnnT expression, some embryos show weak cTnnT localization in

subsets of cells (Figure 1-figure supplement 3A, A'). At a subsequent embryonic

stage (~E7.7), cTnnT signal reveals the cc, which is folding inwards. During folding,

the cTnnT signal increases. cTnnT+ cells are initially columnar epithelial cells and

show apical localization of the tight junction component zona-occluden-1 (ZO-1)

(Figure 1-figure supplement 3B, B'). During differentiation, cardiac precursors switch

to a rounded shape (Linask et al., 1997) (Figure 1C, D) and separate from the

endoderm, while maintaining a basal lamina at the endocardial side (inset in Figure

1D and Figure 2D). Morphogenetic changes starting at ~E8 subsequently lead to the

formation of a hemi-tube whose major axis is transversal to the embryo A-P axis. We

will refer to this stage as transversal HT (Figure 1E). Later, the tube adopts a more

spherical shape, very similar to the linear HT but still open dorsally. We will refer to this stage as open HT (Figure 1F). The HT eventually closes dorsally (Figure 1G, red arrows in Figure 1G'') and a prominent arterial pole (prospective RV) (Zaffran et al., 2004) becomes visible, completing linear HT formation by ~E8.25 (yellow arrows in Figure 1G'', Figure 1H, see also Video 2).

To assess the overall growth of the forming HT, we measured cTnnT+ tissue volume in segmented z-stacks, at the stages described above (Figure 1I and Supplementary file 1). During the first phase of cardiomyocyte differentiation, the cTnnT signal expands resulting in a cardiac crescent rapidly doubling in volume (Figure 1C', D', I). During the subsequent phase of morphogenesis, from cc to open HT stage, growth is less pronounced despite extensive morphological changes (Figure 1E', F', I). The Volume of the HT appears to increase again upon addition of the RV primordium to the arterial pole and dorsal HT closure (Figure 1G', I). HT growth likely reflects an increase in cell number occurring during the formation of the heart tube. Cardiomyocytes are proliferative at this stage (de Boer et al., 2012), and we can indeed observe mitotic figures in the HT (Figure 1J). From this analysis, however, it is unclear how much of the growth observed is due to cardiomyocyte proliferation versus addition of new cardiomyocytes from cardiac progenitor cells located in the splanchnic mesoderm.

3D analysis of cardiac progenitor differentiation

To visualize the boundary where cardiac progenitors abut differentiating cardiomyocytes during HT morphogenesis, we immunostained $Nkx2.5^{cre/+}$ $Rosa26Rtdtomato^{+/-}$ embryos at the transversal-HT stage with the differentiation marker cTnnT and acquired whole-mount images (Figure 2A). Cre activation of tdtomato was detected in both the FHF and SHF, as well as in the endoderm and endocardium (see optical sections in Figure 2A', A'', F, F' and Figure 2-figure supplement 1A) (Stanley et al. 2002). Cardiomyocytes are separated from the endoderm by the endocardium. In contrast, undifferentiated cardiac precursors lie medio-dorsally in direct contact with the endoderm in areas where endocardial cells are not detected (Figure 2A', A''). This is confirmed by the absence of the endothelial marker CD31 (Figure 2-figure supplement 1A, B, B'). We then rendered in 3D the

Nkx2.5^cre-labeled lineages, including both FHF and SHF, and the cTnnT+ tissues,

including only the FHF/cc, which allowed visualizing the boundary between SHF and

cc at the tissue level (Figure 2A, B, B', Figure 2-figure supplement 2A', Video 3 and

see material and methods –non-cardiac tdtomato signal was removed manually–).

To identify the changes associated to cardiomyocyte differentiation at the cellular

level, we labeled single cells with membrane-GFP (see Material and Methods).

cTnnT- progenitors have an epithelial-like columnar cell shape, while the

differentiated cTnnT+ cardiomyocytes are rounder and have lost the columnar

epithelial organization (Figure 2C-E and Figure 2-source data 1). This is reminiscent

of the cell shape transition observed in the distal outflow tract (OFT) at later stages of

heart development, when SHF progenitor-to-cardiomyocyte differentiation takes

place (Francou et al., 2014; Ramsbottom et al., 2014; Sinha et al., 2012). Interestingly,

some cells at the boundary zone exhibit weak cTnnT localization and yet show

columnar shapes typical of mesodermal cardiac precursors (Figure 2F-F'''' and red

arrows in Figure 2F''', F''''). Unlike differentiated cardiomyocytes, these cells do not

show rounded shapes, and therefore they may represent a transient state between

progenitors and differentiated cardiomyocytes; however, the nature of such state

cannot be addressed by static analysis. Differentiation of cardiac progenitors is thus

accompanied by changes in cell shape and detachment from the endoderm.

We next assessed the expression pattern of the *Nkx2.5eGFP* enhancer reporter line, in

which GFP expression is restricted to cardiomyocytes (Lien et al., 1999; Prall et al.,

2007; Wu et al., 2006). To characterize this reporter line in detail, we immunostained

Nkx2.5eGFP embryos against cTnnT, (Figure 3A, B and Figure 3- figure supplement

1A, B) and compared the relative intensities of cTnnT and GFP in manually-

segmented single cells (Figure 3C, D Figure 3-figure supplement 2A and Figure 3-

source data 1). We found that the GFP level varied linearly with cTnnT level (Figure

3D), although considerable variability of GPF levels was observed within each

cTnnT+ and cTnnT- cell population. Scoring of a large number of cells allowed to

reproducibly identify the top 50% GFP-expressing cells as positive for cTnnT+

(Figure 3-source data 1). Genetic tracing experiments using the *Nkx2.5^cre/+*

Rosa26Rtdtomato^+/- line instead show strong tdtomato level in both the FHF and SHF

(Figure 3- figure supplement 3A).

We next characterized the boundary between cardiomyocytes and cardiac precursors

in transversal HT stage embryos of the *Nkx2.5eGFP* reporter line. We measured mean

fluorescent intensity in manually segmented cells at the boundary zone and found that

GFP level and cTnnT signals correlate at the individual-cell level (Figure 3B', E,

Figure 3-figure supplement 2B, C and Figure 3-source data 1). Altogether, these

results indicate that the *Nkx2.5-eGFP* reporter is suitable for tracking cardiomyocytes

in live-imaging and reliably identifies the top 50% GFP-expressing cells as cTnnT-

positive.

2-photon live-imaging of early cardiac development in the mouse embryo reveals

morphogenetic events leading to HT formation

We next established a live-imaging method to dynamically characterize the formation

of the HT in the mouse embryo (Figure 4A) (Chen et al., 2014). We adapted a

previously reported culture system (Nonaka, 2009; Nonaka et al., 2002) in which the

whole mouse embryo is immobilized by inserting the extraembryonic region in a

holder (Figure 4C). After culture, embryos showed normal morphology, their hearts

were beating and circulation was initiated (Figure 4D, E and Video 4). This culture

system in combination with two-photon microscopy enabled the generation of high-

resolution 3D+t videos (Figure 3B and see Methods and Materials). Maximum culture

time and imaging achieved was 24 hours (Figure 4-figure supplement 1A, n=3).

However, typical acquisition times varied in most cases between 3 and 13 hours. The

rate of acquisition varied in most cases between 4 and 9 minutes, with some

exceptions depending on the specific aim of the recording (see video legends).

Imaging *Mesp1^{cre/+}; Rosa26Rtdtomato^{+/-}* embryos allowed tracking the anterior

mesoderm (Saga et al., 1996; Saga et al., 1999), including cardiac lineages, from cc

stage up to HT stage (Figure 4F, F' and Videos 5-6). The time-lapse analysis provided

insight on the formation of the endocardial lumen (Figure 4F', F''). The endocardium

is initially observed as a bilayer of cells and eventually splits into dorsal and ventral

layers, which move apart from each other allowing the lumen to become visible

between the layers (see Videos 6 and 7 for confocal views and video 8 for a bright

field view). Thin cytoplasmic bridges between the endocardial layers persist and

extend as the endocardial layers separate from each other (see white arrows in Figure

4F' and Video 6). Heartbeat becomes detectable around this stage and circulation in

the embryo is then initiated.

Imaging *Nkx2.5*^{cre/+}*; Rosa26RmT/mG*^{+/-} embryos, in which Cre-recombined cells

activate membrane-bound GFP, we could track cells during differentiation, as they

transit from a columnar to a round shape and start contracting (Figure 4G-G'', Video

7 and bright-field Video 8). We found that the endocardial lumen started to appear

while cc cells still remained columnar (compare time points 2h 13m and 3h 48m in

Figure 4G'). Cell rounding is therefore unlikely to initiate the formation of the

cardiovascular lumen.

Interestingly, this analysis also showed that the transformation of the cc into the HT

involves the antero-medial displacement of the splanchnic mesoderm (white arrows in

Figure 4F, F'). This displacement promotes the folding of the cardiac crescent into a

hemi-tube by bringing closer the future borders between the HT and the mesocardium

(see yellow lines in Figure 4F' indicating the distance between the borders at different

times). These borders coincide with the frontiers between cardiomyocytes and

undifferentiated splanchnic mesoderm (see details in Videos 7, 17 and Figure 7A',

A''). Interestingly, the movement of the splanchnic mesoderm is not coherent with

the endoderm but a relative displacement is detected between the two layers (Video 9).

Estimation of splanchnic mesoderm displacement speed towards the midline from

time-lapse analyses (Figure 4F', and Video 6) indicates a range of average speeds

from 12 to 20 μm/hour (15.8 +/-2.4 μm/hour, mean +/- SD, n=3, Figure 4-figure

supplement 2 and Figure 4-source data 1). These measurements estimate that midline

convergence of the splanchnic mesoderm takes approximately 5-7 hours from the late

cc stage until the open HT stage (Figure 4-figure supplement 2).

Our results show the feasibility of live time-lapse analysis of mouse HT formation

and reveal that splanchnic mesoderm displacement, at least in part by sliding over the

endoderm, is an essential aspect of HT morphogenesis.

Live tracking of Cardiomyocyte differentiation in individual cells reveals cardiac

crescent differentiation dynamics.

Next, to specifically track cardiac differentiation, we used the *Nkx2.5eGFP* live

reporter. We first studied the general activation pattern of this reporter in live analysis.

At E7/bud stage, a faint and scattered GFP signal is detected in proximity to the yolk

sack, at the anterior border of the embryo (Figure 5-figure supplement 1A). At neural plate stage, just prior to the ventral folding of the embryo, the GFP signal remains weak but spreads to delineate a crescent in the anterior region of the embryo (Figure 5-figure supplement 1A and Video 10). During about 5-6 hours starting at the EHF stage, the GFP signal increases in intensity, which correlates with the previous observations on cTnnT activation (Figure 4H, Figure 1C and Video 11). From transversal to open HT (5-7 hours) and from the open HT to HT (2-3 hours) the GFP signal remains stable (Figure 5-figure supplement 1B, B', C, C' and Video 12). We conclude that an increase in GFP level in *Nkx2.5eGFP* embryos reports cardiomyocyte differentiation. In addition, these results reveal the timing of the main phases of linear HT development; cc differentiation, formation of the open HT and dorsal closure (Figure 4I).

We next sought to track the trajectories and differentiation of individual cardiac precursors within the entire cardiogenic region by 3D+t live imaging. To this end, we used the *Polr2a^{CreERT2}* (*RERT*) allele (Guerra et al., 2003), which provides ubiquitous tamoxifen-inducible Cre activity in combination with a *Rosa26Rtdtomato* reporter. We then titrated the tamoxifen dose for a labeling density that would allow single cell tracking during prolonged time-lapse analysis and combined this with the *Nkx2.5eGFP* reporter (see materials and methods). Typically, for each video, we acquired z-slices every 3-5μm achieving a total z-depth of 200μm – with some variations depending on the stage considered- and manually tracked in 3D+t for several hours an initial population of ~50 to ~100 cells per video, which represents around 5-10% of the total number of cells present in the cc (de Boer et al., 2012).

We first tracked cells of the cardiac forming region starting at EHF stage –when cardiac precursors are undifferentiated– up to stages in which cardiomyocytes have differentiated in the cc but the transversal HT stage has not been reached yet (Figure 5A and Figure 5-source data 1). At the onset of cardiac differentiation, the cardiac crescent swings ventrally concomitant with foregut pocket formation. During this movement we found that the relative positions of the cardiac progenitors are maintained from the initial stage through the differentiated cc (Video 13). Relative cell positions therefore remain mostly coherent as the embryonic tissues undergo this initial global movement. Differentiation events are detected in some of the tracked cells by cell shape change from columnar to rounded and by the increase in GFP

signal (see example in Figure 5B, D and Figure 5-source data 1). In contrast, other tracked cells located in the splanchnic mesoderm remain in contact with the endoderm, retain a columnar shape and show low GFP level throughout the videos (Figure 5C). These cells are likely to be SHF cardiac progenitors and not endocardial cells, which are not present in this area (Figure 2-figure supplement 2A, B, B'). Endocardial cells are instead present in the cardiac crescent and have typical elongated spindle-like shapes (Figure 4F and Figure 5-figure supplement 3A).

Next, in order to establish a fate map of the cardiac forming region at the EHF stage, we tracked back in time the population of cells that showed high GFP intensity (top 50%) at the end of the video (Figure 5F, H and Figure 5-source data 1). According to our previous analysis (Figure 3D), these cells can reliably be assigned to cardiomyocytes. The initial location of this cell population fated to become cc cardiomyocytes delineates a crescent-shaped domain at EHF stage (Figure 5H). Cells retaining lower GFP intensity level throughout the video initially localize posteriorly and medially to this crescent. Most of the cells that have high GFP levels at the final time point show low GFP levels at the initial time point and increase their GFP level over time (Figure 5E-G and Figure 5-source data 1). These results suggest that cardiomyocytes of the cc differentiate during 5-6 hours starting at the EHF stage, which is consistent with the onset of detectable cTnnT at that stage (Figure 1B and Figure 1-figure supplement 2A, A').

Cells in the cardiac mesoderm do divide during the observation time, so we identified cell division events and tracked the descendant cells. 43% of the tracked cells underwent one division during the 4-5 hour videos. To determine whether cell fate (differentiation versus progenitor) is allocated in the cardiogenic mesoderm at the EHF stage, we tracked GFP levels in dividing cells and their descendants. We found that most sister cell pairs show matched high or low GFP intensity levels at the end of the observation period (38 out of 39, Figure 5I, Figure 5-figure supplement 2A-D and Figure 5-source data 1). This observation suggests that commitment of cardiac precursors to differentiation is already established by the EHF stage and largely transmitted by lineage.

Cardiomyocyte differentiation is not detected during heart tube morphogenesis.

We next studied cardiac differentiation dynamics during subsequent stages when the

cc transforms into the HT by extensive morphogenesis. To do so, we tracked cells

located in the splanchnic mesoderm and cc in *Nkx2.5-eGFP* embryos at successive

periods of around 3 hours covering the 5-7 hours during which the transversal HT

transforms into the open HT (Figure 6A, B, E, Video 14 and Figure 4-source data 1).

This analysis revealed that the trajectories of cc cells move apart from each other over

time as the tissue expands during the transition from the transverse HT to the more

spherical open HT. The HT starts to beat during the observation period, especially at

the later stages, and therefore, in some cases, cardiomyocyte cell shape appears

distorted in single optical sections (Figure 6F), however the GFP level could be

determined. As mentioned above, antero-medial movement of the splanchnic

mesoderm can be observed concomitant with the transformation of the transversal HT

into the open HT (visible also in Video 5 and 12). We found that cells with high GFP

level at the initial time points –differentiated cardiomyocytes– retain rather stable

GFP levels (green tracks in Figure 6C, D, G and Figure 6-source data 1). In addition,

all cells that initially showed low GFP levels did not increase GFP intensity during

time-lapse; thus new events of cardiac differentiation were not detectable by this

approach (red tracks in Figure 6C, D). Tracking cells for longer periods of time,

covering from cc differentiation continuously up to the open HT stage, confirmed that

cc differentiation is followed by a period of time in which no differentiation events

occur (Figure 6- figure supplement1, n=19 cells tracked in one embryo).

To confirm the absence of detectable cardiac differentiation events during this period,

we next focused on the live analysis of cells located at the boundary between

cardiomyocytes and undifferentiated splanchnic mesoderm. As expected, we observed

GFP-low cells located adjacent to GFP-high cells in the boundary zone (Figure 6H

and Video 14). Those cells retain stable GFP levels throughout the tracking time and

did not increase their GFP level (Figure 6I, boundary imaged 20 times in different

locations and in 6 independent embryos). Importantly, they retain a columnar shape

typical of weak cTnnT+ and cTnnT- cells located at the boundary zone (see Figure

2F'' and Figure 3B'). Although these cells migrate antero-medially relative to the

underlying more static endodermal cells (see endodermal cell highlighted by the red

arrow from t=69m in video 9), they do not contribute to the HT during the

observation period. We confirmed this observation in longer time-lapse videos

spanning 7 hours that covered the whole transition from transversal to open HT stage.

Again, progenitors strictly respected the boundary with the HT throughout the entire

time-lapse video (Figure 7A-A''' and Video 15, boundary imaged 5 times in distinct

locations and in 2 independent embryos). All together, these data suggest that during

the transformation of the cc into the dorsally open HT no cardiomyocytes are added to

the HT from the SHF. These observations suggest two distinct phases of early HT

formation: a first phase of differentiation of the FHF into the cc, lasting around 5

hours, and a second phase of HT morphogenesis in which the SHF progenitors remain

undifferentiated, lasting around 5-7 hours. During this second phase, extensive

remodeling of the cardiac crescent is concomitant with the antero-medial splanchnic

mesoderm displacement.

Imaging the Isl1-expressing cell lineage confirms absence of cardiac progenitor

differentiation during early heart tube morphogenesis.

The LIM domain transcription factor Islet1 (Isl1) is a cardiac progenitor marker. Its

expression is transient in the precursors of the cc, while it remains expressed in SHF

progenitors for an extended period (Brade et al., 2007; Prall et al., 2007, Yuan and

Schoenwolf, 2000). Cells of the *Isl1*-expressing lineage detected with Cre reporters

therefore contribute only scarcely to the cc, while extensively to the SHF and its

derivatives (Cai et al., 2003; Ma et al., 2008). To test these observations in live

imaging, we combined *Nkx2.5eGFP* with tracing of the *Isl1* cell lineage using the

Isl1^{cre} driver and the *Rosa26Rtdtomato* reporter. We found that tdtomato labeling is

first detectable in scarce isolated cells of the GFP+ cc during the period when the cc

swings ventrally and differentiates (from t=2h 36m to t=4h in Figure 8A, B). In

contrast, a dense tdtomato labeling appears in the GFP-low cells of the splanchnic

mesoderm as the cardiac crescent fully differentiates (from t=2h24m to t=4h in Figure

8A and C and Video 16). tdtomato is as well detected in the endoderm and

endocardium (not shown). Consistently with previous reports (Cai et al., 2003),

Isl1^{cre}-induced recombination detected in live analysis is thus low in the cc and

complete in the SHF.

Once the cc is formed, if cells of the SHF would continuously differentiate, then

regions of the forming heart tube contributed by the SHF precursors should appear

densely co-labeled with both GFP and tdtomato. Live imaging shows instead that tdtomato+ cells establish a boundary with GFP+ cells, confirming no signs of differentiation of SHF precursors during the observation period (Figure 8A, C and Videos 17). The live imaging, however, did not allow to unambiguously identify all cells located deep inside the live tissue at the final stages recorded. The arterial pole in particular is located deep in the embryo. It is therefore challenging to accurately track cardiac differentiation by the increase of GFP levels there (Figure 8-figure supplement 1A-A'', from around 200μm depth, see next section). To overcome these limitations, we fixed and immunostained embryos against cTnnT after completion of the live-imaging experiments, and imaged them by 3D confocal microscopy. No solid domains containing double-labeled cells were detected, indicating that progenitors located in the SHF did not undergo differentiation in the boundary zone from cc to open HT stage (Figure 8D). These results are consistent with the single-cell tracking analysis and confirm that the SHF does not differentiate during linear HT morphogenesis.

Cardiac differentiation is detected during the late stages of HT development.

We next wanted to determine when cardiac progenitors located in the SHF start to differentiate. In order to address the timing of SHF contribution to the arterial pole, we next fixed and optically cleared *Nkx2.5eGFP*; *Ils1*[cre/+]; *Rosa26Rtdtomato*[+/-] embryos at different stages from cc up to heart looping (n=10) and assessed the appearance of GFP and tdtomato double-positive domains in the HT. In agreement with our previous observations, we found that SHF cells do not differentiate up to the open HT stage, when the dorsal seam of the heart is still open. In contrast, massive appearance of solid domains of double positive cells is observed subsequently in the fully closed HT, reinforcing our previous interpretation (Figure 9A, B and Videos 19-20). At this stage, the primordium of the RV has been added at the arterial pole (Zaffran et al., 2004, Laugwitz et al., 2005, Moretti et al., 2006) and is fully composed of double-positive cells. The dorsal seam of the HT is also densely populated by double-positive cells, indicating a contribution of precursors from the splanchnic mesoderm to the cardiomyocyte population that finalizes the dorsal closure of the linear HT.

In order to capture the initiation of SHF contribution to the HT by live imaging, we

first focused on the venous pole, as this region is more directly exposed than the rest

of the HT and therefore more suitable for live-imaging. In videos that captured the

transition from the open HT to the closed linear HT, some cells at the border between

the FHF and SHF maintain low GFP levels during the first part of the recording and

upregulate GFP to the level of the cardiomyocyte population as the HT forms (Figure

8-figure supplement 2). We next aimed to live-track the activation of SHF

differentiation at the arterial pole. Because of the imaging limitations in this area,

quantitative analysis of the GFP signal was not possible and we instead used the

qualitative detection of SHF cells addition to the HT. To achieve this, we imaged an

Nkx2.5eGFP; *Ils1*[cre/+]; *Rosa26Rtdtomato*[+/-] embryo at successive time points

throughout the transition from open to linear HT and tracked tdtomato+ cells

incorporation to the HT at the arterial pole. This study further confirmed that SHF

differentiation at the arterial pole is initiated when the HT is about to close dorsally

but not before (Figure 9C). Thus, SHF cells do not differentiate during the 5-7 hour

period when morphogenesis of the open HT takes place, but coordinately start

differentiation at different regions of the HT during dorsal closure. Interestingly, these

regions do not only include arterial and venous poles but also the dorsal seam of the

HT.

DISCUSSION

Here we established a whole-embryo live-imaging method based on two-photon

microscopy that allows whole-tissue tracking at cellular resolution. By combining

various genetic tracing tools, we labeled progenitor and differentiated cardiomyocytes

and performed 3D cell tracking over time combined with 3D reconstruction of the HT

at multiple stages. We report three distinct temporal phases of HT formation (Figure

10). During the first phase, the cc differentiates rapidly and morphogenesis, in terms

of changes in the relative position of cells, is minimal. During the second phase,

differentiation is not detected and morphogenetic remodeling gives rise to a dorsally

open HT. During the third phase, cardiac precursor recruitment and differentiation

resumes, contributing to the formation of the RV and the dorsal closure of the HT.

Our results support the early establishment of distinct FHF and SHF cell populations

and show that the morphogenetic changes that transform the cc into a HT largely take
place in the absence of cardiac precursor differentiation. These observations indicate
tissue-level coordination of differentiation and morphogenesis during early
cardiogenesis in the mouse.

The series of 3D reconstructions from fixed embryos was important to establish a
reference staging of HT formation. This allowed us to accurately stage embryos in
live experiments based on morphology and it will also be useful in the future for gene
expression mapping and accurate phenotypic analysis of mutant embryos. The tissue
growth pattern observed in static 3D reconstructions was insightful to suggest
variability in growth rates during different phases of HT formation. Growth of the
differentiated cardiac tissue is relatively paused when the cc undergoes
morphogenesis to form the open HT during the second phase. This is consistent with
previous studies in mouse, chick and human models showing that proliferation drops
in the differentiated myocardium of the forming HT, while proliferation remains high
in the splanchnic mesoderm (Gert van den Berg et al. 2009, de Boer, B.A, 2012,
Sizarov et al. 2010). Our live analysis further showed that SHF cells do not contribute
to the forming HT during the differentiation pause, which correlates with the growth
rate reduction during this phase. This period coincides as well with the onset of
cardiac contractility in the embryo (Tyser et al. 2016).

Following the phase of differentiation pause, growth of the HT is reinitiated by
incorporation of new cells as the HT closes dorsally and the RV precursors are added
at the arterial pole during the third phase. During this third phase, similarities were
found in the differentiation dynamics of SHF precursors and splanchnic precursors
contributing to the dorsal regions of the linear HT. The dorsal aspect of the linear HT
gives rise to the inner curvature of the looped heart, which has an important
contribution to non-chamber myocardium, including atrio-ventricular canal and parts
of the conduction system (Christoffels et al., 2000). Our results suggest that the late
recruitment of progenitors to the dorsal HT could contribute to differences between
inner curvature cardiomyocytes and the rest of the heart tube.

While the live-imaging experiments were essential for the identification of the 5-7
hour hiatus between FHF and SHF differentiation, live imaging of the arterial pole
during the SHF differentiation phase was challenging and was complemented by 3D
reconstructions based on fixed and optically cleared embryos. These experiments

confirmed the pause in differentiation during open HT formation and its reactivation

during linear HT closure. In the future, it will be interesting to explore whether novel

non-toxic index-matching media compatible with embryo viability (Boothe et al.

2017) may alleviate the limitations for deep cardiac imaging during late HT formation.

Regarding the specification of FHF and SHF populations, previous prospective clonal

analyses showed that these lineages diverge around gastrulation (Devine et al., 2014;

Lescroart et al., 2014). In agreement with this, our tracking of cell lineages in the

cardiac forming region shows that sister cells share fates to either the cardiac crescent

or the SHF. In addition, the fact that cells contributing to the SHF do not differentiate

during the period when the cardiac crescent transforms into the primitive heart tube

may contribute to the establishment of the sharp boundary observed between left and

right ventricles later in development (Devine et al., 2014). Further studies will be

required to assess how this temporal pause of cardiac differentiation is regulated. The

molecular analyses of early FHF and SHF precursors suggest that intrinsic molecular

differences between the two lineages appear around or shortly after gastrulation

(Lescroart et al. 2014). These intrinsic differences may contribute to the regulation of

the two distinct differentiation schedules described here. These studies and our

observations, however, cannot discriminate whether this lineage allocation results

from intrinsic differences between these lineages or it is due to their exposure to

position-specific environments, especially as in our studies sister cells remain close

neighbors. Environmental cues thus could also control the sequential differentiation of

FHF and SHF precursors. The Wnt and BMP pathways are well known regulators of

cardiac differentiation (Ai et al., 2007; Jain et al., 2015; Klaus et al., 2007; Kwon et

al., 2007; Marvin et al., 2001; Qyang et al., 2007; Tirosh-Finkel et al., 2010; Ueno et

al., 2007) and specific mechanisms affecting these pathways could be operating

during the formation the HT, whereby the differentiation pathways could be

temporally restrained. Finally, the endoderm may also play a key role in mediating

FHF differentiation. Indeed cardiac mesoderm differentiation is affected in the

absence of Sox17 a transcription factor required for maintenance of the definitive

endoderm- and this is accompanied by the formation of a morphologically abnormal

HT (Pfister et al. 2011). It remains however unclear whether this effect is secondary

to an initial defect in foregut development that is observed in this mutant.

A recent study reported spontaneous calcium transients propagating laterally thought

the cardiac crescent (Tyser et al. 2016). Left/Right (L/R) asymmetry therefore exists within the cardiac crescent, prior to any detectable cardiac contraction, and BMP/SMAD1 signaling in the lateral plate mesoderm may be involved in this asymmetry (Furtado et al. 2008). In our studies, however, we did not detect any L/R difference in cTnnT expression by whole-mount immunodetection or Nkx2.5GFP activation by live analysis in the cc.

Regarding the possible conservation of the temporal differentiation sequence described here, elegant experiments in zebrafish using a cardiac myosin light chain reporter line and a Kaede photo-conversion assay, addressed the temporal order of cardiac differentiation in live embryos (de Pater et al., 2009; Liu and Stainier, 2012). Two distinct phases of cardiomyocyte differentiation were also observed. During a first phase, cardiomyocytes were recruited first into the ventricle and atria at the venous pole. During a second phase, cardiomyocyte differentiation was observed at the arterial pole of the HT. These pulse-chase experiments however do not address whether cardiac differentiation is continuous or includes a differentiation-paused phase, so further studies will be required to establish the conservation of the observations made here for the mouse embryo.

Finally, an important question to address is the functional relevance of the three distinct phases described here; more specifically, what is the role of the observed differentiation pause. An interesting possibility is that this pause would be functionally related to the extensive morphogenetic events that transform the cardiac crescent into the HT. Our study demonstrates coordinated splanchnic mesoderm movements during this phase. These movements involve a very active antero-medial displacement of the splanchnic mesoderm surrounding the cc -mostly fated to the SHF-. This behavior of the splanchnic mesoderm appears essential for transforming the cc into a dorsally closed HT. Importantly, this displacement involves the sliding of the splanchnic layer over the endoderm, suggesting an active role of SHF precursors in this morphogenetic movement. Such displacement of the splanchnic mesoderm over the endoderm during cardiogenesis had been suggested by classical time-lapse studies in the chick embryo (de Haan, 1963) and it is tempting to speculate that a similar phenomenon contributes to the incorporation of SHF cells to both poles of the HT at later stages of HT development (van den Berg et al. 2009; Kelly RG, et al. 2001; Zaffran et al, 2004). These observations suggest that SHF cells do not represent

just a reservoir of cardiac precursors but play a morphogenetic role essential for heart

tube formation. The main consequence of the displacement of the splanchnic

mesoderm layer over the endoderm is the medial convergence of the left and right

frontiers between the cc and SHF to form the dorsal mesocardium and close the HT.

An important consequence of the differentiation pause described here is the stability

of the cc/SHF frontiers during these morphogenetic movements. This stability

prevents further spreading of the differentiation wave from the cc into the

SHF/splanchnic mesoderm, which could interfere the ability of the latter to efficiently

displace over the endoderm. We therefore hypothesize that the stability of the cc/SHF

frontiers –and thus the differentiation pause- would be essential to allow the effective

displacement of SHF/splanchnic mesoderm and elicit HT formation. The temporal

allocation of the morphogenetic phase after cc differentiation allows the formation of

the HT while simultaneously providing an incipient cardiac function essential for the

organization of embryonic circulation. This hypothesis poses a functional basis for the

alternation of differentiation and morphogenesis phases during HT formation.

Our study applies whole-embryo live analysis of cardiac development at tissue level

and with cellular resolution. We expect that extending this experimental approach will

allow to further uncover unexpected and novel mechanisms of organogenesis. While

limited attention had been paid so far to the temporal dynamics of differentiation

during embryonic development, this is an essential aspect of organogenesis

(Gogendeau et al., 2015; Parchem et al., 2015; Yang et al., 2015). Here we show the

relevance of differentiation timing regulation during heart tube formation and its

coordination with morphogenesis at the tissue level. Further understanding of the

molecular and cellular mechanisms underlying these phenomena will help us

expanding pools of cardiac progenitors in vitro or directing them towards

differentiation.

MATERIALS AND METHODS

Mouse Strains:

Mouse alleles used in the manuscript are listed including bibliographic references and

allele   identities   at   the   "Mouse   Genome   Informatics"   data   base   (MGI,

http://www.informatics.jax.org/). *Mesp1^{cre}* (Saga et al., 1999, MGI:2176467), *Isl1^{cre}*

(Cai et al., 2003, MGI:3623159), *Nkx2.5^{cre}* (Stanley et al., 2002, MGI:2448972),

Rosa26Rtdtomato (Madisen et al., 2010, MGI:3809524), *Rosa26RmTmG* (Muzumdar

et al., 2007, MGI:3716464), *Nkx2.5eGFP* (Wu et al., 2006, MGI:5788423),

Polr2a^{CreERT2} (*RERT*) (Guerra et al., 2003, MGI:3772332) and *C57BL/6* (Charles

River). Mice were genotyped as previously described. All animal procedures were

approved by the CNIC Animal Experimentation Ethics Committee, by the

Community of Madrid (Ref. PROEX 220/15) and conformed to EU Directive

2010/63EU and Recommendation 2007/526/EC regarding the protection of animals

used for experimental and other scientific purposes, enforced in Spanish law under

Real Decreto 1201/2005.

Immunostaining and Imaging:

Embryos dissected in Dulbecco's modified Eagle's medium (DMEM, Invitrogen)

were fixed overnight in 2% PFA at 4C, then permeabilized in PBST (PBS containing

0.1% Triton X-100) and blocked (5% goat serum). Embryos were incubated overnight

at 4ºC with antibodies diluted in PBST: mouse anti-cTnnT (1:250, MS-295 Thermo

Scientific), rabbit anti-PH3 (1:250, 06-570 Millipore), CDC31 (553370 BD

Pharmingen™ clone MEC 13.3), SMA (C6198 Sigma) and rabbit anti-Laminin1

(1:500, Sigma, L9393). After washing in freshly prepared PBST at 4ºC, embryos

were incubated with secondary antibodies (Molecular Probes, A21121, A21141,

A11035) coupled to 488, 549 or 649 fluorophores as required at 1:250 and DAPI at

1:500 (Molecular Probes, D3571) overnight at 4ºC. Before imaging, embryos were

washed in PBST at room temperature and cleared with focus clear (Cell Explorer, FC-

101) to enhance the transparency of the embryo. Confocal images were obtained on a

SP8 Leica confocal microscope with a 20X oil objective (0.7 NA) at a 1024 × 1024

pixels dimension with a z-step of 2-4μm. Embryos were systematically imaged

throughout the entire heart tube from top to bottom.

3D Reconstruction and Volumetric Measurement:

For 3D rendering, fluorescent signal in confocal z-stacks was first segmented by

setting intensity thresholds using the trainable Weka segmentation tool plugin

available in Fiji (Arganda-Carreras et al., 2017; Schindelin et al., 2012). The resulting z-stacks were then corrected manually on a slide-by-slide basis to eliminate segmentation mistakes. In case of the cTnnT immunofluorescence images (Figure 1), background signal from the yolk sack was manually masked. The volume of the cTnnT positive myocardium was then computed by multiplying the total segmented area by the z-stack interval using a custom Fiji macro. In the *Nkx2.5$^{cre/+}$; Rosa26tdtomato$^{+/-}$* and *Nkx2.5eGFP* embryos, fluorophore signal present in the endothelium, endocardium and endoderm cells was manually masked prior to segmentation (Figure 1A, Figure 1-figure supplement 1, Figure 2B, Figure 2-figure supplement 2A,A' and, Figure 6A''). For 3D visualization of the 3D segmented image stacks, Imaris software (Bitplane) was used.

Embryo Culture and Two-photon Live-Imaging:

Embryos were dissected at E7.5 in pre-equilibrated DMEM supplemented with 10% foetal bovine serum, 25 mM HEPES-NaOH (pH 7.2), penicillin (50µml21) and streptomycin (50mgml21). Embryos were staged on the basis of morphological criteria (supplementary Figure 1) (Downs and Davies, 1993; Kirstie A. Lawson, 2016), and those between the bud and early somitogenesis stages were used for culture and time-lapse imaging. To track the early phase of cardiac differentiation and subsequent phases of morphogenesis, we used embryos at EHF to transversal HT stage. Embryos were cultured in 50% fresh rat serum, 48% DMEM without phenol red, 1% N-2 neuronal growth supplement (100X, Invitrogen 17502-048) and 1% B-27 supplement (50X Thermo Fisher Scientist 17504044) filter sterilised through a 0.2 mm filter. To hold embryos in position during time-lapse acquisition, we made special plastic holders with holes of different diameters (0.5- 3 mm) to ensure a good fit of embryos similarly to the traps developed by Nonaka et al. (Nonaka, 2009; Nonaka et al., 2002). Embryos were mounted with their anterior side facing up. To avoid evaporation, the medium was covered with mineral oil (Sigma-Aldrich; M8410). Before starting the time-lapse acquisition, embryos were first pre-cultured for at least 2 hours in the microscopy culture set up. The morphology of the embryo was then carefully monitored and if the embryos appeared unhealthy or rotate and move, they were discarded, otherwise, time-lapse acquisition was performed. For the acquisition, we used the Zeiss LSM780 equipped with a 5% CO_2 incubator and a heating chamber

maintaining 37ºC. The objective lens used was a 20X(NA=1) dipping objective,

which allowed a long working distance for imaging mouse embryos and tissues. A

MaiTai laser line at 1000 nm was used for 2-channel 2-photon imaging. Acquisition

was done using Zen software (Zeiss). Typical image settings were: output power:

250mW, pixel dwell time: 7μs, line averaging: 2 and image dimension: 610x610μm

(1024x1024 pixels). To maximize the chance of covering the entire heart tube during

long-term time lapse videos, we allowed 150-200μm of free space between the

objective and the embryo at the beginning of the recording.

Cell labeling, 3D tracking and GFP Intensity Measurement:

For labeling single cells, tamoxifen was administered by oral gavage (2-4 mg/mL) in

RERT;Rosa26R-tdtomato (cell tracking) or *RERT;Rosa26RmTmG* mice (cell shape

study) at E7. Cell shape measurements were done on single cells, imaged in mosaic-

labeled, fixed and immunostained embryos and analyzed with Fiji software (Figure

2D, E). Tracking of tdtomato-labeled cells was done on single cells located within the

cardiogenic mesoderm -excluding endothelial, pericardial and endodermal cells- and

their GFP intensity was measured over time. To track cells manually in 4D stacks, the

MTrackJ Fiji plugin (Meijering et al., 2012) was used. A local square cursor (25x25

pixels) on the cell of interest snaps according to a bright centroid feature on a slice-

by-slice basis. Only tracks lasting for the entire length of the video were kept. When

an ambiguity arises in the tracking between consecutive time points, the track was

discarded. Tracks split at cell divisions. A cell division event is normally clearly

distinguishable over at least 2 time points. In case one of the two daughter cells is not

tractable, the other daughter cell is still tracked. Each track is assigned an ID number

and excel files with all the tracks coordinates in x, y, z and t was generated.

Coordinates of each track were converted into 8-bit 4D images using a custom Fiji

macro in which each cell was represented by a sphere of specific pixel intensity, from

1 to 255, while pixels corresponding to background were set to zero. The 4D images

were then opened with Imaris to perform visualization of the 3D trajectories of each

cell using the "spots" tool, where each object were identified according to pixel

intensity. GFP intensity measurement is performed by segmentation of cell shape. A

Gaussian filter whose radius is adjusted to the typical size of a cell was first applied,

followed by a Laplacian filter. The resulting 32 bits image was next converted to a

mask by thresholding. When objects touched each other, a watershed on the binary mask and manual corrections was applied. Each segmented cell was checked and tracked manually for accuracy. In Figure 3B' nuclei segmentation was performed manually. The mean GFP signal intensity of the segmented objects was then measured using the "analyze particle" tool in Fiji. To quantify the GFP level of tracked cells through time, four to five successive time points were arbitrarily chosen in each video (Figure 5F, Figure 6C, D and Figure 5-figure supplement 1C) except in Figure 5D and Figure 6I, where GFP intensity level was measured in every time point. Background intensities were measured in neural tube cells, which are known to be negative for GFP and cTnnT. Tables containing ID number of tracked cells and GFP intensities were generated and plotted using Prism statistical software.

Statistical Analysis:

For comparisons of two groups, a Man-Whitney U-test was used using Prism statistical software. To find a correlation between GFP and cTnnT levels of 0.8 with an alpha-level f 0.05 and a power of 0.2 at least 10 cells per embryo were required (Figure 3D and Figure 3-figure supplement 2C). Many more cells were computed for each experiment. The linear fit was done using "lm" function from R statistical software (https://www.r-project.org/). To calculate the average speed of splanchnic mesoderm displacement, the shortest distance between the left and right splanchnic mesoderm was calculated in two z-level, using the measure tool in Fiji, and the variation in this distance by time unit was divided by 2, to determine the speed of movement of each sliding side of the splanchnic mesoderm.

Acknowledgments

We thank the CNIC Microscopy unit for help with the live confocal analysis, Fatima Sanchez Cabo for help with the statistical analyses and Florencia Cavodeassi, Miguel Manzanares, Briane Laruy and members of the Torres lab for helpful comments on the manuscript. This work was supported by grants BFU2015-71519-P, BFU2015-70193-REDT and RD16/0011/0019 (ISCIII) from the Spanish Ministry of Economy, Industry and Competitiveness (MEIC). KI was supported by a Human Frontiers Science Program (LT000609/2015) and EMBO (ATL1275-2014) postdoctoral

fellowships. The CNIC is supported by the Spanish MEIC and the Pro CNIC

Foundation, and is a Severo Ochoa Center of Excellence (MINECO award SEV-

2015-0505). The authors declare no conflicts of interest.

REFERENCES:

Abu-Issa, R. (2014). Heart fields: spatial polarity and temporal dynamics. Anat Rec
(Hoboken) 297, 175-182.

Abu-Issa, R., Waldo, K., and Kirby, M.L. (2004). Heart fields: one, two or more? Dev
Biol 272, 281-285.

Ai, D., Fu, X., Wang, J., Lu, M.F., Chen, L., Baldini, A., Klein, W.H., and Martin, J.F.
(2007). Canonical Wnt signaling functions in second heart field to promote right
ventricular growth. Proc Natl Acad Sci U S A 104, 9319-9324.

Arganda-Carreras, I., Kaynig, V., Rueden, C., Eliceiri, K.W., Schindelin, J., Cardona,
A., and Seung, H.S. (2017). Trainable Weka Segmentation: a machine learning tool
for microscopy pixel classification. Bioinformatics.

Biben, C., Weber, R., Kesteven, S., Stanley, E., McDonald, L., Elliott, D.A., Barnett,
L., Koentgen, F., Robb, L., Feneley, M., et al. (2000). Cardiac septal and valvular
dysmorphogenesis in mice heterozygous for mutations in the homeobox gene Nkx2-5.
Circ Res 87, 888-895.

Brade, T., Gessert, S., Kuhl, M., and Pandur, P. (2007). The amphibian second heart
field: Xenopus islet-1 is required for cardiovascular development. Dev Biol 311, 297-
310.

Buckingham, M., Meilhac, S., and Zaffran, S. (2005). Building the mammalian heart
from two sources of myocardial cells. Nat Rev Genet 6, 826-835.

Cai, C.L., Liang, X., Shi, Y., Chu, P.H., Pfaff, S.L., Chen, J., and Evans, S. (2003).
Isl1 identifies a cardiac progenitor population that proliferates prior to differentiation
and contributes a majority of cells to the heart. Dev Cell 5, 877-889.

Chen, C.M., Miranda, A.M., Bub, G., and Srinivas, S. (2014). Detecting cardiac
contractile activity in the early mouse embryo using multiple modalities. Front
Physiol 5, 508.

Christoffels, V.M., Habets, P.E., Franco, D., Campione, M., de Jong, F., Lamers,
W.H., Bao, Z.Z., Palmer, S., Biben, C., Harvey, R.P., et al. (2000). Chamber
formation and morphogenesis in the developing mammalian heart. Dev Biol 223, 266-
278.

Christoffels, V.M., Mommersteeg, M.T., Trowe, M.O., Prall, O.W., de Gier-de Vries,

C., Soufan, A.T., Bussen, M., Schuster-Gossler, K., Harvey, R.P., Moorman, A.F., et
al. (2006). Formation of the venous pole of the heart from an Nkx2-5-negative
precursor population requires Tbx18. Circ Res 98, 1555-1563.

de Boer, B.A., van den Berg, G., de Boer, P.A., Moorman, A.F., and Ruijter, J.M.
(2012). Growth of the developing mouse heart: an interactive qualitative and
quantitative 3D atlas. Dev Biol 368, 203-213.

de Hann RL. (1963) Migration patterns of the precardiac mesoderm in the early chick
embryo. Exp Cell Res.;29:544-60.

de Pater, E., Clijsters, L., Marques, S.R., Lin, Y.F., Garavito-Aguilar, Z.V., Yelon, D.,
and Bakkers, J. (2009). Distinct phases of cardiomyocyte differentiation regulate
growth of the zebrafish heart. Development 136, 1633-1641.

Devine, W.P., Wythe, J.D., George, M., Koshiba-Takeuchi, K., and Bruneau, B.G.
(2014). Early patterning and specification of cardiac progenitors in gastrulating
mesoderm. eLife 2014;3:e03848

Diogo, R., Kelly, R.G., Christiaen, L., Levine, M., Ziermann, J.M., Molnar, J.L.,
Noden, D.M., and Tzahor, E. (2015). A new heart for a new head in vertebrate
cardiopharyngeal evolution. Nature *520*, 466-473.

Downs, K.M., and Davies, T. (1993). Staging of gastrulating mouse embryos by
morphological landmarks in the dissecting microscope. Development 118, 1255-1266.

Evans, S.M., Yelon, D., Conlon, F.L., and Kirby, M.L. (2010). Myocardial lineage
development. Circ Res 107, 1428-1444.

Francou, A., Saint-Michel, E., Mesbah, K., and Kelly, R.G. (2014). TBX1 regulates
epithelial polarity and dynamic basal filopodia in the second heart field. Development
141, 4320-4331.

Furtado MB, Solloway MJ, Jones VJ, Costa MW, Biben C, Wolstein O, Preis JI,
Sparrow DB, Saga Y, Dunwoodie SL, Robertson EJ, Tam PP, Harvey RP. (2008)
BMP/SMAD1 signaling sets a threshold for the left/right pathway in lateral plate
mesoderm and limits availability of SMAD4 Genes Dev. Nov 1;22(21):3037-49.

Galli D, Domínguez JN, Zaffran S, Munk A, Brown NA, Buckingham ME. Atrial
myocardium derives from the posterior region of the second heart field, which
acquires left-right identity as Pitx2c is expressed. Development. 2008
Mar;135(6):1157-67.

Gogendeau, D., Siudeja, K., Gambarotto, D., Pennetier, C., Bardin, A.J., and Basto, R.
(2015). Aneuploidy causes premature differentiation of neural and intestinal stem
cells. Nat Commun 6, 8894.

Guerra, C., Mijimolle, N., Dhawahir, A., Dubus, P., Barradas, M., Serrano, M.,
Campuzano, V., and Barbacid, M. (2003). Tumor induction by an endogenous K-ras
oncogene is highly dependent on cellular context. Cancer Cell 4, 111-120.

Jain, R., Li, D., Gupta, M., Manderfield, L.J., Ifkovits, J.L., Wang, Q., Liu, F., Liu, Y.,
Poleshko, A., Padmanabhan, A., et al. (2015). Integration of Bmp and Wnt signaling
by Hopx specifies commitment of cardiomyoblasts. Science 348, aaa6071.

Kelly, R.G., Brown, N.A., and Buckingham, M.E. (2001). The arterial pole of the
mouse heart forms from Fgf10-expressing cells in pharyngeal mesoderm. Dev Cell 1,
435-440.

Kelly, R.G., Buckingham, M.E., and Moorman, A.F. (2014). Heart fields and cardiac
morphogenesis. Cold Spring Harb Perspect Med 4.

Kirstie A. Lawson, V.W. (2016). A Revised Staging of Mouse Development Before
Organogenesis. Kaufman's Atlas of Mouse Development Supplement.

Klaus, A., Saga, Y., Taketo, M.M., Tzahor, E., and Birchmeier, W. (2007). Distinct
roles of Wnt/beta-catenin and Bmp signaling during early cardiogenesis. Proc Natl
Acad Sci U S A 104, 18531-18536.

Kwon, C., Arnold, J., Hsiao, E.C., Taketo, M.M., Conklin, B.R., and Srivastava, D.
(2007). Canonical Wnt signaling is a positive regulator of mammalian cardiac
progenitors. Proc Natl Acad Sci U S A 104, 10894-10899.

Laugwitz, K.L., Moretti, A., Lam, J., Gruber, P., Chen, Y., Woodard, S., Lin, L.Z.,
Cai, C.L., Lu, M.M., Reth, M., et al. (2005). Postnatal isl1+ cardioblasts enter fully
differentiated cardiomyocyte lineages. Nature 433, 647-653.

Lescroart, F., Chabab, S., Lin, X., Rulands, S., Paulissen, C., Rodolosse, A., Auer, H.,
Achouri, Y., Dubois, C., Bondue, A., et al. (2014). Early lineage restriction in
temporally distinct populations of Mesp1 progenitors during mammalian heart
development. Nat Cell Biol 16, 829-840.

Lescroart, F., Hamou, W., Francou, A., Theveniau-Ruissy, M., Kelly, R.G., and
Buckingham, M. (2015). Clonal analysis reveals a common origin between
nonsomite-derived neck muscles and heart myocardium. Proc Natl Acad Sci U S A
112, 1446-1451.

Lescroart, F., Kelly, R.G., Le Garrec, J.F., Nicolas, J.F., Meilhac, S.M., and
Buckingham, M. (2010). Clonal analysis reveals common lineage relationships
between head muscles and second heart field derivatives in the mouse embryo.
Development *137*, 3269-3279.

Lien, C.L., Wu, C., Mercer, B., Webb, R., Richardson, J.A., and Olson, E.N. (1999).
Control of early cardiac-specific transcription of Nkx2-5 by a GATA-dependent
enhancer. Development 126, 75-84.

Linask, K.K., Knudsen, K.A., and Gui, Y.H. (1997). N-cadherin-catenin interaction:
necessary component of cardiac cell compartmentalization during early vertebrate
heart development. Dev Biol 185, 148-164.

Liu, J., and Stainier, D.Y. (2012). Zebrafish in the study of early cardiac development.
Circ Res 110, 870-874.
849
850 Ma, Q., Zhou, B., and Pu, W.T. (2008). Reassessment of Isl1 and Nkx2-5 cardiac fate
851 maps using a Gata4-based reporter of Cre activity. Dev Biol 323, 98-104.
852
853 Madisen, L., Zwingman, T.A., Sunkin, S.M., Oh, S.W., Zariwala, H.A., Gu, H., Ng,
854 L.L., Palmiter, R.D., Hawrylycz, M.J., Jones, A.R., et al. (2010). A robust and high-
855 throughput Cre reporting and characterization system for the whole mouse brain. Nat
856 Neurosci 13, 133-140.
857
858 Marvin, M.J., Di Rocco, G., Gardiner, A., Bush, S.M., and Lassar, A.B. (2001).
859 Inhibition of Wnt activity induces heart formation from posterior mesoderm. Genes
860 Dev 15, 316-327.
861
862 Meijering, E., Dzyubachyk, O., and Smal, I. (2012). Methods for cell and particle
863 tracking. Methods Enzymol 504, 183-200.
864
865 Meilhac, S.M., Esner, M., Kelly, R.G., Nicolas, J.F., and Buckingham, M.E. (2004a).
866 The clonal origin of myocardial cells in different regions of the embryonic mouse
867 heart. Dev Cell 6, 685-698.
868
869 Meilhac, S.M., Esner, M., Kerszberg, M., Moss, J.E., and Buckingham, M.E. (2004b).
870 Oriented clonal cell growth in the developing mouse myocardium underlies cardiac
871 morphogenesis. J Cell Biol 164, 97-109.
872
873 Mjaatvedt, C.H., Nakaoka, T., Moreno-Rodriguez, R., Norris, R.A., Kern, M.J.,
874 Eisenberg, C.A., Turner, D., and Markwald, R.R. (2001). The outflow tract of the
875 heart is recruited from a novel heart-forming field. Dev Biol 238, 97-109.
876
877 Moorman, A.F., Christoffels, V.M., Anderson, R.H., and van den Hoff, M.J. (2007).
878 The heart-forming fields: one or multiple? Philos Trans R Soc Lond B Biol Sci 362,
879 1257-1265.
880
881 Moretti, A., Caron, L., Nakano, A., Lam, J.T., Bernshausen, A., Chen, Y., Qyang,
882 Y.,Bu, L., Sasaki, M., Martin-Puig, S., et al. (2006). Multipotent embryonic isl1+
883 progenitor cells lead to cardiac, smooth muscle, and endothelial cell diversification.
884 Cell 127, 1151-1165.
885
886 Muzumdar, M.D., Tasic, B., Miyamichi, K., Li, L., and Luo, L. (2007). A global
887 double-fluorescent Cre reporter mouse. Genesis 45, 593-605.
888
889 Nonaka, S. (2009). Modification of mouse nodal flow by applying artificial flow.
890 Methods Cell Biol 91, 287-297.
891
892 Nonaka, S., Shiratori, H., Saijoh, Y., and Hamada, H. (2002). Determination of left-
893 right patterning of the mouse embryo by artificial nodal flow. Nature 418, 96-99.
894
895 Parchem, R.J., Moore, N., Fish, J.L., Parchem, J.G., Braga, T.T., Shenoy, A., Oldham,
896 M.C., Rubenstein, J.L., Schneider, R.A., and Blelloch, R. (2015). miR-302 Is

897 Required for Timing of Neural Differentiation, Neural Tube Closure, and Embryonic
898 Viability. Cell Rep 12, 760-773.
899
900 Pfister S1, Jones VJ, Power M, Truisi GL, Khoo PL, Steiner KA, Kanai-Azuma M,
901 Kanai Y, Tam PP, Loebel DA. (2011) Sox17-dependent gene expression and early
902 heart and gut development in Sox17-deficient mouse embryos. Int J Dev
903 Biol.;55(1):45-58
904
905 Prall, O.W., Menon, M.K., Solloway, M.J., Watanabe, Y., Zaffran, S., Bajolle, F.,
906 Biben, C., McBride, J.J., Robertson, B.R., Chaulet, H., et al. (2007). An Nkx2-
907 5/Bmp2/Smad1 negative feedback loop controls heart progenitor specification and
908 proliferation. Cell 128, 947-959.
909
910 Qyang, Y., Martin-Puig, S., Chiravuri, M., Chen, S., Xu, H., Bu, L., Jiang, X., Lin, L.,
911 Granger, A., Moretti, A., et al. (2007). The renewal and differentiation of Isl1+
912 cardiovascular progenitors are controlled by a Wnt/beta-catenin pathway. Cell Stem
913 Cell 1, 165-179.
914
915 Ramsbottom, S.A., Sharma, V., Rhee, H.J., Eley, L., Phillips, H.M., Rigby, H.F.,
916 Dean, C., Chaudhry, B., and Henderson, D.J. (2014). Vangl2-regulated polarisation of
917 second heart field-derived cells is required for outflow tract lengthening during
918 cardiac development. PLoS Genet 10, e1004871.
919
920 Saga, Y., Hata, N., Kobayashi, S., Magnuson, T., Seldin, M.F., and Taketo, M.M.
921 (1996). MesP1: a novel basic helix-loop-helix protein expressed in the nascent
922 mesodermal cells during mouse gastrulation. Development 122, 2769-2778.
923
924 Saga, Y., Miyagawa-Tomita, S., Takagi, A., Kitajima, S., Miyazaki, J., and Inoue, T.
925 (1999). MesP1 is expressed in the heart precursor cells and required for the formation
926 of a single heart tube. Development 126, 3437-3447.
927
928 Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
929 Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source
930 platform for biological-image analysis. Nat Methods 9, 676-682.
931
Sinha, T., Wang, B., Evans, S., Wynshaw-Boris, A., and Wang, J. (2012). Disheveled
mediated planar cell polarity signaling is required in the second heart field lineage for
outflow tract morphogenesis. Dev Biol 370, 135-144.
Formation of the building plan of the human heart: morphogenesis, growth, and
differentiation. (2011) Sizarov A, Ya J, de Boer BA, Lamers WH, Christoffels VM,
Moorman AF. Circulation.;123(10):1125-35.
938
939 Spater, D., Abramczuk, M.K., Buac, K., Zangi, L., Stachel, M.W., Clarke, J., Sahara,
940 M., Ludwig, A., and Chien, K.R. (2013). A HCN4+ cardiomyogenic progenitor
941 derived from the first heart field and human pluripotent stem cells. Nat Cell Biol 15,
942 1098-1106.
943
Stanley, E.G., Biben, C., Elefanty, A., Barnett, L., Koentgen, F., Robb, L., and
Harvey, R.P. (2002). Efficient Cre-mediated deletion in cardiac progenitor cells
conferred by a 3'UTR-ires-Cre allele of the homeobox gene Nkx2-5. Int J Dev Biol 46,

431-439.

Tirosh-Finkel, L., Zeisel, A., Brodt-Ivenshitz, M., Shamai, A., Yao, Z., Seger, R.,
Domany, E., and Tzahor, E. (2010). BMP-mediated inhibition of FGF signaling
promotes cardiomyocyte differentiation of anterior heart field progenitors.
Development 137, 2989-3000.

Tyser, R.C., Miranda, A.M., Chen, C.M., Davidson, S.M., Srinivas, S., and Riley, P.R.
(2016). Calcium handling precedes cardiac differentiation to initiate the first heartbeat.
Elife 5.

Ueno, S., Weidinger, G., Osugi, T., Kohn, A.D., Golob, J.L., Pabon, L., Reinecke, H.,
Moon, R.T., and Murry, C.E. (2007). Biphasic role for Wnt/beta-catenin signaling in
cardiac specification in zebrafish and embryonic stem cells. Proc Natl Acad Sci U S A
104, 9685-9690.

van den Berg G, Abu-Issa R, de Boer BA, Hutson MR, de Boer PA, Soufan AT,
Ruijter JM, Kirby ML, van den Hoff MJ, Moorman AF. (2009) A caudal proliferating
growth center contributes to both poles of the forming heart tube. Circ Res.
104(2):179-88.

Waldo, K.L., Kumiski, D.H., Wallis, K.T., Stadt, H.A., Hutson, M.R., Platt, D.H., and
Kirby, M.L. (2001). Conotruncal myocardium arises from a secondary heart field.
Development 128, 3179-3188.

Wu, S.M., Fujiwara, Y., Cibulsky, S.M., Clapham, D.E., Lien, C.L., Schultheiss, T.M.,
and Orkin, S.H. (2006). Developmental origin of a bipotential myocardial and smooth
muscle cell precursor in the mammalian heart. Cell 127, 1137-1150.

Yang, S.L., Yang, M., Herrlinger, S., Liang, C., Lai, F., and Chen, J.F. (2015). MiR-
302/367 regulate neural progenitor proliferation, differentiation timing, and survival
in neurulation. Dev Biol 408, 140-150.

Yuan, S., and Schoenwolf, G.C. (2000). Islet-1 marks the early heart rudiments and is
asymmetrically expressed during early rotation of the foregut in the chick embryo.
Anat Rec 260, 204-207.

Zaffran, S., Kelly, R.G., Meilhac, S.M., Buckingham, M.E., and Brown, N.A. (2004).
Right ventricular myocardium derives from the anterior heart field. Circ Res 95, 261-
268.

 FIGURE LEGENDS:

Figure 1. Overview of HT morphogenesis and growth.

(A) Frontal view of an *Nkx2.5cre/+*; *Rosa26Rtdtomato* embryo at EHF stage. (A') 3D

reconstruction of the tdtomato signal in the cardiogenic area. Signal from tdtomato+

endothelial cells identified by shape was manually masked. See also Video 1. (B-G)

Immunostaining for cTnnT (red) and Dapi (blue) showing six consecutive stages

during cardiac differentiation (B-D) and HT morphogenesis (E-G). (B) At EHF cTnnt

is initially not detectable. (C-D) During early somitogenesis, cTnnT signal becomes

detectable in the cc. Insets in (B-D): magnification of single optical sections showing

cTnnT localization and cell shape. (C'-G' and E''-G'') Corresponding 3D renderings

from cTnnT signal reconstruction. Red arrows in (E'-G') highlight the dorsal closure

of the HT. Yellow arrow in G'' highlights the arterial pole (prospective RV). See also

Video 2. (H) Quantification of the arterial pole/RV length in the open HT (41.4

±14.0µm, n=5) and after dorsal closure (109 ±43.44µm, n=7), mean ± SD, p=0.0025.

(I) Quantification of the cardiac volume at the different stages of HT development.

(Initial cc: 1.63.106µm3 ± 0.13, n=4, cc: 2.89.106 ± 0.37µm3, n=3, transversal HT:

3.367. 106µm3 ± 0.95, n=5, open HT: 4.29.106µm3 ± 1.08, n=6, linear HT: 6.37.

106µm3 ± 1.01, n=5, mean ± SD). P-values are indicated on the graph. (J)

Immunostaining of an *Nkx2.5eGFP* embryo for PH3 (red) and Dapi (blue) at HT

stage, showing proliferative cells in the ventricle. Scale bars: 100µm

1009

Figure 1-figure supplement 1. Criteria for embryo staging

We determined the developmental stage of embryos based on morphological

landmarks (Downs and Davies, 1993; Kirstie A. Lawson, 2016). (A) At bud stage/E7,

a small allantois bud is visible. At neural plate stage/E7.5, the allantois is larger and

projects into the yolk sack cavity. The anterior neuroectoderm is enlarged. (B) At

EHF/E7.5, the neural plate starts to form the head fold. During somitogenesis, a

clearly visible foregut pocket appears and the head folds are located dorsally and

anteriorly to the cardiac primordium. The cc has differentiated. It transforms

successively into the transversal HT, the open HT and the linear HT closed dorsally

and with a prominent arterial pole/RV. Scale bars: 100µm.

1020

1021 **Figure 1-figure supplement 2. Lineage tracing using the *Nkx2.5cre* driver shows a**

1022	**contribution of labeled cells to the cardiogenic regions but not to the endoderm**
1023	**at EHF stage**
1024	(A, A') *Nkx2.5$^{cre/+}$; Rosa26Rtdtomato$^{+/-}$* embryo at EHF stage (same embryo as in
1025	Figure 1A. Here a single optical section is shown). Tdtomato labeling is detected in
1026	the cardiac progenitors, endothelial (yellow arrow) and endocardial cells (green
1027	arrows). No labeling is detected in the endoderm (gray arrow). (A'') Segmentation of
1028	cardiac crescent progenitors (shown in dark green) used for the 3D reconstruction
1029	shown in Figure 1A' overlaid with the raw image and. Scale bars: 100μm
1030	
1031	**Figure 1-figure supplement 3. Faint cTnnT signal starts to be detected at EHF**
1032	**stage in apico-basally polarized cc cells.**
1033	(A) A *Mesp1$^{cre/+}$; Rosa26Rtdtomato$^{+/-}$* embryo at EHF stage showing mesodermal
1034	labeling (green)- immunostaining against cTnnT (red) and Dapi (blue). Note that the
1035	embryo is at a slightly more advanced developmental stage than those shown in
1036	(Figure 1B) because the foregut pocket is more invaginated. (A') White inset in (A).
1037	Faint cTnnT signal (red) in few mesodermal cells (arrows) can be detected. (B-B')
1038	*Mesp1$^{cre/+}$; Rosa26Rtdtomato$^{+/-}$* embryo showing the mesoderm in red and
1039	immunostained against the tight junction component zona-occludens-1 (ZO-1) (green)
1040	and Dapi (blue). The cc cells (as seen in transversal sections) have an AB polarized
1041	epithelial morphology. fg: foregut, endo: endoderm, endoc: endocardium. Scale bars:
1042	150μm.
1043	
1044	Figure Source Data 1
1045	Source data for Figure 1H and I
1046	
1047	**Figure 2. Visualization of the boundary between FHF and SHF.**
1048	(A-A'') An *Nkx2.5$^{cre/+}$; Rosa26Rtdtomato$^{+/-}$* embryo immunostained for cTnnT
1049	(green) and Dapi (blue) showing cells of the Nkx2.5+ lineage populating both the
1050	FHF and SHF. (A' and A'') Cross-sections in xz along the dotted lines 1 and 2 shown
1051	in (A), and corresponding schematics highlighting the endoderm (blue), FHF (green),
1052	SHF (red) and endocardium (yellow). Note that tdtomato signal is also detected in the
1053	endoderm and endocardium. (B) 3D reconstruction of the FHF (green) and SHF (red);
1054	(B') 3D drawing of the border between the FHF and SHF (in red) and the FHF and

yolk sack (in green) (based on the embryo shown in (A). For SHF rendering, the

tdtomato+ splanchnic mesoderm was depicted. The FHF was rendered using the

cTnnT signal. See also Video 3. (C) Quantification of the cell roundness (rnd) index

of cardiomyocytes at HT stage on single optical sections. Black bars indicate mean.

Rnd index for cTnnT+ (green) cells is 0.71 ± 0.16 and for cTnnT- (red) cells is $0.38 \pm$

0.16, mean $\pm$ SD, exact p value < 0.0001. (D-E) Membrane-GFP labeling of typical

cTnnT+ FHF (D) and cTnnT-SHF (E) cells at longitudinal HT stage. The specimen is

immunostained for cTnnT (blue), the basement membrane marker Laminin1 (red) and

GFP (green). (F) Single optical section of the same embryo shown in (A, B). (F'-F''')

Inset: red arrows point to cells localizing weak cTnnT signal and have columnar cell

shape in the SHF. White arrows point to cTnnT-negative SHF cells. splan. meso:

splanchnic mesoderm, endo: endoderm, fg: foregut. In all the embryos immunostained

for cTnnT, the yolk sack signal is non-specific background (indicated by an asterisks).

Scale bars: 100μm except in (D-E): 50μm

1069

Figure 2- figure supplement 1. Image segmentation of the FHF and SHF.

(A) Single optical section of an *Nkx2.5$^{cre/+}$; Rosa26Rtdtomato$^{+/-}$* embryo

immunostained for cTnnT (green) and Dapi (blue). Gray arrows highlight tdtomato

signal in the endoderm (A') Segmentation of the FHF and SHF. Segmentation of the

FHF (green) is based on the cTnnT signal while the segmentation of the SHF is based

on the tdtomato positive and cTnnT negative signal (in red). tdtomato labeling in the

endoderm and endocardium has been removed manually.

1077

1078   **Figure 2- figure supplement 2. Endocardium localization in the cardiac crescent.**

1079   (A) Maximum z-projection and (B, B') single optical section of an embryo

1080   immunostained for smooth muscle actin (SMA, labeling differentiated

1081   cardiomyocytes) (red), CD31 (labeling the endocardium) (green) and ZO-1 (blue).

1082   endo: endoderm, splan. meso: splanchnic mesoderm Scale bars: 150μm.

1083

1084   Figure Source Data 2

1085   Source data for Figure 2C

1086

Figure 3. The *Nkx2.5eGFP* transgene is robustly expressed in cTnnT-positive cardiomyocytes

(A) z-maximum projection of an *Nkx2.5eGFP* embryo at transversal HT stage immunostained for cTnnT (red) and Dapi (blue) showing high GFP level in differentiated cardiomyocytes. (B) Single optical section of an *Nkx2.5eGFP* embryo at transversal HT stage immunostained for cTnnT (red) and Dapi (blue). (B') Inset in (B): arrow points the transition between cTnnT+ and cTnnT- domains, corresponding to FHF and SHF, respectively. Dotted line represents the linescan measured in (D) (C) Example of manual segmentation based on Dapi nuclei in cells located in the splanchnic mesoderm and in the neural tube. (D) Linear mixed-effects model to find the relationship between the background-substracted GFP and cTnnT levels adjusted by embryo (GFP=0.95*cTnnT). (E) GFP and cTnnT mean intensities measured within manually segmented cells along the boundary from SHF to cTnnT-positive FHF (B') (n=130 cells analysed in 4 independent embryos). Scale bar: 100µm

Figure 3- figure supplement 1. The *Nkx2.5eGFP* and cTnnT labeling at different z-levels.

(A) z-maximum projection and yz view of a *Nkx2.5eGFP* embryo immunostained against cTnnT showing overlap between eGFP signal and cTnnT localisation along the z dimension. (B) z-maximum projection and optical sections at different z-level of an *Nkx2.5eGFP* embryo, immunostained for cTnnT (red) and Dapi (blue), showing overlapping expression of GFP and cTnnT signal. Scale bar: 100µm

Figure 3- figure supplement 2. High GFP levels are detected in strongly labeled cTnnT cells.

(A,B) Normalized GFP and cTnnT mean level for single segmented cell. Each dot represents a single segmented cell (A) located in the FHF and SHF and (B) located at the boundary zone between the FHF and SHF (n=130 cells analyzed from 8 embryos). (C) Linear mixed-effects model to find the relationship between the background substracted GFP and cTnnT levels adjusted by embryo for cell located at the boundary between the FHF and SHF (GFP=0.94*cTnnT, R^2=0.77, $p<2.2e-16$). Cells are being considered positive for cTnnT when their mean intensity value is above 0. Note that small GFP signal can be detected in the cTnnT-negative SHF cells.

1120

Figure 3- figure supplement 3. *Nkx2.5Cre* genetic tracing labels both the FHF and SHF

(**A**) The GFP expression of the *Nkx2.5eGFP* reporter does not fully overlap with tdtomato expression pattern in *Nkx2.5$^{cre/+}$; Rosa26Rtdtomato$^{+/-}$; Nkx2.5eGFP* embryos at the level of the splanchnic mesoderm/SHF. Scale bar: 100μm.

Figure Source Data 3

Source data for Figure 3D and E

Figure 4. Live-imaging of cardiac differentiation and morphogenesis

(A) Schematic of the set up for imaging live mouse embryos from EHF up to completion of HT formation. (B) Parameters xyzt during live-imaging. (C) An embryonic holder maintains the embryo still during live-imaging. (D-E) After over 10 hours of ex-vivo culture inside the two-photon chamber, the embryo has grown and the cardiac crescent has transformed into a beating heart tube. Arrows in (E) point to the deformation of the heart ventricle due to heart beats. See also Video 4. (F-F'') Time-lapse video sequence of a *Mesp1$^{cre/+}$; Rosa26Rtdtomato$^{+/-}$* embryo –reporting anterior mesoderm–, showing the transition from cardiac crescent stage to heart tube stage. Note that at the initial time point, the foregut pocket is already visible. Arrows indicate the major tissue movements visible (folding of the cardiac crescent, medial movement of the splanchnic mesoderm). The differentiation of cardiomyocytes detaching from the endoderm is visible in (F'). White arrows point to the endocardium. The formation of the endocardial lumen in the transversal HT is visible in F'. By 8h35m, the open HT is fully formed and beating regularly. Yellow lines represent distances between the left and right splanchnic mesoderm. Images are z-max projection of 84 sections covering 180μm (F) and 9 sections covering 36μm (F') acquired every 4μm. See also Videos 5-6. (G-G'') Time-lapse video of an *Nkx2.5$^{cre/+}$; Rosa26RmT/mG$^{+/-}$* embryo during the early stages of cardiac differentiation. Images are z-max projection of 87 sections acquired every 5μm covering 437μm. (G') shows in a single optical section how progenitors change in cell shape and move away from the endoderm during differentiation towards cardiomyocytes (from inset in (G)). (G'')

Cartoon depicting the change in cell shape taking place during cardiac differentiation. See also Video 7. (H) Time-lapse video sequences of *Nkx2.5eGFP* embryos from transversal HT showing low GFP level prior to cardiomyocyte differentiation and increase in GFP intensity level during the stages cardiomyocytes undergo differentiation. (H' and H''), cardiomyocytes in single magnified optical sections. Images are z-maximum projections of 70 sections acquired every 6μm covering 420μm. See also Videos 10. (I) Estimate of the timing between different heart tube development stages. fg: foregut cc: cardiac crescent, endoc: endocardium, ao: aorta, endo: endoderm. Scale bars: 100μm.

Figure 4- figure supplement 1. Embryos can be cultured and imaged under the 2-photon microscope for up to 24 hours

(A) *Nkx2.5eGFP*; *RERT*$^{+/-}$; *Rosa26Rtdtomato*$^{+/-}$ embryo. Starting culture at E7.5, by the end of the 25 hours ex-vivo culture, the HT has formed and looped (experiment repeated 3 times independently). The embryo was subsequently fixed and immunostained for cTnnT (red) and ZO-1 (blue). V: prospective left ventricle, A: prospective atria. Scale bars: 100μm.

Figure 4-figure supplement 2. Quantification of the displacement of the splanchnic mesoderm during HT formation.

(A) z-max projection of a *Mesp1*$^{cre/+}$*; Rosa26Rtdtomato*$^{+/-}$ embryo at transversal HT stage. The yellow line indicates the distance between the left and right splanchnic mesoderm. (B) One example of the changes in left/right distance measured over time.

Figure Source Data 4

Source data for Figure Figure 4-figure supplement 2

Figure 5. Live-imaging of cardiac differentiation at cellular resolution

(A) Time course of an *Nkx2.5eGFP*; *RERT*$^{cre/+}$; *Rosa26Rtdtomato*$^{+/-}$ embryo during stages when cc differentiation takes place -from EHF onwards-. Images are z-maximum projection of 76 sections acquired every 3μm covering 228μm. A bright-

field image of the EHF embryo at the initial time point is also shown. (i, ii and iii) Rationale of the expected evolution of GFP expression: since tracks carry information on reporter expression, if a cell acquires a high level of GFP, we predict that it is committed to differentiate. See also Video 12. (B-D) Examples of time-lapse videos (B and C) and quantification of the GFP level (D) increasing in a single cell during differentiation (B) or remaining low in a cell located in the splanchnic mesoderm (C). A neural tube cell was additionally quantified (blue). Images are single optical sections. (E) Time course of individual cells tracked in the video shown in A. Cells with GFP levels above the median intensity value of all cells at the end of the recording are represented as green spheres and cells with lower GFP levels are shown as red spheres. (F) GFP level through time. Blue dotted line: median value (5249 a.u.). GFP level for each tracked cell was measured at five successive time points. Cell divisions are not represented for simplicity. (G) Normalized GFP level showing the relative increase in GFP levels for the GFP-positive cells at the end of the recording. Average increase: 2,77 fold ± 1.50, mean ± SD, n=95 cells from 3 independent videos. (H) The position of cells that differentiate (green) and cells that do not is shown in the cc at the initial time point (EHF stage). (I) The lineage from cardiac precursors share differentiation fate. Lineages of dividing progenitors (n=55 from 3 independent videos) were identified during early stages of cardiac differentiation. Two daughter cells are defined as sharing the same fate if their GFP intensity levels do not differ by more than 1.5 fold and/or both show levels above or below the threshold value defined. Scale bars: 100μm.

Figure 5-figure supplement 1. Live-imaging of *Nkx2.5eGFP* reporter line

(A-C) Time-lapse video sequences of *Nkx2.5eGFP* embryos from Early Bud (A), cc (B) and open HT stage (C). Images are z-maximum projections of 75 sections acquired every 4μm covering 300μm in (A), 55 sections acquired every 6μm covering 300μm in (B) and 59 sections acquired every 5μm covering 295μm in (C). (B' and C'): z-max projection covering 132μm (B') and 60μm (C') showing the dorsal opening and the dorsal closure of the HT at the end of the videos (representative of 3 embryos). See also Videos 9 and 11. Scale bars: 100μm.

Figure 5- figure supplement 2. Cells divide during cardiac differentiation

1219 (A-C) Time-lapse video of a dividing tdtomato+ cell. Images are single optical

1220 sections (same embryo as shown in Figure 4A). The dividing tdtomato+ cell and

1221 daughter cells are segmented and mean GFP level is measured. (D) Full 4D track of

1222 the dividing cell showing close localization of the progeny in the cardiac crescent.

1223 Scale bars: 100μm.

1224

1225 **Figure 5- figure supplement 3.** Endocardial cells show an elongated shape. (A)

1226 Single optical section of a tamoxifen-induced *Nkx2.5eGFP*; *RERT*[+/-];

1227 *Rosa26Rtdtomato*[+/-] embryo showing a single endocardial labeled cell (white arrow).

1228 Scale bars: 100μm.

1229

1230 Figure Source Data 5

1231 Source data for Figure 5A, D, F, G, I and Figure 5-figure supplement 1C, D

1232

Figure 6. No cardiac differentiation is detected during early HT morphogenesis

(A-B) Initial and final time points of two time-lapse videos of tamoxifen-induced

Nkx2.5eGFP; *RERT*[+/-]; *Rosa26Rtdtomato*[+/-] embryos covering the transformation of

the transversal HT into the open HT. Images are z-maximum projections of 44

sections acquired every 5μm covering 220μm in (A) and of 46 sections acquired

every 6μm covering 276μm in (B). See also Video 13. (C, D) GFP levels over time of

the cells tracked in (A, B). GFP level for each tracked cell was measured at four

successive time points. Green tracks represent cardiomyocytes with GFP intensity

above the median intensity value of all the tracked cells at the last time point, while

red tracks represent cells with lower GFP level. Blue dotted line represents the

median intensity value (10748 a.u. in (C) and 12248a.u. in (D)). Note that when cells

divide, only one of the two daughter cells was represented for simplicity. (E)

Distribution of the tracked cells from the time-lapse video shown in (A) at the initial

and final times of the tracking period. tdtomato+, GFP- cells are represented by red

spheres and tdtomato+, GFP+ cells are represented by green spheres. (F) Example of

tdtomato+ cardiomyocytes tracked in the beating ventricle. (G) Normalized

progression of GFP level in cells classified as GFP+ showing stable GFP levels. Data

collected from 125 tracked cells from four independent videos of 2h52m to 3h12m

duration. Average increase: 1,1 fold ± 0.28, mean ± SD. (H) Magnification extracted

from the video in A (yellow inset) of tdtomato+ cells in the boundary zone between

the undifferentiated and differentiated cells. (I) Quantification of GFP level through

time of the three segmented cells shown in (H). Cells show stable GFP levels and do

not differentiate. Images are z-maximum projections of 6 sections acquired every

5μm covering 30μm. Scale bars: 100μm.

1257

Figure 6- figure supplement 1.

Longer videos can capture both initial cardiac differentiation in the FHF and early HT

morphogenesis. (A) Initial and final images from a time-lapse video of a tamoxifen-

induced *RERT^{+/-}; Rosa26Rtdtomato^{+/-}; Nkx2.5eGFP* embryo. Images are z-maximum

projection of 44 sections acquired every 5μm covering 220μm. (B) GFP levels over

time of the cells tracked in (A). GFP level for each tracked cell was measured at five

successive time points. Green tracks represent cells with GFP intensity above the

median intensity value of all the tracked cells at the last time point, while red tracks

represent cells that maintain lower GFP levels. The blue dotted line represents the

median intensity value. (C) Mean intensity value for all the green and red tracks and

SD. Note that on average the GFP intensity of the green tracks becomes stable from

around the transversal HT stage (around t=300m). Scale bars: 100μm.

1270

1271   Figure Source Data 6

1272   Source data for Figure 6A, B, C, D, G, I, Figure 6-figure supplement 1A, Figure 6-

1273   figure supplement 1B and C.

1274

Figure 7. No cardiac differentiation is detected during early HT morphogenesis.

(A-A'') Time-lapse video of a tamoxifen-induced *RERT^{+/-}; Rosa26Rtdtomato^{+/-};*

Nkx2.5eGFP embryo during the stages at which the transversal HT transforms into an

open HT, showing cells located in the splanchnic mesoderm (white arrows in A' and

A''), respecting the boundary with the HT (yellow arrows in J' show two cells located

in the HT). Lower doses of tamoxifen were injected in order to label only a very small

proportion of cells in red. Images are z-maximum projection of 74 sections (A)

acquired every 4μm covering 296μm and (A') 7 sections covering 28μm. (A'') 3D

reconstruction based of the *Nkx2.5eGFP* signal (green) and tdtomato+ cells located in

the splanchnic mesoderm. See also Video 14. Scale bars: 100μm. Splan. meso:

splanchnic mesoderm.

Figure 8. SHF Isl1-expressing cells do not contribute to the early open-heart tube.
(A) Time-lapse video sequences of an *Isl1*$^{cre/+}$*; Rosa26Rtdtomato*$^{+/-}$*; Nkx2.5eGFP* embryo showing that recombination driven by *Islet1*^{cre} of the *Rosa26* locus is complete in the splanchnic mesoderm and scarce in the FHF (n=3 videos from cc up to open HT stage). (B) Inset from A (yellow frame). Two cells of the Isl1+ lineage located in the cc increase their tdtomato level over time. (C) Inset from A (Red frame); increase of the tdtomato intensity in the splanchnic mesoderm over time. Arrow shows the boundary between SHF and FHF and points to a single cell in the SHF that retains a low level of GFP and a columnar shape. Note that tdtomato signal is also detectable in the endoderm. Images are z-maximum projection of 45 sections acquired every 5μm and covering 225μm in (A) and of 6 sections covering 30μm in (C). See also Videos 15-16. (D) Same embryo as in (A) post-fixed and immunostained against cTnnT after live-imaging, showing that the red cells located in the splanchnic mesoderm are undifferentiated. Scale bars: 100μm.

Figure 8- figure supplement 1. GFP level in deeper z levels cannot be accurately quantified.
(A-A'') *RERT*$^{+/-}$*; Rosa26Rtdtomato*$^{+/-}$*; Nkx2.5eGFP* embryo at transversal HT stage. (A) z-maximum projection of 74 sections acquired every 4μm and covering 296μm. Same embryo as in Figure 6A. (A') Single optical z-section positioned at the white dotted line shown in (A). This z-section shows the green signal decay at deep z positions (z= 0 to -274μm). (A'') An xy frame corresponding to a *-272μm* z-position (indicated by the red dotted line in (A') at different time points during the video. The white arrow indicates td-tomato+ cells.

Figure 8- figure supplement 2. Cells increasing their GPF expression can be detected at the venous pole during activation of SHF contribution to the HT.
(A) Time course of a tamoxifen-induced *RERT*$^{+/-}$*; Rosa26Rtdtomato*$^{+/-}$*; Nkx2.5eGFP* embryo during linear HT formation. Images are z-maximum projection of 44 sections acquired every 5μm covering 220μm. (B) Time course of individual cells tracked within the yellow-framed region in video (A). (C) GFP levels over time of the cells

1318 tracked in (B) Cell 1 shows high GFP level throughout the video while cell 2 and 3

1319 increase their GFP level over time. The blue line represents a neural tube cell. The

1320 dotted line marks the 1h30m time point. Scale bars100μm.

1321

1322 Figure Source Data 8

1323 Source data for Figure 8-figure supplement 2C.

1324

1325

Figure 9. SHF Isl1-expressing cells contribute to the arterial pole during late

linear HT morphogenesis.

(A, B) *Isl1$^{cre/+}$; Rosa26Rtdtomato$^{+/-}$; Nkx2.5eGFP* embryos showing no contribution

of the Isl1 lineage to the transversal HT (A-A'') but contributing robustly to the

arterial pole and dorsal aspect of the linear HT (B- B''). The presence of double-

positive cells in these areas reveals the differentiation of Isl1 lineage cells into

cardiomyocytes (A, B). Images are 74 and 134 optical sections acquired every 2.5μm

and covering 195μm and 335μm, respectively. In (B, B') only the double-positive

cells located in the linear HT are shown, while the tdtomato+ progenitors, located

outside the linear HT are not shown. Lateral views are shown in (A', B') including 3D

reconstructions. (A'') Cross-sections in xy along the dotted lines shown in (A').

Dorsal views are shown in (B'') including the 3D reconstruction of the tdtomato+

cells located in the linear HT. White arrows show the dorsal (mesocardial) regions of

the linear HT. (C) Snapshots of an *Ils1$^{cre/+}$; Rosa26Rtdtomato$^{+/-}$; Nkx2.5eGFP* embryo

at different time points during the third phase showing that the tdtomato+ SHF cells

contribute to the arterial pole of the HT. Images are 52 optical sections acquired every

5μm and covering 260μm. The embryo was imaged continuously every 13m for the

first 6h of culture and then imaged once again at time point 9h30m. Note that at the

last time point, the laser was increased at maximum power in order to reveal more

clearly the red labeled cells at the arterial pole. The contrast had to be enhanced as

well. (C') Images at time points 0, 4h48m and 9h30m. Images at time points 0 and

4h48m are z-maximum projection of 5 sections covering 25 μm. Image at 9h30m is a

single optical section. at: arterial pole. endo: endoderm, splan meso: splanchnic

mesoderm. Scale bars: 100μm.

1350

Figure 10. A model of cardiomyocyte differentiation dynamics during early heart development

We propose that two distinct phases of cardiomyocyte differentiation take place during early heart development. At EHF stage the cc differentiates and starts folding. Cardiomyocytes round up and become contractile while the cardiac progenitors located in the splanchnic mesoderm remain undifferentiated. Subsequently, the cc. undergoes further morphogenesis to transform into a HT, initially open dorsally and no cardiomyocyte differentiation is detected during this transformation. Finally, cardiac differentiation resumes contributing new cardiomyocytes from the splanchnic mesoderm to the arterial pole (prospective RV) and the dorsal closure of the HT.

VIDEO LEGENDS:

Video 1: 3D reconstruction of the cardiac crescent at EHF stage, based on *Nkx2.5$^{cre/+}$ Rosa26Rtdtomato$^{+/-}$* signal. Related to Figure 1A.

Video 2: 3D reconstruction of Mouse HT formation, based on embryos immunostained for cTnnT. Five representative stages are represented. Related to Figure 1C-G.

Video 3: 3D reconstruction of the FHF (green) and SHF (red). Based on *Nkx2.5$^{cre/+}$ Rosa26Rtdtomato$^{+/-}$* embryos immunostained for cTnnT. The border between cTnnT-negative and cTnnT-positive cells can be visualized at the interface between the red and green domains. Related to Figure 2B.

Video 4: z-maximum projection of an *Nkx2.5$^{cre/+}$;Rosa26RmT/mG$^{+/-}$* embryo at cc stage and single-section time-lapse video of the same embryo after 10h 42m of ex-vivo culture (representative analysis from 3 embryos). Interval between frames: 1s. Related to Figure 4E.

Video 5: Time-lapse video of a *Mesp1$^{cre/+}$; Rosa26Rtdtomato$^{+/-}$* embryo from cc up to open HT stage (h:mm:ss) (representative analysis from 4 embryos). Interval between frames: 4m 30. Duration of the video: 13h 16m 30s. Related to Figure 4F.

1384

1385 Video 6: Same embryo as in Video 5. Images are z-maximum projection of 9 sections

1386 acquired every 4μm covering 36μm (h:mm:ss) and allows visualization of the inside

1387 of the cardiac lumen during HT formation. Interval between frames: 4m 30s. Duration

1388 of the video: 8h 10m. Related to Figure 4F'.

1389

1390 Video 7: Time-lapse video of an *Nkx2.5^{cre/+}; Rosa26RmT/mG^{+/-}* embryo (h:mm:ss)

1391 (representative analysis from 2 embryos). Interval between frames: 22m. Duration of

1392 the video: 4h 02m. Related to Figure 4G-G'.

1393

1394 Video 8: Brightfield time-lapse video of a wild type embryo, from cc stage showing

1395 the onset of cardiomyocyte contractility and the formation of the cardiac lumen

1396 (h:mm:ss) (representative analysis from 2 embryos). Interval between frames: 1m 03s.

1397 Duration of the video: 6h 04m 21s. Representative analysis from 2 embryos.

1398

1399 Video 9: Same embryo shown in Video 14 zoomed-in at the level of the splanchnic

1400 mesoderm (mm:ss). SHF cells (green arrows) displace antero-medially relative to the

1401 underlying endoderm (red arrow). In addition, SHF cells move apart from each other

1402 (distance between the green arrows increase from 61 to 83μm over 3h12m). Related

1403 to Figure 6H and Figure 6- figure supplement 6A.

1404

1405 Video 10: Time-lapse video sequences of *Nkx2.5eGFP* embryos from Early Bud/E7.5

1406 stage (h:mm:ss). (representative analysis from 2 embryos). Interval between frames:

1407 12m. Duration of the video: 7h 24m Related to Figure 5-figure supplement 1A.

1408

1409 Video 11: Time-lapse video sequences of *Nkx2.5eGFP* embryos from cc to

1410 transversal HT stage (h:mm:ss). (representative analysis from 3 embryos). Interval

1411 between frames: 24m. Duration of the video: 6h 48m. Related to Figure 4H.

1412

1413 Video 12: Time-lapse video sequences of *Nkx2.5eGFP* embryos from transversal HT

1414 to open HT stage (h:mm:ss) (representative analysis from 3 embryos). Interval

1415 between frames: 8m 09s. Duration of the video: 5h 01m 33s. Related to Figure5 -

1416 figure supplement 1B.

1417

1418   Video 13: Cell tracks in 3D+t during stages when cc differentiation takes place -from

1419   EHF onwards- (h:mm:ss). Cells are represented as green spheres if considered GFP+

1420   at the end of the time-lapse analysis. and as red spheres when considered GFP-

1421   (Representative analysis from 3 embryos). Interval between frames: 19m 18s

1422   Duration of the video: 5h 28m 06s. Related to Figure 5A.

1423

1424   Video 14: Cell tracks in 3D+t during stages the open HT forms. Cells are represented

1425   as green spheres if considered GFP+ at the end of the time-lapse analysis and as red

1426   spheres when considered GFP-(Representative analysis from 3 embryos). Interval

1427   between frames: 7m 40s. Duration of the video: 3h 12m. Related to Figure 6A.

1428

1429   Video 15: Time-lapse video of a *RERT$^{+/-}$; Rosa26Rtdtomato$^{+/-}$; Nkx2.5eGFP* embryo

1430   during the stages at which the transversal HT transforms into the open HT (h:mm:ss).

1431   Interval between frames: 19m 19s. Duration of the video: 7h 24m 35s. (2nd half) 3D

1432   reconstruction at three time points based of the eGFP signal (green) and red-labeled

1433   cells located in the splanchnic mesoderm. Related to Figure 7A.

1434

1435   Video 16: Time-lapse video of an *Isl1$^{cre/+}$; Rosa26Rtdtomato$^{+/-}$; Nkx2.5eGFP* embryo

1436   (h:mm:ss) (representative analysis from 3 embryos). Interval between frames: 6m.

1437   Duration of the video: 7h 18m. Related to Figure 8A.

1438

1439   Video 17: Same embryo shown in Video 16 zoomed-in at the level of the splanchnic

1440   mesoderm (h:mm:ss). Related to Figure 8C.

1441

1442   Video 18: 3D rendering of an *Isl1$^{cre/+}$;Rosa26Rtdtomato$^{+/-}$; Nkx2.5eGFP* embryo at

1443   transversal HT stage. Related to Figure 9A.

1444

1445   Video 19: 3D rendering of an *Isl1$^{cre/+}$;Rosa26Rtdtomato$^{+/-}$; Nkx2.5eGFP* embryo at

1446   linear HT stage. Related to Figure 9B.

Figure 1

differentiation of the cardiac progenitors

A E7.5/EHF

Nkx2.5cre/-
Rosa26Rtdtomato+/-
DAPI

A' ventral view

B E7.5/EHF

cTnnT
DAPI

hf.
fg.

C initial cardiac crescent

hf.
fg.
cc.
endocardium
endoderm

C' ventral view

1.55 106 um3

D cardiac crescent

fg.
cc.
endocardium
endoderm

D' ventral view

3.29 106 um3

HT formation   dorsal closure/outflow formation

E Transversal HT

cTnnT
DAPI
fg.

E' ventral view

3.46 106 um3

E''

F Open HT

fg.

F' ventral view

5.57 106 um3

F''

G Linear HT

fg.

G' ventral view

7.46 106 um3

G'' arterial pole / RV dorsal closure

H

outflow lenght (um)

**

open HT   linear HT

I

volume (um3)

0.0571   ns
ns   0.0476
ns   0.0173

cc   differentiated cc   transversal HT   open HT   linear HT

J

Nkx1.5eGFP
DAPI

Figure 1- figure supplement 1

A

E7/bud stage

1) allantois visible
2) yolk salk and amniotic cavity visible

E7.5/neural plate stage

1) the allantois has elongated
1) no head fold visible
2) no clear anterior neural groove

20X

20X

cardiogenic plate

B

E7.5/EHF

1) head fold visible
2) clear anterior neural groove

Cardiac stages

1) foregut pocket visible
2) headfolds prominently elevated

E8.2

20X

20X

20X

cardiac crescent progenitors

cardiac crescent

transversal HT

open HT

linear HT

Figure 1- figure supplement 2

Figure 1- figure supplement 3

A EHF

A'

CTnnT

B EHF

fg.

Mesp1cre-R26Tdtomato

B'

cc.

endo.

cc.

endoc.

endo.

Mesp1cre/- Rosa26Rtdtomato+/-

cTnnT

Dapi

Figure 2

Figure 2- figure supplement 1

A

Nkx2.5^{Cre}
Rosa26Rtdtomato^{+/+}
DAPI
cTnnT

endoderm

A'

segmentation FHF/SHF

Figure 2- figure supplement 2

Figure 3

Figure 3- figure supplement 1

Figure 3- figure supplement 2

Figure 3- figure supplement 3

A

Figure 4

A E7.5/EHF → HT stage
20X na1
5-6 h
5-7 h

B δt
z
x,y=610um

C

D E7.5 EHF | +10h24m of culture
V
A
A
beating HT

E cardiac crescent | +10h42m of culture
0 | +1s
Nkx2.5^cre/- Rosa26RmT/mG^+/-
100 µm

F z-projection
cardiac crescent | Transversal HT
0 | 1h52m | 3h20m | 6h33m
cc | cc | cc | cc
fg. | fg. | |
splan.meso | splan.meso | splan.meso | splan.meso
nt | nt
mesp1^cre/- Rosa26tdtomato^+/-

F'' single section
cc | cc
endoc | endoc
endoderm | endoderm
0 | 1h52m

F' 36 um z-projection | open HT
0 | 1h52m | 3h20m | 4h48m | 6h33m | 8h35m
myo.
endo.
inflow | inflow
ao. | ao.

G z-projection
differentiation of the cc
cardiac crescent
0 | 2h59m | 5h18m
hf | hf | hf
fg | fg
Nkx2.5^cre/- Rosa26RmT/mG^+/-

G' single section
0 | 2h16m | 3h48m
endo | cc. endo
4h33m | 4h55m | 5h18m

G''
cc
endoc
endoderm
3h48m
cc
5h18m

H z-projection
differentiation of the cc
cardiac crescent
0 | 3h18m | 6h12m
Nkx2.5:eGFP

H' single section

H'' single section

I E7.5/EHF — around 12-16 hours — E8.2
5-6 hours → 5-7 hours → 2-3 hours →
Differentiation of the cc | Formation of the open HT | Dorsal closure
cardiac crescent | transverse HT open HT | Linear HT

Figure 4- figure supplement 1

A

Figure 4- figure supplement 2

A

mesp1^{cre/-}
Rosa26tdtomato^{+/-}

B

Figure 5

Figure 5- figure supplement 1

A

z-projection

E7/bud stage   neural plate stage

Nkx2.5:eGFP   0   yolk sac   7h24m   brightfield

yolk sac

B   morphogenesis   **B'**

z-projection

Transversal HT   open HT

Nkx2.5:eGFP   0   2h33m   5h12m   5h12m

open dorsally

C   **C'**

open HT   linear HT

Nkx2.5:eGFP   0   2h41m   2h41m

closed dorsally

Figure 5- figure supplement 2

A

Nkx2.5 eGFP
RERT^cre/- Rosa26Rtdtomato^+/-
optical section

0 m

B

1h38m   2h 17m   2h 37m   3h 16m   4h 34m

optical section

C

GFP intensity (AU)

daughters 1' and 2'

mother 1

t(mn)

D

Z max proj

Nkx2.5eGFP   5h 53m

time

Figure 5- figure supplement 3

A

RERT*cre/-* Rosa26Rtdtomato*
Nkx2.5 eGFP

Figure 6

Figure 6- figure supplement 1

A

Nkx2.5 eGFP
RERT^{cre/-} Rosa26Rtdtomato^{+/-}

cardiac crescent   0   transversal HT   4h 55m   7h 30m

Figure 7

Figure 8

A z-max proj. cc. *Isl1*^{cre/-} *Rosa26Rtdtomato*^{+/-} *Nkx2.5 eGFP* 0 2h24m 4h00 5h12m 7h00

B single optical section cc. 2h36m 4h00m

C 30um z-max projection 2h24m 4h00 6h00 6h24m 7h00 7h00 endo inflow splan. meso endo splan. meso

D postfixation cTnnT *Isl1*^{cre/-} *Rosa26Rtdtomato*^{+/-} *Nkx2.5 eGFP* Nkx2.5eGFP cTnnT

Figure 8- figure supplement 1

RERT^{cre/-} *Rosa26Rtdtomato*^{+/-}
Nkx2.5 eGFP

Figure 8- figure supplement 2

A

RERT^cre/- Rosa26Rtdtomato^+/-
Nkx2.5 eGFP

0
1h30m
2h00m

B

inflow
1h30m

inflow
1h48m

inflow
2h00m

1 2 3
1 2 3
1 2 3

C

Figure 9

Figure 10

approx 5-6 hours

approx. 2-3 hours

approx. 5-7 hours

differentiation events

time

0
EHF/E7.5

12-16hours
E8.2

cc progenitors   cardiac crescent   transversal HT   open HT   linear HT