Catecholamines Alter the Intrinsic Variability of Cortical Population Activity and Perception

Thomas Pfeffer (1), Arthur-Ervin Avramiea (2), Guido Nolte (1), Andreas K. Engel (1), Klaus Linkenkaer-Hansen (2), Tobias H. Donner (1,3,4)

1 Department of Neurophysiology and Pathophysiology, University Medical Center Hamburg-Eppendorf, Hamburg, Germany
2 Center for Neurogenomics and Cognitive Research, Neuroscience Campus Amsterdam, VU University Amsterdam, The Netherlands
3 Department of Psychology, University of Amsterdam, The Netherlands
4 Amsterdam Brain and Cognition, University of Amsterdam, The Netherlands

Correspondence:
thms.pfffr@gmail.com (T.P.)
t.donner@uke.de (T.H.D.)

Abbreviated title: Catecholamines Alter Cortical and Perceptual Variability

ABSTRACT

The ascending modulatory systems of the brainstem are powerful regulators of

global brain state. Disturbances of these systems are implicated in several major

neuropsychiatric disorders. Yet, how these systems interact with specific neural

computations in the cerebral cortex to shape perception, cognition, and behavior

remains poorly understood. Here, we probed into the effect of two such systems,

the catecholaminergic (dopaminergic and noradrenergic) and cholinergic systems,

on an important aspect of cortical computation: its intrinsic variability. To this end,

we combined placebo-controlled pharmacological intervention in humans,

magnetoencephalographic (MEG) recordings of cortical population activity, and

psychophysical measurements of the perception of ambiguous visual input. A

low-dose catecholaminergic, but not cholinergic, manipulation altered the rate of

spontaneous perceptual fluctuations as well as the temporal structure of "scale-

free" population activity of large swaths of visual and parietal cortex.

Computational analyses indicate that both effects were consistent with an

increase in excitatory relative to inhibitory activity in the cortical areas underlying

visual perceptual inference. We propose that catecholamines regulate the

variability of perception and cognition through dynamically changing the cortical

excitation-inhibition ratio. The combined read-out of fluctuations in perception and

cortical activity we established here may prove useful as an efficient, and easily

accessible marker of altered cortical computation in neuropsychiatric disorders.

INTRODUCTION

The modulatory systems of the brainstem send widespread, ascending

projections to the specialized circuits of the cerebral cortex that mediate

perception, cognition, and goal-directed behavior. These systems regulate

ongoing changes in brain state, even during periods of wakefulness [1–4]. They

are recruited during, and in turn shape, cognitive processes, such as perceptual

inference, learning, and decision-making [5–8]. Because these systems are

implicated in most neuropsychiatric disorders, they are also major targets of

pharmacological therapy of brain disorders [5,9,10]. Taken together,

neuromodulatory systems have remarkably specific effects on cognition, despite

the widespread and diffuse nature of their projections to cortex. An important

challenge for neuroscience is to uncover the mechanistic principles, by which

neuromodulatory systems interact with the cortical computations underlying

cognition.

One key parameter of cortical computation that might be under

neuromodulatory control is the intrinsic variability – i.e., fluctuations that occur

during constant (or absent) sensory input [13,14]. Specifically, it has been

proposed that the catecholaminergic neuromodulators noradrenaline and

dopamine may shift the cortical computations underlying decision-making from a

stable ("exploitative") to a variable ("exploratory") mode [5,15]. A context-

dependent adjustment of the variability of cortical computations may also be

adaptive for perceptual inference in the face of ambiguous sensory input [16].

Animal work has shown that catecholamines and acetylcholine, another

important neuromodulator, alter the intrinsic variability of neural activity [2,11,17–

19] through highly selective interactions with specific elements (pyramidal cells

and/or inhibitory interneurons) of cortical microcircuits [20,21]. But it is unknown

how these changes at the level of cortical microcircuits relate to the intrinsic

variability of perception and cognition.

At the larger scale of cortical mass action that is assessable with non-

invasive recordings in humans, activity also fluctuates intrinsically, in a spatially

and temporally structured manner [22,23]. The temporal structure of these

fluctuations is characteristic of so-called "scale-free" behavior: Power spectra that

scale as a function of frequency according to a power law, $P(f) \propto f^{\beta}$ [24,25],

indicating long-range temporal autocorrelations [26–29]. Some studies have

linked the spatio-temporal structure of the fluctuations in cortical population

activity to specific perceptual and cognitive processes [28,30–32]. But it is

unknown if and how these fluctuations in cortical population activity are

dynamically regulated by neuromodulatory systems.

We aimed to close these gaps by systematically quantifying the effects of

catecholaminergic and cholinergic neuromodulation on the intrinsic variability in

perception and large-scale cortical activity in the healthy human brain. To this

end, we combined placebo-controlled, selective pharmacological interventions,

psychophysical measurements of fluctuations in perception in the face of a

continuously presented and ambiguous visual stimulus, and MEG recordings of

fluctuations in cortical population activity.

Catecholamines, but not acetylcholine, increased both the variability of

perception as well as long-range temporal correlations of intrinsic cortical activity

in visual and parietal cortex. Based on previous theoretical and experimental

work [33–36], we interpreted the increase in perceptual variability in terms of an

increase in the net ratio between cortical excitation and inhibition in those cortical

regions. Simulating a recurrent neural network under synaptic gain modulation

enabled us to show that an analogous mechanism may account for the increase

of long-range temporal correlations of cortical activity under catecholamines.

RESULTS

We tested for changes in intrinsic fluctuations of perception and cortical

population activity under placebo-controlled, within-subjects pharmacological

manipulations of catecholamine (using the noradrenaline reuptake inhibitor

atomoxetine) and acetylcholine (using the cholinesterase inhibitor donepezil)

levels (Fig 1A, see Methods for details on the pharmacological interventions).

Fig 1. Experimental design **(A, B)** Types and time course of experimental sessions. **(A)** Each
subject participated in three sessions, involving administration of placebo, atomoxetine, or
donepezil (session order randomized across subjects). Each session entailed the administration of
two pills, in the order depicted for the different session types. **(B)** Within each session, subjects
alternated between three conditions, Fixation, Task-Counting and Task-Pressing, during which
MEG was recorded (runs of 10 min each). See Materials and Methods for details. **(C)** Group
average power spectrum, averaged across all MEG sensors, for Rest and Task (Placebo condition
only).
114

Fluctuations in cortical activity were measured during two steady-state conditions,

both of which excluded transients in sensory input or motor output (Fig 1B): (i)

fixation of an otherwise gray screen (a condition termed "Fixation"), as in most

studies of human "resting-state" activity [22,23]; and (ii) silent counting of the

spontaneous alternations in the perceptual interpretation of a continuously

presented, ambiguous visual stimulus (dubbed "Task-counting"). In a third

condition that was only used for the analysis of perceptual fluctuations, subjects

immediately reported the perceptual alternations by button press ("Task-

pressing"; i.e. associated with movement-related transients in cortical activity).

This design capitalized on recent insights into the circuit mechanisms underlying

intrinsic perceptual dynamics [33,34,36] which helped constrain the mechanistic

interpretation of the results reported below.

The Results section is organized as follows. We first present the effects of

the "Atomoxetine" and "Donepezil" conditions (each compared against the

"Placebo" condition) on the rate of perceptual fluctuations. These effects were in

line with a boost in the relative strength of excitatory drive of visual cortex under

Atomoxetine. We then show how (i) constant sensory and task drive (i.e., Task-

counting vs. Fixation) and (ii) the pharmacological manipulations affect the

intrinsic fluctuations in cortical activity. We focus on the temporal auto-correlation

structure of intrinsic fluctuations in the amplitude of band-limited cortical

population activity (see Methods and Fig 2). Control analyses showing the drug

effects on other measures of cortical population activity and peripheral

physiological signals support the validity and specificity of our conclusions.

Fig 2. Quantifying the temporal structure of fluctuations in oscillatory cortical activity
(A) *Top.* Time-frequency representation of MEG power fluctuations during Rest (example subject).
Bottom. Filtered signal (10 Hz; black) and the corresponding amplitude envelope (red). **(B)**
Illustration of detrended fluctuation analysis. See main text (Materials and Methods) for details. *Top.*
Cumulative sum of the amplitude envelope. *Bottom.* Detrending of cumulative sum within segments,
shown for two different window lengths N (N_1 = 500 ms and N_2 = 1200 ms). **(C)** Root-mean-square
fluctuation function $<F_N>$. In log-log coordinates, $<F_N>$ increases approximately linearly as a
function of N, with a slope that is the scaling exponent α. **(D)** Illustration of power spectrum
analysis of amplitude envelope. In log-log coordinates, the power spectrum can be approximated by
a straight line, with a slope β (power-law exponent) and an area under the curve (gray) that
quantifies the overall variance of the signal.
150

We close with simulations of a patch of recurrently connected excitatory and

inhibitory integrate-and-fire neurons. The simulations show that the changes in

temporal correlations observed in the MEG data can be explained by a

modulation synaptic gain that altered the net ratio between excitatory and

inhibitory activity.

Atomoxetine increases the rate of bistable perceptual fluctuations

The ambiguous visual stimulus that was continuously presented during both

Task-counting and Task-pressing induced ongoing fluctuations in perception, i.e.,

spontaneous alternations between two apparent rotation directions of 3D-motion

(Fig 1B; see Movie M1), a phenomenon is referred to as multi-stable perception.

The rate of the perceptual alternations reported by the participants provided a

read-out of visual cortical circuit state. Current models explain bistable perceptual

fluctuations in terms of the interplay between feedforward, excitatory drive of

stimulus-selective neural populations in visual cortex, mutual inhibition between

them, stimulus-selective adaptation, and neural "noise" [33,34]. Increases in the

ratio between feedforward, excitatory input to, and mutual inhibition within the

cortical circuit, give rise to faster perceptual alternations. This idea is supported

by convergent evidence from functional magnetic resonance imaging, magnetic

resonance spectroscopy, and pharmacological manipulation of GABAergic

transmission [31,36]. We reasoned that neuromodulators such as noradrenaline

might dynamically change these parameters [12,37], and thereby alter the rate of

perceptual fluctuations.

Atomoxetine increased the rate of perceptual fluctuations compared to

both Placebo and Donepezil conditions (Fig 3A; Atomoxetine vs. Placebo: p =

0.007; t = 2.913; Atomoxetine vs. Donepezil: p = 0.001; t = 3.632; Donepezil vs.

Placebo: p = 0.966; t = -0.043; all paired t-tests, pooled across Task-counting and

Task-pressing). The perceptual alternation rates were highly consistent across

Task-counting and Task-pressing (Fig S1C), supporting the validity of the

counting condition as behavioral read-out of bistable perceptual fluctuations.

Likewise, the Atomoxetine effect on perceptual fluctuation rate was evident for

Task-counting *(p* = 0.045; *t* = 2.103; paired t-test; Fig S1A) and Task-pressing *(p*

= 0.018; *t* = 2.540; paired t-test; Fig S1B) individually.

Fig 3. Atomoxetine, but not Donepezil, increases the rate of perceptual alternations **(A)** Number of
perceptual alternations reported by the subjects per 10 min run, pooled across task conditions
(Task-counting and Task-pressing). **(B)** Same as (A), after regressing out blink and eye movement
data (see Methods and Supplementary Figure S2). Significance was assessed using two-sided
paired t-tests (N=28).
190

These changes in perceptual fluctuations were not explained by an

increase in the rates of eye blinks or fixational eye movements. First, there was

no significant increase during Atomoxetine compared to Placebo in any of five

different eye movement parameters measured here (Fig S2). Second, none of

these parameters correlated with the perceptual alternation rate (Fig S2). Third,

and most importantly, the effect of Atomoxetine on the perceptual dynamics was

also significant after removing (via linear regression) the individual eye movement

parameters (Fig 3B).

In sum, Atomoxetine had an effect on bistable perceptual fluctuations that

was both robust and specific, evident when compared with either Placebo or

Donepezil. This effect was in line with an increase in the strength of excitatory

feedforward drive of visual cortex relative to the strength of mutual inhibition

between the neural sub-populations encoding the competing perceptual

interpretations of the ambiguous stimulus. Such an effect should have occurred in

the motion-sensitive visual cortical areas, which implement the visual competition

induced by the ambiguous structure-from-motion stimulus [38,39].

Atomoxetine increases the scaling exponent of fluctuations in cortical

population activity

We estimated long-range temporal correlations of band-limited amplitude

fluctuations (indicated by the scaling exponent α; see Methods for details) to

quantify intrinsic fluctuations in cortical population activity. Our analyses focused

on amplitude envelope fluctuations in the 8–12 Hz frequency range ("alpha

band"), for two reasons. First, as expected from previous work [40], the cortical

power spectra exhibited a clearly discernible peak in this frequency range, which

robustly modulated with sensory or task drive (suppressed under Task-counting,

Fig 1C). Second, previous studies reported robust long-range temporal

correlations with peaks in the same frequency range [26,29].

We first replicated two previously established observations pertaining to

the scaling exponent α. First, the average across cortical patches and

participants was α = 0.67 (s.d. = ±0.09) during Fixation (Placebo only) and α =

0.64 (s.d. = ±0.07) during Task-counting (Placebo only), indicative of long-range

temporal correlations similar to the ones found in previous work [26,29,41].

Second, the sensory and task drive during Task-counting reliably reduced α

compared to Fixation, again as shown in previous work [27,42]. Across all voxels,

α was significantly larger during Fixation than during Task-counting (p = 0.0062; t

= 2.97; paired t-test; Placebo condition only). This difference was significant

across pharmacological conditions in large parts of cortex including the occipital

and parietal regions that were driven by the motion stimulus (p < 0.05; cluster-

based permutation test; Fig 4D).

Fig 4. Effects of task and pharmacological conditions on long-range temporal correlations of the
amplitude envelope of 8-12 Hz MEG activity. **(A)** Spatial distribution of significant differences in
scaling exponent α between Task-counting and Fixation during Placebo (left), Atomoxetine
(middle), and Donepezil (right). **(B)** Comparison between mean scaling exponents α averaged
across all the entire brain (see Methods) during the different pharmacological conditions. **(C)**
Individual subject differences in scaling exponent α between all drug conditions. **(D, E)** Spatial
distribution of drug-induced changes in scaling exponents. **(D)** Atomoxetine vs. Placebo. **(E)**

Donepezil vs. Placebo. Two-sided permutation tests (N=28); all statistical maps: Threshold at $p =$
0.05, cluster-based. All drug comparisons are averaged across behavioral conditions, i.e., Fixation
and Task-counting.
242

Having verified the validity of our measurements of α we then tested for

changes in α under the pharmacological conditions (Fig 4B-E and Fig 5). There

was a highly significant increase in α for Atomoxetine compared to Placebo when

collapsing across voxels as well as across Fixation and Task-counting ($p =$

0.0068; $t = 2.93$; paired t-test, Fig 4B-C). This effect was widespread, but not

homogenous across cortex, comprising occipital and posterior parietal, as well as

a number of midline regions including the thalamus (Fig 4D, $p = 0.0022$; cluster-

based permutation test). Because it is unclear to which extent intrinsic activations

from deep sources can be recovered using MEG, we focus our description and

conclusions on the effects in cortical regions. Importantly, the Atomoxetine effect

on α was also present at the level of MEG sensors (Fig S4), and hence did not

depend on the source reconstruction method applied here (see Methods).

Fig 5. Atomoxetine increases long-range temporal correlations irrespective of behavioral condition**.**
Spatial distribution of the Atomoxetine-induced changes in scaling exponent α during **(A)** Fixation
and **(B)** Task-counting. **(C)** Conjunction of maps in (A) and (B), highlighting (in green) voxels with
significant increases in both conditions. **(D)** Scaling exponents for Fixation (gray) and Task-counting
(yellow) within conjunction cluster depicted in panel C for Placebo (top), Atomoxetine *(*middle*)* and
Donepezil only (bottom*).*
262

The effect of Atomoxetine on α was subtle, likely due to the low dosage.

But, importantly, the effect was highly reproducible across repeated

measurements. We assessed reproducibility with two complementary

approaches. The first was a region-of-interest (ROI) analysis. We defined a ROI

in terms of a significant cluster for Atomoxetine > Placebo (one-sided paired t-test,

$p < 0.05$, uncorrected) during the first run collected in each session (Fixation and

Task-counting collapsed) and extracted this ROI's mean α from the second run.

We then reversed the procedure and so extracted a second, independent ROI-

based α and averaged the α-estimates. This approach revealed a strong

increase under Atomoxetine ($p = 0.0023; t = 3.365$). The second approach

assessed the reproducibility of the spatial pattern of effects across both runs. To

this end, we correlated the (non-thresholded) individual maps for the Atomoxetine

vs. Placebo difference computed from the first and second run in each session

(again pooling across Task-counting and Fixation) and tested the resulting

correlation coefficients across participants. The average correlation was

significantly different from zero (mean r = 0.29, p < 0.0001; permutation test

against zero).

The Atomoxetine-related increases in scaling exponent α were evident

during both Fixation and Task-counting (Fig 5A, Fixation: $p = 0.0245$; Fig 5B,

Task-counting: $p = 0.0035$; cluster-based permutation test). The effects occurred

in largely overlapping regions of occipital and parietal cortex (Fig 5C). There was

no interaction between the effects of Atomoxetine and Task-counting anywhere in

cortex: a direct comparison of the two Atomoxetine vs. Placebo difference maps,

from Fixation and from Task-counting, yielded no significant clusters ($p > 0.081$

for all clusters; cluster-based permutation test). The same cortical regions in

which α increased during Atomoxetine exhibited decreases during Task-counting:

When testing for the task-dependent change in α (Fig 4A) specifically in the

regions comprising the conjunction cluster of the Atomoxetine effect (Fig 5C), the

reduction during Task-counting was also highly significant (Fig 5D) in all

pharmacological conditions.

In contrast to the robust effect of Atomoxetine on α, there was no

evidence for an effect of Donepezil at the dosage used here. The difference

between Donepezil and Placebo (collapsed across Fixation and Task-counting)

did not reach significance, neither when pooling across voxels ($p = 0.50$; $t = 0.68$;

BF = 0.68; paired t-test; Fig 4B), nor when testing all voxels individually ($p > 0.22$

for all clusters; cluster-based permutation test; Fig 4E; Fig S5). Atomoxetine also

increased the scaling exponents when directly compared to Donepezil during

Task-counting (Fig S6A; $p < 0.05$; two-sided cluster-based permutation test), but

not during Fixation (Fig S6B).

Taken together, the rich experimental design gave rise to a highly specific

and consistent pattern of changes in α under the different experimental

conditions, which helped constrain the mechanistic interpretation of the results.

The Atomoxetine effects were specific, and not just due to the application of any

drug targeting neurotransmitter systems. It is possible that the absence of

detectable Donepezil effects on α was due to the low dosage or short

administration period used here. However, the control analyses presented in the

next section revealed clear effects of Donepezil on both cortical activity as well as

markers of peripheral nervous system activity.

Control analyses for the drug effects on other features of cortical dynamics

or peripheral physiological signals

During Fixation, Atomoxetine and Donepezil both reduced posterior cortical

alpha-band power relative to Placebo in both the 8-12 Hz (Fig 6A; $p < 0.05$ for all

clusters; two-sided cluster-based permutation test) as well as the 2-8 Hz

frequency ranges (Fig S7A). This suppression in low-frequency power under

cholinergic boost is consistent with previous work in rodents [17,18] and humans

[43].

320
321   **Fig 6.** Similar effects of Atomoxetine and Donepezil on 8-12 Hz power. **(A)** Spatial distribution of
322   drug-related alpha power changes during Fixation, thresholded at $p = 0.05$ (two-sided cluster-based
323   permutation test). *Left.* Power changes after the administration of atomoxetine. *Right.* Power
324   changes after the administration of donepezil. **(B)** Same as (A), but for Task-counting. All threshold
325   at $p = 0.05$, cluster-based two-sided permutation tests (N=28).
326
The Atomoxetine-induced changes on 8-12 Hz power exhibited a different

spatial pattern from the one of corresponding change in the scaling exponent α:

within the cluster of the significant main effect of Atomoxetine on α (Fig 4D),

power did not correlate with the changes in α (group average spatial correlation

between pooled difference maps within cluster; $r = 0.073$; $p = 0.129$, *BF* = 1.065).

During Task-counting, neither drug altered MEG-power in the low-frequencies (8-

12 Hz: Fig 6B, p > 0.05 for all clusters; two-sided cluster-based permutation test;

2-8 Hz: Fig S7B), presumably due to the already suppressed power in the 8-12

Hz range in that condition (Fig 1C). Together with the findings reported in the

previous section, the analyses of the mean MEG power indicate that (i) both

drugs reduced the amplitude of cortical low-frequency oscillations and (ii) MEG

power and the scaling exponent α reflected at least partially distinct aspects of

intrinsic cortical dynamics

We also controlled for changes in peripheral physiological signals under

the drugs as potential confounds of the effect on cortical scaling behavior (Fig 7).

As expected, Atomoxetine increased average heart rate (Fig 7A,B). Donepezil

had no detectable effect on average heart rate, during neither Fixation (p =

0.8676; t = 0.16; paired t-test; *BF* = 0.8676; Fig 7A) nor Task-counting (p =

0.3274; t = 1.0; paired t-test; *BF* = 0.3139; Fig 7B). Both drugs altered heart-rate

variability, increasing α computed on the time series of inter-heartbeat-intervals

(see Methods) in both behavioral contexts relative to Placebo (Fixation: p =

0.0012, t = 3.62; Task-counting: p = 0.0167; t = 2.55; Fig 7C; Fixation/Donepezil:

p = 0.0076, t = 2.88; Task-counting/Donepezil: p = 0.0049, t = 3.06; Fig 7D; all

paired t-tests). Critically, the Atomoxetine-induced changes in heart rate showed

no (Task-counting: r = 0.00*; p* = 0.99; Person correlation; *BF* = 0.15) or only weak

and statistically non-significant (Fixation: r = 0.24; p = 0.21; Person correlation;

BF = 0.31) correlations with the changes in cortical activity (Fig 7A/B, right).

Similarly, the Atomoxetine-related changes in the scaling behavior of inter-

heartbeat intervals were not correlated with the changes in cortical scaling

behavior (Fixation: r = 0.22; p = 0.26; *BF* = 0.27; Task-counting: r = 0.26; p =

0.19; *BF* = 0.35; Fig 7C/D, right). Atomoxetine also decreased spontaneous blink

rate during Fixation (p = 0.034; t = 2.24; paired t-test), but not during Task-

counting (p = 0.112; t = 1.645; *BF* = 1.130; paired t-test; Fig S2B). However,

again there was no significant correlation between changes in blink-rate and

changes in cortical scaling behavior due to Atomoxetine (Fixation: r = -0.26; p =

0.19; *BF* = 0.35; Task-counting: r = -0.09; p = 0.64; *BF* = 0.16).

Fig 7. Drug effect on cortical scaling behavior is not explained by systemic drug effects**. (A)** *Left.* Heart rate for Atomoxetine, Placebo and Donepezil during Fixation. *Right.* Correlation of Atomoxetine-related changes in heart rate (x-axis) with Atomoxetine-related changes in MEG scaling exponent *a* (y-axis) (within significant cluster during Fixation). **(B)** As (A), but during Task-counting **(C)** *Right.* Scaling behavior of inter-heartbeat intervals (heart scaling exponent). *Left.* Heart scaling exponent for all pharmacological conditions during Fixation. *Right.* Correlation of Atomoxetine-related changes in heart scaling exponent (x-axis) with Atomoxetine-related changes in MEG scaling exponent *a* (y-axis). **(D)** Same as (C), but during Task-counting. Two-sided t-tests and Pearson correlations (N=28). BF, Bayes factor.

In sum, drug-induced changes in peripheral physiological signals under the drugs, if present, did not account for the Atomoxetine-induced changes in the scaling behavior of the fluctuations in cortical activity (Figs 4 and 5). These controls support our interpretation in terms of a specific effect on cortical net E/I ratio rather than non-specific secondary effects due to the systemic drug effects or changes in retinal input due to blinks.

Change in scaling exponent under Atomoxetine is consistent with increase in net E/I ratio in cortical circuits

Atomoxetine had an effect on perceptual fluctuations that was in line with a relative increase in excitation in cortical circuits of occipital and posterior parietal cortex that processed the ambiguous visual motion stimulus. We reasoned that

this change in circuit state might have also produced the observed change in

scaling behavior of intrinsic cortical activity fluctuations under Atomoxetine. In

order to solidify this intuition, we simulated the activity of a neural network model

made up of recurrently connected excitatory and inhibitory integrate-and-fire units

(Fig 8). In what follows, we use the term "E/I ratio" to refer to the ratio of

excitatory and inhibitory activity across the circuit [44] and "E/I balance" to refer to

a specific regime of E/I ratios, in which excitation and inhibition changes in a

proportional way [45–48].

394
Fig 8. Changes in scaling behavior in a neural network under gain modulation. **(A)-(I)** Dynamic
modulation of excitation-inhibition ratio alters long-range temporal correlations in recurrent network
model. **(A)** Model architecture. The network consisted of 2500 excitatory and inhibitory integrate-
and-fire units and random, local (within an area of 7x7 units) connectivity (magnified within the red
square). **(B)** Neuromodulation was simulated as a gain modulation term multiplied with excitatory
synaptic weights (w_{ee} and w_{ie}). **(C)** Power spectrum of the simulated neural mass activity, with a
peak in the alpha range. **(D)** κ as a function of excitatory and inhibitory connectivity (with a spacing
of 2.5%; means across 10 simulations per cell). The region of $\kappa \sim 1$, overlaps with the region of $\alpha >$
0.5 and splits the phase space into an excitation-dominant ($\kappa > 1$) and an inhibition-dominant region
($\kappa < 1$). The black square depicts the network configuration that was chosen for assessing the effects
of neuromodulation **(E)** Scaling exponent α as a function of excitatory and inhibitory connectivity.
(F) Same as (D) and (E), but for mean firing rate. **(G)** κ as a function of independent synaptic gain
modulation. Red square, baseline state of critical network before synaptic gain modulation. White
line, axis corresponding to largest change in ratio **(H)** Same as (D), but for scaling exponent α. **(I)**
Same as (G) and (H), but for firing rate.

We started from a network (Figure 8A) that was similar to the one

developed and analyzed in a previous study [49]. The basic features of the model

were as follows. The model was built to generate oscillations of neural mass

activity (summed across all units) in the alpha-band (8-12 Hz; Figure 8B). The

amplitude envelope of these oscillations fluctuated over time, with scale-free

long-range temporal correlations. Those scale-free intrinsic fluctuations in cortical

activity were sensitive to variations in the percentage of excitatory and inhibitory

connections in the circuit (i.e., microstructure). Our previous work [49],

reproduced here (Figure 8D-F) showed that such a model accounts for the joint

emergence of two scale-free phenomena at different spatial scales (single unit

activity vs. mass activity) and temporal scales (tens of milliseconds vs. hundreds

of seconds): (i) neuronal avalanches with an event size distribution following a

power-law; and (ii) long-range temporal correlations of the amplitude envelope

fluctuations of the circuits mass activity. Both phenomena have been established

in empirical measurements of cortical population activity [26,50]. Neuronal

avalanches are activity deflections (i.e., exceeding a certain threshold) that

propagate through the cortical network [50], with an "event size" corresponding to

the number of activated units. In line with (Shew et al., 2009) we quantified the

power-law scaling of the size distributions of avalanches in the model with the

kappa-index (κ): the similarity between the actual event size distribution and a

power-law distribution with an exponent of -1.5; A κ of 1 indicates perfect match

between the two.

We extended this model by means of a multiplicative modulation of

synaptic gain [37,51] (Fig 8B). This allowed us to explore how catecholaminergic

effects on neural circuits might change the two phenomena of scale-free neural

population activity described above. We first determined the structural

connectivity (small squares in Fig 8D-F) and the time scale parameters of the

model such that the network generated intrinsic alpha-band oscillations (Fig 8C)

with amplitude fluctuations that exhibited neuronal avalanches with scale-free

event size distributions (Fig 8D) as well as long-range temporal correlations (with

$\alpha \sim 0.85$). We then independently modulated specific excitatory or inhibitory

connections through the multiplicative scaling of the corresponding synaptic

weights, in two ways. In the version shown in Fig 8, we modulated only excitatory

synapses, but independently on excitatory as well as inhibitory neurons (EE and

IE), thus producing asymmetries in the circuits net E/I ratio as in recent modeling

work on a cortical circuit for perceptual decision-making [44]. In the second

version (Fig S8A), we co-modulated EE and IE and independently modulated

inhibitory synapses on excitatory neurons (EI). This was intended to specifically

simulate glutamate receptors (AMPA or NMDA) in former two cases (mediating

the effects of excitatory neurons) as opposed to modulations of GABA receptors

in the latter case (mediating the effects of inhibitory neurons on others). N_{EE} and

N_{IE} were co-modulated by the same factor for simplicity, because we did not

assume that excitatory (glutamatergic) synapses would be differentially

modulated depending on whether they were situated on excitatory or inhibitory

target neurons.

Both types of changes in net E/I ratio robustly altered κ (Fig 8G and Fig

S8B), α (Fig 8H and Fig S8C), and the mean firing rate (Fig 8I). The effect of

changes in E/I ratio on the scaling exponent α were non-monotonic, dependent

on the starting point: increases in excitation led to increases in α when starting

from an inhibition-dominant point, but to decreases in α when starting from an

excitation-dominant point (Fig 8H, white line). The effects of excitatory and

inhibitory gain modulation on the temporal correlation structure of the simulated

population activity were qualitatively similar to the effects of changes in the

fraction of excitatory and inhibitory synapses simulated (as shown in Fig 8D-F).

The latter simulated individual differences in cortical anatomical microstructure,

and the former simulated state-dependent changes in cortical circuit interaction,

which occur within an individual brain.

In the model, the scaling exponent α exhibited a non-monotonic

dependence on E/I ratio (see the white diagonal line in Fig 8G-I and schematic

depiction in Fig 9). Consequently, without knowing the baseline state, any change

in α was ambiguous with respect to the direction of the change in E/I ratio (i.e.,

towards excitation- or inhibition-dominance). Thus, the observed increase in α

under Atomoxetine during Fixation could have been due to either an increase or

a decrease in E/I ratio.

Fig 9. Schematic of inference from observed change in scaling exponents to net E/I ratio (see
Results for details). The non-monotonic dependence of scaling exponent α on E/I ratio
(corresponding to white line in panel H) is replotted schematically. **(A)** Measured scaling exponent
α during Fixation (gray) can result from both, inhibition- or excitation-dominant regimes; the
baseline is unknown. Assuming that sensory drive (Task-counting; yellow dot) either decreases or
does not change E/I ratio, the observed decrease in scaling exponent during Task-counting (yellow)
reflects a shift towards the inhibition-dominance (blue arrows), consistent with animal physiology
[52,53]. **(B)** This constrains the baseline state for the interpretation of the Atomoxetine-induced
increase in scaling exponent during Task-counting (red): The latter increase likely reflects an
increase in E/I ratio (red arrow).

Importantly, insights from animal physiology helped constrain the baseline

state during Task-counting: In the awake state, visual drive decreases E/I ratio in

visual cortex V1, due to the recruitment of inhibitory mechanisms that outweigh

the excitatory sensory drive [52,53]. We assumed that the same held for the

Task-counting condition (constant visual stimulation) of our study.

This condition enabled us to infer the change in net E/I ratio under

Atomoxetine. The rationale is illustrated in Fig 9. The animal physiology results

referred to above indicte that the observed decrease in α during Task-counting

was due to a shift towards inhibition-dominance (yellow point in Fig 9A). Under

this assumption, the Atomoxetine-induced increase in α during was due to an

increase in net E/I ratio (Fig 9B). Because the effects of Atomoxetine on α were

the same during Task-counting and Fixation, it is likely that the same mechanism

was at play during Fixation.

In sum, under certain, the simulations provided a mechanistic explanation

for the observed MEG effects: effective changes in the cortical E/I ratio, due to

multiplicative changes of synaptic gain [37] or other mechanisms [12,20] – the

same conclusion inferred from the increase in the rate of perceptual alternations

above.

DISCUSSION

Neuromodulators regulate ongoing changes in the operating mode of cognitive

processes [1,5,6,10,54] as well as of cortical microcircuits [11,12,17,18,20,21].

Here, we unraveled the effect of two major classes of neuromodulators,

catecholamines and acetylcholine, on the intrinsic variability of cortical

computation, an important parameter shaping the operating mode. We used two

separate read-outs of this parameter: (i) the rate of fluctuations in perception

induced by ambiguous visual input and (ii) the temporal correlation structure of

(scale-free) fluctuations of the amplitude envelope of band-limited cortical mass

activity. Catecholamines, but not acetylcholine, altered both read-outs.

Simulations of a recurrent neural network revealed that, under well supported

physiological assumptions,, the observed changes in the temporal structure of

fluctuations in cortical activity is indicative of an increase in the cortical E/I ratio.

Earlier modeling and empirical data show that such an increase in net E/I ratio in

visual cortex is also consistent with the increased rate of perceptual fluctuations

under the catecholaminergic boost.

Cortical distribution of Atomoxetine effects on cortical activity fluctuations

The Atomoxetine effects on the scaling exponent were widespread across cortex,

but not entirely homogenous. They were pronounced across occipital and parietal

cortex, but not robust in frontal cortex (see Fig 5B). This distribution might point to

a noradrenergic, rather than dopaminergic origin. Atomoxetine increases the

levels of both catecholamines, noradrenaline and dopamine, but the cortical

projection zones differ substantially between both systems: Dopaminergic

projections mainly target prefrontal cortex [55] and only sparsely to occipital

cortex [56,57], whereas the noradrenergic projections are more widespread and

strong to occipital and parietal cortex [58]. Alternatively, this distribution may

reflect the different receptor composition across cortical regions [58,59]. The

relative frequency of the different noradrenaline receptors differs between pre-

frontal and posterior cortex [58], which might translate a homogenous

noradrenaline release into a heterogeneous effect on the activity in these

different cortical regions. An important next step will be to investigate the

differential role of different noradrenaline receptors, and different regional

receptor profiles, in shaping the cortex-wide effect of noradrenaline on long-range

temporal correlations.

Opposite effects of external drive and catecholamines on long-range

temporal correlations

Consistent with our current results, previous studies also found a decrease in

temporal autocorrelations of cortical activity due to external drive [27,42]. The

observation is consistent with the insight from intracellular recordings of cortical

neurons in animals, that cortical responses to sensory stimulation in the awake

state are dominated by inhibition [52,53,60,61]. One candidate source of this

sensory-driven state change is thalamocortical inhibition [62], but intracortical

feedback inhibition might also contribute [63]. Modeling work shows that the

driven state is associated with shortened temporal autocorrelations as well as a

decrease in the entropy of activity states in large-scale cortical networks [64].

Correspondingly, the increase in long-range temporal autocorrelations under

catecholaminergic modulation may be associated with an increase in entropy –

that is, a tendency to explore a larger set of cortical activity states. It is tempting

to link this to the prominent idea that high sustained noradrenaline levels promote

an exploratory mode of cortical computation and behavior [5].

Convergent evidence for catecholaminergic increase in cortical E/I ratio

Cortical circuits maintain a tight balance between excitation and inhibition, which

is largely preserved across contexts and levels of the cortical hierarchy [46,48].

However, even in the absence of changes in sensory input, neuromodulators

such as noradrenaline and acetylcholine can change the cortical E/I ratio [65,66].

The E/I ratio, in turn, shapes the computational properties of cortical circuits

[67,68], and thereby the behavior of the organism [37,44,69]. Substantial

evidence already points to significant changes in E/I ratio in schizophrenia and

autism [70–72]. Similar changes might be at play in other brain disorders as well

[73].

Our simulations indicated that the temporal correlation structure of neural

population activity, as measured with the scaling exponent α, is sensitive to

changes in E/I ratio, produced through synaptic gain modulation (see the white

line in Fig 8H). In both versions of our model, the neuromodulatory effects were

not perfectly symmetric (see the deviations of peak scaling exponents from main

diagonal in Fig 8H). While the latter effect was small and may be specific to the

details of the model, it remains possible that the subtle changes in scaling

exponents we observed were produced through symmetric gain modulations that

maintained the net E/I balance (i.e., along the main diagonal). However, two

additional lines of evidence converge on our conclusion that catecholamines (in

particular noradrenaline) boosted E/I ratio. First, in the same participants, the

catecholaminergic manipulation had a reliable effect on the perceptual switch rate,

which is also indicative of cortical E/I ratio [33,34,36]. Second, results invasive

rodent work also point to an increase in cortical E/I ratio under noradrenaline:

Noradrenaline was found to decrease spontaneous inhibition in auditory cortex

[66] and mediate a tonic depolarization of visual cortical neurons during

locomotion [20].

No evidence for donepezil effects on cortical or perceptual fluctuations

The absence of an effect of Donepezil on either perceptual fluctuations or long-

range temporal correlations of cortical activity may be due to the small dosage or

the single administration of the drug in our study. Even so, our donepezil

manipulation was sufficient to robustly change heart rate variability and, more

importantly, low-frequency power of cortical activity, an established marker of

593 cholinergic action in cortex [17,18,43,74]. The lack of the effects of donepezil on

594 perceptual fluctuations and cortical scaling behavior might also be due to the

595 specific properties of cholinergic action on the cortical net E/I ratio. Invasive

596 evidence indicates that acetylcholine can rapidly disinhibit pyramidal cells by

597 activating a chain of two inhibitory interneurons [21], a mechanism that may alter

598 E/I ratio mainly during stimulus-evoked responses [65]. By contrast,

599 noradrenaline also alters the levels of tonic inhibition of pyramidal cells occurring

600 spontaneously [66]. This might explain the dissociation between the effects of

601 atomoxetine and donepezil under the current steady-state conditions, which

602 excluded (or minimized) stimulus-evoked transients .

604 **Functional consequences of changes in net cortical E/I ratio**

605 We observed a selective increase in the rate of spontaneous perceptual

606 alternations under catecholaminergic adding to evidence that these dynamics are

607 under neuromodulatory control [75]. Such a change could be due to an increase

608 in cortical "noise" [33]. Future invasive studies should relate chatecholaminergic

609 changes in the variability of the spiking activity [76] of neurons contributing

610 directly to the contents of multi-stablke perception.

611  We suspect that an increase in cortical E/I ratio will have particularly

612 strong effects on behavior when affecting parietal and prefrontal cortical circuits

613 characterized by slow intrinsic timescales [30,32,77] and involved in persistent

614 activity during working memory and the slow accumulation of information over

615 time [69]. It is possible that the catecholaminergic effects on parietal cortex we

616 observed here reflects an increase in the recurrent excitation, which, is essential

617 for sustained processes such as working memory [78] as well as information

618 integration during decision-making [32,79]. Future work should assess this

619   through the use of tasks probing into network reverberation and information

620   accumulation in association cortex.

A control parameter for critical network dynamics

In our model, long-range temporal correlations in the fluctuations of neural mass

activity (i.e., activity summed across the entire local network) [26] and avalanches

within the neuronal network [50] jointly emerge at the same E/I ratio. Both

phenomena are commonly interpreted as hallmarks of "criticality" [26,29,50,80] –

a state of a complex dynamical system poised between order and chaos [81–83].

It has been proposed that the cortex operates in a narrow regime around

criticality [83,84], potentially optimizing its computational capacities [80,85–88]. A

number of reports showed that cortical dynamics may continuously vary around

the critical state [89–92], but the source of these fluctuations has, so far,

remained unknown. Here, we have identified catecholaminergic neuromodulation

as an endogenous factor controlling these spontaneous variations in critical

dynamics.

In complex systems, critical dynamics can emerge in a self-organized

fashion [81], or through an external control parameter that fine-tunes the system.

The tuning of temperature in the Ising model of spin magnetization [83] is a

common example for the latter case. It is tempting to speculate that

catecholaminergic tone serves as such a control parameter in the cerebral cortex.

Link between catecholaminergic effects on fluctuations in perception and

cortical mass activity

We here used two read-outs of catecholaminergic effects, constituting two distinct

expressions of the resulting changes in cortical circuit state. The envelope of

cortical alpha-band oscillations collapsed across large chunks of cortex is unlikely

to encode the contents of perception in the phenomenon studied here. The

perceived direction of 3D-motion, which fluctuates spontaneously, is encoded in

fine-grained patterns of neural population activity within motion-sensitive visual

cortical areas [39,93]. The power of alpha-band oscillations is a more global

feature of cortical population activity, which is likely insensitive to the fine-grained,

within-area patterns of neural population activity. The widespread release of

neuromodulators changes also the cortical circuit state, specifically E/I ratio, in a

widespread manner. Such changes, in turn, alter the highly specific (fine-grained)

interactions between percept-selective populations of visual cortical neurons that

give rise to the perceptual dynamics [33,34,36]. Thus, although both read-outs

likely tap into similar changes in global cortical circuit state, there is no one-to-to

mapping between them.

Limitations of the current modeling approach

While our model simulations provided important mechanistic insights, the model

has limitations that should be addressed in future work. First, different from the

MEG data, the power of alpha-band oscillations behaves similarly to the scaling

exponents in the model (Fig S8E). This is because the model oscillations emerge

from the same recurrent neuronal interactions within the patch that also shape

the long-range temporal correlations in the envelopes of the amplitude envelopes

of these oscillations. By contrast, in the brain, alpha-band power of local cortical

mass signals is likely affected by a variety of sources other than local circuits, for

instance alpha-frequency modulated input from the thalamus [94]. This might

lead to dissociations between changes in MEG power and long-range temporal

correlations of the power fluctuations which the model does not capture in its

present form. Second, the model lacks long-range excitatory connections, which

are prominent in the real cortex, and whose effects on the correlation structure of

cortical fluctuations are largely unknown.

Conclusion

The combined measurement of fluctuations in bistable perception as well as in

cortical mass activity under steady-state conditions provides an easily assessable,

multi-level read-out of pharmacological effects on cortical computation. In our

study, this read-out supported the idea that catecholamines boost the intrinsic

variability of perception and behavior, an effect that might be mediated by an

increase in the net E/I ratio in the visual cortical system. This read-out may be

useful for inferring changes in cortical E/I ratio in neuropsychiatric disorders, or in

their pharmacological treatment in future work.

METHODS

Pharmacological MEG experiment

Participants

30 healthy human participants (16 females, age range 20-36, mean 26.7)

participated in the study after informed consent. The study was approved by the

Ethical Committee responsible for the University Medical Center Hamburg-

Eppendorf. Two participants were excluded from analyses, one due to excessive

MEG artifacts, the other due to not completing all 3 recording sessions. Thus, we

report results from N=28 participants (15 females). In one of those participants,

one Task-counting run (during the Atomoxetine condition) was not recorded due

to a software problem with the data acquisition computer.

General design

We pharmacologically manipulated the levels of catecholamines (noradrenaline

and dopamine) and acetylcholine in a double-blind, randomized, placebo-

controlled, and cross-over experimental design (Fig 1A, B). Each participant

completed three experimental sessions, consisting of drug (or placebo) intake at

two time points, a waiting period of 3 hours, and an MEG recording. During each

MEG session, participants were seated on a chair inside a magnetically shielded

MEG chamber. Each session consisted of 6 runs of different tasks, each of which

was 10 minutes long and followed by breaks of variable duration.

Pharmacological intervention

We used the selective noradrenaline reuptake inhibitor atomoxetine (dose: 40

mg) to boost the levels of catecholamines, specifically noradrenaline and (in

prefrontal cortex) dopamine [10]. We used the cholinesterase inhibitor donepezil

(dose: 5 mg) to boost acetylcholine levels. Atomoxetine is a relatively selective

inhibitor of the noradrenaline transporter, which is responsible for the natural

reuptake of noradrenaline that has been released into the extracellular space.

Consequently, atomoxetine acts to increase the extracellular levels of

noradrenaline, an effect that has been confirmed experimentally in rats prefrontal

cortex [95]. The same study showed that atomoxetine also increases the

prefrontal levels of dopamine, which has a molecular structure very similar to the

one of noradrenaline and is, in fact, a direct precursor of noradrenaline.

Atomoxetine has smaller affinity to the serotonin transporter, and there are

discrepant reports about the quantitative relevance of these effects: While one

study found no increases in serotonin levels under atomoxetine [95], a recent

study reports a significant atomoxetine-related occupancy of the serotonin

transporter in non-human primates [96] at dosages which would correspond to

human dosages of 1.0 – 1.8 mg/kg. Note, that these dosages are substantially

higher than the administered dosage in this study (40 mg, independent of body

weight). It is, therefore, unclear to which extent our Atomoxetine condition

affected cortical serotonin levels.

Donepezil is a selective inhibitor of the enzyme acetylcholinesterase,

which breaks up all the extracellular acetylcholine to terminate its synaptic action.

Consequently, donepezil acts to increase the extracellular levels of acetylcholine.

Donepezil is also an agonist of the endoplasmatic $sigma_1$-receptor, which

modulates intracellular calcium signaling.

A mannitol-aerosil mixture was administered as placebo. All substances

were encapsulated identically in order to render them visually indistinguishable.

Peak plasma concentrations are reached ~3-4 hours after administration for

donepezil [97] and 1-2 hours after administration for atomoxetine [98],

respectively. We adopted the following procedure to account for these different

pharmacokinetics (Fig 1A): participants received two pills in each session, one 3

h and another 1.5 h before the start of MEG recording. In the Atomoxetine

condition, they first received a placebo pill (t = -3 h) followed by the atomoxetine

pill (t = -1.5 h). In the Donepezil condition, they first received the donepezil pill (t =

-3 h), followed by placebo (t = -1.5 h). In the Placebo condition, they received a

placebo at both time points. The half-life is ~ 5 h for atomoxetine [98] and ~ 82 h

for donepezil, respectively [97]. In order to allow plasma concentration levels to

return to baseline, the three recording sessions were scheduled at least 2 weeks

apart. This design ensured maximum efficacy of both pharmacological

manipulations, while effectively blinding participants as well as experimenters.

Stimuli and behavioral tasks

In each session, participants alternated between three different task conditions (2

runs à 10 minutes per condition) referred to as Fixation, Task-counting, and

Task-pressing in the following (Fig 1B). All conditions entailed overall constant

sensory input. Fixation and Task-counting also entailed no overt motor responses

and are, therefore, referred to as "steady-state" conditions in the following. We

used these steady-state conditions to quantify intrinsic fluctuations in cortical

activity. Task-pressing entailed motor responses and was used for reliable

quantification of perceptual dynamics. All instructions and stimuli were projected

onto a screen (distance: 60 cm) inside the MEG chamber. The individual

conditions are described as follows.

Fixation. Participants were asked to keep their eyes open and fixate a

green fixation dot (radius = 0.45° visual angle) presented in the center of an

otherwise gray screen. This is analogous to eyes-open measurements of

"resting-state" activity widely used in the literature on intrinsic cortical activity

fluctuations.

Task-counting. Participants viewed a seemingly rotating sphere giving rise

to the kinetic depth effect [99,100]: spontaneous changes in the perceived

rotation direction (Fig 1B). The stimulus subtended 21° of visual angle. It

consisted of 1000 dots (500 black and 500 white dots, radius: 0.18° of visual

angle) arranged on a circular aperture presented on a mean-luminance gray

background, with the green fixation dot in the center. In order to minimize tracking

eye movements, the sphere rotation was along the horizontal axis, either "forward"

(towards the observer) or "backward" (away from the observer), and the dot

density decreased along the horizontal axis towards the center of the stimulus.

Participants were instructed to count the number of perceived changes in rotation

direction and report the total number of perceived transitions at the end of the run.

Just like during Fixation, Task-counting minimized any external (sensory or

motor) transients. Subjects silently counted the alternations in perceived rotation

direction and verbally reported the total count after the end of the 10 minutes run.

Task-pressing. This condition was identical to Task-counting, except that

participants were instructed to press and hold one of two buttons with their index

finger to indicate the perceived rotation direction of the sphere. Thus, each

perceptual alternation was accompanied by a motor response leading to change

in the button state. This allowed for a more reliable quantification of participants'

perceptual dynamics. On two sessions (Atomoxetine condition), button presses

were not registered. Hence, the corresponding analyses were performed on 26

participants.

Data acquisition

MEG was recorded using a whole-head CTF 275 MEG system (CTF Systems,

Inc., Canada) at a sampling rate of 1200 Hz. In addition, eye movements and

pupil diameter were recorded with an MEG-compatible EyeLink 1000 Long

Range Mount system (SR Research, Osgoode, ON, Canada) at a sampling rate

of 1000 Hz. In addition, electrocardiogram (ECG) as well as vertical, horizontal

and radial EOG were acquired using Ag/AgCl electrodes (sampling rate 1200 Hz).

Data analysis

Eye data

Eye blinks were detected using the manufacturer's standard algorithm with

default settings. Saccades and microsaccades were detected using the saccade

detection algorithm described in [101], with a minimum saccade duration of 4

samples (= 4 ms) and a threshold velocity of 6. For 18 out of 28 participants, only

horizontal eye movements were recorded.

EOG data

EOG events (blinks and saccades) were extracted using semi-automatic artifact

procedures as implemented in FieldTrip [102]. In short, EOG traces were

bandpass filtered using a third-order butterworth filter (1 – 15 Hz) and the

resulting signal was z-scored. All time points where the resulting signal exceeded

a z-score of 4 were marked as an EOG event.

MEG data

Preprocessing. First, all data were cleaned of strong transient muscle artifacts

and squid jumps through visual inspection and manual as well as semi-automatic

artifact rejection procedures, as implemented in the FieldTrip toolbox for MATLAB

[102]. To this end, data segments contaminated by such artifacts (+/- 500 ms)

were discarded from the data (across all channels). Subsequently, data were

downsampled to 400 Hz split into low (2-40 Hz) and high (>40 Hz) frequency

components, using a 4th order (low- or high-pass) Butterworth filter. Both signal

components were separately submitted to independent component analysis [103]

using the FastICA algorithm [104]. Artifactual components (eye blinks/movements,

muscle artifacts, heartbeat and other extra-cranial artifacts) were identified based

on three established criteria [105]: power spectrum, fluctuation in signal variance

over time (in bins of 1s length), and topography. Artifact components were

reconstructed and subtracted from the raw signal and low- and high frequencies

were combined into a single data set. On average, 20 (+/- 14) artifact

components were identified for the low frequencies and 13 (+/- 7) artifactual

components were identified for the high frequencies.

Spectral analysis. Sensor-level spectral estimates (power spectra and cross

spectral density matrices) were computed by means of the multi taper method

using a sequence of discrete prolate Slepian tapers [106]. For the power

spectrum shown in Fig 1C, power spectra were computed using a window length

of 5s and a frequency smoothing of 2 Hz, yielding 19 orthogonal tapers. The

focus of this paper was on the fluctuations of the amplitude envelopes, rather

than on the (oscillatory) fluctuations of the carrier signals *per se*. The temporal

correlation structure of the amplitude envelope fluctuations of cortical activity

seems similar across different carrier frequency bands [29]. We focused on

amplitude envelope fluctuations in the alpha-band because (i) the cortical power

spectra exhibited a clearly discernible alpha-peak, which robustly modulated with

task, as expected from previous work [40] (Fig 1C); and (ii) the computational

model used to study the effect of synaptic gain modulation on cortical activity

fluctuations was tuned to produce alpha-band oscillations (see above and [49]).

Source reconstruction: general approach. The cleaned sensor level signals (N

sensors) were projected onto a grid consisting of $M = 3000$ voxels covering gray

matter of the entire brain (mean distance: 6.3 mm) using the exact low-resolution

brain electromagnetic tomography (eLORETA; [107] method.   The grid was

constructed from the ICBM152 template [108], covering gray matter across the

brain. The magnetic leadfield was computed, separately for each subject and

session, using a single shell head model constructed from the individual

structural MRI scans and the head position relative to the MEG sensors at the

beginning of the run [109]. In case no MRI was available (4 subjects), the

leadfield was computed from a standard MNI template brain transformed to an

estimate of the individual volume conductor using the measured fiducials (located

at the nasion, the left and the right ear).

In order to depict the source-level results, we interpolated the voxel-level

results onto the surface of the brain. Activations from structures distant to the

surface are not shown and were exponentially attenuated.

Source level estimates of amplitude envelopes and power. For comparing

amplitude envelope and power estimates between experimental conditions in

source space we aimed to select a single direction of the spatial filter for each

voxel across pharmacological conditions (i.e., MEG sessions), but separately for

Fixation and Task-Counting conditions. The rationale was to avoid filter-induced

biases in the comparisons between the pharmacological conditions, while

allowing that external task drive might systematically change the dipole

orientations.

To this end, we first computed the mean source-level cross-spectral

density matrix $C(r,f)$ for each frequency band, f, averaged across the three

MEG sessions, as follows:

$C(r,f) = \frac{1}{3}\sum_{i=1}^{3}\left(A_i^T(r)C_i(f)A_i(r)\right)$ (1)

whereby i indicated the MEG session, $C_i(f)$ was the (sensor-level) session- and

frequency-specific cross-spectral density matrix and A_i is the spatial filter for

session i. We then extracted the first eigenvector $u_1(r,f)$ of the session-average

matrix $C(r,f)$, by means of singular value decomposition, and computed the

unbiased filter selective for the dominant dipole orientation, $B_i(r,f)$, as:

$B_i(r,f) = A_i(r)u_1(r,f)$ (2)

This procedure ensures that, for each voxel, dipole orientation was

chosen such that power is maximized. Please note that this filter was now

frequency-specific, whereas the previous filters, $A_i(r)$, were not. To obtain

instantaneous estimates of source-level amplitudes, the sensor-level signal for

session i, $X_i(t)$, was band-pass filtered (using a finite impulse response filter)

and Hilbert-transformed, yielding a complex-valued signal $H_i(f, t)$ for each

frequency band. This signal was projected into source space through

multiplication with the unbiased spatial filter, $B_i(r, f)$, and the absolute value was

taken:

$Env_i(r, f, t) = |(H_i(f, t)\, B_i(r, f))|$ (3)

where $Env_i(r, f, t)$ was the estimated amplitude envelope time course of source

location r and frequency f. Next, for each session, unbiased source-level cross

spectral density estimates were obtained from the sensor-level cross-spectral

density matrix $C_i(f)$ and the frequency-specific, unbiased spatial filter $B_i(f)$. The

main diagonal of the resulting matrix contains source-level power estimates for all

source locations:

$S_i(f) = diag(B_i^T(f)_i C_i(f) B_i(f))$ (4)

These computations where repeated separately for the Task-counting and

Fixation conditions, session by session. The differences in amplitude envelope

fluctuations and power estimates between pharmacological and task conditions

reported in this paper were robust with respect to the specifics of the analysis

approach. In particular, we obtained qualitatively similar pharmacological effects

in sensor space, as reported in an earlier conference abstract [110].

Detrended fluctuation analysis. The source-level amplitude envelopes

$Env_i(r, f, t)$ were submitted to detrended fluctuation analysis [111,112] in order to

quantify long-range temporal correlations. Detrended fluctuation analysis

quantifies the power law scaling of the fluctuation (root-mean-square) of a locally

detrended, cumulative signal with time-window length. Different from the analysis

of the more widely known autocorrelation function [30,77], detrended fluctuation

analysis provides robust estimates of the autocorrelation structure for stationary

and non-stationary time series. The procedure of the detrended fluctuation

analysis is illustrated in Fig 2.

For simplicity, in the following, we re-write the amplitude envelope

$Env_i(r, f, t)$ as x of length T. First, we computed the cumulative sum of the

demeaned x, (Fig 2B):

$X(t) = \sum_{t'=1}^{t}(x(t') - \langle x \rangle)$ (5)

where t' and t denote single time points up to length T. The cumulative signal X

was then cut into $i = 1 \dots k$ segments Y_i of length N (overlap: 50%), where $k =$

$floor[(T - N)/(0.5\,N)]$ (Fig 2B, top). Within each segment Y_i of equal length N,

the linear trend Y_{i_trend} (least squares fit) was subtracted from Y_i (Fig 2B, bottom,

blue vs. red lines), and the root-mean-square fluctuation for a given segment was

computed as:

$F_{N\_i} = \left[\frac{1}{N}\sum_{n=1}^{N}(Y_i(n) - Y_{i\_trend}(n))^2\right]^{\frac{1}{2}}$ (6)

where n indicates the individual time points. The fluctuation was computed for all

k segments of equal length N and the average fluctuation was obtained through:

924                    $<F_N> = \frac{1}{k}\sum_{i=1}^{k}F_{N\_i}$ (7)

The procedure was repeated for 15 different logarithmically spaced window

lengths N, ranging from 3 s to 50 s, which yields a fluctuation function (Fig 2C).

As expected for scale-free time series (103), this fluctuation function follows a

power-law of the form:

$<F_N> \propto N^{\alpha}$ (8)

The "scaling exponent" α was computed through a linear regression fit in log-log

coordinates (Fig 2C). The longest and shortest window lengths were chosen

according to guidelines provided in [112].

A scaling exponent of $\alpha \sim= 0.5$ indicates a temporally uncorrelated ("white

noise") process. Scaling exponents between $0.5 < \alpha < 1$ are indicative of scale-

free behavior and long-range temporal correlations [112], whereas exponents of

$\alpha < 0.5$ indicate long-range anti-correlations ("switching behavior") and $\alpha > 1$ are

indicative of an unbounded process [112]. The scaling exponents for alpha-band

MEG amplitude envelopes estimated in this study ranged (across experimental

conditions, MEG sensors and participants) from 0.40 and 1.04, with 99.4% of all

estimates in the range from 0.5 to 1. This is indicative of scale-free behavior and

consistent with previous human MEG work [26–29,42,113].

Relationship between measures of cortical variability. Scale-free behavior of

neural time series has also been quantified via analysis of the power spectrum

[24,25]. There is a straightforward relationship between both approaches, which

we explain below, to help appreciate our results in the context of these previous

studies. The power spectrum of the amplitude envelope of cortical activity is

typically well approximated by the power law $p(f) \propto f^{-\beta}$, where β is referred to

as the power-law exponent (Fig 2D). For power-law decaying autocorrelations,

the relationship between the power-law exponent β and the scaling exponent α

(estimated through DFA) of a time series is:

$\beta = 2\alpha - 1$ (9)

Analysis of ECG data. ECG data were used to analyze two measures of

peripheral autonomic activity: average heart rate and heart rate variability. For

both measures, we used an adaptive threshold to detect the R-peak of each

QRS-complex in the ECG. Heart rate was then computed by dividing the total

number of R-components by time. Heart rate variability was quantified by means

of the detrended fluctuations analysis described for MEG above, but now applied

to the time series of the intervals between successive R-peaks [28,29]. In line

with the MEG analyses, we used windows ranging from 3 to 50 heartbeats

(roughly corresponding to 3–50 s).

Statistical tests

Statistical comparisons of all dependent variables between conditions were,

unless stated otherwise, performed using paired t-tests.

Null effects are difficult to interpret using regular null hypothesis

significance testing. The Bayes Factor addresses this problem by quantifying the

strength of the support for the null hypothesis over the alternative hypothesis

provided by the data, taking effect size into account. Wherever null effects were

conceptually important, results obtained from a regular (paired) t-test [114] and

Pearson correlations [115] were converted into corresponding Bayes Factors.

To map significant changes of scaling exponents α across the brain, we

computed a non-parametric permutation test based on spatial clustering

[116,117]. This procedure has been shown to reliably control for Type I errors

arising from multiple comparisons. First, a paired t-test was performed to identify

voxels with significant changes (voxel with $p < 0.05$). Subsequently, significant

voxels are combined into clusters based on their spatial adjacency. Here, a voxel

was only included into a cluster when it had at least two significant neighbors.

Subsequently, the t-values of all voxels comprising a cluster were summed,

which yields a cluster statistic (i.e., a cluster t-value) for each identified cluster.

Next, a randomization null distribution was computed using a permutation

procedure (N = 10.000 permutations). On each permutation, the experimental

labels (i.e., the pharmacological conditions) were randomly re-assigned within

participants and the aforementioned procedure was repeated. For each iteration,

the maximum cluster statistic was determined and a distribution of maximum

cluster statistics was generated. Eventually, the cluster statistic of all empirical

clusters was compared to the values obtained from the permutation procedure.

All voxels comprising a cluster with a cluster statistic smaller than 2.5% or larger

than 97.5% of the permutation distribution were labeled significant, corresponding

to a corrected threshold of $\alpha = 0.05$ (two-sided).

Model simulations

To simulate the effects of synaptic gain modulation on cortical activity fluctuations,

we extended a previously described computational model of a local cortical patch

[49] by means of multiplicative modulation of synaptic gain. All features of the

model were identical to those of the model by [49], unless stated otherwise. The

model consisted of 2500 integrate-and-fire neurons (75% excitatory, 25%

inhibitory) with local connectivity within a square (width = 7 units) and a

connection probability that decayed exponentially with distance (Fig 8A). The

dynamics of the units were governed by:

$I_i = I_i + \sum_j N_{ij} W_{ij} S_j$ (10)

$\tau_i \frac{dI_i}{dt} = I_0 - I_i$ (11)

where subscripts i, j indicated different units, N_{ij} was a multiplicative gain factor,

W_{ij} were the connection weights between two units, and S_j a binary spiking

vector representing whether unit *j* did or did not spike on the previous time step,

and $I_0 = 0$. For all simulations reported in this paper, we optimized the

connection weights using Bonesa [118], a parameter tuning algorithm, such that

the network exhibited alpha-band oscillations, long-range temporal correlations,

and neuronal avalanches (see below). The optimized values for the connection

weights   were $W_{EE} = 0.0085$, $W_{IE} = 0.0085$, $W_{EI} = -0.569$ and $W_{II} = -2$

whereby subscript E indicated excitatory, subscript I indicated inhibitory, and the

first and second subscript referred to the receiving and sending unit, respectively.

On each time step (dt = 1 ms), I_i was updated for each unit i, with the

summed input from all other (connected) units j and scaled by a time constant

$\tau_i = 9\,\text{ms}$, which was the same for excitatory and inhibitory units. The probability

of a unit generating a spike output was given by:

$P_{si} = P_{si} + I_i$ (12)

$\tau_P \frac{dP_{si}}{dt} = P_0 - P_{si}$ (13)

with the time constant for excitatory units $\tau_P = 6\,ms$ and for inhibitory $\tau_P = 12\,ms$.

P_0 was the background spiking probability, with $P_0(exc.) = 0.000001\,[1/ms]$ and

$P_0(inh.) = 0\,[1/ms]$. For each time step, it was determined whether a unit did or

did not spike. If it did, the probability of that unit spiking was reset to

$P_r(excitatory) = -2\,[1/ms]$ and $P_r(inhibitory) = -20\,[1/ms]$.

We used this model to analyze the dependency of two quantities on E/I

ratio: (i) the power-law scaling of the distributions of the sizes of neuronal

avalanches [50] estimated in terms of the kappa-index κ which quantifies the

difference between an empirically observed event size distribution and a

theoretical reference power-law distribution with a power-law exponent -1.5 [86],

and (ii) the scaling behavior (scaling exponent α) of the amplitude envelope

fluctuations of the model's local field potential. To this end, we summed the

activity across all (excitatory and inhibitory) neurons to obtain a proxy of the local

field potential. We band-pass filtered the local field potential in the alpha-band (8–

12 Hz) and computed long-range temporal correlations in the alpha-band

amplitude envelopes following the procedure described above (see *Detrended*

fluctuation analysis of MEG data), using windows sizes ranging from 5 s to 30 s.

In order to assess the influence of structural excitatory and inhibitory

connectivity on network dynamics (Figs 4D-F), we varied the percentage of units

(excitatory and inhibitory) a given excitatory or inhibitory unit connects to within a

local area (7 units x 7 units; Fig 8A). These percentages were varied

independently for excitatory and inhibitory units with a step size of 2.5%.

The gain factor N_{ij} was the main difference to the model described by [49].

It was introduced to simulate the effects of neuromodulation on synaptic

interactions in the cortical network [37]. For this, we kept all the structural

parameters fixed (42.5% excitatory connectivity, 75% inhibitory connectivity;

small square in Figs 4D-F), in a range where the model exhibits both robust long-

range temporal correlations as well as neuronal avalanches. Note that any other

combination of parameters would yield similar results, as long as the model

exhibits these two phenomena. From the chosen starting point, we systematically

varied the synaptic gain factors, in two different ways. In the first version, we only

varied N_{EE} and N_{IE} to dynamically modulate the circuit's net E/I ratio (Fig 8B), in a

way consistent with recent modeling of the effects of E/I ratio on a cortical circuit

for perceptual decision-making [44]. In the second version, we varied N_{EE}, N_{IE},

and N_{EI} (Fig S8A). Here, N_{EI} was modulated independently from N_{EE}, and N_{IE},

which in turn were co-modulated by the same factor.

Per parameter combination, we ran 10 simulations, using the Brian2

spiking neural networks simulator [119]. Each simulation was run for 1000

seconds, with a random initialization of the network structure and the probabilistic

spiking.

1060

ACKNOWLEDGEMENTS

The authors thank Christiane Reissmann for help with the data collection, as well

as Sander Nieuwenhuis and Peter Murphy for helpful comments on the

manuscript. This work was supported by the German Research Foundation

(DFG): Heisenberg Professorship DO 1240/3-1 (to T.H.D.), and the Collaborative

Research Center SFB 936 (Projects A2/A3, A7, Z3, to A.K.E., T.H.D., G.N.,

respectively); by the BMBF (Project 161A130, to A.K.E.); by the EU (ERC-2010-

AdG-269716, to A.K.E.), by a grant from the state of Hamburg (CROSS FV25, to

A.K.E.); and by the Netherlands Organization for Scientific Research (NWO,

dossiernummer 406-15-256 to K.L.-H. and A.-E.A.)

1071

AUTHOR CONTRIBUTIONS

Conceptualization: T.P., A.K.E., and T.H.D.; Experimental design: T.P. and

T.H.D.; Model design: T.P., A-E.A., K.L-H., and T.H.D.; Investigation: T.P.;

Formal analysis: T.P.; Model simulations: A.-E.A.; Writing - Original draft: T.P.

and T.H.D.; Writing – Review & Editing: T.P., A-E.A., G.N., A.K.E., K.L-H., and

T.H.D. - Funding Acquisition: K.L-H., A.K.E., and T.H.D.; Supervision: G.N., K.L-

H., and T.H.D.

1079

1080 **COMPTETING FINANCIAL INTERESTS**

1081 The authors declare no competing financial interests.

1082

REFERENCES

1. Harris KD, Thiele A. Cortical state and attention. Nat Rev Neurosci.
2011;12: 509–523. doi:10.1038/nrn3084

2.   McGinley MJ, Vinck M, Reimer J, Batista-Brito R, Zagha E, Cadwell CR, et
al. Waking State: Rapid Variations Modulate Neural and Behavioral
Responses. Neuron. 2015;87: 1143–1161.
doi:10.1016/j.neuron.2015.09.012

3.   Guedj C, Monfardini E, Reynaud AJ, Farnè A, Meunier M, Hadj-Bouziane F.
Boosting Norepinephrine Transmission Triggers Flexible Reconfiguration
of Brain Networks at Rest. Cereb Cortex. 2016; doi:10.1093/cercor/bhw262

4.   van den Brink RL, Pfeffer T, Warren CM, Murphy PR, Tona K-D, van der
Wee NJA, et al. Catecholaminergic Neuromodulation Shapes Intrinsic MRI
Functional Connectivity in the Human Brain. J Neurosci. 2016;36: 7865–
7876. doi:10.1523/JNEUROSCI.0744-16.2016

5.   Aston-Jones G, Cohen JD. An integrative theory of locus coeruleus-
norepinephrine function: adaptive gain and optimal performance. Annu Rev
Neurosci. 2005;28: 403–450.
doi:10.1146/annurev.neuro.28.061604.135709

6.   Yu AJ, Dayan P. Uncertainty, neuromodulation, and attention. Neuron.
2005;46: 681–692. doi:10.1016/j.neuron.2005.04.026

7.   Nelson A, Mooney R. The Basal Forebrain and Motor Cortex Provide
Convergent yet Distinct Movement-Related Inputs to the Auditory Cortex.
Neuron. 2016;90: 635–48. doi:10.1016/j.neuron.2016.03.031

8.   de Gee JW, Colizoli O, Kloosterman NA, Knapen T, Nieuwenhuis S,
Donner TH. Dynamic modulation of decision biases by brainstem arousal
systems. eLife. 2017;6. doi:10.7554/eLife.23232

9.   Berridge CW. Noradrenergic modulation of arousal. Brain Res Rev.
2008;58: 1–17. doi:10.1016/j.brainresrev.2007.10.013

10.   Robbins TW, Arnsten AFT. The neuropsychopharmacology of fronto-
executive function: monoaminergic modulation. Annu Rev Neurosci.
2009;32: 267–287. doi:10.1146/annurev.neuro.051508.135535

11.   Lee S-H, Dan Y. Neuromodulation of brain states. Neuron. 2012;76: 209–
222.

12.   Froemke RC. Plasticity of Cortical Excitatory-Inhibitory Balance. Annu Rev
Neurosci. 2015;38: 195–219. doi:10.1146/annurev-neuro-071714-034002

13.   Glimcher PW. Indeterminacy in brain and behavior. Annu Rev Psychol.
2005;56: 25–56. doi:10.1146/annurev.psych.55.090902.141429

14.   Renart A, Machens CK. Variability in neural activity and behavior. Curr
Opin Neurobiol. 2014;25: 211–220. doi:10.1016/j.conb.2014.02.013

15.   Frank MJ, Doll BB, Oas-Terpstra J, Moreno F. Prefrontal and striatal
dopaminergic genes predict individual differences in exploration and
exploitation. Nat Neurosci. 2009;12: 1062–1068. doi:10.1038/nn.2342

16. Moreno-Bote R, Knill DC, Pouget A. Bayesian sampling in visual
perception. Proc Natl Acad Sci. 2011;108: 12491–12496.
doi:10.1073/pnas.1101430108

17. Pinto L, Goard MJ, Estandian D, Xu M, Kwan AC, Lee S-HH, et al. Fast
modulation of visual perception by basal forebrain cholinergic neurons.
2013;16: 1857–63. doi:10.1038/nn.3552

18. Chen N, Sugihara H, Sur M. An acetylcholine-activated microcircuit drives
temporal dynamics of cortical activity. Nat Neurosci. 2015;18: 892–902.
doi:10.1038/nn.4002

19. Minces V, Pinto L, Dan Y, Chiba AA. Cholinergic shaping of neural
correlations. Proc Natl Acad Sci. 2017;114: 5725–5730.
doi:10.1073/pnas.1621493114

20. Polack P-O, Friedman J, Golshani P. Cellular mechanisms of brain state-
dependent gain modulation in visual cortex. Nat Neurosci. 2013;16: 1331–
1339. doi:10.1038/nn.3464

21. Fu Y, Tucciarone JM, Espinosa JS, Sheng N, Darcy DP, Nicoll RA, et al. A
cortical circuit for gain control by behavioral state. Cell. 2014;156: 1139–
1152. doi:10.1016/j.cell.2014.01.050

22. Fox MD, Raichle ME. Spontaneous fluctuations in brain activity observed
with functional magnetic resonance imaging. Nat Rev Neurosci. 2007;8:
700–711. doi:10.1038/nrn2201

23. Deco G, Jirsa VK, McIntosh AR. Emerging concepts for the dynamical
organization of resting-state activity in the brain. Nat Rev Neurosci.
2011;12: 43–56. doi:10.1038/nrn2961

24. Miller KJ, Sorensen LB, Ojemann JG, den Nijs M. Power-law scaling in the
brain surface electric potential. PLoS Comput Biol. 2009;5: e1000609.
doi:10.1371/journal.pcbi.1000609

25. He BJ, Zempel JM, Snyder AZ, Raichle ME. The temporal structures and
functional significance of scale-free brain activity. Neuron. 2010;66: 353–
369. doi:10.1016/j.neuron.2010.04.020

26. Linkenkaer-Hansen K, Nikouline VV, Palva JM, Ilmoniemi RJ. Long-range
temporal correlations and scaling behavior in human brain oscillations. J
Neurosci Off J Soc Neurosci. 2001;21: 1370–1377.

27. He BJ. Scale-Free Properties of the Functional Magnetic Resonance
Imaging Signal during Rest and Task. J Neurosci. 2011;31: 13786–13795.
doi:10.1523/JNEUROSCI.2111-11.2011

28. Palva JM, Zhigalov A, Hirvonen J, Korhonen O, Linkenkaer-Hansen K,
Palva S. Neuronal long-range temporal correlations and avalanche
dynamics are correlated with behavioral scaling laws. Proc Natl Acad Sci.
2013;110: 3585–3590. doi:10.1073/pnas.1216855110

29. Zhigalov A, Arnulfo G, Nobili L, Palva S, Palva JM. Relationship of fast- and slow-timescale neuronal dynamics in human MEG and SEEG. J Neurosci Off J Soc Neurosci. 2015;35: 5385–5396. doi:10.1523/JNEUROSCI.4880-14.2015

30. Honey CJ, Thesen T, Donner TH, Silbert LJ, Carlson CE, Devinsky O, et al. Slow Cortical Dynamics and the Accumulation of Information over Long Timescales. Neuron. 2012;76: 423–434. doi:10.1016/j.neuron.2012.08.011

31. Donner TH, Sagi D, Bonneh YS, Heeger DJ. Retinotopic Patterns of Correlated Fluctuations in Visual Cortex Reflect the Dynamics of Spontaneous Perceptual Suppression. J Neurosci. 2013;33: 2188–2198. doi:10.1523/JNEUROSCI.3388-12.2013

32. Chaudhuri R, Knoblauch K, Gariel M-A, Kennedy H, Wang X-J. A Large-Scale Circuit Mechanism for Hierarchical Dynamical Processing in the Primate Cortex. Neuron. 2015;88: 419–431. doi:10.1016/j.neuron.2015.09.008

33. Moreno-Bote R, Rinzel J, Rubin N. Noise-induced alternations in an attractor network model of perceptual bistability. J Neurophysiol. 2007;98: 1125–1139. doi:10.1152/jn.00116.2007

34. Noest AJ, van Ee R, Nijs MM, van Wezel RJA. Percept-choice sequences driven by interrupted ambiguous stimuli: A low-level neural model. J Vis. 2007;7: 10. doi:10.1167/7.8.10

35. Deco G, Romo R. The role of fluctuations in perception. Trends Neurosci. 2008;31: 591–598. doi:10.1016/j.tins.2008.08.007

36. van Loon AM, Knapen T, Scholte HS, St. John-Saaltink E, Donner TH, Lamme VAF. GABA Shapes the Dynamics of Bistable Perception. Curr Biol. 2013;23: 823–827. doi:10.1016/j.cub.2013.03.067

37. Eckhoff P, Wong-Lin KF, Holmes P. Optimality and Robustness of a Biophysical Decision-Making Model under Norepinephrine Modulation. J Neurosci. 2009;29: 4301–4311. doi:10.1523/JNEUROSCI.5024-08.2009

38. Parker AJ, Krug K, Cumming BG. Neuronal activity and its links with the perception of multi-stable figures. Philos Trans R Soc Lond B Biol Sci. 2002;357: 1053–1062. doi:10.1098/rstb.2002.1112

39. Brouwer GJ, van Ee R. Visual cortex allows prediction of perceptual states during ambiguous structure-from-motion. J Neurosci Off J Soc Neurosci. 2007;27: 1015–1023. doi:10.1523/JNEUROSCI.4593-06.2007

40. Donner TH, Siegel M. A framework for local cortical oscillation patterns. Trends Cogn Sci. 2011;15: 191–199. doi:10.1016/j.tics.2011.03.007

41. Linkenkaer-Hansen K, Smit DJA, Barkil A, van Beijsterveldt TEM, Brussaard AB, Boomsma DI, et al. Genetic contributions to long-range temporal correlations in ongoing oscillations. J Neurosci Off J Soc Neurosci. 2007;27: 13882–13889. doi:10.1523/JNEUROSCI.3083-07.2007

42. Linkenkaer-Hansen K, Nikulin VV, Palva S, Ilmoniemi RJ, Palva JM.
Prestimulus oscillations enhance psychophysical performance in humans.
J Neurosci Off J Soc Neurosci. 2004;24: 10186–10190.
doi:10.1523/JNEUROSCI.2584-04.2004

43. Bauer M, Kluge C, Bach D, Bradbury D, Heinze HJ, Dolan RJ, et al.
Cholinergic enhancement of visual attention and neural oscillations in the
human brain. Curr Biol CB. 2012;22: 397–402.
doi:10.1016/j.cub.2012.01.022

44. Lam NH, Borduqui T, Hallak J, Roque AC, Anticevic A, Krystal JH, et al.
Effects of Altered Excitation-Inhibition Balance on Decision Making in a
Cortical Circuit Model. bioRxiv. 2017; doi:http://dx.doi.org/10.1101/100347

45. van Vreeswijk C, Sompolinsky H. Chaos in neuronal networks with
balanced excitatory and inhibitory activity. Science. 1996;274: 1724–1726.

46. Shadlen MN, Newsome WT. The variable discharge of cortical neurons:
implications for connectivity, computation, and information coding. J
Neurosci Off J Soc Neurosci. 1998;18: 3870–3896.

47. Okun M, Lampl I. Balance of excitation and inhibition. Scholarpedia.
2009;4: 7467. doi:10.4249/scholarpedia.7467

48. Isaacson JS, Scanziani M. How inhibition shapes cortical activity. Neuron.
2011;72: 231–243. doi:10.1016/j.neuron.2011.09.027

49. Poil S-S, Hardstone R, Mansvelder HD, Linkenkaer-Hansen K. Critical-
State Dynamics of Avalanches and Oscillations Jointly Emerge from
Balanced Excitation/Inhibition in Neuronal Networks. J Neurosci. 2012;32:
9817–9823. doi:10.1523/JNEUROSCI.5990-11.2012

50. Beggs JM, Plenz D. Neuronal avalanches in neocortical circuits. J
Neurosci Off J Soc Neurosci. 2003;23: 11167–11177.

51. Servan-Schreiber D, Printz H, Cohen J. A network model of catecholamine
effects: gain, signal-to-noise ratio, and behavior. Science. 1990;249: 892–
895. doi:10.1126/science.2392679

52. Haider B, Häusser M, Carandini M. Inhibition dominates sensory
responses in the awake cortex. Nature. 2013;493: 97–100.
doi:10.1038/nature11665

53. Adesnik H. Synaptic Mechanisms of Feature Coding in the Visual Cortex of
Awake Mice. Neuron. 2017;95: 1147–1159.e4.
doi:10.1016/j.neuron.2017.08.014

54. Sara SJ. The locus coeruleus and noradrenergic modulation of cognition.
Nat Rev Neurosci. 2009;10: 211–223. doi:10.1038/nrn2573

55. Morrison JH, Foote SL. Noradrenergic and serotoninergic innervation of
cortical, thalamic, and tectal visual structures in Old and New World
monkeys. J Comp Neurol. 1986;243: 117–138.
doi:10.1002/cne.902430110

56.   Pennartz CM. The ascending neuromodulatory systems in learning by
reinforcement: comparing computational conjectures with experimental
findings. Brain Res Brain Res Rev. 1995;21: 219–245.

57.   Roelfsema PR, van Ooyen A, Watanabe T. Perceptual learning rules
based on reinforcers and attention. Trends Cogn Sci. 2010;14: 64–71.
doi:10.1016/j.tics.2009.11.005

58.   Salgado H, Treviño M, Atzori M. Layer- and area-specific actions of
norepinephrine on cortical synaptic transmission. Brain Res. 2016;1641:
163–176. doi:10.1016/j.brainres.2016.01.033

59.   Ramos BP, Arnsten AFT. Adrenergic pharmacology and cognition: focus
on the prefrontal cortex. Pharmacol Ther. 2007;113: 523–536.
doi:10.1016/j.pharmthera.2006.11.006

60.   Crochet S, Poulet JFA, Kremer Y, Petersen CCH. Synaptic Mechanisms
Underlying Sparse Coding of Active Touch. Neuron. 2011;69: 1160–1175.
doi:10.1016/j.neuron.2011.02.022

61.   Zhou M, Liang F, Xiong XR, Li L, Li H, Xiao Z, et al. Scaling down of
balanced excitation and inhibition by active behavioral states in auditory
cortex. Nat Neurosci. 2014;17: 841–850. doi:10.1038/nn.3701

62.   Swadlow HA. Thalamocortical control of feed-forward inhibition in awake
somatosensory "barrel" cortex. Philos Trans R Soc B Biol Sci. 2002;357:
1717–1727. doi:10.1098/rstb.2002.1156

63.   Kepecs A, Fishell G. Interneuron cell types are fit to function. Nature.
2014;505: 318–326. doi:10.1038/nature12983

64.   Ponce-Alvarez A, He BJ, Hagmann P, Deco G. Task-Driven Activity
Reduces the Cortical Activity Space of the Brain: Experiment and Whole-
Brain Modeling. Graham LJ, editor. PLOS Comput Biol. 2015;11:
e1004445. doi:10.1371/journal.pcbi.1004445

65.   Froemke RC, Merzenich MM, Schreiner CE. A synaptic memory trace for
cortical receptive field plasticity. Nature. 2007;450: 425–429.
doi:10.1038/nature06289

66.   Martins ARO, Froemke RC. Coordinated forms of noradrenergic plasticity
in the locus coeruleus and primary auditory cortex. Nat Neurosci. 2015;18:
1483–1492. doi:10.1038/nn.4090

67.   Murphy BK, Miller KD. Multiplicative gain changes are induced by
excitation or inhibition alone. J Neurosci. 2003;23: 10040–10051.

68.   Denève S, Machens CK. Efficient codes and balanced networks. Nat
Neurosci. 2016;19: 375–382. doi:10.1038/nn.4243

69.   Wang X-J. Decision making in recurrent neuronal circuits. Neuron.
2008;60: 215–234. doi:10.1016/j.neuron.2008.09.034

70. Yizhar O, Fenno LE, Prigge M, Schneider F, Davidson TJ, O'Shea DJ, et al.
Neocortical excitation/inhibition balance in information processing and
social dysfunction. Nature. 2011;477: 171–178. doi:10.1038/nature10360

71. Lisman J. Excitation, inhibition, local oscillations, or large-scale loops: what
causes the symptoms of schizophrenia? Curr Opin Neurobiol. 2012;22:
537–544. doi:10.1016/j.conb.2011.10.018

72. Nelson SB, Valakh V. Excitatory/Inhibitory Balance and Circuit
Homeostasis in Autism Spectrum Disorders. Neuron. 2015;87: 684–698.
doi:10.1016/j.neuron.2015.07.033

73. Fuchs T, Jefferson SJ, Hooper A, Yee P-H, Maguire J, Luscher B.
Disinhibition of somatostatin-positive GABAergic interneurons results in an
anxiolytic and antidepressant-like brain state. Mol Psychiatry. 2017;22:
920–930. doi:10.1038/mp.2016.188

74. Eggermann E, Kremer Y, Crochet S, Petersen CCH. Cholinergic Signals in
Mouse Barrel Cortex during Active Whisker Sensing. Cell Rep. 2014;9:
1654–1660. doi:10.1016/j.celrep.2014.11.005

75. Carter OL, Pettigrew JD, Hasler F, Wallis GM, Liu GB, Hell D, et al.
Modulating the rate and rhythmicity of perceptual rivalry alternations with
the mixed 5-HT2A and 5-HT1A agonist psilocybin. Neuropsychopharmacol
Off Publ Am Coll Neuropsychopharmacol. 2005;30: 1154–1162.
doi:10.1038/sj.npp.1300621

76. Noudoost B, Moore T. Control of visual cortical signals by prefrontal
dopamine. Nature. 2011;474: 372–375. doi:10.1038/nature09995

77. Murray JD, Bernacchia A, Freedman DJ, Romo R, Wallis JD, Cai X, et al.
A hierarchy of intrinsic timescales across primate cortex. Nat Neurosci.
2014;17: 1661–1663. doi:10.1038/nn.3862

78. Wang X-J. Synaptic reverberation underlying mnemonic persistent activity.
2001;24. doi:10.1016/s0166-2236(00)01868-3

79. Wang X-JJ. Probabilistic decision making by slow reverberation in cortical
circuits. 2002;36: 955–68. doi:10.1016/s0896-6273(02)01092-9

80. Beggs JM. The criticality hypothesis: how local cortical networks might
optimize information processing. Philos Transact A Math Phys Eng Sci.
2008;366: 329–343. doi:10.1098/rsta.2007.2092

81. Bak P, Tang C, Wiesenfeld K. Self-organized criticality: An explanation of
the 1/f noise. Phys Rev Lett. 1987;59: 381–384.
doi:10.1103/PhysRevLett.59.381

82. Bak P. How nature works: the science of self-organized criticality [Internet].
New York, NY, USA: Copernicus; 1996. Available:
http://catalog.hathitrust.org/api/volumes/oclc/34623628.html

83. Chialvo DR. Emergent complex neural dynamics. Nat Phys. 2010;6: 744–
750. doi:10.1038/nphys1803

84.   Hesse J, Gross T. Self-organized criticality as a fundamental property of
1328   neural   systems.   Front   Syst   Neurosci.   2014;8:   166.
1329   doi:10.3389/fnsys.2014.00166

1330   85.   Kinouchi O, Copelli M. Optimal dynamical range of excitable networks at
1331   criticality. Nat Phys. 2006;2: 348–351. doi:10.1038/nphys289

1332   86.   Shew WL, Yang H, Petermann T, Roy R, Plenz D. Neuronal avalanches
1333   imply maximum dynamic range in cortical networks at criticality. J Neurosci
1334   Off   J   Soc   Neurosci.   2009;29:   15595–15600.
1335   doi:10.1523/JNEUROSCI.3864-09.2009

1336   87.   Shew W, Yang H, Yu S, Roy R. Information capacity and transmission are
1337   maximized in balanced cortical networks with neuronal avalanches. J ….
1338   2011;

1339   88.   Shriki O, Yellin D. Optimal Information Representation and Criticality in an
1340   Adaptive Sensory Recurrent Neuronal Network. PLoS Comput Biol.
1341   2016;12: e1004698. doi:10.1371/journal.pcbi.1004698

1342   89.   Priesemann V, Valderrama M, Wibral M, Le Van Quyen M. Neuronal
1343   avalanches differ from wakefulness to deep sleep--evidence from
1344   intracranial depth recordings in humans. PLoS Comput Biol. 2013;9:
1345   e1002985. doi:10.1371/journal.pcbi.1002985

1346   90.   Arviv O, Goldstein A, Shriki O. Near-Critical Dynamics in Stimulus-Evoked
1347   Activity of the Human Brain and Its Relation to Spontaneous Resting-State
1348   Activity. J Neurosci Off J Soc Neurosci. 2015;35: 13927–13942.
1349   doi:10.1523/JNEUROSCI.0477-15.2015

1350   91.   Fagerholm ED, Lorenz R, Scott G, Dinov M, Hellyer PJ, Mirzaei N, et al.
1351   Cascades and Cognitive State: Focused Attention Incurs Subcritical
1352   Dynamics.   J   Neurosci.   2015;35:   4626–4634.
1353   doi:10.1523/JNEUROSCI.3694-14.2015

1354   92.   Shew WL, Clawson WP, Pobst J, Karimipanah Y, Wright NC, Wessel R.
1355   Adaptation to sensory input tunes visual cortex to criticality. Nat Phys.
1356   2015;11: 659–663. doi:10.1038/nphys3370

1357   93.   Krug K, Cicmil N, Parker AJ, Cumming BG. A Causal Role for V5/MT
1358   Neurons Coding Motion-Disparity Conjunctions in Resolving Perceptual
1359   Ambiguity. 2013;23: 1454–9. doi:10.1016/j.cub.2013.06.023

1360   94.   Bollimunta A, Mo J, Schroeder CE, Ding M. Neuronal mechanisms and
1361   attentional modulation of corticothalamic α oscillations. J Neurosci Off J
1362   Soc   Neurosci.   2011;31:   4935–4943.   doi:10.1523/JNEUROSCI.5580-
1363   10.2011

1364   95.   Bymaster FP, Katner JS, Nelson DL, Hemrick-Luecke SK, Threlkeld PG,
1365   Heiligenstein JH, et al. Atomoxetine increases extracellular levels of
1366   norepinephrine and dopamine in prefrontal cortex of rat: a potential
1367   mechanism   for   efficacy   in   attention   deficit/hyperactivity   disorder.

Neuropsychopharmacol Off Publ Am Coll Neuropsychopharmacol.
2002;27: 699–711. doi:10.1016/S0893-133X(02)00346-9

96. Ding Y-S, Naganawa M, Gallezot J-D, Nabulsi N, Lin S-F, Ropchan J, et al.
Clinical doses of atomoxetine significantly occupy both norepinephrine and
serotonin transports: Implications on treatment of depression and ADHD.
NeuroImage. 2014;86: 164–171. doi:10.1016/j.neuroimage.2013.08.001

97. Tiseo, Rogers, Friedhoff. Pharmacokinetic and pharmacodynamic profile of
donepezil HCl following evening administration: Evening administration of
donepezil HCl. Br J Clin Pharmacol. 1998;46: 13–18. doi:10.1046/j.1365-
2125.1998.0460s1013.x

98. Sauer J-M, Ring BJ, Witcher JW. Clinical pharmacokinetics of atomoxetine.
Clin   Pharmacokinet.   2005;44:   571–590.   doi:10.2165/00003088-
200544060-00002

99. Wallach H, O'connell DN. The kinetic depth effect. J Exp Psychol. 1953;45:
205–217.

100. Sperling G, Dosher BA, Landy MS. How to study the kinetic depth effect
experimentally. J Exp Psychol Hum Percept Perform. 1990;16: 445–450.

101. Engbert R, Kliegl R. Microsaccades uncover the orientation of covert
attention. Vision Res. 2003;43: 1035–1045.

102. Oostenveld R, Fries P, Maris E, Schoffelen J-M. FieldTrip: Open source
software   for   advanced   analysis   of   MEG,   EEG,   and   invasive
electrophysiological data. Comput Intell Neurosci. 2011;2011: 156869.
doi:10.1155/2011/156869

103. Bell AJ, Sejnowski TJ. An information-maximization approach to blind
separation and blind deconvolution. Neural Comput. 1995;7: 1129–1159.

104. Hyvarinen A. Fast and robust fixed-point algorithms for independent
component analysis. IEEE Trans Neural Netw. 1999;10: 626–634.
doi:10.1109/72.761722

105. Hipp JF, Siegel M. Dissociating neuronal gamma-band activity from cranial
and ocular muscle activity in EEG. Front Hum Neurosci. 2013;7.
doi:10.3389/fnhum.2013.00338

106. Mitra PP, Pesaran B. Analysis of dynamic brain imaging data. Biophys J.
1999;76: 691–708. doi:10.1016/S0006-3495(99)77236-X

107. Pascual-Marqui RD, Lehmann D, Koukkou M, Kochi K, Anderer P, Saletu
B, et al. Assessing interactions in the brain with exact low-resolution
electromagnetic tomography. Philos Transact A Math Phys Eng Sci.
2011;369: 3768–3784. doi:10.1098/rsta.2011.0081

108. Fonov V, Evans AC, Botteron K, Almli CR, McKinstry RC, Collins DL, et al.
Unbiased   average   age-appropriate   atlases   for   pediatric   studies.
NeuroImage. 2011;54: 313–327. doi:10.1016/j.neuroimage.2010.07.033

109. Nolte G. The magnetic lead field theorem in the quasi-static approximation
and its use for magnetoencephalography forward calculation in realistic
volume conductors. Phys Med Biol. 2003;48: 3637–3652.

110. Pfeffer T, Linkenkaer-Hansen K, Avramiea A-E, Engel AK, Donner TH.
Noradrenaline increases long-range temporal correlations of neuronal
alpha oscillations in the human cortex. 2015. p. 393.27.

111. Peng CK, Buldyrev SV, Havlin S, Simons M, Stanley HE, Goldberger AL.
Mosaic organization of DNA nucleotides. Phys Rev E Stat Phys Plasmas
Fluids Relat Interdiscip Top. 1994;49: 1685–1689.

112. Hardstone R, Poil S-S, Schiavone G, Jansen R, Nikulin VV, Mansvelder
HD, et al. Detrended Fluctuation Analysis: A Scale-Free View on Neuronal
Oscillations. Front Physiol. 2012;3. doi:10.3389/fphys.2012.00450

113. Montez T, Poil S-S, Jones BF, Manshanden I, Verbunt JPA, van Dijk BW,
et al. Altered temporal correlations in parietal alpha and prefrontal theta
oscillations in early-stage Alzheimer disease. Proc Natl Acad Sci.
2009;106: 1614–1619. doi:10.1073/pnas.0811699106

114. Rouder JN, Speckman PL, Sun D, Morey RD, Iverson G. Bayesian t tests
for accepting and rejecting the null hypothesis. Psychon Bull Rev. 2009;16:
225–237. doi:10.3758/PBR.16.2.225

115. Wetzels R, Wagenmakers E-J. A default Bayesian hypothesis test for
correlations and partial correlations. Psychon Bull Rev. 2012;19: 1057–
1064. doi:10.3758/s13423-012-0295-x

116. Nichols TE, Holmes AP. Nonparametric permutation tests for functional
neuroimaging: A primer with examples. Hum Brain Mapp. 2002;15: 1–25.
doi:10.1002/hbm.1058

117. Maris E, Oostenveld R. Nonparametric statistical testing of EEG- and
MEG-data.   J   Neurosci   Methods.   2007;164:   177–190.
doi:10.1016/j.jneumeth.2007.03.024

118. Eiben AE, Smit SK. Parameter tuning for configuring and analyzing
evolutionary   algorithms.   Swarm   Evol   Comput.   2011;1:   19–31.
doi:10.1016/j.swevo.2011.02.001

119. Goodman DF, Stimberg M, Yger P, Brette R. Brian 2: neural simulations on
a variety of computational hardware. BMC Neurosci. 2014;15: P199.
doi:10.1186/1471-2202-15-S1-P199

1442

SUPPLEMENTAL MATERIAL

S1 Fig. Similar Atomoxetine-related effects in both Task-counting and Task-resting conditions. **(A)** Number of perceptual alternations reported by the subjects per 10 min run for Task-counting condition. **(B)** Same as (A), but for Task-pressing condition. **(C)** Relation between the number of reported alternations during Task-counting (x-axis) and Task-pressing (y-axis). The blue line depicts a linear relation with slope 1 as a reference. Two-sided t-tests and Pearson correlations (N=28).

S2 Fig. Change in perceptual alternation rate is not due to change in blinks or fixational eye movements. **(A)** Number of EOG events for during Task-counting (left), Task-pressing (middle) and pooled across both conditions (right). Scatter plots depict the relation between the number of EOG events (x-axis) and the number of reported perceptual alternations (y-axis). **(B)** Same as (A), but for the number of detected eye blinks. **(C)** Same as (A) and (B), but for the number of saccades (horizontal and vertical). **(D)** Same as (C), but for horizontal saccades only. **(E)** Same as (D), but for vertical saccades only. Two-sided t-tests and Pearson correlations (N=28). BF, Bayes factor. These control analyses demonstrate that the change in perceptual dynamics under Atomoxetine is not explained by changes in ocular parameters.

S3 Fig. Power spectra, averaged across all MEG sensors, during Fixation **(A)** and Task-counting. **(B)** Black bar denotes significant differences assessed using a paired cluster-based permutation test (p < 0.05).

S4 Fig. Sensor-level scaling exponent for the Atomoxetine condition, pooled across Fixation and Task-counting conditions. Thresholded at $p = 0.05$, two-sided cluster-based permutation test.

S5 Fig. No Donepezil-related changes in scaling exponent in neither behavioral contexts. **(A)** Spatial distribution of Donepezil-induced changes in scaling exponent α during Fixation, thresholded at $p = 0.05$ (two-sided cluster-based permutation test). **(B)** As (A), but for Task-counting.

S6 Fig. Direct comparison of the drug effects on scaling exponent α. **(A)** Comparison of the effects of the two drugs conditions (i.e., Atomoxetine vs. Donepezil) during Fixation. **(B)** Same as (A), but during Task-counting. All thresholds at *p* = 0.05, cluster-based two-sided permutation tests (N=28).

S7 Fig. Similar effects of Atomoxetine and Donepezil on low-frequency (2-8 Hz) power. **(A)** Spatial distribution of drug-related low-frequency power changes during Fixation, thresholded at *p* = 0.05 (two-sided cluster-based permutation test). *Left.* Power changes after the administration of atomoxetine. *Right.* Power changes after the administration of donepezil. **(B)** Same as (A), but for Task-counting. The changes in low-frequency power, in combination with the reported decreases in alpha-band power, demonstrate a robust effect of both drugs on cortical dynamics.

S8 Fig. Different version of modulation of E/I ratio in cortical patch model **(A)** Neuromodulation was simulated as a gain modulation term multiplied with excitatory (EE and IE) and/or inhibitory (EI only) synaptic weights. **(B)** κ as a function of excitatory and inhibitory connectivity (with a spacing of 2.5%; means across 10 simulations per cell). The region of $\kappa\sim1$, overlaps with the region of $\alpha > 0.5$ and splits the phase space into an excitation-dominant ($\kappa>1$) and an inhibition-dominant region ($\kappa<1$). **(C)** Same as (B), but for scaling exponent α. **(D)** Same as (B) and (C), but for firing rate. In sum, the alternative version of modulation of E/I ratio yields comparable results to the version presented in Figure 8. **(E)** Model power spectra under different levels of synaptic gain modulation (neuromodulation).