1 **Supervised machine learning reveals introgressed loci in the genomes of *Drosophila***
2 ***simulans* and *D. sechellia***
3

4 Daniel R. Schrider[*,†,1], Julien Ayroles[§], Daniel R. Matute[‡], Andrew D. Kern[*,†]
5
6 [*]Department of Genetics, and [†]Human Genetics Institute of New Jersey, Rutgers University,
7 Piscataway, New Jersey 08554
8
9 [§]Ecology and Evolutionary Biology Department; Lewis Sigler Institute for Integrative Genomics,
10 Princeton University, Princeton, New Jersey, 08540
11
12 [‡]Biology Department, University of North Carolina, Chapel Hill, NC 27510.
13
14 [1]Corresponding author. Current address: Department of Genetics, University of North Carolina
15 at Chapel Hill, 120 Mason Farm Rd, Chapel Hill, NC 27514. E-mail: drs@unc.edu

16
17 **ABSTRACT**
18

Hybridization and gene flow between species appears to be common. Even though it is clear that
hybridization is widespread across all surveyed taxonomic groups, the magnitude and
consequences of introgression are still largely unknown. Thus it is crucial to develop the
statistical machinery required to uncover which genomic regions have recently acquired
haplotypes via introgression from a sister population. We developed a novel machine learning
framework, called FILET (Finding Introgressed Loci via Extra-Trees) capable of revealing
genomic introgression with far greater power than competing methods. FILET works by
combining information from a number of population genetic summary statistics, including
several new statistics that we introduce, that capture patterns of variation across two populations.
We show that FILET is able to identify loci that have experienced gene flow between related
species with high accuracy, and in most situations can correctly infer which population was the
donor and which was the recipient. Here we describe a data set of outbred diploid *Drosophila*
sechellia genomes, and combine them with data from *D. simulans* to examine recent
introgression between these species using FILET. Although we find that these populations may
have split more recently than previously appreciated, FILET confirms that there has indeed been
appreciable recent introgression (some of which might have been adaptive) between these
species, and reveals that this gene flow is primarily in the direction of *D. simulans* to *D.*
sechellia.

37
38
39
40

AUTHOR SUMMARY

Understanding the extent to which species or diverged populations hybridize in nature is crucially important if we are to understand the speciation process. Accordingly numerous research groups have developed methodology for finding the genetic evidence of such introgression. In this report we develop a supervised machine learning approach for uncovering loci which have introgressed across species boundaries. We show that our method, FILET, has greater accuracy and power than competing methods in discovering introgression, and in addition can detect the directionality associated with the gene flow between species. Using whole genome sequences from *Drosophila simulans* and *Drosophila sechellia* we show that FILET discovers quite extensive introgression between these species that has occurred mostly from *D. simulans* to *D. sechellia*. Our work highlights the complex process of speciation even within a well-studied system and points to the growing importance of supervised machine learning in population genetics.

INTRODUCTION

Up to 10% of animal [1] and plant [2] species have the ability to hybridize with other species. Our recent ability to collect large-scale genomic data has confirmed that hybridization is common in nature. Indeed the ubiquity of hybridization upon secondary contact raises the question of how large a role hybridization plays in the emergence or collapse of new lineages [3].

Three general patterns have emerged from recent efforts to search for introgression in genomic data. First, whole-genome sequencing has shown that introgression occurs in all taxa for which its signature has been systematically sought (primates reviewed in [4], plants in [5, 6], fungi [7] and oomycetes in [8]). In general, genetic exchange between species through fertile hybrids might be common between closely related species [9-13] but can also occur between divergent species [14-17].

Second, introgression is heterogeneously distributed across the genome. For instance, mitochondrial genome exchange is surprisingly common (e.g., [18-20] among many) between species, whereas sex chromosomes are less likely to cross species-boundaries, perhaps due to their disproportionate role in hybrid incompatibilities [17, 21-24]. Generally it seems that functional regions of the genome might be less likely to participate in introgression. This is perhaps best known from the case of Neanderthal hybridization with non-African human populations [25, 26], which has left modern human genomes distinct gradients of introgression across different functional compartments of the genome.

Finally, the mode and intensity of natural selection acting on introgressed DNA can vary substantially. Loci that contribute to reproductive isolation, and as such to the persistence of species in the face of hybridization, should be less likely to be introgressed [27] as a result of purifying selection in hybrids. Additionally, introgressed haplotypes containing mildly deleterious variants may be purged after migrating into a population with a larger effective size where selection is more effective [28, 29]. On the other hand, much of the genome may be porous to introgression between closely related species if the net effect of such introgressed variation is fitness neutral. Of course genetic exchange between populations can also provide a source of adaptive alleles that may facilitate adaptation to new environments (reviewed in ref. [30]). Introgressions have indeed been shown to be involved in adaptation in animals (e.g. [31-33]), plants (e.g. [34]) and fungi [35]. For instance, adaptation to high altitude in Tibetans appears to have been caused by introgression of alleles from an archaic Denisovan-like source into modern humans [36]. Another particularly well-studied system of adaptive introgression comes from *Heliconius* butterflies where gene exchange has facilitated the origin and maintenance of mimetic rings [32] and even of hybrid species [37, 38].

Clearly, hybridization and introgression play an important role in shaping the landscape of genetic variation, thus if we wish to fully understand its evolutionary role a reliable framework for the inference of introgressed alleles is needed. Approaches to detect introgression in the genome fall into a few different camps. Genome-wide approaches can identify whether

admixture has occurred in a set of populations. These include clustering methods which seek to
infer which individuals are admixed and to assign a proportion of admixture to each individual
without previous knowledge of the parental populations [39-41]. Some genome-wide approaches
instead attempt to infer the directionality of introgression by examining allele frequency
differences among populations [25, 42]. The main limitation of this class of methods is that they
cannot identify which regions of the genome are likely to have crossed species boundaries.

On the other hand, locus-specific ancestry approaches (e.g. [43-47]) seek to uncover the
mosaic of ancestry for each sampled haplotype, and thus can also identify portions of haplotypes
that have been introgressed between species or populations. These fine-resolution approaches are
powerful but often have assumptions and requirements that cannot be fulfilled in many taxa
which range from the need of phased haplotypes to recombination maps. The main limitation of
these approaches is that many require a set of reference haplotypes—individuals known to be
unadmixed representatives of either population—in order to properly infer the origin of each
allele in each (non-reference) sample haplotype.

The last family of approaches designed to uncover introgressed loci has focused on the
use of relative and absolute levels of divergence measured in genomic windows. Largely such
methods have consisted of new summary statistics that capture elements of the expected
coalescent genealogy under a model of recent introgression between species. These approaches
have the advantage that no donor or recipient populations must be identified a priori. Among the
measurements of divergence, F_{ST} [48] is most commonly used. Another popular point of
departure has been the d_{xy} statistic of Nei and Li [49] which measures the average pairwise
distance between alleles sampled from two populations. For instance, Joly et al. [50], Geneva et
al. [51] and Rosenzweig et al. [52] use the minimum rather than the mean of these pairwise
divergence values, termed d_{min}. d_{min} is sensitive to abnormally short branch lengths between
alleles drawn from two populations, as would be expected when introgression is recent. Each of
these statistics has attractive properties and adequate power in some instances, however no one
statistic has perfect sensitivity in every scenario.

Here we introduce a new method for finding introgressed loci based on supervised
machine learning that we call FILET (Finding Introgressed Loci using Extra Trees Classifiers).
FILET combines a large number of summary statistics (Materials and Methods) that provide
complementary information about the shape of the genealogy underlying a region of the genome.
These summary statistics include both previously developed statistics (including, but not limited
to, those based on d_{min} and d_{xy}) as well as 5 new summary statistics that we describe below. Our
reasoning for this approach was that by combining many statistics for detecting introgression we
should achieve sensitivity to introgression across a larger range of scenarios than accessible to
any individual statistic. Buoyed by our recent work showing the power and flexibility of Extra
Trees classifiers [53] for population genomic inference [54, 55], we leveraged this machine
learning paradigm for identification of introgression. Using simulations we show that FILET is
far more powerful and versatile than competing methods for identifying introgressed loci.

Further we apply FILET to examine patterns of introgression between *Drosophila simulans* and
its island endemic sister taxon *Drosophila sechellia*.
The speciation event that gave rise to the island endemic *Drosophila sechellia* from a
Drosophila simulans-like ancestor is a textbook example of a specialist species that evolved
from a presumably generalist ancestor [56, 57]. Indeed, *D. sechellia* has quite remarkably
specialized to breed on the toxic fruit of *Morinda citrifolia* [58], while *D. simulans* (and *D.*
mauritiana) do not tolerate the organic volatile compounds in the ripe fruit [59-61]. The genetic
and neurological underpinnings of this key ecological difference have been identified, at least in
part [62-67] making the *D. simulans*/*D. sechellia* pair one of the most successful cases of genetic
dissection of the causes of an ecologically relevant trait. While this is so, the population genetics
of divergence between these species has only been examined in the context of either population
samples from a handful of loci [68-71] or in the absence of population data [72]. These studies
estimated population divergence time between *D. simulans* and *D. sechellia* to be as early as
~250,000 years ago [72] or as old as ~413,000 years ago [70]. All population genomic surveys
demonstrate that *D. sechellia* harbors little genetic variation in comparison to *D. simulans*,
perhaps as a result of a population size crash/founder event from which the population has not
recovered [68, 71]. Moreover it has been suggested that what little variation there is in *D.*
sechellia shows little population genetic structure among separate island populations in the
Seychelles archipelago [71]. Lastly there is some evidence of introgression between each pair of
species within the *D. simulans* complex [72], and *D. simulans* and *D. sechellia* have been found
to hybridize in the field [73]. Here we characterize the population genetics of divergence
between *D. sechellia* and *D. simulans,* combining existing whole-genome sequences from a
mainland population of *D. simulans* [74] with newly generated genome sequences from *D.*
sechellia. Applying FILET to these data confirms previous reports of introgression between
these species and reveals that this gene flow is primarily in the direction of *D. simulans* to *D.*
sechellia. Finally, the success of our approach underscores the potential power of supervised
machine learning for evolutionary and population genetic inference.
162
163 **MATERIALS AND METHODS**
164
165 **Statistics capturing the population genetic signature of introgression**
166
We set out to assemble a set of statistics that could be used in concert to reliably determine
whether a given genomic window had experienced recent gene flow. Several statistics that have
been designed to this end ask whether there is a pair of samples exhibiting a lower than expected
degree of sequence divergence within the window of interest. The most basic of these is d_{min}, the
minimum pairwise divergence across all cross-population comparisons (S1 Fig; [50]). The
reasoning behind d_{min} is that even if only a single sampled individual contains an introgressed
haplotype, d_{min} should be lower than expected and the introgression event may be detectable. A
related statistic is G_{min}, which is equal to d_{min}/d_{xy} [51]; the presence of this term in the

denominator is meant to control for variation in the neutral mutation rate across the genome.
RND_{min} accomplishes this by dividing d_{min} by the average divergence of all sequences from either
species to an outgroup sequence [52]. The name of this statistic is derived from its constituent
parts, d_{min}, and RND [75].
As described in the following section, we incorporated a number of previously devised
statistics into our classification approach, including some of those based on d_{min}. We also
included some novel statistics that we designed to have improved sensitivity to particularly
recent introgression. The first of these is defined as:
$d_{d1} = d_{min}/\pi_1$
where π_1 is nucleotide diversity [49] in population 1. Similarly, $d_{d2} = d_{min}/\pi_2$. d_{d1} and d_{d2} statistics
are so named because they compare d_{min} to diversity within populations 1 and 2, respectively.
The rationale behind these statistics is that, if there has been recent introgression from population
1 into population 2, and at least one sampled chromosome from population 2 contains the
introgressed haplotype, then the cross-population pair of individuals yielding the value of d_{min}
should both trace their ancestry to population 1. Thus, the sequence divergence between these
two individuals should on average be equal to π_1. Similarly, if there was introgression in the
reverse direction d_{min} would be on the order of π_2. Following similar rationale, we devised two
related statistics: $d_{d-Rank1}$ and $d_{d-Rank2}$. $d_{d-Rank1}$ is the percentile ranking of d_{min} among all pairwise
divergences *within* population 1; the value of this statistic should be lower when there has been
introgression from population 1 into population 2. $d_{d-Rank2}$ is the analogous statistic for
introgression from population 2 into population 1. We also included a statistic comparing
average linkage disequilibrium within populations to average LD within the global population
(i.e. lumping all individuals from both species together), as follows:
$Z_X = (Z_{nS1} + Z_{nS2}) / (2 \times Z_{nSG})$
where Z_{nS1}, and Z_{nS2} measure average LD [76] between all pairs of variants within the window in
population 1 and population 2, respectively, and Z_{nSG} which measures LD within the global
population. The reasoning behind this statistic is based on the assumption that, in the presence of
gene flow, LD may be elevated within the recipient population(s) but not in the global
population. S2 Fig shows that the distributions of these statistics do indeed differ substantially
between genealogies with and without introgression (simulation scenarios described below),
especially when this introgression occurred recently. In addition to these and other statistics
summarizing diversity across the two population samples, we also incorporated several single-
population statistics into our classifier (see below), as these may also contain information about
recent introgression. For example, separate measures of nucleotide diversity in our two
population samples would contain useful information because introgression is expected to
increase diversity in the recipient population, especially if the source population was large or if
the two populations split long ago.
212
213 **Description of FILET classifier**
214

We used a supervised machine learning approach to assign a genomic window to one of three
distinct classes on the basis of a "feature vector" consisting of a number of statistics
summarizing patterns of variation within the window from two closely related populations.
These three classes are: introgression from population 1 into population 2, introgression from
population 2 into population 1, and the absence of introgression. Specifically, we used an Extra-
Trees classifier [53], which is an extension of random forests [77], an ensemble learning
technique that creates a large ensemble of semi-randomly generated binary decision trees [78],
before taking a vote among these decision trees in order to decide which class label should be
assigned to a given data instance (i.e. genomic window in our case). In an Extra-Trees classifier,
the tree building process is even more randomized than in typical random forests: in addition to
selecting a random subset of features when generating a tree, the separating threshold for each
feature is randomly chosen, rather than selected the threshold that optimally separates the data
classes. We require example regions for each class in order to train the Extra-Trees classifier, so
we used coalescent simulations to generate these training examples (described below). Our
ultimate goal was to detect introgression within 10 kb windows in *Drosophila*, so to train our
classifier properly we simulated chromosomal regions approximating this length (simulation
details are given below). The target window size could easily be altered by changing the length
of the regions simulated for training (i.e. by adjusting the recombination and mutation rates, θ
and ρ).

FILET's feature vector contains a number of single-population summaries of per-base
pair genetic variation: π, the variance in pairwise distances among individuals, the density of
segregating sites, the density of polymorphisms private to the population, Fay and Wu's H and θ_H
statistics [79], and Tajima's D [80]. The feature vector also includes two single-population
summary statistics that are not normalized per base pair: Z_{nS} (which is averaged across all pairs
of SNPs), and the number of distinct haplotypes observed in the window. Each feature vector
included values of these 9 statistics for each population, yielding 18 single-population statistics
in total. In addition, the two-population statistics included in FILET's feature vector were as
follows: F_{ST} (following ref. [81]), Hudson's S_{nn} [82], per-bp d_{xy}, per-bp d_{min}, G_{min}, d_{d1}, d_{d2}, $d_{d\text{-}Rank1}$, $d_{d\text{-}Rank2}$, Z_X, IBS_{MaxB} (the length of the maximum stretch of identity by state, or IBS, among
all pairwise between-population comparisons), and IBS_{Mean1} and IBS_{Mean2} which capture the
average IBS tract length when comparing all pairs of sequences within populations 1 and 2,
respectively. These IBS statistics are calculated by examining all pairs of individual sequences
within a population (or across populations in the case of IBS_{MaxB}), noting the positions of
differences, and examining the distribution of lengths between these positions (as well as
between the first position and the beginning of the window and between the last position and the
end of the window). Note that we did not include RND_{min} or other measures such as Patterson's
D and F_4 statistics [83] that require alignment to one or more additional species along with the
focal pair, because we designed FILET so that it would not require outgroup information. We
note however that through its use of supervised machine learning, FILET could easily be
extended to incorporate such data. Instead, in order to improve robustness to mutational

variation, we adopted the approach of drawing the mutation rate from a wide range of values
when generating training examples to train FILET to classify data from our *Drosophila* samples
(see below). All code necessary to run the FILET classifier (including calculating summary
statistics on both simulated and real data sets, training, and classification) along with the full
results of our application to *D. simulans* and *D. sechellia* (described below) are available at
https://github.com/kern-lab/FILET/.
261
262 **Simulated test scenarios**
263
Following Rosenzweig et al. [52], we used the coalescent simulator msmove
(https://github.com/geneva/msmove) to simulate data for testing FILET's power to detect
introgression in populations with four different values of T_D (the time since divergence):
$0.25\times4N$, $1\times4N$, $4\times4N$, and $16\times4N$ generations ago, where N is the population size. For each of
these simulations the population size was held constant (i.e. the ancestral population size equals
that of both daughter populations). We developed a classifier for each of these scenarios of
population divergence. Supervised machine learning techniques such as the Extra-Trees
classifier require training data consisting of examples from each of the three classes, but in
practice a large number of example loci known to have experienced introgression may not be
available. We therefore simulated training data sets for each of the four values of T_D. Again
following Rosenzweig et al. [52], the relevant parameters for each of these simulations include:
T_M, the time since the introgression event, which we drew from $\{0.01\times T_D, 0.05\times T_D, 0.1\times T_D,$
$0.15\times T_D, \ldots, 0.9\times T_D\}$ (i.e. multiples of $0.05\times T_D$ up to 0.9, and also including $0.01\times T_D$); and P_M,
the probability that a given lineage would migrate from the source population to the sink
population during the introgression event, which we drew from $\{0.05, 0.1, 0.15, \ldots, 0.95\}$. We
simulated an equal number of training examples for each combination of these two parameter
values for both directions of gene flow, yielding 10^4 simulations in total for both of these classes,
conditioning that each of these instances must have contained at least one migrant lineage.
Finally, we simulated an equivalent number of samples without introgression, yielding a
balanced training set (10^4 examples for each class).
Next, we computed feature vectors as described above for each of these training
examples, and proceeded with training our Extra-Trees classifiers by conducting a grid search of
all training parameters precisely as described in Schrider and Kern [54], and setting the number
of trees in the resulting ensemble to 100. All training and classification with the Extra-Trees
classifier was performed using the scikit-learn Python library (http://scikit-learn.org; [84]). We
also calculated feature importance and rankings thereof by training an Extra-Trees classifier of
500 decision trees on the same training data (using scikit-learn's defaults for all other learning
parameters), and then using this classifier's "feature_importances_" attribute. Briefly, this
feature importance score is the average reduction in Gini impurity contributed by a feature across
all trees in the forest, always weighted by the probability of any given data instance reaching the
feature's node as estimated on the training data [85]; this measure thus captures both how well a

feature separates data into different classes and how often the feature is given the opportunity to
split (i.e. how often it is visited in the forest). The values of these scores are then normalized
across all features such that they sum to one.
For each T_D, we evaluated the appropriate classifier against a larger set of 10^4 simulations
generated for each parameter combination along a grid of values of T_M and P_M. The values of P_M
were drawn from the same set as those in training as described above, while one additional
possible value of T_M was included: $0.001 \times T_D$. Also note that for these simulations we did not
require at least one migrant lineage as we had done for training (information that is recorded by
msmove). For test simulations with bidirectional migration, we did not require each replicate
sample to contain at least one migrant lineage, though we modified msmove to record which
samples did in fact experience migration. In addition to test examples for each direction of gene
flow, we simulated 10^4 examples where no migration occurred in order to assess false positive
rates. In order to examine the performance of FILET when an unsampled ghost lineage was the
source of introgression, we generated test simulations with the same values of T_D, T_M, and P_M as
above, but where the source of the introgression was a third, unsampled population that separated
from the two sampled populations at time T_D. In all of our simulations, both for training and
testing, we set locus-wide population mutation and recombination rates θ and ρ to 50 and 250,
respectively, similar to autosomal values in 10 kb windows in *D. melanogaster* [86] and sampled
15 individuals from each population. We also experimented with different window sizes,
simulating training and test data (1,000 replicates for each class for each set) with window sizes
corresponding to 1 kb, 10 kb, 5 kb, and 50 kb by multiplying θ and ρ by the appropriate scalar.
When testing the sensitivity of our method on these data, we considered a window to be
introgressed if FILET's posterior probability of the no-introgression class was <0.05, except for
the scenario with T_D equal to $16 \times 4N$ generations ago in which case we used a posterior
probability cutoff of 0.01, as we found that this step mitigated the elevated false positive rate
under this scenario (reducing the rate from >10% to the estimate of 6% shown in S3 Fig). In
windows labeled as introgressed, the direction of gene flow was determined by asking which of
the two introgression classes had a higher posterior probability. Note that we used the same
demographic scenario for both the training and test data for each T_D, and discuss the implications
of demographic model misspecification in the Results and Discussion.
In order to compute receiver operator characteristic (ROC) curves we constructed
balanced binary training sets composed of 10^4 examples with no introgression, and 10^4 examples
allowing for introgression (with equal representation to each combination of T_M, P_M, and
direction of introgression. The score that we obtained for each test example in order to compute
the ROC curve was one minus the posterior probability of no introgression as generated by the
Extra-Trees classifier (i.e. the classifier's estimated probability of introgression, regardless of
directionality).
332
333   **Comparison to ChromoPainter**
334

We compared FILET's accuracy to that of ChromoPainter [46], a software program that utilizes a variant of the copying model first proposed by Li and Stephens [87]. In this model each sample haplotype is a mosaic composed of chromosomal segments chosen from a set of possible ancestral haplotypes, allowing for differences caused by mutation and the potential for changes in ancestry at recombination breakpoints. Such an approach can thus be used to predict the ancestry of each individual at each polymorphism—these predictions are referred to as "paintings" by ChromoPainter. To this end we repeated our simulations above but increased the size of the chromosomal segments to 1 Mb by increasing θ and ρ to 5000 and 25000. In these simulations only gene flow from population 2 to population 1 was allowed, and we modified msmove to record the coordinates of introgressed segments, and to restrict introgression events to those involving segments falling entirely within the central 100 kb of the chromosome. For each combination of T_M and P_M we generated 10 replicate simulations, including 10 replicates without introgression.

We ran ChromoPainter with the following parameters: the "-a 0 0" switch to model each individual haplotype as a mosaic of each other individual rather than using a set of predefined reference haplotypes, "-i 10" and "-ip" options to estimate copying proportions over 10 Expectation-Maximization (EM) iterations, and the default "-s 10" switch to stochastically draw 10 chromosome paintings for each individual from the HMM following EM. We then used the output from ChromoPainter to predict introgressed chromosomal segments as follows:

For each polymorphism, we examine each haplotype i among our n haplotypes, and record which of the other n-1 haplotypes serves as the best ancestor for i in each of our 10 paintings. We then examine each individual in population 2 (the recipient population), and count the number of paintings for which the ancestral haplotype is from population 1. If this number is > 5 (i.e. a majority) for any of our individuals in population 2, then we consider this focal polymorphism to be introgressed. If two adjacent polymorphisms are predicted to be introgressed, all sites between them are also considered to be introgressed. If only 1 polymorphism is predicted, then just that one site is considered introgressed. We also produced a more stringent version of these predictions by only retaining introgressed segments consisting of at least 25 consecutive introgressed polymorphisms. Note that ChromoPainter requires base pair positions, and msmove uses an infinite sites model where polymorphisms are located in a continuous space between zero and one. Thus in order to perform this analysis we had to map msmove's continuous locations to physical locations, which we accomplished by multiplying by 10^6 and rounding to the nearest available position.

We compared ChromoPainter to a sliding-window application of FILET's classifier for 10 kb windows (with 1 kb step sizes). We also produced finer-scale FILET predictions using a 1 kb classifier (with 100 bp step sizes) to refine predictions made by the 10 kb classifier: only sliding windows predicted as introgressed by the 1 kb classifier and lying within introgressed segments predicted by the 10 kb classifier were retained as candidates by this version. For the refinement step, FILET's posterior probability cutoff for introgression was relaxed to 0.5 (i.e. introgression more probable than not); a more lenient cutoff is appropriate here because this

375 classifier was only applied within regions already predicted to be introgressed by the 10 kb
376 classifier.
377
378 ***Drosophila sechellia* collection**
379
Drosophila sechellia flies were collected in the islands of Praslin, La Digue, Marianne and Mahé
with nets over fresh *Morinda* fruit on the ground. All flies were collected in January of 2012.
Flies were aspirated from the nets by mouth (1135A Aspirator – BioQuip; Rancho Domingo,
CA) and transferred to empty glass vials with wet paper balls (to provide humidity) where they
remained for a period of up to three hours. Flies were then lightly anesthetized using FlyNap
(Carolina Biological Supply Company, Burlington, NC) and sorted by sex. Females from the
melanogaster species subgroup were individualized in plastic vials with instant potato food
(Carolina Biologicals, Burlington, NC) supplemented with banana. Propionic acid and a pupation
substrate (Kimwipes Delicate Tasks, Irving TX) were added to each vial. Females were allowed
to produce progeny and imported using USDA permit P526P-15-02964. The identity of the
species was established by looking at the taxonomical traits of the males produced from
isofemale lines (genital arches, number of sex combs) and the female mating choice (i.e.,
whether they chose *D. simulans or D. sechellia* in two-male mating trials).
393
394 **Sequence data and variant calling and phasing**
395
We obtained sequence data from 20 *D. simulans* inbred lines [74] from NCBI's Short Read
Archive (BioProject number PRJNA215932). We also sequenced wild-caught outbred *D.*
sechellia females (see above) from Praslin (*n*=7 diploid genomes), La Digue (*n*=7), Marianne
(*n*=2), and Mahé (*n*=7). These new *D. sechellia* genomes are available on the Short Read
Archive (BioProject number PRJNA395473). For each line we then mapped all reads with bwa
0.7.15 using the BWA-MEM algorithm [88] to the March 2012 release of the *D. simulans*
assembly produced by Hu et al. [89] and also used the accompanying annotation based on
mapped FlyBase release 5.33 gene models [90]. Next, we removed duplicate fragments using
Picard (https://github.com/broadinstitute/picard), before using GATK's HaplotypeCaller (version
3.7; [91-93]) in discovery mode with a minimum Phred-scaled variant call quality threshold (-
stand_call_conf) of 30. We then used this set of high-quality variants to perform base quality
recalibration (with GATK's BaseRecalibrator program), before again using the HaplotypeCaller
in discovery mode on the recalibrated alignments. For this second iteration of variant calling we
used the --emitRefConfidence GVCF option in order to obtain confidence scores for each site in
the genome, whether polymorphic or invariant. Finally, we used GATK's GenotypeGVCFs
program to synthesize variant calls and confidences across all individuals and produce genotype
calls for each site by setting the --includeNonVariantSites flag, before inferring the most
probable haplotypic phase using SHAPEIT v2.r837 [94]. The genotyping and phasing steps were
performed separately for our *D. simulans* and *D. sechellia* data, and for each of step in the

pipeline outlined above we used default parameters unless otherwise noted. In order to remove
potentially erroneous genotypes (at either polymorphic or invariant sites), we considered
genotypes as missing data if they had a quality score lower than 20, or were heterozygous in *D.*
simulans. After throwing out low-confidence genotypes, we masked all sites in the genome
missing genotypes for more than 10% of individuals in either species' population sample, as well
as those lying within repetitive elements as predicted by RepeatMasker
(http://www.repeatmasker.org). Only SNP calls were included in our downstream analyses (i.e.
indels of any size were ignored). These phased and masked data are available at
https://zenodo.org/record/1166021.
424
425 **Demographic inference**
426
Having obtained genotype data for our two population samples, we used $\partial a \partial i$ [95] to model their
shared demographic history on the basis of the folded joint site frequency spectrum
(downsampled to $n=18$ and $n=12$ in *D. simulans* and *D. sechellia*, respectively); using the folded
spectrum allowed us to circumvent the step of producing whole genome alignments to outgroup
species in *D. simulans* coordinate space in order to attempt to infer ancestral states. We used an
isolation-with-migration (IM) model that allowed for continual exponential population size
change in each daughter population following the split. This model includes parameters for the
ancestral population size (N_{anc}), the initial and final population sizes for *D. simulans* (N_{sim_0} and
N_{sim}, respectively), the initial and final sizes for *D. sechellia* (N_{sech_0} and N_{sech}, respectively), the
time of the population split (T_D), the rate of migration from *D. simulans* to *D. sechellia*
($m_{sim \rightarrow sech}$), and the rate of migration from *D. sechellia* to *D. simulans* ($m_{sech \rightarrow sim}$). We also fit our
data to a pure isolation model that was identical to our IM model but with $m_{sim \rightarrow sech}$ and $m_{sech \rightarrow sim}$
fixed at zero. Finally, we fit our data to an admixture model identical to the isolation model but
with the addition of two parameters: the time of a pulse admixture event from *D. simulans* into
D. sechellia (T_{AD}) and the proportion of individuals in *D. sechellia* migrating from *D. simulans*
during this event (P_{AD}). Our optimization procedure consisted of an initial optimization step
using the Augmented Lagrangian Particle Swarm Optimizer [96], followed by a second step of
optimization refining the initial model using the Sequential Least Squares Programming
algorithm [97], both of which are included in the pyOpt package for optimization in Python
(version 1.2.0; [98]) as in Schrider et al. [99]. We performed ten optimization runs fitting both of
these models to our data, each starting from a random initial parameterization, and assessed the
fit of each optimization run using the AIC score. Code for performing these optimizations can be
obtained from https://github.com/kern-lab/miscDadiScripts, wherein 2popIM.py,
2popIsolation.py, 2popIsolation_admixture.py fit the IM, isolation, and admixture models
described above, respectively. For scaling times by years rather than numbers of generations, we
assumed a generation time of 15 gen/year as has been estimated in *D. melanogaster* [100].
453
Training FILET to detect introgression between *D. simulans* and *D. sechellia*

Having obtained a demographic model that provided an adequate fit to our data, we set out to
simulate training examples under this demographic history for each of our three classes (i.e. no
migration, migration from *D. simulans* to *D. sechellia*, and from *D. sechellia* to *D. simulans*). For
training examples including introgression, T_M was drawn uniformly from the range between zero
generations ago and $T_D/4$, while P_M raged uniformly from (0, 1.0]. In addition, in order to make
our classifier robust to uncertainty in other parameters in our model, for each training example
we drew values of each of the remaining parameters from [$x−(x/2)$, $x+(x/2)$], where x is our point
estimate of the parameter from ∂a∂i. In addition to the parameters from our demographic model
(T_D, ρ, N_{anc}, N_{sim}, and N_{sech}), these include the population mutation rate $\theta=4N\mu$ (where μ was set
to 3.5×10^{-9}), and the ratio of θ to the population recombination rate ρ (which we set to 0.2).
Continuous migration rates were set to zero (i.e. the only migration events that occurred were
those governed by the T_M and P_M parameters, and the no-migration examples were truly free of
migrants). In total, this training set comprised of 10^4 examples from each of our three classes.

As described above, we masked genomic positions having too many low confidence
genotypes or lying within repetitive elements (described above) before proceeding with our
classification pipeline. While doing so, we recorded which sites were masked within each 10 kb
window in the genome that we would later attempt to classify. In order to ensure that our
masking procedure affected our simulated training data in the same manner as our real data, for
each simulated 10 kb window we randomly selected a corresponding window from our real
dataset and masked the same sites in the simulated window that had been masked in the real one.
We then trained our classifier in the same manner as described above.

In order to ensure that this classifier would indeed be able to reliably uncover loci
experiencing recent gene flow between our two populations, we assessed its performance on
simulated test data. First, we applied the classifier to test examples simulated under this same
model (again, 10^4 for each class). Next, to address the effect of demographic model
misspecification, we applied our classifier to an isolation model with a different parameterization
and no continuous size change in the daughter populations. For this model we simply set N_{sim}
and N_{sech} to $\pi_{sim}/4\mu$ and $\pi_{sech}/4\mu$, respectively, where π for a species is the average nucleotide
diversity among all windows included in our analysis after filtering, and μ was again set to
3.5×10^{-9}. We then set N_{anc} to be equal to N_{sim}, and set T to $d_{xy}/(2\mu) − 2N_{anc}$ generations where d_{xy}
is the average divergence between *D. simulans* and *D. sechellia* sequences across all windows.
This latter value is simply the expected TMRCA for cross-species pairs of genomes, minus the
expected waiting time until coalescence during the one-population (i.e. ancestral) phase of the
model. This simple model thus produces samples with similar levels of nucleotide diversity for
the two daughter populations as those produced under our IM model, but that would differ in
other respects (e.g. the joint site frequency spectrum and linkage disequilibrium, which would be
affected by continuous population size change after the split). For both test sets we masked sites
in the same manner as for our training data before running FILET.

Classifying genomic windows with FILET

We examined 10 kb windows in the *D. simulans* and *D. sechellia* genomes, summarizing diversity in the joint sample with the same feature vector as used for classification (see above), ignoring sites that were masked as described above. We omitted from this analysis any window for which >25% of sites were masked, and then applied our classifier to each remaining window, calculating posterior class membership probabilities for each class. We then used a simple clustering algorithm to combine adjacent windows showing evidence of introgression into contiguous introgressed elements: we obtained all stretches of consecutive windows with >90% probability of introgression as predicted by FILET (i.e. the probability of no-introgression class <10%), and retained as putatively introgressed regions those that contained at least one window with >95% probability of introgression. In order to test for enrichment of these introgressed regions for genic/intergenic sequence or particular Gene Ontology (GO) terms from the FlyBase 5.33 annotation release [90], we performed a permutation test in which we randomly assigned a new location for each cluster or introgressed windows (ensuring the entire permuted cluster landed within accessible windows of the genome according to our data filtering criteria). We generated 10,000 of these permutations.

RESULTS AND DISCUSSION

FILET detects introgressed loci with high sensitivity and specificity

We sought to devise a bioinformatic approach capable of leveraging population genomic data from two related population samples to uncover introgressed loci with high sensitivity and specificity. In the Materials and Methods, we describe several previous and novel statistics designed to this end. However, rather than preoccupying ourselves with the search for the ideal statistic for this task, we took the alternative approach of assembling a classifier leveraging many statistics that would in concert have greater power to discriminate between introgressed and non-introgressed loci. Supervised machine learning methods have proved highly effective at making inferences in high-dimensional datasets and are beginning to make inroads in population genetics [101]. In this vein, we designed FILET, which uses an extension of random forests called an Extra-Trees classifier [53]. We previously succeeded in applying Extra-Trees classifiers for a separate population genetic task—finding recent positive selection and discriminating between hard and soft sweeps [54, 55]—though other methods such as support vector machines [102] or deep learning [103] could also be applied to this task.

Briefly, FILET assigns a given genomic window to one of three distinct classes—recent introgression from population 1 into population 2, introgression from population 2 into 1, or the absence of introgression—on the basis of a vector of summary statistics that contain information about the two-population sample's history. This feature vector contains a variety of statistics summarizing patterns of diversity within each population sample, as well as a number of

statistics examining cross-population variation (see Materials and Methods for a full description).
FILET must be trained to distinguish among these three classes, which we accomplish by
supplying 10,000 simulated example genomic windows of each class, with each example
represented by its feature vector. Because we expect that the majority of introgression events to
be non-adaptive, these simulations did not include natural selection. Once this training is
complete, FILET can then be used to infer the class membership of additional genomic windows,
whether from simulated or real data.

We began by assessing FILET's power on a number of simulated datasets, examining
windows roughly equivalent to 10 kb in length in *Drosophila* (Materials and Methods). In
particular, because the power to detect introgression depends on the time since their divergence,
T_D, we measured FILET's performance under four different values of T_D, training a separate
classifier for each. In Figure 1 ($T_D=0.25\times4N$) and S3 Fig (T_D values of 1, 4, and $16\times4N$), we
compare FILET's power to that of two related statistics that have been devised to detect
introgressed windows, d_{min} and G_{min} (Materials and Methods). These figures show that FILET
has high sensitivity to introgression across a much wider range of introgression timings (T_M) and
intensities (P_M) than either of these statistics under each value of T_D, and that this disparity is
amplified dramatically for smaller values of T_D. Furthermore, these figures demonstrate that
FILET infers the correct directionality of recent introgression with high accuracy, whereas d_{min}
and G_{min} contain no information about the direction of gene flow. Finally, FILET does not appear
especially sensitive when the source of gene flow is an unsampled ghost population rather than
one of the two sequenced populations (S4 Fig), though it could potentially be trained to detect
such cases if desired.

We also note that for d_{min} and G_{min} we established 95% significance thresholds from our
simulated training data without introgression, thereby achieving a false positive rate of 5%. In
order to assess FILET's false positive rate, we classified a set of test simulations without
introgression and found that FILET's false positive rate was considerably lower (Figure 1 and S3
Fig) except for our largest value of T_D, where it was initially higher (0.4% for $T_D=0.25\times4N$ but
>10% for $T_D=16\times4N$), despite our selection of a posterior probability cutoff of 95% (Methods).
This illustrates an important issue with posterior probability estimates produced by supervised
machine learning methods: they may occasionally be miscalibrated. We therefore adjusted the
cutoff for the $T_D=16\times4N$ scenario (to 99% probability of introgression) which lowered our false
positive rate to 6% as shown in S3 Fig. Thus, when an appropriate posterior probability cutoff is
chosen—a task that can be performed in a straightforward manner by testing on simulated data—
FILET achieves much greater sensitivity to introgression than d_{min} and G_{min} often at a much
lower false positive rate. We also demonstrate the FILET's greater power than these statistics via
ROC curves (S5 Fig), where it outperforms each statistic under each scenario. Specifically, the
difference in power between FILET and d_{min} is dramatic for smaller values of T_D (area under
curve, or AUC, of 0.85 versus 0.73 when $T_D=0.25\times4N$ for FILET and d_{min}, respectively) but
comparatively miniscule for our largest T_D (AUC of 0.94 versus 0.93 when $T_D=16\times4N$). It
therefore appears that FILET's performance gain relative to single statistics is highest for the

more difficult task of finding introgression between very recently diverged populations, while for
the easier case of detecting introgression between highly diverged populations some single
statistics may perform nearly as well.
We also experimented with smaller training sets, finding similar classification power
(measured by AUC) as above when we trained FILET using only 1000 or even 100 simulated
examples per class (S6 Fig), though in the latter case estimated class posterior probabilities were
far less accurate. In addition, we examined the effect of altering the target window size used
when training and testing FILET (S7 Fig).
583
584 **A comparison of the power and resolution of FILET and ChromoPainter**
585
Methods designed to uncover changes in ancestry along a recombining chromosome within
admixed populations can also be used to recover introgressed regions. To this end we used
ChromoPainter [46] which has the advantage of not requiring a set of "reference haplotypes"
known to be free of introgression from each population, and can instead predict for each
haplotype, which of all the other haplotypes in the sample (from either population) is most
closely related. We simulated two-population samples for 1 Mb chromosomes where
introgression from population 2 to population 1 was allowed in the central 100 kb window, and
used ChromoPainter to identify introgressed loci (see Methods). We then ran FILET on these
simulations, this time using a sliding-window approach to detect introgressed segments
(Methods).
Figure 2 shows that FILET has substantially higher sensitivity than ChromoPainter—
summing across the entire parameter space (including many scenarios where introgression is
quite difficult to detect) FILET recovered 27.7% of introgressed base pairs compared to 19.4%
for ChromoPainter—while having a roughly 20-fold lower false positive rate (0.42% for FILET
versus 9.31% for ChromoPainter). For scenarios with more ancient and less intense
introgression, we did observe somewhat higher sensitivity in ChromoPainter's predictions.
However, this seems to be driven largely by ChromoPainter's propensity to identify a larger
fraction of base pairs as introgressed regardless of their true ancestry, as evidenced by its higher
false positive rate. To demonstrate this further we show for the positive predictive value (the
number of base pairs correctly predicted to be introgressed divided by the total number of base
pairs predicted to be introgressed) for each method in S8 Fig. This figure shows that FILET's
positive predictive is consistently far higher than ChromoPainter's. We sought to improve this by
adopting a more stringent threshold for ChromoPainter's predictions, requiring at least 25
adjacent polymorphisms to be called introgressed in order to retain the candidate region. This
approach did succeed at reducing the false positive rate to 1.15%, though this is still substantially
higher than FILET's, but this improvement came at the cost of ChromoPainter's sensitivity,
which was reduced to 8.6%, roughly one-third that of FILET (Figure 2 and S8 Fig). We also
tried an intermediate threshold (5 polymorphisms), but did not observe a substantial increase in
specificity compared to our initial more lenient approach (8.49% false positive rate). Thus, while

we cannot rule out that it may be possible to devise a method to leverage ChromoPainter's model
to predict introgression that exceeds the performance of our application of ChromoPainter here,
our results suggest that it is unlikely that such an approach would eclipse the performance of
FILET. We note that ChromoPainter does have the advantage of not requiring simulated training
data. ChromoPainter also has the potential to identify donor and recipient haplotypes, which
FILET currently does not, but the far greater power of FILET demonstrated above will make it
preferable to many researchers who are interested in identifying introgressed regions. Moreover
our above results imply that predictions of the span and origin of introgressed haplotypes made
directly from ChromoPainter's output may not always be particularly reliable.

It is important to note that in the above simulations many introgressed regions will be
considerably smaller than our 10 kb window size. This fact, combined with our use of accuracy
measurements counting the number of individual base pairs correctly classified, makes the
results presented above useful for evaluating FILET's resolution and the impact of window size
on our predictions. By these measures FILET outperforms ChromoPainter, which does not use
windows and is only limited in scale by the density of polymorphisms. This suggests that when
using sliding windows FILET is able to achieve adequate resolution regardless of its use of a
predefined window size. Nonetheless we sought to improve our resolution further by using a
finer-scale FILET classifier trained on 1 kb windows as described above to refine the location of
putatively introgressed regions identified by the 10 kb classifier (see Methods). While this did
marginally reduce our false positive rates and increase our positive predictive values (see Figure
2 and S8 Fig), sensitivity was also somewhat reduced (to 25.7%; Figure 2). The relatively minor
effect of adding this refinement step reinforces the notion that a predefined window size is not a
major hindrance to our methods' effectiveness. Thus for most applications a window size that
yields enough polymorphisms to reliably calculate the statistics included in our feature vectors
may suffice.

Overall, FILET detects introgressed regions with greater power and resolution than
ChromoPainter, a method designed to detect ancestry tracks along recombining chromosomes.
However we note that many methods of this class exist, and it is possible that some may achieve
greater accuracy in some circumstances (e.g. if reference haplotype panels are used).

Sensitivity to continuous gene flow

While FILET is designed for identifying particular genomic windows that experienced
introgression as a result of a pulse migration event, genomic windows with genealogies that
include introgression may of course also be produced by continuous migration, with the timing
of geneflow varying from window to window. We therefore simulated genomic windows
experiencing a variety of bidirectional migration rates under each of our values of T_D and
recorded the fraction of windows for which our sampled individuals contained at least one
migrant lineage. Next, for each simulated window we applied the FILET classifier trained under
the appropriate divergence time as described above, recording the fraction of windows with at

least one migrant lineage that were classified as introgressed by FILET. The results of these tests
(S1 Table) show that for each value of T_D, the lowest bidirectional migration rates that we tested
do not produce migrant lineages, while higher rates will produce a small to modest fraction
migrants, most of which are undetected (e.g. when m=0.01, 23% and 59% of windows contain at
least one migrant when T_D=0.25 and T_D=1, respectively, but <5% are detected by FILET). Thus,
FILET, as currently trained, may not be sensitive to gene flow produced by low levels of
continuous migration. However, as the migration rate increases further, more and more of these
migrant windows will be detected (e.g. when m=1, 70% and 100% of windows are detected as
migrants by FILET when T_D=0.25 and T_D=1, respectively).
664

665 **Ranking the importance of statistics for detecting introgression**
666

Although our goal was to use a set of statistics to perform more accurate inference than possible
using individual ones, another benefit of our Extra-Trees approach is that it also allows for a
natural way to evaluate the extent to which different statistics are informative under different
scenarios of introgression. To this end, we used the Extra-Trees classifier to calculate feature
importance, which captures the ability of each statistic to separate the data into its respective
classes (Materials and Methods). We find that for our lowest T_D (a split N generations ago) the
top four features, all with similar importance, are the density of private alleles in population 1,
the density of private alleles in population 2, $d_{d-Rank1}$, and $d_{d-Rank2}$. For our next-lowest T_D ($4N$
generations ago), the top four, again with similar importance score estimates, are F_{ST}, Z_X, d_{d1}, and
d_{d2}. Thus our newly devised d_d and Z_X statistics seem to be particularly informative in the case of
recent introgression between closely related populations. For the larger values of T_D, d_{xy} and d_{min}
rise to prominence. The complete lists of feature importance for each T_D are shown in S2 Table.
The exceptional accuracy with which FILET uncovers introgressed loci underscores the
potential for machine learning methods to yield more powerful population genetic inferences
than can be achieved via more conventional tools which are often based on a single statistic.
Indeed, machine learning tools have been successfully leveraged in efforts to detect recent
positive selection [54, 104-107], to infer demographic histories [108], or even to perform both of
these tasks concurrently [109].
685

686 **Joint demographic history of *D. simulans* and *D. sechellia***
687

As described in the Materials and Methods, we used publically available *D. simulans* sequence
data [74], and collected and sequenced a set of *D. sechellia* genomes. We mapped reads from
these genomes to the *D. simulans* assembly [89], obtaining high coverage >28× for each
sequence (see sampling locations, mapping statistics, and Short Read Archive identifier
information listed in S3 Table). We do not expect that our reliance on the *D. simulans* assembly
resulted in any appreciable bias, as reads from our *D. sechellia* genomes were successfully
mapped to the reference genome at nearly the same rate as reads from *D. simulans* (S3 Table).

After completing variant calling and phasing (Materials and Methods), we performed
principal components analysis on intergenic SNPs at least 5 kb away from the nearest gene in
order to mitigate the bias introduced by linked selection [99, 110, 111]. While this is unlikely to
completely eliminate the confounding effect of linked selection in *Drosophila*, the fraction of
mutations that are deleterious is far greater in coding regions than in intergenic regions (~90%
versus <50% according to [112]); thus it is reasonable to presume that the impact of linked
selection is reduced several kilobases away from genes [113]. We observed evidence of
population structure within *D. sechellia*. In particular, the samples obtained from Praslin
clustered together, while all remaining samples formed a separate cluster (S9 FigA). Running
fastStructure [114] on this same set of SNPs yielded similar results: when the number of
subpopulations, *K*, was set to 2 (the optimal value for *K* selected by fastStructure's chooseK.py
script), our data were again subdivided into Praslin and non-Praslin clusters. Indeed, across all
values of *K* between 2 and 8, fastStructure's results were suggestive of marked subdivision
between Praslin and non-Praslin samples, and comparatively little subdivision within the non-
Praslin data (S9 FigB). This surprising result differs qualitatively from previous observations
from smaller numbers of loci [71, 115], and underscores the importance of using data from many
loci—preferably intergenic and genome-wide—in order to infer the presence or absence of
population structure.

Next, we examined the site frequency spectra of the Praslin and non-Praslin clusters,
noting that both had an excess of intermediate frequency alleles in comparison to that of the *D.*
simulans dataset (S10 Fig), in line with our expectations of *D. sechellia's* demographic history.
We also note that the Praslin samples contained far more variation (50,243 sites were
polymorphic within Praslin) than non-Praslin samples (4,108 SNPs within these samples). This
difference in levels of variation may reflect a much lesser degree of population structure and/or
inbreeding on the island of Praslin than across the other islands, or may result from other
demographic processes. Additional samples from across the Seychelles would be required to
address this question. In any case, in light of this observation we limited our downstream
analyses of *D. sechellia* sequences to those from Praslin.

Because we required a model from which to simulate training data for FILET, we next
inferred a joint demographic history of our population samples using ∂a∂i [95]. In particular, we
fit three demographic models to our dataset: an isolation-with-migration (IM) model allowing for
continuous population size change and migration following the population divergence, an
isolation model with the same parameters but fixing migration rates at zero, and an isolation
model with one burst of pulse migration from *D. simulans* into *D. sechellia* (Materials and
Methods). In S4 Table we show our model optimization results, which show clear support for the
IM model over the other models. The IM model that provided the best fit to our data (Figure 3A)
includes a much larger population size in *D simulans* than *D. sechellia* (a final size of 9.3×10^6
for *D. simulans* versus 2.6×10^4 for *D. sechellia*), consistent with the much greater diversity
levels in *D. simulans* [10, 71]. This model also exhibits a modest rate of migration, with a
substantially higher rate of gene flow from *D. simulans* to *D. sechellia* ($2\times N_{anc}m$=0.086) than

vice-versa ($2 \times N_{anc}m$=0.013). Thus, the results of our demographic modeling are consistent with
the observation of hybrid males in the Seychelles [73], and the possibility of recent introgression
between these two species across a substantial fraction of the genome (see refs. [72, 116]).

An interesting characteristic of the model shown in Figure 3A is that, assuming 15
generations per year, the estimated time of the *D. simulans*- *D. sechellia* population split is ~86
kya, or 1.3×10^6 generations ago. This contrasts with a recent estimate of 2.5×10^6 generations ago
from Garrigan et al. [72] which was based on single genomes rather than population genomic
data, but did account for ancestral polymorphism, as did estimates from Obbard et al. [117]
which yielded even older split times. Supporting our inference, we note that our average
intergenic cross-species divergence of 0.017 yields an average TMRCA of $\sim 2.5 \times 10^6$ generations
ago, assuming a mutation rate of 3.5×10^{-9} mutations per generation as observed in *D.*
melanogaster [112, 118], and this estimate would include the time before coalescence in the
ancestral population. Unless the mutation rate the *D. simulans* species complex is substantially
lower than in *D. melanogaster*, a population split time of 2.5×10^6 generations ago therefore
seems unlikely given that the ancestral population size (and therefore the period of time between
the population divergence and average TMRCA) was probably large (>500,000 by our estimate).
Thus, we conclude that the *D. simulans* and *D. sechellia* populations may have diverged more
recently than previously appreciated, perhaps within the last 100,000 years.

Although the specific parameterization of our model should be regarded as a preliminary
view of these species' demographic history that is adequate for the purposes of training FILET,
future efforts with larger sample sizes will be required to refine this model. That being said, the
basic features of this model—a much larger *D. simulans* population size than *sechellia*, and a
fairly large ancestral population size—are unlikely to change qualitatively.

Widespread introgression from *D. simulans* to *D. sechellia*

Accuracy and robustness of FILET under estimated model: Having obtained a suitable model of
the *D. simulans*- *D. sechellia* joint demographic history, we proceeded to simulate training data
and train FILET for application to our dataset (Materials and Methods). After training FILET
and applying it to simulated data under the estimated demographic model, we find that we have
good sensitivity to introgression (56% of windows with introgression are detected, on average),
and a false positive rate of only 0.2% (Figure 3B). Thus, while we may miss some introgressed
loci, we can have a great deal of confidence in the events that we do recover. Our feature
rankings for this classifier are included in S2 Table—under this scenario the most informative
feature is our newly devised d_{d-sim}. Note that we achieve high accuracy despite some of the
difficulties presented by the demographic model in Figure 3A, most notably the asymmetry in
effective population sizes between our two species. Indeed, because our method is trained under
this demographic history, such characteristics of genealogies produced under the assumed
demographic history (such as asymmetry in π) with and without introgression become the signals
used by FILET to make its classifications.

As shown in Figure 3B we find that this classifier has greater sensitivity to introgression
from *D. sechellia* to *D. simulans* than vice-versa. The cause of a stronger signal of *D. sechellia→*
D. simulans introgression can be understood from a consideration of the d_{min} statistic under each
of our three classes. When there is no introgression, d_{min} will be similar to the expected
divergence between *D. simulans* and *D. sechellia*; when there is introgression from *D. simulans*
to *D. sechellia*, we may expect d_{min} to be proportional to π_{sim}, which may only be a moderate
reduction relative to the no-introgression case given the large population size in *D. simulans*;
when there is introgression from *D. sechellia* to *D. simulans* then d_{min} is proportional to π_{sech}
which is dramatically lower than the expectation without introgression. While many of our
statistics do not rely on d_{min}, this example illustrates an important property of the genealogies
which include introgression from *D. sechellia* to *D. simulans* that would make them easier to
detect than gene flow in the reverse direction.
We also tested this classifier's performance on a different demographic scenario (S4
Table) in order to examine the effect of model misspecification during training. In particular, we
devised a simple island model with two population sizes: a larger size for *D. simulans* and the
ancestral population (7.6×10^5), and a smaller size for *D. sechellia* (5.7×10^4) with a split time of
~59 kya. Our simple procedure for estimating these values is described in the Materials and
Methods. Again, we find that we have good power to detect introgression with a very low false
positive rate (0.28%; S11 Fig). Although there are myriad incorrect models that we could test
FILET against, this example suggests that FILET's performance is robust to at least some
scenarios of demographic misspecification.
796
Application to population genomic data: We applied FILET to 10,185 non-overlapping 10 kb
windows that passed our data quality filters (101.85 Mb in total, or 86.7% of the five major
chromosome arms; Materials and Methods). FILET classified 267 windows as introgressed with
high-confidence, which we clustered into 94 contiguous regions accounting for 2.93% of the
accessible portion of the genome (2.99 Mb in total; Materials and Methods). This finding is
qualitatively similar to a previous estimate (4.6%) by Garrigan et al. [72] based on comparisons
of single genomes from each species in the *D. simulans* complex. Unlike this previous effort,
FILET is able to infer the directionality of introgression with high confidence (Figure 3B), and
we find evidence that the majority of this introgression has been in the direction of *D. simulans*
to *D. sechellia*: only 21 of the 267 (7.9%) putatively introgressed windows were classified as
introgressed from *sechellia* to *D. simulans*. This finding is not a result of a detection bias, as we
have greater power to detect gene flow from *D. sechellia* to *D. simulans* than in the reverse
direction. Given that our *D. simulans* sequences are from the mainland, one interpretation of this
result is that although there has been recent gene flow from *D. simulans* into the Seychelles,
where *D. simulans* and *D. sechellia* occasionally hybridize, there does not appear to be an
appreciable rate of back-migration to the mainland of *D. simulans* individuals harboring
haplotypes donated from *D. sechellia*. On the other hand, *D. sechellia* alleles may often be
purged from *D. simulans* by natural selection. This may be in part due to the reduced ecological

niche size of *D. sechellia*, such that any alleles which may introgress into *D. simulans* and lead to preference for or resistance to *Morinda* fruit may prove deleterious in other environments. More generally, *D. sechellia* haplotypes introgressing into *D. simulans* may harbor more deleterious alleles due to their smaller population size, which will be more effectively purged in the larger *D. simulans* population if mutations are not fully recessive [28]. Tests of these hypotheses will have to wait for a population sample of genomes from *D. simulans* collected in the Seychelles.

We asked whether our candidate introgressed loci were enriched for particular GO terms using a permutation test (Materials and Methods), finding no such enrichment. We did observe a deficit in the number of genes either partially overlapping or contained entirely within introgressed regions in our true set versus the permuted set; although a paucity of introgressed genes would be consistent with introgressed functional sequence often being deleterious, this difference was not significant (297 vs. 373.2, respectively; *P*=0.083; one-sided permutation test).

One notable introgressed region on 3R that FILET identified had been previously found by Garrigan et al. as containing a 15 kb region of introgression. We show that gene flow in this region actually extends for over 200 kb (Figure 4). When Brand et al. [119] sequenced the 15 kb region originally flagged by Garrigan et al. in a number of *D. simulans* and *D. sechellia* individuals, they also uncovered evidence of a selective sweep in *D. sechellia* originating from an adaptive introgression from *D. simulans*. Our data set also supports the presence of an adaptive introgression event at this locus: a 10 kb window lying within the putative sweep region (highlighted in Figure 4) is in the lower 5% tail of both d_{min} (consistent with introgression) and π_{sech} (consistent with a sweep in *sechellia*); this is the only window in the genome that is in the lower 5% tail for both of these statistics. This region contains two ionotropic glutamate receptors, *CG3822* and *Ir93a*, which may be involved in chemosensing among other functions [120], and the latter of which appears to play a role in resistance to entomopathogenic fungi [121]. Also near the trough of variation within *D. sechellia* are several members of the *Turandot* gene family, which are involved in humoral stress responses to various stressors including heat, UV light, and bacterial infection [122, 123], and perhaps parasitoid attack as well [124]. On the other hand, Brand et al. [119] hypothesize that the target of selection may be a transcription factor binding hotspot between *RpS30* and *CG15696*, and the phenotypic target of this sweep remains unclear.

Interestingly, this particular window is the only one in this region that is classified by FILET as having recent gene flow from *D. sechellia* to *D. simulans*. However this classification may be erroneous as one might expect FILET, which was not trained on any examples of adaptive introgression, to make an error in such a scenario because rather than gene flow increasing polymorphism in the recipient population, diversity is greatly diminished if the introgressed alleles rapidly sweep toward fixation. We note that this window is immediately flanked by a large number of windows classified as introgressed from *D. simulans* to *D. sechellia* and which show a large increase in diversity in the recipient population as expected. Moreover, Brand et al.'s phylogenetic analysis of introgression in this region also supported gene flow in this direction. Brand et al. also found evidence suggesting that the introgressed haplotype began

sweeping to higher frequency in *D. simulans* (though it has not reached fixation in this species)
prior to the timing of the introgression and subsequent sweep in *D. sechellia*. Thus we conclude
that the adaptive allele probably did indeed originate in *D. simulans* before migrating to *D.*
sechellia, and FILET's apparent error in this case underscores the genealogical differences
between adaptive gene flow and introgression events involving only neutral alleles.

Concluding remarks

Here we present a novel machine learning approach, FILET, that leverages population genomic
data from two related populations in order to determine whether a given genomic window has
experienced gene flow between these populations, and if so in which direction. We applied
FILET to a set of *D. simulans* genomes as well as a new set of whole genome sequences from the
closely related island endemic *D. sechellia*, confirming widespread introgression and also
inferring that this introgression was largely in the direction of *D. simulans* to *D. sechellia*. Future
work leveraging *D. simulans* data sampled from the Seychelles will be required to determine
whether this asymmetry is a consequence of low rate of migration of *D. simulans* back to
mainland Africa (where our *D. simulans* data were obtained), or whether the directionality of
gene flow is biased on the islands themselves. In addition to creating FILET, we devised several
new statistics, including the d_d statistics and Z_X which our feature rankings show to be quite
useful for uncovering gene flow.

Despite the success of FILET on both simulated data sets and real data from *Drosophila*,
there are several improvements that could be made. First, by framing the problem as one of
parameter estimation (i.e. regression) rather than classification, we may be able to precisely infer
the values of relevant parameters of introgression events (i.e. the time of the event and the
number of migrant lineages). Deep learning methods, which naturally allow for both
classification and regression, may prove particularly useful for this task [103]. Indeed, Sheehan
and Song [109] used deep learning to infer demographic parameters (regression) while
simultaneously identifying selective sweeps (classification). Another step we have not taken is to
explicitly handle adaptive introgression, which could potentially greatly improve our approach's
power to detect such events.

While population genetic inference has traditionally relied on the design of a summary
statistic sensitive to the evolutionary force of interest, a number of highly successful supervised
machine learning methods have been put forth within the last few years [54, 104-109]. These
methods are often thought of as black boxes, a characterization that may not always be fair [125].
Indeed in the context of evolutionary genetics such machine learning approaches are easily
interpreted as we have strong generative models that guide our intuition. Nonetheless, classical
statistical estimation from parametric models may often be more interpretable. Hybrid
approaches combining machine learning techniques with Bayesian approaches to estimate
posterior distributions of evolutionary parameters (e.g. [127]) thus represent an attractive
alternative to either approach in their "pure" form. As genomic data sets continue to grow, we

argue that machine learning approaches—in whatever shape they eventually take—leveraging high dimensional feature spaces have the potential to revolutionize evolutionary genomic inference.

ACKNOWLEDGMENTS

We thank Michael Lan for his work on an early iteration of this project. We also thank Joshua Schraiber and two anonymous reviewers for comments on the manuscript.

REFERENCES

1. Mallet J. Hybridization as an invasion of the genome. Trends in ecology & evolution. 2005;20(5):229-37.

2. Whitney KD, Ahern JR, Campbell LG, Albert LP, King MS. Patterns of hybridization in plants. Perspectives in Plant Ecology, Evolution and Systematics. 2010;12(3):175-82.

3. Barton NH. The role of hybridization in evolution. Mol Ecol. 2001;10(3):551-68.

4. Tung J, Barreiro LB. The contribution of admixture to primate evolution. Current opinion in genetics & development. 2017;47:61-8.

5. Baack EJ, Rieseberg LH. A genomic view of introgression and hybrid speciation. Current opinion in genetics & development. 2007;17(6):513-8.

6. Goulet BE, Roda F, Hopkins R. Hybridization in plants: old ideas, new techniques. Plant Physiol. 2017;173(1):65-78.

7. Gladieux P, Ropars J, Badouin H, Branca A, Aguileta G, Vienne DM, et al. Fungal evolutionary genomics provides insight into the mechanisms of adaptive divergence in eukaryotes. Mol Ecol. 2014;23(4):753-73.

8. Schardl C, Craven K. Interspecific hybridization in plant-associated fungi and oomycetes: a review. Mol Ecol. 2003;12(11):2861-73.

9. Brandvain Y, Kenney AM, Flagel L, Coop G, Sweigart AL. Speciation and introgression between Mimulus nasutus and Mimulus guttatus. PLoS Genet. 2014;10(6):e1004410.

10. Begun DJ, Holloway AK, Stevens K, Hillier LW, Poh Y-P, Hahn MW, et al. Population genomics: whole-genome analysis of polymorphism and divergence in Drosophila simulans. PLoS Biol. 2007;5(11):e310.

11. Kulathinal RJ, Stevison LS, Noor MA. The genomics of speciation in Drosophila: diversity, divergence, and introgression estimated using low-coverage genome sequencing. PLoS Genet. 2009;5(7):e1000550.

12. Martin SH, Dasmahapatra KK, Nadeau NJ, Salazar C, Walters JR, Simpson F, et al. Genome-wide evidence for speciation with gene flow in Heliconius butterflies. Genome Res. 2013;23(11):1817-28.

13.   Fontaine MC, Pease JB, Steele A, Waterhouse RM, Neafsey DE, Sharakhov IV, et al.
Extensive introgression in a malaria vector species complex revealed by phylogenomics.
Science. 2015;347(6217):1258524.

14.   Nürnberger B, Lohse K, Fijarczyk A, Szymura JM, Blaxter ML. Para-allopatry in
hybridizing fire-bellied toads (Bombina bombina and B. variegata): Inference from
transcriptome-wide coalescence analyses. Evolution. 2016;70(8):1803-18.

15.   Rothfels CJ, Johnson AK, Hovenkamp PH, Swofford DL, Roskam HC, Fraser-Jenkins
CR, et al. Natural hybridization between genera that diverged from each other
approximately 60 million years ago. The American Naturalist. 2015;185(3):433-42.

16.   Nadeau NJ, Ruiz M, Salazar P, Counterman B, Medina JA, Ortiz-Zuazaga H, et al.
Population genomics of parallel hybrid zones in the mimetic butterflies, H. melpomene
and H. erato. Genome Res. 2014;24(8):1316-33.

17.   Turissini DA, Matute DR. Fine scale mapping of genomic introgressions within the
Drosophila yakuba clade. bioRxiv. 2017:152421.

18.   Bachtrog D, Thornton K, Clark A, Andolfatto P. Extensive introgression of
mitochondrial DNA relative to nuclear genes in the Drosophila yakuba species group.
Evolution. 2006;60(2):292-302.

19.   Leavitt DH, Marion AB, Hollingsworth BD, Reeder TW. Multilocus phylogeny of
alligator lizards (Elgaria, Anguidae): Testing mtDNA introgression as the source of
discordant molecular phylogenetic hypotheses. Mol Phylogenet Evol. 2017;110:104-21.

20.   Sarver BA, Demboski JR, Good JM, Forshee N, Hunter SS, Sullivan J. Comparative
phylogenomic assessment of mitochondrial introgression among several species of
chipmunks (Tamias). Genome Biol Evol. 2016;9(1):7-19.

21.   Carneiro M, Albert FW, Afonso S, Pereira RJ, Burbano H, Campos R, et al. The genomic
architecture of population divergence between subspecies of the European rabbit. PLoS
Genet. 2014;10(8):e1003519.

22.   Maroja LS, Larson EL, Bogdanowicz SM, Harrison RG. Genes with restricted
introgression in a field cricket (Gryllus firmus/Gryllus pennsylvanicus) hybrid zone are
concentrated on the X chromosome and a single autosome. G3: Genes, Genomes,
Genetics. 2015;5(11):2219-27.

23.   Muirhead CA, Presgraves DC. Hybrid incompatibilities, local adaptation, and the
genomic distribution of natural introgression between species. The American Naturalist.
2016;187(2):249-61.

24.   Phifer-Rixey M, Bomhoff M, Nachman MW. Genome-wide patterns of differentiation
among house mouse subspecies. Genetics. 2014;198(1):283-97.

25.   Green RE, Krause J, Briggs AW, Maricic T, Stenzel U, Kircher M, et al. A draft
sequence of the Neandertal genome. Science. 2010;328(5979):710-22.

26.   Sankararaman S, Mallick S, Dannemann M, Prüfer K, Kelso J, Pääbo S, et al. The
genomic landscape of Neanderthal ancestry in present-day humans. Nature.
2014;507(7492):354-7.

27. Turner TL, Hahn MW, Nuzhdin SV. Genomic islands of speciation in Anopheles gambiae. PLoS Biol. 2005;3(9):e285.

28. Harris K, Nielsen R. The genetic cost of Neanderthal introgression. Genetics. 2016;203(2):881-91.

29. Juric I, Aeschbacher S, Coop G. The strength of selection against Neanderthal introgression. PLoS Genet. 2016;12(11):e1006340.

30. Hedrick PW. Adaptive introgression in animals: examples and comparison to new mutation and standing variation as sources of adaptive variation. Mol Ecol. 2013;22(18):4606-18.

31. Norris LC, Main BJ, Lee Y, Collier TC, Fofana A, Cornel AJ, et al. Adaptive introgression in an African malaria mosquito coincident with the increased usage of insecticide-treated bed nets. Proceedings of the National Academy of Sciences. 2015;112(3):815-20.

32. Pardo-Diaz C, Salazar C, Baxter SW, Merot C, Figueiredo-Ready W, Joron M, et al. Adaptive introgression across species boundaries in Heliconius butterflies. PLoS Genet. 2012;8(6):e1002752.

33. Song Y, Endepols S, Klemann N, Richter D, Matuschka F-R, Shih C-H, et al. Adaptive introgression of anticoagulant rodent poison resistance by hybridization between old world mice. Curr Biol. 2011;21(15):1296-301.

34. Bechsgaard J, Jorgensen TH, Schierup MH. Evidence for Adaptive Introgression of Disease Resistance Genes Among Closely Related Arabidopsis Species. G3: Genes, Genomes, Genetics. 2017;7(8):2677-83.

35. Cheeseman K, Ropars J, Renault P, Dupont J, Gouzy J, Branca A, et al. Multiple recent horizontal transfers of a large genomic region in cheese making fungi. Nature Communications. 2014;5:2876.

36. Huerta-Sánchez E, Jin X, Bianba Z, Peter BM, Vinckenbosch N, Liang Y, et al. Altitude adaptation in Tibetans caused by introgression of Denisovan-like DNA. Nature. 2014;512(7513):194-7.

37. Melo MC, Salazar C, Jiggins CD, Linares M. Assortative mating preferences among hybrids offers a route to hybrid speciation. Evolution. 2009;63(6):1660-5.

38. Salazar C, Baxter SW, Pardo-Diaz C, Wu G, Surridge A, Linares M, et al. Genetic evidence for hybrid trait speciation in Heliconius butterflies. PLoS Genet. 2010;6(4):e1000930.

39. Alexander DH, Novembre J, Lange K. Fast model-based estimation of ancestry in unrelated individuals. Genome Res. 2009;19(9):1655-64.

40. Anderson E, Thompson E. A model-based method for identifying species hybrids using multilocus genetic data. Genetics. 2002;160(3):1217-29.

41. Pritchard JK, Stephens M, Donnelly P. Inference of population structure using multilocus genotype data. Genetics. 2000;155(2):945-59.

42.   Pickrell JK, Pritchard JK. Inference of population splits and mixtures from genome-wide
allele frequency data. PLoS Genet. 2012;8(11):e1002967.

43.   Guan Y. Detecting structure of haplotypes and local ancestry. Genetics. 2014;196(3):625-
42.

44.   Tang H, Coram M, Wang P, Zhu X, Risch N. Reconstructing genetic ancestry blocks in
admixed individuals. The American Journal of Human Genetics. 2006;79(1):1-12.

45.   Sohn K-A, Ghahramani Z, Xing EP. Robust estimation of local genetic ancestry in
admixed populations using a nonparametric Bayesian approach. Genetics.
2012;191(4):1295-308.

46.   Lawson DJ, Hellenthal G, Myers S, Falush D. Inference of population structure using
dense haplotype data. PLoS Genet. 2012;8(1):e1002453.

47.   Price AL, Tandon A, Patterson N, Barnes KC, Rafaels N, Ruczinski I, et al. Sensitive
detection of chromosomal segments of distinct ancestry in admixed populations. PLoS
Genet. 2009;5(6):e1000519.

48.   Wright S. The genetical structure of populations. Ann Hum Genet. 1949;15(1):323-54.

49.   Nei M, Li W-H. Mathematical model for studying genetic variation in terms of restriction
endonucleases. Proceedings of the National Academy of Sciences. 1979;76(10):5269-73.

50.   Joly S, McLenachan PA, Lockhart PJ. A statistical approach for distinguishing
hybridization and incomplete lineage sorting. The American Naturalist. 2009;174(2):E54-
E70.

51.   Geneva AJ, Muirhead CA, Kingan SB, Garrigan D. A new method to scan genomes for
introgression in a secondary contact model. PLoS ONE. 2015;10(4):e0118621.

52.   Rosenzweig BK, Pease JB, Besansky NJ, Hahn MW. Powerful methods for detecting
introgressed regions from population genomic data. Mol Ecol. 2016;25(11):2387-97. doi:
10.1111/mec.13610.

53.   Geurts P, Ernst D, Wehenkel L. Extremely randomized trees. Machine Learning.
2006;63(1):3-42.

54.   Schrider DR, Kern AD. S/HIC: Robust Identification of Soft and Hard Sweeps Using
Machine Learning. PLoS Genet. 2016;12(3): e1005928.

55.   Schrider DR, Kern AD. Soft sweeps are the dominant mode of adaptation in the human
genome. Mol Biol Evol. 2017;34(8):1863–77. Epub 2017/05/10. doi:
10.1093/molbev/msx154. PubMed PMID: 28482049.

56.   Jones CD. The genetic basis of Drosophila sechellia's resistance to a host plant toxin.
Genetics. 1998;149(4):1899-908.

57.   Jones CD. The genetics of adaptation in Drosophila sechellia. Genetica. 2005;123(1-
2):137.

58.   Louis J, David J. Ecological specialization in the Drosophila melanogaster species
subgroup: a case study of D. sechellia. Acta oecologica Oecologia generalis.
1986;7(3):215-29.

59. Farine J-P, Legal L, Moreteau B, Le Quere J-L. Volatile components of ripe fruits of
Morinda citrifolia and their effects on Drosophila. Phytochemistry. 1996;41(2):433-8.

60. Legal L, Chappe B, Jallon JM. Molecular basis ofMorinda citrifolia (L.): Toxicity on
drosophila. J Chem Ecol. 1994;20(8):1931-43.

61. Legal L, Moulin B, Jallon JM. The relation between structures and toxicity of oxygenated
aliphatic compounds homologous to the insecticide octanoic acid and the chemotaxis of
two species of Drosophila. Pestic Biochem Physiol. 1999;65(2):90-101.

62. Andrade López J, Lanno S, Auerbach J, Moskowitz E, Sligar L, Wittkopp P, et al.
Genetic basis of octanoic acid resistance in Drosophila sechellia: functional analysis of a
fine-mapped region. Mol Ecol. 2017;26(4):1148-60.

63. Dekker T, Ibba I, Siju K, Stensmyr MC, Hansson BS. Olfactory shifts parallel
superspecialism for toxic fruit in Drosophila melanogaster sibling, D. sechellia. Curr
Biol. 2006;16(1):101-9.

64. Huang Y, Erezyilmaz D. The genetics of resistance to Morinda fruit toxin during the
postembryonic stages in Drosophila sechellia. G3: Genes, Genomes, Genetics.
2015;5(10):1973-81.

65. Hungate EA, Earley EJ, Boussy IA, Turissini DA, Ting C-T, Moran JR, et al. A locus in
Drosophila sechellia affecting tolerance of a host plant toxin. Genetics.
2013;195(3):1063-75.

66. Matsuo T, Sugaya S, Yasukawa J, Aigaki T, Fuyama Y. Odorant-binding proteins
OBP57d and OBP57e affect taste perception and host-plant preference in Drosophila
sechellia. PLoS Biol. 2007;5(5):e118.

67. Shiao M-S, Chang J-M, Fan W-L, Lu M-YJ, Notredame C, Fang S, et al. Expression
divergence of chemosensory genes between Drosophila sechellia and its sibling species
and its implications for host shift. Genome Biol Evol. 2015;7(10):2843-58.

68. Hey J, Kliman RM. Population genetics and phylogenetics of DNA sequence variation at
multiple loci within the Drosophila melanogaster species complex. Mol Biol Evol.
1993;10(4):804-22.

69. Kern AD, Jones CD, Begun DJ. Molecular population genetics of male accessory gland
proteins in the Drosophila simulans complex. Genetics. 2004;167(2):725-35.

70. Kliman RM, Andolfatto P, Coyne JA, Depaulis F, Kreitman M, Berry AJ, et al. The
population genetics of the origin and divergence of the Drosophila simulans complex
species. Genetics. 2000;156(4):1913-31.

71. Legrand D, Tenaillon MI, Matyot P, Gerlach J, Lachaise D, Cariou M-L. Species-wide
genetic variation and demographic history of Drosophila sechellia, a species lacking
population structure. Genetics. 2009;182(4):1197-206.

72. Garrigan D, Kingan SB, Geneva AJ, Andolfatto P, Clark AG, Thornton KR, et al.
Genome sequencing reveals complex speciation in the Drosophila simulans clade.
Genome Res. 2012;22(8):1499-511.

73.   Matute D, Ayroles J. Hybridization occurs between Drosophila simulans and D. sechellia
in the Seychelles archipelago. J Evol Biol. 2014;27(6):1057-68.

74.   Rogers RL, Cridland JM, Shao L, Hu TT, Andolfatto P, Thornton KR. Landscape of
standing variation for tandem duplications in Drosophila yakuba and Drosophila
simulans. Mol Biol Evol. 2014;31(7):1750-66.

75.   Feder JL, Xie X, Rull J, Velez S, Forbes A, Leung B, et al. Mayr, Dobzhansky, and Bush
and the complexities of sympatric speciation in Rhagoletis. Proceedings of the National
Academy of Sciences. 2005;102(suppl 1):6573-80.

76.   Kelly JK. A test of neutrality based on interlocus associations. Genetics.
1997;146(3):1197-206.

77.   Breiman L. Random forests. Machine Learning. 2001;45(1):5-32.

78.   Quinlan JR. Induction of decision trees. Machine Learning. 1986;1(1):81-106.

79.   Fay JC, Wu C-I. Hitchhiking under positive Darwinian selection. Genetics.
2000;155(3):1405-13.

80.   Tajima F. Statistical method for testing the neutral mutation hypothesis by DNA
polymorphism. Genetics. 1989;123(3):585-95.

81.   Hudson RR, Slatkin M, Maddison W. Estimation of levels of gene flow from DNA
sequence data. Genetics. 1992;132(2):583-9.

82.   Hudson RR. A new statistic for detecting genetic differentiation. Genetics.
2000;155(4):2011-4.

83.   Patterson N, Moorjani P, Luo Y, Mallick S, Rohland N, Zhan Y, et al. Ancient admixture
in human history. Genetics. 2012;192(3):1065-93.

84.   Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O, et al. Scikit-
learn: Machine learning in Python. Journal of Machine Learning Research.
2011;12(Oct):2825-30.

85.   Breiman L, Friedman J, Stone CJ, Olshen RA. Classification and regression trees: CRC
press; 1984.

86.   Chan AH, Jenkins PA, Song YS. Genome-wide fine-scale recombination rate variation in
Drosophila melanogaster. PLoS Genet. 2012;8(12):e1003090.

87.   Li N, Stephens M. Modeling linkage disequilibrium and identifying recombination
hotspots using single-nucleotide polymorphism data. Genetics. 2003;165(4):2213-33.

88.   Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
arXiv. 2013. doi: 1303.3997.

89.   Hu TT, Eisen MB, Thornton KR, Andolfatto P. A second-generation assembly of the
Drosophila simulans genome provides new insights into patterns of lineage-specific
divergence. Genome Res. 2013;23(1):89-98.

90.   Gramates LS, Marygold SJ, Santos Gd, Urbano J-M, Antonazzo G, Matthews BB, et al.
FlyBase at 25: looking to the future. Nucleic Acids Res. 2017;45(D1):D663-71.

91. McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, et al. The
Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA
sequencing data. Genome Res. 2010;20(9):1297-303.

92. DePristo MA, Banks E, Poplin R, Garimella KV, Maguire JR, Hartl C, et al. A
framework for variation discovery and genotyping using next-generation DNA
sequencing data. Nat Genet. 2011;43(5):491-8.

93. Auwera GA, Carneiro MO, Hartl C, Poplin R, del Angel G, Levy-Moonshine A, et al.
From FastQ data to high-confidence variant calls: the genome analysis toolkit best
practices pipeline. Current protocols in bioinformatics. 2013;43:11.0. 1-.0. 33.

94. Delaneau O, Zagury J-F, Marchini J. Improved whole-chromosome phasing for disease
and population genetic studies. Nat Methods. 2013;10(1):5-6.

95. Gutenkunst RN, Hernandez RD, Williamson SH, Bustamante CD. Inferring the joint
demographic history of multiple populations from multidimensional SNP frequency data.
PLoS Genet. 2009;5(10):e1000695.

96. Jansen PW, Perez RE. Constrained structural design optimization via a parallel
augmented Lagrangian particle swarm optimization approach. Computers & Structures.
2011;89(13):1352-66.

97. Kraft D. A software package for sequential quadratic programming: DFVLR
Obersfaffeuhofen, Germany; 1988.

98. Perez RE, Jansen PW, Martins JR. pyOpt: a Python-based object-oriented framework for
nonlinear constrained optimization. Structural and Multidisciplinary Optimization.
2012;45(1):101-18.

99. Schrider DR, Shanku AG, Kern AD. Effects of Linked Selective Sweeps on
Demographic Inference and Model Selection. Genetics. 2016;204(3):1207-23. doi:
10.1534/genetics.116.190223.

100. Pool JE. The mosaic ancestry of the Drosophila genetic reference panel and the D.
melanogaster reference genome reveals a network of epistatic fitness interactions. Mol
Biol Evol. 2015;32(12):3236-51.

101. Schrider DR, Kern AD. Supervised Machine Learning for Population Genetics: A New
Paradigm. Trends Genet. 2018:10.1016/j.tig.2017.12.005.

102. Cortes C, Vapnik V. Support-vector networks. Machine Learning. 1995;20(3):273-97.

103. LeCun Y, Bengio Y, Hinton G. Deep learning. Nature. 2015;521(7553):436-44.

104. Lin K, Li H, Schlötterer C, Futschik A. Distinguishing positive selection from neutral
evolution: boosting the performance of summary statistics. Genetics. 2011;187(1):229-
44.

105. Pavlidis P, Jensen JD, Stephan W. Searching for footprints of positive selection in whole-
genome SNP data from nonequilibrium populations. Genetics. 2010;185(3):907-22.

106. Pybus M, Luisi P, Dall'Olio GM, Uzkudun M, Laayouni H, Bertranpetit J, et al.
Hierarchical boosting: a machine-learning framework to detect and classify hard selective
sweeps in human populations. Bioinformatics. 2015;31(24):3946-52.

107. Ronen R, Udpa N, Halperin E, Bafna V. Learning natural selection from the site
frequency spectrum. Genetics. 2013;195(1):181-93.

108. Pudlo P, Marin J-M, Estoup A, Cornuet J-M, Gautier M, Robert CP. Reliable ABC model
choice via random forests. Bioinformatics. 2016;32(6):859-66.

109. Sheehan S, Song YS. Deep learning for population genetic inference. PLoS Comput Biol.
2016;12(5):e1004845.

110. Gazave E, Ma L, Chang D, Coventry A, Gao F, Muzny D, et al. Neutral genomic regions
refine models of recent rapid human population growth. Proceedings of the National
Academy of Sciences. 2014;111(2):757-62.

111. Ewing GB, Jensen JD. The consequences of not accounting for background selection in
demographic inference. Mol Ecol. 2016;25(1):135-41.

112. Schrider DR, Houle D, Lynch M, Hahn MW. Rates and genomic consequences of
spontaneous mutational events in Drosophila melanogaster. Genetics. 2013;194(4):937-
54.

113. Langley CH, Stevens K, Cardeno C, Lee YCG, Schrider DR, Pool JE, et al. Genomic
variation in natural populations of *Drosophila melanogaster*. Genetics. 2012;192(2):533-
98.

114. Raj A, Stephens M, Pritchard JK. fastSTRUCTURE: variational inference of population
structure in large SNP data sets. Genetics. 2014;197(2):573-89.

115. Legrand D, Vautrin D, Lachaise D, Cariou M-L. Microsatellite variation suggests a
recent fine-scale population structure of Drosophila sechellia, a species endemic of the
Seychelles archipelago. Genetica. 2011;139(7):909.

116. Navascués M, Legrand D, Campagne C, Cariou M-L, Depaulis F. Distinguishing
migration from isolation using genes with intragenic recombination: detecting
introgression in the Drosophila simulans species complex. BMC Evol Biol.
2014;14(1):89.

117. Obbard DJ, Maclennan J, Kim K-W, Rambaut A, O'Grady PM, Jiggins FM. Estimating
divergence dates and substitution rates in the Drosophila phylogeny. Mol Biol Evol.
2012;29(11):3459-73.

118. Keightley PD, Trivedi U, Thomson M, Oliver F, Kumar S, Blaxter M. Analysis of the
genome sequences of three Drosophila melanogaster spontaneous mutation accumulation
lines. Genome Res. 2009;19:1195-201.

119. Brand CL, Kingan SB, Wu L, Garrigan D. A selective sweep across species boundaries in
Drosophila. Mol Biol Evol. 2013;30(9):2177-86.

120. Benton R, Vannice KS, Gomez-Diaz C, Vosshall LB. Variant ionotropic glutamate
receptors as chemosensory receptors in Drosophila. Cell. 2009;136(1):149-62.

121. Lu H-L, Wang JB, Brown MA, Euerle C, Leger RJS. Identification of Drosophila
mutants affecting defense to an entomopathogenic fungus. Scientific reports. 2015;5.

122. Ekengren S, Hultmark D. A family of Turandot-related genes in the humoral stress
response of Drosophila. Biochem Biophys Res Commun. 2001;284(4):998-1003.

123.   Ekengren S, Tryselius Y, Dushay MS, Liu G, Steiner H, Hultmark D. A humoral stress
1209   response in Drosophila. Curr Biol. 2001;11(9):714-8.

1210   124.   Salazar-Jaramillo L, Jalvingh KM, de Haan A, Kraaijeveld K, Buermans H, Wertheim B.
1211   Inter-and intra-species variation in genome-wide gene expression of Drosophila in
1212   response to parasitoid wasp attack. BMC Genomics. 2017;18(1):331.

1213   125.   Breiman L. Statistical modeling: The two cultures (with comments and a rejoinder by the
1214   author). Statistical science. 2001;16(3):199-231.

1215   126.   Baehrens D, Schroeter T, Harmeling S, Kawanabe M, Hansen K, MÃžller K-R. How to
1216   explain individual classification decisions. Journal of Machine Learning Research.
1217   2010;11(Jun):1803-31.

1218   127.   Blum MG, François O. Non-linear regression models for Approximate Bayesian
1219   Computation. Statistics and Computing. 2010;20(1):63-73.

1220

Fig. 1. Heatmaps showing several methods' sensitivity to detect introgression. We show the fraction of simulated genomic regions with introgression occurring under various combinations of migration times (T_M, shown as a fraction of the population divergence time T_D) and intensities (P_M, the probability that a given lineage will be included in the introgression event) that are detected successfully by each method. (*A*) Accuracy of d_{min} and G_{min} statistics, where a simulated region is classified as introgressed if the values of these statistics are found in the lower 5% tail of the distribution under complete isolation (from simulations). Thus, the false positive rate is fixed at 5%. (*B*) The accuracy of FILET on these same simulations. On the left we show the fraction of regions correctly classified as introgressed (compare to panel *A*). On the right, we show the fraction of all simulated regions that are not only classified as introgressed, but also for which the direction of gene flow was correctly inferred (i.e. if the direction is inferred with 100% accuracy for a given cell in the heatmap, the color shade of that cell will be identical to that in the heatmap on the left). The false positive rate, as determined from applying FILET to a simulated test set with no migration, is also shown.

Fig. 2. A comparison of the power and resolution of FILET and ChromoPainter using simulations of a 1 Mb chromosome where introgression was allowed within the central 100 kb region. As in Figure 1, the population split time was set to N generations ago, and the darkness of the heatmap shows sensitivity to introgression. Unlike Figure 1, here we are measuring sensitivity at the level of the individual base pair rather than evaluating the question of whether a window at large was recovered as containing introgressed alleles. The "coarse" version of FILET refers to a FILET classifier trained to detect introgression in 10 kb windows, which was applied to sliding windows (1 kb step size) across the chromosome. The "fine" version of FILET applied a classifier trained on 1 kb windows to sliding windows (100 bp step size) within those regions classified as introgressed by the FILET classifier. The lenient version of ChromoPainter required evidence of introgression at a single SNP to identify introgression, while the stringent version required candidate regions to contain at least 25 consecutive SNPs supporting introgression.

Fig. 3. Inferred joint population history of *D. simulans* and *D. sechellia*, and power to detect introgression under this model. (*A*) The parameterization of our best-fitting demographic model. Migration rates are shown by arrows, and are in units of $2 \times N_{anc}m$, where m is the probability of migration per individual in the source population per generation. (*B*) Confusion matrix showing FILET's classification accuracy under this model as assessed on an independent simulated test set. Perfect accuracy would be 100% along the entire diagonal from top-left to bottom-right, and the false positive rate is the sum of top-middle and top-right cells.

Fig. 4. A large genomic region on 3R classified by FILET as introgressed from *D. simulans* to *D. sechellia*. Values of the $d_{d\text{-}sim}$ and d_{min} (upper two panels) within each 10 kb window in the region are shown, along with the posterior probability of introgression from FILET (i.e. $1 - P$(no introgression)). Clustered regions classified as introgressed are shown as gray rectangles superimposed over these probabilities. Also shown are windowed values of π in *D. sechellia*, with the sweep region highlighted in red, and the locations of annotated genes with associated FlyBase identifiers [90].

SUPPLEMENTAL FIGURE AND TABLE LEGENDS

S1 Fig. Illustration of the difference in values of the d_{min} statistic calculated from joint population samples with and without introgression.

S2 Fig. Violin plots showing the values of d_{min}, all four d_d statistics, and Z_X under simulated scenarios including introgression or lacking it for each values of T_D. The values of these statistics were obtained from the training data sets described in the Materials and Methods.

S3 Fig. Heatmaps showing several methods' sensitivity to detect introgression. Same as Figure 1, but for other values of T_D. (*A*) Accuracy for d_{min} and G_{min} when $T_D = 1 \times 4N$ generations. (*B*) Accuracy of FILET when $T_D = 1 \times 4N$. (*C*) and (*D*) show the same when $T_D = 4 \times 4N$. (E) and (F) show the same when $16 \times 4N$.

S4 Fig. Heatmaps showing FILET's sensitivity to introgression from an unsampled ghost population. (*A*) Sensitivity when $T_D = 0.25 \times 4N$ generations. (*B*) Sensitivity when $T_D = 1 \times 4N$ generations. (*C*) $T_D = 4 \times 4N$ generations. (*D*) $T_D = 16 \times 4N$.

S5 Fig. ROC curves showing power of FILET, d_{min} and G_{min} under each value of T_D. In order to generate these curves we transformed the classification task into a binary one: discriminating between isolation and introgression in either direction. (*A*) $T_D = 0.25 \times 4N$ generations. (*B*) $T_D = 1 \times 4N$ generations. (*C*) $T_D = 4 \times 4N$. (*D*) $T_D = 16 \times 4N$. Training and test sets for these problems contained equal numbers of examples of introgression from population 1 into 2 and introgression from population 2 into 1.

S6 Fig. ROC curves showing power of versions of FILET trained with decreasing numbers of training instances (ranging from 100 to 10000 for each class). (*A*) $T_D = 0.25 \times 4N$ generations. (*B*) $T_D = 1 \times 4N$ generations. (*C*) $T_D = 4 \times 4N$. (*D*) $T_D = 16 \times 4N$.

S7 Fig. ROC curve showing FILET's power when trained and tested on simulated examples with different window sizes with $T_D = 0.25 \times 4N$ generations.

S8 Fig. A comparison of the positive predictive value of FILET and ChromoPainter on the same simulated data used for Fig 2. The "coarse" and "fine" versions of FILET, and lenient and stringent versions of ChromoPainter's predictions, are as defined for Fig 2. In cases where the positive predictive value is undefined (i.e. no base pairs were predicted to be introgressed), it is displayed as zero (i.e. a white cell in the heatmap).

S9 Fig. Population structure within *D. sechellia.* (*A*) The top three principal components of all *D. sechellia* diploid genomes. The cluster on the left shows the individuals from Praslin, while the

cluster on the right shows all other individuals. Note that the cluster on the right is far less dispersed due to the very small amount of polymorphism among these individuals. The numbers in parentheses on each axis show the fraction of the variance explained by each principal component. (*B*) Results of running fastStructure on our *D. sechellia* samples with the number of subpopulations (*K*) ranging from 2 to 8.

S10 Fig. Site frequency spectra of *D. sechellia* samples from Praslin, *D. sechellia* samples from all other locations, and *D. simulans* samples. The *D. sechellia* samples were both downsampled to *n*=12 as described in the text, while *D. simulans* was downsampled to *n*=18 (i.e. the same sample sizes used for our demographic inference). These SFS show the fraction of all polymorphisms found in each bin rather than the raw number of polymorphisms, and thus do not contain information about the total number of SNPs. As described in the text, there is >12-fold more polymorphism in the Praslin samples than in the non-Praslin samples.

S11 Fig. Confusion matrix showing FILET's classification accuracy when trained under out inferred model of the *simulans-sechellia* joint demographic history, but applied to test data generated under a different model (described in Materials and Methods and shown in S4 Table). under this model as assessed on an independent simulated test set. Perfect accuracy would be 100% along the entire diagonal from top-left to bottom-right, and the false positive rate is the sum of top-middle and top-right cells.

S1 Table. Results when applying FILET to simulations with constant bidirectional migration. 10000 simulated replicates were tested for each parameter combination.

S2 Table. Feature importance and rankings for each classifier used in this study.

S3 Table. Sampling location, sequencing/mapping statistics, and SRA identifiers for each genome included in this study.

S4 Table. Demographic parameter estimates inferred by ∂a∂i, along with a simple naïve model.