qDSB-Seq: quantitative DNA double-strand break sequencing

Yingjie Zhu[1,9], Anna Biernacka[2,9], Benjamin Pardo[3], Norbert Dojer[1,4], Romain Forey[3], Magdalena Skrzypczak[3], Bernard Fongang[1], Jules Nde[1], Raziyeh Yousefi[1], Philippe Pasero[3], Krzysztof Ginalski[2] and Maga Rowicka[1,5, 6,7,8]

[1] Department of Biochemistry and Molecular Biology, University of Texas Medical Branch at Galveston, Galveston, Texas, USA.

[2] Laboratory of Bioinformatics and Systems Biology, Centre of New Technologies, University of Warsaw, Warsaw, Poland.

[3] Institute of Human Genetics, Montpellier, France.

[4] Institute of Informatics, University of Warsaw, Warsaw, Poland.

[5] Institute for Translational Sciences, University of Texas Medical Branch at Galveston, Galveston, Texas, USA.

[6] Sealy Center for Molecular Medicine, University of Texas Medical Branch at Galveston, Galveston, Texas, USA.

[7] Sealy Center for Structural Biology and Molecular Biophysics, University of Texas Medical Branch at Galveston, Galveston, Texas, USA.

[8] Correspondence should be addressed to M.R. (Maga.Rowicka@utmb.edu)

[9] These authors contributed equally

Short title: Quantitative DSB sequencing

Keywords: DNA double-stranded break, quantitative DSB sequencing, restriction enzyme, Zeocin, hydroxyurea, camptothecin, replication fork, fork barrier

Abstract

Sequencing-based methods for mapping DNA double-strand breaks (DSBs) allow measurement only of relative frequencies of DSBs between loci, which limits our understanding of the physiological relevance of detected DSBs. We propose quantitative DSB sequencing (qDSB-Seq), a method providing both DSB frequencies per cell and their precise genomic coordinates. We induced spike-in DSBs by a site-specific endonuclease and used them to quantify labeled DSBs (e.g. using i-BLESS). Utilizing qDSB-Seq, we determined numbers of DSBs induced by a radiomimetic drug and various forms of replication stress, and revealed several orders of magnitude differences in DSB frequencies. We also measured for the first time Top1-dependent absolute DSB frequencies at replication fork barriers. qDSB-Seq is compatible with various DSB labeling methods in different organisms and allows accurate comparisons of absolute DSB frequencies across samples.

Introduction

There is tremendous interest in precisely measuring DNA double-strand breaks (DSBs) genome-wide since such measurement can give key insights into DNA damage and repair,

cancer development[1], radiation biology, and also increasingly popular genome editing
techniques[2]. Starting with our BLESS method[3], several high-resolution and direct methods
to label DSBs genome-wide have recently been developed[4-7], which have opened up new
possibilities for sensitive and specific detection of DSBs. For example, BLESS was applied
in identifying the on-target and off-target cutting sites of Cas9 endonuclease[8] and studying
DSB repair[9]. However, we still lack an effective strategy to both precisely detect DSB
distribution genome-wide and quantify their absolute frequencies per cell, which is crucial
to assess physiological relevance of detected DSBs. Immunofluorescence microscopy in
combination with γ-H2AX and 53BP1 antibodies was used to count breaks per cell[10], but
does not allow determining their precise locations. Moreover, counting discrete nuclear
foci is an imprecise way to estimate DSB numbers per cell both due to DSB clustering and
limited specificity of antibodies. Quantitative Polymerase Chain Reaction (qPCR) based
methods can estimate absolute break frequency but only at selected loci[11]. An approach
was developed recently to quantify breaks globally based on amount of radiolabeled DNA
and locally based on DNA break immunocapture[12], but its accuracy in detecting
physiological DSBs was not tested. BLISS[7] quantifies DSBs by utilizing unique molecular
identifiers (UMIs) to identify unique DSB ends and counting cells in the sample. BLISS is
designed for detecting DSBs in samples with low number of cells and thus shares
challenges of single-cell sequencing, such as low genome coverage and over-amplification.
Moreover, employment of UMIs is challenging. Short UMIs may lead to UMI collisions[13]
(i.e. observing two reads with the same sequence and the same UMI barcode but originating
from two different genomic molecules), especially in case of DSBs enriched in specific
locations. Long UMIs may interfere with primer sequence binding and accumulate
sequencing errors, which may lead to severe overestimation of DSBs[14].

This lack of a general method and computational solution to simultaneously determine
DSB frequencies per cell and their precise genomic loci limits our understanding of the
physiological relevance of observed DSBs and hinders comparisons between experiments.
Here, we propose quantitative DSB sequencing (qDSB-Seq), an approach that allows
measuring DSB frequencies per cell genome-wide, and a computational solution to achieve
accurate quantification. Our approach relies on inducing spike-in DSBs by a site-specific
endonuclease, which are used to quantify DSBs detected by a DSB labeling method e.g. i-
BLESS[15] and can be combined with any DSB labeling technique. We present a
comprehensive validation and several applications of qDSB-Seq: quantifying DSBs
induced by a radiomimetic drug, occurring during replication stress and caused by natural
replication fork barriers.

Results

qDSB-Seq implementation, computational method and validation. qDSB-Seq is a
combination of genome-wide high-resolution DSB-labeling (i-BLESS[15], BLESS[3], END-
seq[6], etc.) and inducing DSBs (spike-ins) in pre-determined loci using a site-specific
endonuclease (**Fig. 1a-c**). Quantification is based on an assumption (verified below) that
the number of labeled reads at a given genomic locus resulting from DSB sequencing is
proportional to the underlying DSB frequency (proportionality coefficient α in **Fig. 2a**).

To estimate this coefficient α, we induce spike-in DSBs at pre-determined genomic loci
and, relying on knowledge of their exact genomic locations, quantify their frequency using

genomic DNA sequencing (gDNA) or qPCR (**Fig. 1a, Fig. 2a**). The spike-in DSBs are
created by digestion with a restriction endonuclease before DSB labeling (**Fig. 1b,c**). Next,
the frequency of induced spike-in DSBs, B_{cut}, is calculated from enzyme cutting efficiency,
f_{cut}, that is calculated from gDNA sequencing data based on numbers of cut and uncut DNA
fragments covering cutting sites in gDNA (**Fig. 2a, Methods**), or qPCR data
(**Supplementary Fig. 1, Methods**).

Finally, the absolute frequency of studied DSBs, $B_{studied}$, is estimated from DSB sequencing
data:

$$B_{studied} = \frac{R_{studied}}{\alpha} \text{ , where } \alpha = \frac{R_{cut}}{B_{cut}} \tag{1}$$

and $R_{studied}$ and R_{cut} are the numbers of labeled reads originating from studied DSBs and
from enzyme cutting sites (spike-ins), respectively, and $B_{cut} \sim f_{cut}$.

Reproducibility and accuracy of cutting efficiency estimation. The number of labeled
reads per DSB (coefficient α) which is used for the final DSB quantification, as explained
above, is computed from enzyme cutting efficiency, f_{cut} (**Equation (1), Methods**).
Therefore, to calculate α accurately, we need to be able to estimate enzyme cutting
efficiency accurately. Commonly qPCR is used for precise measurement of cutting
efficiencies, however, this technique is inconvenient to use for multiple cutting sites. Thus,
we propose to use gDNA sequencing to determine spike-in cutting efficiencies (**Fig. 2a,
Methods**). To verify the accuracy and reproducibility of our proposed approach, we treated
immobilized and deproteinized yeast DNA with NotI enzyme and compared cutting
efficiencies at its recognition sites calculated using gDNA sequencing data and qPCR. The
cutting efficiencies for the selected NotI cutting site were highly consistent: 61% for gDNA
sequencing and 62% for qPCR. To examine if our approach can also be applied to breaks
introduced *in vivo*, which can be subjected to repair and resection, we used a yeast strain
engineered to produce a single site-specific DSB by I-SceI endonuclease *in vivo*. Cutting
efficiencies calculated based on gDNA sequencing and based on qPCR (**Supplementary
Fig. 1, Methods**) were again very consistent: 71% and 73%, respectively (**Fig. 2b**). We
therefore conclude that our method of estimating enzymes cutting efficiency based on
gDNA sequencing yields accurate and precise results.

Dependence of quantification on enzyme choice and types of breaks induced. DSBs
occurring *in vivo* are subject to DNA damage repair and therefore might be labeled with
different efficiencies than breaks induced *in vitro*. Moreover, different types of double-
stranded DNA ends (blunt or sticky) could also be detected more or less efficiently by a
given DSB labeling method. We therefore asked whether any restriction enzyme and any
manner of digestion can be applied to create spike-in DSBs that would lead to accurate
quantification. First, to test if restriction enzyme choice or the types of double-stranded
DNA ends influences our quantification results, we determined the spontaneous DSB
frequencies in yeast G_1 phase cells using NotI or SrfI spike-ins, which create sticky and
blunt ends, respectively. The number of spontaneous breaks in G_1 phase cells estimated
using these enzymes was consistent: 0.9 ± 0.3 DSBs per cell for NotI spike-in and $1.0 \pm$
0.6 DSBs per cell for SrfI spike-in (**Fig. 2c**). Then, to test if the results are affected by the
manner of digestion, we compared DSB estimations based on quantification using NotI (5'
overhangs) *in vitro* digestion and I-SceI (3' overhangs) *in vivo* digestion in HU-treated
wild-type cells (described below). Again, results were highly similar: 137 ± 12 and 153

± 52 DSBs per cell (**Fig. 2d**). In conclusion, qDSB-Seq provided consistent results in all
tested cases irrespective of the restriction enzyme used, types of DNA ends created by that
enzyme, or the manner of digestion.

Dependence of accurate quantification on adequate cutting efficiency. For accurate
quantification of studied DSBs, it is necessary that the relationship between the number of
labeled reads and DSB frequencies at different genomic locations is linear (**Equation (1)**,
Fig. 2a). This relationship could be affected by the frequencies of spike-in DSBs, which is
determined by an enzyme cutting efficiency. Therefore, we asked whether any frequency
of induced spike-in DSBs (i.e. any enzyme cutting efficiency) can be employed. To test
the influence of enzyme cutting efficiency on the quantification results, we performed 35
digestions for 25 samples using enzymes with multiple cutting sites (NotI, SrfI, AsiSI, and
BamHI) and then tested the linear relationship between the labeled reads and cutting
efficiencies for each digestion using Pearson Correlation Coefficient. We observed that
strong correlation ($R > 0.5$) (e.g. **Fig. 2e**) was always achieved for cutting efficiencies
between 12% and 62% (**Supplementary Fig. 2**, **Supplementary Table 2**) and for some
lower cutting efficiencies (4-12%). However, for the extreme cutting efficiencies (higher
than 84% or lower than 4%) the correlation was always weak (**Supplementary Fig. 3**). In
such cases, the number of observed cut or uncut fragments was low, making our estimates
less accurate, which likely decreased the correlation. Moreover, small variations in f_{cut}
between sites contributed to the decreased correlation (**Supplementary Fig. 3**).
Additionally, in samples for which digestion efficiencies are very high, the elevated level
of reads at spike-in sites ($> 75\%$) (**Supplementary Table 1**) can potentially disrupt (due
to low initial sequence diversity) Illumina sequencing[16]. Taken together, we conclude that
adequate cutting frequencies (4% to 84%) lead to a constant ratio between the labeled reads
and the cutting efficiencies for accurate quantification.

Stability of estimation of DSB frequencies per cell. We next asked whether our method
generates reproducible results. To test this, we calculated DSB frequencies in untreated G_1
cells based on different spike-ins. In spite of the various enzymes used (NotI, SrfI) we
obtained a very consistent number of DSBs (**Fig. 2f, Supplementary Fig. 4**,
Supplementary Table 1). Based on our calculations the frequency of spontaneous DSBs
in untreated G_1 wild-type cells is 1.0 ± 0.4 DSBs per cell, both the average and the range
(0.6-1.7 DSBs per cell) are consistent with previous studies[17, 18] (**Supplementary Table
1**). Further, we quantified DSBs based on the individual cutting sites in each of these
samples. The variation of the DSB quantification results depending on the individual
cutting sites used was lower than the average value (**Supplementary Table 1**). Similarly,
in *pif1* mutants, where stability of some DNA secondary structures is affected and we
observed increase in DSB numbers related to G-quadruplex[15] structures, we obtained
average DSB number 2.1 DSBs per cell. DSB quantification was consistent between the
samples (s.d. 0.3 DSBs per cell) (**Supplementary Fig. 4**, **Supplementary Table 1**).

Applications of qDSB-Seq

Quantification of DSBs induced by a radiomimetic drug, Zeocin. Some DSB-inducing
agents affect only particular sequences and structures, while others cause DNA damage
throughout the genome, e.g. irradiation. As DSB sequencing data inform only about read
distribution in the genome and is primarily used to identify regions enriched in reads, even
very large but global DSB induction will be undetectable using typical normalization

methods, e.g. normalization to the background. Therefore, to test application of qDSB-Seq
to such a challenging case, we used the radiomimetic agent Zeocin[19], a member of the
bleomycin drug family. After performing DSB sequencing, no apparent difference in raw
read counts between Zeocin-treated (ZEO) and untreated G_1 phase (G_1) cells was observed
(**Fig. 3a, Supplementary Fig. 5**). In contrast, after quantification (using qDSB-Seq with
NotI spike-in) we concluded that 1.1 ± 0.3 DSBs/cell were present in the G_1 sample and
7.4 ± 1.7 in ZEO, indicating that Zeocin induced 6.3 ± 2.0 DSBs per cell. Strikingly, Zeocin
significantly increased the number of DSBs (1.7- to 13-fold) in 99.8% of 5 kb genomic
intervals (p-value < 2e-12, hypergeometric test, **Methods**).

Interestingly, we observed that Zeocin-induced DSBs are especially enriched (3.0-fold) in
nucleosome-depleted regions (NDR) and reduced (0.4-fold) in nucleosome-protected
regions (both $p < 10^{-3}$, permutation test, **Methods**). Specifically, DSBs in the Zeocin-
treated sample occur 1.8 times as often between predicted nucleosome positions[20] as within
nucleosomes (**Fig. 3b**). Moreover, the preference for DSB location between nucleosomes
is even higher (4.1-fold) for long (> 100 nt) NDR regions (**Fig. 3c,d**). However, we do not
observe a 10 bp periodicity corresponding to the rotational positioning of the DNA helix
on the nucleosome. These results are consistent with previous findings that Zeocin-induced
cleavage is most suppressed in nucleosome-bound DNA and that this suppression is not
dependent on inaccessibility of the minor groove, but is caused by inability of the
nucleosome-bound DNA to undergo a conformational change that is required for Zeocin
binding[21]. Zeocin-induced DSBs are also enriched in DNA regions capable of forming very
stable DNA secondary structures (**Fig. 3e**), including G-quadruplexes (G4s)[22]. Further
studies will be necessary to elucidate this phenomenon. Nevertheless, increased DNA
damage on G4 structures could be related to nucleosome remodeling on G4s[23], consistent
with our finding that Zeocin prefers to cleave nucleosome-free DNA.

Quantification of DSBs induced by replication stress. We next used qDSB-Seq to
quantify replication-associated DSBs under hydroxyurea-induced replication stress (**Fig.
4a**). Hydroxyurea (HU) inhibits ribonucleotide reductase, resulting in decreased dNTP
levels and subsequent replication fork stalling and the slowing down of S phase[24]. Without
the protection of replication checkpoints, stalled forks may undergo catastrophic collapse
at high concentration or prolonged HU treatment[25], such as we used.

Using NotI spike-in, we observed that one hour treatment with 200 mM HU induced on
average 137.6 ± 12.0 DSBs per cell in wild-type yeast cells (WT +HU sample), which
represents a 9-fold increase relative to untreated S phase cells (15.4 ± 3.2 DSBs per cell).
The detected breaks showed a clear replication-related pattern: a significant enrichment of
DSB signal around replication origins (**Fig. 4b,c**). To further analyze the HU-induced
DSBs we classified them into two-ended DSBs and one-ended DSBs (**Supplementary Fig.
6**). Two-ended DSBs arise when two strands of DNA double helix are damaged (by i.e.
endonucleases, radiation or chemical compounds), while broken replication forks result in
one-ended DSBs. We identified one-ended DSBs using our method based on comparing
the number of reads between Watson and Crick strands (**Supplementary Fig. 6**, **Methods**)
and discovered that among all DSBs detected in HU-treated WT cells 71.7 ± 6.2 DSBs
were one-ended (**Fig. 4d**). Of those, 85% (60.6 ± 5.2 DSBs) were located within +/-10 kb
regions of active origins, resulting in an average of 0.4 one-ended DSB (broken fork) per
origin (**Fig. 4d**). Such one-ended DSBs would not be detected by some other DSB detecting

methods, such as pulse-field gel electrophoresis, which explains some earlier reports that
wild-type yeast cells are not sensitive to HU[25]. The observed one-ended DSBs might
correspond to broken forks resulting from transient DNA breaks occurring on the leading
strand, as reported by Sasaki *et al*[26]. In agreement with this theory, we discovered that two
hours after removal of HU, the number of one-ended DSBs decreased dramatically (by
86%) (**Fig. 4d**), indicating that replication-associated DNA damage present during HU
treatment is not permanent.

Quantification of DSBs at ribosomal replication fork barriers. Replication fork barriers
(RFBs) are natural barrier that blocks replication forks to protect nearby, highly expressed
rRNA genes from collisions between transcription and replication complexes[26, 27] (**Fig. 5a**).
DSBs occurring at the ribosomal replication fork barriers (RFBs) have been observed using
Southern blot in the budding yeast[28-31]. However, precise frequencies and genomic
locations of these DSBs were not established due to lack of a quantitative and sensitive
DSB detection method[26]. Using qDSB-Seq, here we both precisely quantified DSB
frequencies near RFBs and identified their genomic coordinates.

It was reported that Fob1 proteins bound to an RFB site block replication fork progression,
resulting in generation of a one-ended DSBs[30]. Indeed, in unperturbed S-phase cells, we
observed 1.1 DSBs per cell (0.0055 DSBs per rDNA repeat) on rDSB-1 and rDSB-2 sites
upstream of RFB1 and RFB2 (two closely spaced RFB loci) (**Fig. 5b,c and**
Supplementary Table 3). As expected, we did not detect any DSBs at these sites in G_1-
arrested cells confirming that the observed DSBs at RFBs are replication-dependent.

It was previously shown that Top1 in the presence of Fob1 specifically cleaves defined
sequences in the RFB region[32]. When we inhibited the religation step of Top1 by adding
100 μM camptothecin (CPT) for 45 min treatment, we observed a CPT-dependent DSB
site (rDSB-3), exactly at the same location as the previously identified Top1-dependent
cleavage site (**Fig. 5c**). In addition, this site also colocalizes with a Fob1 binding region, in
agreement with a previous discovery that the recruitment and stabilization of Top1 requires
the binding of Fob1 protein[32]. Our quantification shows the DSB frequency at rDSB-3 site
was 0.1 DSB per cell, lower than at rDSB-1 and rDSB-2. Finally, our results agree with
previous work[26] in which approximately one DSB arises in an rDNA array during
replication in a yeast cell (**Fig. 5b**); such low frequencies are caused by recombination in
the rDNA array[26]. Based on the results above, qDSB-Seq fills the need to enable detection
of these rare breaks at replication fork barriers and allowed us for the first time to quantify
the frequency of cleavage of Topoisomerase 1 (Top1) at RFBs.

Discussion

We propose qDSB-Seq, a general framework that allows estimating both absolute DSB
frequencies (per cell) and their precise genomic coordinates. qDSB-Seq combines a DSB-
labeling method with a quantification technique; quantification is achieved by inducing
easy-to-measure spike-in DSBs via restriction enzyme digestion.

Due to increasing evidence of a relationship between emergence of DSBs and human
diseases such as cancer[1], there is growing interest in precise detection of DSBs. Several
general genome-wide methods for detection of DSBs with single-nucleotide resolution
have recently been developed[3-6], however their usefulness is limited because they only
allow comparison of DSB levels between genomic loci within the same sample.

Normalization to the total number of reads is often employed to enable comparison
between different samples, but this method is not always applicable. For example, it cannot
be used if DSBs are induced throughout the whole genome or if the DSB background varies,
which is common[33]. Therefore, in case of agents that create such DSB patterns, e.g. by
irradiation or radiomimetic drugs, data normalized to the total reads number will not reveal
global induction of breaks as shown in **Fig. 3a**. In contrast, our approach allows not only
estimation of relative increases of DSB signal between samples (regardless of signal
distribution), but also quantification of absolute DSB numbers per cell. For example, we
discovered that 1 hour treatment with 100 µg/ml Zeocin results in 6.7-times increase in
DSBs, namely from 1.1 ± 0.3 to 7.4 ± 1.7 DSBs per cell. Additionally, we discovered that
Zeocin significantly increases DSB levels in 99.8% of 5kb genomic intervals, but with
differences in ratios: from 1.7- to 13-fold. qDSB-Seq opens up new possibilities in studying
the impact of DSB inductors or gene mutations on genome instability, i.e. it may potentially
allow determining the outcomes of different doses of anticancer drugs in healthy and tumor
cells. Moreover, qDSB-Seq allows assessing DSB frequencies not only for the whole
genome, but also for a specific locus. For instance, using our approach, for the first time
we quantified changes of DSB frequency at RFBs between wild-type and CPT-treated cells,
thus revealing the frequency of Top1-dependent DSBs in RFB region.

Key innovation of qDSB-Seq is spike-in DSBs used for normalization. Such spike-in DSBs
can be introduced both *in vivo* and *in vitro*; each manner of digestion has its strengths and
weaknesses. *In vivo* digestion requires organism-specific constructs, such as the I-SceI
yeast strain we used, while *in vitro* digestion can be applied to any organism. Moreover,
for *in vitro* digestion, since spike-in DSBs are never repaired and thus there are no resected
DNA ends. Resected DNA ends may result in spike-in related reads located up to several
kilobases from the cutting sites, which may complicate data analysis. On the other hand,
for *in vivo* digestion it is possible to determine enzyme cutting frequency before addition
of spike-in cells to the sample of interest, which facilitates obtaining final cutting efficiency
in the desired range by selecting desired mixing proportions. *In vivo* digestion can be also
used to study the DNA damage response in systems such as DivA[34].

Enzyme cutting efficiency is a key parameter influencing qDSB-Seq accuracy. As shown
above, using extremely low or high cutting efficiencies may result in inaccurate
quantification results, while within an adequate range (4% to 84%), the number of labeled
reads per DSB (proportionality coefficient α) remains constant, which allows for
consistently accurate quantification. If spike-in DSBs are introduced *in vivo*, to achieve
desired cutting efficiency one needs to mix in appropriate proportions cells in which full
digestion (or digestion with known efficiency) was performed with the studied cells. In
case of *in vitro* digestion, the studied cells should be treated with a dose of an enzyme much
lower than recommended for full digestion. The enzyme cutting efficiency can be then
estimated by performing qPCR and, if needed, the dose can be adjusted before sequencing.

To facilitate choice of a restriction enzyme for qDSB-Seq experiments we provide lists of
restriction enzymes sorted according to their cutting efficiencies per Mb in the yeast,
human, mouse and fruit fly genomes (**Supplementary Table 4**), as well as Genome-wide
Restriction Enzyme Digestion STatistical Analysis Tool, GREDSTAT, at
http://bioputer.mimuw.edu.pl:23456. Enzymes with multiple cutting sites should yield best
quantification results, since estimation of the enzyme cutting frequency will be less

influenced by a potential local bias. Constructs with a single enzyme cutting site, such as
the I-SceI strain we employed, allow convenience of using qPCR to determine an enzyme
cutting frequency. Therefore, for enzymes with multiple cutting sites, we developed a
method to estimate enzyme cutting efficiency from gDNA sequencing data, and proved its
accuracy by comparing with qPCR results. On the other hand, usage of rare cutting
enzymes is preferable, since they allow for optimal cutting efficiencies at individual sites
without unnecessarily increasing percentage of spike-ins in total reads. There is no benefit
to using a higher spike-in percentage than necessary; high spike-in percentages, especially
exceeding 30-50% of total reads, may cause quality issues with Illumina sequencing[16].
Unlike enzyme cutting efficiency, percentage of spike-in reads cannot be determined
before sequencing, since it depends both on enzyme cutting efficiency and number of DSBs
present in the data. Therefore, if there is a probability that high level of spike-ins may be
achieved unintentionally (e.g. during pilot experiments), we recommend using our
modified protocols for generation of high-quality sequencing data from low-diversity
samples[16].

qDSB-Seq is compatible with any DSB labeling technique, but will also share limitations
of the used method. For example, we tested that the type of generated DNA ends will not
determine quantification results when using i-BLESS for DSB labeling. However, as we
discussed in[15], some DSB sequencing technologies cannot detect all types of DNA ends.
Therefore, qDSB-Seq, when used in combination with such technology, will also exhibit
bias in quantifying DSBs with these types of DNA ends.

When interpreting qDSB-Seq results, it is important to keep in mind that qDSB-Seq relies
on sequencing data derived from a population of cells. Therefore, it only yields an average
number of DSBs per cell, which may or may not be representative of a typical single cell.
This problem can be solved by combining qDSB-Seq with a complementary method,
giving insight into population-distribution of DSBs, as we proposed elsewhere[33].

In summary, qDSB-Seq is a novel approach, which allows absolute DSB quantification
genome-wide and accurate cross-sample comparison and can be applied to any organism,
for which a DSB labeling method is available. qDSB-Seq relies on a key innovation, using
spike-in DSBs induced by a restriction enzyme for normalization. Using qDSB-Seq, we
quantified the numbers of DSBs induced by a radiomimetic drug and replication stress;
measured for the first time Top1-dependent DSB frequencies at replication fork barriers
and revealed several orders of magnitude differences in DSB frequencies. Such high
variability in genome breakage highlights the importance of quantification and shows how
challenging data interpretation would be without the normalization provided by qDSB-Seq.

Acknowledgements

This research was supported by the NIH grant R01GM112131 to M.R. (Y.Z., N.D., B.F.,
J.N., R.Y. and M.R.), Polish National Science Centre grant to M.S.
(2015/17/D/NZ2/03711), and Foundation for Polish Science grant TEAM/2016-2 to K.G.
This work was also supported by Ligue contre le Cancer (Equipe labelisee), Agence
Nationale pour la Recherche (ANR) and Institut National du Cancer (INCa) grants to P.P.
(B.P., R.F. and P.P.), National Science Center grant 2016/21/B/ST6/01471 to N.D., and a
training fellowship from the Gulf Coast Consortia on the Computational Cancer Biology

Training Program (CPRIT Grant No. RP170593) to Y.Z. The authors are grateful to
Heather Lander of the Sealy Center for Structural Biology and Molecular Biophysics at
UTMB, for editorial services for the manuscript.
357

Author contributions

Competing Financial interests

The authors declare no competing financial interests.

References

1. Khanna, K.K. & Jackson, S.P. DNA double-strand breaks: signaling, repair and
the cancer connection. *Nature genetics* **27**, 247-254 (2001).
2. Slaymaker, I.M. et al. Rationally engineered Cas9 nucleases with improved
specificity. *Science* **351**, 84-88 (2016).
3. Crosetto, N. et al. Nucleotide-resolution DNA double-strand break mapping
by next-generation sequencing. *Nat Methods* **10**, 361-365 (2013).
4. Hoffman, E.A., McCulley, A., Haarer, B., Arnak, R. & Feng, W. Break-seq reveals
hydroxyurea-induced chromosome fragility as a result of unscheduled
conflict between DNA replication and transcription. *Genome Res* **25**, 402-412
(2015).
5. Lensing, S.V. et al. DSBCapture: in situ capture and sequencing of DNA breaks.
Nature methods **13**, 855-857 (2016).
6. Canela, A. et al. DNA Breaks and End Resection Measured Genome-wide by
End Sequencing. *Molecular cell* **63**, 898-911 (2016).
7. Yan, W.X. et al. BLISS is a versatile and quantitative method for genome-wide
profiling of DNA double-strand breaks. *Nature communications* **8**, 15058
(2017).
8. Ran, F.A. et al. In vivo genome editing using Staphylococcus aureus Cas9.
Nature **520**, 186-191 (2015).
9. Aymard, F. et al. Genome-wide mapping of long-range contacts unveils
clustering of DNA double-strand breaks at damaged active genes. *Nature*
structural & molecular biology **24**, 353-361 (2017).

10.   Popp, H.D., Brendel, S., Hofmann, W.K. & Fabarius, A. Immunofluorescence
Microscopy of gammaH2AX and 53BP1 for Analyzing the Formation and
Repair of DNA Double-strand Breaks. *J Vis Exp* (2017).

11.   Chailleux, C. et al. Quantifying DNA double-strand breaks induced by site-
specific endonucleases in living cells by ligation-mediated purification. *Nat*
Protoc **9**, 517-528 (2014).

12.   Gregoire, M.C. et al. Quantification and genome-wide mapping of DNA
double-strand breaks. *DNA repair* **48**, 63-68 (2016).

13.   Clement, K., Farouni, R., Bauer, D.E. & Pinello, L. AmpUMI: design and analysis
of unique molecular identifiers for deep amplicon sequencing. *Bioinformatics*
34, i202-i210 (2018).

14.   Smith, T., Heger, A. & Sudbery, I. UMI-tools: modeling sequencing errors in
Unique Molecular Identifiers to improve quantification accuracy. *Genome Res*
27, 491-499 (2017).

15.   Biernacka, A. et al. i-BLESS is an ultra-sensitive method for detection of DNA
double-strand breaks. *Commun Biol* **1**, 181 (2018).

16.   Mitra, A., Skrzypczak, M., Ginalski, K. & Rowicka, M. Strategies for achieving
high sequencing accuracy for low diversity samples and avoiding sample
bleeding using illumina platform. *PLoS One* **10**, e0120520 (2015).

17.   Lobrich, M. et al. gammaH2AX foci analysis for monitoring DNA double-
strand break repair: strengths, limitations and optimization. *Cell Cycle* **9**, 662-
669 (2010).

18.   Thongsroy, J. et al. Replication-independent endogenous DNA double-strand
breaks in Saccharomyces cerevisiae model. *PLoS One* **8**, e72706 (2013).

19.   Shimada, K. et al. TORC2 signaling pathway guarantees genome stability in
the face of DNA strand breaks. *Molecular cell* **51**, 829-839 (2013).

20.   Lee, W. et al. A high-resolution atlas of nucleosome occupancy in yeast.
Nature genetics **39**, 1235-1244 (2007).

21.   Povirk, L.F. DNA damage and mutagenesis by radiomimetic DNA-cleaving
agents: bleomycin, neocarzinostatin and other enediynes. *Mutat Res* **355**, 71-
89 (1996).

22.   Bochman, M.L., Paeschke, K. & Zakian, V.A. DNA secondary structures:
stability and function of G-quadruplex structures. *Nature reviews. Genetics*
13, 770-780 (2012).

23.   Hershman, S.G. et al. Genomic distribution and functional analyses of
potential G-quadruplex-forming sequences in Saccharomyces cerevisiae.
Nucleic acids research **36**, 144-156 (2008).

24.   Koc, A., Wheeler, L.J., Mathews, C.K. & Merrill, G.F. Hydroxyurea arrests DNA
replication by a mechanism that preserves basal dNTP pools. *The Journal of*
biological chemistry **279**, 223-230 (2004).

25.   Singh, A. & Xu, Y.J. The Cell Killing Mechanisms of Hydroxyurea. *Genes (Basel)*
7 (2016).

26.   Sasaki, M. & Kobayashi, T. Ctf4 Prevents Genome Rearrangements by
Suppressing DNA Double-Strand Break Formation and Its End Resection at
Arrested Replication Forks. *Molecular cell* **66**, 533-545 e535 (2017).

27.   Kobayashi, T. The replication fork barrier site forms a unique structure with
439   Fob1p and inhibits the replication fork. *Molecular and cellular biology* **23**,
440   9178-9188 (2003).
441   28.   Kobayashi, T., Horiuchi, T., Tongaonkar, P., Vu, L. & Nomura, M. SIR2
442   regulates recombination between different rDNA repeats, but not
443   recombination within individual rRNA genes in yeast. *Cell* **117**, 441-453
444   (2004).
445   29.   Weitao, T., Budd, M. & Campbell, J.L. Evidence that yeast SGS1, DNA2, SRS2,
446   and FOB1 interact to maintain rDNA stability. *Mutat Res* **532**, 157-172
447   (2003).
448   30.   Burkhalter, M.D. & Sogo, J.M. rDNA enhancer affects replication initiation and
449   mitotic recombination: Fob1 mediates nucleolytic processing independently
450   of replication. *Molecular cell* **15**, 409-421 (2004).
451   31.   Weitao, T., Budd, M., Hoopes, L.L. & Campbell, J.L. Dna2 helicase/nuclease
452   causes replicative fork stalling and double-strand breaks in the ribosomal
453   DNA of Saccharomyces cerevisiae. *The Journal of biological chemistry* **278**,
454   22513-22522 (2003).
455   32.   Di Felice, F., Cioci, F. & Camilloni, G. FOB1 affects DNA topoisomerase I in vivo
456   cleavages in the enhancer region of the Saccharomyces cerevisiae ribosomal
457   DNA locus. *Nucleic acids research* **33**, 6327-6337 (2005).
458   33.   Zhu, Y., Biernacka, A., Pardo, B., Forey, R., Dojer, N., Yousefi, R., Nde, J.,
459   Fongang, B., Mitra, A., Li, J., Skrzypczak, M., Kudlicki, A., Pasero, P., Ginalski, K.,
460   Rowicka, M. Integrated analysis of patterns of DNA breaks reveals break
461   formation mechanisms and their population distribution during replication
462   stress. *BioRxiv* (2017).
463   34.   Caron, P. et al. Non-redundant Functions of ATM and DNA-PKcs in Response
464   to DNA Double-Strand Breaks. *Cell Rep* **13**, 1598-1609 (2015).
465   35.   Schmittgen, T.D. & Livak, K.J. Analyzing real-time PCR data by the
466   comparative C-T method. *Nat Protoc* **3**, 1101-1108 (2008).
467   36.   Langmead, B., Trapnell, C., Pop, M. & Salzberg, S.L. Ultrafast and memory-
468   efficient alignment of short DNA sequences to the human genome. *Genome
469   Biol* **10**, R25 (2009).
470   37.   Markham, N.R. & Zuker, M. DINAMelt web server for nucleic acid melting
471   prediction. *Nucleic acids research* **33**, W577-581 (2005).
472   38.   Kudlicki, A.S. G-Quadruplexes Involving Both Strands of Genomic DNA Are
473   Highly Abundant and Colocalize with Functional Sites in the Human Genome.
474   *PLoS One* **11**, e0146174 (2016).
475   39.   Yabuki, N., Terashima, H. & Kitada, K. Mapping of early firing origins on a
476   replication profile of budding yeast. *Genes to cells : devoted to molecular &
477   cellular mechanisms* **7**, 781-789 (2002).
478

ONLINE METHODS

Strains and growth conditions. Yeast strains used in this study are listed in

Supplementary Table 5. Cells were grown in YPD medium at 25°C until early log phase
and were then arrested in G_1 for 170 min with 8 μg/ml α-factor. For exposure to Zeocin
cells were treated with 100 μg/ml Zeocin (Invivogen) for 1 hour. The I-SceI strain was
cultured in YPR medium, galactose was added for 2 h to induce I-SceI cutting. For
exposure to hydroxyurea, cells were released from G_1 arrest by addition of 75 μg/ml
Pronase (Sigma) and 200 mM HU was added 20 min before Pronase release followed by 1
h incubation. Collected cells were washed with cold SE buffer (5M NaCl, 500 mM EDTA,
pH 7.5) and immediately subjected to DSB labeling.

DSB sequencing. DSB labeling was performed using our i-BLESS method as described
in[15]. Zeocin treated cells were additionally subjected to reaction with NEBNext® FFPE
DNA Repair Mix prior to proximal adapter ligation. Sequencing libraries for i-BLESS and
respective gDNA samples were prepared using ThruPLEX DNA-seq Kit (Rubicon
Genomics). i-BLESS libraries were prepared without prior fragmentation and further size
selection. Quality and quantity of the libraries were assessed on a 2100 Bioanalyzer using
HS DNA Kit, and on a Qubit 2.0 Fluorometer using Qubit dsDNA HS Assay Kit (Life
Technologies). The libraries were sequenced (2x70bp) on Illumina HiSeq2500/HiSeq4000
platforms, according to our modified experimental and software protocols for generation
of high-quality data from low-diversity samples[16].

qDSB-Seq with NotI, SrfI, AsiSI, and BamHI digestion. In addition to DSB sequencing,
as described above, a digestion with a restriction enzyme was performed before DSB
labeling. Samples were treated with NotI (NEB, Thermo Scientific), SrfI (NEB), AsiSI
(NEB), or BamHI (Thermo Scientific) for 1 h at 37ºC. The dose and incubation time of
these restriction enzymes were listed in **Supplementary Table 6**.

qDSB-Seq with I-SceI spike-in. For I-SceI spike-in we used a yeast strain (I-SceI strain)
with GAL inducible I-SceI endonuclease and a single I-SceI cutting site integrated at the
ADH4 locus on chromosome VII. To measure the cleavage efficiency of I-SceI, cell
aliquots were taken pre- (RAFF) and 2 h post- (GAL) cleavage induction, and total
genomic DNA was extracted. DNA was serially diluted and amplified for 25 cycles with
primers spanning the I-SceI cutting site. Cleavage efficiency was inferred by comparing
the amount of amplified DNA in GAL (cut) vs. RAFF (uncut) conditions. We used CASY
Cell Counter (Roche Applied Science) to mix this spike-in with our sample of interest
(wild-type cells with replication stress induced by hydroxyurea treatment) in proportion
2:98. The cutting ratio of the I-SceI endonuclease expressed in the I-SceI strain was
estimated using an unmixed I-SceI strain and **Equation (1)** below.

Quantitative PCR. To validate cutting efficiency for NotI, input gDNA was analyzed by
real-time PCR using primers flanking a selected NotI site at chrI: 114016-114023 (forward:
AGAGTTGGGAATGTGTGCCC, reverse: GGGCAGCAACACAAAGTGTC) and
KAPA SYBR® FAST kit (Life Technologies). Four technical replicates using two
different concentrations of input DNA were performed. We compared the amount of PCR
product amplified in untreated (C) vs. NotI treated cells (T) by data analysis based on the
ΔC_T method[35], where the ΔC_T value was obtained by subtraction of the C_T value in sample
C from the C_T value in sample T. Final cutting efficiency was calculated as mean efficiency
for all dilutions according to the formula below:

527
$$f_{cut} = 1 - \frac{1}{2^{\Delta C_T}}$$

We used calibration data to empirically correct ΔC_T.

Sequencing data analysis. We used *iSeq* (http://breakome.eu/software.html) to ensure
sequencing data quality before mapping. Next, *iSeq* was used to remove i-BLESS proximal
and distal barcodes (TCGAGGTAGTA and TCGAGACGACG, respectively). Reads
labeled with the proximal barcode, which are directly adjacent to DSBs, were selected and
mapped to the version of the yeast S288C genome sacCer3 (we manually corrected
common polymorphisms) using bowtie[36] v0.12.2 with the alignment parameters '-m1 –v1'
(to exclude ambiguous mapping and low-quality reads). For ribosomal DNA mapping in
replication fork barrier analysis, we mapped sequencing reads using the parameter '-v1' to
allow multiple mapped reads. The end base pairs of the reads were trimmed using bowtie
'-3' parameter. The parameter choice was based on the *iSeq* quality report. For calculation
of the absolute number of DSBs per cell only mapped reads were retained. Further, the
reads identified as originating from telomere ends were removed. The telomeric reads were
identified as those exhibiting the CAC motif in the whole AC-rich strand; regular
expression C{0,3}AC{1,10} in the PERL language was used to identify them.

Calculation of DSB frequencies per cell. Paired-end sequencing of gDNA or qPCR was
used to measure the cutting efficiency of the endonuclease. For an enzyme with a single
cutting site (e.g. I-SceI), we used the following procedure to calculate cutting efficiency
(f_{cut}) from whole genome paired-end sequencing data:

548
$$f_{cut} = \frac{N_{cut}}{N_{cut} + 2N_{uncut}} - f_{bg} \tag{1}$$

where, N_{cut} is the number of fragments cut by an enzyme, N_{uncut} is the number of uncut
fragments covering the cutting site, and f_{bg} is the background level of breaks (e.g. resulting
from sonication). N_{cut} fragments were counted in empirically determined, several
nucleotide vicinities of the canonical cutting sites, based on visual examination of the read
distribution. For enzyme with multiple cutting sites, reads mapped to each cutting site were
first classified as "cut" or "uncut" and the results were summed over all cutting sites:

555
$$f_{cut} = \frac{\sum_{i=1}^{N_{sites}} N_{cut}^i}{\sum_{i=1}^{N_{sites}} N_{cut}^i + 2 \sum_{i=1}^{N_{sites}} N_{uncut}^i} - f_{bg}$$

To estimate cutting efficiency, we used only cutting sites to which > 100 paired-end reads
were mapped and their cutting efficiency was larger than 0. To estimate f_{bg}, we randomly
selected genomic windows of the same size as those used to count cut and uncut fragments
and estimated "cutting efficiency" in those intervals using the left part of **Equation (1)**. For
clarity, these errors are omitted in **Equations (2)** to (**4**).

Next, we calculated the number of spike-in DSBs induced at restriction sites, B_{cut}:

562
$$B_{cut} = f_{cut} N_{sites} p \tag{2}$$

563   where f_{cut} is the cutting efficiency in undiluted samples, N_{sites} is the number of used enzyme
564   restriction sites (e.g. 39 for NotI) and p is the proportion of digested cells ($p = 1$ unless
565   mixing with an *in vivo* digested construct is used).

566   Then we computed the number of mapped sequencing reads per DSB or the coefficient, α:

$$567 \qquad\qquad\qquad\qquad \alpha = \frac{R_{cut}}{B_{cut}} \qquad\qquad\qquad\qquad (3)$$

568   where R_{cut} is the number of labeled reads mapped to the cutting sites and B_{cut} is the total
569   number of induced DSBs.

570   Finally, we computed studied DSBs per cell ($B_{studied}$) using the following formula:

$$571 \qquad\qquad\qquad\qquad B_{studied} = \frac{R_{studied}}{\alpha} \qquad\qquad\qquad\qquad (4)$$

where $B_{studied}$ is the number of studied DSBs per cell in the whole genome, or in a specific
region (eg. a replication region), or at a specific location (eg. an enzyme cutting site). In
this study, we calculated the studied breaks per cell for the whole genome after subtracting
reads generated from enzyme cutting sites, telomeres, and ribosomal DNA. Errors for
$B_{studied}$ are the standard deviation of breaks calculated from different cutting sites for
enzymes with multiple cutting sites (**Supplementary Table 1**). Based on replicates, we
concluded that thus calculated errors are conservatively estimated. For an enzyme with a
single cutting site in a given genome, errors for B_{cell} were assigned using computed errors
of the cutting efficiencies from f_{bg}.
581

Background estimation and removal. To quantify DSBs likely resulting from broken
forks near origins, we first removed background not related to replication. To define such
background, we calculated DSB density in a 500 bp sliding window with a 50 bp step; the
peak of this distribution was assumed to be background DSB frequency. This background
was subtracted from the data at each position, resulting negative values were assigned to
zero.
588

Analysis of fragile regions and enrichment. Hygestat_BLESS v1.2.3 in the *iSeq* package
(http://breakome.eu/software.html) was used to identify fragile regions (i.e. regions with
significant increase of the read numbers in treatment versus control samples), which were
defined using the hypergeometric probability distribution and Benjamini-Hochberg
correction. To evaluate the enrichment of fragile regions on nucleosomes, we used
hygestat_annotations v2.0, which computed the proportion of mappable nucleotides
belonging to both the fragile regions and the nucleosomes, and the proportion of mappable
nucleotides belonging to both genomic regions and the nucleosomes. To estimate the p-
value for the feature enrichment inside fragile regions, we used 1000 permutations to
calculate the empirical distribution of the ratio under the null hypothesis.
599

Estimation of one-ended DSBs. To estimate the total number of one-ended DSBs, we
performed hypergeometric test based on the number of i-BLESS sequencing reads from
Watson and Crick strands using Hygestat_BLESS v1.2.3 in the *iSeq* package with a 500 nt
window size. Regions with $P < 1e\text{-}10$ were classified as one-ended DSB regions, *P* value
was corrected by the Bonferroni correction. The subtraction between reads from Watson
and Crick was treated as the number of one-ended reads used to calculate one-ended DSBs
using the DSB calculation method.
607

Comparison of DSB levels between ZEO and G$_1$ samples. We used read counts for 5000
nt mappable intervals produced by hygestat_BLESS; ZEO read numbers were normalized

using qDSB-Seq quantification. We evaluated the null hypothesis that the number of DSBs
in G_1 cells is the same or lower than in ZEO using very conservative 5 standard deviation
confidence intervals (assuming Poisson distribution of reads). All genomic windows
with >17 reads in 5 kb were significantly enriched in DSBs in ZEO as compared with G_1
cells (P < 2e-12, calculated using the hypergeometric probability distribution and the
Bonferroni correction).
616
DNA secondary structure and G-quadruplex prediction. DNA secondary structures
were defined by free energy at 37°C using UNAFold[37] v3.8 in a 50 bp sliding window with
a 25 bp step along the whole yeast genome. We predicted G-quadruplexes (both canonical
intrastrand and non-canonical inter-strand) in the budding yeast genome using AllQuads[38]
software, with the standard 7-nt threshold on loop length.

Statistical analysis. Results of quantification are shown as mean ± s.d. To conduct
enrichment analysis, the P values were first calculated using the hypergeometric
distribution function as implemented in the GNU Scientific Library for C++ and then
corrected for multiple hypothesis testing using the Benjamini-Hochberg method. The
threshold for statistical significance was $P < 0.05$.

Code availability. Custom code used in this study is available upon request from authors
or http://breakome.eu/software.html.
629
630   **Data availability**. The DSB sequencing data will be available upon publication at
631   Sequence Read Archive.

Figures

Figure 1. qDSB-Seq method. (a) In qDSB-Seq protocol after DSBs induction, cells are treated with a restriction enzyme to introduce site-specific, infrequent DSBs (spike-ins). Next, DSBs are labeled (using e.g. i-BLESS) and sequenced. Simultaneously, gDNA sequencing (or qPCR) is performed and used to estimate the cutting efficiency of the enzyme, and thus frequency of induced DSB spike-ins, which is then used to quantify the absolute DSB frequency (per cell) of studied DSBs in the sample (**Methods**). **(b-c)** Spike-in DSBs were induced in two different ways: **(b)** the studied cells were digested using the NotI, SrfI, AsiSI, or BamHI restriction enzyme *in vitro*; **(c)** cells expressing the restriction enzyme I-SceI *in vivo* (the I-SceI strain) were mixed with the studied cells.

Figure 2. qDSB-Seq computation and validation. (a) Computation of the labeled reads per DSB and DSBs per cell. The ratio of the labeled reads and cutting efficiency at enzyme cutting sites was calculated and then used for DSB quantification in the studied genomic loci. **(b)** The estimation of I-SceI cutting efficiency based on gDNA sequencing and qPCR (error was calculated from technical replicates). **(c-d)** Dependence of qDSB-Seq quantification on the restriction enzyme used. DSB levels obtained for **(c)** untreated WT G_1 phase cells, and for **(d)** HU-treated WT S phase cells quantified using NotI and SrfI digestion *in vitro* and I-SceI digestion *in vivo* (errors of the estimated DSB frequencies were calculated as described in **Methods**). **(e)** Correlation between the number of labeled reads at cutting sites and their cutting efficiencies in an untreated G_1 phase sample, digested with NotI enzyme with average cutting efficiency of 18%. *R*: Pearson correlation coefficient. **(f)** Quantification of DSBs in untreated G_1 phase cells. The dashed lines and the stripes are the mean value and 95% confidence interval, respectively. Mean ± s.d. is shown for each sample.

Figure 3. Quantification of Zeocin-induced DSBs. **(a)** Raw read density and quantified DSB density (500 bp sliding window with 50 bp step) in a representative fragment of chromosome XII for untreated and Zeocin-treated wild-type G_1 phase cells. Raw read density was normalized to the total number of reads. **(b-c)** Density of Zeocin-induced DSBs in **(b)** all and **(c)** ≥100 bp low nucleosome occupancy regions. Nucleosome locations from Lee *et al.*[20] were used, DSB densities, expressed as DSBs per million cells, were calculated in a 50 bp sliding window with a 5 bp step. **(d)** Comparison of DSB densities in high nucleosome occupancy regions (High) and low nucleosome occupancy regions (Low). **(e)** Enrichment of Zeocin-induced DSBs in regions prone to form very stable DNA secondary structures (e.g. hairpins), as defined by free energy in a 50 bp sliding window as described in **Methods**. Zeocin-induced DSBs were defined as regions with significant enrichment of DSB-labeled reads in ZEO sample compared with G_1 phase control, as identified using Hygestat_BLESS. Enrichment analysis was performed using hygestat_annotations (**Methods**). *P* values: * *P* < 0.05, ** *P* < 0.01, *** *P* < 0.001.

Figure 4. Quantification of replication-associated DSBs. (a) Schematic representation of HU experiments. Cells were arrested in G₁ phase with α-factor, treated with HU before release to S phase, harvested after 1 hour or resuspended in fresh medium and harvested 2 hours after removal of HU. **(b)** Example of quantified DSB data from HU-treated wild-type and 2-hour recovery cells. Replication origins are marked with green triangles, absolute frequencies of DSBs for a fragment of chromosome IV are shown in a million cells. As a control, replication profile (values of DNA copy number) in WT +HU sample is shown, for which the number of gDNA reads in a 500 bp window in WT +HU sample was normalized by G₁ sample. **(c)** Meta-profile of DSBs around active replication origins under HU treatment, defined as 144 origins with firing time < 25 min (early origins, firing time according to Yabuki *et al.*[39]}). Median of DSB densities, expressed as DSBs per million cells in 2 kb window around each early origin, was calculated, the background was removed as described in **Methods**. **(d)** Quantification of one-ended DSBs. Errors of the estimated one-ended DSB frequencies were calculated as described in **Methods**.

Figure 5. DNA double-strand breaks at replication fork barriers. (a) Scheme of Replication Fork Barriers (RFBs) at yeast rDNA locus. **(b)** The total number of RFB-related one-ended DSBs (peaks as defined in panel **c)** calculated from the difference of Watson and Crick strand reads (**Methods**); **(c)** Quantified DSBs signal in RFB region. RFB1 and RFB2 are indicated by the red boxes on the top. The green boxes mark Fob1 protein binding sites mapped *in vitro*. The yellow star indicates Top1 cleavage site. The red arrows point out the observed ribosomal DSB sites, rDSB-1, rDSB-2, and rDSB-3.