Dopaminergic drug effects on probability weighting during risky decision-making

Karita E. Ojala[1], Lieneke K. Janssen[1], Mahur M. Hashemi[1], Monique H. M. Timmer[1,2], Dirk E. M. Geurts[1,3], Niels P. ter Huurne[1,4], Roshan Cools[1,3], Guillaume Sescousse[1]

[1] Donders Institute for Brain, Cognition and Behaviour, Radboud University, Nijmegen, the Netherlands

[2] Department of Neurology, Radboud University Medical Centre, Nijmegen, the Netherlands

[3] Department of Psychiatry, Radboud University Medical Centre, Nijmegen, the Netherlands

[4] Karakter Child and Adolescent Psychiatry University Centre, Nijmegen, the Netherlands

Corresponding authors:

Karita E. Ojala
Division of Clinical Psychiatry Research
Psychiatric Hospital, University of Zurich
Lenggstrasse 31, 8032 Zurich
Switzerland
karita.ojala@uzh.ch

Guillaume Sescousse
Donders Centre for Cognitive Neuroimaging
PO Box 9101, 6500 HB Nijmegen
The Netherlands
g.sescousse@donders.ru.nl

Pages (22)
Figures (3), Tables (3)
Words: Abstract (224), Introduction (649), Discussion (1416)

Author contributions: R.C. and G.S. designed research, L.K.J., M.H., M.H.M.T., D.E.M.G., N.P.H. and G.S. performed research, K.E.O. AND G.S. analysed the data, and K.E.O, R.C. and G.S. wrote the paper.

Conflict of interest: The authors declare no competing financial interests.

Acknowledgements: GS was supported by a Veni grant from the Netherlands Organisation for Scientific Research (NWO). We would like to thank Romain Ligneul, Payam Piray and Bram Zandbelt for helpful discussions on data analysis and visualization.

Current author affiliations: Karita E. Ojala: Division of Clinical Psychiatry Research, Psychiatric University Hospital, University of Zurich, Zurich, Switzerland. Lieneke K. Janssen: Department of Neurology, Max Planck Institute for Human Cognitive and Brain Sciences; Leipzig University Medical Center, IFB Adiposity Diseases, Leipzig, Germany.

Abstract

Dopamine has been associated with risky decision-making, as well as with pathological gambling, a behavioural addiction characterized by excessive risk-taking behaviour. However, the specific mechanisms through which dopamine might act to foster risk-taking and pathological gambling remain elusive. Here we test the hypothesis that this might be achieved, in part, via modulation of subjective probability weighing during decision-making. Healthy controls ($n = 21$) and pathological gamblers ($n = 16$) played a decision-making task involving choices between sure monetary options and risky gambles both in the gain and loss domains. Each participant played the task twice, either under placebo or the dopamine D_2/D_3 receptor antagonist sulpiride, in a double-blind, counter-balanced, design. A prospect theory modelling approach was used to estimate subjective probability weighting and sensitivity to monetary outcomes. Consistent with prospect theory, we found that participants presented a distortion in the subjective weighting of probabilities, i.e. they overweighted low probabilities and underweighted moderate to high probabilities, both in the gain and loss domains. Compared with placebo, sulpiride attenuated this distortion in the gain domain. Across drugs, the groups did not differ in their probability weighting, although in the placebo condition, gamblers consistently underweighted losing probabilities. Overall, our results reveal that dopamine D_2/D_3 receptor antagonism modulates the subjective weighting of probabilities in the gain domain, in the direction of more objective, economically rational decision-making.

Introduction

A wealth of animal and human studies has implicated dopamine in risk-taking behaviour. Pharmacological studies in rats have shown that risk preferences are sensitive to manipulations of the dopaminergic system (Zeeb et al., 2009; Simon et al., 2009; St Onge and Floresco, 2009). Similarly, in humans, drugs enhancing or blocking dopamine have been shown to modulate risk-taking behaviour (Cools et al., 2003; Norbury et al., 2013; Rutledge et al., 2015; Rigoli et al., 2016). Furthermore, various genetic polymorphisms influencing dopamine metabolism or receptors have been associated with individual differences in risk preferences (Dreber et al., 2009; Kuhnen and Chiao, 2009; Mata et al., 2012). Yet, the specific neurocognitive mechanisms through which dopamine modulates risk-taking behaviour remain partly elusive. Some studies have suggested an influence via reward valuation mechanisms (Zhong et al., 2009) while other studies have shown that this influence is exerted via a change in value-independent gambling propensity (Rigoli et al., 2016; Rutledge et al., 2015; Timmer et al., 2017). Here we focus on a less well-investigated hypothesis, which is the role of dopamine on the subjective weighting of probabilities, both in healthy participants and individuals suffering from pathological gambling, a psychiatric disorder characterized by excessive risk-taking.

A useful and popular framework for examining how dopamine influences probability weighting is prospect theory (Kahneman and Tversky, 1979). Prospect theory posits that the departure of human agents from rational economic decision-making (i.e., expected value maximization) results from diminishing sensitivity to outcome value on the one hand, and non-linear weighting of probabilities on the other hand. People typically overweight low probabilities and underweight moderate to high probabilities, which results in an inverted-S-shaped probability weighting function and a diminished sensitivity to changes in probabilities in the medium range (Fig. 1*B*). A previous PET study in humans has shown that the degree of non-linear probability weighting in the gain domain is correlated with striatal dopamine D_1 receptor availability across subjects (Takahashi et al., 2010). Work with fMRI has also shown that probability distortion is accompanied by similarly distorted patterns of striatal BOLD activity (Hsu et al., 2009). Here, we aimed to establish a causal link between dopamine and probability distortion by adopting a within-subject, placebo-controlled, cross-over pharmacological design using the selective dopamine D_2/D_3 receptor antagonist sulpiride.

Dopamine has been linked to pathological gambling (PG, also called gambling disorder), an addictive disorder characterized by excessive financial risk-taking in the face of negative consequences. Initial evidence came from the clinical observation that a subset of patients with Parkinson's disease develop PG symptoms after receiving dopaminergic replacement therapy, including dopamine D_2/D_3

receptor agonists (Voon et al., 2009). This concurs with recent evidence showing that PG is characterized by a hyper-dopaminergic state (Boileau et al., 2014; van Holst et al., 2017). However, the specific mechanisms through which dopamine may act to foster PG remain elusive. In a previous study (Ligneul et al., 2013), we have shown that pathological gamblers show an elevation in their probability weighting function compared with healthy controls, reflecting an increased preference for risk or "optimism bias" in the gain domain (Gonzalez and Wu, 1999). Based on this observation, we aimed to test whether sulpiride could normalize risk-taking behaviour in pathological gamblers, by decreasing subjective probability weighting.

In the present pharmaco-behavioural study, pathological gamblers and healthy controls made choices between safe and risky options, both following intake of placebo and sulpiride. We used prospect theory modelling to estimate subjective probability weighting and sensitivity to outcome value, separately in the gain and loss domains. Our main objective was to assess the effect of sulpiride on the two main characteristics of the probability weighting function, i.e. non-linear distortion (sensitivity to changes in probability) and elevation (optimism bias). At a more exploratory level, we were also interested in comparing those effects in the gain and loss domains, given extensive literature showing differential effects of dopamine on gains versus losses (Frank et al., 2004; Pessiglione et al., 2006).

Materials and Methods

Participants.

We recruited 22 healthy controls and 22 pathological gamblers, following an in-depth structured psychiatric interview administered by a medical doctor (MINI Plus; Sheehan et al., 1998). One gambler was excluded because his data was accidentally not written to the log file for one drug session. One control participant and five gamblers were excluded due to extreme behaviours violating core assumptions of prospect theory (see *Statistical analysis* for more details). Therefore, the reported results are based on data from 21 controls and 16 gamblers. The present task was part of a larger study for which the participants were paid €50 on each session. The other tasks in the study were a reversal learning task (Janssen et al., 2015), a slot machine task measuring sensitivity to near-misses (Sescousse et al., 2016), and a mixed gamble task measuring loss aversion. All participants provided written informed consent, which was approved by the regional research ethics committee (Commissie Mensgebonden Onderzoek, region Arnhem-Nijmegen).

Pathological gamblers were recruited through advertisement (N = 13) and addiction clinics (N = 3). None of the gamblers was in treatment at the time of testing, except for one of them who was just starting

a cognitive behavioural therapy for his gambling problems. Controls were recruited through advertisement. All gamblers, with the exception of one, qualified as pathological gamblers ($\geqslant$ 5 DSM-IV criteria for pathological gambling; American Psychological Association, 2000). One gambler qualified as problem gambler as he met only four DSM-IV criteria. The severity of gambling symptoms was assessed using the South Oaks Gambling Screen (SOGS; Lesieur and Blume, 1987). All gamblers had a minimum SOGS score of 6 (range = 6–18), whereas controls, with the exception of two participants, had a SOGS score of 0 (range = 0–2).

The two groups were matched for age, net income, body mass index, and verbal IQ (Table 1). Participants were excluded if they consumed more than four alcoholic beverages daily; were using psychotropic medication; had a lifetime history of schizophrenia, bipolar disorder, attention deficit hyperactivity disorder, autism, eating disorder, anxiety disorder, or obsessive compulsive disorder; or had a past 6 months history of major depressive episode. Given the high co-morbidity between pathological gambling and other psychiatric disorders (Lorains et al., 2011), gamblers with the following co-morbidities were included: past cannabis dependence (> 5 months; N = 1); lifetime history of dysthymia (N = 1); and remitted post-traumatic stress disorder (remitted > 4 years; N = 1). One gambler also used cannabis weekly in the past 6 months, but did not meet the DSM-IV criteria for abuse/dependence. The control participants did not have any history of substance abuse or dependence. A number of self-report questionnaires were further used to characterize the participants (Table 1).

Pharmacological manipulation.

The dopamine D_2/D_3 receptor antagonist sulpiride (400 mg) or a placebo was administered to the participants before the experiment, using a within-subject, double-blind, counter-balanced design (placebo-sulpiride: 10 controls, 8 gamblers; sulpiride-placebo: 11 controls, 8 gamblers). The test sessions were separated by at least 1 week. Sulpiride was chosen as the dopamine-modulating drug in this study based on a few reasons. First, it is one of the more selective agents, acting selectively on dopamine D_2/D_3 receptors. As mentioned earlier, D_2/D_3 agents are known to cause pathological gambling symptoms in a subset of patients with Parkinson's disease. Moreover, sulpiride has been shown to modulate the sensitivity to reward and punishment during learning in human studies (Eisenegger et al., 2014; van der Schaaf et al., 2014). Background neuropsychological functioning, physiological measures and subjective mood were measured at several time points during the protocol, in order to check for non-specific effects of sulpiride; no such effects were observed. The risky decision-making task was performed approximately 3 h 15 min after drug intake, thus coinciding with high plasma concentrations of sulpiride (Mehta et al., 2003).

Experimental design and statistical analysis.

Experimental task. We used a "certainty equivalent" procedure (Fig. 1*A*) based on the protocol developed by Abdellaoui and colleagues (2008, 2011). Participants made several series of decisions between a sure amount of money (either a gain or a loss) and a gamble (either a pure-gain or pure-loss gamble). In each series of decisions, the gamble was fixed and the sure amount was iteratively adjusted in order to converge towards a "certainty equivalent" corresponding to the sure amount that felt subjectively equivalent to the gamble. There were 10 series of decisions (i.e., 10 different gambles) in the gain domain and 10 series of decisions in the loss domain (Table 2).

In each series of choices, the sure amount offered on the first trial corresponded to the expected value of the gamble. On subsequent trials, the sure amount was adjusted based on the previous choice according to the bisection (midpoint) method (Abdellaoui et al., 2011):

$$\text{sure amount} = \text{ lower bound} + \frac{(\text{upper bound} - \text{lower bound})}{2} \qquad (1)$$

where the lower bound was initialized with the amount *y* offered with probability 1–*p* in the gamble, and the upper bound was initialized with the amount *x* offered with probability *p* (Table 2). If the gamble was chosen, the lower bound was replaced with the value of the sure amount on that trial. If the sure option was chosen, it was the upper bound that was replaced with the value of the sure amount. This algorithm thus increased the sure amount in the subsequent trial if the gamble was chosen but decreased it if the sure option was chosen, driving the participant toward their "certainty equivalent", that is, the indifference point between the risky and safe options. The decision for each trial was self-paced, after which the participant's choice was highlighted on the screen. The participant did not receive any feedback.

Each series of decisions consisted of six trials, which is considered enough to provide reliable certainty equivalent estimates (Abdellaoui et al., 2011). In order to check for errors and random responses, each series ended with two control trials that required choosing between the gamble and a sure amount slightly above or below the estimated certainty equivalent. If the participant's response was not consistent with previous choices, the series was repeated. Participants were not explicitly informed about these control trials. We checked that the number of repetitions was not significantly different between healthy controls and pathological gamblers (gain domain: $Z = 0.55$, $p = .60$; loss domain: $Z = 1.31$, $p = .20$), between the placebo and sulpiride drug conditions (gain domain: $Z = 1.66$, $p = .098$; loss domain: $Z = 0.36$, $p = .72$), or between gains and losses in general ($Z = 1.47$, $p = .14$).

In total, participants went through a minimum of 160 experimental trials (10 series * [6 choices + 2 control trials] * 2 [gain/loss]). The task was the same in the loss domain but with negative amounts of money. Gain and loss trials were presented in separate blocks and the order of the blocks was counter-balanced across participants and drug sessions. The order of the gambles within gain and loss blocks was randomized. The task was performed on a computer and the task presentation was created with the Psychophysics Toolbox 2 (Brainard, 1997) for Matlab (www.mathworks.com).

Behavioural modelling. We used the semi-parametric method introduced by Abdellaoui et al. (2008, 2011; see also Fox and Poldrack, 2014) in order to estimate the value and probability weighting functions of prospect theory. This procedure was employed separately for gains and losses and for the drug and placebo conditions, within each individual participant.

In the first step of the procedure, the certainty equivalents of the gambles with varying amounts of money but a fixed probability of 2/6 (gamble indices i = 2, ..., 7 in Table 2) were used to estimate the probability weight $w(2/6)$ as well as the curvature of a parametrically defined version of the value function $v(\bullet)$. By definition, the utility of each gamble is equal to the utility of its certainty equivalent and, based on prospect theory, we can write:

$$v(CE) = w(p)\,v(x) + (1-w(p))\,v(y) \tag{2}$$

where CE is the certainty equivalent, x is the amount of money to be won with probability p and y is the amount of money to be won with probability $1-p$. Assuming a power function x^α for $v(\bullet)$ (Fox and Poldrack, 2014), where α quantifies sensitivity to outcome values, we can further write:

$$CE = [w(p)(x^\alpha - y^\alpha) + y^\alpha]^{\frac{1}{\alpha}} \tag{3}$$

Using a non-linear least squares procedure (*lsqcurvefit* function in Matlab), we could then estimate the optimal parameter values α and $w(2/6)$ that minimized the least squares $|CE(i) - \widehat{CE}(i)|$, where $\widehat{CE}(i)$ are the measured certainty equivalents for gambles indices i = 2, ...,7, expressed as:

$$\widehat{CE}(i) = [w(2/6)((x_i)^\alpha - (y_i)^\alpha) + (y_i)^\alpha]^{\frac{1}{\alpha}} \tag{4}$$

In the second step of the procedure, non-parametric estimates of the remaining probability weights $w(1/6)$, $w(3/6)$, $w(4/6)$ and $w(5/6)$ were derived from the certainty equivalents of the corresponding gambles (gamble indices i = 1, 8, 9 and 10 in Table 2). Since y = 0 in these gambles, based on equation (3) each probability weight can be calculated as follows:

$$CE = [w(p)x^\alpha]^{\frac{1}{\alpha}} \Leftrightarrow w(p) = \frac{CE^\alpha}{x^\alpha} \tag{5}$$

Based on those probability weights, we further derived a parametric estimation of the probability weighting function. We used a non-linear least squares procedure to estimate the two-parameter function proposed by Lattimore and colleagues (Lattimore et al., 1992), in which the sensitivity to changes in probabilities is quantified with distortion parameter γ, and the optimism about risk is quantified with elevation parameter δ:

$$w(p) = \frac{\delta p^\gamma}{\delta p^\gamma + (1-p)^\gamma} \tag{6}$$

In order to avoid local minima in our least squares estimations, we used an approach with randomized starting values. The two-step estimation procedure was run 200 times with starting values randomly drawn from [0, 5] for parameters α, δ, and γ, and from [0, 1] for $w(2/6)$. The resulting prospect theory parameters with the smallest squared norm of the residuals ('resnorm'), reflecting the goodness-of-fit between the model and the data, were selected for the subsequent statistical analysis.

Statistical analysis. One control participant and four pathological gamblers were excluded from subsequent group analyses based on the estimation of their individual probability weights. Indeed, for all these participants $w(1/6)$ was higher than $w(5/6)$ (in either the gain or loss domain), which violates the basic assumption of positive monotonicity in the evaluation of probabilities. One pathological gambler was further excluded due to extremely risk averse behaviour (α value over three standard deviations away from the mean) that seemed to result from highly self-controlled behaviour (based on debriefing with the participant).

Prospect theory parameters α, δ, and γ were compared across groups and drug conditions, separately in the gain and loss domains, using non-parametric statistics due to the non-normal distribution of the data. Main effects of the within-subject Drug factor were assessed using Wilcoxon tests. Main effects of between-subject Group factor were assessed using Mann-Whitney U tests, after parameters were averaged across drug sessions. Drug-by-Group interactions were examined with Mann-Whitney U

tests comparing sulpiride minus placebo values between the two groups. Bonferroni correction was used to correct for the six comparisons performed for each dependent variable (parameters α, δ, and γ): the two main effects of Drug and Group as well as their interaction, times the two contexts (gains and losses). Therefore, the corrected p-values correspond to the uncorrected p-values multiplied by 6. For effect sizes, we use the Common Language Effect sizes (CLE; Wuensch, 2015; Grissom and Kim, 2012) for intuitive interpretation. For the Mann-Whitney U tests, the CLE was calculated as U divided by the product of the two groups' sample sizes. For the Wilcoxon tests, the CLE was calculated as the number of positive differences (in favour of sulpiride over placebo) divided by the number of comparisons, that is, the total sample size. Therefore, the CLE represents the probability of a randomly selected value from one group/condition being higher than a randomly sampled value from the other group/condition. For both tests, there is no difference between the groups or conditions at CLE = .5.

Results

Table 3 reports group estimates for parameters α, δ, and γ in the study. Figure 3 illustrates the shape of the probability weighting function separately for gains and losses and both drug conditions in each group.

Sensitivity to changes in probabilities (distortion parameter γ)

A change in the distortion parameter γ of the probability weighting function represents a change in the non-linear weighting and thus the sensitivity to changes in probability. The distortion parameter γ did not significantly differ between control participants and pathological gamblers either in the gain domain, (Z = 1.47, p_{uncorr} = .15, CLE = .64) or the loss domain (Z = −1.13, p_{uncorr} = .27, CLE = .39).

However, there was a significant effect of the drug on γ in the gain domain (Z = 2.96, p_{uncorr} = .003, p_{corr} = .018, CLE = .70). Specifically, participants had higher values of γ under sulpiride (*Mdn* = 0.69) than under placebo (*Mdn* = 0.58), indicating lower levels of distortion of the probability weighting function in the sulpiride condition (Fig. 2). In the loss domain, there was no difference between placebo and sulpiride (Z = 0.36, p_{uncorr} = .72, CLE = .41). Drug effect did not interact with drug order in either the gain (Z = 1.46, p = .15, CLE = .65) or the loss domain (Z = 0.58, p_{uncorr} = .58, CLE = .56), indicating no reliable session effects. The drug effect (sulpiride–placebo) was not significantly different between control participants and pathological gamblers in the gain domain (Z = 0.55, p_{uncorr} = .60, CLE = .55) or in the loss domain (Z = −2.02, p_{uncorr} = .044, p_{corr} = .26, CLE = .30).

Optimism about risk (elevation parameter δ)

A change in the elevation parameter δ of the probability weighting function represents a shift in the weighting of the entire probability range, thus reflecting overall optimism or pessimism about risk. The elevation parameter δ did not significantly differ between control participants and pathological gamblers either in the gain domain ($Z = -1.41$, $p_{uncorr} = .17$, CLE = .36) or in the loss domain ($Z = -1.96$, $p_{uncorr} = .051$, $p_{corr} = .31$, CLE = .31). Moreover, there was no effect of drug either in the gain domain ($Z = -0.31$, $p_{uncorr} = .76$, CLE = .43) or in the loss domain ($Z = 0.39$, $p_{uncorr} = .70$, CLE = .59). Finally, the drug effect (sulpiride–placebo) was not significantly different between control participants and pathological gamblers in the gain domain ($Z = -0.74$, $p = .48$, CLE = .43) or in the loss domain ($Z = 1.57$, $p = .12$, CLE = .65).

For optimal comparison with our previous study in which we found a group difference in δ in the gain domain (Ligneul et al., 2013), we further compared the groups in the placebo condition alone. This analysis did not reveal a significant group difference in δ the gain domain ($Z = .03$, $p_{uncorr} = 1.0$, CLE = .50) but did reveal a significant difference in the loss domain ($Z = -2.9$, $p_{uncorr} = .003$, $p_{corr} = .018$, CLE = .22). Specifically, pathological gamblers had lower values of δ (*Mdn* = 0.42) than control participants (*Mdn* = 1.08), indicating lower elevation of the probability weighting function in the loss domain (Fig. 3*C*, *D*).

Sensitivity to outcomes (curvature parameter α)

Since our procedure also enabled to measure the curvature parameter of the value function, we also examined potential effects of group and drug. Non-parametric tests indicated that there was no significant difference between control participants and pathological gamblers either in the gain domain ($Z = 0.86$, $p_{uncorr} = .40$, CLE = .58) or in the loss domain ($Z = 1.17$, $p_{uncorr} = .25$, CLE = .61). Moreover, there was no effect of drug in the gain domain ($Z = 1.53$, $p_{uncorr} = .13$, CLE = .62) or in the loss domain ($Z = -1.21$, $p_{uncorr} = .23$, CLE = .41). Finally, the drug effect (sulpiride–placebo) was not significantly different between control participants and pathological gamblers in the gain domain ($Z = 1.96$, $p_{uncorr} = .051$, $p_{corr} = .31$, CLE = .69) or in the loss domain ($Z = -1.69$, $p_{uncorr} = .10$, CLE = .34).

Sensitivity analyses

In order to confirm the pattern of our main result on probability distortion, we performed an analysis of the probability weights themselves, which were obtained using a semi-parametric procedure, as opposed to the parametric estimation of γ. Specifically, we performed a 2 (Groups) x 2 (Drugs) x 5 (Probability levels: 1/6, 2/6, 3/6, 4/6 and 5/6) ANOVA on the probability weights $w(p)$ in the gain domain. We observed a significant interaction of Drug and Probability level on the $w(p)$ ($F(2.7, 94.495) = 3.21$, $p = .031$, $\eta^2 = .084$), thus strengthening our main result that sulpiride differentially modulates small versus medium-to-large

probability weights. However, matched samples post-hoc t-tests between the $w(p)$ for the two drug conditions failed to reach significance ($w(1/6)$: $t(36) = 1.15$, $p = .26$, $w(2/6)$: $t(36) = 1.39$, $p = .17$, $w(3/6)$: $t(36) = 0.62$, $p = .54$, $w(4/6)$: $t(36) = -0.15$, $p = .26$, $w(5/6)$: $t(36) = -1.41$, $p = .17$).

We also repeated our estimation procedure with different variations to check the robustness of our results despite small changes in the way the parameters were estimated. First, we estimated parameters δ and γ using the Prelec version of the probability weighing function (Prelec, 1998), instead of the Lattimore version (Lattimore et al., 1992). The Prelec function is defined by the following equation:

$$w(p) = e^{-\delta(-\ln p)^{\gamma}} \tag{7}$$

The parameters δ and γ have the same interpretation as in the Lattimore function, except that the degree of elevation decreases when the parameter δ increases. When the same analysis was conducted on the parameter estimates obtained with the Prelec function, the drug effect on the distortion parameter γ remained significant ($Z = 2.71$, $p = .007$, CLE = .70), emphasizing that sulpiride decreases the distortion of the probability weighting function (i.e., increases the parameter γ) compared with placebo.

In addition, the drug effect on the distortion parameter γ remained significant ($Z = 2.96$, $p = .003$, CLE = .70) when we used a linear value function ($\alpha = 1$) instead of a power function (x^{α}), a common assumption that we had made in our previous study (Ligneul et al., 2013).

Furthermore, the drug effect remained significant when we excluded the one participant with past cannabis dependence ($Z = 2.83$, $p = .005$, CLE = .69), as well as when all possible participants were included (i.e., no exclusion based on behavioural criteria, leading to 22 healthy controls and 21 pathological gamblers) ($Z = 3.50$, $p < .001$, CLE = .72).

Finally, in order to assess the accuracy of the parameter estimation, we ran a parameter recovery procedure (Heathcote et al., 2015). First, we used the parameter values from the original estimation to simulate new data. Specifically, we generated synthetic certainty equivalents for every gamble (i.e., 10 gambles in the gain the domain and 10 gambles in the loss domain) for each participant and each drug condition, using equation (3). We then used these simulated certainty equivalents in combination with the previously described semi-parametric procedure (Abdellaoui *et al.*, 2011), in order to estimate α, δ, and γ parameters again. Across-subject correlation coefficients between the original and the recovered parameter values were $r = 1$ in the case of α, δ and γ (in the case of $w(2/6)$: $r = .93$ in the gain domain and $r = .89$ in the loss domain), indicating efficient parameter recovery and high accuracy in the original parameter estimation. As expected, our main result indicating a significant drug effect on the elevation parameter γ retained the same effect size with the recovered parameter values, i.e. CLE = .70.

Discussion

This study investigated the effect of a dopaminergic manipulation on probability weighting during risk-taking in pathological gamblers and healthy participants. In line with our first hypothesis, we found that blocking dopamine D_2/D_3 receptors attenuated probability distortion in the gain domain. However, in contrast to our second hypothesis, the elevation of the probability weighting function was not affected by the dopaminergic manipulation and did not differ between pathological gamblers and healthy controls in the gain domain, even though a group difference was observed in the loss domain under placebo. Similarly, we did not find evidence for differences in sensitivity to outcomes between pathological gamblers and healthy controls, as well as no effect of the drug on the sensitivity to outcomes.

Our results demonstrate that the degree of non-linear probability weighting during decision-making is modulated by dopamine. More specifically, blocking D_2/D_3 receptors decreased probability distortion in the gain domain; this made participants more linear – or rational – in their overall assessment of probabilities, and thus more sensitive to changes in probabilities in the medium range. Such a differential effect of a dopaminergic agent on low versus high probabilities is consistent with several previous studies. For instance, Norbury et al. (2013) have shown that, in low sensation-seeking participants, the dopamine D_2/D_3 receptor agonist cabergoline increases risk-taking for high winning probabilities, while decreasing it for low winning probabilities. Similarly, Stopper et al. (2013) have shown that in rats, the administration of a dopamine D_1 receptor agonist increases risk-taking behaviour in the context of high winning probabilities, but decreases it in the context of low winning probabilities. Interestingly, in all these studies including ours, the interaction of dopaminergic drug effects with probability level led to more rational behaviour maximizing long-term expected value. Thus, it could be that, instead of inducing a shift in risk-taking, modulating dopamine might induce a shift in the adherence to the principle of expected value maximization. This is an intriguing hypothesis that would deserve to be formally tested in future studies.

Particularly relevant for the current study is the work of Takahashi et al. (2010), which to our knowledge is the only study that has explicitly investigated the role of dopamine in probability weighting. In their PET study, Takahashi et al. (2010) reported that lower dopamine D_1, but not D_2, receptor binding in the striatum was associated with higher levels of probability distortion. This seems partly at odds with the current results, which suggest that that D_2 receptor stimulation also plays a role in probability weighting. One hypothesis is that the drug effect observed in the current study could reflect a change in the balance between D_1- and D_2-receptor mediated activity in the direct and indirect pathways of the basal ganglia respectively, with sulpiride-induced D_2/D_3 receptor blockade being associated with a shift towards

Dopaminergic drug effects on probability weighting

D_1-receptor-dependent Go-pathway activity (Frank and O'Reilly, 2006; van der Schaaf et al., 2014; Jocham et al., 2011). Accordingly, we observed that D_2/D_3 receptor blockade decreases distortion, which is in line with the observation of Takahashi et al. that higher D_1 receptor binding in the striatum is also associated with less distortion.

A number of previous studies have shown that dopaminergic manipulations induce a global shift in risk attitudes, both in humans (Rigoli et al., 2016; Rutledge et al., 2015; Djamshidian et al., 2010; Riba et al., 2008; Timmer et al., 2017) and animals (Zeeb et al., 2009; Cocker et al., 2012; St Onge and Floresco, 2009). As mentioned previously, the lack of such an effect in our study could stem from the fact that, in contrast to most of these studies, which only manipulated one probability (or a limited range of probabilities), we considered the whole range of probabilities and observed opposite effects for high and low probabilities. Another distinctive feature of our experimental design is the absence of monetary feedback, which was meant to avoid contamination of the decision-making process by previous outcomes (Schonberg et al., 2011). This is important since risk attitudes, and in particular probability distortion, have been shown to differ when making decisions from description (as is the case in our study) versus from experience (i.e., based on feedback) (Hertwig and Erev, 2009). In addition, recent evidence in rats suggests that the influence of the dopamine D_2 pathway on risky behaviour is exerted via the signalling of prior outcomes (Zalocusky et al., 2016). Thus, the absence of feedback in our task could explain why the blockade of dopamine D_2 receptors failed to produce a global effect on risk attitudes. Interestingly, the vast majority of human studies reporting a global shift in risk-taking following a dopaminergic manipulation have used dopamine-enhancing agents such as L-Dopa. We are not aware of any studies reporting similar effects following dopamine D_2/D_3 receptor blockade.

We were not able to replicate our previous result showing an elevation of the probability weighting function in the gain domain (i.e., increased preference for risk) in pathological gamblers compared with healthy controls (Ligneul et al., 2013). One important methodological difference is that the monetary amounts used in the current study were much higher than in our previous study (300-1200€ versus 2-20€). It has been observed that people tend to be more risk-seeking for low-stake gambles than large-stake gambles, an observation referred to as the "peanuts effect" (Prelec and Loewenstein, 1991; see also Weber and Chapman, 2005). It is possible that the gamblers in our previous study were particularly sensitive to the peanuts effect and engaged in particularly high risk-seeking behaviour in the presence of low-stake gambles. It is also possible that the control participants in the current study happened to be more risk-seeking than average. The qualitative comparison of median values for the elevation parameter in the gain domain (Ligneul et al. 2013: $\delta_{Controls} = 0.74$, $\delta_{Gamblers} = 1.03$; current study: $\delta_{Controls} = 0.99$, $\delta_{Gamblers} = 0.90$) with typical values reported in the literature (Fox and Poldrack, 2014, Table A.3: median $\delta = 0.77$)

lends credence to these hypotheses: it seems that the control participants in the current study were more risk-seeking than average, while the gamblers were less-risking than in our previous study.

Another difference is that in our previous study we assumed a linear value function, whereas in the current study we estimated it empirically based on the certainty equivalents. Given the trade-off between prospect theory parameters α (curvature parameter of the value function) and δ (elevation parameter of the probability weighting function) in accounting for risk attitudes (Fox and Poldrack, 2014), it could be that part of the risk-seeking behaviour was absorbed by the α parameter in our current modelling procedure, whereas all of it was absorbed by the δ parameter in the previous study. Note however that our current results remained qualitatively unchanged when the estimation procedure was run with a linear value function.

While no group difference was observed in the gain domain, analyses restricted to the placebo condition revealed that, in the loss domain, pathological gamblers showed a significant decrease in the elevation of their probability weighting function compared with heathy controls (Fig. 3*C*, *D*). This observation implies a general underweighting of losing probabilities, which could contribute to the optimism bias and excessive risk-taking behaviour observed in pathological gamblers. However, given that this result was not predicted and only applies to the placebo condition, we prefer to refrain from speculating further before it is replicated.

This study is not without limitations. First, we had a modest sample size, due partly to the complexities of running pharmacological studies in patients, and the exclusion of several participants based on outlying behaviour and violations of basic prospect theory assumptions. Yet, in order to mitigate the increased likelihood of false positives (Poldrack et al., 2017), we implemented stringent Bonferroni correction for multiple comparisons, and demonstrated the convergence of results across various sensitivity analyses. Another limitation is the moderate test-retest reliability of decision-making measures in addictive disorders such as pathological gambling (Kräplin et al., 2016). This might have limited our ability to replicate our previous result (Ligneul et al., 2013) and more generally our ability to uncover true differences between groups or drugs.

In summary, this study provides evidence supporting the hypothesis that modulating dopamine affects how humans weight winning probabilities during decision-making. Dopamine D_2/D_3 receptor antagonism shifts probability weighting in the direction of more objective, economically rational decision-making. In future studies, it will be important to replicate this result, and further compare the contributions of D_1 and D_2/D_3 receptors with the same method, since the effect has now been observed in relation to both receptors (see Takahashi et al., 2010).

References

Abdellaoui M, Bleichrodt H, L'Haridon O (2008) A tractable method to measure utility and loss aversion under prospect theory. J Risk Uncertain 36:245–266.

Abdellaoui M, Diecidue E, Öncüler A (2011) Risk preferences at different time periods: an experimental investigation. Manage Sci 57:975–987.

Boileau I, Payer D, Chugani B, Lobo DSS, Houle S, Wilson AA, Warsh J, Kish SJ, Zack M (2014) In vivo evidence for greater amphetamine-induced dopamine release in pathological gambling: a positron emission tomography study with [(11)C]-(+)-PHNO. Mol Psychiatry 19:1305–1313.

Brainard DH (1997) The Psychophysics Toolbox. Spat Vis 10:433–436.

Cocker PJ, Dinelle K, Kornelson R, Sossi V, Winstanley CA (2012) Irrational choice under uncertainty correlates with lower striatal D2/3 receptor binding in rats. J Neurosci 32:15450–15457.

Cools R, Barker RA, Sahakian BJ, Robbins TW (2003) L-Dopa medication remediates cognitive inflexibility, but increases impulsivity in patients with Parkinson's disease. Neuropsychologia 41:1431–1441.

Djamshidian A, Jha A, O'Sullivan SS, Silveira-Moriyama L, Jacobson C, Brown P, Lees A, Averbeck BB (2010) Risk and learning in impulsive and nonimpulsive patients with Parkinson's disease. Mov Disord 25:2203–2210.

Dreber A, Apicella CL, Eisenberg DTA, Garcia JR, Zamore RS, Lum JK, Campbell B (2009) The 7R polymorphism in the dopamine receptor D4 gene (DRD4) is associated with financial risk taking in men. Evol Hum Behav 30:85–92.

Eisenegger C, Naef M, Linssen A, Clark L, Gandamaneni PK, Müller U, Robbins TW (2014) Role of dopamine D2 receptors in human reinforcement learning. Neuropsychopharmacology 39:2366–2375.

Fox CR, Poldrack RA (2008) Appendix: Prospect Theory and the Brain. In: Neuroeconomics: Decision Making and the Brain (Glimcher PW, Camerer CF, Fehr E, Poldrack RA, eds), pp 533–567. New York, NY: Academic Press.

Frank MJ, Seeberger LC, O'Reilly RC (2004) By carrot or by stick: cognitive reinforcement learning in Parkinsonism. Science 306:1940–1943.

Frank MJ, O'Reilly RC (2006) A mechanistic account of striatal dopamine function in human cognition: Psychopharmacological studies with cabergoline and haloperidol. Behav Neurosci 120:497–517.

Gonzalez R, Wu G (1999) On the shape of the probability weighting function. Cogn Psychol 38:129–166.

Heatherton TF, Kozlowski LT, Frecker RC, Fagerström K-O (1991) The Fagerström Test for Nicotine Dependence: a revision of the Fagerström Tolerance Questionnaire. Br J Addict 86:1119–1127.

Heathcote A, Brown SD, Wagenmakers E-J (2015) An Introduction to Good Practices in Cognitive Modeling. In: An Introduction to Model-Based Cognitive Neuroscience (Forstmann BU, Wagenmakers E-J, eds), pp 25–48. New York, NY: Springer New York.

Hertwig R, Erev I (2009) The description-experience gap in risky choice. Trends Cogn Sci 13:517–523.

Hsu M, Krajbich I, Zhao C, Camerer CF (2009) Neural response to reward anticipation under risk is nonlinear in probabilities. J Neurosci 29:2231–2237.

Janssen LK, Sescousse G, Hashemi MM, Timmer MHM, Ter Huurne NP, Geurts DEM, Cools R (2015) Abnormal modulation of reward versus punishment learning by a dopamine D2-receptor antagonist in pathological gamblers. Psychopharmacology 232:3345–3353.

Jocham G, Klein TA, Ullsperger M (2011) Dopamine-mediated reinforcement learning signals in the striatum and ventromedial prefrontal cortex underlie value-based choices. J Neurosci 31:1606–1613.

Kahneman D, Tversky A (1979) Prospect theory: an analysis of decision under risk. Econometrica 47:263–291.

Kräplin A, Scherbaum S, Bühringer G, Goschke T (2016) Retest reliabilities of decision-making and cognitive control measures in addictive disorders. Sucht 62:191–202.

Kuhnen CM, Chiao JY (2009) Genetic determinants of financial risk taking. PLoS One 4.

Lattimore PK, Baker JR, Witte AD (1992) The influence of probability on risky choice: A parametric examination. J Econ Behav Organ 17:377–400.

Lesieur H, Blume S (1987) The South Oaks Gambling Screen (SOGS): a new instrument for the identification of pathological gamblers. Am J Psychiatry 144:1184–1188.

Ligneul R, Sescousse G, Barbalat G, Domenech P, Dreher J-C (2013) Shifted risk preferences in pathological gambling. Psychol Med 43:1–10.

Lorains FK, Cowlishaw S, Thomas SA (2011) Prevalence of comorbid disorders in problem and pathological gambling: Systematic review and meta-analysis of population surveys. Addiction 106:490–498.

Mata R, Hau R, Papassotiropoulos A, Hertwig R (2012) DAT1 polymorphism is associated with risk taking in the Balloon Analogue Risk Task (BART). PLoS One 7: e39135.

Mehta MA, McGowan SW, Lawrence AD, Aitken MRF, Montgomery AJ, Grasby PM (2003) Systemic sulpiride modulates striatal blood flow: Relationships to spatial working memory and planning. Neuroimage 20:1982–1994.

Norbury A, Manohar S, Rogers RD, Husain M (2013) Dopamine modulates risk-taking as a function of baseline sensation-seeking trait. J Neurosci 33:12982–12986.

Pessiglione M, Seymour B, Flandin G, Dolan RJ, Frith CD (2006) Dopamine-dependent prediction errors underpin reward-seeking behaviour in humans. Nature 442:1042–1045.

Poldrack RA, Baker CI, Durnez J, Gorgolewski KJ, Matthews PM, Munafò M, Nichols TE, Poline J-B, Vul E, Yarkoni T (2017) Scanning the horizon: towards transparent and reproducible neuroimaging research. Nat Rev Neurosci 18:115–126.

Prelec D (1998) The probability weighting function. Econometrica 66:497–527.

Prelec D, Loewenstein G (1991) Decision making over time and under uncertainty: a common approach. Manage Sci 37:770–786.

Riba J, Krämer UM, Heldmann M, Richter S, Münte TF (2008) Dopamine agonist increases risk taking but blunts reward-related brain activity. PLoS One 3: e2479.

Rigoli F, Rutledge RB, Chew B, Ousdal OT, Dayan P, Dolan RJ (2016) Dopamine increases a value-independent gambling propensity. Neuropsychopharmacology 41:2658–2667.

Rutledge RB, Skandali N, Dayan P, Dolan RJ (2015) Dopaminergic modulation of decision making and subjective well-being. J Neurosci 35:9811–9822.

Saunders JB, Aasland OG, Babor TF, De la Fuente JR, Grant M (1993) Development of the Alcohol Use Disorders Identification Test (AUDIT): WHO collaborative project on early detection of persons with harmful acohol consumption‐II. Addiction 88:791‐804.

Schonberg T, Fox CR, Poldrack RA (2011) Mind the gap: bridging economic and naturalistic risk-taking with cognitive neuroscience. Trends Cogn Sci 15:11–19.

Sescousse G, Janssen LK, Hashemi MM, Timmer MHM, Geurts DEM, ter Huurne NP, Clark L, Cools R (2016) Amplified striatal responses to near-miss outcomes in pathological gamblers. Neuropsychopharmacology 41:2614–2623

Simon NW, Gilbert RJ, Mayse JD, Bizon JL, Setlow B (2009) Balancing risk and reward: a rat model of risky decision making. Neuropsychopharmacology 34:2208–2217.

St Onge JR, Floresco SB (2009) Dopaminergic modulation of risk-based decision making. Neuropsychopharmacology 34:681–697.

Stopper CM, Khayambashi S, Floresco SB (2013) Receptor-specific modulation of risk-based decision making by nucleus accumbens dopamine. Neuropsychopharmacology 38:715–728.

Takahashi H, Matsui H, Camerer CF, Takano H, Kodaka F, Ideno T, Okubo S, Takemura K, Arakawa R, Eguchi Y, Murai T, Okubo Y, Kato M, Ito H, Suhara T (2010) Dopamine D1 receptors and nonlinear probability weighting in risky choice. J Neurosci 30:16567–16572.

Timmer MHM, Sescousse G, van der Schaaf ME, Esselink RAJ, Cools R (2017) Reward learning deficits in Parkinson's disease depend on depression. Psychol Med:1–10.

Trepel C, Fox CR, Poldrack RA (2005) Prospect theory on the brain? Toward a cognitive neuroscience of decision under risk. Cogn Brain Res 23:34–50.

Van Der Schaaf ME, Van Schouwenburg MR, Geurts DEM, Schellekens AFA, Buitelaar JK, Verkes RJ, Cools R (2014) Establishing the dopamine dependency of human striatal signals during reward and punishment reversal learning. Cereb Cortex 24:633–642.

Van Holst RJ, Sescousse G, Janssen LK, Janssen M, Berry AS, Jagust WJ, Cools R (2017) Increased striatal dopamine synthesis capacity in gambling addiction. Biol Psychiatry 1–8.

Voon V, Fernagut PO, Wickens J, Baunez C, Rodriguez M, Pavon N, Juncos JL, Obeso JA, Bezard E (2009) Chronic dopaminergic stimulation in Parkinson's disease: from dyskinesias to impulse control disorders. Lancet Neurol 8:1140–1149.

Weber BJ, Chapman GB (2005) Playing for peanuts: why is risk seeking more common for low-stakes gambles? Organ Behav Hum Decis Process 97:31–46.

Zalocusky KA, Ramakrishnan C, Lerner TN, Davidson TJ, Knutson B, Deisseroth K (2016) Nucleus accumbens D2R cells signal prior outcomes and control risky decision-making. Nature 531:642–646.

Zeeb FD, Robbins TW, Winstanley CA (2009) Serotonergic and dopaminergic modulation of gambling behavior as assessed using a novel rat gambling task. Neuropsychopharmacology 34:2329–2343.

Zhong S, Israel S, Xue H, Sham PC, Ebstein RP, Chew SH (2009) A neurochemical approach to valuation sensitivity over gains and losses. Proc R Soc London B Biol Sci 276:4181–4188.

Zigmond AS, Snaith RP (1983) The Hospital Anxiety and Depression Scale. Acta Psychiatr Scand 67:361–370.

Figures and Tables

Figure 1. The gambling task and the probability weighting function of prospect theory. **A**, Each trial consisted of a self-paced choice between a sure option (on the left) and a risky gamble (on the right), followed by visual confirmation of the choice (a frame around the chosen option) and fixation. The sure amount in the next trial was adjusted based on the choice (increased if gamble was chosen, decreased if the sure option was chosen), with the gamble being fixed. After six choices, the sure amount that was reached provided an indifference point between the two options, defined as the 'certainty equivalent' of the gamble. A new series of choices involving a new gamble was then started (in total: 10 gambles in the gain domain and 10 gambles in the loss domain). No feedback was provided on the outcome of the choices. **B**, The solid black line represents a typical probability weighting function, with overweighting of low probabilities and underweighting of moderate to high probabilities. The dashed diagonal line represents neutrality with regard to sensitivity to probabilities.

Figure 2. Dopaminergic modulation of probability distortion. **A**, Boxplot illustrating the drug effect (sulpiride−placebo) on the distortion parameter *γ* of the probability weighting function in the gain domain, across all participants. Box height represents the interquartile range (IQR), black line represents the median, and whiskers represent the largest and smallest values no further than 1.5*IQR. Single data points are values are located outside the whiskers. **B**, Within-subject paired observations of *γ* estimates in the placebo and sulpiride conditions for both experimental groups (different illustration of the result presented in Fig. 2*A*). **C**, Fitted probability weighting function, based on the median estimates of *δ* (elevation) and *γ* (distortion) parameters across all participants. The shaded areas illustrate the variance of *γ* across participants, with the boundaries corresponding to the probability weighting function plotted with median *δ*, and 25th and 75th percentile *γ*.

Figure 3. Fitted probability weighting function based on group median estimates of δ (elevation) and γ (distortion). Across groups, sulpiride decreased probability distortion in the gain domain compared with placebo (panels **A**, **B**). When examining the placebo condition alone, pathological gamblers showed a decreased elevation of their probability weighting function in the loss domain compared with healthy controls (panels **C, D**).

Table 1. Demographic characteristics and questionnaire scores.

	Healthy controls (n = 21)			Pathological gamblers (n = 16)			
Variable	Range	M	SD	Range	M	SD	p
Age	18–52	32.1	11.4	21–50	35.8	8.8	.29
Net income (€)	0–3570	1691	1123	750–3250	1750	949	.87
Body mass index	18.3–30.9	23.1	3.2	20.8–26.9	23.9	2.0	.38
SOGS	0–2	0.2	0.5	6–18	12.4	3.9	<.001
FTND	0–5	0.6	1.4	0–8	2.5	2.9	.014
Number of current smokers	–	10	–	–	10	–	.37
AUDIT	0–14	6.2	3.8	0–15	7.7	4.6	.27
HADS anxiety	0–12	2.7	2.8	1–12	4.9	3.4	.035
HADS depression	0–10	1.6	2.3	0–15	4.9	4.4	.006
Verbal IQ	85–128	106	9.5	77–123	103	12.3	.43

M = Mean. SD = Standard Deviation. SOGS: South Oaks Gambling Screen; FTND: Fagerstrom Test for Nicotine Dependence (Heatherton et al., 1991); AUDIT: Alcohol Use Disorders Identification Test (Saunders et al., 1993); HADS: Hospital Anxiety and Depression Scale (HADS; Zigmond and Snaith, 1983).

Table 2. Gambles with varying outcomes and probabilities.

	Gamble index i									
	1	2	3	4	5	6	7	8	9	10
x	1200	1200	600	1200	600	1000	1200	1200	1200	1200
p	1/6	2/6	2/6	2/6	2/6	2/6	2/6	3/6	4/6	5/6
y	0	0	0	600	300	400	900	0	0	0

x is the larger amount of money in the gamble that could be won or lost with probability p. y is the smaller amount of money in the gamble that could be won or lost with probability 1−p. x and y are in €. For losses, the amounts of money were the same but negative.

Table 3. Estimates of prospect theory parameters.

Parameter	Controls				Gamblers			
	Placebo		Sulpiride		Placebo		Sulpiride	
	Median	IQR	Median	IQR	Median	IQR	Median	IQR
α_{gains}	0.74	0.66	0.80	0.42	0.80	0.37	1.16	0.81
α_{losses}	1.02	1.14	1.28	0.64	1.92	1.46	1.36	0.95
δ_{gains}	0.99	0.86	0.93	0.85	0.90	0.51	0.68	0.61
δ_{losses}	1.08	0.80	0.83	0.61	0.42	0.31	0.78	0.99
γ_{gains}	0.55	0.50	0.66	0.54	0.64	0.35	0.89	0.49
γ_{losses}	0.97	0.61	1.06	0.60	0.88	0.78	0.72	0.72

IQR = Interquartile range.