Adiposity QTL *Adip20*

5   Adiposity QTL *Adip20* decomposes into at least four loci when dissected using congenic strains

Cailu Lin, Brad D. Fesi, Michael Marquis, Natalia P. Bosak, Anna Lysenko, Mohammed Amin
Koshnevisan, Fujiko F. Duke, Maria L. Theodorides, Theodore M. Nelson, Amanda H.
McDaniel, Mauricio Avigdor, Charles J. Arayata, Lauren Shaw, Alexander A. Bachmanov and
Danielle R. Reed

Monell Chemical Senses Center, 3500 Market St, Philadelphia PA

Please address correspondence:
Danielle R. Reed, Ph.D.
Monell Chemical Senses Center
3500 Market St
Philadelphia PA 19104
reed@monell.org

Abstract

An average mouse in midlife weighs between 25 and 30 g, with about a gram of tissue in the

largest adipose depot (gonadal), and the weight of this depot differs between inbred strains.

Specifically, C57BL/6ByJ mice have heavier gonadal depots on average than do 129P3/J mice.

To understand the genetic contributions to this trait, we mapped several quantitative trait loci

(QTLs) for gonadal depot weight in an F_2 intercross population. Our goal here was to fine-map

one of these QTLs, *Adip20* (formerly *Adip5*), on mouse chromosome 9. To that end, we

analyzed the weight of the gonadal adipose depot from newly created congenic strains. Results

from the sequential comparison method indicated at least four rather than one QTL; two of the

QTLs were less than 0.5 Mb apart, with opposing directions of allelic effect. Different types of

evidence (missense and regulatory genetic variation, human adiposity/body mass index

orthologues, and differential gene expression) implicated numerous candidate genes from the

four QTL regions. These results highlight the value of mouse congenic strains and the value of

this sequential method to dissect challenging genetic architecture.

An average mouse at six months of age weighs about 25 to 30 g. Some of this weight is due to

adipose tissue located in several depots, the largest of which is the gonadal (**Figure 1**). In

rodents, depots can differ in weight up to 20-fold, depending on the particular strain, and while

adipose depot weight is highly heritable [1, 2], specific genetic determinants for each depot

reflect its specialized functions [3]. Investigators have interbred specific strains and looked for

chromosomal regions that affect the weight of individual adipose depots, and this procedure has

identified hundreds of influential genomic regions, or quantitative trait loci (QTLs), which

curators catalog in the Mouse Genome Database [4]. While it has been difficult to identify the

underlying genetic variants for many complex traits, including adiposity [5], it remains an

important scientific goal because this knowledge may better explain genetic risks and suggest

new biological pathways and therapeutic targets [6].

Our specific goal here was to map QTLs for gonadal adipose depot weight in mice and

to identify the underlying genetic variants using a pair of contrasting inbred strains. We began

QTL mapping by intercrossing the C57BL/6ByJ (B6) strain with the 129P3/J (129) strain [7-9]

and found a QTL on chromosome 9, which we originally called *Adip5* (at ~50 Mb) but was

renamed *Adip20*. Chromosomal mapping using intercross populations did not provide sufficient

resolution to narrow the genomic interval to specific candidate genes, so our next step was to

refine the location of this QTL by creating and studying mouse congenic strains. We assayed or

imputed genotypes for a common set of markers on mouse chromosome 9 and analyzed the

genotype-phenotype associations using several computational approaches, including a

sequential method of strain comparison [10]. Maternal effects and imprinting affect body

composition, including adipose depot weight, so we incorporated several experimental design

features to control these influences. After the mapping studies, we used several methods to

identify candidate genes.

Method

Animal husbandry. We bred all mice in the animal vivarium at the Monell Chemical

Senses Center, located in Philadelphia, Pennsylvania (USA), except for the inbred B6 and 129

mice used for the first stages of breeding, purchased from the Jackson Laboratory. During this

time, husbandry practices were stable, owing in part to having the same vivarium personnel and

in part to using a consistent type of bedding (Aspen Shavings, Northeastern Products Corp,

Warrensburg, NY). We fed mice Rodent Diet 8604 (Harlan Teklad, Madison, WI), and they lived

in a 12:12 light cycle, with lights off at 6 pm, barring unusual circumstances (e.g., weather-

related power outages). The Monell Institutional Animal Care and Use Committee approved

these procedures.

We started branching congenic mice by backcrossing N_6 (F_1 x B6) males (from

heterozygous consomic B6.129-Chr9 mice [11] with a partial donor region from chromosome 9)

with B6 females to produce mice with 129-derived donor chromosomal regions of various

lengths. We named the strains with codes that reflect their lineage; for instance, all strains with

the prefix 1 (e.g., 1.1) descended from the same male. The breeding goal was to identify male

breeders with a donor region that overlapped the QTL location discovered in the original F_2

intercross.

We bred congenic mice that had either zero or one copy of the 129-derived donor region

by mating inbred B6 female mice with heterozygous congenic males. This approach reduced

maternal effects (all mothers were the same genotype) and reduced imprinting effects (only

fathers contributed the donor region). Each congenic mouse was potentially genetically unique

(because the paternal donor region could shorten due to meiotic recombination). Therefore, we

genotyped each congenic mouse to ensure we could define the donor region breakpoints. While

most mice had full-length donor regions, some had shorter regions, which we refer to as

"partial."

Genotyping. We evaluated simple sequence-length polymorphism markers (e.g.,

D9Mit2) by polyacrylamide gel electrophoresis after polymerase chain reaction amplification by

locus-specific primers [12] in our laboratories. We assayed single nucleotide polymorphisms

(SNPs) at three locations: at the Genotyping and RNA Analysis Core at the Monell Chemical

Senses Center; at the Center for Inherited Disease Research (see Electronic Resources) as

part of an NIH-funded genotyping supplement; or by a commercial vendor (LGC, Beverly, MA;

formerly KBiosciences) as a fee-for-service. When assaying variants in the Monell genotyping

core, we used primers and allele-specific dye-labeled probes (Life Technologies, Carlsbad, CA).

Irrespective of genotype location, controls (blank samples, and genomic DNA from inbred

progenitors and their F_1 hybrids) were included in all genotyping assays, and we retested

unlikely genotypes as needed. We list all markers tested and their base pair positions in **S1**

Table. We imputed missing data by tracing the parental origin of the marker alleles and

assuming that no double recombination occurred between markers 26 Mb or less apart. We

selected this 26 Mb criterion as the smallest distance between two observed recombinant

events in the most densely genotyped populations we have studied [11]. We also genotyped 12

representative congenic mice to evaluate residual genetic background using the Mouse

Universal Genotyping Array (www.neogen.com) with 7,851 SNP markers built on the Illumina

platform. We tallied the numbers of heterozygous markers in the genetic background (excluding

the targeted donor regions) and expressed them as a fraction of the total valid markers for an

estimate of residual heterozygosity.

Phenotyping measures. After euthanasia by carbon dioxide asphyxiation, we weighed

the body of each mouse to the nearest 0.1 g. During necropsy, we removed the gonadal

adipose depot using anatomic landmarks [13, 14] and weighed the wet tissue to the nearest 0.1

g.

Gonadal adipose depot gene expression. We conducted two microarray experiments to

quantify gonadal depot adipose tissue gene expression between congenic mice with and

without the donor region (Experiments 1 and 2). In Experiment 1, we selected tissue from male

mice from congenic strain 4. We chose this strain because the donor region captured the

Adip20 peak we previously identified [15] (42.6 to 58.3 Mb). Half of the mice were heterozygous

for the donor region (129/B6; N=7), and half were littermates homozygous for the donor region

(B6/B6; N=7). Experiment 2 replicated Experiment 1 except there were N=6 mice in each

genotype group, for a total of 12 mice. We isolated RNA from each gonadal adipose depot and

measured genome-wide exon-by-exon expression using the Affymetrix Mouse Gene 1.0 ST

Array. Members of the University of Pennsylvania Microarray Core facility prepared the RNA,

performed the chip hybridization, and did preliminary quality control steps following

manufacturer directions.

Data analysis. We took three steps to prepare for the statistical analysis of the gonadal

adipose depot weights. First, we eliminated data from mice that (a) appeared to be sick (e.g.,

hunched posture) or (b) had tumors or other obvious gross morphology found during necropsy

(**S2 Table**). Second, we determined whether the data were normally distributed using the

fitdistrplus R package [16], log-transformed the data (**S1 Figure)** if needed, and confirmed this

transformation was effective using the Kolmogorov-Smirnov test (**S3 Table**). Third, we

evaluated the importance of body weight and age as covariates using Pearson correlation

coefficients. We observed that body weight was highly correlated with adipose depot weight but

that age, owing to the tight range (measured in days), was not (**S2 Figure**). Therefore, we used

body weight but not age as a covariate for the subsequent analyses:

1. Marker association analysis. We treated all congenic mice as one mapping population and

conducted a general linear model analysis of the log-transformed gonadal adipose depot weight

for each marker, using marker genotype and strain as fixed factors and body weight as a

covariate, and using a type 1 (sequential) sum of squares. For the presentation of the data, we

(a) computed marker association test statistics and converted the associated p-values to the

negative logarithm with base 10, (b) calculated the genotype mean, and (c) calculated the effect

sizes using Cohen's *D* [17] for the peak linked marker in the mapping population. We defined a

'significant' statistical threshold as an α level of 0.05 after Bonferroni correction for the number

of markers (N=148; $-\log_{10}(\alpha/N)$= 3.47) and a suggestive threshold as an α level of 0.63 ($-$

$\log_{10}(\alpha/N) = 2.37$) [18].

2. Common segment method. The common segment method is based on the assumption that a

chromosomal region of interest harbors one QTL. Therefore, in our case, if the QTL is within a

congenic donor region, there is a difference in gonadal adipose depot weight between host and

donor mice, whereas congenic strains that do not share that QTL-containing donor region have

no phenotype. Using this method, we conducted two analyses, one narrow and one broad. For

the narrow analysis, we considered only a subset of congenic mice, those mice with full-length

donor regions that were also from large genotype groups (N mice ≥ 12/strain/genotype; 12

strains). For the broad analysis, we used nearly all congenic mice as described below. For the

narrow and broad analyses, we grouped mice within each congenic strain by whether they had

a host or donor region, and compared these two groups using a general linear model with the

presence of the donor region and strain as fixed factors and body weight as a covariate. We

conducted the post hoc Fisher's least significance difference to compare the phenotype

between genotype groups for each congenic strain, and p<0.05 was used as a significance

level. Those strains with a significant difference between genotype groups we classified as

Adip20 positive (+), and those that were not different we classified as *Adip20* negative (-).

3. Sequential method. The sequential method can detect single and multiple QTLs [19] using an

established procedure to compare congenic strains in a predetermined order. To establish the

strain comparison order, we constructed a minimum spanning tree (MST) based on the donor

region size and location with a custom R-script following the previously published pseudocode

[19] supplemented with the R software package *optrees* [20] and *igraph.* We compared

heterozygous congenic mice (129/B6) to the group of all homozygous littermates (B6/B6). To

make comparisons between mice of different genotypes, we used a general linear model

followed by least-squares difference post hoc analysis with a statistical threshold of $p < 0.05$.

DNA sequence variants and human homology. Using an online database [21], we

identified all genes residing within the genomic coordinates of the QTLs defined in the

sequential congenic analysis. We next used another online database [22] containing genomic

variants among inbred mouse strains from a large-scale genome sequencing project [23, 24] to

compare the genomic sequences within the defined QTL regions for strains (129P2/OlaHsd and

C57BL/6J) closely related to parental inbred strains used in our study. We formatted these

variants using an online tool [25], and we identified regulatory variants as well as coding

variants (missense and stop codon gain/loss) with the potential to cause functional changes. We

determined which coding sequence variants were likely to be functional using computer

algorithms in the program Sorting Intolerant From Tolerant [26]. Eight SNPs were excluded from

this analysis because we could find no unique identifier for the variant in the dbSNP database

(Build 138) [27]. We also identified human genes and their variants associated with obesity that

are located in the regions of conserved synteny with the mouse QTL regions by searching an

online catalog of human genome-wide association results [28] with key words "adipose" and

"body mass index".

Analysis of microarray data. We performed gene expression analysis using Partek®

Genomics Suite® (version 6.6, build 6.16.0419) by importing files of raw intensities for each

probe set, performing the post-import quality control analyses, and comparing genotype groups

(host versus donor region) for differential gene expression [29-31]. We report the differential

expression as the log2-fold change (corresponding to a 1.5-fold change) with an associated p-

value (p<0.05) corrected for false discovery using the number of genes examined [32]. We

selected this relaxed fold-change cutoff to capture the anticipated small differences in gene

expression that often arise from allelic variation [33]. We present data from the relevant genes

using a customized volcano plot [34] annotating the resulting graph using the *ggrepel* [35] and

ggplot2 [36] R packages.

We compared the results between Experiment 1 and Experiment 2 and report only those

genes that consistently differed by genotype. We included genes within the entire congenic

donor region (not just the smaller QTL regions as defined below), reasoning that gene

regulatory elements can be located millions of base pairs from their protein-coding regions [37].

We also examined the genome-wide pattern of differential expression beyond the donor

fragment region of the congenic strain 4. We took this additional step to confirm our expectation

that more genes would differ inside than outside the congenic region. To that end, we identified

the top 200 differentially expressed genes (those with the lowest p-values) and used their

chromosomal locations to create an ideogram [38].

For all analyses listed above, we computed the statistical tests with R (version 3.2.0) and

R-studio (version 0.99.489) and graphed the results using GraphPad Prism 6 (version 6.05;

GraphPad Software, La Jolla, CA). All data are available for download on Github

(https://github.com/DanielleReed/Adip20) and the Center for Open Science (osf.io/yeqjf). All

mentions of mouse genomic coordinates refer to GRCm38 (mm10).

Results

Characteristics of congenic strains. We studied 1,293 congenic mice in total from 22

strains (**Table 1, S4 Table**) and confirmed the residual heterozygosity of these newly created

strains was low (<0.2%; **S4 Figure**). The average adipose depot weight of the adult congenic

mice ranged from 0.32±0.14 grams to 0.99±0.58 grams, depending on strain.

Marker association analysis. In an initial general linear model analysis of a combined

group of mice from all congenic strains, we estimated effect of marker genotype on gonadal

adipose depot weight (**Figure 2**). The result confirmed a linkage with a large effect size at ~52.5

Mb; this effect size was 0.5 unit of pooled standard deviation for two independent genotype

groups (**Figure 2C**), and 129-derived allele increase the phenotype (**Figure 2B**). However, the

result also indicated possible additional peaks (**Figure 2A**).

Common segment method. As expected, the common segment method was

uninformative for both the narrower and broader analyses, failing to point to one region with a

single QTL. In the narrow analysis, of the 12 congenic strains we evaluated, three were

Adip20(+) and nine were *Adip20(-)* (**S5-6 Tables**). The positive strains had no single region in

common. For the broad analysis, we used mice from 19 of the 22 strains (three strains had too

few mice of a particular genotype for statistical analysis; **S4, S8 Table**); of these strains, five

were *Adip20(+)* and fourteen were *Adip20(-)* (**S7-8 Tables)**. Like the narrow analysis, the broad

analysis indicated that the positive strains had no single region in common.

Sequential method. Using the sequential method, we analyzed the congenic strains with

group sizes N ≥ 12/genotype/strain (**S5 Table**). We put the 12 congenic strains that met this

sample-size criterion into one of three groups for comparison; the groupings and comparison

order (described below) were suggested by the results of the MST analysis (**Figure 3A**). Within

these groups, we compared pairs of strains, in the predetermined order, starting from

comparison of the host strain (all mice without the donor fragment) and the strain with the

smallest donor region (**Figure 3B**). The sequential analysis results indicated the presence of

four QTLs (**Figure 3C**), which we ordered from proximal to distal, starting with QTL1. For most

of the QTLs, the 129-derived allele increased the weight of the gonadal adipose depot, but

QTL3 had the opposite direction of effect (**Figures 3C and 4).**

Genetic variations. We analyzed genes residing in the QTL regions estimated in the

sequential congenic analysis: QTL1, 9:42779420-44142555; QTL2, 9:48789608-53884418;

QTL3, 9:54242579-54286770; and QTL4, 9:58262383-124595110. We refined the poorly

localized QTL4, drawing on marginally significant mapping results from the original F_2 intercross

[15], to marker *rs3721068* with a 10-Mb flanking region (QTL4, 9:108915010-118915010).

There were 44,009 variants between the 129P2/OlaHsd and C57BL/6J strains (closely related

to the B6 and 129 parent strains used in our study), which we extracted (**S3 Figure**). Of these

variants, the *in silico* analysis predicted that 2% change some aspect of mRNA regulation (**S9**

Table). There are 429 protein-coding genes [21], and 31% of the variants that map within the

coding sequence are missense variants. Specifically, 48 of these genes contain either

missenses or nonsenses variants (**S10 Table**).

Previously reported mouse and human QTLs. Using these QTL co-ordinates, we

surveyed the mouse genome database [4] for other adiposity QTLs on chromosome 9. Of the

eight previously reported QTLs (*Adip5, Adip14, Dob2, Mobq8, Obq5, Carfhg2, Plbcq5*, and

Tdmq1), seven QTLs had overlapping confidence intervals with the QTLs reported here (**S11**

Table). We also identified the regions of conserved synteny of human chromosomes 3, 11, and

15. We next examined results of recent human genome-wide association studies to find genes

associated with body mass index and adiposity [28] and residing in these regions (**Table 2**), and

to examine mouse orthologous genes with sequence variants. Four human genes (*FDX1,*

ARHGAP20, PDCD6IP, and *TRAK)* associated with adiposity and six (*C11orf53, NCAM1,*

DMXL2, *OSBPL10*, *LOC101928114,* and *CCK*) associated with body mass index (a commonly

used but indirect measure of human adiposity) resided in these regions. Among mouse

orthologues of these 10 genes, at least one, *Osbpl10*, has a protein-coding variant between

129P2/OlaHsd and C57BL/6J strains closely related to the B6 and 129 parental strains used in

our study.

Microarray results. The donor region of the congenic strain 4 (42.6 to 58.3 Mb; **Figure**

3C) has 277 genes, and in each microarray experiment, ~15% of these genes passed the

microarray differential expression filters with a 1.5-fold change and/or a false discovery rate of

p<0.05 (**Figure 5**). More than half of these genes were differentially expressed in both

microarray Experiments 1 and 2 (*Abcg4, Acsbg1, Alg9, Arcn1, C230081A13Rik, Cd3g, Cryab,*

Dixdc1, Fam55d, Fxyd6, Hmbs, Il10ra, Isl2, Mcam, Mpzl2, Mpzl3, Nnmt, Ptpn9, Sik2, and Sik3;

Figure 5 and **S12 Table**). Two genes that were differentially expressed had two missense

variants apiece (*Dixdc1* and *Sik2;* **S10 Table**). Five genes including two with the missense

variants contained putative cis-regulator variants (*Alg9, Cryab, Dixdc1, Mcam*, and *Sik2*; **S9**

Table). We also confirmed the expectation that differentially expressed genes cluster in the

congenic region rather than elsewhere in the genome (**S4 Figure**).

Integrating all methods above, there was no convergence to a single candidate gene

(**Figure 6**). Several genes did not survive the candidate gene filters used here but are

noteworthy near misses because their human orthologues are associated with similar traits in

genome-wide association studies (*Fdx1* and *Ncam1*; **Table 2**). While no single gene was

implicated consistently, several had more evidence than did others, for instance, *Dixdc1* and

Sik2, as mentioned above, were differentially expressed in gonadal adipose depots and

contained several putative regulatory and missense variants.

Discussion

Here we bred congenic strains with the expectation that we could narrow the genomic

location of an adiposity QTL, *Adip20*, to a region with only one or a very few genes on mouse

chromosome 9. The successful identification of several obesity genes, including leptin [39], the

leptin receptor [40], tubby [41, 42], and agouti [43], suggested this strategy would be useful

even when effect sizes differ by genetic context [44]. These early successes spawned other

efforts like ours to map genetic variants that contribute to obesity [45-62], and investigators have

identified hundreds of QTLs [4]. When we fine-mapped the *Adip20* locus, we detected effects of

at least four QTLs on chromosome 9. These results illustrate both the complexity of body

composition genetics and the utility of congenic strains to better characterize the underlying

genetic architecture and, further, suggest that decomposition of a single QTL into multiple QTLs

is commonplace for traits like adipose depot weight and obesity [63-65].

In general, the study of complex traits is easier when we reduce or eliminate extraneous

sources of trait variation. For obesity, sex-by-genotype interactions are a common feature of

natural populations, including humans [66], as well as mouse intercross studies [8, 67, 68].

Thus, we chose to study only male mice to reduce the potential interaction of a particular

genotype with sex, while acknowledging this choice makes the results less generalizable [69]. In

addition to reducing variation in adiposity due to sex, we controlled age as a source of potential

trait variation to reduce genotype-by-age interactions [70]. We also tried to reduce parent-of-

origin and maternal effects because of their known influence on adiposity [71, 72]. We controlled

some maternal effects in the sense that all congenic mice had mothers of the same genotype,

and we controlled some imprinting and other parent-of-origin effects because all congenic mice

inherited the donor region from the father (i.e., paternal transmission). We acknowledge that

maternal effects are a composite of many factors, not just maternal genotype, and so the

offspring may vary based on maternal body weight and parenting behavior. However, generally,

these experimental choices were useful in reducing certain types of variation.

Obesity as a trait is both simple and complex: it is simple conceptually because its single

hallmark is an excess of adipose tissue. However, it is complex because not all adipose tissue

is the same. It is stored in depots that differ in location, size, and function [13, 14, 73], and each

depot has a different genetic architecture [15] and pattern of gene expression [74, 75]. Further

complexities arise because adipose tissue is hard to measure directly, and each measurement

method yields different results [76]. Here we chose to study one adipose depot, which we

measured directly by necropsy. This is a simple measurement method, and our expectation was

that its genetic architecture would be less complex than for whole-body obesity. We specifically

chose the gonadal over other depots because it is the heaviest [13, 14, 73] and a proxy for

overall adiposity [77, 78]. However, despite all these complexities, both mouse and human

genetic approaches point to the same underlying mammalian biology: QTL regions found here

contain many mouse orthologues for obesity genes identified by human genome-wide

association studies.

We began the construction of congenic mice guided by mapping results from the

intercross population [15]. We also bred reciprocal consomics, which helped us define the

genetic architecture [11], and provided a source of breeder mice for constructing the congenic

strains. Building on information from the consomic mice study [11], we learned that a single

Adip20 QTL on chromosome 9 could not explain the observed results from the reciprocal strains

[11]. Study of the congenic mice here confirmed and extended those more general results,

uncovering the location of the multiple QTL and identifying their opposing effects.

Within the four QTL regions we identified, there are many candidate genes, and we see

a technical gap between available and ideal testing methods. The most common available

method is the total elimination of each gene (e.g., gene knockout) and quantification of knockout

versus control mice. However, this method has a serious limitation as an approach to study

obesity: large-scale knockout surveys indicate that inactivating mutations affect early embryonic

growth for nearly half of genes tested [79], and about a third of genes affect body composition in

adult mice [80, 81]. This method has poor specificity, so we cannot assume that if knockouts

differ from controls in body composition (e.g., [82]), then the target gene is causal. Likewise,

bioinformatics methods to evaluate functional variants are also not ideal because so many

genes have potentially functional variants. Furthermore, we cannot completely exclude any of

the thousands of observed variants because evaluating potential function with currently

available informatics methods remains imperfect. For complex traits, the challenge is now to find

new ways to winnow a large number of variants to only a few or to find a way to test many

variants and their combinations rapidly. While there are statistical methods to winnow the

candidate gene by trying to parse causal relationships among genotype gene expression and

trait [83-85], ultimately we need direct tests of individual genotypes on traits.

While congenic mapping methods were useful in identifying additional adiposity QTLs on

mouse chromosome 9, there are at least three limitations to consider when interpreting the

results. The study of male mice exclusively we mentioned earlier. A second limitation is the

residual heterozygosity within the congenic strains that we confirmed through genotyping. The

amount was low relative to the number of backcross cycles because we used marker-assisted

breeding, but still it may have contributed to experimental noise. To put this limitation in a

broader context, we have learned recently from whole-genome sequencing studies that this

approach will not ensure that all members of a particular inbred strain are completely genetically

identical [86]. Whether variants among members of the same inbred strain are technical artifacts

or represent true diversity is controversial, but if true this may explain in part the rapidly

divergence in body weight during selective breeding of inbred mouse strains [87]. The third

limitation is that the congenic breakpoints were concentrated more in the middle regions of the

chromosome, and we may have missed additional QTLs in the proximal region where there

were fewer breakpoints.

We used complementary methods to evaluate causal genes, but no single gene

emerged as a highly prioritized candidate within any QTL region. This lack of convergence

among methods may be due in part to the differing assumptions of each. For instance, the

microarray approach we used here assumes that causal variation affects gene expression in the

gonadal adipose depot of adult mice, but the actual causal variants may not affect gene

expression at all or may act in a different tissue or at a different time in development. Obviously,

the limitation of microarray experiments is the target tissue selection, and we acknowledge that

genotype effects in other tissue could be causal (e.g., liver tissue [84, 88]). These points made,

there was some convergence among methods. Specifically, the *OSBPL10* gene was identified

by a human genome-wide association study of body mass index [89], and the mouse orthologue

of this gene, which is within the QTL4 region, had a missense variant between the 129 and B6

mouse strains. Similarly, mouse gonadal adipose tissue expresses different amounts of *Dixdc1*

and *Sik2* mRNA, depending on QTL2 genotype, and each of the corresponding genes has two

predicted missense and several regulatory variants. These results provide direction in the future

evaluation of candidate genes.

Acknowledgements

We gratefully acknowledge the assistance with animal breeding of Rebecca James,

Liang-Dar (Daniel) Hwang, Zakiyyah Smith, Matt Kirkey, Amy Colihan, and Laurie Pippett. We

also acknowledge Richard Copeland and the consistent high-quality assistance of the animal

care staff at the Monell Chemical Senses Center and thank them for their service. Michael G.

Tordoff and Gary K. Beauchamp commented on a draft of the manuscript. Joseph H. Nadeau

provided valuable advice on analytical approaches and manuscript preparation. We thank the

two anonymous reviewers for their constructive comments on this manuscript.

<div align="center">Funding</div>

388   Accession ID

389   We used mice for these experiments, NCBI Taxon ID 10090.

390   Electronic Resources

391   http://www.cidr.jhmi.edu/nih/Completed_Projects_Report_June15.pdf

Adiposity QTL *Adip20*

Table 1. Characteristics of congenic mice by strains

Strain[a]	N	Age at necropsy (days)		Gonadal adipose (grams)	Body weight (grams)
		Range	Mean±SD[b]	Mean±SD	Mean±SD
1	113	175-187	180±2	0.54±0.24	31.77±2.93
1.1	176	162-184	180±2	0.55±0.28	31.44±2.96
1.1.1	3	180-180	180±0	0.99±0.58	36.13±4.72
1.2	14	180-182	181±1	0.61±0.28	33.43±4.24
3	175	177-190	180±2	0.56±0.24	31.18±3.19
3.1	120	178-182	180±1	0.55±0.25	31.64±3.21
3.1.1	106	178-186	180±1	0.50±0.23	31.55±3.11
3.1.1.1	127	179-184	181±1	0.39±0.17	31.10±2.26
3.1.1.2	34	179-181	180±1	0.40±0.17	31.21±2.58
3.1.1.3	14	179-181	180±1	0.56±0.38	31.23±3.77
3.1.1.4	63	103[c]-193	160±29	0.32±0.14	29.57±2.63
3.1.2	21	180-183	181±1	0.51±0.24	31.70±2.68
3.1.3	6	178-183	181±2	0.40±0.14	30.03±2.46
3.1.4	11	179-181	180±1	0.75±0.28	31.79±3.82
3.1.4.1	26	179-191	181±3	0.51±0.22	32.28±2.20
4	119	175-188	180±2	0.52±0.22	31.09±3.05
4.1	44	179-231	182±8	0.46±0.18	30.50±2.20
4.1a	22	177-190	180±3	0.51±0.19	31.31±2.42
4.2	16	180-181	180±0	0.52±0.18	30.90±2.51
4.3	18	179-183	181±1	0.51±0.18	31.11±2.55
4.4	61	178-193	181±3	0.48±0.28	30.95±2.89
4.5	4	179-181	180±1	0.90±0.45	34.89±4.53

[a]Substrain IDs. [b]SD=0 indicates less than one day. [c]We necropsied 10 younger mice ranging from 103 to 124 days of age.

Adiposity QTL *Adip20*

Table 2. Human genome-wide associations within mouse QTLs for adiposity and body mass index

QTL	Mouse QTL location	Human region	Adiposity	Body mass index
1	chr9:42779420-44142555	chr11:119308720-120693788	None	None
2	chr9:48789608-53884418	chr11:107691025-114109156	*FDX1, ARHGAP20*	*C11orf53, NCAM1*
3	chr9:54242579-54286770	chr15:51295990-51341931	None	*DMXL2*
4	chr9:108915010-118915010	chr3:27712199-48621406	*PDCD6IP, TRAK1*	*OSBPL10, LOC101928114, CCK*

This table lists chromosome: bp locations for mouse (GRCm38; mm10) and human (GRCh38). We obtained human orthologous regions by the UCSC genome converter [90]. "Adiposity" and "Body mass index" columns contain Human Gene Nomenclature Committee gene symbols.

Figure captions

Figure 1. Anatomical location of the gonadal adipose depot in a male mouse.

Figure 2. Detection of gonadal adipose depot weight QTLs on chromosome 9 by association analyses of a pooled congenic population. (**A**) Location of the markers in Mb on mouse chromosome 9 (mChr9); the y-axis is the –log p-values (black line) obtained in a general linear regression model analysis with body weight and strain as covariates (red line, significant threshold; gray line, suggestive). We defined the QTL confidence intervals by two units of –log10 p-value drop from the peak (blue line). For QTL1-QTL4 (Q1-Q4), the shaded peak areas correspond to QTL regions defined by the sequential method. (**B and C**) Average weight of the gonadal adipose depot by genotype (B6/B6, B6/129; **B**) and Cohen's *D* effect size (**C**) of *rs3723670*, the most associated marker (i.e., the marker with the lowest p value). *p<0.0001, difference between genotypes.

Figure 3. Dissection of QTLs with the congenic strains and the sequential method. (**A**) Minimum spanning tree (MST). (**B**) Mean and standard error of the mean (SEM) by strain for gonadal adipose depot weight in grams. Host = combined group of homozygous littermates without the donor fragment from all congenic strains; N = number of mice in each group. P-values are for least-significant-difference t-tests for each sequential comparison; red text indicates p-values that met the threshold (p<0.05). For the comparisons of two strains, the first strain was either greater (>), less than (<), or equal to (=) the second strain in gonadal adipose depot weight. "Model" lists QTLs within the donor fragment of each congenic strain suggested by the sequential analysis. (**C**) Congenic strains and genotypes for informative markers: A = B6/B6, shown as white bars; H = 129/B6, shown as dark gray bars; the recombinant ends correspond to both sides of the donor fragment for each congenic strain between H and A (or chromosome end), shown as light gray bars. QTLs are shown at the bottom as black boxes; arrows

show the direction of effect (e.g., ↑ indicates that the QTL allele from the 129 strain

increases the trait value). The data support the presence of four QTLs; we reported the

detailed logic for this conclusion in **S1Text**.

Figure 4. QTL effects. The least square means (dark red lines) and 95% lower and

upper confidence limits (blue lines) for QTL genotypes were obtained from the general

linear model with body weight as covariate in the congenic mouse data used for the

sequential analyses. We selected one marker in the middle of each QTL to plot each

QTL effect. The host (A) comprised littermates homozygous for B6/B6 from all strains;

we selected appropriate congenic mice (H) as suggested by the results of the sequential

analysis.

Figure 5. Extraction of differentially expressed genes: filtering on a volcano plot of the

microarray data analysis of the Experiment 1 (top) and Experiment 2 (bottom). In

Experiment 1, we selected the gonadal adipose tissue from male mice from congenic

strain 4 with the donor region captured QTL1, QTL2, and QTL3 (42.6 to 58.3 Mb). Half of

the mice were heterozygous for the donor region (129/B6; N=7), and half were

littermates homozygous for the donor region (B6/B6; N=7). Experiment 2 replicated

Experiment 1 except with N=6 mice in each genotype group, for a total of 12 mice. The

customized volcano plots depict the estimated $\log_2$-fold change (x-axis) and statistical

significance ($-\log_{10}$ p-value; y-axis). Each dot represents one gene. The green dots

show genes with absolute $\log_2$-fold change > 0.58 (i.e., 1.5-fold change) and false

discovery rate < 0.05; the red dots show genes with false discovery rate < 0.05; the

yellow dots show genes with absolute $\log_2$-fold change > 0.58; the black dots show

genes that did not reach these statistical thresholds. We label genes that passed any

filters by name. Twenty differentially expressed genes are reproducible between

Experiment 1 and 2, labeled with blue text and square boxes. Red stars show two genes

that passed the filters in one but not both experiments yet are noteworthy because their

human orthologues are associated with adiposity or body mass index in genome-wide

association studies (**Table 2**). We summarize the additional experimental details in **S12**

Table.

Figure 6. The overlap of candidate gene by nomination method. We extracted

candidate genes based on evidence from missense genetic variation, human adiposity,

or body mass index orthologues and differential gene expression.

References

1.   Reed DR, Bachmanov AA, Tordoff MG. Forty mouse strain survey of body composition. Physiol Behav. 2007;91(5):593-600. PubMed PMID: 17493645.

2.   Reed DR, Duke FF, Ellis HK, Rosazza MR, Lawler MP, Alarcon LK, et al. Body fat distribution and organ weights of 14 common strains and a 22-strain consomic panel of rats. Physiol Behav. 2011;103:523-9. Epub 2011/04/21. doi: 10.1016/j.physbeh.2011.04.006. PubMed PMID: 21504752.

3.   Fried SK, Lee MJ, Karastergiou K. Shaping fat distribution: New insights into the molecular determinants of depot- and sex-dependent adipose biology. Obesity (Silver Spring). 2015;23(7):1345-52. doi: 10.1002/oby.21133. PubMed PMID: 26054752.

4.   Eppig JT, Bult CJ, Kadin JA, Richardson JE, Blake JA, Anagnostopoulos A, et al. The Mouse Genome Database (MGD): from genes to mice--a community resource for mouse biology. Nucleic Acids Res. 2005;33(Database issue):D471-5. PubMed PMID: 15608240.

5.   Drinkwater NR, Gould MN. The long path from QTL to gene. PLoS Genet. 2012;8(9):e1002975. doi: 10.1371/journal.pgen.1002975. PubMed PMID: 23049490; PubMed Central PMCID: PMCPMC3462162.

6.   El-Sayed Moustafa JS, Froguel P. From obesity genetics to the future of personalized obesity therapy. Nat Rev Endocrinol. 2013;9(7):402-13. doi: 10.1038/nrendo.2013.57. PubMed PMID: 23529041.

7.   Reed DR, Li X, McDaniel AH, Lu K, Li S, Tordoff MG, et al. Loci on chromosomes 2, 4, 9, and 16 for body weight, body length, and adiposity identified in a genome scan of an F2 intercross between the 129P3/J and C57BL/6ByJ mouse strains. Mamm Genome. 2003;14(5):302-13. PubMed PMID: 12856282.

8.   Reed DR, McDaniel AH, Li X, Tordoff MG, Bachmanov AA. Quantitative trait loci for individual adipose depot weights in C57BL/6ByJ x 129P3/J F(2) mice. Mamm Genome. 2006;17(11):1065-77. PubMed PMID: 17103053.

9.   Bachmanov AA, Reed DR, Tordoff MG, Price RA, Beauchamp GK. Nutrient preference and diet-induced adiposity in C57BL/6ByJ and 129P3/J mice. Physiol Behav. 2001;72:603-13.

10.   Shao H, Sinasac DS, Burrage LC, Hodges CA, Supelak PJ, Palmert MR, et al. Analyzing complex traits with congenic strains. Mamm Genome. 2010;21(5-6):276-86. doi: 10.1007/s00335-010-9267-5. PubMed PMID: 20524000; PubMed Central PMCID: PMCPMC3805105.

11.   Lin C, Fesi BD, Marquis M, Bosak NP, Theodorides ML, Avigdor M, et al. Body composition qtls identified in intercross populations are reproducible in consomic mouse strains. PLoS One. 2015;10(11):e0141494. doi: 10.1371/journal.pone.0141494. PubMed PMID: 26551037; PubMed Central PMCID: PMCPMC4638354.

12.   Dietrich W, Katz H, Lincoln SE, Shin HS, Friedman J, Dracopoli NC, et al. A genetic map of the mouse suitable for typing intraspecific crosses. Genetics. 1992;131(2):423-47. Epub 1992/06/01. PubMed PMID: 1353738; PubMed Central PMCID: PMC1205016.

13.   Cinti S. The adipose organ. Milano: Editrice Kurtis; 1999.

14.   Hayakawa T, Yamasita H, Iwaki T. A Color Atlas of Sectional Anatomy of the Mouse: Braintree Scientific Inc; 2001.

15.   McDaniel AH, Li X, Tordoff MG, Bachmanov AA, Reed DR. A locus on mouse Chromosome 9 (Adip5) affects the relative weight of the gonadal but not retroperitoneal adipose depot. Mamm Genome. 2006;17(11):1078-92. PubMed PMID: 17103052.

16.   Delignette-Muller M, Pouillot, R., Denis, J., and Dutang, C. . fitdistrplus: Help to Fit of a Parametric Distribution to Non-Censored or Censored Data. R package. 2014.

17.   Cohen J. Statistical power analysis for the behavioral sciences. 2nd ed. Hillsdale, N.J.: L. Erlbaum Associates; 1988. xxi, 567 p. p.

18.   Lander E, Kruglyak L. Genetic dissection of complex traits: guidelines for interpreting and reporting linkage findings. Nat Genet. 1995;11:241-7.

19.   Shao H, Sinasac DS, Burrage LC, Hodges CA, Supelak PJ, Palmert MR, et al. Analyzing complex traits with congenic strains. Mammalian genome : official journal of the International Mammalian Genome Society. 2010;21(5-6):276-86. Epub 2010/06/05. doi: 10.1007/s00335-010-9267-5. PubMed PMID: 20524000.

20.   Fontenla M. optrees: Optimal Trees in Weighted Graphs. 1.0 ed. CRAN2014.

21.   Mus musculus (laboratory mouse) genome view [Internet]. National Center for Biotechnology Information 2015 [cited 10/7/2016]. Available from: https://www.ncbi.nlm.nih.gov/projects/mapview/map_search.cgi?taxid=10090&build=105.0.

22.   Mouse Genomes Project - Query SNPs, indels or SVs [Internet]. 2011 [cited 10/7/2016]. Available from: http://www.sanger.ac.uk/sanger/Mouse_SnpViewer/rel-1505.

23.   Keane TM, Goodstadt L, Danecek P, White MA, Wong K, Yalcin B, et al. Mouse genomic variation and its effect on phenotypes and gene regulation. Nature. 2011;477(7364):289-94. Epub 2011/09/17. doi: 10.1038/nature10413. PubMed PMID: 21921910.

24.   Yalcin B, Wong K, Agam A, Goodson M, Keane TM, Gan X, et al. Sequence-based characterization of structural variation in the mouse genome. Nature. 2011;477(7364):326-9. Epub 2011/09/17. doi: 10.1038/nature10432. PubMed PMID: 21921916.

25.   McLaren W, Gil L, Hunt S, Riat H, Ritchie G, Thormann A, et al. The Ensembl

Variant Effect Predictor. Genome Biology. 2016;17(1):122. doi: 10.1186/s13059-016-

0974-4.

26.   Anonymous. SIFT web server: predicting effects of amino acid substitutions on

proteins: Nucleic Acids Research; 2012 [cited 2016 10/7/2016]. August 15th

2015:[W542-7 (Link)]. Available from: http://siftdna.org/www/SIFT_dbSNP.html.

27.   Anonymous. dbSNP, Single Nucleotide Polymorphisms;

http://www.ncbi.nlm.nih.gov/SNP/index.html: National Center for Biotechnology

Information; 2016 [cited 2016 10/7/2016]. Available from:

http://www.ncbi.nlm.nih.gov/SNP/index.html.

28.   GWAS Catalog: the NHGRI-EBI Catalog of published genome-wide association

studies [Internet]. European Molecular Biology Laboratory. 2015 [cited 10/7/2016].

Available from: http://www.ebi.ac.uk/gwas/.

29.   de Buhr MF, Mahler M, Geffers R, Hansen W, Westendorf AM, Lauber J, et al.

Cd14, Gbp1, and Pla2g2a: three major candidate genes for experimental IBD identified

by combining QTL and microarray analyses. Physiol Genomics. 2006;25(3):426-34. doi:

10.1152/physiolgenomics.00022.2005. PubMed PMID: 16705022.

30.   de Ledesma AM, Desai AN, Bolivar VJ, Symula DJ, Flaherty L. Two new

behavioral QTLs, Emo4 and Reb1, map to mouse Chromosome 1: Congenic strains and

candidate gene identification studies. Mamm Genome. 2006;17(2):111-8. doi:

10.1007/s00335-005-0107-y. PubMed PMID: 16465591.

31.   Stylianou IM, Clinton M, Keightley PD, Pritchard C, Tymowska-Lalanne Z,

Bunger L, et al. Microarray gene expression analysis of the Fob3b obesity QTL identifies

positional candidate gene Sqle and perturbed cholesterol and glycolysis pathways.

Physiol Genomics. 2005;20(3):224-32. PubMed PMID: 15598878.

32.   Benjamini Y, Hochberg Y. Controlling the false discovery rate: A practical and powerful approach to multiple testing. Journal of the Royal Statistical Society Series B (Methodological). 1995;57(1):289-300.

33.   Zhuang Y, Adams KL. Extensive allelic variation in gene expression in populus F1 hybrids. Genetics. 2007;177(4):1987-96. doi: 10.1534/genetics.107.080325. PubMed PMID: 18073418; PubMed Central PMCID: PMCPMC2219518.

34.   Yick CY, Zwinderman AH, Kunst PW, Grunberg K, Mauad T, Dijkhuis A, et al. Transcriptome sequencing (RNA-Seq) of human endobronchial biopsies: asthma versus controls. Eur Respir J. 2013;42(3):662-70. doi: 10.1183/09031936.00115412. PubMed PMID: 23314903.

35.   J SKaI. Repulsive Text and Label Geoms for 'ggplot2'. CRAN2016. 2016.

36.   Wickham H CW, RStudio ggplot2: Create Elegant Data Visualisations Using the Grammar of Graphics. CRAN2016. 2016.

37.   Dixon JR, Selvaraj S, Yue F, Kim A, Li Y, Shen Y, et al. Topological domains in mammalian genomes identified by analysis of chromatin interactions. Nature. 2012;485(7398):376-80. doi: 10.1038/nature11082. PubMed PMID: 22495300; PubMed Central PMCID: PMCPMC3356448.

38.   Kin T, Ono Y. Idiographica: a general-purpose web application to build idiograms on-demand for human, mouse and rat. Bioinformatics. 2007;23(21):2945-6. doi: 10.1093/bioinformatics/btm455.

39.   Zhang Y, Proenca R, Maffei M, Barone M, Leopold L, Friedman JM. Positional cloning of the mouse obese gene and its human homologue. Nature. 1994;372(6505):425-32. PubMed PMID: 7984236.

40.   Tartaglia LA, Dembski M, Weng X, Deng N, Culpepper J, Devos R, et al. Identification and expression cloning of a leptin receptor, OB-R. Cell. 1995;83(7):1263-71. PubMed PMID: 8548812.

41.   Kleyn PW, Fan W, Kovats SG, Lee JJ, Pulido JC, Wu Y, et al. Identification and characterization of the mouse obesity gene tubby: a member of a novel gene family. Cell. 1996;85(2):281-90. PubMed PMID: 8612280.

42.   Noben-Trauth K, Naggert JK, North MA, Nishina PM. A candidate gene for the mouse mutation tubby. Nature. 1996;380(6574):534-8. PubMed PMID: 8606774.

43.   Bultman SJ, Michaud EJ, Woychik RP. Molecular characterization of the mouse agouti locus. Cell. 1992;71(7):1195-204. PubMed PMID: 1473152.

44.   Coleman DL. Obese and diabetes: two mutant genes causing diabetes-obesity syndromes in mice. Diabetologia. 1978;14(3):141-8. Epub 1978/03/01. PubMed PMID: 350680.

45.   Jerez-Timaure NC, Kearney F, Simpson EB, Eisen EJ, Pomp D. Characterization of QTL with major effects on fatness and growth on mouse chromosome 2. Obes Res. 2004;12(9):1408-20. Epub 2004/10/16. doi: 10.1038/oby.2004.177. PubMed PMID: 15483205.

46.   Ishimori N, Li R, Kelmenson PM, Korstanje R, Walsh KA, Churchill GA, et al. Quantitative trait loci that determine plasma lipids and obesity in C57BL/6J and 129S1/SvImJ inbred mice. J Lipid Res. 2004;45(9):1624-32. PubMed PMID: 15210844.

47.   Cheverud JM, Ehrich TH, Hrbek T, Kenney JP, Pletscher LS, Semenkovich CF. Quantitative trait loci for obesity- and diabetes-related traits and their dietary responses to high-fat feeding in LGXSM recombinant inbred mouse strains. Diabetes. 2004;53(12):3328-36. PubMed PMID: 15561968.

48.   Taylor BA, Wnek C, Schroeder D, Phillips SJ. Multiple obesity QTLs identified in an intercross between the NZO (New Zealand obese) and the SM (small) mouse strains. Mamm Genome. 2001;12(2):95-103. PubMed PMID: 11210195.

49.   Ogino T, Wei S, Wei K, Moralejo DH, Kose H, Mizuno A, et al. Genetic evidence for obesity loci involved in the regulation of body fat distribution in obese type 2 diabetes rat, OLETF. Genomics. 2000;70(1):19-25. PubMed PMID: 11087657.

50.   Horvat S, Bunger L, Falconer VM, Mackay P, Law A, Bulfield G, et al. Mapping of obesity QTLs in a cross between mouse lines divergently selected on fat content. Mamm Genome. 2000;11(1):2-7. PubMed PMID: 10602985.

51.   Brockmann GA, Kratzsch J, Haley CS, Renne U, Schwerin M, Karle S. Single QTL effects, epistasis, and pleiotropy account for two-thirds of the phenotypic F(2) variance of growth and obesity in DU6i x DBA/2 mice. Genome Res. 2000;10(12):1941-57. PubMed PMID: 11116089.

52.   Taylor BA, Tarantino LM, Phillips SJ. Gender-influenced obesity QTLs identified in a cross involving the KK type II diabetes-prone mouse strain. Mamm Genome. 1999;10(10):963-8. PubMed PMID: 10501964.

53.   Suto J, Matsuura S, Imamura K, Yamanaka H, Sekikawa K. Genetics of obesity in KK mouse and effects of A(y) allele on quantitative regulation. Mamm Genome. 1998;9(7):506-10. Epub 1998/07/10. PubMed PMID: 9657845.

54.   Keightley PD, Morris KH, Ishikawa A, Falconer VM, Oliver F. Test of candidate gene--quantitative trait locus association applied to fatness in mice. Heredity. 1998;81 (Pt 6):630-7. PubMed PMID: 9885188.

55.   Brockmann GA, Haley CS, Renne U, Knott SA, Schwerin M. Quantitative trait loci affecting body weight and fatness from a mouse line selected for extreme high growth. Genetics. 1998;150(1):369-81. PubMed PMID: 9725853.

56. York B, Lei K, West DB. Inherited non-autosomal effects on body fat in F2 mice derived from an AKR/J x SWR/J cross. Mamm Genome. 1997;8(10):726-30. Epub 1997/10/08. PubMed PMID: 9321464.

57. Taylor BA, Phillips SJ. Obesity QTLs on mouse chromosomes 2 and 17. Genomics. 1997;43(3):249-57. PubMed PMID: 9268627.

58. Lembertas AV, Perusse L, Chagnon YC, Fisler JS, Warden CH, Purcell-Huynh DA, et al. Identification of an obesity quantitative trait locus on mouse chromosome 2 and evidence of linkage to body fat and insulin on the human homologous region 20q. J Clin Invest. 1997;100(5):1240-7. PubMed PMID: 9276742.

59. York B, Lei K, West DB. Sensitivity to dietary obesity linked to a locus on chromosome 15 in a CAST/Ei x C57BL/6J F2 intercross. Mamm Genome. 1996;7(9):677-81. Epub 1996/09/01. PubMed PMID: 8703121.

60. Taylor BA, Phillips SJ. Detection of obesity QTLs on mouse chromosomes 1 and 7 by selective DNA pooling. Genomics. 1996;34(3):389-98. Epub 1996/06/15. doi: 10.1006/geno.1996.0302. PubMed PMID: 8786140.

61. West DB, Waguespack J, York B, Goudey-Lefevre J, Price RA. Genetics of dietary obesity in AKR/J x SWR/J mice: segregation of the trait and identification of a linked locus on chromosome 4. Mamm Genome. 1994;5(9):546-52. PubMed PMID: 8000138.

62. West DB, Goudey-Lefevre J, York B, Truett GE. Dietary obesity linked to genetic loci on chromosomes 9 and 15 in a polygenic mouse model. J Clin Invest. 1994;94(4):1410-6. PubMed PMID: 7929816.

63. Gularte-Merida R, Farber CR, Verdugo RA, Islas-Trejo A, Famula TR, Warden CH, et al. Overlapping mouse subcongenic strains successfully separate two linked body fat QTL on distal MMU 2. BMC Genomics. 2015;16:16. Epub 2015/01/24. doi:

10.1186/s12864-014-1191-8. PubMed PMID: 25613955; PubMed Central PMCID: PMC4308015.

64.   Farber CR, Medrano JF. Dissection of a genetically complex cluster of growth and obesity QTLs on mouse chromosome 2 using subcongenic intercrosses. Mamm Genome. 2007;18(9):635-45. PubMed PMID: 17694346.

65.   Mollah MB, Ishikawa A. Intersubspecific subcongenic mouse strain analysis reveals closely linked QTLs with opposite effects on body weight. Mamm Genome. 2011. Epub 2011/04/01. doi: 10.1007/s00335-011-9323-9. PubMed PMID: 21451961.

66.   Randall JC, Winkler TW, Kutalik Z, Berndt SI, Jackson AU, Monda KL, et al. Sex-stratified genome-wide association studies including 270,000 individuals show sexual dimorphism in genetic loci for anthropometric traits. PLoS genetics. 2013;9(6):e1003500. Epub 2013/06/12. doi: 10.1371/journal.pgen.1003500. PubMed PMID: 23754948; PubMed Central PMCID: PMC3674993.

67.   Farber CR, Medrano JF. Fine mapping reveals sex bias in quantitative trait loci affecting growth, skeletal size and obesity-related traits on mouse chromosomes 2 and 11. Genetics. 2007;175(1):349-60. PubMed PMID: 17110492.

68.   Zhao W, Ma C, Cheverud JM, Wu R. A unifying statistical model for QTL mapping of genotype x sex interaction for developmental trajectories. Physiol Genomics. 2004;19(2):218-27. PubMed PMID: 15304622.

69.   Clayton JA, Collins FS. Policy: NIH to balance sex in cell and animal studies. Nature. 2014;509(7500):282-3. PubMed PMID: 24834516.

70.   Cheverud JM, Routman EJ, Duarte FA, van Swinderen B, Cothran K, Perel C. Quantitative trait loci for murine growth. Genetics. 1996;142(4):1305-19. Epub 1996/04/01. PubMed PMID: 8846907; PubMed Central PMCID: PMC1207127.

71.   Wolf JB, Cheverud JM, Roseman C, Hager R. Genome-wide analysis reveals a complex pattern of genomic imprinting in mice. PLoS Genet. 2008;4(6):e1000091. PubMed PMID: 18535661.

72.   Lawson HA, Cheverud JM, Wolf JB. Genomic imprinting and parent-of-origin effects on complex traits. Nat Rev Genet. 2013;14(9):609-17. doi: 10.1038/nrg3543. PubMed PMID: 23917626; PubMed Central PMCID: PMCPMC3926806.

73.   Pond CM. The fats of life. Cambridge ; New York: Cambridge University Press; 1998. 337 p.

74.   Yamamoto Y, Gesta S, Lee KY, Tran TT, Saadatirad P, Kahn CR. Adipose depots possess unique developmental gene signatures. Obesity (Silver Spring). 2010;18(5):872-8. PubMed PMID: 20111017.

75.   Gesta S, Bluher M, Yamamoto Y, Norris AW, Berndt J, Kralisch S, et al. Evidence for a role of developmental genes in the origin of obesity and body fat distribution. Proc Natl Acad Sci USA. 2006;103(17):6676-81. PubMed PMID: 16617105.

76.   Halldorsdottir S, Carmody J, Boozer CN, Leduc CA, Leibel RL. Reproducibility and accuracy of body composition assessments in mice by dual energy x-ray absorptiometry and time domain nuclear magnetic resonance. Int J Body Compos Res. 2009;7(4):147-54. PubMed PMID: 21909234; PubMed Central PMCID: PMCPMC3169293.

77.   Fesi B, Marquis M, Lysenko A, James R, Lin C, Bachmanov A, et al. Refining fat phenotypes: multiple measures of body composition in 1,386 mice. Kroc Lecture and Symposium; University of Pennsylvania2013.

78.   Eisen EJ, Coffey MT. Correlated responses in body composition based on selection for different indicator traits in mice. J Anim Sci. 1990;68(11):3557-62. Epub 1990/11/01. PubMed PMID: 2262408.

79.   Dickinson ME, Flenniken AM, Ji X, Teboul L, Wong MD, White JK, et al. High-throughput discovery of novel developmental phenotypes. Nature. 2016;537(7621):508-14. doi: 10.1038/nature19356. PubMed PMID: 27626380.

80.   White JK, Gerdin AK, Karp NA, Ryder E, Buljan M, Bussell JN, et al. Genome-wide generation and systematic phenotyping of knockout mice reveals new roles for many genes. Cell. 2013;154(2):452-64. Epub 2013/07/23. doi: 10.1016/j.cell.2013.06.022. PubMed PMID: 23870131; PubMed Central PMCID: PMC3717207.

81.   Reed DR, Lawler MP, Tordoff MG. Reduced body weight is a common effect of gene knockout in mice. BMC Genet. 2008;9(1):4. PubMed PMID: 18182103.

82.   Koscielny G, Yaikhom G, Iyer V, Meehan TF, Morgan H, Atienza-Herrero J, et al. The International Mouse Phenotyping Consortium Web Portal, a unified point of access for knockout mice and related phenotyping data. Nucleic Acids Res. 2014;42(Database issue):D802-9. doi: 10.1093/nar/gkt977. PubMed PMID: 24194600; PubMed Central PMCID: PMCPMC3964955.

83.   Schadt EE, Lamb J, Yang X, Zhu J, Edwards S, Guhathakurta D, et al. An integrative genomics approach to infer causal associations between gene expression and disease. Nat Genet. 2005;37(7):710-7. doi: 10.1038/ng1589. PubMed PMID: 15965475; PubMed Central PMCID: PMCPMC2841396.

84.   Ishikawa A. Identification of a Putative Quantitative Trait Gene for Resistance to Obesity in Mice Using Transcriptome Analysis and Causal Inference Tests. PLoS One. 2017;12(1):e0170652. doi: 10.1371/journal.pone.0170652. PubMed PMID: 28114323; PubMed Central PMCID: PMCPMC5256930.

85.   Olsson AH, Volkov P, Bacos K, Dayeh T, Hall E, Nilsson EA, et al. Genome-wide associations between genetic and epigenetic variation influence mRNA expression and

insulin secretion in human pancreatic islets. PLoS Genet. 2014;10(11):e1004735. doi: 10.1371/journal.pgen.1004735. PubMed PMID: 25375650; PubMed Central PMCID: PMCPMC4222689.

86.   Oey H, Isbel L, Hickey P, Ebaid B, Whitelaw E. Genetic and epigenetic variation among inbred mouse littermates: identification of inter-individual differentially methylated regions. Epigenetics Chromatin. 2015;8:54. doi: 10.1186/s13072-015-0047-z. PubMed PMID: 26692901; PubMed Central PMCID: PMCPMC4676890.

87.   Keightley PD. Genetic basis of response to 50 generations of selection on body weight in inbred mice. Genetics. 1998;148(4):1931-9. PubMed PMID: 9560406.

88.   Raab RM, Bullen J, Kelleher J, Mantzoros C, Stephanopoulos G. Regulation of mouse hepatic genes in response to diet induced obesity, insulin resistance and fasting induced weight reduction. Nutr Metab (Lond). 2005;2:15. doi: 10.1186/1743-7075-2-15. PubMed PMID: 15985155; PubMed Central PMCID: PMCPMC1201568.

89.   Boardman JD, Domingue BW, Blalock CL, Haberstick BC, Harris KM, McQueen MB. Is the gene-environment interaction paradigm relevant to genome-wide studies? The case of education and body mass index. Demography. 2014;51(1):119-39. doi: 10.1007/s13524-013-0259-4. PubMed PMID: 24281739; PubMed Central PMCID: PMCPMC4035460.

90.   Kent WJ, Sugnet CW, Furey TS, Roskin KM, Pringle TH, Zahler AM, et al. The human genome browser at UCSC. Genome Res. 2002;12(6):996-1006. PubMed PMID: 12045153.

Supporting Information

S1 Table. Markers used for genotyping and their physical positions on chromosome 9

S2 Table. List of mice with tumors or otherwise excluded from analysis

Weight = Body weight in grams; Length, = body length in centimeters; Gonadal = gonadal adipose depot weight at the time of dissection in grams; Tumor location includes, weights if available

S3 Table. Statistics for descriptive and normality tests of gonadal adipose depot weight in congenic populations

S4 Table. Weight of gonadal depot and body weight (BW) of congenic mice with donor regions of different genotypes of donor regions

[a]Donor region genotype: Host = B6/B6; Donor = 129/B6.

[b]Donor region size: NA = mice without donor region; Partial = mice that inherited a partial-length (due to a recombination event) donor region from father; Full = mice that inherited full-length donor region from the father

S5 Table. Genotypes and phenotypes of the 12 congenic strains (strains with numbers of mice ≥12 mice/strain, and all mice having full-length donor regions)

Column headings show IDs of congenic strains. We show here only markers informative for breakpoints of the donor regions. A=B6/B6 genotype; H=129/B6 genotype; X=data are not available. Based on analysis of gonadal depot weight (**S4,6 Table**), we identified strains as Adip20(+) (shaded; strains with significant differences between mice with B6/B6 and 129/B6 genotypes) or Adip20(-) (no shading; strains with no significant differences between mice with B6/B6 and 129/B6 genotypes). The far right column ("Shared region") indicates whether a QTL-containing region is shared only by Adip20(+) but not by Adip20(-) strains (i.e., whether a marker genotype is H in all Adip20(+) strains

and A in all Adip20(-) strains). None of the regions of chromosome 9 met this single QTL only criterion.

S6 Table. Narrow analysis of gonadal depot weight in the 12 congenic strains (with numbers of mice ≥12 mice/strain, and all mice having full-length donor regions): general linear model (GLM), least square means, and post hoc Fisher's least significance difference (LSD) tests. N=Numbers of mice in each group with and without a donor region (donor genotype 129/B6; host genotype B6/B6). The donor group includes only mice with full-length length donor regions.

S7 Table. Broad strain inclusion criterion. Genotypes and phenotypes of the congenic strains of any sample size, except as noted below, including mice with partial or full-length donor regions. However, we still excluded three strains from this analysis (shaded in light grey) because they had very few mice (1.1.1, 3.1.3, 4.5). For other details, see **S5 Table**.

S8 Table. Broad analysis of gonadal depot weight using the criterion from **S7 Table**.

S9 Table. Regulatory variants within the four QTL regions predicted with an on-line tool Variant Effect Prediction[25]

S10 Table. Sorting Intolerant From Tolerant (SIFT) prediction of missense and stop codon gain/loss sequence variants of genes residing within the 4 adiposity QTL regions on chromosome 9

S11 Table. Summary of the mouse adiposity QTLs on chromosome 9

Type = mapping population; Con = congenic; Overlap refers to QTL1- QTL4 as identified by the sequential method. Approach = method of estimating confidence interval (LDR = logarithmic transformed probability; LOD = logarithm of the odds); Ref=reference.

S12 Table. Filtering of microarray data to identify reproducible candidate genes

[a]There are 277 genes within the donor region of strain 4 (42.6 to 58.3 Mb). Genes that passed the filters for differential expression in both Experiment 1 and 2 we display with blue font. Two genes in red font miss this criterion but their human orthologues are associated with adiposity or body mass index in GWAS studies (see **Table 2**). We show the remaining genes in black font.

S1 Text. Logic of the sequential method for each of the three branches of the minimum spanning tree (MST; central, left, and right) in **Figure 3**. Arrows after QTLs indicate the direction of effect of the 129-derived allele**.**

S1 Figure. Cullen and Frey graphs and empirical and theoretical data for untransformed and log-transformed congenics data. We tested the distribution of the gonadal adipose depot weight data for normality; **S3 Table** contains the parameters of the model-fitting untransformed and log-transformed data. CDF = cumulative distribution function.

S2 Figure. Pearson correlations between (A) gonadal adipose depot and body weight, (B) gonadal adipose depot and age as well as (C) body weight and age of all congenic mice. Gonadal = Gonadal adipose depot in grams; BW = body weight in grams; Age in days.

S3 Figure. Predicted effect for variants within four QTLs. In total, we uploaded 44,009 variants for the variant function analyses to the online tool Variant Effect Predictor; we found data from nearly all variants (99.8%) using this tool; of these variants, 2% were regulatory variants, and 31% of variants in coding regions were missense variants.

S4 Figure. Genome-wide overview of the microarray data from Experiment 1 with differentially expressed genes (blue dots) in adipose tissue between genotypes (129/B6 vs B6/B6) in congenic male mice. We observed potential heterozygosity (green dots) of this congenic strain in at least one of the samples at 25 of the 2,715 polymorphic markers. Of these, 19 were located in the congenic donor region (red dots), leaving 7

markers (green dots) with potential residual heterozygosity that may have produced

differential gene expression independent of the donor region.

Gonadal

Click here to access/download

Supporting Information

Supplemental Figure 1.docx

Click here to access/download
Supporting Information
Supplemental Figure 2.docx

Click here to access/download
Supporting Information
Supplemental Figure 3.docx

Click here to access/download
Supporting Information
Supplemental Figure 4.docx

Click here to access/download
Supporting Information
Supplemental Table 1.xlsx

Click here to access/download
Supporting Information
Supplemental Table 2.docx

Click here to access/download
Supporting Information
Supplemental Table 3.xlsx

Click here to access/download
Supporting Information
Supplemental Table 4.docx

Click here to access/download
Supporting Information
Supplemental Table 5.docx

Click here to access/download
Supporting Information
Supplemental Table 6.xlsx

Click here to access/download
Supporting Information
Supplemental Table 7.docx

Click here to access/download

Supporting Information

Supplemental Table 8.xlsx

Click here to access/download
Supporting Information
Supplemental Table 9.xlsx

Click here to access/download

Supporting Information

Supplemental Table 10.xlsx

Click here to access/download
Supporting Information
Supplemental Table 11.docx

Click here to access/download
Supporting Information
Supplemental Table 12.docx

Click here to access/download
Supporting Information
Supplemental Text 1.doc