Global expansion and redistribution of *Aedes*-borne virus transmission risk with climate change

Sadie J. Ryan[1,2,3*†], Colin J. Carlson[4,5*], Erin A. Mordecai[6], Leah R. Johnson[7]

Affiliations:

[1]Department of Geography, University of Florida, Gainesville, Florida, United States of America

[2] Emerging Pathogens Institute, University of Florida, Gainesville, Florida, United States of America

[3] School of Life Sciences, University of KwaZulu-Natal, Durban, South Africa

[4] Department of Biology, Georgetown University, Washington, DC 2007, U.S.A.

[5] National Socio-Environmental Synthesis Center, University of Maryland, Annapolis, Maryland 21401, U.S.A.

[6] Department of Biology, Stanford University, 371 Serra Mall, Stanford, California, United States of America

[7] Department of Statistics, Virginia Polytechnic and State University, 250 Drillfield Drive, Blacksburg, Virginia, United States of America

*These authors share lead author status.

[†] **Corresponding author:** sjryan@ufl.edu

Abstract: Forecasting the impacts of climate change on *Aedes*-borne viruses—especially dengue, chikungunya, and Zika—is a key component of public health preparedness. We apply an empirically parameterized Bayesian transmission model of *Aedes*-borne viruses for the two vectors *Aedes aegypti* and *Aedes albopictus* as a function of temperature to predict cumulative monthly global transmission risk in current climates, and compare with projected risk in 2050 and 2080 based on general circulation models (GCMs). Our results show that if mosquito range shifts track optimal temperatures for transmission (26-29 °C), we can expect poleward shifts in *Aedes*-borne virus distributions. However, the differing thermal niches of the two vectors produce different patterns of shifts under climate change. More severe climate change scenarios produce proportionally worse population exposures from *Aedes aegypti*, suggesting a strong coupling of climate change and continued emergence of dengue virus; but not from *Aedes albopictus* in the most extreme cases, suggesting a less straightforward climate link for chikungunya. Expanding risk of transmission from both mosquitoes will likely be a serious problem, even in the short term, for most of Europe; but significant reductions are also expected for *Aedes albopictus*, most noticeably in southeast Asia and west Africa. Within the next century, nearly a billion people are threatened with new exposure to both *Aedes* spp. in the worst-case scenario; but massive net losses in risk are noticeable for *Aedes albopictus*, especially in terms of year-round transmission, marking a global shift towards more seasonal risk across regions. Many other complicating factors (like mosquito range limits and viral evolution) exist, but overall our results indicate that while climate change will lead to both increased and new exposures to vector-borne disease, partial mitigation may accrue major unanticipated health costs from diseases that are mainly spread by *Aedes albopictus*.

Author Summary: The established scientific consensus indicates that climate change will

severely exacerbate the risk and burden of *Aedes*-transmitted viruses, including dengue,

chikungunya, Zika, West Nile virus, and other significant threats to global health security. Here,

we show that the story is more complicated, first and foremost due to differences between the

more heat-tolerant *Aedes aegypti* and the more heat-limited *Aedes albopictus.* Almost a billion

people could face their first exposure to viral transmission from either mosquito in the worst-case

scenario, especially in Europe and high-elevation tropical and subtropical regions. On the other

hand, while year-round transmission from *Ae. aegypti* is likely to expand (especially in south

Asia and sub-Saharan Africa), *Ae. albopictus* loses significant ground in the tropics, marking a

global shift towards seasonal risk as the tropics eventually become too hot for transmission.

Complete mitigation of climate change to a pre-industrial baseline could protect almost a billion

people from arbovirus range expansions; but middle-of-the-road mitigation may actually increase

exposure, especially for viruses mainly transmitted by *Aedes albopictus* (such as chikungunya).

Mitigating climate change also shifts the burden of both dengue and chikungunya (and

potentially other *Aedes* viruses) from higher-income regions back onto the tropics where

transmission might otherwise be curbed by rising temperatures.

Introduction

Climate change will almost certainly have a profound effect on the global distribution and

burden of infectious diseases [1–3]. Current knowledge suggests that the range of mosquito-

borne diseases could expand dramatically in response to climate change [4,5]. However, the

physiological and epidemiological relationships between mosquito vectors and the environment

are complex and often non-linear, and experimental work has showed an idiosyncratic

relationship between warming temperatures and disease transmission [6,7]. In addition,

pathogens can be vectored by related species, which may be sympatric, or several pathogens may

be transmitted by the same vector. Accurately forecasting the potential impacts of climate change

on *Aedes*-borne viruses—which include widespread threats like dengue and yellow fever, as well

as several emerging threats like chikungunya and Zika—thus becomes a key problem for public

health preparedness [4,8,9]. In this paper, we compare the roles and impact of two vectors, *Aedes*

aegypti (the primary vector of dengue virus around the world) and *Aedes albopictus* (the primary

vector of chikungunya virus in its recent emergence), in their contribution to potential

transmission landscapes in a changing climate.

The intensification and expansion of vector-borne disease is likely to be one of the most

significant threats posed by climate change to human health [2,10]. Mosquito vectors are of

special concern, due to the global morbidity and mortality from diseases like malaria and dengue

fever, as well as the prominent public health crises caused by (or feared from) several recently-

emergent viral diseases like West Nile, chikungunya, and Zika. The relationship between climate

change and mosquito-borne disease is perhaps best studied, in both experimental and modeling

work, for malaria and its associated *Anopheles* vectors. While climate change could exacerbate

the burden of malaria at local scales, more recent evidence challenges the "warmer-sicker world"

expectation [11,12]. The optimal temperature for malaria transmission has recently been

demonstrated to be much lower than previously expected [13], likely leading to net decreases in

optimal habitat at continental scales in the coming decades [12].

Relative to malaria, less is known about the net impact of climate change on *Aedes*-borne

diseases. At a minimum, the distribution of *Aedes* mosquitoes is projected to shift in the face of

climate change, with a mix of expansions in some regions and contractions in others, and no

overwhelming net global pattern of gains or losses [3,8]. Ecophysiological differences between

Aedes vector species are likely to drive differences in thermal niches, and therefore different

distributions of transmission risk [6,14], now and in the future. The consequences of those range

shifts for disease burden are therefore likely to be important, but are challenging to summarize

across landscapes and pathogens. Of all *Aedes*-borne diseases, dengue fever has been most

frequently modeled in the context of climate change, and several models of the potential future

of dengue have been published over the last two decades, with some limited work building

consensus among them [4]. Models relating temperature to vectorial capacity (the number of new

infectious mosquito bites generated from a human case), and applying general circulation models

(GCMs) to predict the impacts of climate change, date back to the late 1990s [5]. A study from

2002 estimated that the population at risk (PAR) from dengue would rise from 1.5 billion in

1990, to 5-6 billion by 2085, as a result of climate change [15]. A more recent study suggested

that climate change alone should increase the global dengue PAR by 0.28 billion by 2050, but

accounting for projected changes in global economic development (using GDP as a predictor for

dengue risk) surprisingly reduces the projected PAR by 0.12 billion over the same interval [16].

Mechanistic models have shown that increases or decreases in dengue risk can be predicted for

the same region based on climate models, scenario selection, and regional variability [17].

Chikungunya and Zika viruses, which have emerged more recently as a public health

crisis, are less well-studied in the context of climate change. A monthly model for chikungunya

in Europe, constrained by the presence of *Ae. albopictus*, found that the A1B and B1 scenarios

both correspond to substantial increases in chikunguya risk surrounding the Mediterranean [18].

A similar modeling study found that dengue is likely to expand far more significantly due to

climate change than Zika [9] (though epidemiological differences among these three viruses

remain unresolved [19–21]). However, the combined role of climate change and El Niño has

already been suggested as a possible driver of the 2016 Zika pandemic's severity [9]. Global

mechanistic forecasts accounting for climate change are all but nonexistent for both chikungunya

and Zika, given how recently both emerged as public health crises, and how much critical

information is still lacking in the basic biology and epidemiology of both pathogens.

In this study, we apply a new mechanistic model of the spatiotemporal distribution of

Aedes-borne viral outbreaks to resolve the role climate change could play in the emergence of

diseases like dengue, chikungunya, and Zika. Whereas other mechanistic approaches often rely

on methods like dynamic energy budgets to build complex biophysical models for *Aedes*

mosquitoes [22,23], and subsequently (sometimes) extrapolate potential epidemiological

dynamics [5], our approach uses a single basic cutoff for the thermal interval where viral

transmission is possible. The simplicity and transparency of the method masks a sophisticated

underlying model that links the basic rate of reproduction R_0 for *Aedes*-borne viruses to

temperature, via experimentally-determined physiological response curves for traits like biting

rate, fecundity, mosquito lifespan, extrinsic incubation rate, and transmission probability [6]. The

model is easily projected into geographic space by defining model-based measures of suitability

and classifying each location in space as suitable or not; we take a Bayesian approach in order to

take into account uncertainty in the experimental data. This threshold condition defines the

temperatures at which transmission is not prevented, rather than the more familiar threshold at

which disease invasion is expected ($R_0 > 1$, which cannot be predicted in the absence of

assumptions about vector and human population sizes and other factors). We then classify each

location by suitability in each month based on already published projections for current climates

in the Americas [6].

Here, we expand the framework for both *Ae. aegypti* and *Ae. albopictus* to project

cumulative months of suitability in current and future (2050 and 2080) climates, and further

examine how global populations at risk might change in different climate change scenarios. We

explore variation among both climate model selection (general circulation models; GCMs), and

potential emissions pathways described in the IPCC 5 (representative concentration pathways;

RCPs). In doing so, we provide the first mechanistic forecast for the potential future transmission

risk of chikungunya and Zika, which have been forecasted primarily via phenomenological

methods (like ecological niche modeling [9]). Our study is also the first to address the seasonal

aspects of population at risk for *Aedes*-borne diseases in a changing climate.

Methods

The Bayesian Model

Our study presents geographic projections of published experimentally-derived mechanistic

models of viral transmission by *Aedes aegypti* and *Aedes albopictus*. The approach to fit the

thermal responses in a Bayesian framework and combine them to obtain the posterior

distribution of R_0 as a function of these traits is described in detail in Johnson *et al.* [7] and the

particular traits and fits for *Aedes aegypti* and *Ae. albopictus* are presented in Mordecai *et al.*

[24]. In the original modeling study, the underlying data was compiled on transmission of

dengue virus by both mosquito species, and the models for each mosquito were subsequently validated on data compiled for three viruses (dengue, chikungunya, and Zika). Once we obtain our posterior samples for R_0 as a function of temperature we can evaluate the probability that $R_0 > 0$ (Prob($R_0 > 0$)) at each temperature, giving a distinct curve for each mosquito species. We then define cutoff of Prob($R_0 > 0$) = α to determine our estimates of the thermal niche; here, we use $\alpha = 0.975$. This very high probability allows us to isolate a temperature window for which transmission is almost certainly not excluded; this provides a conservative approach. For *Aedes aegypti*, these bounds are 21.3—34.0 °C, and for *Aedes albopictus*, 19.9—29.4 °C.

Current & Future Climates

Current mean monthly temperature data was derived from the WorldClim dataset (www.worldclim.org) [25] For future climates, we selected four general circulation models (GCMs) that are most commonly used by studies forecasting species distributional shifts, at a set of four representative concentration pathways (RCPs) that account for different global responses to mitigate climate change. These are the Beijing Climate Center Climate System Model (BCC-CSM1.1); the Hadley GCM (HadGEM2-AO and HadGEM2-ES); and the National Center for Atmospheric Research's Community Climate System Model (CCSM4). Each of these can respectively be forecasted for RCP 2.6, RCP 4.5, RCP 6.0 and RCP 8.5. RCP numbers correspond to increased radiation in W/m^2 by the year 2100, therefore expressing scenarios of increasing severity. (However, even these scenarios are nonlinear over time; for example, in 2050, RCP 4.5 is a more severe change than 6.0**.**) Climate model output data for future scenarios were acquired from the research program on Climate Change, Agriculture, and Food Security (CCAFS) web portal (http://ccafs-climate.org/data_spatial_downscaling/), part of the Consultative Group for International Agricultural Research (CGIAR). We used the model outputs created using the delta downscaling method, from the IPCC AR5. For visualizations presented in the main paper (Figure 2), we used the HadGEM2-ES model, the most commonly used GCM. The mechanistic transmission model was projected onto the climate data using the 'raster' package in R 3.1.1. Subsequent visualizations were generated in ArcMap.

Population at Risk

To quantify a measure of risk, comparable between current and future climate scenarios, we used
population count data from the Gridded Population of the World, version 4 (GPW4) [26],
predicted for the year 2015. We selected this particular population product as it is minimally
modeled *a priori*, ensuring that the distribution of population on the earth's surface has not been
predicted by modeled covariates that would also influence our mechanistic vector-borne disease
model predictions. These data are derived from most recent census data, globally, at the smallest
administrative unit available, then extrapolated to produce continuous surface models for the
globe for 5-year intervals from 2000-2020. These are then rendered as globally gridded data at
30 arc-seconds; we aggregated these in R ('raster'[30]) to match the climate scenario grids at 5
minute resolution (approximately 10 km^2 at the equator). We used 2015 population count as our
proxy for current, and explored future risk relative to the current population counts. This
prevents arbitrary demographic model-imposed patterns emerging, possibly obscuring climate-
generated change. We note that these count data reflect the disparities in urban and rural patterns
appropriately for this type of analysis, highlighting population dense parts of the globe.
Increasing urbanization would likely amplify the patterns we see, as populations increase overall,
and the lack of appropriate population projections at this scale for 30-50 years in the future
obviously limits the precision of the forecasts we provide. We thus opted for a most conservative
approach. We finally subdivide global populations into geographic and socioeconomic regions as
used by the Global Burden of Disease studies (**Figure S1**) [28].
203

Results
The current pattern of suitability suggested by our model based on mean monthly temperatures
(**Figure 1**) reproduces the known or projected distributions of dengue [29], chikungunya [30],
and Zika [9,31,32] well. For both *Ae. aegypti* and *Ae. albopictus,* most of the tropics is currently
optimal for viral transmission year-round, with suitability declining along latitudinal gradients.
Many temperate regions are suitable for up to 6 months of the year currently, but outside the
areas mapped as "suitable" by previous disease-specific distribution models, or where *Aedes*
mosquitoes are established; in some cases, limited outbreaks may only happen when cases are
imported from travelers (e.g. in northern Australia, where dengue is not presently endemic but
outbreaks happen in suitable regions [17]; or in mid-latitude regions of the United States, where
it has been suggested that traveler cases could result in limited autochthonous transmission

[31,33]). In total, our model predicts that 6.01 billion people currently live in areas suitable for

Ae. aegypti transmission at least part of the year (i.e., 1 month or more) and 6.33 billion in areas

suitable for *Ae. albopictus* transmission (using mean temperatures).

Even by 2050, warming temperatures are expected to produce dramatic expansions of

Aedes transmission risk (**Figure 2**). For *Aedes aegypti*, the pattern is (visually, even) fairly

straightforward: major expansions of one- or two-month transmission risk in temperate regions

are paired with expansion of year-round transmission in the tropics, even into the high-elevation

regions that were previously protected. *Aedes albopictus* transmission risk similarly expands

majorly (more so, even) into temperate regions, especially high latitude parts of Eurasia and

North America. But the upper thermal limits to *Aedes albopictus* transmission are passed in

many places, producing major reductions in regions of seasonal risk (like North Africa) and

year-round suitability (northern Australia, the Amazon basin, central Africa and southern Asia).

Whereas the conventional tropical-temperate gradient of mosquito-borne transmission is

preserved for *Aedes aegypti*, warming becomes so severe in the tropics that year-round *Aedes*

albopictus starts to look more unfamiliar, especially in the more extreme climate pathways. By

2080, year-round transmission by *Ae. albopictus* is mostly confined to high elevation regions,

southern Africa, and the Atlantic coast of Brazil; and even *Aedes aegypti* has begun to lose some

core area of year-round transmission in the Amazon basin especially.

Globally, our models suggest a net increase in population at risk from *Aedes*-borne virus

exposure, closely tracking the global rise in mean temperatures (**Figure 3**). For both mosquitoes,

populations at risk of any exposure will experience a major net increase by 2050, on the order of

roughly half a billion people; but even then, increases are more severe for *Ae. aegypti* than for

Ae. albopictus. But by 2080, the differences between the mosquitoes produce a different result:

while more severe warming continues to increase exposure for *Ae. aegypti*, up to nearly a billion

net new exposures, the most extreme expansions for *Ae. albopictus* are in middle of the road

scenarios (RCP 4.5 and 6.0). For year-round exposure, net changes tell an even more different

story between the two mosquitoes. For *Ae. aegypti*, warming temperatures lead to a net increase

of roughly 100-300 million people in areas of year-round transmission; though in fact, in RCP

8.5 by 2080, some parts of the tropics become so warm that even *Ae. aegypti* is no longer able to

transmit. But even by 2050 in the mildest scenarios, there are drastic net losses of year-round

transmission for *Aedes albopictus*, and these only become more severe – approaching roughly

700 million – in the warmest timelines.

Breaking these results down by region (**Table 1 & 2**) highlights just how much regional

velocity of climate change is likely to determine future disparities in global health. For *Aedes*

aegypti, the most notable net increases in all transmission risk are in all regions of Europe, with

additional notable gains in east Asia, high-elevation parts of central America and east Africa, and

the United States and Canada. But increases are expected across the board except in the

Caribbean, where minor net losses are expected across scenarios and years. In contrast, for *Aedes*

albopictus, there are much more regionally-variable changes anticipated. Major gains in Europe

are again expected across the board, as well as less significant increases in central America, east

Africa and east Asia, and the U.S. and Canada. But major net losses are also expected in several

regions, including tropical Latin America, western Africa, south Asia and most of all southeast

Asia, with a net loss of nearly 125 million people at risk by 2080 in RCP 8.5. In some cases, as

for western Africa and southeast Asia, the difference between RCP 6.0 and 8.5 is on the order of

50 and 100 million people respectively, highlighting just how large a (negative) difference minor

levels of climate mitigation might make in those regions.

For year-round transmission, the patterns are again less straightforward (**Table S1 & S2**),

but overall, they highlight a global shift towards more seasonal risk for both mosquitoes,

especially in the warmest scenarios. For *Aedes aegypti*, some of the largest net gains in people at

risk are expected in southern Africa, with additional notable increases expected in Latin

America. But even for *Ae. aegypti,* warming temperatures exceed levels suitable for year-round

transmission in some cases; for example, of all pathways, RCP 4.5 leads to the most severe

increases in southern Asia. Overall, almost 600 million people will move into areas with year-

round suitability for transmission, though the net increase will be much less (**Table S3**). For

Aedes albopictus, major net losses are expected in south and southeast Asia (totaling more than

400 million with the most extreme warming), and additional losses are expected in east Africa

and Latin America. Only the southern part of sub-Saharan Africa consistently experiences net

gains in year-round transmission risk; but gross increases are also expected in several regions,

most of all east Africa, moving roughly 250 million people into areas of year-round transmission

despite nearly triple that number in net losses.

We finally consider the idea of "first exposures" separately (gross gains, not accounting

for losses), which may be the most epidemiologically significant form of exposure. As the 2005

epidemic of chikungunya in India and the 2015 pandemic of Zika virus in the Americas

highlight, arboviral introductions into naïve populations can produce atypically severe outbreaks

on the order of millions of infections. We rank regions by these first exposures (**Table 3**), and we

find that consistently the most significant new exposures are expected in Europe and east Africa

for both mosquitoes, confirming fears that both regions may—as a consequence of climate

change, if not currently—be at risk from these types of black swan outbreaks. With an outbreak

of chikungunya virus in Kenya concurrent to the development of this study, these results seem

especially prescient.

Discussion

The dynamics of mosquito-borne illnesses are climate driven, and current work mostly suggests

that climate change will create massive opportunities for the expansion and intensification of

Aedes-borne illnesses within the next century. Especially since the emergence of Zika in the

Americas, many modeling studies have anticipated climate-driven emergence of dengue and

chikungunya at higher latitudes [34,35] and higher elevations [36,37]. Within this literature,

there have been several global studies of potential expansion [9,17,38], as well as significant

focused interest in North America and Europe (perhaps reflecting geographic biases in research

priorities and research institutions) [39]. The vast majority of this work has suggested that

climate change will probably increase the global burden of morbidity and mortality from dengue

and chikungunya, and therefore, that mitigation will likely improve global health outcomes

[40,41]. Perhaps most concerning are fears that *Aedes*-borne viruses will be introduced into

regions that have previously been unsuitable for transmission, given the potential for explosive

outbreaks (like Zika in the Americas, or chikungunya in India) when viruses are first introduced

into naïve populations [42]. The emergence of a Zika pandemic in the Old World [43], the

establishment of chikungunya in Europe beyond small outbreaks [18], or introduction of dengue

anywhere the virus (or any given serotype) has not recently been found, is still a critical concern.

Overall, our findings support the general view that climate change will produce major

expansions of *Aedes*-borne viral transmission risk. However, we also find more nuanced patterns

emerging between the two species, and among different climate pathways. The largest increases
in population at risk are consistently in Europe, with additional increases in high altitude regions
in the tropics (eastern Africa and the northern Andes) and in the United States and Canada.
These increases are expected not only for occasional exposure, but also for longer seasons of
transmission, especially for *Ae. aegypti*. But mosquitoes are adapted to their existing climatic
range, and while viral transmission will surely track warming temperatures into new places over
some intervals, there is no reason to think warming temperatures would produce a unilateral and
indefinite increase in disease transmission. Here we show that in the tropics, for *Aedes albopictus*
in particular, more extreme climate pathways produce warming temperatures that exceed the
suitable range for transmission in many parts of the world; and in the long term, even though
total exposure may increase from both mosquitoes in our study, we predict a global shift towards
seasonal regimes of exposure from *Aedes albopictus* and therefore, probably, chikungunya.
As warming temperatures may begin to exceed the upper thermal bounds of transmission,
this produces an unexpected problem in terms of climate change mitigation. Total mitigation
(down to pre-industrial baselines) would presumably prevent this redistribution of global risk.
But partial mitigation of climate change, i.e. less severe RCPs, could keep *Ae. albopictus*
mosquitoes especially within optimal thermal ranges for more of the year, and thereby produce
worse clinical outcomes. If our model continues to appropriately predict thermal bounds of viral
transmission into the future, this suggests a potential tradeoff in global health priorities, where
the scenarios that mitigate global dengue exposure, for example, might come with expanded risk
of chikungunya. But it is also unlikely that decisions about climate change mitigation will strictly
be decided based on global health priorities, given the already insufficient response to curb
carbon emissions and keep temperatures below the 2° C target [44]. In light of this, models such
as the ones we present here are probably most useful as a means to anticipate possible futures,
depending on the degree of mitigation achieved.
These global disease futures are inherently and incredibly stochastic, and the degree to
which our models correspond to reality depend not only on uncertainty about climate change, but
uncertainty about the biotic homogenization process for disease. For example, reductions in
transmission may be less prevalent than we expect here, as—even accounting for the velocity of
climate change—viruses will probably have sufficient time to adapt to warming temperatures
(within whatever evolvability they possess). Increases in transmission risk are also complicated

by many factors, such as the presence or absence of *Aedes* mosquitoes, which are also undergoing their own semi-independent range shifts facilitated by both climate change and human movement; our model already describes areas where *Aedes albopictus* and *Ae. aegypti* are absent but could be present in the future (and even now the ranges of these mosquitoes are not static, by any standard). Whether expanding transmission risk leads to future establishment and viral outbreaks depends not only on disease introduction, but also on land use patterns and urbanization at regional scales, a fact that may ultimately buffer some high-elevation regions like the Andes from increased disease risk [45,46].

In practice, these models are a first step towards an adequate understanding of potential global health futures, and the forecast horizon of these models will ultimately be determined by a number of confounding factors [47,48]. In particular, the link from transmission risk to clinical outcomes is confounded by other health impacts of climate change, including changing precipitation patterns, socioeconomic development, changing patterns of land use and urbanization, potential vector (and virus) evolution and adaptation to warming temperatures, and changing healthcare and vector management landscapes, all of which covary strongly (potentially leading to nonlinearities). Moreover, human adaptation to climate change will matter just as much as mitigation in determining how risk patterns shift; for example, increased drought stress will likely encourage water storage practices that increase proximity to *Aedes* breeding habitat [49]. Together these will determine the burden of *Aedes*-borne outbreaks, in ways that determine the eventual relevance of the forecasts we present here.

These models are certainly not the only ones to address this pressing topic, and different approaches control for data limitations, confounding processes, climate model uncertainty and disease model uncertainty, different concepts of population at risk, and preferences towards experimental, mechanistic, or phenomenological approaches differently. While climate change poses perhaps the most serious growing threat to global health security, the relationship between climate change and worsening clinical outcomes for *Aedes*-borne diseases is unlikely to be straightforward, and no single model will accurately predict the complex process of a global regime shift in *Aedes*-borne viral transmission. Our models only set an outer spatiotemporal bound to where transmission is thermally plausible; climate change is likely to change the risk-burden relationship at fine scales within those zones of transmission in nonlinear ways, such that areas with shorter seasons of transmission could still experience worse overall disease burdens,

or vice versa. Combining broad spatial models with finer-scale models of attack rates or outbreak size is a critical step towards bridging scales [43,50], but more broadly, research building consensus between all available models is of paramount importance [51]. This task is not limited to research on dengue and chikungunya; with several emerging flaviviruses on the horizon [52,53], and countless other emerging arboviruses likely to test the limits of public health infrastructure in coming years [54], approaches like ours that bridge the gap between experimental biology and global forecasting can be one of the foundational methods of anticipating and preparing for the "Next Big One."

Acknowledgements

This work was funded by the National Science Foundation (DEB-1518681 to SJR, LRJ, EAM, NSF DEB-1641145 to SJR, and DEB-1640780 to EAM), and CDC grant 1U01CK000510-01: Southeastern Regional Center of Excellence in Vector-Borne Diseases: the Gateway Program, to SJR. This publication was supported by the Cooperative Agreement Number above from the Centers for Disease Control and Prevention. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the Centers for Disease Control and Prevention. Support was also provided by the Stanford Woods Institute for the Environment (https:// woods.stanford.edu/research/environmental- venture-projects), and the Stanford Center for Innovation in Global Health (http://globalhealth. stanford.edu/research/seed-grants.html). Van Savage, Naveed Heydari, Jason Rohr, Matthew Thomas, and Marta Shocket provided helpful discussions on modeling approaches.

Author Information

The authors declare no competing interests. Correspondence and requests for materials should be addressed to S.J.R. (sjryan@ufl.edu).

Author Contributions

SJR and LJ ran the models, and CJC and SJR ran the analyses. CJC, SJR, EAM, and LJ wrote the manuscript. CJC and SJR made the figures.

References

1.   Hoberg EP, Brooks DR. Evolution in action: climate change, biodiversity dynamics and emerging
infectious disease. Phil Trans R Soc B. 2015;370: 20130553.

2.   Lafferty KD. The ecology of climate change and infectious diseases. Ecology. 2009;90: 888–900.

3.   Escobar LE, Romero-Alvarez D, Leon R, Lepe-Lopez MA, Craft ME, Borbor-Cordova MJ, et al.
Declining Prevalence of Disease Vectors Under Climate Change. Sci Rep. 2016;6.

4.   Messina JP, Brady OJ, Pigott DM, Golding N, Kraemer MU, Scott TW, et al. The many projected
futures of dengue. Nat Rev Microbiol. 2015;13: 230–239.

5.   Patz JA, Martens W, Focks DA, Jetten TH. Dengue fever epidemic potential as projected by general
circulation models of global climate change. Environ Health Perspect. 1998;106: 147.

6.   Mordecai E, Cohen J, Evans MV, Gudapati P, Johnson LR, Lippi CA, et al. Detecting the impact of
temperature on transmission of Zika, dengue, and chikungunya using mechanistic models. PLoS
Negl Trop Dis. 2017;11: e0005568.

7.   Johnson LR, Ben-Horin T, Lafferty KD, McNally A, Mordecai E, Paaijmans KP, et al.
Understanding uncertainty in temperature effects on vector-borne disease: a Bayesian approach.
Ecology. 2015;96: 203–213.

8.   Campbell LP, Luther C, Moo-Llanes D, Ramsey JM, Danis-Lozano R, Peterson AT. Climate
change influences on global distributions of dengue and chikungunya virus vectors. Phil Trans R
Soc B. 2015;370: 20140135.

9.   Carlson CJ, Dougherty ER, Getz W. An ecological assessment of the pandemic threat of Zika virus.
PLoS Negl Trop Dis. 2016;10: e0004968.

10. Githeko AK, Lindsay SW, Confalonieri UE, Patz JA. Climate change and vector-borne diseases: a
regional analysis. Bull World Health Organ. 2000;78: 1136–1147.

11. Ibelings B, Gsell A, Mooij W, Van Donk E, Van Den Wyngaert S, Domis D, et al. Chytrid
infections and diatom spring blooms: paradoxical effects of climate warming on fungal epidemics in
lakes. Freshw Biol. 2011;56: 754–766.

12. Ryan SJ, McNally A, Johnson LR, Mordecai EA, Ben-Horin T, Paaijmans K, et al. Mapping
physiological suitability limits for malaria in Africa under climate change. Vector-Borne Zoonotic
Dis. 2015;15: 718–725.

13. Mordecai EA, Paaijmans KP, Johnson LR, Balzer C, Ben-Horin T, Moore E, et al. Optimal
temperature for malaria transmission is dramatically lower than previously predicted. Ecol Lett.
2013;16: 22–30.

14. Brady OJ, Golding N, Pigott DM, Kraemer MU, Messina JP, Reiner Jr RC, et al. Global
temperature constraints on Aedes aegypti and Ae. albopictus persistence and competence for dengue
virus transmission. Parasit Vectors. 2014;7: 338.

15. Hales S, De Wet N, Maindonald J, Woodward A. Potential effect of population and climate changes
on global distribution of dengue fever: an empirical model. The Lancet. 2002;360: 830–834.

16.   Åström C, Rocklöv J, Hales S, Béguin A, Louis V, Sauerborn R. Potential distribution of dengue
436   fever under scenarios of climate change and economic development. Ecohealth. 2012;9: 448–454.

437   17.   Williams C, Mincham G, Faddy H, Viennet E, Ritchie S, Harley D. Projections of increased and
438   decreased dengue incidence under climate change. Epidemiol Infect. 2016; 1–10.

439   18.   Fischer D, Thomas SM, Suk JE, Sudre B, Hess A, Tjaden NB, et al. Climate change effects on
440   Chikungunya transmission in Europe: geospatial analysis of vector's climatic suitability and virus'
441   temperature requirements. Int J Health Geogr. 2013;12: 51.

442   19.   Funk S, Kucharski AJ, Camacho A, Eggo RM, Yakob L, Murray LM, et al. Comparative analysis of
443   dengue and Zika outbreaks reveals differences by setting and virus. PLoS Negl Trop Dis. 2016;10:
444   e0005173.

445   20.   Bastos L, Villela DA, Carvalho LM, Cruz OG, Gomes MF, Durovni B, et al. Zika in Rio de Janeiro:
446   assessment of basic reproductive number and its comparison with dengue. BioRxiv. 2016; 055475.

447   21.   Riou J, Poletto C, Boëlle P-Y. A comparative analysis of Chikungunya and Zika transmission.
448   Epidemics. 2017;

449   22.   Kearney M, Porter WP, Williams C, Ritchie S, Hoffmann AA. Integrating biophysical models and
450   evolutionary theory to predict climatic impacts on species' ranges: the dengue mosquito Aedes
451   aegypti in Australia. Funct Ecol. 2009;23: 528–538.

452   23.   Hopp MJ, Foley JA. Global-scale relationships between climate and the dengue fever vector, Aedes
453   aegypti. Clim Change. 2001;48: 441–463.

454   24.   Mordecai EA, Cohen JM, Evans MV, Gudapati P, Johnson LR, Lippi CA, et al. Detecting the
455   impact of temperature on transmission of Zika, dengue, and chikungunya using mechanistic models.
456   PLoS Negl Trop Dis. 2017;11: e0005568.

457   25.   Hijmans RJ, Cameron SE, Parra JL, Jones PG, Jarvis A. Very high resolution interpolated climate
458   surfaces for global land areas. Int J Climatol. 2005;25: 1965–1978.

459   26.   Center for International Earth Science Information Network (CIESIN), Columbia University.
460   Gridded Population of the World, Version 4 (GPWv4). [Internet]. US NASA Socioeconomic Data
461   and Applications Center (SEDAC); 2016. Available: http://sedac.ciesin.columbia.edu/data/set/gpw-
462   v4-population-count-adjusted-to-2015-unwpp-country-totals

463   27.   Hijmans RJ, van Etten J. raster: Geographic analysis and modeling with raster data. [Internet]. 2012.
464   Available: http://CRAN.R-project.org/package=raster

465   28.   Moran AE, Oliver JT, Mirzaie M, Forouzanfar MH, Chilov M, Anderson L, et al. Assessing the
466   global burden of ischemic heart disease: part 1: methods for a systematic review of the global
467   epidemiology of ischemic heart disease in 1990 and 2010. Glob Heart. 2012;7: 315–329.

468   29.   Bhatt S, Gething PW, Brady OJ, Messina JP, Farlow AW, Moyes CL, et al. The global distribution
469   and burden of dengue. Nature. 2013;496: 504–507.

30. Nsoesie EO, Kraemer M, Golding N, Pigott DM, Brady OJ, Moyes CL, et al. Global distribution
and environmental suitability for chikungunya virus, 1952 to 2015. Euro Surveill Bull Eur Sur Mal
Transm Eur Commun Dis Bull. 2016;21.

31. Samy AM, Thomas SM, Wahed AAE, Cohoon KP, Peterson AT. Mapping the global geographic
potential of Zika virus spread. Mem Inst Oswaldo Cruz. 2016;111: 559–560.

32. Messina JP, Kraemer MU, Brady OJ, Pigott DM, Shearer FM, Weiss DJ, et al. Mapping global
environmental suitability for Zika virus. Elife. 2016;5: e15272.

33. Bogoch II, Brady OJ, Kraemer M, German M, Creatore MI, Kulkarni MA, et al. Anticipating the
international spread of Zika virus from Brazil. Lancet Lond Engl. 2016;387: 335–336.

34. Ng V, Fazil A, Gachon P, Deuymes G, Radojević M, Mascarenhas M, et al. Assessment of the
probability of autochthonous transmission of Chikungunya virus in Canada under recent and
projected climate change. Environ Health Perspect. 2017;125.

35. Butterworth MK, Morin CW, Comrie AC. An analysis of the potential impact of climate change on
dengue transmission in the southeastern United States. Environ Health Perspect. 2017;125: 579.

36. Acharya BK, Cao C, Xu M, Khanal L, Naeem S, Pandit S. Present and Future of Dengue Fever in
Nepal: Mapping Climatic Suitability by Ecological Niche Model. Int J Environ Res Public Health.
2018;15: 187.

37. Equihua M, Ibáñez-Bernal S, Benítez G, Estrada-Contreras I, Sandoval-Ruiz CA, Mendoza-Palmero
FS. Establishment of Aedes aegypti (L.) in mountainous regions in Mexico: increasing number of
population at risk of mosquito-borne disease and future climate conditions. Acta Trop. 2017;166:
316–327.

38. Tjaden NB, Suk JE, Fischer D, Thomas SM, Beierkuhnlein C, Semenza JC. Modelling the effects of
global climate change on Chikungunya transmission in the 21 st century. Sci Rep. 2017;7: 3813.

39. Tjaden NB, Caminade C, Beierkuhnlein C, Thomas SM. Mosquito-borne diseases: advances in
modelling climate-change impacts. Trends Parasitol. 2017;

40. O'Neill BC, Done JM, Gettelman A, Lawrence P, Lehner F, Lamarque J-F, et al. The benefits of
reduced anthropogenic climate change (BRACE): a synthesis. Clim Change. 2018;146: 287–301.

41. Colón-González FJ, Harris I, Osborn TJ, São Bernardo CS, Peres CA, Hunter PR, et al. Limiting
global-mean temperature increase to 1.5–2° C could reduce the incidence and spatial spread of
dengue fever in Latin America. Proc Natl Acad Sci. 2018;115: 6243–6248.

42. Lucey DR, Gostin LO. The emerging Zika pandemic: enhancing preparedness. Jama. 2016;315:
865–866.

43. Siraj AS, Perkins TA. Assessing the population at risk of Zika virus in Asia–is the emergency really
over? BMJ Glob Health. 2017;2: e000309.

44. Hagel K, Milinski M, Marotzke J. The level of climate-change mitigation depends on how humans
assess the risk arising from missing the 2 C target. Palgrave Commun. 2017;3: 17027.

45.   Grau HR, Aide TM, Zimmerman JK, Thomlinson JR, Helmer E, Zou X. The ecological
507   consequences of socioeconomic and land-use changes in postagriculture Puerto Rico. AIBS Bull.
508   2003;53: 1159–1168.

509   46.   Li Y, Kamara F, Zhou G, Puthiyakunnon S, Li C, Liu Y, et al. Urbanization increases Aedes
510   albopictus larval habitats and accelerates mosquito development and survivorship. PLoS Negl Trop
511   Dis. 2014;8: e3301.

512   47.   Getz WM, Marshall CR, Carlson CJ, Giuggioli L, Ryan SJ, Romañach SS, et al. Making ecological
513   models adequate. Ecol Lett. 2018;21: 153–166.

514   48.   Petchey OL, Pontarp M, Massie TM, Kéfi S, Ozgul A, Weilenmann M, et al. The ecological
515   forecast horizon, and examples of its uses and determinants. Ecol Lett. 2015;18: 597–611.

516   49.   Beebe NW, Cooper RD, Mottram P, Sweeney AW. Australia's dengue risk driven by human
517   adaptation to climate change. PLoS Negl Trop Dis. 2009;3: e429.

518   50.   Perkins TA, Siraj AS, Ruktanonchai CW, Kraemer MU, Tatem AJ. Model-based projections of
519   Zika virus infections in childbearing women in the Americas. Nat Microbiol. 2016;1: 16126.

520   51.   Carlson CJ, Dougherty E, Boots M, Getz W, Ryan S. Consensus and conflict among ecological
521   forecasts of Zika virus outbreaks in the United States. Sci Rep. 2018;8: 4921.

522   52.   Evans MV, Murdock CC, Drake JM. Anticipating Emerging Mosquito-borne Flaviviruses in the
523   USA: What Comes after Zika? Trends Parasitol. 2018;

524   53.   Olival K, Willoughby A. Prioritizing the "Dormant" Flaviviruses. EcoHealth. 2017;14: 1–2.

525   54.   Gould EA, Higgs S. Impact of climate change and other factors on emerging arbovirus diseases.
526   Trans R Soc Trop Med Hyg. 2009;103: 109–121.

 Figures and Tables

529 **Figure 1 | Mapping current transmission risk.** Maps of current monthly suitability based on
530 mean temperatures for a temperature suitability threshold corresponding to the posterior
531 probability that scaled $R_0 > 0$ is 97.5% for (a) *Aedes aegypti* and (b) *Aedes albopictus*, and (c) the
532 number of people at risk (in billions) as a function of their months of exposure for *Aedes aegypti*
533 and *Aedes albopictus.*

535
536

Figure 2 | Mapping future transmission risk scenarios for *Aedes aegypti and Aedes albopictus*. Maps of monthly suitability based on a temperature threshold corresponding to the posterior probability that scaled $R_0 > 0$ is greater or equal to 97.5%, for transmission by *Aedes aegypti* and *Aedes albopictus* for predicted mean monthly temperatures under current climate and future scenarios for 2050: RCP 2.6 and RCP 8.5 in HadGEM2-ES.

544 **Figure 3 | Projected net changes in population at risk**. Projections are given as the net
545 difference in billions at risk, for *Aedes aegypti* and *Aedes albopictus* transmission, between
546 current maps and 2050 (top row) or 2080 (bottom row). Results are further broken down by
547 representative climate pathways (RCPs), each averaged across 4 general circulation models.
548

549
550
551

Table 1. Changing population at risk patterns for *Aedes aegypti*. All values are given in millions; future projections are averaged across GCMs, broken down by year (2050, 2080) and RCP (2.6, 4.5, 6.0, 8.5), and are given as net change from current population at risk. 0+/0- denote the sign of smaller non-zero values that rounded to 0.0, whereas "0" denotes true zeros.

Region	Current	2050				2080			
		2.6	4.5	6.0	8.5	2.6	4.5	6.0	8.5
Asia (Central)	69.9	8.4	10.5	9.9	12.2	8.1	11.8	12.5	15.6
Asia (East)	1,321.9	42.5	49.2	46.4	58.9	38.8	56.7	61.9	72.7
Asia (High Income Pacific)	164.0	-0.5	0+	-0.5	0.7	-0.6	0.6	1.0	1.7
Asia (South)	1,666.4	-0.1	1.6	0.7	3.7	-0.5	3.4	4.3	8.2
Asia (Southeast)	593.9	-2.1	0+	-0.6	2.3	-2.4	1.6	2.6	5.5
Australasia	12.9	3.6	5.7	5.3	6.7	4.3	6.2	6.9	8.0
Caribbean	40.4	-1.8	-1.7	-1.7	-1.6	-1.8	-1.6	-1.6	-1.5
Europe (Central)	22.7	44.2	71.8	69.0	83.3	59.0	79.3	85.5	90.6
Europe (Eastern)	41.3	57.9	110.4	93.5	133.9	80.0	124.7	130.7	156.2
Europe (Western)	114.6	47.2	132	112.0	166.8	90.3	156.4	180.8	220.9
Latin America (Andean)	31.3	2.8	3.4	3.3	4.0	2.6	3.9	4.1	5.5
Latin America (Central)	160.3	20.4	24.6	23.4	36	18.4	34.6	39.0	61.1
Latin America (Southern)	42.8	8.1	8.9	8.8	9.9	7.6	9.6	10.2	12.8
Latin America (Tropical)	181.8	19.2	19.5	19.5	19.6	18.9	19.6	19.7	19.8
North Africa & Middle East	439.5	19.7	24.1	23.8	27.2	19.3	25.9	27.3	30.3
North America (High Income)	281.9	36.2	48.3	42.6	55.0	37.8	53.6	57.1	62.8
Oceania	6.2	0.3	0.6	0.5	0.8	0.2	0.8	0.9	1.5
Sub-Saharan Africa (Central)	115.6	5.7	6.8	6.5	7.8	5.3	7.7	8.3	9.5
Sub-Saharan Africa (East)	274.8	48.8	63.7	59.1	72.2	44.7	70.8	76.6	90.9
Sub-Saharan Africa (Southern)	46.1	23.6	25.8	25.6	26.7	23.4	26.7	27.1	28.0
Sub-Saharan Africa (West)	384.0	-0.9	-0.7	-0.8	-0.7	-0.9	-0.6	-0.6	-0.4

Table 2. Changing population at risk patterns for *Aedes albopictus*. All values are given in millions; future projections are averaged across GCMs, broken down by year (2050, 2080) and RCP (2.6, 4.5, 6.0, 8.5), and are given as net change from current population at risk. 0+/0- denote the sign of smaller non-zero values that rounded to 0.0, whereas "0" denotes true zeros.

Region	Current	2050				2080			
		2.6	4.5	6.0	8.5	2.6	4.5	6.0	8.5
Asia (Central)	75.7	5.0	6.9	6.4	8.8	4.7	8.1	9.1	11.2
Asia (East)	1,367.0	16.1	20.8	18.9	25.2	15.0	24.0	26.5	32.4
Asia (High Income Pacific)	167.7	-2.6	-2.3	-2.6	-2.0	-2.7	-2.1	-1.9	-2.8
Asia (South)	1,673.8	-3.2	-1.7	-2.3	0+	-3.5	-0.5	-0.3	-19.1
Asia (Southeast)	602.5	-5.3	-3.8	-4.0	-6.7	-5.4	-8.5	-20.1	-124.8
Australasia	16.6	3.2	3.9	3.8	4.5	3.3	4.2	4.7	5.3
Caribbean	40.8	-1.8	-1.8	-1.8	-1.8	-1.9	-1.8	-1.8	-2.3
Europe (Central)	44.8	51.3	65.0	65.1	68.3	60.6	67.8	68.9	70.7
Europe (Eastern)	70.4	84.0	116.6	104.2	123.1	101.4	122.0	123.3	129.9
Europe (Western)	135.3	98.5	179.8	161.4	208.9	149.2	199.4	215.3	243.0
Latin America (Andean)	33.9	1.6	2.2	2.0	2.6	1.6	2.6	2.7	2.5
Latin America (Central)	179.1	21.9	27.0	27.9	31.1	17.6	29.7	30.3	23.6
Latin America (Southern)	50.4	3.2	3.6	3.6	4.8	2.8	4.2	4.9	7.6
Latin America (Tropical)	203.0	-1.5	-2.0	-1.6	-6.0	-1.5	-5.6	-8.0	-26.3
North Africa & Middle East	455.0	10.6	13.0	12.9	14.2	10.4	13.5	14.1	11.8
North America (High Income)	311.6	20.6	28.4	26.0	32.1	22.6	31.6	32.3	34.7
Oceania	6.8	0.5	0.8	0.6	1.0	0.4	0.9	1.0	0.8
Sub-Saharan Africa (Central)	120.8	2.8	3.5	3.3	4.2	2.5	4.1	4.4	-3.8
Sub-Saharan Africa (East)	320.2	30.3	39.1	36.3	42.4	27.9	41.8	42.8	34.2
Sub-Saharan Africa (Southern)	70.1	3.4	3.8	3.8	3.9	3.4	4.0	4.0	4.3
Sub-Saharan Africa (West)	384.9	-1.4	-1.5	-1.5	-2.0	-1.4	-1.9	-3.5	-59.0

Table 3. Top 10 regional increases in overall transmission risk (one or more months).
Regions are ranked based on millions of people exposed for the first time to any transmission risk; parentheticals give the net change (first exposures minus populations escaping transmission risk). All values are given for the worst-case scenario (RCP 8.5) in the longest term (2080).

Aedes aegypti		*Aedes albopictus*	
1. Europe (Western)	224 (220.9)	1. Europe (Western)	246.2 (243)
2. Europe (Eastern)	156.4 (156.2)	2. Europe (Eastern)	130.1 (129.9)
3. Sub-Saharan Africa (East)	92.8 (90.9)	3. Europe (Central)	71 (70.7)
4. Europe (Central)	90.9 (90.6)	4. Sub-Saharan Africa (East)	58.1 (34.2)
5. Asia (East)	81.7 (72.7)	5. Latin America (Central)	51.9 (23.6)
6. North America (High Income)	65.7 (62.8)	6. Asia (East)	41.4 (32.4)
7. Latin America (Central)	62 (61.1)	7. North America (High Income)	37.7 (34.7)
8. North Africa & Middle East	34.3 (30.3)	8. North Africa & Middle East	19.4 (11.8)
9. Sub-Saharan Africa (Southern)	28 (28)	9. Asia (South)	12.1 (-19.1)
10. Latin America (Tropical)	21.7 (19.8)	10. Asia (Central)	11.2 (11.2)
Total (across all 21 regions)	**951.3 (899.7)**	**Total** (across all 21 regions)	**721.1 (373.9)**

     Supplementary Figures & Tables

573   **Figure S1. Global health regions.** We adopt the same system as the Global Burden of Disease
574   Study in our regional breakdown.

Table S1. Changing year-round (12 month) population at risk patterns for *Aedes aegypti.*
All values are given in millions; future projections are averaged across GCMs, broken down by
year (2050, 2080) and RCP (2.6, 4.5, 6.0, 8.5), and are given as net change from current
population at risk. 0+/0- denote the sign of smaller non-zero values that rounded to 0.0, whereas
"0" denotes true zeros. (Losses do not indicate loss of any transmission, only to reduction 11 or
fewer months.).

Region	Current	2050				2080			
		2.6	4.5	6.0	8.5	2.6	4.5	6.0	8.5
Asia (Central)	0	0	0	0	0	0	0	0	0
Asia (East)	0+	1.4	1.8	1.1	3.7	1.6	4.5	4	8.3
Asia (High Income Pacific)	3.6	-0.2	-0.2	-0.2	-0.2	-0.2	-0.2	-0.2	-0.2
Asia (South)	286.4	21.8	71.8	13.7	73.6	12.1	89.7	72.6	29.6
Asia (Southeast)	499.4	19.2	22.4	19.9	25.1	18.9	26.3	15.4	-10.3
Australasia	0.2	0+	0+	0+	0.1	0+	0.2	0.2	0.3
Caribbean	34.8	1.8	2.2	2.1	2.8	1.7	2.6	2.9	3.3
Europe (Central)	0	0	0	0	0	0	0	0	0
Europe (Eastern)	0	0	0	0	0	0	0	0	0
Europe (Western)	0	0	0	0	0	0	0	0	0+
Latin America (Andean)	14.0	3.9	4.8	4.6	5.7	3.5	5.4	5.8	7.5
Latin America (Central)	88.1	13.0	18.8	17.0	25.8	12.0	24.4	27.4	34.1
Latin America (Southern)	0	0	0	0	0	0	0	0	0.2
Latin America (Tropical)	67.5	27.2	34.5	30.8	41.5	27.3	39	42.9	54.9
North Africa & Middle East	12.5	-5.2	-5.5	-6.0	-5.6	-4.7	-5.4	-5.4	-3.9
North America (High Income)	0.5	0.3	0.9	0.6	1.5	0.3	1.9	1.6	5.5
Oceania	0	0	0	0	0	0	0	0	0
Sub-Saharan Africa (Central)	5.3	0.3	0.6	0.4	0.9	0.3	0.7	0.9	1.7
Sub-Saharan Africa (East)	79.0	19.1	23.1	22.4	26.8	16.6	25.4	28.0	36.1
Sub-Saharan Africa (Southern)	126.9	43.8	60.7	56.7	78.3	37.9	74.3	85.5	110.3
Sub-Saharan Africa (West)	0	0+	0.1	0.1	0.3	0+	0.2	0.6	4.4

Table S2. Changing year-round (12 month) population at risk patterns for *Aedes albopictus*. All values are given in millions; future projections are averaged across GCMs, broken down by year (2050, 2080) and RCP (2.6, 4.5, 6.0, 8.5), and are given as net change from current population at risk. 0+/0- denote the sign of smaller non-zero values that rounded to 0.0, whereas "0" denotes true zeros. (Losses do not indicate loss of any transmission, only to reduction 11 or fewer months).

Region	Current	2050				2080			
		2.6	4.5	6.0	8.5	2.6	4.5	6.0	8.5
Asia (Central)	0	0	0	0	0	0	0	0	0
Asia (East)	1.3	1.4	-0.4	-0.5	-1	1	-0.9	-1	-1.2
Asia (High Income Pacific)	3.6	-0.3	-0.5	-0.4	-2.9	-0.2	-2.2	-3.1	-3.6
Asia (South)	98.3	-73	-80.3	-78.9	-87.3	-67.7	-86.1	-88.5	-92.6
Asia (Southeast)	435.3	-133.9	-213.3	-190.9	-277.4	-131.9	-254.8	-282.7	-343.6
Australasia	0.2	0+	0+	0+	0-	0+	0-	0-	0-
Caribbean	39.3	-5.9	-11.7	-9.4	-17.5	-4.5	-16.1	-18.1	-28.0
Europe (Central)	0	0	0	0	0	0	0	0	0
Europe (Eastern)	0	0	0	0	0	0	0	0	0
Europe (Western)	0	0	0	0	0+	0	0+	0+	0.1
Latin America (Andean)	17.9	2	0.1	0.5	-3	1.8	-2	-3.1	-5
Latin America (Central)	97.2	-23.2	-26.8	-25.3	-31.0	-20.8	-29.3	-31.4	-33.6
Latin America (Southern)	0	0	0	0	0+	0	0	0+	0+
Latin America (Tropical)	93.6	-0.8	-5.2	-6.1	-9.7	-2.9	-10.1	-12.9	-37.1
North Africa & Middle East	2.6	0+	0+	-0.2	-0.2	0.1	0-	-0.2	-0.1
North America (High Income)	1	2.8	1.4	1	0+	1.6	0.1	-0.2	-0.2
Oceania	0	0	0	0	0	0	0	0	0
Sub-Saharan Africa (Central)	5.9	0.4	0.4	0.5	0.1	0.4	0.1	0-	-1.4
Sub-Saharan Africa (East)	96.6	8.0	5.1	6.8	-4.5	7.5	-9.1	-9.9	-45.8
Sub-Saharan Africa (Southern)	133.5	31.9	38.8	38.8	43.4	29.7	39.5	43.9	39.2
Sub-Saharan Africa (West)	0+	0+	0.5	0.4	1.8	0+	0.9	2	6.2

596 **Table S3. Top 10 regional increases in year-round transmission risk (12 months).** All
597 calculations are otherwise the same as in Table 3.

Aedes aegypti		*Aedes albopictus*	
1. Asia (South)	209.9 (29.6)	1. Sub-Saharan Africa (East)	114.3 (39.2)
2. Sub-Saharan Africa (East)	152.6 (110.3)	2. Latin America (Tropical)	39.7 (-37.1)
3. Latin America (Tropical)	63.2 (54.9)	3. Latin America (Central)	38.1 (-33.6)
4. Asia (Southeast)	44 (-10.3)	4. Sub-Saharan Africa (Central)	23.1 (-45.8)
5. Latin America (Central)	40.7 (34.1)	5. Asia (Southeast)	16.3 (-343.6)
6. Sub-Saharan Africa (Central)	36.6 (36.1)	6. Latin America (Andean)	8.6 (-5)
7. Sub-Saharan Africa (West)	8.7 (-130.2)	7. Sub-Saharan Africa (Southern)	6.2 (6.2)
8. Asia (East)	8.3 (8.3)	8. Sub-Saharan Africa (West)	2.5 (-194)
9. Latin America (Andean)	8 (7.5)	9. North Africa & Middle East	2.4 (-0.1)
10. North Africa & Middle East	7.4 (-3.9)	10. Oceania	2 (-1.4)
Total (across all 21 regions)	**597.2 (151.6)**	**Total** (across all 21 regions)	**256.5 (-740.8)**

598