Linear relationship between peak and season-long abundances in insects

Ksenia S. Onufrieva[1] & Alexey V. Onufriev[2,3,4]

[1]Department of Entomology, Virginia Tech, Blacksburg, VA 24061, USA

[2]Department of Computer Science, Virginia Tech, Blacksburg, VA, USA, [3]Department of
Physics, Virginia Tech, Blacksburg, VA, USA, [4]Center for High End Computer Systems,
Virginia Tech, Blacksburg, VA, USA,

*Correspondence: Ksenia Onufrieva, Department of Entomology, 216A Price Hall, Virginia
Tech, Blacksburg, VA 24061-0319, USA. Phone: +1 540- 250-7428; Fax: +1 540- 231-9131
Email: ksenia@vt.edu

1. **Abstract** An accurate quantitative relationship between key characteristics of an insect population, such as season-long and peak abundances, can be very useful in pest management programs. To the best of our knowledge, no such relationship has been established.
2. Here we establish a predictive linear relationship between insect count M_{pw} during the week of peak abundance, the length of seasonal flight period, F (number of weeks) and season-long cumulative abundance $A \approx 0.41 M_{pw}$. The derivation of the equation is based on several general assumptions, and does not involve fitting to experimental data, which implies generality of the result.
3. The equation was tested using extensive data collected on captures of male gypsy moth *Lymantria dispar* (L.) (Lepidoptera: Erebidae), in pheromone-baited traps during 15 years.
4. The model was also tested using trap catch data for two species of mosquitoes, *Culex pipiens* (L.) (Diptera: Culicidae) and *Aedes albopictus* (Skuse) (Diptera: Culicidae), in Gravid and BG-sentinel mosquito traps.
5. The simple, parameter-free equation approximates experimental data points with relative error of 13% and R^2 = 0.997, across all of the species tested.
6. For gypsy moth, we also related season-long and weekly trap catches to the daily trap catches during peak flight.
7. We describe several usage scenarios, in which the derived relationships are employed to help link results of the small-scale field study to the operational Slow the Spread of the gypsy moth program.

Key words: gypsy moth, mosquito, trap catches, flight period, sampling, pest management, mathematical model

Introduction

Surveys are crucial for monitoring insect activity, crop pest levels, local movement, long-range migration, feeding and reproduction, and are widely used in pest management programs. Various tools are used for insect surveys, including several types of traps, which are deployed extensively to detect and monitor insect population levels, presence of invading populations and phenological development, for purposes of both applied pest management and research (Abell et al., 2015; El-Sayed et al., 2006; Elkinton & Cardé, 1981; Liebhold & Tobin, 2008; Poland & McCullough, 2006; Sharov et al., 1995; Thorpe et al., 2006; Tobin et al., 2009; Wall & Perry, 1987; Witzgall et al., 2010; Witzgall et al., 2008). At least 20 million pheromone lures are produced for monitoring and mass trapping annually worldwide (Witzgall et al., 2010). In the United

States, over 100,000 pheromone-baited traps are deployed annually just to monitor gypsy moth (Roberts & Ziegler, 2007).

Insect surveys also play an important role in research on population dynamics, seasonal phenology, mating success and mating disruption in various species, including but not limited to gypsy moth, codling moth, light brown apple moth and oriental fruit moth (Casado et al., 2014; Epstein et al., 2006; Knight et al., 2015; Onufrieva et al., 2013; Onufrieva et al., 2008; Onufrieva et al., 2010; Pfeiffer, 1993; Sharov et al., 2002; Stelinski et al., 2013; Stelinski et al., 2004; Stelinski et al., 2010; Stelinski et al., 2007; Tcheslavskaia et al., 2005a; Tcheslavskaia et al., 2005b; Thorpe et al., 2006; Thorpe et al., 1999; Thorpe et al., 2007). Many of the studies are conducted using releases of laboratory reared insects rather than wild populations, which helps to ensure similar population densities among experimental plots and can provide a longer season of flight for data collection (Casado et al., 2014; Onufrieva et al., 2013; Onufrieva et al., 2008; Onufrieva et al., 2010; Stelinski et al., 2013; Tcheslavskaia et al., 2005a; Tcheslavskaia et al., 2005b). Due to logistics and experimental design, researchers often work with daily or weekly insect counts (Onufrieva et al., 2008; Onufrieva et al., 2010; Pfeiffer, 1993; Sharov et al., 2002; Tcheslavskaia et al., 2005b; Thorpe et al., 2006; Thorpe et al., 1999; Thorpe et al., 1993), while large-scale management programs normally have access to season-long counts only (Tobin & Blackburn, 2007). To the best of our knowledge, no method currently exists to relate daily or weekly counts to season-long counts. Relating daily and weekly counts to season-long counts as well as the converse problem of predicting maximum daily counts from the known season-long ones is important for interpretation of research results and their application in the management programs. Since collecting daily or weekly data in a large-scale management program is very costly and impractical, a model that relates daily and weekly population characteristics to the easy to obtain season-long counts should be of practical benefit.

Materials and Methods

Experimental design. We used data from standard USDA milk carton pheromone-baited traps (Webb, 1982) deployed in 2000, 2001, 2013 and 2014 in Goshen Wildlife Management Area, VA and in 2001, 2002, 2004 and 2006 in Appomattox-Buckingham and Cumberland State Forests, VA. In 2000 – 2006. At each location, 5 to 20 USDA pheromone-baited traps were deployed and checked weekly to monitor the flight period of wild gypsy moth populations as part of ongoing research experiments. Captures from all traps in a given location were averaged and considered a single observation. Since 2013, we began to monitor the traps on a daily basis: in 2013, traps were monitored from June 14 to August 4; in 2014, traps were monitored from June 30 to August 8. In both 2013 and 2014, 2 sentinel traps were deployed and

checked daily for 5 days each week. In 2016, a trap was placed in Blacksburg, VA and
monitored every day from June 20 to August 10.

Season-long trap catch for gypsy moth. Season-long trap counts were highly
variable, showing strong dependence of the season-long cumulative catch (abundance)
A on both the length of the flight period F and the trap catch during the week of peak
flight, M_{pw} (Figure 1).

[Figure 1]

A standard assumption often made by others (Hokyo & Kiritani, 1967; Robinet et
al., 2007; Tobin et al., 2009) is that the distribution of trap captures as a function of time
is Gaussian. We have verified this assumption explicitly for two randomly selected
years (2006 and 2014) of gypsy moth trap catches, (Figure 2).

[Figure 2]

Robustness of the model to insect and trap type. To test robustness of our main
result (Equation 1) to insect and trap type, we used mosquito trapping data collected by
the Mosquito and Forest Pest Management program, Prince William County, VA in
2015. We used *Culex pipiens* catches in gravid traps (Irish et al., 2013) and *Aedes*
albopictus catches in BG-Sentinel traps (Maciel-de-Freitas et al., 2006). The trap data
were collected weekly from May 3 to October 23, 2015.

Daily trap catch for gypsy moth. Daily gypsy moth male trap catches were not
available to us prior to 2013. Once daily trap catches became available in 2013,
followed by 2014 and 2016, we also sought to relate maximum daily trap catch at peak
flight to the season-long trap catch. For this relationship, we used daily trap catches
collected in 2014 as a training set, and daily trap catches collected in 2013 and 2016
served as test sets. Unlike the weekly counts, in which day-to-day fluctuations of insect
counts are automatically averaged out to produce a smooth distribution that deviates
little from the expected Gaussian distribution (Figure 2), individual daily values are too
variable for a single characteristic such as daily average to be useful in practice.
Instead, here we seek to establish a predictive relationship for the range of the daily
counts, that is its conservative upper and lower bounds. It is probably hopeless to try to
derive the range from first principles; instead, we deduce it from our experimental data
for gypsy moth, which we re-interpret as follows.   Since the beginning of peak flight
week assignment is arbitrary and depends on the day a trap was checked, we simulated
different week assignments using a sliding window of 7 days to determine the week of
peak flight, making sure that the maximum observed value of daily catch is always
included in the "peak flight" week. In effect, the procedure simulates 7 different
experimental outcomes. The analysis yielded a range of proportionality coefficients
between trap catch during the peak flight week and the peak daily value.

Results

The model and its derivation. The model describes the relationship between
insect counts M_{pw} during the week of peak activity, the flight period *F*, and season-long
cumulative counts A, which is a direct measure of the insect abundance. It is known that
pheromone traps achieve detection probability of nearly 100% even for relatively sparse
populations (Larsson, 2016). It is therefore reasonable to assume high sensitivity of the
traps. Specifically, we assume that traps are sensitive enough to start catching insects
when their population reaches 1% of the maximum population, consistent with our own
experiments, Figure 2. We also demonstrate below that the specific value of "trap
sensitivity" is not very important for as long as it is high enough – the mathematical
structure of the resulting equation is such that it is insensitive to the value of the
"sensitivity threshold". For a symmetric distribution, the flight period F is defined as
follows: the traps begin to catch insects at t = t_p – F/2, when the insect density rises
above the 1% trap sensitivity threshold, and stop at t = t_p + F/2, once the insect density
drops below this threshold, Figure 2. We model the distribution of insects caught as a
function of time as a Gaussian function $M_{pw}e^{-(t-t_p)^2/2\sigma}$, centered around the peak flight
time point t_p, Figure 2. Then, the cumulative catch (A)

$$A \ = \ \int_{-F/2}^{+F/2} M_{pw}e^{-(t-t_p)^2/2\sigma}dt \ \approx \ M_{pw}\int_{-\infty}^{+\infty} e^{-(t-t_p)^2/2\sigma}dt = \ M_{pw} \times \sigma\sqrt{2\pi}$$

145   The full width *F* of a Gaussian function at 1/100 (1%) of the maximum is related
146 to σ via

$$F = 2\sigma\sqrt{2ln\,100}$$

147   Therefore,

148                $A = \dfrac{2\pi}{2\sqrt{2ln\,100}}M_{pw}F \approx 0.41 M_{pw}F$ (Equation 1)

which is the main result of this work. Its robustness to the assumptions is discussed
below, and is demonstrated experimentally.

We stress that the derivation of Eq. 1 did not involve any fitting to experimental
data. The relationship is expected to hold as long as the following conditions are
satisfied: (1) The seasonal change in the insect population as a function of time (weekly
averages) can be reasonably approximated by a Gaussian function. (2) The trap
method is sensitive enough to detect insects during most of the flight period. (3) Traps
do not saturate. The importance of (1) and (2) for our model is self-evident. To
understand the importance of the last assumption, consider a situation where it breaks

down completely: at very high population density during several weeks, high enough so
the traps are completely saturated. Obviously, if this were the case, the weekly catch
estimates would be flat around the peak, no longer representative of the true insect
abundance, and the model that assumes a Gaussian distribution would fail.

Robustness of the model. Assumptions (2) and (3) essentially state that the catch
method is, in some sense, "well designed". As long as this is the case, Equation 1 is not
very sensitive to details. Indeed, consider a trap that faithfully represents the true
population density in the following sense: catch = α*(true population), where the
constant α stays the same within a given year and experiment, but may vary for different
species and years. Now suppose a different trap is used, which changes the value of α
by a factor of 2 for a different species/year. Both A and M_{pw} in Equation 1 will change
by a factor of 2, without affecting the linear nature of the relationship. There will be a
slight effect on the proportionality constant K in A = K*M_{pw}*F due to the use of more/less
sensitive trap (e.g. one that can sense 1/200 instead of 1/100 of the maximum
population, which will affect the trap sensitivity threshold, Figure 2), but note that the
functional dependence of K on the trap sensitivity threshold is extremely weak,
$K \sim \sqrt{\log\left(1/threshold\right)}$. In the above example of the threshold changing from 1/100 to
1/200, the difference in the proportionality constant in Equation 1 is $\sqrt{2ln100}$ vs.
$\sqrt{2ln\,200}$, or 3.03 vs. 3.25, which is small.

While assumption (1) is likely to hold for many species, it implies that the insects
are short-lived relative to their entire developmental periods. Otherwise, even if the
developmental period itself is normally distributed (which is approximately true even for
humans), the abundance can reach a long plateau during the season, inconsistent with
the Gaussian shape. However, as we shall see below, the simple and parameter-free
model described by Equation 1 works surprisingly well for completely different insect
species and trapping methods in the temperate climate zones. We note that tropical
species may not have a distinct start and end of flight points, and so the model may not
be applicable.

Experimental verification of the model. In our test data set, see "Methods",
season-long male gypsy moth trap catches ranged from 7.3 to 462 males/trap/week;
flight duration ranged from 3.7 to 7 weeks; and maximum trap catch during peak week
ranged from 4.4 to 172 males/trap. Season-long trap catches of *C. pipiens* and *A.
albopictus* ranged from 6.22 to 60.72 and from 0.12 to 31.12, respectively. Flight
duration of both species was 24 weeks. The data strongly confirms the proposed
functional dependence of season-long trap catch on flight duration and trap catch during
peak flight (Figures 3 and 4).

[Figure 3]

The predicted season-long trap catches from trap catches during the week of
peak flight correlate strongly with the actual, measured season-long trap catches for all
three species tested ($R^2 = 0.9977$, SD = 0.13, Figure 4).

It is remarkable that the accuracy of Equation 1 is the same for all of the species
tested, even though the biology, catch methods, and the geographical areas are
different. Moreover, for the mosquito species the catch distributions (not shown) are not
as close to idealized Gaussian as they are for the gypsy moth, Figure 2, although in all
cases there is a clear peak, the beginning and the end of flight points. This observation
suggests strong robustness of the proposed model, and its potential to work for other
species as well.

[Figure 4]

Relationship to daily trap catch for gypsy moth. For reasons discussed in
"Methods", here we seek to relate the range of daily trap catches during the peak flight
week to the other measurable parameters such as the maximum weekly and the total
catch, M_{pw} and A. Experimental daily trap catches ranged from 0 to 13.5 males/trap/day
in 2013, from 0 to 10 males/trap/day in 2014, and from 0 to 15 in 2016. In an idealized
scenario, without random day-to-day variation, trap catches during the peak week of
flight period can be calculated as:

$$M_{pw} = \int_{t_p-3.5}^{t_p+3.5} M_{pd} \times e^{\frac{-(t-t_p)^2}{2\sigma^2}} \approx 7 \times M_{pd}$$

where M_{pd} is the daily peak, that is the absolute maximum of the distribution. In
the idealized case this simple relationship holds because for the species discussed
here, σ is considerably larger than 7 days, and so the integrand varies little over the
integration range in the above expression. However, in reality, daily values fluctuate
significantly around the predicted Gaussian peak, resulting in two effects. First, the
proportionality coefficient K in $M_{pw} = K M_{pd}$ becomes less than 7, with daily peak values
that deviate stronger from the Gaussian-based expectation resulting in lower K. Second,
K will vary from year to year. In 2014, K ranged from 2.35 to 4.66. Below we estimate
maximum and minimum bounds on K, based on the variation in daily trap catches
inferred from our analysis of the experimental data on gypsy moth from 2014, see
"Methods" for details. Our estimate is that $2.35M_{pd} < M_{pw} < 7M_{pd}$, or conversely, $0.14M_{pw}$
$< M_{pd} < 0.43M_{pw}$. We use data points from 2013 and 2016 to validate the estimate
(Figure 5). Clearly, the maximum daily catch during the week of peak flight is within the
estimated lower and upper bounds.

[Figure 5]

Using our main result $A \approx 0.41 M_{pw} F$, we arrive at another useful relationship:

$0.96 M_{pd} F < A < 2.87 M_{pd} F$ (Equation 2)

Discussion

In this paper, we have established a predictive linear relationship between insect
count M_{pw} during the week of peak abundance, the length of seasonal flight period, F
(number of weeks) and season-long cumulative catch $A \approx 0.41 M_{pw} F$. This relationship
was derived without any fitting to experiment, based on very general assumptions,
which likely explains its remarkable accuracy and the fact that it works across the three
tested species with very different biology, behavior, and trapping methods: *Lymantria*
dispar (Lepidoptera: Lymantriidae), *Culex pipiens* (Diptera: Culicidae) and *Aedes*
albopictus (Diptera: Culicidae). Although the model has only been validated using three
species of insects, we expect the relationship, Equation 1, between maximum weekly
counts and season-long counts to hold true for many other species, as long as their
emergence is reasonably approximated by Gaussian distribution, and the collection
method is sensitive and faithfully reflects variation in abundance. The expectation is
based on the generality of the arguments used in the derivation, absence of fitting to a
specific species or trap type, and the already established agreement with experiment
across three species.

Two interrelated scenarios of the model use in practice are exemplified below:

Example Usage Scenario I.

The formula $A \approx 0.41 M_{pw} F$ is used to estimate the hard-to-measure peak
abundance M_{pw} from the other two population parameters, *A* and *F,* which themselves
do not require intense monitoring. Thus, data collected as part of monitoring in the pest
management program can be analyzed using this formula and appropriate method of
control can be chosen based on the maximum daily value. This is important in large-
scale control/monitoring programs, in which a method of control depends on the
population density. For example, currently, in the Slow The Spread of the gypsy moth
program (STS), the low dosage (6 g AI/acre) of mating disruptant is recommended for
use in low-density gypsy moth populations, in which trap catches do not exceed 30
males/trap/season (Thorpe et al., 2006). Ability to directly relate weekly trap catches
from experimental plots to the season-long trap catches used by STS decision algorithm
(Tobin & Sharov, 2007; Tobin et al., 2004) may allow us to improve the method of
mating disruption used against gypsy moth by accurately estimating population
densities during peak flight and using more appropriate control measures.

Example Usage Scenario II

In this situation, the unknown season-long trap catch for a model (artificially
created) population used in a research study is estimated from the peak abundance of
that model population, which in this scenario is easily available by experimental design.
In most cases, researchers are not trying to simulate flight, they simply repeatedly
release similar numbers of insects to assess treatment efficacy. Therefore, the total of
all captured insects in an experimental plot cannot be interpreted as the true season-
long catch, which, therefore, would be unknown. Instead, each trap catch can be treated
as the catch during peak flight and a season-long trap catch can be estimated using the
proposed formula. This approach allows to estimate the population density for which the
treatment efficacy is assessed, to better interpret research results and to appropriately
apply them in the management programs.

The data collected on gypsy moth phenology over the past 16 years allowed us
to conduct a comprehensive analysis and relate season-long and weekly trap catches
and flight duration to the daily trap catches. The data collectively indicate a significant
variability in flight duration, which in turn causes significant variability in peak trap
catches in populations with the same density as measured by male moth catches in
pheromone-baited traps.

To account for this variability, the model provides a range for a daily peak value,
to allow researches and managers to estimate best and worst case scenarios, predict
efficacy of control tactic, and make decisions to ensure optimal results. In addition to the
immediate application in gypsy moth management programs, this model may be utilized
to predict mating success of gypsy moth females and likelihood of persistence of
isolated low-density populations (Contarini et al., 2009; Sharov et al., 1995; Tobin et al.,
2013).

Our main result – the uncovered linear relationship between insect counts during
the week of peak abundance, length of activity period and season-long trap catches –
has already been verified for three very different insect species. The fact that the
relationship holds, quantitatively and accurately, for species with quite different biology,
is indicative of the generality of the relationship. It may therefore not be surprising if
future investigations identify other species for which the same relationship holds true.

Acknowledgements

We thank Slow the Spread of the gypsy moth Program Technical committee for helpful
discussions and support for the research that lead to this analysis; Andrea Hickman,
Samuel Newcomer, and Michael Merz for field assistance; Timothy McGonegal for
providing mosquito trapping data. We also thank Andrew Liebhold (USDA Forest
Service) for his valuable comments on an earlier version of the manuscript.

References

ABELL, K., POLAND, T. M., COSSE, A. & BAUER, L. S. 2015. Trapping techniques for
emerald ash borer and its introduced parasitoids. *In:* VAN DRIESCHE, R. G. &
REARDON, R. C. (eds.) *Biology and control of emerald ash borer. FHTET-2014-*
09. Morgantown, WV U.S. Department of Agriculture, Forest Service, Forest
Health Technology Enterprise Team.
CASADO, D., CAVE, F. & WELTER, S. 2014. Puffer Ū-CM dispensers for mating
disruption of codling moth: Area of influence and impacts on trap finding success
by males. *IOBC-WPRS Bulletin,* 99**,** 25-31.
CONTARINI, M., ONUFRIEVA, K. S., THORPE, K. W., RAFFA, K. F. & TOBIN, P. C.
2009. Mate-finding failure as an important cause of Allee effects along the
leading edge of an invading insect population. *Entomologia Experimentalis et*
Applicata, 133**,** 307-314.
EL-SAYED, A. M., SUCKLING, D. M., WEARING, C. H. & BYERS, J. A. 2006. Potential
of mass trapping for long-term pest management and eradication of invasive
species. *Journal of Economic Entomology,* 99**,** 1550-1564.
ELKINTON, J. S. & CARDÉ, R. T. 1981. The use of pheromone traps to monitor the
distribution and population trends of the gypsy moth. *In:* MITCHELL, E. D. (ed.)
Management of insect pests with semiochemicals. New York: Plenum.
EPSTEIN, D. L., STELINSKI, L. L., REED, T. P., MILLER, J. R. & GUT, L. J. 2006.
Higher Densities of Distributed Pheromone Sources Provide Disruption of
Codling Moth (Lepidoptera: Tortricidae) Superior to That of Lower Densities of
Clumped Sources. *Journal of Economic Entomology,* 99**,** 1327-1333.
HOKYO, N. & KIRITANI, K. 1967. A method for estimating natural survival rate and
mean fecundity of an adult insect population by dissecting the female
reproductive organs. *Researches on Population Ecology,* 9**,** 130-142.
IRISH, S. R., MOORE, S. J., DERUA, Y. A., BRUCE, J. & CAMERON, M. M. 2013.
Evaluation of gravid traps for the collection of Culex quinquefasciatus, a vector of
lymphatic filariasis in Tanzania. *Transactions of The Royal Society of Tropical*
Medicine and Hygiene, 107**,** 15-22.
KNIGHT, A. L., BASOALTO, E. & WITZGALL, P. 2015. Improving the Performance of
the Granulosis Virus of Codling Moth (Lepidoptera: Tortricidae) by Adding the
Yeast Saccharomyces cerevisiae with Sugar. *Environmental Entomology,* 44**,**
252-259.
LARSSON, M. C. 2016. Pheromones and Other Semiochemicals for Monitoring Rare
and Endangered Species. *Journal of Chemical Ecology,* 42**,** 853-868.
LIEBHOLD, A. M. & TOBIN, P. C. 2008. Population ecology of insect invasions and
their management. *Annual Review of Entomology,* 53**,** 387-408.
MACIEL-DE-FREITAS, R., EIRAS, Á. E. & LOURENÇO-DE-OLIVEIRA, R. 2006. Field
evaluation of effectiveness of the BG-Sentinel, a new trap for capturing adult

Aedes aegypti (Diptera: Culicidae). *Memorias do Instituto Oswaldo Cruz,* 101**,** 321-325.

ONUFRIEVA, K., THORPE, K., HICKMAN, A., LEONARD, D., ROBERTS, E. & TOBIN, P. 2013. Persistence of the gypsy moth pheromone, disparlure, in the environment in various climates. *Insects,* 4**,** 104 - 116.

ONUFRIEVA, K. S., BREWSTER, C. C., THORPE, K. W., SHAROV, A. A., LEONARD, D. S., REARDON, R. C., MASTRO, V. C., SELLERS, P. & ROBERTS, E. A. 2008. Effects of the 3M (TM) MEC Sprayable Pheromone (R) formulation on gypsy moth mating success. *Journal of Applied Entomology,* 132**,** 461-468.

ONUFRIEVA, K. S., THORPE, K. W., HICKMAN, A. D., TOBIN, P. C., LEONARD, D. S. & ROBERTS, E. A. 2010. Effects of SPLAT (R) GM sprayable pheromone formulation on gypsy moth mating success. *Entomologia Experimentalis et Applicata,* 136**,** 109-115.

PFEIFFER 1993. Mating Disruption for Control of Damage by Codling Moth in Virginia Apple Orchards. *Entomologia experimentalis et applicata,* 67**,** 57.

POLAND, T. M. & MCCULLOUGH, D. G. 2006. Emerald ash borer: invasion of the urban forest and the threat to North America's ash resource. *Journal of Forestry,* 104**,** 118-124.

ROBERTS , E. A. & ZIEGLER, A. H. 2007. Gypsy Moth Population Monitoring and Data Collection. *In:* TOBIN, P. & BLACKBUM, L. M. (eds.) *Slow the Spread: A National Program to Manage the Gypsy Moth.* Newtown Square, PA: USDA Forest Service Northern Research Station.

ROBINET, C., LIEBHOLD, A. & GRAY, D. 2007. Variation in developmental time affects mating success and Allee effects. *Oikos,* 116**,** 1227-1237.

SHAROV, A. A., LIEBHOLD, A. M. & RAVLIN, F. W. 1995. Prediction of gypsy moth (Lepidoptera: Lymantriidae) mating success from pheromone trap counts. *Environmental Entomology,* 24**,** 1239-1244.

SHAROV, A. A., THORPE, K. W. & TCHESLAVSKAIA, K. 2002. Effect of synthetic pheromone on gypsy moth (Lepidoptera : Lymantriidae) trap catch and mating success beyond treated areas. *Environmental Entomology,* 31**,** 1119-1127.

STELINSKI, L. L., GUT, L. J. & MILLER, J. R. 2013. An Attempt to Increase Efficacy of Moth Mating Disruption by Co-Releasing Pheromones With Kairomones and to Understand Possible Underlying Mechanisms of This Technique. *Environmental Entomology,* 42**,** 158-166.

STELINSKI, L. L., GUT, L. J., PIERZCHALA, A. V. & MILLER, J. R. 2004. Field observations quantifying attraction of four tortricid moths to high-dosage pheromone dispensers in untreated and pheromone-treated orchards. *Entomologia Experimentalis et Applicata,* 117**,** 187-196.

STELINSKI, L. L., MILLER, J. R., LEDEBUHR, R., SIEGERT, P. GUT, L. J. 2007. Season-long mating disruption of Grapholita molesta (Lepidoptera: Tortricidae)

by one machine application of pheromone in wax drops (SPLAT-OFM). *Journal of Pest Science*, 80: 109 – 117.

STELINSKI, L. L., LAPOINTE, S. L. & MEYER, W. L. 2010. Season-long mating disruption of citrus leafminer, Phyllocnistis citrella Stainton, with an emulsified wax formulation of pheromone. *Journal of Applied Entomology,* 134**,** 512-520.

TCHESLAVSKAIA, K., BREWSTER, C., THORPE, K., SHAROV, A., LEONARD, D. & ROBERTS, A. 2005a. Effects of intentional gaps in spray coverage on the efficacy of gypsy moth mating disruption. *Journal of Applied Entomology,* 129**,** 475-480.

TCHESLAVSKAIA, K. S., THORPE, K. W., BREWSTER, C. C., SHAROV, A. A., LEONARD, D. S., REARDON, R. C., MASTRO, V. C., SELLERS, P. & ROBERTS, E. A. 2005b. Optimization of pheromone dosage for gypsy moth mating disruption. *Entomologia Experimentalis et Applicata,* 115**,** 355-361.

THORPE, K., REARDON, R., TCHESLAVSKAIA, K., LEONARD, D. & MASTRO, V. 2006. *A review of the use of mating disruption to manage gypsy moth, Lymantria dispar (L.)*, FHTET-2006–13. U.S. Department of Agriculture, Forest Service, Forest Health Technology Enterprise Team.

THORPE, K. W., MASTRO, V. C., LEONARD, D. S., LEONHARDT, B. A., MCLANE, W., REARDON, R. C. & TALLEY, S. E. 1999. Comparative efficacy of two controlled-release gypsy moth mating disruption formulations. *Entomologia Experimentalis et Applicata,* 90**,** 267-277.

THORPE, K. W., RIDGWAY, R. L. & LEONHARDT, B. A. 1993. Relationship between Gypsy-Moth (Lepidoptera, Lymantriidae) Pheromone Trap Catch and Population-Density - Comparison of Traps Baited with 1 and 500 Mu-G (+)-Disparlure Lures. *Journal of Economic Entomology,* 86**,** 86-92.

THORPE, K. W., TCHESLAVSKAIA, K. S., TOBIN, P. C., BLACKBUM, L. M., LEONARD, D. S. & ROBERTS, E. A. 2007. Persistent effects of aerial applications of disparlure on gypsy moth: trap catch and mating success. *Entomologia Experimentalis et Applicata,* 125**,** 223-229.

TOBIN, P. & SHAROV, A. 2007. The Decision Algorithm: Selection of and Recommendation for Potential Problem Areas. *In:* TOBIN, P. & BLACKBUM, L. M. (eds.) *Slow the Spread: A National Program to Manage the Gypsy Moth.* Newtown Square, PA: USDA Forest Service.

TOBIN, P. C. & BLACKBURN, L. M. 2007. Slow the spread: a national program to manage the gypsy moth. *General Technical Report NRS-6.* Newtown Square, PA: USDA Forest Service

TOBIN, P. C., KLEIN, K. T. & LEONARD, D. S. 2009. Gypsy moth (Lepidoptera: Lymantriidae) flight behavior and phenology based on field-deployed automated pheromone-baited traps. *Environmental Entomology,* 38**,** 1555-1562.

TOBIN, P. C., ONUFRIEVA, K. S. & THORPE, K. W. 2013. The relationship between
male moth density and female mating success in invading populations of
Lymantria dispar. *Entomologia Experimentalis et Applicata,* 146**,** 103-111.
TOBIN, P. C., SHAROV, A. A., LIEBHOLD, A. A. & LEONARD, D. S. 2004.
Management of the gypsy moth through a Decision Algorithm under the STS
Project. *American entomologist (Lanham, Md.),* 50**,** 200-200.
WALL, C. & PERRY, J. 1987. Range of action of moth sex-attractant sources.
Entomologia experimentalis et applicata, 44**,** 5-14.
WEBB, R. E. 1982. Mass trapping of the gypsy moth. *In:* KYDONIEUS, A. F. &
BEROZA, M. (eds.) *Insect suppression with controlled release pheromone*
systems. Boca Raton, Fla.: CRC Press.
WITZGALL, P., KIRSCH, P. & CORK, A. 2010. Sex Pheromones and Their Impact on
Pest Management. *Journal of Chemical Ecology,* 36**,** 80-100.
WITZGALL, P., STELINSKI, L., GUT, L. & THOMSON, D. 2008. Codling moth
management and chemical ecology. *Annual Review of Entomology,* 53**,** 503-522.

Figure 1: Season-long trap catches depend strongly both on trap catches during peak

flight and flight duration. Conversely, trap catches during peak flight cannot be predicted

from the flight period alone.

Figure 2: Weekly male gypsy moth catches in pheromone baited traps, 2006 (A) and

2014 (B) and list-square fit of experimental data points to a normal distribution. In both

cases, the correlation $R^2 > 0.99$. Grey arrows indicate the time points where the insect

population reaches the trap sensitivity threshold.

Figure 3: Season-long trap catches depend linearly on the product of trap catch during

the peak flight week and the length of the flight period. The relationship holds true for

several different insect species and catch methods.

Figure 4: Predicted (solid and checkered black columns for gypsy moth and mosquito

data, respectively) and actual (solid and checkered grey columns for gypsy moth and

mosquito data, respectively) season-long trap catches vs trap catches during peak flight

and flight duration.

Figure 5: Daily gypsy moth male catches in pheromone-baited traps in Virginia, 2013

(A) and 2016 (B). Lines indicate estimated minimum and maximum possible daily

values estimated from the maximum observed weekly peak value.

459   Figure 1

465   Figure 2

469   Figure 3

Trap catch during the peak flight (M_{pw})

473   Figure 4

475   Figure 5