Domain-Specific Patterns of Activity Support Metacognition in Human Prefrontal Cortex

Jorge Morales[1*], Hakwan Lau[2,3], Stephen M. Fleming[4]

[1] Department of Philosophy, Columbia University, New York, NY, USA.
jorge.morales@columbia.edu
[2] Department of Psychology, UCLA, Los Angeles, CA, USA. hlau@psych.ucla.edu
[3] Brain Research Institute, UCLA
[4] Wellcome Trust Centre for Neuroimaging, University College London, London, United Kingdom. stephen.fleming@ucl.ac.uk

* *Corresponding author:* jorge.morales@columbia.edu (J. Morales)

Abstract: Metacognition is the capacity to evaluate the success of one's own cognitive processes in various domains, e.g. memory and perception. It remains controversial whether metacognition relies on a domain-general resource that is applied to different tasks, or whether self-evaluative processes are domain-specific. Here we directly investigated this issue by examining the neural substrates engaged when metacognitive judgments were made during perceptual and memory tasks matched for stimulus and performance characteristics. By comparing patterns of functional magnetic resonance imaging (fMRI) activity while subjects evaluated their performance, we revealed both domain-specific and domain-general metacognitive representations. Multi-voxel activity patterns in right frontopolar cortex predicted levels of confidence in a domain-specific fashion, whereas domain-general signals predicting confidence and accuracy were found in a widespread network in the frontal and posterior midline. Moreover, individual differences in perceptual and memory metacognitive ability were predicted by the degree of confidence decoding in frontal and parietal cortex, respectively. The demonstration of domain-specific metacognitive representations suggests the presence of a content-rich mechanism available to introspection and cognitive control.

Keywords: metacognition; memory; perception; multi-voxel pattern analysis; fMRI; prefrontal cortex

Major subject area: Neuroscience

INTRODUCTION

Metacognition is the capacity to evaluate the success of one's cognitive processes in various domains, e.g. perception or memory (1-4). Metacognitive ability can be assessed in the laboratory by quantifying the trial-by-trial correspondence between objective performance and subjective confidence (5-8). Anatomical (9-11), functional (12-16) and neuropsychological (1-4,17-19) evidence indicates specific neural substrates (especially in frontolateral, frontomedial, and parietal regions) contribute to metacognition across a range of task domains, including memory and perception. However, the neurocognitive architecture supporting metacognition remains controversial. Does metacognition rely on a common, domain-general resource that is recruited to evaluate performance on a variety of tasks? Or is metacognition supported by domain-specific components?

Current computational perspectives (20,21) suggest both domain-general and domain-specific representations may be important for guiding behavior. On the one hand, one needs to be able to compare confidence estimates in a "common currency" across a range of arbitrary decision scenarios (22). One solution to this problem is to maintain a global resource with access to arbitrary sensorimotor mappings (23-25). Candidate neural substrates for a domain-general resource are the frontoparietal and cingulo-opercular networks, known to be involved in arbitrary control operations (23). In particular, the dorsomedial prefrontal cortex (dmPFC; encompassing the paracingulate cortex and pre-supplementary motor area) has been implicated both in representing confidence and detecting errors across a range of tasks (12,26). On the other hand, if the system only had access to generic confidence signals, appropriate switching between particular tasks or strategies on the basis of their expected success would be compromised. Functional imaging evidence implicates human frontopolar cortex in tracking the reliability of specific alternative strategies during decision-making (27), and such regions may also support domain-specific representations of confidence.

Current behavioral evidence of a shared resource for metacognition is ambiguous, in part due to the difficulty of distilling metacognitive processes from those supporting primary task performance (5-7). Some studies have found that efficient metacognition in one task predicts good metacognition in another (9,28,29), whereas others indicate the independence of metacognitive abilities (14,30,31). Recent studies employing bias-free measures of metacognition have identified differences in the neural correlates of memory and perceptual metacognition in both healthy subjects (9,14) and neuropsychological patients (19). However, the study of behavioral individual differences provides only indirect evidence of the neural and computational architecture supporting metacognition.

Here, we directly investigate this ontology by examining neural substrates engaged when metacognitive judgments are made during perceptual and memory tasks matched for stimulus and performance characteristics. We employ a combination of univariate and multivariate analyses of functional magnetic resonance imaging (fMRI) data to identify domain-specific and domain-general neural substrates engaged during metacognitive judgments. We also distinguish

activations engaged by a metacognitive judgment from neural activity which parametrically tracks confidence level. Together, our findings reveal generic and specific confidence representations co-exist, consistent with a computational hierarchy underpinning effective metacognition.

RESULTS

We analyzed the data of 24 subjects who underwent hemodynamic neuroimaging while performing two-alternative forced-choice discrimination tasks in perceptual (PER) and memory (MEM) domains (Figure 1A). In the perceptual task, subjects were asked to indicate the brighter of two stimuli (words or abstract shapes). In the memory task, subjects were asked to memorize exemplars of the same stimulus types, and then select the previously-learned stimulus from two stimuli presented on each trial. By using a 2×2 design (TASK DOMAIN × STIMULUS TYPE), we were able to ensure differences in neural activations across perception and memory tasks were not confounded by stimulus type. In half of the trials (Confidence condition) subjects performed a metacognitive evaluation after the discrimination task by rating their confidence in the correctness of their response by selecting a number from a 1-to-4 scale (1=not confident; 4=very confident). In order to differentiate metacognitive judgment-related activity from visuomotor activity engaged by use of the confidence scale, in the other half of the trials (Follow condition) subjects were asked to respond according to a highlighted number without evaluating confidence in their response.

Behavior

We first compared task performance, measured by percentage of correct responses, across condition, task, and stimulus type. A 2×2×2 repeated measures ANOVA (CONFIDENCE/FOLLOW × PERCEPTION/MEMORY × SHAPES/WORDS) showed that performance was well-matched across rating conditions (Confidence vs Follow) ($F_{1,23}$=3.036, P=0.095; Figure S1). Matching performance across stimulus type was more difficult to achieve because subjects' memory for words was expected to be considerably higher than that for abstract shapes trials based on pilot data (see Methods for details). Instead, we aimed to match subjects' performance for each stimulus type across task domains by titrating the difficulty of the perceptual task to approximate the performance expected for each stimulus type in the memory task (shapes: PER M=73%, MEM M=67%; words: PER M=81%, MEM M=89%; Figure 1B). Critically, this ensured performance was matched across task domains when averaging stimulus types across participants (PER: M=77%, MEM: M=78%; paired t-test T_{23}=0.38, P=0.70; Figure 1C).

Figure 1. Task design & performance results. (A) Subjects performed two-alternative forced-choice discrimination tasks about perception and memory. In perception blocks, subjects selected the brighter of two stimuli. Memory blocks started with an encoding period and then subjects indicated in each trial which of two stimuli appeared during the encoding period. Abstract shapes and words were used as stimuli in both tasks. In Confidence blocks, subjects rated their confidence and in Follow blocks they pressed the highlighted number. (B) Scatter plot of percentage correct responses per block type in the Confidence condition. Each marker represents a subject. (C) Mean percentage correct responses by domain, averaged over subjects and stimulus types. Dotted lines indicate chance performance. Bars indicate standard error of the mean (s.e.m.). See Figure S1 for performance in the Follow condition. P=perception M=memory

Figure 2. Metacognitive measures. (A) Mean number of correct and incorrect trials per confidence rating. **(B)** Metacognitive efficiency measured by log M-ratio (meta-d'/d'). Zero indicates that metacognitive sensitivity (meta-d') is equal to task sensitivity d' (i.e. the d' that would have been predicted to give rise to the observed confidence rating data assuming a signal detection theoretic ideal observer). Group-level hierarchical Bayesian estimates differed significantly between domains. Error bars indicate 95% high-density interval (HDI) from posterior samples. **(C)** Subjects' mean individual hierarchical Bayesian estimates of metacognitive efficiency scores were not correlated across the perceptual and memory domains. **(D)** To estimate the degree of generality of metacognitive efficiency between domains, we calculated a domain-general index (DGI) for each subject that quantifies the similarity between their log M-ratio scores in each domain (see main text for details). Greater DGI scores indicate less metacognitive consistency across domains. Mean log M-ratios are shown for reference. Bars in (A) and (D) indicate s.e.m. *** $P_{\theta>0} \sim 1$. P=perception M=memory.

As expected, subjects gave higher confidence ratings after correct decisions than after incorrect decisions (Figure 2A), and mean confidence ratings were similar across task domains (PER M=2.62, MEM M=2.47; paired t-test T_{23}=1.26, P=0.22). Reaction times for confidence ratings were not different between domains (PER M=518ms, MEM M=516ms; paired t-test T_{23}=0.16, P=0.87). We next estimated log M-ratio, a metacognitive efficiency measure derived from signal detection theory that assays the degree to which confidence ratings distinguish between correct and incorrect trials (5,6,32). Metacognitive efficiency in the perceptual task was significantly lower than in the memory task ($P_{\theta>0} \sim 1$; see Figure 2B & Methods for details), consistent with

117   previous findings (19). We did not find a correlation between metacognitive efficiency scores in
118   the perceptual and memory domains (r_{22}=-0.076; P=0.72; Figure 2C). To further assess a poten-
119   tial covariation between metacognitive abilities in each domain, we calculated for each subject a
120   domain-generality index (DGI) that quantifies the similarity between their scores in each domain
121   (19):
122
123        $$\text{DGI} = |\log M_P - \log M_M|$$
124
where M_P = perceptual M-ratio and M_M = memory M-ratio. Lower DGI scores indicate more
similar metacognitive efficiencies between domains (DGI=0 indicates identical scores). Mean
DGI for shapes (1.42), words (0.66) and collapsed by stimulus type (0.95) were higher than zero
(Figure 2D). Metacognition for words was behaviorally more stable across domains as the DGI
was smaller than for shapes (paired t-test: T_{23}=2.86; P=0.009). Together, these results suggest
domain-specific constraints on metacognitive ability.
131
fMRI analyses
We next turned to our fMRI data to assess the overlap between neural substrates engaged when
metacognitive judgments are made during perceptual and memory tasks. In standard univariate
analyses, we found elevated activity in dACC/pre-SMA and bilateral insulae when comparing
the Confidence against the Follow condition (Figure S2A). We also replicated negative correla-
tions between confidence and activation in dACC/pre-SMA, parietal cortex and PFC that have
been reported in several previous studies (12-14,33) (Figure S2B). However, we reasoned that
while metacognition-related activity may show a consistent univariate profile across tasks (see
Figures S3-S5 for univariate analyses), distributed activity patterns in frontal and parietal areas
may carry task-specific information (33,34). To investigate the domain-specificity of activity
patterns in frontoparietal and cingulo-opercular regions we therefore performed a series of multi-
voxel pattern analyses (MVPA). We focused on two features of metacognition-related activity in
the analysis of functional imaging data - judgment-related (JR) activity (the difference between
Confidence trials requiring a metacognitive judgment and the visuomotor control condition) and
confidence level-related (CLR) activity (the parametric relationship between confidence and ac-
tivity). To separately quantify these effects for each task domain, we focused on four a priori re-
gions of interest (ROI) in dACC/pre-SMA, bilateral rostrolateral prefrontal cortex (rlPFC) and
precuneus (PCUN) (Figure S4), which previous studies have found to be recruited by perceptual
and memory metacognition (9,10,12-14), together with whole-brain searchlight analyses.
151
152
153

Figure 3. MVPA Results. *Judgment Related-Activity* **(A)** ROI results for across-domain (light colors) and mean within-domain (dark colors) classification accuracy of Confidence vs Follow trials. Inset: As a control, classification accuracy in the ventricles was not different from chance (across: $T_{23}=0.66$, $P=0.52$; within: $T_{23}=1.04$, $P=0.31$). **(B)** Searchlight analysis for same classifications as in (A) (within-domain classifications uncollapsed). *Confidence Level Related-Activity* **(C)** Same as in (A) but for low versus high confidence classification accuracy. Inset: Classification accuracy in the ventricles was not different from chance (across: $T_{21}=-0.24$, $P=0.81$; within: $T_{21}=0.86$, $P=0.40$). **(D)** Searchlight analysis for same classifications as in (C), with within-domain classifications uncollapsed and visuomotor activity exclusively masked (see Figure S8 for visuomotor mask and Figure S9 for whole-brain maps of raw within-domain classification accuracy values). Bars in (A) and (C) indicate means and error bars s.e.m. Dashed lines indicate chance classification (50%). See Figures S6-S7 for ROI results split by domain. All clusters in (B) and (D) are significant at cluster-defining threshold $P<0.001$, corrected for multiple comparisons at $P_{FWE}<0.05$. Image displayed at $P<0.001$, uncorrected. Color bars indicate T-scores. See Tables S4-S6. L=left; R=right. *** $P\leq0.001$ ** $P\leq0.01$ * $P<0.05$.

ROI analysis of judgment-related activity. If metacognitive judgments are based on domain-
general processes (i.e. shared across perceptual and memory tasks), a decoder trained to classify
JR-activity in perceptual trials should accurately discriminate JR-activity when tested on
memory trials (and vice versa). Alternatively, domain-specific activity profiles would be indicat-
ed by significant within-domain classification of JR-activity in the absence of across-domain
transfer. To adjudicate between these hypotheses, we performed a support vector machine
(SVM) decoding analysis using as input vectors the run-wise beta images pertaining to Confi-
dence and Follow trials obtained from a general linear model (GLM). For within-domain classi-
fication we used standard leave-one-out cross-validations for each domain and we tested for
across-domain generalization using a cross-classification (see Methods for details). Chance clas-
sification in both analyses was 50%.
Mean within-domain classification results were significantly above chance in all ROIs
(one-sample *t*-tests Bonferroni corrected for multiple comparisons α=0.05/4=0.0125: dACC/pre-
SMA T_{23}=5.77, *P*=6.99×10^{-6}; L rlPFC T_{23}=3.27, *P*=0.003; R rlPFC T_{23}=4.47, *P*=0.0002; PCUN
T_{23}=2.98, *P*=0.007; dark colors in Figure 3A; see Figure S6 for results split by domain). In con-
trast, across-domain generalizations were not significantly different from chance in any ROI (one
sample *t*-test Bonferroni corrected: dACC/pre-SMA, T_{23}=1.95, *P*=0.063; L rlPFC, T_{23}=0.76,
P=0.44; R rlPFC, T_{23}=1.24, *P*=0.228; PCUN T_{23}=1.40, *P*=0.175; light colors in Figure 3A). This
suggests that the patterns of activity that distinguish metacognitive judgments from the visuomo-
tor control condition in one domain are distinct from similar patterns in the other domain. In par-
ticular, within-domain classification accuracy was significantly different from across-domain
classification accuracy (dACC/pre-SMA: T_{23}=2.88, *P*=0.008: right rlPFC: T_{23}=2.24, *P*=0.035).
These results are consistent with the hypothesis that metacognitive judgments recruit domain-
specific patterns of cortical activity in anterior frontal and medial-parietal regions.
196
Searchlight analysis of judgment-related activity. We ran a similar decoding analysis using an
exploratory whole-brain searchlight, obtaining a classification accuracy value per voxel (35).
Consistent with our ROI results, we observed significant within-domain classification in large
swathes of bilateral PFC for both perception (red) and memory (blue) (Figure 3B & Table S4).
Within-perception classification was also successful in parietal regions—precuneus in particu-
lar—and within-memory activity patterns were classified accurately in occipital regions. We also
identified clusters showing significant across-domain generalization (yellow) in dACC, pre-
SMA, SFG (BA 9), supramarginal gyrus (BA 40), and bilateral IFG/insula, consistent with uni-
variate results (Figure S2A).
206
207 *ROI analysis of confidence level-related activity.* We next asked whether confidence is encoded
208 in a domain-general or domain-specific fashion by applying a similar approach to discriminate
209 low versus high confidence trials. Note that in this case, and unlike for judgment-related activity
210 (Figure S5B), ROI univariate analyses did not reveal any differences in confidence-related ac-
211 tivity between domains (Figure S5A). We hypothesized that if confidence level is encoded by

domain-general neural activity patterns, it should be possible to train a decoder to discriminate low (1-2) from high (3-4) confidence rating patterns in the perceptual task and then accurately classify confidence on the memory task (and vice versa). In the absence of across-domain classification, within-domain classification above chance is indicative of confidence level-related domain-specific activity patterns. ROI cross-classifications and cross-validations were performed in a similar fashion as above. One-sample t-tests (Bonferroni-corrected) showed across-domain classification was significantly above chance in dACC/pre-SMA (T_{21}=2.83, P=0.010) and precuneus (T_{21}=4.69, P=9.14×10^{-5}) but not in rlPFC (left: T_{21}=1.36, P=0.19; right: T_{21}=0.97, P=0.34), indicative of a generic confidence signal (Figure 3C, light colors). In contrast, within-domain cross-validation classification accuracy was significantly above chance in right rlPFC (T_{21}=3.74, P=0.001; Bonferroni-corrected). Importantly, classification accuracy in this ROI differed from the corresponding across-domain classification accuracy (paired t-test T_{21}=2.37, P=0.027; Figure 3C, dark colors; see Figure S7 for results split by domain). Together, these results suggest the co-existence of two kinds of CLR neural activity: dACC/pre-SMA and precuneus encode a generic confidence signal, whereas patterns of activity in right rlPFC were modulated by task, allowing within-domain classification of confidence level.

Searchlight analysis of confidence level-related activity. Here we leveraged the Follow trials as a control for low-level visuomotor confounds by exclusively masking out activity patterns associated with usage of the confidence scale (Figure S8 & Table S5). The remaining activity patterns can therefore be ascribed to confidence level-related signals that do not encode visual or motor features of the rating (Figure 3D). We found widespread across-domain classification of confidence level in a predominantly midline network including a large cluster encompassing dACC/pre-SMA, ventromedial PFC (vmPFC), and striatum (yellow). Domain-specific confidence activity was successfully decoded from right PFC (insula, IFG, BA 9, 46) in memory trials (blue) and was also independently decoded in both domains from dACC/pre-SMA.

Generalization of CLR-activity. We reasoned that even in the absence of confidence reports, confidence activity patterns may predict decision accuracy, a proxy of confidence on 'no-report' Follow trials (though we note that the neural correlates of confidence and performance are also likely to be partially distinct; 36,37). We trained a decoder with CLR-beta images on Confidence trials and tested it on beta images indexing accurate/inaccurate decisions on Follow trials as proxies of low/high confidence (and vice versa). This analysis confirmed that activity patterns in dACC/pre-SMA (T_{21}=2.578, P=0.0175) and right rlPFC (T_{21}=2.48, P=0.0215) could be used to predict accuracy levels in Follow trials above chance (Figure S10).

Metacognitive efficiency and confidence level-related activity classification. Finally, we reasoned that if confidence-related patterns of activation are contributing to metacognitive judgments, they should track individual differences in metacognitive efficiency. To test for such a relation, we asked whether individual metacognitive efficiency scores predicted searchlight clas-

sification accuracy of confidence level. Memory metacognitive efficiency predicted memory
confidence classification accuracy in a cluster in right precuneus (cluster defining threshold
P<0.001). This cluster was significant after small-volume correction for multiple comparisons at
P_{FWE}<0.05 using an anatomical mask of bilateral precuneus (Figure 4). Perceptual metacognitive
efficiency predicted perceptual confidence classification accuracy in bilateral vmPFC and cere-
bellum (cluster-defining threshold *P*<0.001, uncorrected). These results are in agreement with
functional (10), anatomical (9) and lesion studies (19) that have linked individual variation in
metacognitive accuracy to precuneus and PFC in memory and perception, respectively.
260

Figure 4. Metacognitive efficiency & confidence classification. Log M-ratio scores predicted
within-domain classification accuracy of confidence level-related activity. Individual metacogni-
tive efficiency in memory trials (blue) predicted classification accuracy in a cluster in right
precuneus and perceptual metacognitive efficiency predicted confidence classification in percep-
tual trials (red) in bilateral ventral frontomedial areas and in the cerebellum. Images displayed at
P<0.001, uncorrected. See main text for statistical analyses. Color bars indicate T-statistic. See
Table S7.
269
270

271 **DISCUSSION**
272

When performing a cognitive task, confidence estimates allow for comparisons of performance
across a range of different scenarios (22). Such estimates must also carry information about the
task context if they are to be used in decision-making. Here we investigated the domain-
generality and domain-specificity of representations that support metacognition of perception
and memory.
Unlike previous studies (9), subjects' performance was matched between domains for
two different types of stimulus, thereby eliminating potential performance and stimulus con-
founds. Subjects' confidence ratings were also matched between domains and followed expected
patterns of higher ratings after correct decisions than after incorrect decisions. Using MVPA, we

showed the existence of both domain-specific and domain-general metacognition-related activity
during perceptual and memory tasks. We report three main findings, and discuss each of these in
turn.

First, we obtained convergent evidence from both univariate and multivariate analyses
that a cingulo-opercular network centered on dACC/pre-SMA encodes a generic signal predic-
tive of confidence level across memory and perceptual tasks. Previous studies of metacognition
have implicated the cingulo-opercular network in tracking confidence level (12,13,15,33,38)
. However, we go beyond these previous studies to provide evidence that these signals generalize
to predict confidence across two distinct cognitive domains. This finding is consistent with pos-
terior medial frontal cortex as a nexus for monitoring the fidelity of generic sensorimotor map-
pings, building on previous findings that error-related event-related potentials originating from
this region are sensitive to variation in subjective certainty (39,40). The activity in this region
was also consistently elevated by the requirement for a metacognitive judgment (12), although
the pattern of these increases did not generalize across tasks.

Second, in anterior frontal regions we found activity patterns that tracked both the re-
quirement for metacognitive judgments and level of confidence. Critically, however, confidence-
related activity patterns were selective for domain in right rlPFC: they differed according to
whether the subject was engaged in a metacognitive judgment about perception or memory. Such
signals may support the "tagging" of confidence with contextual information, thereby facilitating
the use of confidence for behavioral control (27,41). It is possible that anterior prefrontal regions
combine generic confidence signals with domain-specific information to fine-tune decision-
making and action selection in situations in which subjects need to regularly switch between
tasks or strategies on the basis of their reliability (41). An alternative hypothesis, also compatible
with our data, is that PFC first estimates the confidence level specifically for the current task,
which is then relayed to medial areas to recruit the appropriate resources for cognitive control in
a task-independent manner. Processing dynamics may also unfold simultaneously in both areas.
These possibilities echo a longstanding debate in the cognitive control literature on the relative
primacy of medial and lateral PFC in the hierarchy of control (42,43). Further inquiry and devel-
opment of computational models of the hierarchical or parallel functional coupling of these net-
works in metacognition is necessary.

Third, we obtained convergent evidence that precuneus plays a specific role in
metamemory judgments. In univariate fMRI analyses, we found the requirement for a metacog-
nitive judgment recruited precuneus only on memory, but not perceptual, trials (Figure S5B). In-
dividual metacognitive efficiency scores in memory trials predicted classification accuracy in
precuneus (Figure 4), and individual differences in metacognitive efficiency scores in perceptual
trials predicted classification accuracy in ventromedial prefrontal cortex (albeit at an uncorrected
threshold). These findings are consistent with previous results that the medial parietal cortex
makes a disproportional contribution to memory metacognition (9,14), and offer a potential ex-
planation for a decrease in perceptual, but not memory, metacognitive efficiency seen in patients
with frontal lesions (12). However, we do not wish to conclude that precuneus involvement is

specific to metamemory. We note that univariate negative correlations with confidence were
found also on perceptual trials, and multivariate classification results in precuneus indicated the
presence of both perceptual and memory-related signals. This dual involvement of the precuneus
in perception and memory metacognition is consistent with previous studies which suggest a re-
lationship between precuneus structure and visual perceptual metacognition (9,10).

Our experimental design assumes that visual perception and memory are distinct do-
mains. We acknowledge that distinguishing between cognitive domains or individuating percep-
tual modalities is not straightforward (44). For instance, different modalities (e.g. vision, audi-
tion, touch, etc.) could be part of a unified perceptual domain for metacognitive purposes. Recent
findings suggest metacognitive efficiency in one perceptual modality predicts metacognitive ef-
ficiency in others and that they share electrophysiological markers (29). However, it is an open
question whether modality-specific patterns of metacognitive activity could be decoded using
multivariate approaches. More research is needed on the neural architecture of metacognition in
other cognitive domains, and whether this architecture changes in a graded or discrete fashion as
a function of task or stimulus.

In summary, our results provide evidence for the co-existence of content-rich metacogni-
tive representations in frontopolar cortex with generic confidence-related signals in frontoparietal
and cingulo-opercular regions. Such an architecture may be appropriate for "tagging" lower-level
feelings of confidence with higher-order contextual information to allow effective behavioral
control. Our findings reconcile previously conflicting results on the domain-specificity/generality
of metacognition, and lay the groundwork for a mechanistic understanding of reflective judg-
ments of cognition.

344
345
346 **MATERIALS AND METHODS**
347
Participants
Thirty healthy subjects (ages 18-33, mean 24.97 years old; SD=4.44; 14 males) with normal or
corrected-to-normal vision were monetarily compensated and gave written informed consent to
participate in the study at the Center for Neural Science at New York University. The number of
participants was determined a priori at n=30, which is in line with recent guidelines on neuroim-
aging sample sizes (45). The study protocols were approved by the local Institutional Review
Board.
355
356 *Task*
357 Experiments were programmed in MATLAB (MathWorks) using Psychtoolbox (46). Prior to
358 entering the scanner, participants were familiarized with the tasks and the confidence rating
359 scale. During practice trials, subjects got feedback on their responses but did not rate confidence.

The brightness of a randomly located reference stimulus was fixed (mid-gray). The difference in brightness (Δb) between the two stimuli was titrated to calibrate the required brightness of the non-reference stimulus using a staircase procedure similar to previous experiments (10,12,19). We used a fixed large step size 2-down/1-up procedure where subjects practiced until they reached 15 reversals or 90 trials following recommended ratios (47) to target performance levels similar as the expected ones in the memory blocks. Subjects began the experiment with a Δb value determined by the average of the Δb values at each reversal, excluding the first one. Throughout the experiment, we kept a small step size staircase running to account for learning or tiredness. After computing independent thresholds for words and abstract shapes, subjects practiced one whole block (9 trials) in each condition that mimicked experimental blocks (72 trials total). Instructions emphasized that confidence ratings should reflect relative confidence and participants were encouraged to use all ratings.

The experiment consisted of 432 trials of two-alternative forced-choice perception and memory tasks divided into six scanner runs. Task domains were interleaved across runs and counterbalanced across subjects. Runs consisted of four alternations of Confidence and Follow blocks (nine trials each), in which subjects had to either rate their confidence by selecting a number from a 1-to-4 scale, or "follow the computer" by pressing the button corresponding to the highlighted number irrespective of their confidence. The highlighted number was yoked to the previous Confidence block ratings (randomized presentation order) to ensure subjects pressed the same buttons in both conditions during any given pair of blocks. Words and abstract shapes were used as stimuli (interleaved blocks, order counterbalanced across runs). Subjects were reminded at the beginning of each block of the condition, task, and stimulus type that would follow. Subjects used two fingers of their right hand to respond on an MRI-compatible button box: left stimulus (index) and right stimulus (middle). For confidence ratings, they used four fingers: 1 (index), 2 (middle), 3 (ring) and 4 (small). If subjects failed to provide either type of response within the allotted time (see Figure 1A for details), the trial was missed and an exclamation mark was displayed for the remainder of the trial. Failing to press the highlighted number counted as a missed trial. The whole experiment lasted ~1.5 hours.

Stimuli

Stimuli were presented on a black background. Throughout trials, before responses, a mid-gray fixation cross subtending 0.3 degrees of visual angle was presented between stimuli. The reference stimulus in the perceptual task and all stimuli in the memory task were mid-gray. All stimuli were surrounded by an isoluminant blue bounding box separated from the stimulus by a gap of at least 0.15 degrees of visual angle.

22- or 28-line abstract shapes were randomly created by specifying an (invisible) grid of 6×6 squares that subtended 4 degrees of visual angle where lines could connect two vertices horizontally, vertically, or diagonally. The first line always stemmed from the central vertex of the invisible grid randomly connecting one of the surrounding eight vertices to ensure shape centrali-

ty within the grid. The remaining lines were drawn sequentially, ensuring all lines were connect-
ed. Orientation and originating vertices were selected randomly.

All words were nouns of 6 to 12 letters with 1 to 4 syllables obtained from the Medical
Research Council Psycholinguistic Database (48). In the perceptual task, words had high famili-
arity, concreteness, and imageability ratings (400-700). In the memory task, words had low rat-
ings (100-400) to increase task difficulty. Each word and each shape was presented once
throughout the experiment (across perceptual and memory blocks, including practice trials). All
subjects were tested on the same words and shapes (counterbalanced across Confidence and Fol-
low conditions across subjects). Words and rating scales were presented using DS-Digital font
(40 points) to make their visual features similar to the abstract shapes.

To obtain stimulus sets of similar difficulty for shapes and words, we ran a series of pilot
studies where participants rated abstract shapes' distinctiveness and then performed both tasks
(171 Amazon Mechanical Turk participants [73 for shapes; 98 for words] and 6 subjects in the
laboratory). Based on these results, we expected a mean performance of ~71% correct responses
when 22- and 28-line distinctive shapes were used in the same block, and ~83% correct when
long words (6-12 letters) with low concreteness, imageability and familiarity ratings (100-400)
were used. To further increase difficulty, we created pairs of old and new words split between
Confidence and Follow conditions (counterbalanced across subjects), blocked by similar seman-
tic category (e.g. finance, argumentation, character traits, etc.), such that each new word within a
block was freely associated with one old word (and when possible, vice versa) according to the
University of South Florida free association normed database (49).

Behavioral data analysis

Data analysis was performed in MATLAB and statistical analysis in RStudio (50). We
estimated metacognitive efficiency by computing log M-ratio (meta-d'/d'). d' is a signal detection
theoretic measure of type 1 sensitivity, while meta-d' is a measure of type 2 sensitivity (i.e. the
degree to which a subject discriminates correct form incorrect responses) expressed in the same
units as type 1 sensitivity (d') (5,6). We used hierarchical Bayesian estimation to incorporate sub-
ject-level uncertainty in group-level parameter estimates (32). Certainty on this parameter was
determined by computing the 95% high-density interval (HDI) from the posterior samples (51).
Two subjects were discarded for missing more than 10% of the trials (i.e. >1 standard deviation
from the average missed trials, which was 5%). Missed trials were not analyzed.

Multi-voxel pattern analysis

MVPAs were carried out in MATLAB using The Decoding Toolbox (35). We classified beta
images from GLMs modeling JR- and CLR-activity in ROI and whole-brain searchlight analyses
(see Supplemental Methods for fMRI data acquisition and preprocessing). ROI MVPAs were
performed on normalized, smoothed images. Previous work has shown that these preprocessing
steps have minimal impact on SVM classification accuracy, while allowing meaningful compari-

son across subject-specific differences in anatomy, as in standard fMRI analyses (52,53). A single accuracy value per subject, per condition was extracted and used for group analysis.

Searchlight analyses used 12mm-radius spheres centered around a given voxel, for all voxels, on spatially realigned and slice-timing corrected images from each subject to create whole-brain accuracy maps. For group-level analyses, these individual searchlight maps were spatially normalized and smoothed using a Gaussian kernel (8mm FWHM) and entered into one-sample *t*-tests against chance accuracy (33,35). Whole-brain cluster inference was carried out in the same manner as in univariate analysis (see Supplementary Methods).

Prior to decoding, for JR-activity classification we modeled the rating periods in the Confidence and Follow conditions in two regressors of interest per run. For CLR-activity, we collapsed ratings 1 and 2 into a low confidence regressor and ratings 3 and 4 into a high confidence regressor to allow binary classification. Despite being encouraged to use the four levels of confidence in every run, two subjects did not provide ratings for one of the classes in at least one run and were left out from the analysis to avoid entering unbalanced training data into the classifier. The main result of the ROI analysis (Figure 3C) is not affected when the unbalanced data of these two subjects are included: within-domain classification accuracy in right rlPFC (M=62.22%) is significantly above chance (T_{23}= 4.22; *P*=0.0003) and different from across-domain classification accuracy (M=52.84%; paired *t*-test: T_{23}= 2.54; *P*=0.018). The remaining parameters were specified as in the univariate case (see Supplementary Methods).

In independent across-domain classifications, we used the run-wise beta images reflecting JR- and CLR-activity as pattern vectors in a linear support vector classification model (as implemented in LIBSVM). We assigned each vector from each domain a label corresponding to Confidence (1) and Follow (-1) in the JR-analysis and Low Confidence (-1) and High Confidence (1) in CLR-analysis. We trained a support vector machine (SVM) with the vectors from one domain (3 per class, 6 in total) and tested the decoder on the 6 vectors from the other domain (and vice versa). We report the average classification accuracies of these two-way cross-classifications.

For within-domain classifications, we ran independent leave-one-run-out cross-validations for each domain on JR-activity (Confidence vs Follow) and CLR-activity (Low vs High confidence). The pattern vectors from two of the three runs in each domain were used to train an SVM to predict the same classes in the vectors from the left-out run. We compared the true labels of the left-out run with the labels predicted by the model and iterated this process for the other two runs to calculate a mean cross-validated accuracy independently for each domain.

Metacognitive scores (log M-ratio) were estimated independently for perceptual and memory blocks as well as a single score collapsing both domains. These scores were inserted as a covariate in second level analyses for within- and across-domain classifications of confidence level-related activity. The covariate contrast was then obtained for each domain map and for the across-domain results. An anatomical map of the precuneal region was extracted from the Harvard-Oxford probabilistic atlas of human cortical brain areas, and used for performing a small-volume correction for the within-memory results.

478 **ACKNOWLEDGEMENTS**
479 We thank Aurelio Cortese for helping with the MVPA design. This work was partially supported
480 by the National Institute of Neurological Disorders and Stroke of the National Institutes of
481 Health (Grant No. R01NS088628) to H.L. and S.M.F., by the Templeton Foundation (Grant No.
482 21569) to H.L., and by a Sir Henry Wellcome Fellowship to S.M.F. (Grant No. WT096185).
483

484 **REFERENCES**
485

1. Metcalfe J, Shimamura AP, editors. Metacognition. Cambridge, MA: MIT Press; 1994.
2. Flavell JH. Metacognition and cognitive monitoring: A new area of cognitive–
developmental inquiry. American Psychologist. American Psychological Association;
1979;34(10):906–11.
3. Nelson TO, Narens L. Metamemory: A Theoretical Framework and New Findings. In:
Bower G, editor. The Psychology of Learning and Motivation. New York: Academic
Press; 1990. pp. 125–73.
4. Fleming SM, Dolan RJ, Frith CD. Metacognition: computation, biology and function. Phil
Trans R Soc B. The Royal Society; 2012;367(1594):1280–6.
5. Maniscalco B, Lau H. A signal detection theoretic approach for estimating metacognitive
sensitivity from confidence ratings. Conscious Cogn. 2012;21:422–30.
6. Fleming SM, Lau H. How to measure metacognition. Front Hum Neurosci. 2014 Jul;8.
7. Galvin SJ, Podd JV, Drga V, Whitmore J. Type 2 tasks in the theory of signal detectabil-
ity: Discrimination between correct and incorrect decisions. Psychon Bull Rev. Springer;
2003;10(4):843–76.
8. Overgaard M, Sandberg K. Kinds of access: different methods for report reveal different
kinds of metacognitive access. Phil Trans R Soc B. 2012 Apr 9;367(1594):1287–96.
9. McCurdy LY, Maniscalco B, Metcalfe J, Liu KY, de Lange FP, Lau H. Anatomical Cou-
pling between Distinct Metacognitive Systems for Memory and Visual Perception. J Neu-
rosci. 2013 Jan 30;33(5):1897–906.
10. Fleming SM, Weil RS, Nagy Z, Dolan RJ, Rees G. Relating Introspective Accuracy to
Individual Differences in Brain Structure. Science. 2010 Sep 16;329(5998):1541–3.
11. Maniscalco B, McCurdy LY, Odegaard B, Lau H. Limited Cognitive Resources Explain a
Trade-Off between Perceptual and Metacognitive Vigilance. J Neurosci. 2017 Feb
1;37(5):1213–24.
12. Fleming SM, Huijgen J, Dolan RJ. Prefrontal Contributions to Metacognition in Perceptu-
al Decision Making. J Neurosci. 2012 May 2;32(18):6117–25.
13. Fleck MS, Daselaar SM, Dobbins IG, Cabeza R. Role of Prefrontal and Anterior Cingu-
late Regions in Decision-Making Processes Shared by Memory and Nonmemory Tasks.
Cereb Cortex. 2006;16(11):1623–30.
14. Baird B, Smallwood J, Gorgolewski KJ, Margulies DS. Medial and Lateral Networks in
Anterior Prefrontal Cortex Support Metacognitive Ability for Memory and Perception. J

518   Neurosci. 2013 Oct 16;33(42):16657–65.

519   15.   Hilgenstock R, Weiss T, Witte OW. You'd Better Think Twice: Post-Decision Perceptual
520   Confidence. Neuroimage. 2014 Oct 1;99:323–31.

521   16.   Yokoyama O, Miura N, Watanabe J, Takemoto A, Uchida S, Sugiura M, et al. Right fron-
522   topolar cortex activity correlates with reliability of retrospective rating of confidence in
523   short-term recognition memory performance. Neurosci Res. 2010 Nov;68(3):199–206.

524   17.   Shimamura AP, Squire LR. Memory and metamemory: a study of the feeling-of-knowing
525   phenomenon in amnesic patients. J Exp Psychol Learn. 1986 Jul;12(3):452–60.

526   18.   Schnyer DM, Verfaellie M, Alexander MP, LaFleche G, Nicholls L, Kaszniak AW. A role
527   for right medial prefrontal cortex in accurate feeling-of-knowing judgments: evidence
528   from patients with lesions to frontal cortex. Neuropsychologia. 2004 Jan;42(7):957–66.

529   19.   Fleming SM, Ryu J, Golfinos JG, Blackmon KE. Domain-specific impairment in meta-
530   cognitive accuracy following anterior prefrontal lesions. Brain. 2014 Aug
531   6;137(10):2811–22.

532   20.   Pouget A, Drugowitsch J, Kepecs A. Confidence and certainty: distinct probabilistic quan-
533   tities for different goals. Nature Neurosci. 2016 Feb 23;19(3):366–74.

534   21.   Fleming SM, Daw ND. Self-evaluation of decision-making: A general Bayesian frame-
535   work for metacognitive computation. Psychol Rev. 2017;124(1):91–114.

536   22.   de Gardelle V, Mamassian P. Does Confidence Use a Common Currency Across Two
537   Visual Tasks? Psychol Sci. 2014 Apr 3.

538   23.   Cole MW, Reynolds JR, Power JD, Repovs G, Anticevic A, Braver TS. Multi-task con-
539   nectivity reveals flexible hubs for adaptive task control. Nature Neurosci. Nature Re-
540   search; 2013 Sep 1;16(9):1348–55.

541   24.   Heekeren HR, Marrett S, Ruff DA, Bandettini PA, Ungerleider LG. Involvement of hu-
542   man left dorsolateral prefrontal cortex in perceptual decision making is independent of re-
543   sponse modality. Proc Natl Acad Sci USA. 2006 Jun 27;103(26):10023–8.

544   25.   Holroyd CB, Yeung N, Coles MGH, Cohen JD. A Mechanism for Error Detection in
545   Speeded Response Time Tasks. J Exp Psychol Gen. 2005;134(2):163–91.

546   26.   Gehring WJ, Goss B, Coles MGH, Meyer DE, Donchin E. A Neural System for Error De-
547   tection and Compensation. Psychol Sci. Sage Publications, Inc; 1993 Nov 1;4(6):385–90.

548   27.   Donoso M, Collins AGE, Koechlin E. Foundations of human reasoning in the prefrontal
549   cortex. Science. 2014 Jun 27;344(6191):1481–6.

550   28.   Ais J, Zylberberg A, Barttfeld P, Sigman M. Individual consistency in the accuracy and
551   distribution of confidence judgments. Cognition. 2016 Jan;146:377–86.

552   29.   Faivre N, Filevich E, Solovey G, Kuhn S, Blanke O. Behavioural, modeling, and electro-
553   physiological evidence for domain-generality in human metacognition. bioRxiv. 2016.

554   30.   Kelemen WL, Frost PJ, Weaver CA. Individual differences in metacognition: evidence
555   against a general metacognitive ability. Mem Cognit. 2000 Jan;28(1):92–107.

556   31.   Vo VA, Li R, Kornell N, Pouget A, Cantlon JF. Young Children Bet on Their Numerical
557   Skills: Metacognition in the Numerical Domain. Psychol Sci. 2014 Sep 10;25(9):1712–21.

32.   Fleming SM. Hierarchical Bayesian estimation of metacognitive efficiency from confi-
dence ratings. Neurosci Consc. 2017 Apr 22;3(1).

33.   Hebart MN, Schriever Y, Donner TH, Haynes JD. The Relationship between Perceptual
Decision Variables and Confidence in the Human Brain. Cereb Cortex. 2014;26(1):118–
30.

34.   Cole MW, Ito T, Braver TS. The Behavioral Relevance of Task Information in Human
Prefrontal Cortex. Cereb Cortex. 2016 May 17;26(6):2497–505.

35.   Hebart MN, Görgen K, Haynes JD. The Decoding Toolbox (TDT): a versatile software
package for multivariate analyses of functional imaging data. Front Neuroinform. 2015
Jan;8.

36.   Rounis E, Maniscalco B, Rothwell JC, Passingham RE, Lau H. Theta-burst transcranial
magnetic stimulation to the prefrontal cortex impairs metacognitive visual awareness.
Cogn Neurosci. 2010 Aug 18;1(3):165–75.

37.   Cortese A, Amano K, Koizumi A, Kawato M, Lau H. Multivoxel neurofeedback selec-
tively modulates confidence without changing perceptual performance. Nat Commun.
2016 Dec 7;7.

38.   De Martino B, Fleming SM, Garrett N, Dolan RJ. Confidence in value-based choice. Na-
ture Neurosci. 2013;16:105–10.

39.   Boldt A, Yeung N. Shared Neural Markers of Decision Confidence and Error Detection. J
Neurosci. 2015 Feb 25;35(8):3478–84.

40.   Scheffers MK, Coles M. Performance monitoring in a confusing world: error-related brain
activity, judgments of response accuracy, and types of errors. J Exp Psychol Human.
2000;26(1):141–51.

41.   Purcell BA, Kiani R. Hierarchical decision processes that operate over distinct timescales
underlie choice and changes in strategy. Proc Natl Acad Sci USA. 2016 Aug
2;113(31):E4531–40.

42.   Tang H, Yu H-Y, Chou C-C, Crone NE, Madsen JR, Anderson WS, et al. Cascade of neu-
ral processing orchestrates cognitive control in human frontal cortex. eLife. 2016 Feb
18;5.

43.   Kerns JG, Cohen JD, Stenger VA, Carter CS. Prefrontal cortex guides context-appropriate
responding during language production. Neuron. 2004 Jul 22;43(2):283–91.

44.   Macpherson F. Introduction: Individuating the Senses. In: Macpherson F, editor. The
Senses. Oxford: Oxford University Press; 2011. pp. 3–42.

45.   Poldrack RA, Baker CI, Durnez J, Gorgolewski KJ, Matthews PM, Munafò MR, et al.
Scanning the horizon: towards transparent and reproducible neuroimaging research. Nat
Rev Neurosci. 2017 Feb;18(2):115–26.

46.   Brainard DH. The Psychophysics Toolbox. Spat Vis. 1997 Jan 1;10(4):433–6.

47.   García-Pérez MA. Forced-choice staircases with fixed step sizes: asymptotic and small-
sample properties. Vision Res. 1998 Jun;38(12):1861–81.

48.   Wilson M. MRC Psycholinguistic Database: Machine-usable dictionary, version 2.00. Be-

hav Res Methods Instrum Comput. 1988;20(1):6–10.

49. Nelson DL, McEvoy CL, Schreiber TA. The University of South Florida free association,
rhyme, and word fragment norms. Behav Res Methods Instrum Comput. 2004
Aug;36(3):402–7.

50. R Studio Team. RStudio: Integrated Development for R. Boston, MA: RStudio, Inc; 2015.
Available from: http://www.rstudio.com

51. Kruschke J. Doing Bayesian Data Analysis. Academic Press; 2010.

52. Kamitani Y, Sawahata Y. Spatial smoothing hurts localization but not information: Pitfalls
for brain mappers. Neuroimage. 2010 Feb 1;49(3):1949–52.

53. Op de Beeck HP. Against hyperacuity in brain reading: spatial smoothing does not hurt
multivariate fMRI analyses? Neuroimage. 2010 Feb 1;49(3):1943–8.

Supplementary Information

Results from univariate analysis of fMRI data

Judgment-related activity. In a Confidence>Follow contrast (Figure S1A), we found significant activations across domains in dorsal anterior cingulate cortex (dACC; BA 24 & 32), pre-supplementary motor area (pre-SMA), bilateral insulae, superior and middle frontal gyri (SFG & MFG), and lingual gyrus, consistent with previous findings (1).

Confidence level-related activity. We found positive correlations between confidence level and activation in left primary motor cortex and SMA, middle and anterior cingulate cortex (BA 11, 25), as well as in the left hippocampus, and right cerebellum and lingual gyrus (Figure S1B, hot colors). Negative correlations with confidence were found in dACC/ pre-SMA, bilateral middle and inferior frontal gyrus (BA 9, 46), bilateral parietal cortex (BA 40), and left cerebellum and lingual gyrus (Figure S1B, cool colors). Similar negative correlations between confidence and activation in dACC/pre-SMA, parietal cortex and PFC have been reported in several previous studies (1-4).

Confidence level-related activity by domain. We further assessed these data splitting them by domain. Positive correlations with confidence in perception and memory trials were shared in pre- and post-central gyri, consistent with subjects using different fingers from their right hand to rate confidence (green in Figure S2A). Shared negative correlations with confidence were found in right dorsal PFC (green in Figure S2B). In perceptual trials, a negative correlation with confidence was significant in dACC/pre-SMA, bilateral parietal regions and left inferior and middle frontal gyri (BA9, 10, 46), whereas in memory trials the same negative correlation was found in lingual gyrus and right superior frontal gyrus (BA6).

ROI analysis. To separately quantify these effects for each task domain, we focused on four a priori regions of interest (ROI) in dACC/pre-SMA, bilateral rostrolateral prefrontal cortex (rlPFC) and precuneus (PCUN) (Figure S3A), which previous studies have found to be recruited by perceptual and memory metacognition (1,2,4-6). An analysis of confidence level-related activity recapitulated the whole-brain results, with negative relationships between confidence and activity in dACC/pre-SMA. Importantly, no significant differences in beta values were found between domains in any of the ROIs (all paired *t*-tests: $T_{23}<1.6$; $P>0.05$; Figure 4). Unlike the whole-brain results by domain, the ROI analysis was suggestive of a lack of domain-specificity in confidence level-related activity. This ambiguity suggests a potential insensitivity of univariate analyses to putative domain-specific differences in confidence encoding.

With respect to judgment-related activity, in a series of repeated measures 2×2 ANOVAS (CONDITION × TASK) we found greater activity on Confidence compared to Follow trials in both tasks in all ROIs except precuneus, where this increase was observed only for memory trials

678 (Figure S3B). These results are compatible with previous findings indicating a distinctive role of
679 precuneus during memory metacognition (4,6).
680
681 **Supplementary Methods**
682
fMRI data acquisition
Brain images were acquired using a 3T Allegra scanner (Siemens). BOLD-sensitive functional
images were acquired using a T2*-weighted gradient-echo echo-planar images (42 transverse
slices, interleaved acquisition; TR, 2.34s; TE, 30ms; matrix size: 64×64; 3×3mm in-plane resolu-
tion; slice thickness: 3mm; flip angle: 90°; FOV: 126mm). The main experiment consisted of
three runs of 210 volumes and three runs of 296 volumes for the perceptual and memory tasks,
respectively. We collected a T1-weighted MPRAGE anatomical scan (1×1×1mm voxels; 176
slices) and local field maps for each subject.
691
fMRI data preprocessing
Imaging analysis was carried out using SPM12 (Statistical Parametric Mapping;
www.fil.ion.ucl.ac.uk/spm). The first five volumes of each run were discarded to allow for T1
stabilization. Functional images were realigned and unwarped using local field maps (7) and then
slice-timing corrected (8). Each participant's structural image was segmented into gray matter,
white matter, cerebral spinal fluid, bone, soft tissue, and air/background images using a nonlinear
deformation field to map it onto template tissue probability maps (9). This mapping was applied
to both structural and functional images to create normalized images to Montreal Neurological
Institute (MNI) space. Normalized images were spatially smoothed using a Gaussian kernel
(8mm FWHM). We set a within-run 1mm rotation and 4mm affine motion cut-off criterion,
which led to the exclusion of 4 subjects, leaving a total of 24 subjects whose functional and be-
havioral data were fully analyzed.
Univariate analysis
All our GLMs focus on the rating period (from initial response to confidence rating or number
following) and motion correction parameters were entered as covariates of no interest along with
a constant term per run. Regressors were convolved with a canonical hemodynamic response
function (HRF). Low-frequency drifts were excluded with a 1/128Hz high-pass filter. Missed
trials were not modeled. For JR-analyses, we created a GLM with two regressors of interest per
run to estimate BOLD response amplitudes in each voxel during the rating period in each trial of
the Confidence and Follow blocks. For the CLR-parametric modulation analysis, a GLM was
used to estimate BOLD responses in Confidence blocks. There were two regressors of interest in
each run, one modeling the confidence rating period and another that encoded a parametric mod-
ulation by confidence rating.
715
Statistical inference

For the JR-analysis, single-subject contrast images of the Confidence and Follow regressors were
entered into a second-level random effects analysis using one-sample *t*-tests against zero to as-
sess group-level significance. For the CLR-parametric modulation analysis, single-subject con-
trast images of the parametric modulator were entered into a similar second-level random effects
analysis. Activations were visualized using Surf Ice (https://www.nitrc.org/projects/surface/) and
MRIcro (http://www.mccauslandcenter.sc.edu/crnl/mricro).
723
ROI analysis
To define regions of interest, 12mm spheres were centered at MNI coordinates identified from
previous literature (Figure S4). Beta values were extracted from subjects' Confidence>Follow
contrast and confidence parametric modulator contrast images for the JR and CLR analyses, re-
spectively.

Supplementary Figures

730
Figure S1. Task performance (percentage correct) was matched across Confidence and Follow
conditions for each combination of task and stimulus type. In a 2×2×2 repeated measures ANO-
VA (CONFIDENCE/FOLLOW × PERCEPTION/MEMORY × SHAPES/WORDS), there was no three-way
interaction ($F_{1,23}=2.573$, $P=0.12$) and no effect of rating condition ($F_{1,23}=3.036$, $P=0.095$). None
of the four paired t-tests comparing performance between the Confidence and Follow conditions
returned a significant difference ($P>0.05$).

A Judgment-related activity

B Confidence-related activity

Figure S2. fMRI univariate analysis results. (A) Activity from the Confidence>Follow contrast. **(B)** Activity parametrically modulated by level of reported confidence. Hot colors indicate a positive correlation with confidence and cool colors a negative correlation. All displayed activations are significant at cluster-defining threshold $P<0.001$, corrected for multiple comparisons at $P_{FWE}<0.05$. Images displayed at $P<0.001$, uncorrected. Color bars reflect T-statistic (see Tables S1 & S2). L=left; R=right.

A **Positively correlated confidence level-related activity**

z = 48   y = -19   x = -6

3.5 ███████ 6.4   3.5 ███████ 8.0   ███
perception   memory   conjunction

B **Negatively correlated confidence level-related activity**

z = 41   y = -85   x = 50

-3.5 ███████ -7.3   -3.5 ███████ -5.6   ███
perception   memory   conjunction

Figure S3. Confidence level-related fMRI univariate analysis by domain. (A) Activity positively correlated by level of reported confidence in perceptual (red) and memory (trials). Green areas indicate significant voxels in both domains. Common regions were active only in left sensorimotor areas. **(B)** Activity negatively correlated by level of reported confidence; colors indicate same as in (A). Areas common to both domains were found in right dorsolateral PFC. Confidence was anti-correlated in perception with bilateral parietal regions, dACC/pre-SMA, and left dorsolateral PFC; in memory, left lingual gyrus was active. All displayed activations are significant at cluster-defining threshold $P<0.001$, corrected for multiple comparisons $P_{FWE}<0.05$. Images displayed at $P<0.001$, uncorrected. Graded color bars reflect T-statistic (see Table S3).

763
764
Figure S4. Regions of Interest. We created 12mm-radius spherical binary masks for four re-
gions of interest (ROI) based on previously reported peak MNI coordinates. ROIs in left ros-
trolateral prefrontal cortex (L rlPFC) [-33, 44, 28], right rlPFC (R rlPFC) [27, 53, 25] and dorsal
anterior cingulate cortex/pre-supplementary motor area (dACC/pre-SMA) [0, 17, 46] were creat-
ed based on (1). The mask for precuneus (PCUN) [0, -64, 24] was based on (6). The MNI x-
coordinates for the dACC/pre-SMA and PCUN masks were set to 0 to ensure bilaterality. As a
control, we also created a 6mm-radius sphere centered at the ventricles [0 2 15].

A **Confidence level -related activity**

B **Judgment-related activity**

Figure S5. Univariate ROI analysis results. (A) Estimated mean beta values of confidence lev-
el-related activity. No significant difference between domains was found in any ROI. **(B)** Esti-
mated mean beta values of judgment-related activity. Repeated measures 2×2 ANOVA (CONDI-
TION × TASK) on each ROI revealed that in dACC/pre-SMA there was a greater activity on Con-
fidence compared to Follow trials ($F_{1,23}$=19.34, P=0.0002) and this difference was greatest on
memory trials (interaction: $F_{1,23}$=12.16, P=0.0019; post hoc paired t-test, T_{23}=5.472, P=0.0001).
In frontopolar cortex there was a consistently heightened activity on Confidence compared to
Follow trials in both left rlPFC ($F_{1,23}$=6.62, P=0.017) and right rlPFC ($F_{1,23}$=9.282, P=0.0057).
Finally, in precuneus, activity was overall greater on memory compared to perception trials
($F_{1,23}$=36.78, P=0.000003), and differentiated between Confidence and Follow trials in the
memory task (interaction: $F_{1,23}$=15.86, P=0.0006; paired t-test, T_{23}=2.431, P=0.023). Error bars
indicate s.e.m. *** P<0.001 ** P<0.01 *P<0.05

Figure S6. Judgment-related MVPA ROI results split by domain. ROI results for across-
domain (left bars; same as in Figure 5A and presented here for comparison) and mean within-
perception (middle bars) and within-memory (right bars) classification accuracy of judgement-
related activity. In dACC/pre-SMA both within-perception (T_{23}=4.66, P=0.0001) and within-
memory (T_{23}=4.28; P=0.0003) classifications were different from chance, and each was signifi-
cantly different from across-domain classification accuracy (paired t-tests perception: P=0.0261;
memory: P=0.026). In left rlPFC, classification accuracy within-perception was different from
chance (T_{23}=3.29; P=0.003) and from across-domain classification accuracy (paired t-test:
P=0.043). In right rlPFC within-perception (T_{23}=3.19; P=0.004) and within-memory (T_{23}=3.32;
P= 0.003) classifications were significantly above chance. All reported t-tests against chance are
significant after Bonferroni correction (α=0.05/4=0.0125). Inset: Classification accuracy in the
ventricles, tested as a control, was not different from chance (perception: P=0.72; memory:
P=0.34). *** P≤0.001 ** P≤0.01 * P<0.05

Figure S7. Confidence level-related MVPA ROI results split by domain. ROI results for across-domain (left bars; same as in Figure 5C and presented here for comparison) and mean within-perception (middle bars) and within-memory (right bars) classification accuracy of confidence level-related activity. In right rlPFC, within-memory ($T_{21}=3.47$; $P=0.002$) classification accuracy was significantly above chance (Bonferroni corrected). It was also different from across-domain classification accuracy (paired t-test $T_{21}=2.39$; $P=0.026$). Within-perception classification accuracy was significantly above chance, uncorrected ($T_{21}=2.32$; $P=0.03$). Inset: Classification accuracy in the ventricles was not different from chance (perception: $P=0.28$; memory: $P=0.79$). *** $P\leq0.001$ ** $P\leq0.01$ * $P<0.05$

810
Figure S8. Low-level visuomotor features of the rating scale. We trained a classifier with sub-
jects' low (1-2) and high (3-4) ratings from the Confidence condition and tested on trials from
the Follow condition (and vice versa). Since rating-scale numbers in Follow trials were not relat-
ed to confidence level, any accurately classified voxel could be expected to reflect visuomotor
requirements of using the confidence scale rather than evaluating confidence. Accordingly, this
analysis identified regions of visual cortex, parietal cortex and primary motor areas. This group-
level accuracy map was used as an exclusive mask in whole-brain searchlight analyses of CLR-
activity to eliminate visuomotor-related activations. All clusters are significant at $P_{FWE}<0.05$
cluster-level corrected for multiple comparisons using a cluster-defining threshold of $P<0.001$.
Image displayed at $P<0.001$, uncorrected. The bar indicates T-score (Table S5).

Figure S9. Mean raw confidence level-related classification accuracy scores. For illustration purposes, we present here whole-brain maps of group mean raw accuracy scores for within-domain confidence level-related classifications (Figure 5D), exclusively masked by visuomotor activity (Figure S8). Color bars indicate mean classification accuracy. Circles indicate the four selected ROIs (Figure 4 – figure supplement 1). **(A)** Within-perception. **(B)** Within-memory.

Figure S10. Classification accuracy of confidence level-related activity using accuracy as
proxy. Cross-classification accuracy results of training a decoder on CLR-activity and testing it
on activity from the Follow condition using accurate vs inaccurate trials as proxies of Low and
High confidence (and vice versa). Classification accuracy was above chance in dACC/pre-SMA
(T_{21}=2.58, P=0.0175) and right rlPFC (T_{21}=2.48, P=0.0215). Cross-classification of activity in
left rlPFC (T_{21}=-0.13, P=0.898) and precuneus (T_{21}=-0.38, P= 0.71) was at chance. Bars indicate
group means and bars s.e.m. * P<0.05

Table S1. Judgment-related univariate activity for Confidence>Follow contrast. Significant
activations at cluster-defining threshold *P*<0.001, corrected for multiple comparisons at
P_{FWE}<0.05

Label	Voxels at *P<0.001*	P_{FWE} cluster-corrected	Peak *z*-score	Peak voxel MNI coordinates	Laterality
Anterior cingulate gyrus, dACC, pre-SMA (BA 6, 24,32)	802	<0.001	4.97	6, 29, 26	L/R
Insula	323	<0.001	4.31	-27, 17, 8	L
Superior and middle frontal gyri	75	0.018	4.07	30, 41, 17	R
Cuneus, lingual gyrus	86	0.010	3.97	9, -67, -10	R
Insula, superior temporal gyrus	117	0.002	3.80	39, 11, 2	R

841 **Table S2. Confidence level-related univariate activity parametric correlations.** Significant
842 activations at cluster-defining threshold $P<0.001$, corrected for multiple comparisons at
843 $P_{FWE}<0.05$

Correlation	Label	Voxels at $P<0.001$	P_{FWE} cluster-corrected	Peak z-score	Peak voxel MNI coordinates	Laterality
Positive	Post- and pre-central gyri	848	<0.001	5.30	-33, -16, 71	L
	BA 11, 25	445	<0.001	5.02	-6, 17, -4	L/R
	Lingual gyrus, cerebellum	323	<0.001	4.85	12, -58, -10	R
	SMA, BA 31	228	<0.001	4.82	-3, -10, 53	L/R
	Hippocampus, para-hipocampal gyrus	59	0.019	4.43	-27, -16, -16	L
	Post- and pre-central gyri	107	0.001	4.41	42, -22, 47	R
	Cuneus	53	0.030	4.12	18, -82, 35	R
	Posterior cingulate	51	0.035	3.81	-6, -55, 14	L
Negative	dACC, pre-SMA	553	<0.001	5.12	6, 29, 44	L/R
	Cerebellum, lingual gyrus, cuneus	287	<0.001	4.85	-39, -61, -43	L
	Middle and inferior frontal gyrus, BA 9	272	<0.001	4.55	-42, 17, 26	L
	Middle and inferior frontal gyrus, BA 46, 9	498	<0.001	4.39	48, 17, 29	R
	Parietal cortex, BA 40	135	<0.001	3.97	36, -52, 44	R
	Parietal cortex	71	0.008	3.81	-30, -61, 47	L

845
846

Table S3. Confidence level-related univariate activity parametric correlations by domain.
Significant activations at cluster-defining threshold $P<0.001$, corrected for multiple comparisons at $P_{FWE}<0.05$

Correlation	Label	Voxels at $P<0.001$	P_{FWE} cluster-corrected	Peak z-score	Peak voxel MNI coordinates	Laterality
Positive						
Perception	Precentral gyrus, BA6	231	<0.001	4.78	-30, -25, 53	L
	Paracentral lobule (supplementary motor area), BA6	184	<0.001	4.55	-3, -16, 50	L/R
	Precuneus, posterior cingulate	139	<0.001	4.30	15, -73, 20	L/R
Memory	Precentral & postcentral gyri, BA6	723	<0.001	5.46	-30, -16, 71	L
	Cerebellum	55	0.037	4.54	-15, -37,-13	L
	Cerebellum	65	0.018	4.41	24, -46, -25	R
	Precuneus	63	0.021	4.40	21, -49, -17	R
	Middle Temporal Gyrus	52	0.045	4.08	-45, -73, 2	L
	Anterior cingulate, BA25	61	0.024	4.05	-6, 20, 1	L/R
Conjunction	Precentral & postcentral gyri, BA6	167			-30, -25, 53; -30, -16, 71	L
Negative						
Perception	Inferior parietal lobule, precuneus, BA7, BA40	434	<0.001	5.22	33, -61, 32	R
	Inferior & middle frontal gyri, BA9, BA10, BA46	364	<0.001	4.46	48, 5, 23	R
	Inferior & middle frontal gyri, BA9	214	<0.001	4.36	-45, 5, 29	L
	dACC, pre-SMA, BA8	239	<0.001	4.25	6, 29, 44	L/R
	Superior and inferior parietal lobules, precuneus, BA7, BA40	121	<0.001	4.16	-30, -61, 47	L
Memory	Lingual gyrus, cuneus	106	0.001	4.42	-6, -88, 2	L
	Superior & middle frontal gyri, BA6	94	0.003	4.04	36, 14, 56	R
	Inferior & middle frontal gyri	87	0.004	3.89	51, 17, 32	R
Conjunction	Middle frontal gyrus, BA9	29			51, 17, 32	R

852 **Table S4. Judgment-related activity in whole-brain searchlight classification accuracy map.**
853 Significant activations at cluster-defining threshold *P*<0.001, corrected for multiple comparisons
854 at P_{FWE}<0.05

Classification	Label	Voxels at *P*<0.001	P_{FWE} cluster-corrected	Peak *z*-score	Peak voxel MNI coordinates	Laterality
Across-domain	Insula/IFG	592	0.001	4.31	-33, 23, 5	L
	SFG & BA9 dACC/pre-SMA	928	<0.001	4.29 4.19 4.01	24, 53, 38 -12, 14, 41 12, 44, 23	R L R
	Supramarginal gyrus, BA 40	207	0.039	3.93	-51, -55, 29	L
	IFG/Insula/STG	275	0.016	3.61	48, 5, -13	R
Within-perception	Superior, Middle, Medial, Inferior FG, dACC, pre-SMA, BA 8, 10, 32	4699	<0.001	5.10	-12, 14, 59	L/R
	Parietal cortex, precuneus, supramarginal gyrus, BA 7, 40	1141	<0.001	4.30	51, -46, 44	R
	Precuneus, BA 40	339	0.010	4.21	-36, -46, 47	L
Within-memory	Middle & Superior FG, BA 9, 10	325	0.008	4.54	39, 41, 38	R
	Lingual gyrus, cuneus, calcarine	510	0.001	4.53	9, -91, 2	R
	Superior FG, dACC, pre-SMA	1066	<0.001	4.33	6, -4, 62	L/R
	Middle, Inferior FG, BA 9, 10, 45, 46	1003	<0.001	4.09	-42, 38, 2	L

857 **Table S5. Low-level visuomotor activity (Confidence/Follow ratings) in whole-brain search-**
858 **light classification accuracy map.** Significant activations at cluster-defining threshold $P<0.001$,
859 corrected for multiple comparisons at $P_{FWE}<0.05$
860

Label	Voxels at $P<0.001$	P_{FWE} cluster-corrected	Peak z-score	Peak voxel MNI coordinates	Laterality
Parietal lobe, occipital lobe, right insula & right middle FG	11240	<0.001	6.14	-39, -37, 47	L/R
Cerebellum	195	0.039	4.88	30, -70, -28	R

862 **Figure S6. Confidence level-related activity in whole-brain searchlight classification accu-**
863 **racy map.** Accuracy maps were masked to exclude low-level visuomotor regions (see Figure S8
864 & Table S5). Significant activations at cluster-defining threshold $P<0.001$, corrected for multiple
865 comparisons at $P_{FWE}<0.05$
866

Classification	Label	Voxels at $P<0.001$	P_{FWE} cluster-corrected	Peak z-score	Peak voxel MNI coordinates	Laterality
Across-domain	Inferior FG, ACC, BA 11, 32	2434	<0.001	5.20	-21, 44, 2	L/R
	Superior and middle temporal gyri	176	<0.001	4.99	-57, -52, 20	L
	Middle temporal gyrus, anterior cerebellum	1193	<0.001	4.92	3, -46, -34	R
	Middle cingulate gyrus	46	0.017	4.85	-6, -10, 47	L/R
	Superior temporal gyrus	142	<0.001	4.68	66, -7, 5	R
	Precuneus, BA 7, 19	543	<0.001	4.68	-12, -61, 65	L/R
	Superior temporal gyrus	110	<0.001	4.68	-42, 26, -28	L
	Posterior cerebellum	96	<0.001	4.54	-9, -85, -22	L
	SMA, BA 6	84	0.001	4.53	-9, -10, 62	L/R
	Superior FG, dACC, pre-SMA	482	<0.001	4.39	-3, 11, 56	L/R
	Post-central gyrus	186	<0.001	4.33	-57, -19, 23	L
	Middle FG1	143	<0.001	4.28	36, 5, 26	R
	Fusiform and parahippocampal gyri	63	0.003	4.27	36, -19, -34	R
	Middle cingulate cortex	64	0.003	4.25	0, -31, 53	L/R
	Posterior cerebellum	53	0.008	4.10	-48, -58, -37	L
	Inferior FG	38	0.039	3.84	27, 23, -25	R
Within-perception	Superior FG, dACC/pre-SMA	185	0.032	3.88	-6, 29, 47	L/R
Within-memory	Inferior and middle FG, insula	503	<0.001	4.35	57, 23, 14	R
	dACC, pre-SMA	211	0.021	3.79	-9, 2, 47	L/R

868 **Figure S7. Metacognitive efficiency scores predictive of confidence level-related activity in**
869 **whole-brain searchlight classification accuracy maps.** Significant activations in the memory-
870 domain at cluster-defining threshold $P<0.001$, small-volume corrected for multiple comparisons
871 at $P_{FWE}<0.05$ using an anatomical precuneus mask. Significant activations in the perceptual do-
872 main at cluster-defining threshold $P<0.001$ (uncorrected).

Domain	Label	Voxels at $P<0.001$	P_{FWE} small-volume cluster-corrected	Peak z-score	Peak voxel MNI coordinates	Laterality
Memory	Precuneus, BA7	54	0.028	4.39	9, -70, 56	R

Domain	Label	Voxels at $P<0.001$	P cluster-uncorrected	Peak z-score	Peak voxel MNI coordinates	Laterality
Perception	Middle frontal gyrus, BA11	103	0.041	4.28	-15, 32, 11	L
	Cerebellum	111	0.035	4.16	-9, -76, -43	L/R
	Medial frontal gyrus, BA11	127	0.026	3.98	15, 50, -10	R

897 ***Supplementary References***
898
1. Fleming SM, Huijgen J, Dolan RJ. Prefrontal Contributions to Metacognition in Perceptual
Decision Making. J Neurosci. 2012 May 2;32(18):6117–25.
2. Fleck MS, Daselaar SM, Dobbins IG, Cabeza R. Role of Prefrontal and Anterior Cingulate
Regions in Decision-Making Processes Shared by Memory and Nonmemory Tasks. Cereb
Cortex. 2006;16(11):1623–30.
3. Hebart MN, Schriever Y, Donner TH, Haynes JD. The Relationship between Perceptual
Decision Variables and Confidence in the Human Brain. Cereb Cortex. 2014;26(1):118–30.
4. Baird B, Smallwood J, Gorgolewski KJ, Margulies DS. Medial and Lateral Networks in
Anterior Prefrontal Cortex Support Metacognitive Ability for Memory and Perception. J
Neurosci. 2013 Oct 16;33(42):16657–65.
5. Fleming SM, Weil RS, Nagy Z, Dolan RJ, Rees G. Relating Introspective Accuracy to Indi-
vidual Differences in Brain Structure. Science. 2010 Sep 16;329(5998):1541–3.
6. McCurdy LY, Maniscalco B, Metcalfe J, Liu KY, de Lange FP, Lau H. Anatomical Cou-
pling between Distinct Metacognitive Systems for Memory and Visual Perception. J Neuro-
sci. 2013 Jan 30;33(5):1897–906.
7. Andersson JLR, Hutton C, Ashburner J, Turner R, Friston K. Modeling Geometric Defor-
mations in EPI Time Series. Neuroimage. 2001 May;13(5):903–19.
8. Sladky R, Friston KJ, Tröstl J, Cunnington R, Moser E, Windischberger C. Slice-timing ef-
fects and their correction in functional MRI. Neuroimage. 2011 Sep 15;58(2):588–94.
9. Ashburner J, Friston KJ. Unified segmentation. Neuroimage. 2005 Jul;26(3):839–51.
919