Anisotropic Cellular Mechanoresponse for

Radial Size Maintenance of Developing Epithelial Tubes

Tsuyoshi Hirashima*[1] and Taiji Adachi[2]

1. Department of Pathology and Biology of Diseases, Graduate School of Medicine,
Kyoto University

2. Institute for Frontier Life and Medical Sciences, Kyoto University

Correspondence to:

Tsuyoshi Hirashima, Ph.D.

Department of Pathology and Biology of Diseases, Graduate School of Medicine, Kyoto

University, Yoshida-konoe-cho, Sakyo-ku, Kyoto 606-8315, Japan.

Tel/Fax: +81-75-753-9450

E-mail: hirashima.tsuyoshi.2m@kyoto-u.ac.jp

Abstract

Cellular behaviors responding to mechanical forces control the size of multicellular tissues as demonstrated in isotropic size maintenance of developing tissues. However, how mechanoresponse systems work to maintain anisotropic tissue size including tube radial size remains unknown. Here we reveal the system underlying radial size maintenance of the murine epididymal tubule by combining quantitative imaging, mathematical modeling, and mechanical perturbations. We found that an oriented cell intercalation making the tubule radial size smaller counteracts a cell tension reduction due to neighbor cell division along the tubule circumferential axis. Moreover, we demonstrated that the tubule cells enhance actomyosin constriction driving the cell intercalation in response to mechanical forces anisotropically applied on the cells. Our results suggest that epididymal tubule cells have endogenous systems for responding as active cell movement to mechanical forces exclusively along the circumferential axis, and the anisotropic cellular mechanoresponse spontaneously controls the tubule radial size.

Keywords: Anisotropy; Epithelial tube; Mathematical model; Mechanoresponse; Quantitative imaging; Size maintenance; Tissue mechanics

Introduction

Cell-cell communication via cellular behaviors is necessary to regulate tissue maintenance, development, and size. The underlying principle for this communication is a cellular response produced in reaction to mechanical forces generated within the tissues [1,2]. In tissue volume maintenance, for example, cell proliferation is counterbalanced by cell death, and these coordinated cellular behaviors are the mechanoresponse to cell-crowding pressure due to proliferation, which leads to cell death within the tissue [3–5]. On the other hand, in tissue length maintenance along an axis, such as radial size maintenance of a tube during elongation [6–8], the anisotropic mechanoresponse system in tissues might organize cooperative cellular behaviors to distribute a net increase in the tissue volume not along the axis, but in other axes. However, the system responsible for this anisotropic tissue maintenance is unclear. Here we utilized epididymal tubules in murine embryo as an experimental model system for a proliferating single-layered epithelial tube [7,9] to reveal anisotropic cellular response to mechanical forces for radial size maintenance of proliferating tubes.

Results

Actomyosin constriction generated within epithelial cells is required for the maintenance of tubule radial size

In the epididymis, phosphorylated myosin regulatory light chain (pMRLC), a marker of active tension generation via actomyosin constriction, localizes only to apical cellular junctions of the epithelial tubules at 15.5 and E16.5 (Fig. 1A). To examine the mechanical roles of actomyosin constriction in the epididymal tubules, we applied Blebbistatin/Y-27632 for inhibiting actomyosin constriction of non-muscle myosin II [10,11] or Calyculin A for inhibiting the protein phosphatases of pMRLC [12] to the epididymides in explant cultures, and observed the tubule morphology 1 day after the inhibitor treatment (Fig. 1B). The inhibitor assay revealed that inhibiting actomyosin-mediated apical constriction resulted in tubule expansion whereas constitutive activation of the tension led to tubule shrinkage (Fig. 1B´; Materials and Methods (II-i)).

     To clarify the inhibitor effects at the cell level, we examined morphological changes of the tubule cells by immunolabeling for the E-cadherin (Fig. 1C). For the Blebbistatin treatment as in the case of Y-27632, the tubule cells became flattened, i.e., those became more extended parallel to the tubule surface and the cell length along the apico-basal axis

became shorter than those in control. For Calyculin A treatment, each cell shape became

rounded due to strong cellular constriction, leading to the disintegration of cell-cell adhesion;

as a result of the aberrant constriction, some cells were excluded from the tubules and

lumenized structure of the epididymal tubules was disrupted (Fig. 1C).

For quantitative analysis at single cell resolution, we measured the cell height along the

apico-basal axis and the cell area at the middle point of the cell height, and calculated the cell

volume and the aspect ratio of cell shape as a measure for the cell flatness (Fig. 1D; Materials

and Methods (II-ii)). Interestingly, we found the cell volume was preserved even though the

aspect ratio became larger for the increase of tubule diameter in the case of Blebbistatin

treatment (Fig. 1D´). Based on this data, we performed mathematical analysis to examine

how the force balance in epithelial cells within mono-layered tissues influences the cellular

morphology at mechanical equilibrium and clarified that the cell area/height monotonically

increases/decreases for the attenuation of apical constriction (Materials and Methods (III-i)),

which can explain the variation of cell morphology for the inhibitor treatments. Taken

together, these results indicate that the force balance within the epithelial layer of the tubule

determines its radial size.

Cell division along the circumferential axis of tubules produces mechanical forces to

expand the tubule

Next, we examined the impact of cell division orientation in the tubule at E15.5 and E16.5

because the cell division orientation is a key determinant for the morphogenesis of

developing tubules [13,14]. We used phospho-histon H3 (pHH3) and γ-tubulin each as a

marker of mitotic cells and microtubule organizing centers (MTOCs). We measured spindle

orientation in the mitotic cells within the tubule by extracting the axis connecting between the

two MTOCs in each pHH3-positive cell without destructing the tissue structure (Fig. 2A-A").

We manually measured three angles α_1 , α_2, and α_3 (Fig. 2B) to define a local coordinate

system O' (Fig. 2C), origin of which corresponds to the center position of mitotic cell (white

arrows in Fig. 2B). In the local sphere coordinates, we obtained statistical distributions for

two angles of the cell division orientation: φ and θ (Fig. 2C; Materials and Methods

(II-iii)). The distribution of φ shows that most of the mitotic cells divide parallel to the

epithelial layer (The Rayleigh test, $P<0.001$ at E15.5 and E16.5); however, the distribution of

θ shows that the orientation is not biased towards the longitudinal or circumferential axis

(The Rayleigh test, $P>0.05$ at E15.5 and E16.5) (Fig. 2D). Furthermore, the distance between

MTOCs shows no correlations with the angle θ (r=-0.07 at E15.5, r=-0.02 at E16.5 where r

is the Pearson's correlation coefficient), denying a possibility that the spindle orientation

might turn to a specific orientation when the later stage of mitosis (Fig. 2E).

We then investigated how the cell division orientation affects the tubule morphology

using numerical simulations of a computational model (Mov. 1). We here employed the

vertex dynamics model, a type of cell-oriented model, to represent multi-cellular dynamics in

epithelial tubes [15–17]. See the Materials and Methods (III-ii) - (III-iv) for more details of

the mathematical model and simulation settings (Fig. S1). In the simulation, virtual epithelial

tubes grew along the division orientation, suggesting that cell division along the

circumferential axis is attributed to continuous radial growth (Fig. 2F and F′). In addition, we

found that the cell division orientation affects mechanical stress in each cell of the virtual

tubes (Fig. 2F). To evaluate the mechanical stress in the cells related to the tube

morphogenesis, we introduced a scalar quantity, i.e., cell tension anisotropy, which becomes

larger when the cell tension along the circumferential axis becomes weaker than that along

the longitudinal axis. See Eq. (v-2) in the Materials and Methods (III-v) for the definition of

cell tension anisotropy. As shown in Figure 2F-F′′, the cell tension anisotropy in the tubes

depends on the cell division orientation and corresponds with the growth direction of the

tubes. The simulation analysis indicates that the circumferential cell division leading to the

tube expansion entails the increase of cell tension anisotropy locally in the tissue. This

implies that cells should actively counteract the changes in cell tension anisotropy for tubule

radial size maintenance.

Oriented cell intercalation and circumferential cell division should be coupled for the

maintenance of tubule radial size

To explore how the cells react, we examined the pMRLC spatial distribution in the tubule at

sub-cellular resolution through automatic extraction for each apical cell junction (Fig. 3A).

We performed whole-tissue coimmunostaining for pMRLC and Zonula occludens-1 (ZO-1),

an apical tight junction marker, and quantified the pMRLC mean intensity at junctions,

junction angle, and length through digital image processing (Fig. 3B; Materials and Methods

(II-iv)). We found that the distribution of pMRLC intensity was well fitted to a log-normal

distribution (Fig. 3C), indicating that enriched pMRLC localization can be observed in an

unignorable number of apical junctions (Fig. 3D). To evaluate the relationship between the

junction angle and the pMRLC intensity, we categorized the junction angle into three groups;

longitudinal (long.): $0 \le \vartheta \le \pi/6$, intermediate (intm.): $\pi/6 < \vartheta < \pi/3$, and circumferential

(circ.): $\pi/3 \le \vartheta \le \pi/2$ (Fig. 3E). As shown in Figure 3E′, the pMRLC tends to localize more

on junctions that orient more in the circumferential axis (One-way ANOVA, $P<0.001$). As

well as the junction angle, we also categorized the junction length into three groups: short:

$0 \le \ell \le 2$ μm, middle: $2 < \ell < 4$ μm, long: $4 \le \ell \le 10$ μm (Fig. 3F), and found that the

pMRLC tends to localize more on shorter junctions (One-way ANOVA, $P<0.001$) (Fig. 3F′).

These results suggest that the circumferential junctions shrink through the

actomyosin-mediated constriction.

Because pMRLC localization on the apical junctions triggers cell intercalation mediated

through junction shrinkage [1,18], the polarized pMRLC localization on the circumferential

junctions can drive the oriented cell intercalation leading to longitudinal elongation and radial

shrinkage of the tubule. The present data suggest that the oriented cell intercalation causes the

tubule shrinkage; on the other side, the tubule expansion occurs due to the circumferential

cell division shown in the previous section. Therefore, the maintenance of tubule radial size

requires a proper balance between myosin activity driving intercalation and cell division

frequency.

To evaluate this relationship, we measured the pMRLC intensity along the tubule at the

whole organ scale at E15.5 and E16.5 (Materials and Methods (II-v)), and found that the

intensity in the head region of epididymis is larger than that in the tail region (Fig. 3G). This

intensity profile is similar to the spatial profile of the number of mitotic cells in the

epididymal tubules previously reported [19] (Fig. 3H). Indeed, the pMRLC intensity and the

mitotic cell numbers in the tubules show positive correlation (r=0.49 in E15.5 and r=0.56 in

E16.5 where r is the Pearson's correlation coefficient) (Fig. 3 H′; Materials and Methods

(II-vi)). Western blotting analysis also shows the myosin activity, defined as the ratio of

pMRLC to the total MRLC (tMRLC), in the head region is greater than that in the tail region

(Fig. 3I and I′). Moreover, cell morphology differences corroborate junctional tension due to

the pMRLC in each region (Fig. 3J and S2; Materials and Methods (II-vii)). The apical

junction length along the circumferential axis in the head region is shorter than that in the tail

region (Kruskal-Wallis test, $P<0.001$) (Fig. 3J′), and the apical surface area in the head

region are smaller than that in the tail region (Kruskal-Wallis test, $P<0.001$) (Fig. 3J′′). Thus,

a regulatory mechanism balancing the oriented cell intercalation driven by pMRLC

localization and cell division should exist.

Cell division provides mechanical stimulations to the neighboring cells

Cell division can generate mechanical perturbations in multi-cellular tissues [20]; hence, we

anticipated that the cell divisions load the mechanical forces to the adjacent cells of divided

ones and this stimulus would trigger the oriented cell intercalation in the epididymal tubule.

We then investigated the change of the cell tension anisotropy against the cell division using

extensive numerical simulations of the vertex dynamics model.

We examined the dependency of cell division anisotropy against the two factors: relative

position of cells to the neighbors cell division axis (Fig. 4A) and normalized elapsed time

from the cell division $\tau_\alpha'^{\text{cycle}}$, ranged from 0 to 1 (see the Materials and Methods (III-iv) for

the definition). In addition, we focused on the two cases of cell division orientation, i.e.,

longitudinal ($0 \leq \theta \leq \pi/6$ where θ is defined in Fig. 2C) and circumferential

($\pi/3 \leq \theta \leq \pi/2$), and made histograms of the cell tension anisotropy for each case (Fig. 4B).

The results shown in Figure 4B´ indicate that the cell division mechanically affects the cell

tension anisotropy in the neighboring cells of divided cells on the division axis (Rows in Fig.

4B´). This is intuitive because the divided cells push along the division axis to the

neighboring cells and hence, tension in the neighboring cells along the division axis can be

attenuated. Furthermore, the histograms show that the attenuation effect due to the cell

division becomes smaller along with the elapsed time from the cell division $\tau_\alpha'^{\text{cycle}}$. (Columns

in Fig. 4B´).

Would the cell division be the only source to produce the tension attenuation of cells

within multicellular tissues? To answer this question, we next plotted histograms of the

elapsed time from the cell division $\tau_\alpha'^{\text{cycle}}$ when the cell tension anisotropy ρ_α is over

various thresholds $\rho^* = 0.039, 1, \text{and } 0.2$ in the cells on the circumferential axis of divided

cells (Fig. 4C-C´´; $\rho^* = 0.0385$ is the mean of cell tension anisotropy in the upper left case

of Fig. 4B´). Our numerical investigations revealed that the cell tension anisotropy could

become over the given thresholds at timing elapsed enough from the cell division although

the cell tension anisotropy in most of the cells (~70%) are affected immediately following to

the cell division.

These results led to the two conclusions. First, most of the cells that receive significant

mechanical impact from the cell division should be on the cell division axis and can be found

soon after the cell division. Second, cell tension can be varied independent to the cell division,

rationalizing the introduction of mechano-responsive regime into our model argued in the

later section.

Mechanical stimulations provided by neighbor cell divisions trigger the onset of cell

intercalation

To directly observe cellular responses to the mechanical forces generated by the cell divisions,

we performed live imaging analysis for epididymal tubule cells dissected at E15.5 using ex

vivo organ culture systems. For visualization of the tubule cell membrane, a conditional

fluorescence reporter line, crossed the R26R-Lyn-Venus line [21] and the Pax2-Cre line [22],

was used. Because the epididymal tubules were located in deep region of the epididymis,

~100 μm away from the capsule of epididymides (Fig. 5A), confocal microscopy was not

appropriate for this case; hence, we employed an incubator-integrated multiphoton excitation

microscope system for stable deep tissue live imaging in explant cultures (Fig. 5B).

We focused on the behavior of cells neighboring to mitotic cells on the cell division axis

as those cells are more likely to show cellular responses against the mechanical perturbations

according to the first conclusion in the previous section. Careful observations led to the

discovery of characteristic behaviors in those cells; the oriented cell intercalation for

elongating the tubule occurs more frequently against the circumferential cell division

compared against the longitudinal one (Fig. 5C and Mov. 2). For quantification of the

oriented cell intercalation behavior, we measured the cell intercalation strain rate [23] for the

circumferential ($0 \leq \theta \leq \pi/6$) or longitudinal cell division ($\pi/3 \leq \theta \leq \pi/2$) (Fig. 5C and S3),

and calculated the cell intercalation rate for tube elongation (CITE) as a measure of the

oriented cell intercalation in the tubules. Positive/negative values of the CITE indicate that

cell intercalations undergo for the tubule elongation/shortening (Fig. 5D; see the Materials

and Methods (II-viii) for more details of CITE). From the quantitative analysis, we found that

the oriented cell intercalation tends to occur 60 min after the circumferential cell division

(*t*-test P<0.05 in the time range: 50-60 min in the case of circumferential cell division) (Fig.

5E).

Considering the cellular movement as an output response, the oriented cell intercalation

should be operated through rapid biochemical process such as phosphorylation. In addition to

this inference, we demonstrated by the extensive numerical simulations that the cell division

within the epithelial tissues could attenuate the cell tension in adjacent cells along the

division orientation owing to pressure of the dividing cells to the adjacent ones (Fig. 2F and

4B′). Therefore, we hypothesized that MRLC would be phosphorylated as a cellular response

to the reduction of circumferential cell tension provided by the neighboring cells, ultimately

driving the oriented cell intercalation.

Polarized cellular responses to mechanical simulations lead to the maintenance of the

tubule radial size

To check whether this hypothesis is rational, we carried out the computer simulation of the

vertex dynamics model including a mechano-responsive regime and its counterpart regimes

(Mov. 3). We implemented the mechano-responsive regime into the mathematical model as

the cells keep constricting one of the circumferential junctions for a parameterized period

since the cell tension anisotropy ρ_α reached at the threshold value ρ^* due to the reception

of intense pressure (Fig. 6A and S4; Materials and Methods (III-v)). Through the numerical

analysis, we found that the mechano-responsive regime in the simulation realized

spatiotemporally-constant tubes with smaller radial size compared to the case of no

regulation (Fig. 6B and D).

On the other hand, we performed simulations in counterpart regime, the 'scheduled'

regime, where the timing for junction constriction was predetermined and the constricting

cells were randomly selected. For determining the schedule of the constriction, we sampled

the ratio of constricting cells along the course of normalized simulation time τ' in the

mechano-responsive regime and obtained the function by fitting the mean curve data (Fig.

6C; see Eq. (v-3) in the Materials and Methods (III-v)). In the scheduled regime, the

simulation generated tubes with irregular surfaces and non-constant radial size along the

longitudinal axis whereas the radial size was smaller as well as the case of

mechano-responsive regime (Fig. 6B, D and D′).

In the scheduled regime, the oriented cell intercalation that leads to the local shrinkage

of tubules occurs independently to the variations of cell tension, resulting in the irregular

surface of tubules. In the mechano-responsive regime, however, the mechanical stress at the

supra-cellular scale can be modulated due to the active multicellular movement immediately

responding to the mechanical perturbations; therefore, the smooth-curved surface throughout

the longitudinal axis of tubules can be realized in this regime. These simulation analyses

suggest that the mechano-responsive regime, as an embodiment of our hypothesis, explain

the radial size maintenance of proliferating tubes.

The epididymal tubule cells exhibit cellular responses as myosin activation to anisotropic mechanical perturbations

To verify the hypothesis, we applied uniaxial compression to isolated epididymides embedded in collagen gel using a flexible polydimethylsiloxane (PDMS) chamber, and evaluated the myosin activity as a cellular response to the compression (Fig. 7A). We performed the compression assay for 3 groups, i.e., no uniaxial compression as the control, lateral, and longitudinal compression, to examine the cellular response to different mechanical perturbations (Fig. 7B). After 10 min of continuous load, we quantified the cell shape and configuration altered to the uniaxial compression. For the quantification, we extracted the shape of epididymal cells from the immunofluorescence images for E-cadherin (Fig. 7C), and calculated the aspect ratio and the major axis angle from best-fit ellipses of the cell shapes. The shape and configuration of cells in the uniaxial compression treatments were significantly different from those in the no uniaxial compression, confirming that the epididymal tubule cells were deformed along the compression axis (The Kruskal-Wallis test, $P<0.001$) (Fig. 7D-D´´).

After 20 min, we measured the myosin activity as the relative intensity of pMRLC to the total MRLC (tMRLC) and found that the myosin activity is larger in the lateral compression compared in other cases, indicating that the MRLC phosphorylation is exclusively enhanced due to the lateral compression (The Friedman test, $P<0.05$) (Fig. 7E and E´). Moreover, we quantified the level of MRLC kinase as the ratio of the Rho-associated protein kinase 1 (ROCK1) to the tMRLC, and found that it was not significantly different between treatments (The Friedman test, $P>0.05$) (Fig. 7E and E´´). These results suggest that the epididymal tubule cells possess an intrinsic system in which MRLC phosphorylation is triggered in response to the cell tension reduction along the tubule circumferential axis without changing the total amount of ROCK.

Discussion

Altogether, we showed that the cells trigger myosin activation leading to oriented cell intercalation by responding to mechanical forces anisotropically in the tubule. Therefore, we propose a system in which the spontaneous mechanoresponsive behavior of cells at the supra-cellular scale organizes the maintenance of the tubule radial size at the tissue scale (Fig. 4F). In developmental processes, anisotropic growth of proliferating tissues while

maintaining an axis, such as tube elongation due to the radial maintenance specifically dealt
with in this study, causes generation of compressive forces in the growing axis, eventually
leading to a buckling-induced wavy pattern formation [7,24,25]. Thus, the proposed polarized
mechanoresponse system might work as a regulatory principle underlying mechanisms for
self-organized tissue morphogenesis [26,27].

To understand the system further, future work will need to identify what molecular
machineries sense the anisotropic mechanical force in tissues and how the received
mechanical stimuli are chemically transduced to activate the myosin in apical junctions. It is
tempting to speculate that the directional preference of cells would be provided by regulators
of planar cell polarity (PCP) that promote the downstream myosin phosphorylation [1,28,29],
such as Vang-like (VANGL) and Tyrosine-protein kinase-like 7 (PTK7). These proteins are
well aligned circumferentially in junctions of epididymal tubule cells and loss of PCP
disrupts the actomyosin-mediated cell intercalation resulting in tubule expansion [30] (Fig.
S5) as seen in kidney development [6]. Our findings reveal structural components and
dynamic properties in the mechanoresponse system that controls tissue morphogenesis.

Materials and Methods

This section includes (I) Experiments, (II) Quantifications, (III) Mathematical modeling, and (IV) Others (Statistical tests, etc.).

(I) Experiments

(i) Animals

For live imaging analysis, we used fluorescence reporter mice produced by crossing the Pax2-Cre line[22] (a gift from T. Furukawa in Osaka Univ.) and R26R-Lyn-Venus line[21]. Otherwise, we used imprinting control region (ICR) mice purchased from Japan SLC, Inc. We designated the midnight preceding the plug as embryonic day 0.0 (E0.0), and all mice were sacrificed by cervical dislocation to minimize suffering at E15.5 or E16.5. All the animal experiments were approved by the local Ethical Committee for Animal Experimentation of Institute for Frontier Life and Medical Sciences, Kyoto University, and were performed in compliance with the Guide for the Care and Use of Laboratory Animals at Kyoto University.

(ii) Antibodies and Fixative Solutions for Immunostaining

We used the following primary antibodies with dilution ratio 1:200 as a standard condition: rabbit monoclonal anti-E-cadherin (Cell Signaling Technology, #3195), mouse monoclonal anti-γ-Tubulin, (Sigma-Aldrich, #5326), rabbit polyclonal anti-phospho histon H3 (pHH3)(Merck Millipore, #06-570), rat monoclonal anti-pHH3 (Abcam, #ab10543), rabbit polyclonal anti-phosho Myosin light chain (pMRLC) (Abcam, #ab2480), rabbit polyclonal anti-Vangl1 (Atlas Antibody, #HPA025235), goat polyclonal anti-Vangl2 (Santa Cruz Biotechnology, #sc-46561), and mouse monoclonal anti-ZO-1 (Thermo Fischer Scientific, #33-9100). We used the paraformaldehyde (PFA) for anti-E-cadherin, anti-γ-Tubulin, and anti-pHH3; we used the trichloroacetic acid (TCA) for anti-pMRLC, anti-Vangl1, anti-Vangl2, and anti-ZO-1.

(iii) Whole-tissue Immunofluorescence and Fluorescent Dye

Dissected epididymides were fixed with 4% PFA in phosphate buffered saline (PBS) for 2 h at 4°C or for 20 min at 37°C, or with 2% TCA in Ca^{2+}- and Mg^{2+}-free PBS for 20 min at 4°C, depending on target epitopes. The samples were then blocked by incubation in 10% normal goat serum (NGS) (Abcam, #ab156046) or 10% normal donkey serum (Abcam, #ab166643)

diluted in 0.1% Triton X-100/PBS, depending on derived animals of secondary antibody, for

3 h at 37°C. The samples were treated with primary antibodies overnight at 4°C, and

subsequently incubated with secondary antibodies conjugated to either the AlexaFluor 546 or

the AlexaFluor 647 (1:1000, Thermo Fischer Scientific) overnight at 4°C. We used

TRITC-conjugated phalloidin (0.3 μg/ml, Merck Millipore, #FAK100) and Hoechst 33342 (5

μg/ml, Thermo Fischer Scientific, #H3570) for the visualization of F-actin and DNA,

respectively.

(iv) Volumetric Fluorescence Imaging

We mounted samples with 1% agarose gel of 10 μl on a glass-based dish (Greiner Bio-One,

#627871) for stable imaging. Then, the samples were immersed with the BABB solution

(benzyl-alcohol and benzyl-benzoate, 1:2) or CUBIC2 solution for optical clearing[19,31,32].

Finally, we obtained 8-bit volumetric fluorescence images using the confocal laser scanning

platform Leica TCS SP8 equipped with the hybrid detector Leica HyD. We used the

objective lenses magnifications of ×20 (numerical aperture (NA) = 0.75, working distance

(WD) = 680 μm, HC PL APO CS2, Leica), ×40 (NA = 1.3, WD = 240 μm, HC PL APO CS2,

Leica), or ×60 (NA = 1.4, WD = 140 μm, HC PL APO CS2, Leica).

(v) Explant Organ Culture

The epididymides were surgically excised at a region of the vas deferens close to the tail of

epididymis, and those with testis were dissected from the embryonic body. Samples were

then placed on a hydrophilic polytetrafluoroethylene organ culture insert with a pore size of

0.4 μm (Merck Millipore, #PICM01250), which was preset in a 35 mm petri dish (Thermo

Fischer Scientific, #153066) filled with a culture medium. The culture medium we used was

Dulbecco's Modified Eagle's medium (Nacalai Tesque, #08489-45), containing 10% fetal

bovine serum (Thermo Fischer Scientific, #12483) and 1% penicillin-streptomycin mixed

solution (Nacalai Tesque, #26253-84) at 37°C under 5% CO_2. The samples were cultured in

an air-liquid interface; the total amount of culture medium was 800 μl for the petri dish.

(vi) Small-molecule Inhibitors

We used the following inhibitors to suppress or activate the generation of

actomyosin-mediated tension: Blebbistatin (Merck Millipore, #203391), Y-27632 (Merck

Millipore, #SCM075), and Calyculin A (Cell Signaling Technology, #9902).

(vii) Live Imaging

(vii) Live Imaging

The organ culture setting was according to the condition as described above with the

following 2 minor modifications. First, we used a 35 mm glass-based dish (Iwaki,

#3910-035) with 700 μl of the culture medium in the dish. Second, we adhered each

supporting leg of the organ culture insert on the dish with 20 μl of 1% agarose gel. This

preparation prevents the culture insert from sliding during the live imaging. We used an

incubator-integrated multiphoton fluorescence microscope (Olympus) using a ×25

water-immersion lens (NA = 1.05, WD = 2mm, Olympus). Imaging conditions were as

follows; excitation wavelength: 945nm (Mai-Tai DeepSee eHP, Spectra-Physics), AOTF:

3.5%-1.5%, scan size: 1024 × 320 pixels, scan speed: 10 μsec/pixel, z-axis interval: 1 μm,

and time interval: 5 min, magnification: ×2.

(viii) Tissue Compression Assay

For the tissue compression assay, we used a manual stretch/compression device (STREX,

#STB-10) with a polydimethylsiloxane (PDMS) chamber (STREX, #STB-CH-04), and

conducted in the following manner. First, we exposed the PDMS chamber to plasma arc for 1

min using a desktop vacuum plasma processing device (STREX, PC-40) for hydrophilic

treatment to enhance the adhesion between the chamber and collagen gel into which the

epididymides were embedded. Next, we immediately placed isolated epididymides onto the

hydrophilized PDMS chamber in its 50% stretched state, and fulfilled with 500 μl of type I

collagen (Nitta Gelatin, Cellmatrix Type I-A), followed by gelation process for 10 min at

37°C. Then, we applied 1 ml of the culture medium into the chamber, and the epididymides

with the collagen gel were compressed by relaxing from the 50% stretched state around 0.2

mm/sec. No relaxing the stretched chamber was regarded as control treatment. Finally, the

samples were incubated for 10 min for the morphological confirmation analysis and 20 min

for the Western blotting analysis at 37°C under 5% CO_2. In the lateral compression assay, the

epididymal tubules were deformed along the circumferential axis in any peripheral region

and the circumferential length of individual cells became smaller. We consider this is due to

gravity of the culture medium filled on the upper layer of gel as this does not happen without

the culture media.

For the immunoblotting assay, we extracted total protein from each treatment, and

adjusted the amount to the same among the treatments for the SDS-PAGE as described above.

We first confirmed that there were no significant differences of GAPDH and those of MRLC

among the treatments. Then, we evaluated the myosin activity and the level of MRLC kinase

(ROCK1); even when the ROCK1 level in the epididymides was evaluated as the ratio of

ROCK1 to the GAPDH, it showed qualitatively the same results. There was no ROCK2

expression in the epididymides. Finally, the activities were each normalized by mean values

of the control group.

(ix) Total Protein Extraction and Western Blotting

For total protein extraction, dissected epididymides were immersed in the SDS-free RIPA

buffer with protease inhibitor cocktail (Nacalai Tesque, #08714) supplemented with the

EDTA-free phosphatase inhibitor cocktail (1:100, Nacalai Tesque, #07575), and were

disrupted by an ultrasonic cell disruptor (Microson). The lysates were placed onto the ice for

20 min and were centrifuged at 13,000 ×g for 10 min at 4°C. Protein concentration of the

supernatant was determined by the bicinchoninic acid assay.

The lysates were prepared for SDS-PAGE by adding 2×Laemmli sample buffer

(Bio-Rad, #161-0737) with 2-mercaptoethanol (Bio-Rad, #161-0710) and by boiling at 96°C

for 5 min. Next, the lysates containing approximately 5 μg of proteins were loaded into each

lane of the Mini-PROTEAN precast gels (Bio-Rad, #4569035), and electrophoresis was

carried out in Tris/glycine/SDS running buffer (Bio-Rad, #1610732) at constant 150V for 35

min. Then, the proteins were blotted onto 0.2 μm polyvinylidene difluoride membrane

(Bio-Rad, #1704272) in HIGH MW mode (1.3A, 25V for 10 min) of the Trans-Blot$^{®}$

Turbo^{TM} Transfer System (Bio-Rad, #170-4155) for ROCK1 detection and in the LOW MW

mode (1.3A, 25V for 5 min) for others.

The blotted membranes were then immersed in 15% H_2O_2/Tris-buffered saline (TBS)

solution for 30 min at room temperature for blocking endogenous peroxidase followed by

blocking with 5% NGS at 37°C for 60 min. For immunoblotting, the membranes were

incubated with primary antibodies diluted in 0.1% TBS/Tween-20 at 4°C over night. The

concentrations of antibodies used were 1:100,000 for mouse monoclonal anti-GAPDH (Wako,

#015-25473), 1:500 for rabbit polyclonal anti-myosin light chain 2 (Cell Signaling, #3672)

and mouse monoclonal anti-phospho-myosin light chain 2 (Cell Signaling, #3675), and

1:2000 for rabbit monoclonal anti-ROCK1 (Abcam, #ab45171). Then, the membranes were

incubated with diluted secondary antibody solutions in TBS/Tween-20 at 37°C for 60 min;

the concentration was 1:50,000 for goat anti-rabbit IgG conjugatd to horseradish peroxidase

(HRP) (Santa Cruz Biotechnology, sc-2004) and goat anti-mouse IgG conjugated to HRP

(Santa Cruz Biotechnology, sc-2005). Finally, protein bands were detected using the

Amersham ECL Select Western Blotting Detection Reagent (GE Healthcare, #RPN2235),

and were scanned with ImageQuant LAS 4000 (GE Healthcare).

(II) Quantifications

(i) Tubule Diameter in Inhibitor Assay (Fig. 1B and B')

First, we performed immunostaining for the E-cadherin of 8 different epididymides for each

inhibitor treatment to visualize the epididymal tubule as described above, and obtained the

volumetric images of 512 × 512 pixels using ×20 lens with z-axis interval of 5 μm. We

utilized the built-in module of a software platform, the Leica Suite X, (Leica), to create a

montage image of whole epididymis. Then, we manually extracted epididymal tubules from

maximum intensity projection images by erasing the efferent ductules and noise pixels due to

non-specific staining. From the preprocessed images, we obtained the centerline of

epididymal tubule by applying the built-in skeletonization algorithm in MATLAB for the

binary images obtained with arbitrarily determined thresholds, and finally calculated the

diameter along the centerline of tubule.

(ii) Aspect Ratio and Cell Volume in Inhibitor Assay (Fig. 1C-D')

First, we performed immunostaining for the E-cadherin of dissected epididymides to

visualize shape of individual epididymal cells as described above, and obtained the

volumetric images of 512 × 512 pixels using ×40 lens with z-axis interval of 1 μm. We

collected all samples from 3 different epididymides for each treatment. Then, we chose a

sampling point along the centerline of an epididymal tubule randomly, and collected 5

samples for the cell height and 20 cell samples for the characteristic cell length along the

circumferential direction of the epididymal tubule. For each sampling point, we manually

measured the cell height on an image plane that exposed the apical-basal axis of the cells

(right column of the Fig. 1C). In addition, we manually traced peripheral of the cells at the

middle point of cell height and measured cell area (left column of the Fig. 1C); we then

calculated the square root of the cell area for the characteristic length of the cells. Note that

the cell areas at any point of the cell height are not significantly different from those at

apical/basal sides. Utilizing the cell height and the characteristic length, we finally calculated

the aspect ratio between the two measures as defined in Fig. 1D, and the cell volume by the

multiplication of the two. Because the values of cell height are almost same within each

sampling point, we used only the mean value of cell height for the calculation of the aspect

ratio and that of the cell volume. Therefore, the standard deviations for the aspect ratio and

the cell volume represent the variations in the characteristic cell length.

(iii) Cell Division Orientation (Fig. 2A-D)

We first performed coimmunolabeling for γ-Tubulin and pHH3 of 6 different epididymides

to detect microtubule organizing centers of mitotic cells as described in this section (I)-(iii)

with a minor modification. That is, the samples were incubated in 10 mM sodium citrate

buffer including 0.1% Triton X-100/PBS for 20 min at 80°C for heat-induced retrieval of

γ-Tubulin epitope before blocking treatment. Next, we obtained volumetric images of 1024 ×

256 pixels as described above using ×40 lens with z-axis interval of 1 μm. Third, we

manually measured three angles α_1, α_2, and α_3 shown in Fig. 2B for each mitotic cell to define

a local coordinate system (O'; see Fig. 2C), origin of which corresponds to the center position

of mitotic cell. Each orthogonal basis of O'-coordinate is defined as follows: longitudinal

direction of epididymal tubule (x'), surface normal of epididymal tubule (z'), and orthogonal

direction for both x' and z' according to the right-handed system (y'). Then, we manually

measured positions of the two γ-Tubulin-positive dots (microtubule organizing centers:

MTOCs) in a pHH3-positive cell, and obtained two angles φ and θ in a local sphere

coordinate (Fig. 2D) resulted from coordinate transformation into the O'-system. Finally, we

selected samples under a criterion that the distance between the two γ-Tubulin-positive dots

is more than 1.5 μm. We regarded the samples whose φ is less than 40 degree as cells

dividing in parallel to the surface of epididymal tubule.

(iv) Apical Junction Morphology and pMRLC/VANGL Signal Intensity (Fig. 3A-F', Fig.

S2, and Fig. S5)

We performed coimmunostainining for ZO-1 and either pMRLC or VANGL2 of 8 different

epididymides as described above with a minor modification: the dilution ratio of

anti-pMRLC was 1:400 for preventing saturation binding. We obtained volumetric images of

1024 × 256 as described above using ×60 lens with z-axis interval of 0.3 μm. Then, we

separated the images into smaller ones around 200 pixels (≈36 μm) on a side so that the slope

of epididymal tubule can be regarded as a linear line.The longitudinal The digital image

processing was performed as follows. First, images for ZO-1 were filtered slice by slice with

a 2D median filter (3 × 3 pixels) followed by a 3D Gaussian filter with a standard deviation

of 3 pixels to reduce undesired noise. Next, the filtered images were binarized using the 3D

maximum entropy thresholding method [33], and the maximum one among the binarized component of 26-connected pixels was regarded as an apical junction network. Then, we performed 3D skeletonization process for the images of apical junctions, and obtained a graph network including nodes and edges of skeletonized voxels utilizing distributed MATLAB codes with modifications [34]. Finally, to extract the skeleton of apical cellular junctions (Fig. 3A), we screened out the skeletonized edges to avoid over-skeletonization results in the following 3 criteria; (1) The edge length between two nodes is less than 10 μm, (2) the edge length is shorter than two-fold linear length between two nodes linked with the corresponding edge, (3) the number of edges from a node is more than 3. The screening process under the criteria produced 96% left from the total number of raw skeletonized edges. The relative angle of apical junction edges against the longitudinal axis of epididymal tubule was calculated as $\vartheta_j = \left| \cos^{-1}\left(\hat{\mathbf{g}}_j \cdot \hat{\mathbf{l}} \right) \right|$ where j is an index of apical edges, $\hat{\mathbf{g}}$ is an unit vector of apical edges, and $\hat{\mathbf{l}}$ is that of the longitudinal axis of tubule, as shown in Figure 3B. For the signal quantification, we extracted signal intensity of pMRLC or that of VANGL2 on the apical cellular junctions by using a thickened skeleton of apical junctions with a dilation operator (3 × 3 × 3 voxels), and calculated average values for each edge of thickened apical junction skeletons.

(v) pMRLC Signal Intensity along Epididymal Tubule (Fig. 3G and H)

First, we performed immunostaining for the pMRLC of 12 different epididymides as described above with a minor modification: the dilution ratio of anti-pMRLC was 1:100. Second, we obtained the volumetric images of 512 × 512 as described above using ×20 lens with z-axis interval of 1 μm; we manually made montage images to avoid an overlap region of composite images. Third, we manually erased signal intensity on pixels corresponding to non-apical cellular region from the MIP images to obtain the centerline of epididymal tubule as described above. Then, we calculated average values of pMRLC intensity within a circle with a radius of 9 um along the epididymal tubule. Finally, we obtained the pMRLC signal distribution along the epididymal tubule by smoothing the averaged signal intensity with a moving average filter with a span of 40 μm – data in 20 μm range from the both ends of tubule were omitted. The maximum value of vertical axis in Fig. 3H is 1 which corresponds to the maximum intensity of 8 bit images. The horizontal axis of Fig. 3H was normalized by the tubule length of each sample.

(vi) Correlation between the Number of Mitotic Cells and pMRLC Signal Intensity (Fig. 3H')

We used the data on the number of mitotic cells from our previous work [19] and one on the pMRLC intensity obtained in this study. We divided 30 bins along the epididymal tubule and calculated maximum likelihood estimates with the Poisson distribution for the number of mitotic cells and with the Gaussian distribution for the pMRLC intensities in each bin. The Pearson's correlation coefficients are 0.49 (*P*-value=0.006) for E15.5 and 0.56 (*P*-value=0.001) for E16.5.

(vii) Apical Surface Area (Fig. 3J and 3J")

We obtained 1024 × 256 volumetric images of fluorescently-labeled F-actin in epididymal tubules as described above using ×40 lens with z-axis interval of 1 μm, and randomly collected samples of apical cell surface from image planes of 3 different epididymal volumetric data on which the apical cell surfaces exhibited. Then, we calculated the apical surface area by tracing the peripheral of apical cell surfaces manually.

(viii) Cell Intercalation Rate for Tube Elongation (CITE)

We first obtained time-lapsed volumetric images for epithelial dynamics from the head region of 4 different samples at E15.5 through the live imaging as described above, and collected time series data for a group of cells around mitotic cells. In particular, we sampled those cells on an image plane perpendicular to the optical axis from the image data, and focused on cross sections in-between the apical and basal part of cells. We defined cytokinesis as an origin of time for our analysis, and analyzed data on cell intercalation dynamics with time interval of 10 min ($\Delta t = 10 \, \text{min}$).

Next, based on an earlier study, we calculated the cell intercalation strain rate tensor $\mathbf{L}_{\text{I}}^{t}$ at time *t* as follows:

$$\mathbf{L}_{\text{I}}^{t} = \mathbf{L}_{\text{T}}^{t} - \mathbf{L}_{\text{C}}^{t},$$

where $\mathbf{L}_{\text{T}}^{t}$ is a local tissue velocity gradient tensor, and $\mathbf{L}_{\text{C}}^{t}$ is a cell shape strain rate tensor [23]. Because our analysis was performed in planar, the tensors are 2 × 2 matrices. For the calculation of a local tissue velocity gradient tensor $\mathbf{L}_{\text{T}}^{t}$, we specified one or two neighboring cells of a daughter cell from a mitotic cell that located an extension of the division axis as central cells, and marked the adjacent cells of the central cells including the central cells and the daughter cell as domain cells. Then, we manually traced each one of *n* domain cells to

obtain cell areas, cell shapes with the best fit ellipses, and cell centroids $\mathbf{r}_j^t$, where j is an

index of domain cells (see marked cells by yellow lines in Fig. S4A). Using data of the cell

centroids $\mathbf{r}_j^t$ and that of velocities $\mathbf{u}_j^t$, the local velocity gradient tensor was obtained by the

least squares approximations according to the following equation,

$\qquad \mathbf{u}_j^t \approx \left\langle \mathbf{u}^t \right\rangle + \mathbf{L}_{\mathrm{T}}\left(\mathbf{r}_j^t - \left\langle \mathbf{r}^t \right\rangle\right),$

where $\left\langle \mathbf{r}^t \right\rangle = \sum_j \mathbf{r}_j^t / n$, $\mathbf{u}_j^t = \left(\mathbf{r}_j^{t+\Delta t} - \mathbf{r}_j^t\right)/\Delta t$ and $\left\langle \mathbf{u}^t \right\rangle = \left(\left\langle \mathbf{r}^{t+\Delta t} \right\rangle - \left\langle \mathbf{r}^t \right\rangle\right)/\Delta t$. $\left\langle \cdot \right\rangle$ indicates an

average within the domain.

$\qquad$ For the calculation of the cell shape strain rate tensor $\mathbf{L}_{\mathrm{C}}^t$, we used lengths of principal

axes of a fitted ellipse and angle of the major axis from the longitudinal axis of epididymal

tubule to obtain a matrix $\mathbf{C}_j^t$ that represents the cell ellipse. Then, we calculated $\mathbf{L}_{\mathrm{C},j}^t$ from

$\mathbf{C}_j^t$ and $\mathbf{C}_j^{t+\Delta t}$, and $\mathbf{L}_{\mathrm{C}}^t$ was calculated by taking the area-weighted average of $\mathbf{L}_{\mathrm{C},j}^t$ within

the domain cells. Note that we normalized $\mathbf{L}_{\mathrm{C}}^t$ to satisfy that a trace of $\mathbf{L}_{\mathrm{C}}^t$ was equal to

that of $\mathbf{L}_{\mathrm{T}}^t$, realizing that the cell intercalation strain rate tensor $\mathbf{L}_{\mathrm{I}}^t$ had zero dilation.

$\qquad$ Finally, we obtained the CITE at time t (L_{ll}^t) from an element of the cell intercalation

strain rate tensor:

$\qquad \mathbf{L}_{\mathrm{I}}^t = \begin{pmatrix} L_{ll}^t & L_{lc}^t \\ L_{cl}^t & L_{cc}^t \end{pmatrix}.$

Note that the $L_{ll}^t = -L_{cc}^t$.

$\qquad$ The data were classified into two groups depending on the relative angle of cell division

axis to the longitudinal axis of epididymal tubule – we defined the relative division angle of

0-30 degrees as longitudinal division and that of 60-90 degrees as circumferential division.

(III) Mathematical Modeling

(i) Mathematical Analysis for Tube Morphology at Mechanical Equilibrium in Inhibitor

Assay (Fig. 1D)

We here argue a point that the results in the inhibitor assays can be explained by mechanical

force balance in the tubule tissue. To do so, we defined a potential energy of single epithelial

cells in tissues based on our experimental results, and analyzed how actomyosin-based

constriction on apical junctions affects epithelial morphology at mechanical equilibrium. We

referred to an earlier study [35] for the analysis in this section. The following notations are

valid only in this subsection.

We considered an epithelial sheet, each cell within which has the volume V and

adheres neighboring cells with the area S. Each single cell generates contractile forces at the

apical junctions whose perimeter P. Then, we defined the potential energy as a minimum

form from the epithelial morphology as follows:

$$U_{\text{Cell}} = \lambda_{\text{Adh}} S + \lambda_{\text{P}} P^2 + \lambda_{\text{V}} \left(V - V_0 \right)^2,$$ Eq. (i-1)

where V_0 is a target cell volume, λ_{Adh}, λ_{P}, and λ_{V} are coefficient of cell-cell adhesion,

that of apical constriction, and that of the cell volume constraint, respectively. To derive a

potential energy in the simplest form, we assumed the geometry of single epithelial cells as

hexagonal prisms of base length r and height h. Since $S = 2\sqrt{3}rh$ and $P = 2\sqrt{3}r$, we

have

$$U_{\text{Cell}} = 2\sqrt{3}rh\lambda_{\text{Adh}} + 12\lambda_{\text{P}}r^2 + \lambda_{\text{V}} \left(V - V_0 \right)^2,$$ Eq. (i-2)

instead of Eq. (i-1). As shown in Figure 1D', the cell volume is almost constant even though

the cell shapes change with the inhibitor, i.e., $V = V_0$; therefore, we have $h = 2V_0 / \sqrt{3}r^2$

since $V = \sqrt{3}r^2 h / 2$. Substituting these into Eq. (i-2) leads to

$$\frac{r}{4} U_{\text{Cell}} = V_0 \lambda_{\text{Adh}} + 3\lambda_{\text{P}} r^3.$$ Eq. (i-3)

To remove prefactors provided by the assumption of hexagonal prism and make a

non-dimensional form, we substituted $r = r'\left(V_0 / 3 \right)^{1/3}$ and $U_{\text{Cell}} = U'_{\text{Cell}} 4 \left(1/3 \right)^{-1/3} \lambda_{\text{Adh}} V_0^{2/3}$ into

Eq. (i-3), and then derived

$$U'_{\text{Cell}} = \frac{1}{r'} + \frac{\lambda_{\text{P}}}{\lambda_{\text{Adh}}} r'^2.$$ Eq. (i-4)

To obtain the apical cell area proportional to the square of cell base length at minimum of the

potential energy, we calculated $r'_{\text{eq}} = \left(r' \big| \partial U'_{\text{Cell}} / \partial r' = 0 \right)$, and then obtained

$$r'^2_{\text{eq}} \sim \lambda_{\text{P}}^{-2/3}.$$ Eq. (i-5)

Eq. (i-5) indicates that apical cell area/cell height becomes larger/shorter when the apical

constriction becomes weaker in the inhibitor (note $h = 2V_0 / \sqrt{3}r^2$), which is consistent with

our experimental results. Furthermore, this shows that the base length of cells r'_{eq} becomes

larger in the inhibitor, meaning the tubule composed of the cells should expand by the

inhibitors. Therefore, it is reasonable that mechanical force balance only within cells in the

tubule determines the morphological change of tubule observed in our experiments.

(ii) Basic Formalization of the Vertex Dynamics Model

We employed a vertex dynamics model (VDM), a type of cell-oriented model, to represent

multi-cellular dynamics in epithelial tissues [15,16,36–41]. In this model, a cell is

geometrically regarded as a polygon/polyhedron, of which vertices are elementary points that

constitute the cell shape, and a group of cells can be represented by a set of

polygons/polyhedrons shared by neighboring cells [17,42–44]. In this study, we paid notice

to apical junctions of the epididymal tubule on which a marker of tension generator pMRLC

is only localized (Fig. 1A), and modeled the developing single-layered epithelial tube as a

flexible multicellular membrane in 3D space according to earlier studies [45,46].

In the VDM, the dynamics of position of vertex i, $\mathbf{r}_i$, obey the equation of motion based

on the principle of least potential energy U as follows:

654
$$\eta\left(\frac{d\mathbf{r}_i}{dt} - \langle \mathbf{v}_i \rangle\right) = -\nabla_i U \,, \qquad \text{Eq. (ii-1)}$$

655 where η is a viscosity coefficient, and $\langle \mathbf{v}_i \rangle$ is a local velocity of vertex i, defined as

656
$$\langle \mathbf{v}_i \rangle = \frac{\sum_{\alpha_i}^{n_{\alpha_i}} d\mathbf{r}_{\alpha_i}/dt}{n_{\alpha_i}} \,, \qquad \text{Eq. (ii-2)}$$

657 where α_i is an index for cells contacting to vertex i, n_{α_i} is a number of the cells

658 contacting to vertex i, and $\mathbf{r}_{\alpha_i}$ is a centroid of cell α_i [38,47,48]. For a potential energy as a

659 minimum expression to represent epididymal tubule cells, we defined as

661
$$U = \sum_{\alpha}\left\{\frac{\lambda_{\text{A}}}{2}\left(A_\alpha - A_\alpha^{\text{O}}\right)^2 + \frac{\lambda_{\text{P}}}{2}P_\alpha^2\right\} + \sum_i \frac{\lambda_{\text{B}}}{2}\kappa_i^2 A_i + \sum_j\left\{\gamma_j^{\text{cst}}L_j - \gamma_j^{\text{ext}}H(L^{\text{O}} - L_j)\right\}, \quad \text{Eq. (ii-3)}$$

according to earlier studies [16,36,49]. α, i, and j each denote an index of cells, that of

vertices, and that of junctions/edges. The first term represents the cell elasticity; λ_{A} is its

coefficient, A_α is area of cell α, and A_α^O is a target area of cells. The second term

represents isotropic actomyosin contractility at the periphery of apical junctions; λ_P is its

coefficient and P_α is perimeter of the cell α. The third term represents bending energy of

the flexible epithelial membrane embedded in 3D space; λ_B is its coefficient, κ_i is a

discrete mean curvature defined at vertex i of triangular meshes [50], and A_i is summation

of an area of triangular mesh fractions around vertex i, each explained one by one. For

calculation of the discrete mean curvature at a vertex, let surface of the modeled epithelial

membrane be a set of triangular meshes consisting of both the vertex of cells and the centroid

of cells (Fig. S1A). Then, we defined the discrete mean curvature at vertex i as

$$\kappa_i = \frac{1}{2S_i} \left| \sum_{k \in V_i} \left(\cot\theta_{i,k}^1 + \cot\theta_{i,k}^2 \right) (\mathbf{r}_i - \mathbf{r}_k) \right|,$$ Eq. (ii-4)

where k is an index of the triangular elements included in the set of 1-ring neighbor vertices

of the vertex i, V_i, S_i is total area of triangular meshes in V_i, $\theta_{i,k}^1$ and $\theta_{i,k}^2$ are each angle

depicted in Figure S1A'. See [50] for more details. For calculation of A_i, let us consider

triangles consisting a vertex and its 1-ring neighbor cell centroids illustrated in Figure S1A'',

and then sum the all triangles up around the vertex. In earlier studies, bending energy was

defined as a summation of inner product of unit normal vectors to the surfaces between the

adjacent cells [45,46]. This is valid when all mesh size is ideally equal [51,52]; therefore, we

introduced the bending energy function based on the discrete curvature. The last term

represents anisotropic junction constriction/extension; $\gamma_j^{\text{cst/ext}}$ is its coefficient assigned to

each edge, L_j is length of edge, L^O is a target edge length, and H is the Heaviside step

function. For the simulation in Fig. 2F-F'', 4, and S1, $\gamma_j^{\text{cst/ext}}$ =0. The details of this term are

described later.

(iii) Normalized form of the model

By introducing a characteristic cell size A_0, the equations were transformed into

nondimensionalized form to reduce the number of parameters as

$$\frac{d\mathbf{r}_i'}{dt'} - \left\langle \mathbf{v}_i' \right\rangle = -\nabla_i' U',$$ Eq. (iii-1)

where $t' = t\lambda_A A_0 / \eta$, $\mathbf{r}_i' = \mathbf{r}_i / \sqrt{A_0}$, $\mathbf{r}_{\alpha_i}' = \mathbf{r}_{\alpha_i} / \sqrt{A_0}$, $\nabla' = \nabla\sqrt{A_0}$, $U' = U / (\lambda_A A_0^2)$, and

$$U' = \sum_{\alpha}\left\{\frac{1}{2}\left(A'_{\alpha} - A'^{O}_{\alpha}\right)^2 + \frac{\lambda'_{P}}{2}P'^{2}_{\alpha}\right\} + \sum_{i}\frac{\lambda'_{B}}{2}\kappa'^{2}_{i}A'_{i} + \sum_{j}\left\{\gamma'^{cst}_{j}L'_{j} - \gamma'^{ext}_{j}H(L^{O} - L'_{j})\right\},$$

Eq. (iii-2)

where $A'_{\alpha} = A_{\alpha}/A_0$, $A'^{O}_{\alpha} = A^{O}_{\alpha}/A_0$, $P'_{\alpha} = P_{\alpha}/\sqrt{A_0}$, $\kappa'_{i} = \kappa_{i}\sqrt{A_0}$, $A'_{i} = A_{i}/A_0$, $L'_{j} = L_{j}/\sqrt{A_0}$,

$L^{O} = L^{O}/\sqrt{A_0}$, $\lambda'_{P} = \lambda_{P}/(\lambda_{A}A_0)$, $\lambda'_{B} = \lambda_{B}/(\lambda_{A}A_0^2)$, $\gamma'^{cst/ext}_{j} = \gamma^{cst/ext}_{j}/(\lambda_{A}A_0^{-3/2})$. In addition, we

introduced a normalized simulation time defined as $\tau' = t'/T^{cycle}$, where T^{cycle} is an average

cell cycle length, which will be explained. We used these variables and equations, Eq. (iii-1)

and (iii-2), for the numerical simulations in this study.

(iv) Cell intercalation, Proliferation, and Simulation Conditions

As for the cell intercalation, we assumed that edge rearrangement occurred when the

normalized length of edge became smaller than 0.05 and the edge length was set to be 0.1

after the edge rearrangement. See [44] for details of the edge rearrangement.

As for the cell proliferation, we introduced the average cell cycle length T^{cycle} , and set

cell cycle length of each cell T^{cycle}_{α} obey the normal distribution of which the mean is T^{cycle}

and the standard deviation is 5% of the mean, i.e., $T^{cycle}_{\alpha} \sim N(T^{cycle}, 0.05T^{cycle})$. Also, we

introduced normalized cell cycle time of each cell as $\tau'^{cycle}_{\alpha} = t'^{cycle}_{\alpha}/T^{cycle}_{\alpha}$, where t'^{cycle}_{α} is

nondimensionalized cell cycle time of cell α . τ'^{cycle}_{α} at $t'=0$ was assigned within 0 to 0.95

in uniform distribution. As the average cell cycle length in the epididymal tubule can be

roughly estimated as 24 hours and cell volume tends to get increasing within the apical

surface of epididymal tubule since 50-100 min before the cytokinesis, we simply modeled the

cell dynamics during cell cycle as $A'^{target}_{\alpha} = 1$ for $0 \leq \tau'^{cycle}_{\alpha} < 0.95$, $A'^{target}_{\alpha} = 20\tau'^{cycle}_{\alpha} - 18$ for

$0.95 \leq \tau'^{cycle}_{\alpha} < 1$, and the cell divides with returning A'^{target}_{α} of daughter cells to 1.The type of

cell division orientation was set as follows – circumferential/longitudinal division: uniform

distribution in a range of $\pm 0.05\pi$ from the circumferential/longitudinal axis of tube, and

non-biased division: uniform distribution in all range of θ defined in the local coordinate at

cell (Fig. 2C).

As initial condition for the simulation, we set that a virtual tube was composed of 14

cells along the circumferential direction based on our observation and of 30 cells along the
longitudinal axis as for extensive numerical simulations. The cross-sectional center of the
tube was set at $y=0$ and $z=0$ of global coordinate and the tube was put along the x axis. As
boundary condition, the following energy was added to the original potential energy U' at
tip cells of the tube:

$$\sum_{\alpha_{\text{tip}}} \lambda'_{\text{bnd}} \left(\sqrt{(r'^{y}_{\alpha_{\text{tip}}})^2 + (r'^{z}_{\alpha_{\text{tip}}})^2} - R'_0 \right)^2 ,$$ Eq. (iv-1)

where α_{tip} is an index for the tip cells of tube, λ'_{bnd} is its coefficient, R'_0 is a normalized
mean radius of the tube to a characteristic cell length $\sqrt{A_0}$ in the initial setting, $r'^{y}_{\alpha_{\text{tip}}}$, and $r'^{z}_{\alpha_{\text{tip}}}$
each represent the relative position of center of cells α_{tip} on y and z axis to $\sqrt{A_0}$. This
energy restricts displacement of the cells α_{tip} in radial direction of the tube, which imitates
physical constraints by the vas deferens and the efferent ducts as boundary tissues of the
epididymal tubule. We set $\lambda'_{\text{bnd}} = 0.1$ in the simulations.

(v) Mechano-responsive and Scheduled regimes
In the following, we explain the last two terms of Eq. (iii-2): the anisotropic edge constriction
and extension after the cell intercalation. We here introduce a state variable assigned to each
cell, $\Theta_\alpha = \{0,1,2\}$; $\Theta_\alpha = 0,1,2$ represent a naïve state (colored in white, Fig. 6B),
edge-constriction state (red, Fig. 6B), and edge-extension state (blue, Fig. 6B), respectively.
For all cells, $\Theta_\alpha = 0$, $\gamma'^{\text{cst}}_j = 0$, and $\gamma'^{\text{ext}}_j = 0$, unless otherwise noted. In our simulation, the
following procedures were conducted for both 'mechano-responsive' and 'scheduled' regime.
First, cells α' that might become a constricting cell were selected and let the state variable
of the cell $\Theta_{\alpha'} = 1$; the way of selecting cells depends on each regime, i.e, the
mechano-responsive or the scheduled regime. Second, one of the edges that orients to
circumferential axis of the tube (the angle ϑ_j is within $70° - 90°$) was selected among all
edges composing the cells α', and index the edge j'. Then, state of cells including the
edge j' turned to $\Theta_{\alpha'} = 1$. Third, the constriction of edge j' was performed according to a
schedule as shown in Figure S4A; we set the edge kept constricting in a period τ'^{cst} via
transition period τ'^{trns} from the starting point of edge constriction by an assumption with

phosphorylation process of pMRLC. The maximum value of $\gamma_{j'}^{\prime\text{cst}}$ was set to be $\gamma^{\prime\text{cst}}$. $\Theta_{\alpha'}$

and $\gamma_{j'}^{\prime\text{cst}}$ return to 0 at the end of edge constriction process. In this study, parameter

dependence of $\tau^{\prime\text{cst}}$ and that of $\gamma^{\prime\text{cst}}$ were examined and determined (Fig. S4B), and we set

$\tau^{\prime\text{trms}} = 0.02$. Finally, if the edge arrangement occurred during the process of edge

constriction ($\Theta_{\alpha'} = 1$), then the state of cells having the rearranged edge changes to $\Theta_{\alpha'} = 2$

and the edge becomes an extension mode, in which the rearranged edge converges to a

constant target length; $\gamma_{j'}^{\prime\text{cst}} = 0$, $\gamma_{j'}^{\prime\text{ext}} = 0.01$, and $L^O = 0.1$. We set the period of edge

extension as $\tau^{\prime\text{ext}} = 0.02$. $\Theta_{\alpha'}$ and $\gamma_{j'}^{\prime\text{ext}}$ return to 0 after the process of edge extension. Our

implementation for the edge extension process is based on an earlier report that the junction

after the cell intercalation grows due to local polarized force driven by medial actomyosin

contraction [49].

As for the way of selecting the cells α' in the first step described above, we adopted

the two: mechano-responsive and scheduled regime. On the mechano-responsive regime, we

supposed that the epididymal tubule cells could receive mechanical stress provided by

adjacent cells, and a stress tensor in cells α in the local cell coordinate system O' was

defined as

$$\sigma_\alpha = \left(A_\alpha' - A_\alpha'^O\right)\mathbf{I} + \sum_{j_\alpha}\left(\lambda_P' + \gamma_{j_\alpha}^{\prime\text{cst}} - \gamma_{j_\alpha}^{\prime\text{ext}}\right)\frac{\mathbf{a}_{j_\alpha} \otimes \mathbf{a}_{j_\alpha}}{L_{j_\alpha}' A_\alpha'},$$ Eq. (v-1)

where $\mathbf{I}$ is a unit tensor, j_α is an index of edges composing the cell α, $\mathbf{a}$ is a vector

representing the relative position of two vertices composing the edge j_α [53–55].

Denoting a diagonal element of the tensor σ_α corresponding to the longitudinal axis of the

tube and that to the circumferential axis as $\sigma_\alpha^{\text{long}}$ and $\sigma_\alpha^{\text{circ}}$, we then defined the cell tension

anisotropy as

$$\rho_\alpha = \sigma_\alpha^{\text{long}} - \sigma_\alpha^{\text{circ}}.$$ Eq. (v-2)

Note that the cell tension anisotropy ρ_α is a time-dependent variable. In the

mechano-responsive regime, the cells were assigned as α' when ρ_α was over a threshold

$\rho*$, of which parameter dependency for the resulting tube morphology was examined (Fig.

S4B).

On the scheduled regime, the cells α' were randomly selected but timing of the

selection was predetermined according to the schedule shown in Figure 6C; Figure 6C shows

the fraction of cells having constricting edges to all tube cells in the mechano-responsive

regime (n=10, gray), and its mean (black). In our simulation, we used the following logistic

function (blue) by fitting the mean curve with the least-squared method:

$$f(\tau') = k_1\left(1+\exp\left[-k_2(\tau'-k_3)\right]\right)^{-1} \qquad \text{Eq. (v-3)}$$

with $k_1 = 0.1$, $k_2 = 0.01$, and $k_3 = 0.17$.

(vi) Morphological and Mechanical Quantities in Simulation

We used the following morphological quantities in the simulation; [1] tube length, i.e.,

longitudinal linear length between distal edges of tube, [2] tube diameter, i.e., mean diameter

of circumferentially-averaged diameter through the centerline of tube, and [3] curvedness, i.e.,

averaged curvature of the vertices throughout the tube defined as $\sum_i \kappa_i' A_i' \big/ \sum_i A_i'$.

For the cell tension anisotropy ρ_α, refer to the Eq. (v-2). For the tube tension

anisotropy in Figure 2F, the stress tensor in the tube was defined as

$$\sigma_{\text{tube}} = \left[\sum_\alpha \left(A_\alpha' - A_\alpha'^{O}\right)A_\alpha'\mathbf{I} + \sum_j \left(\lambda_P' + \gamma_j'^{\text{cst}} - \gamma_j'^{\text{ext}}\right)\frac{\mathbf{a}_j \otimes \mathbf{a}_j}{L_j'}\right]\Big/\sum_\alpha A_\alpha', \qquad \text{Eq. (vi-1)}$$

according to earlier studies [53–55], and using its diagonal element $\sigma_{\text{tube}}^{\text{long}}$ and $\sigma_{\text{tube}}^{\text{circ}}$, we

introduced

$$\rho_{\text{tube}} = \sigma_{\text{tube}}^{\text{long}} - \sigma_{\text{tube}}^{\text{circ}} \qquad \text{Eq. (vi-2)}$$

as the tube tension anisotropy. $\sigma_{\text{tube}}^{\text{long}}$ and $\sigma_{\text{tube}}^{\text{circ}}$ each correspond to the tube longitudinal

tension and the tube circumferential tension appeared in Figure S1.

(vii) Determination of Parameter Values (Fig. S1 and Fig. S4)

Parameter values used in this study were (i) $\lambda_P'=10^{-2}$, $\lambda_B'=10^{0.75}$, (ii) $T^{\text{cycle}}=10^{3.75}$, (iii)

$\rho* = 0.1$, $\gamma'^{\text{cst}} = 1$, and $\tau'^{\text{cst}} = 0.167$ unless otherwise noted. As for (i), we determined the

values, at which there were less variations at $t' = 1000$ from the initial configuration (Fig.

S1B and S1B'). Note that no proliferation was implemented for this analysis. As for (ii), we

examined the parameter dependence of cell proliferation rate $1/T^{\text{cycle}}$ to the morphological

and the mechanical quantities at $\tau' = 1$ (Fig. S1C). The results show that the cell division

orientation less affected to the tube shape (length and diameter) in faster cell proliferation

rate. In addition, smooth surface of tube was not maintained if the cell proliferation rate was

larger than $10^{-3.5}$ (see curvedness in Fig. S1C). This is because the cell proliferation rate is

faster than relaxation time of tube in the dynamics. We determined as $T^{\text{cycle}} = 10^{3.75}$. As for

(iii), we examined the parameter dependence of $\rho*$, γ'^{cst}, and τ'^{cst} to the tube shape and

its variance at $\tau' = 1$ (Fig. S4B). We first determined $\rho* = 0.1$ because the coefficient of

variation (CV) became larger at $\rho* = 0.08$ and there were less differences of the mean

values to the variation of γ'^{cst} at $\rho* = 0.125$. Then, we determined $\gamma'^{\text{cst}} = 1$ because this

point is at a boundary for both the mean and the CV. Finally, as for τ'^{cst}, we arbitrarily set

the value because there were less differences in any values.

(IV) Others

(i) Statistical Hypothesis Testing

Exact information of the statistical tests, sample sizes, test statistics, and *P*-values were

described in the Supplemental Information. Statistical hypothesis testing was performed

according to [56]. *P*-values of less than 0.05 were considered to be statistically significant in

two-tailed tests, and were classified as 4 categories; * (*P*<0.05), ** (*P*<0.01), *** (*P*<0.001),

and n.s. (not significant, i.e., $P \geq 0.05$).

(ii) Replicates and Sample Number

All replicates in this study are biological replicates. The *n* in the figure legends indicates the

number of samples used for each analysis as described below.

Figure 1B′: *n* indicates the number of epididymides collected from 3 pregnant mice.

Figure 1D′: *n* indicates the number of cells randomly chosen from 3 epididymides collected

from a pregnant mouse.

Figure 2D and E: *n* indicates the number of cells sampled from 6 epididymides collected

from a pregnant mouse.

Figure 3C-F′: *n* indicates the number of apical edges sampled from 8 epididymides collected

from 2 pregnant mice.

Figure 3H and H′: *n* indicates the number of epididymides collected from 3 pregnant mice.

Figure J′ and J′′: *n* indicates the number of cells randomly chosen from 3 epididymides

collected from a pregnant mouse.

Figure 5E: *n* indicates the number of a group of cells sampled from 4 epididymides each

collected from 4 pregnant mice.

Figure 7D-D′′: *n* indicates the number of cells randomly chosen from 10 epididymides

collected from 3 pregnant mice.

Figure 7E′ and E′′: *n* indicates the number of pregnant mice.

(iii) Graph

Except for boxplot, we used mean as representative values, and error bars represent standard

deviation. Boxplots were drawn using R (GNU project).

(iv) Software

For digital image processing, we used MATLAB (MathWorks) and Image J (National

Institute of Health). For graphics, we used MATLAB (MathWorks), R (GNU project),

Gnuplot (Free Software), and Paraview (Kitware). For statistical analysis, we used MATLAB

(MathWorks) and R (GNU project).

Additional files

Additional files include full information of statistical hypothesis testing, and supplementary

five figures.

Author Contributions

T.H. performed experiments and simulations, and analyzed all data; T.H. and T.A. wrote the

manuscript.

Acknowledgements

This work was supported by the Platform Project for Supporting in Drug Discovery and Life
Science Research (Platform for Dynamic Approaches to Living System) from the MEXT and
the AMED, and by the JSPS KAKENHI grant 15K18541. We would like to thank T.
Furukawa of Osaka Univ. for kindly providing the Pax2-Cre mice.

Competing Interests

883 The authors declare that no competing interests exist.

References

[1] Guillot C, Lecuit T. Mechanics of epithelial tissue homeostasis and morphogenesis.
Science 2013;340:1185–9. doi:10.1126/science.1235249.

[2] Mammoto T, Mammoto A, Ingber DE. Mechanobiology and Developmental Control.
Annu Rev Cell Dev Biol 2013;29:27–61.
doi:10.1146/annurev-cellbio-101512-122340.

[3] Eisenhoffer GT, Rosenblatt J. Bringing balance by force: Live cell extrusion controls
epithelial cell numbers. Trends Cell Biol 2013;23:185–92.
doi:10.1016/j.tcb.2012.11.006.

[4] Hufnagel L, Teleman AA, Rouault H, Cohen SM, Shraiman BI. On the mechanism of
wing size determination in fly development. Proc Natl Acad Sci U S A
2007;104:3835–40. doi:10.1073/pnas.0607134104.

[5] Shraiman BI. Mechanical feedback as a possible regulator of tissue growth. Proc Natl
Acad Sci U S A 2005;102:3318–23. doi:10.1073/pnas.0404782102.

[6] Karner CM, Chirumamilla R, Aoki S, Igarashi P, Wallingford JB, Carroll TJ. Wnt9b
signaling regulates planar cell polarity and kidney tubule morphogenesis. Nat Genet
2009;41:793–9. doi:10.1038/ng.400.

[7] Hirashima T. Pattern Formation of an Epithelial Tubule by Mechanical Instability
during Epididymal Development. Cell Rep 2014;9:866–73.
doi:10.1016/j.celrep.2014.09.041.

[8] Andrew DJ, Ewald AJ. Morphogenesis of epithelial tubes: Insights into tube formation,
elongation, and elaboration. Dev Biol 2010;341:34–55.
doi:10.1016/j.ydbio.2009.09.024.

[9] Joseph A, Yao H, Hinton BT. Development and morphogenesis of the
Wolffian/epididymal duct, more twists and turns. Dev Biol 2009;325:6–14.
doi:10.1016/j.ydbio.2008.10.012.

[10] Watanabe T, Hosoya H, Yonemura S. Regulation of Myosin II Dynamics by
Phosphorylation and Dephosphorylation of Its Light Chain in Epithelial Cells. Mol
Biol Cell 2007;18:605–16. doi:10.1091/mbc.E06-07-0590.

[11] Kovács M, Tóth J, Hetényi C, Málnási-Csizmadia A, Seller JR. Mechanism of
blebbistatin inhibition of myosin II. J Biol Chem 2004;279:35557–63.
doi:10.1074/jbc.M405319200.

[12] Peterson LJ, Rajfur Z, Maddox AS, Freel CD, Chen Y, Edlund M, et al. Simultaneous

stretching and contraction of stress fibers in vivo. Mol Biol Cell 2004;15:3497–508.

doi:10.1091/mbc.E03-09-0696.

[13]   Tang N, Marshall WF, McMahon M, Metzger RJ, Martin GR. Control of mitotic

spindle angle by the RAS-regulated ERK1/2 pathway determines lung tube shape.

Science (80-) 2011;333:342–5. doi:10.1126/science.1204831.

[14]   Fischer E, Legue E, Doyen A, Nato F, Nicolas J-F, Torres V, et al. Defective planar

cell polarity in polycystic kidney disease. Nat Genet 2006;38:21–3.

doi:10.1038/ng1701.

[15]   Fletcher AG, Osterfield M, Baker RE, Shvartsman SY. Vertex models of epithelial

morphogenesis. Biophys J 2014;106:2291–304. doi:10.1016/j.bpj.2013.11.4498.

[16]   Rauzi M, Verant P, Lecuit T, Lenne P-F. Nature and anisotropy of cortical forces

orienting Drosophila tissue morphogenesis. Nat Cell Biol 2008;10:1401–10.

doi:10.1038/ncb1798.

[17]   Nagai T, Honda H. A dynamic cell model for the formation of epithelial tissues. Philos

Mag Part B 2001;81:699–719. doi:10.1080/13642810108205772.

[18]   Walck-Shannon E, Hardin J. Cell intercalation from top to bottom. Nat Rev Mol Cell

Biol 2014;15:34–48. doi:10.1038/nrm3723.

[19]   Hirashima T, Adachi T. Procedures for the Quantification of Whole-Tissue

Immunofluorescence Images Obtained at Single-Cell Resolution during Murine

Tubular Organ Development. PLoS One 2015;10:e0135343.

doi:10.1371/journal.pone.0135343.

[20]   Stewart MP, Helenius J, Toyoda Y, Ramanathan SP, Muller DJ, Hyman A a.

Hydrostatic pressure and the actomyosin cortex drive mitotic cell rounding. Nature

2011;469:226–30. doi:10.1038/nature09642.

[21]   Abe T, Kiyonari H, Shioi G, Inoue KI, Nakao K, Aizawa S, et al. Establishment of

conditional reporter mouse lines at ROSA26 locus for live cell imaging. Genesis

2011;49:579–90. doi:10.1002/dvg.20753.

[22]   Ohyama T, Groves AK. Generation of Pax2-Cre Mice by Modification of a Pax2

Bacterial Artificial Chromosome. Genesis 2004;38:195–9. doi:10.1002/gene.20017.

[23]   Blanchard GB, Kabla AJ, Schultz NL, Butler LC, Sanson B, Gorfinkiel N, et al. Tissue

tectonics: morphogenetic strain rates, cell shape change and intercalation. Nat Methods

2009;6:458–64. doi:10.1038/nmeth.1327.

[24]   Savin T, Kurpios NA, Shyer AE, Florescu P, Liang H, Mahadevan L, et al. On the

growth and form of the gut. Nature 2011;476:57–62. doi:10.1038/nature10277.

[25]   Dong B, Hannezo E, Hayashi S. Balance between Apical Membrane Growth and

Luminal Matrix Resistance Determines Epithelial Tubule Shape. Cell Rep 2014.

doi:10.1016/j.celrep.2014.03.066.

[26]   Beloussov L V. Morphomechanics of Development. Springer; 2015.

doi:10.1007/978-3-319-13990-6.

[27]   Sasai Y. Cytosystems dynamics in self-organization of tissue architecture. Nature

2013;493:318–26. doi:10.1038/nature11859.

[28]   Nishimura T, Honda H, Takeichi M. Planar cell polarity links axes of spatial dynamics

in neural-tube closure. Cell 2012;149:1084–97. doi:10.1016/j.cell.2012.04.021.

[29]   Bosveld F, Bonnet I, Guirao B, Tlili S, Wang Z, Petitalot A, et al. Mechanical Control

of Morphogenesis by Fat/Dachsous/Four-Jointed Planar Cell Polarity Pathway.

Science (80-) 2012;336:724–7. doi:10.1126/science.1221071.

[30]   Xu B, Washington AM, Domeniconi RF, Ferreira Souza AC, Lu X, Sutherland A, et al.

Protein tyrosine kinase 7 is essential for tubular morphogenesis of the Wolffian duct.

Dev Biol 2016;412:219–233.

[31]   Yokomizo T, Yamada-Inagawa T, Yzaguirre AD, Chen MJ, Speck NA, Dzierzak E.

Whole-mount three-dimensional imaging of internally localized immunostained cells

within mouse embryos. Nat Protoc 2012;7:421–31. doi:Doi 10.1038/Nprot.2011.441.

[32]   Susaki EA, Tainaka K, Perrin D, Kishino F, Tawara T, Watanabe TM, et al.

Whole-brain imaging with single-cell resolution using chemical cocktails and

computational analysis. Cell 2014;157:726–39. doi:10.1016/j.cell.2014.03.042.

[33]   Kapur JN, Sahoo PK, Wong AKC. A new method for gray-level picture thresholding

using the entropy of the histogram. Comput Vision, Graph Image Process 1985;29:140.

doi:10.1016/S0734-189X(85)90156-2.

[34]   Kerschnitzki M, Kollmannsberger P, Burghammer M, Duda GN, Weinkamer R,

Wagermaier W, et al. Architecture of the osteocyte network correlates with bone

material quality. J Bone Miner Res 2013;28:1837–45. doi:10.1002/jbmr.1927.

[35]   Hannezo E, Prost J, Joanny J-F. Theory of epithelial sheet morphology in three

dimensions. Proc Natl Acad Sci 2014;111:27–32. doi:10.1073/pnas.1312076111.

[36]   Farhadifar R, Röper J-C, Aigouy B, Eaton S, Jülicher F. The influence of cell

mechanics, cell-cell interactions, and proliferation on epithelial packing. Curr Biol

2007;17:2095–104. doi:10.1016/j.cub.2007.11.049.

[37]   Bielmeier C, Alt S, Weichselberger V, La Fortezza M, Harz H, J??licher F, et al.

Interface Contractility between Differently Fated Cells Drives Cell Elimination and

Cyst Formation. Curr Biol 2016;26:563–74. doi:10.1016/j.cub.2015.12.063.

[38]   Mao Y, Tournier AL, Hoppe A, Kester L, Thompson BJ, Tapon N. Differential

proliferation rates generate patterns of mechanical tension that orient tissue growth.

EMBO J 2013;32:2790–803. doi:10.1038/emboj.2013.197.

[39]   Honda H, Motosugi N, Nagai T, Tanemura M, Hiiragi T. Computer simulation of

emerging asymmetry in the mouse blastocyst. Development 2008;135:1407–14.

doi:10.1242/dev.014555.

[40]   Trichas G, Smith AM, White N, Wilkins V, Watanabe T, Moore A, et al.

Multi-cellular rosettes in the mouse visceral endoderm facilitate the ordered migration

of anterior visceral endoderm cells. PLoS Biol 2012;10.

doi:10.1371/journal.pbio.1001256.

[41]   Aegerter-Wilmsen T, Heimlicher MB, Smith AC, de Reuille PB, Smith RS, Aegerter

CM, et al. Integrating force-sensing and signaling pathways in a model for the

regulation of wing imaginal disc size. J Cell Sci 2012;125:e1–e1.

doi:10.1242/jcs.120758.

[42]   Honda H, Eguchi G. How much does the cell boundary contract in a monolayered cell

sheet? J Theor Biol 1980;84:575–88. doi:10.1016/S0022-5193(80)80021-X.

[43]   Honda H, Tanemura M, Nagai T. A three-dimensional vertex dynamics cell model of

space-filling polyhedra simulating cell behavior in a cell aggregate. J Theor Biol

2004;226:439–53. doi:10.1016/j.jtbi.2003.10.001.

[44]   Fletcher AG, Osborne JM, Maini PK, Gavaghan DJ. Implementing vertex dynamics

models of cell populations in biology within a consistent computational framework.

Prog Biophys Mol Biol 2013;113:299–326. doi:10.1016/j.pbiomolbio.2013.09.003.

[45]   Osterfield M, Du X, Schüpbach T, Wieschaus E, Shvartsman SY. Three-Dimensional

Epithelial Morphogenesis in the Developing Drosophila Egg. Dev Cell 2013;24:400–

10. doi:10.1016/j.devcel.2013.01.017.

[46]   Du X, Osterfield M, Shvartsman SY. Computational analysis of three-dimensional

epithelial morphogenesis using vertex models. Phys Biol 2014;11:66007.

doi:10.1088/1478-3975/11/6/066007.

[47]   Ishimoto Y, Morishita Y. Bubbly vertex dynamics: A dynamical and geometrical

model for epithelial tissues with curved cell shapes. Phys Rev E - Stat Nonlinear, Soft

Matter Phys 2014;90. doi:10.1103/PhysRevE.90.052711.

[48]   Okuda S, Inoue Y, Eiraku M, Adachi T, Sasai Y. Vertex dynamics simulations of

viscosity-dependent deformation during tissue morphogenesis. Biomech Model

1020   Mechanobiol 2014;14:413–25. doi:10.1007/s10237-014-0613-5.

1021 [49] Collinet C, Rauzi M, Lenne P, Lecuit T. Local and tissue-scale forces drive oriented

1022   junction growth during tissue extension. Nat Cell Biol 2015;17:1247–58.

1023   doi:10.1038/ncb3226.

1024 [50] Meyer M, Desbrun M, Schr P, Barr AH. Discrete Differential-Geometry Operators for

1025   Triangulated 2-Manifolds. Vis Math III 2003:113–34. doi:10.1016/j.cagd.2007.07.005.

1026 [51] Kantor Y, Nelson DR. Crumpling transition in polymerized membranes. Phys Rev Lett

1027   1987;58:2774–7. doi:10.1103/PhysRevLett.58.2774.

1028 [52] Seung HS, Nelson DR. Defects in flexible membranes with crystalline order. Phys Rev

1029   A 1988;38:1005–18. doi:10.1103/PhysRevA.38.1005.

1030 [53] Batchelor GK. The stress system in a suspension of force-free particles. J   Fluid

1031   Mech 1970;41:545–70. doi:10.1017/S0022112070000745.

1032 [54] Ishihara S, Sugimura K. Bayesian inference of force dynamics during morphogenesis.

1033   J Theor Biol 2012;313:201–11. doi:Doi 10.1016/J.Jtbi.2012.08.017.

1034 [55] Sugimura K, Ishihara S. The mechanical anisotropy in a tissue promotes ordering in

1035   hexagonal cell packing. Development 2013;140:4091–101. doi:10.1242/dev.094060.

1036 [56] Zar JH. Biostatistical Analysis, 5th Edition. 2014.

1037

1038

Figure Legends

1043 Figure 1 Actomyosin-mediated apical constriction is required for the maintenance of
1044 the tubule radial size.

(A) Maximum intensity projection (MIP) of immunofluorescence images for the pMRLC in
epididymis at E16.5. Magenta dotted lines represent the basal of epididymal tubule. Scale
bar: 50 μm. (B and B′) Morphological change of tubules in the inhibitor assays. (B) MIP of
immunofluorescence images for the E-cadherin. The head region of epididymis to the tail
from left to right. Scale bar: 200 μm. (B′) The tubule diameter along the tubule. *n*=8. (C-C′′)
Morphological change of tubule cells in the inhibitor assays. (C) Immunofluorescence images
for the E-cadherin at sections of the tubule. Yellow/magenta lines represent apical/basal of
the cells. Scale bar: 10 μm. (D) Schematic for the cell morphology quantities; the aspect ratio
is defined as the squared cell area over the cell height and the cell volume is defined as the
cell area times the cell height. (D′) The cell morphology quantities versus the tubule diameter
in the control and blebbistatin treatment. *n*=20.

Figure 2 Cell division orientation determines the mechanical stress and tubule radial

size.

(A) MIP of immunostained images for the pHH3 (mitotic cells, red) and γ-tubulin (MTOCs,

white). (A′ and A′′) Magnified view of the dotted square in (A), displaying the two cells

divide in different orientations. Scale bars: 10 μm. (B) Measurement of cell division

orientation in the tubule. Arrows indicate the center position of the mitotic cell. Using the

three angles α_1, α_2, and α_3, the observation coordinate O is transformed into the local

coordinate $O′$ in Fig. 2C. (C) Illustration of the local coordinates. (D) The angle distributions

for the cell division orientation. Colors in the θ distribution represent samples, of which φ

ranges from 0-40 degrees (orange) and from 40-90 degrees (gray). (E) Scatter plot for the

distance between MTOCs and the angle θ. The sample number is the same as in (D). (F-F′′)

Model simulations in different cell division orientations. (F) The vertex dynamics model

simulation of proliferating tubes. The cell division orientation determines the mechanical

stress in the cells and tubule radial size. The color represents the cell tension anisotropy. (F′

and F′′) Time course of morphological and mechanical quantities along the normalized

simulation time $\tau′$. See the Materials and Methods (III-v) and (III-vi) for those quantities

and Materials and Methods (III-iii) for the definition of $\tau′$. n=15.

1077

Figure 3 Polarized apical constriction is coupled to cell division.

(A) Digital processing from immunofluorescence images for the ZO-1 detects nodes and junctions of apical surfaces. (B) Illustration of the junction angle ϑ and length ℓ. ϑ measures the angle from the longitudinal axis of tubules. (C) Histogram of pMRLC intensity (gray) fitted with a log-normal distribution (green). The small window displays the tail of distribution, showing the well-fitting. n=22,037. (D) MIP of co-immunostained images of pMRLC (green) and ZO-1 (blue). Arrows indicate pMRLC localization at circumferential junctions. Scale bar: 10 μm. (E-F′) Relationship between the pMRLC intensity and the junction angle/length. The log-scale histograms are shown in three divided groups. Black arrows represent the mean intensity in the longitudinal/short group and white arrows represent that in the circumferential/long group. (G) MIP of pMRLC immunostaining at the whole-tissue scale. Scale bar: 100 μm. (H) Spatial profile of pMRLC intensity (blue) and the number of mitotic cells (orange) along the tubule. n=12. (H′) The pMRLC intensity versus the mitotic cell number. The color indicates position of data in the tubule. (I and I′) Myosin activity quantification by immunoblotting. (J-J″) Apical junction morphology visualized by

1093　　　staining for the F-actin and E-cadherin. n=200. Scale bar: 10 μm.

1094

1095

1096

Figure 4 Computational analysis of cell stress influenced by cell division in tubes.

(A) The schematics for glossary. The cell tension anisotropy was calculated in cells

neighboring to the divided cells. (B and B′) Histograms of cell tension anisotropy against the

cell divisions. (B) The explanatory illustration for the histograms. Red indicates the case of

circumferential cell division and blue indicates the case of longitudinal cell division. (B′)

Histograms of the cell tension anisotropy in various cases of the relative angle to the cell

division axis (vertical) and in those of the elapsed time of neighbors' cell division $\tau_\alpha'^{\text{cycle}}$

(horizontal). (C-C′′) Histograms of the elapsed time after the cell division $\tau_\alpha'^{\text{cycle}}$ when the

cell tension anisotropy is over various thresholds (ρ^* =0.039, 0.1, and 0.2) in neighboring

cells on the circumferential axis. The tail of histograms is magnified in the small windows.

1107

1108

1109

1110

1111

1112

Figure 5 Live imaging revealed that the oriented cell division occurs following to the

circumferential cell division.

(A) The dimensions of epididymis. The rendering image from the immunofluorescence for

the ZO-1 (magenta) (left) and its schematics (right). (B) The schematics for the

incubator-integrated multiphoton excitation microscopy system. (C) Snapshots from live

imaging data in the case of the circumferential and longitudinal cell divisions. Time origin is

at the timing of cytokinesis completion. Asterisks represent daughters of divided cell, one of

which is colored artificially in green. Some cells are also colored for easy visualization. The

scale and sign for cell intercalation strain rate is shown in the bottom. Scale bar: 10 μm. (D)

Schematics for the CITE. (E) The time series data for the case of circumferential cell division

(red, *n*=20) and that of the longitudinal division (blue, *n*=23). Time zone in which the mean

value is not statistically zero is highlighted.

1125

1126

1127

Figure 6 Model simulations suggest that the mechano-responsive cell behavior explains

the maintenance of tubule radial size.

(A) Schematics for the mechano-responsive regime. In this regime, the circumferential

junction constriction starts when the cell tension anisotropy becomes larger than the threshold

ρ^* due to the pressure from circumferentially-neighboring cells. (B) Generated tubes in

different regimes: no constriction, mechano-responsive, and scheduled. Red color in the tubes

represents cells constricting the circumferential junction and blue represents cells that have

completed the cell intercalation. (C) The ratio of constricting cells to the total cells over the

normalized simulation time τ' in the mechano-responsive regime: data in each simulation

(gray), the mean value (black) and fitting with the logistic function (blue). n=10. (D and D′)

Resulting morphological quantities of tubes, the tube diameter and the tube curvedness, in

different regimes. See the Materials and Methods (III-vi) for the definition of those

morphological quantities. n=10.

1141

1142

1143

1144
1145

Figure 7 Mechanical perturbation assays uncovered the polarized mechano-response
system for the maintenance of tubule radial size.

(A and B) Schematics for the uniaxial compression assays. (C) Morphology of the tubule
cells following the mechanical treatments. The cell shape was recognized by
immunofluorescence for the E-cadherin. Scale bar: 5 μm. (D-D´´) The aspect ratio and the
major axis angle for each treatment: control (gray), lateral compression (red), and
longitudinal compression (blue). The data are shown as scatter plot and histogram in (D) and
as boxplot in (D´ and D´´). *n*=500. (E-E´´) Immunoblotting and quantification of myosin
activity and ROCK1 for the mechanical treatments. The tubule cells activate myosin as a
cellular response to lateral compression. (E´ and E´´) The data are normalized to the mean
value in the control. *n*=9. (F) Model for the maintenance of tubule radial size.

1157

1158 # Additional files (Supplemental Information)

1159 Anisotropic Cellular Mechanoresponse for Radial Size Maintenance of

1160 Developing Epithelial Tubes

1161

1162 Tsuyoshi Hirashima and Taiji Adachi

1163

1164

1165 **This file includes:**

1166

1167 Statistical Hypothesis Testing

1168 Supplementary Figures S1 to S5

1169 Still images and captions for Movies 1 to 3

1170

 Statistical Hypothesis Testing

1172 The statistical tests, sample sizes, test statistics, and *P*-values were described below. *P*-values

1173 of less than 0.05 were considered to be statistically significant in two-tailed tests, and were

1174 classified as 4 categories; * (*P*<0.05), ** (*P*<0.01), *** (*P*<0.001), and n.s. (not significant,

1175 i.e., $P \geq 0.05$).

1176

1177 ➤ Figure 2D, φ at E15.5

1178 Note: The range of data was transformed to 0-360º for circular distributions.

1179 H_0: The population of φ at E15.5 is uniformly distributed around the circle.

1180 H_A: The population of φ at E15.5 is not uniformly distributed around the circle.

1181 The Rayleigh test. *n*=277, Test statistic: *Z*=35.51, *P*<0.001; Reject H_0.

1182

➤ Figure 2D, θ at E15.5

Note: The range of data was transformed to 0-360º for circular distributions.

H_0: The population of θ at E15.5 is uniformly distributed around the circle.

H_A: The population of θ at E15.5 is not uniformly distributed around the circle.

The Rayleigh test. *n*=197 (0-40º), Test statistic: *Z*=0.40, *P*=0.67; Do not reject H_0.

The Rayleigh test. *n*=277 (0-90º), Test statistic: *Z*=0.25, *P*=0.78; Do not reject H_0.

1189

1190 ➤ Figure 2D, φ at E16.5

1191 Note: The range of data was transformed to 0-360º for circular distributions.

1192 H_0: The population of φ at E16.5 is uniformly distributed around the circle.

1193 H_A: The population of φ at E16.5 is not uniformly distributed around the circle.

1194 The Rayleigh test. *n*=358, Test statistic: *Z*=27.32, *P*<0.001; Reject H_0.

1195

➤ Figure 2D, θ at E16.5

Note: The range of data was transformed to 0-360º for circular distributions.

H_0: The population of θ at E16.5 is uniformly distributed around the circle.

H_A: The population of θ at E16.5 is not uniformly distributed around the circle.

The Rayleigh test. *n*=240 (0-40º), Test statistic: *Z*=1.95, *P*=0.14; Do not reject H_0.

The Rayleigh test. *n*=358 (0-90º), Test statistic: *Z*=0.57, *P*=0.56; Do not reject H_0.

1202

➤ Figure 3E'

Note: The data were transformed on logarithmic scale.

H_0: The mean intensities of pMRLC among the three groups of junction angle are all equal.

H_A: The mean intensities of pMRLC among the three groups of junction angle are not all

equal.

One-way analysis of variance. $n=6,259$ (Longitudinal), $n=8,457$ (Intermediate), $n=7,323$

(Circumferential), Test statistic: $F=165.47$, $P<0.001$; Reject H_0.

Multiple comparisons using the Tukey-Kramer Test with unequal sample sizes: $P<0.001$

(Long. vs. Intm.), $P<0.001$ (Long. vs. Circ.), and $P<0.001$ (Intm. vs. Circ.).

1212

➢ Figure 3F'

Note: The data were transformed on logarithmic scale.

H_0: The mean intensities of pMRLC among the three groups of junction length are all equal.

H_A: The mean intensities of pMRLC among the three groups of junction length are not all

equal.

One-way analysis of variance. $n=8,691$ (Short), $n=8,749$ (Middle), $n=3,719$ (Long), Test

statistic: $F=306.65$, $P<0.001$; Reject H_0.

Multiple comparisons using the Tukey-Kramer test with unequal sample sizes: $P<0.001$

(Short. vs. Middle), $P<0.001$ (Short vs. Long), and $P<0.001$ (Middle vs. Long).

1222

➢ Figure 3J'

H_0: The circumferential edge length is the same in all three regions of epididymal tubule.

H_A: The circumferential edge length is not the same in all three regions of epididymal tubule.

The Kruskal-Wallis test, $n=200$ for all groups, Test statistic: $H=14.84$, $P<0.001$; Reject H_0.

Steel-Dwass multiple comparisons: $P=0.016$ (Head vs. Body), $P=0.001$ (Head vs. Tail),

$P=0.61$ (Body vs. Tail).

1229

➢ Figure 3J"

H_0: The apical surface area is the same in all three regions of epididymal tubule.

H_A: The apical surface area is not the same in all three regions of epididymal tubule.

The Kruskal-Wallis test, $n=200$ for all groups, Test statistic: $H=89.58$, $P<0.001$; Reject H_0.

Steel-Dwass multiple comparisons: $P<0.001$ (Head vs. Body), $P<0.001$ (Head vs. Tail),

$P=0.71$ (Body vs. Tail).

1236

➢ Figure 5C

H_0: The mean value of CITE in a certain time zone from the cell division is equal to zero.

H_A: The mean value of CITE in a certain time zone from the cell division is not equal to zero.

t test, $n=20$ in the case of circumferential cell division and $n=23$ in that of longitudinal one,

Test statistic: absolute value of t: $|t|$

[Circ., 10-20 min]: $|t|=0.97$, $P=0.34$; Do not reject H_0

[Circ., 20-30 min]: $|t|=1.01$, $P=0.32$; Do not reject H_0

[Circ., 30-40 min]: $|t|=1.12$, $P=0.29$; Do not reject H_0

[Circ., 40-50 min]: $|t|=1.06$, $P=0.30$; Do not reject H_0

[Circ., 50-60 min]: $|t|=2.37$, $P=0.03$; Reject H_0

[Circ., 60-70 min]: $|t|=1.50$, $P=0.15$; Do not reject H_0

[Circ., 70-80 min]: $|t|=1.09$, $P=0.29$; Do not reject H_0

[Circ., 80-90 min]: $|t|=0.54$, $P=0.59$; Do not reject H_0

[Circ., 90-100 min]: $|t|=1.75$, $P=0.10$; Do not reject H_0

[Circ., 100-110 min]: $|t|=0.04$, $P=0.97$; Do not reject H_0

[Circ., 110-120 min]: $|t|=0.35$, $P=0.73$; Do not reject H_0

[Long., 10-20 min]: $|t|=2.02$, $P=0.06$; Do not reject H_0

[Long., 20-30 min]: $|t|=0.51$, $P=0.61$; Do not reject H_0

[Long., 30-40 min]: $|t|=1.05$, $P=0.30$; Do not reject H_0

[Long., 40-50 min]: $|t|=0.98$, $P=0.34$; Do not reject H_0

[Long., 50-60 min]: $|t|=0.21$, $P=0.83$; Do not reject H_0

[Long., 60-70 min]: $|t|=1.03$, $P=0.31$; Do not reject H_0

[Long., 70-80 min]: $|t|=1.08$, $P=0.29$; Do not reject H_0

[Long., 80-90 min]: $|t|=0.31$, $P=0.76$; Do not reject H_0

[Long., 90-100 min]: $|t|=0.03$, $P=0.97$; Do not reject H_0

[Long., 100-110 min]: $|t|=1.75$, $P=0.09$; Do not reject H_0

[Long., 110-120 min]: $|t|=0.24$, $P=0.81$; Do not reject H_0

1264

➢ Figure 6D

H_0: The mean values of tube diameter among the three groups of regime are all equal.

H_A: The mean values of tube diameter among the three groups of regime are not all equal.

One-way analysis of variance, $n=10$ for all groups, $F=791.10$, $P<0.001$; Reject H_0.

Tukey multiple comparisons: $P<0.001$ (No constriction vs. Mech-resp.), $P<0.001$ (No

constriction vs. Scheduled), and $P=0.002$ (Mech-resp vs. Scheduled).

1271

➢ Figure 6D'

H_0: The mean values of curvedness among the three groups of regime are all equal.

H_A: The mean values of curvedness among the three groups of regime are not all equal.

One-way analysis of variance, $n=10$ for all groups, $F=575.25$, $P<0.001$; Reject H_0.

Tukey multiple comparisons: $P<0.001$ (No constriction vs. Mech-resp.), $P<0.001$ (No

constriction vs. Scheduled), and $P<0.001$ (Mech-resp vs. Scheduled).

1278

➢ Figure 7D'

H_0: The aspect ratio of cells is the same in all three treatments of compression assay.

H_A: The aspect ratio of cells is not the same in all three treatments of compression assay.

The Kruskal-Wallis test, $n=500$ for all groups, Test statistic: $H=42.10$, $P<0.001$; Reject H_0.

Steel multiple comparisons: $P<0.001$ (Control vs. Lateral), $P=0.043$ (Control vs.

Longitudinal).

1285

➢ Figure 7D"

H_0: The angle of cells is the same in all three treatments of compression assay.

H_A: The angle of cells is not the same in all three treatments of compression assay.

The Kruskal-Wallis test, $n=500$ for all groups, Test statistic: $H=366.04$, $P<0.001$; Reject H_0.

Steel multiple comparisons: $P<0.001$ (Control vs. Lateral), $P<0.001$ (Control vs.

Longitudinal).

1292

➢ Figure 7E'

H_0: The ratio of pMRLC to tMRLC is the same in all three treatments of compression assay.

H_A: The ratio of pMRLC to tMRLC is not the same in all three treatments of compression

assay.

The Friedman test, $n=9$ for all groups, Test statistic: $Q=6$, $P=0.049$; Reject H_0.

Steel multiple comparisons: $P=0.039$ (Control vs. Lateral), $P=1.82$ (Control vs.

Longitudinal).

1300

➢ Figure 7E'

H_0: The ratio of ROCK1 to tMRLC is the same in all three treatments of compression assay.

H_A: The ratio of ROCK1 to tMRLC is not the same in all three treatments of compression

assay.

The Friedman test, $n=9$ for all groups, Test statistic: $Q=1.56$, $P=0.46$; Do not reject H_0.

Steel multiple comparisons: P=0.50 (Control vs. Lateral), P=0.60 (Control vs. Longitudinal).

1307

1308

Figure S1 Preparation for the simulation without anisotropic junction constriction,

corresponding to the Figure 2

(A-A") Schematics for the explanation of discrete curvature in the vertex model. See the

Materials and Methods (III-ii) for the notations. (B and B') Diagram of generated tube

morphology for parameters λ'_B and λ'_P, and relative variation of morphological quantities in

the absence of cell proliferation. Tubes are visualized in the four corners of parameter space

(filled circles). Open circles represent parameter values used in this study. The axes labels are

1316   shown in B. (C) Parameter dependence of morphological and mechanical quantities for cell

1317   division orientation and cell proliferation rate. Color represents the type of cell division

1318   orientation. (D and D') Time course of tube circumferential tension and that of tube

1319   longitudinal tension. See the Materials and Methods (III-vi) for the definition of tube

1320   circumferential/longitudinal tension.

1321

1322

Figure S2 Apical junction length and angle, corresponding to the Figure 3

Frequency map on junction length ℓ and angle θ (left). Relative frequency in each region of epididymal tubule (Head, Body, and Tail) and angle category (Longitudinal, Intermediate, and Circumferential) is shown as histograms (right). The red/blue arrow indicates median/mean. The axis label of histograms is represented in the bottom. The circumferential junction length is shorter than the longitudinal junction length in any regions. In addition, the circumferential junction length in the head region is shorter than that in the tail region.

1330

1331

1332

1333

Figure S3 Measurement of cell intercalation strain rate from the live imaging data,

corresponding to the Figure 5

(A) Illustration for target cells to calculate the cell intercalation strain rate. See the Materials

and Methods (II-viii) for the details. (B) Graph representations for the cell intercalation strain

rate. Orthogonal lines in the left graph represent principal strain rate. The amplitude and

angle deviation of the positive principal strain rate (blue) in the left are expressed in the right

angle-amplitude graph. (C) Time course of the cell intercalation strain rate as in the form of

angle-amplitude graph for each type of cell division orientation: circumferential cell division

and longitudinal one.

1349

Figure S4 Preparation for the simulations in the mechano-responsive regime,

corresponding to the Figure 6

(A) Schematics for the model. See the Materials and Methods (III-v) for the details. (B)

Parameter dependence of tube morphology. The referenced graph is shown in the upper right.

CV: coefficient of variance. See the Materials and Methods (III-vii) for determining the

parameter values.

1356

1357

1358

1359

Figure S5 Spatial distribution of polarity proteins Van like protein (VANGL) 1 and 2,

corresponding to the Discussion

(A) Immunofluorescence for VANGL1 and VANGL2 demonstrates that their

co-localizations are on circumferential junctions of epididymal tubule. (B-C') Relation of

junction angle v.s. the fluorescence intensity, and that of junction length v.s. the intensity as

with the form of Fig. 3E-F'. These graphs indicate the VANGL localization on the apical

junctions of tubule shows significant polarization.

1367

1368

1369 **Movie 1 Model Simulation in Different Cell Division Orientation**

1370

1371

1372
1373

1374 **Movie 2 Live Imaging for Multicellular Behaviors against Cell Division**

1375

1376

No constriction

Mechano-responsive

Scheduled

1377
1378

Movie 3 Model Simulation in Different Regimes

A pMRLC
Apical
Basal
50 μm

B E-cadherin
DMSO (Control)
Head   Tail
Blebbistatin 30 μM
Calyculin A 20 nM
200 μm

B' n=8

Tubule diameter [μm]
Head   Tail

— Control
— Blebbistatin (30 μM)
— Y27632 (10 μM)
— Calyculin A (20 nM)

C
DMSO (Control)
Blebbistatin 30 μM
Calyculin A 20 nM
10 μm   E-cadherin
Apical   Basal

D
Height
Area

Aspect ratio = $\frac{Area^{1/2}}{Height}$

Cell volume = Area × Height

D' n=20

Aspect ratio

Cell volume [μm³]
1.2 (×10³)
0.8
0.4
0

■ ● Control
■ ● Blebbistatin

40   90   140   190   240
Tubule diameter [μm]

A

Longitudinal axis

MTOC
Mitotic cells

10 μm

A'

Cell 1
Cell 2

10 μm

A"

Orientation of cell division

Cell 1

Cell 2

Longitudinal axis

B

α_1

α_2

α_2

z

x
O
y

C

Surface normal

Longitudinal

z'

x'
O
y'

MTOC

Circumferential

MTOC

Cell division axis

z'

φ

O
y'

x'

θ

MTOC

D

E15.5

$\varphi=[0°, 40°]$: n=197
$\varphi=[0°, 90°]$: n=277

90°

φ

0-40°

0 20 40 60 80

90°

θ

■ $\varphi=[0°, 40°)$
□ $\varphi=[40°, 90°]$

0 10 20 30 40
# of samples

E16.5

$\varphi=[0°, 40°]$: n=240
$\varphi=[0°, 90°]$: n=358

90°

φ

0-40°

0 20 40 60 80

90°

θ

■ $\varphi=[0°, 40°)$
□ $\varphi=[40°, 90°]$

0 10 20 30 40 50
# of samples

E

● $\varphi=[0°, 40°)$
● $\varphi=[40°, 90°]$

E15.5

θ 60

E16.5

θ 60

0 2 4 6 8 10 12 14
Distance between MTOCs [μm]

F

Circumferential division

Non-biased division

Longitudinal division

Cell tension anisotropy
-0.2   0.2

F'

n=15

Circumferential
Non-biased
Longitudinal

Tube diameter 6

F"

0.03

Tube tension anisotropy

-0.03

0   Normalized time τ'   1

A ZO-1

3D image processing

● Node ▬ Junction

B ● Node ▬ Junction
Length: ℓ
Longitudinal axis
Angle: ϑ

C ■ Data (n=22,037) ― Fitting (log-normal)
Frequency
1200
700
0.13 (log-scale)
0   pMRLC intensity   0.8

D pMRLC ZO-1
10 µm

E Long. | Intm. | Circ.
pMRLC Intensity [au]
0.8
0   Junction angle ϑ [rad]   π/2

E' Long. n=6,259 | Intm. n=8,457 | Circ. n=7,323
pMRLC intensity (log-scale) [au]
0.6
0.03
0 600   0 600   0 600

F Short⊢Middle⟶Long
pMRLC Intensity [au]
0.8
0   Junction length ℓ [µm]   10

F' Short n=8,691 | Middle n=8,749 | Long n=3,719
pMRLC intensity (log-scale) [au]
0.6
0.03
0 600   0 600   0 400

G Head   Body   Tail
pMRLC intensity Low ― High
100 µm

H E15.5   n=12
pMRLC intensity [au] 0.8
# of mitotic cells 14
Head   Tail

E16.5   n=12
pMRLC intensity [au] 0.7
# of mitotic cells 25
Head   Tail

H' Head ― Tail
E15.5
pMRLC intensity [au] 0.7
1.0 # of mitotic cells 2.6

E16.5
pMRLC intensity [au] 0.5
1.5 # of mitotic cells 5.5

I E15.5   E16.5
Head-side Tail-side Head-side Tail-side
pMRLC
tMRLC
GAPDH

I' E15.5   E16.5
pMRLC/tMRLC
1.2   1.2
Head-side Tail-side Head-side Tail-side

J Head   Tail
10 µm
Actin E-cadherin

J' n=200
Circ. junction length [µm]
10
**
* n.s
Head Body Tail

J" n=200
Apical surface area [µm²]
20
***
*** n.s
Head Body Tail

A
Relative angle to division axis
Division axis
Divided cells

B
# of samples
450
0
n=3000
Long.
Circum.
-0.3   0.3
Cell tension anisotropy

B'
Relative angle to division axis
0-10
10-20
20-30
30-40
40-50
50-60
60-70
70-80
80-90

$10^0/10^{3.75}$ $10^1/10^{3.75}$ $10^2/10^{3.75}$ $10^3/10^{3.75}$

Time from cell division τ'^{cycle}_α

C ρ^*=0.039   n=14,924
Relative frequency
0.7
0
0.05
0
0   1
Time from cell division τ'^{cycle}_α

C' ρ^*=0.1   n=9,972
Relative frequency
0.7
0
0.05
0
0   1
Time from cell division τ'^{cycle}_α

C" ρ^*=0.2   n=1,316
Relative frequency
0.7
0
0.05
0
0   1
Time from cell division τ'^{cycle}_α

A

z

100 μm

0 μm

Epididymal tubule

Capsule

B

CO₂ gas incubator

Epididymis with testis at E15.5

Culture insert

25x

Water supply system

Multiphoton excitation

C

Circumferential division

circum.

long.

10 μm

0 min   40 min   80 min   120 min

Cell intercalation strain rate [min⁻¹]

10 min-   30 min-   50 min-   70 min-   90 min-   110 min-

Longitudinal division

0 min   40 min   80 min   120 min

Cell intercalation strain rate [min⁻¹]

10 min-   30 min-   50 min-   70 min-   90 min-   110 min-

Cell intercalation strain rate [min⁻¹]

0.01

-0.01

-0.01   0.01

positive
negative

D

Cell Intercalation rate for Tube Elongation (CITE)

Positive

circum.

long.

Negative

E

Circumferential division (n=20)
Longitudinal division (n=23)

0.015

0.01

0.005

CITE [min⁻¹]

0.005

0.01

-0.015

0   20-30   40-50   60-70   80-90   100-110

Time from cell division [min]

A *Mechano-responsive regime*

Pressure $\rho_\alpha \geq \rho^*$ Junctional constriction Cell intercalation

B No constriction

Mechano-responsive

Scheduled

Constrict-ing cells
Intercalat-ed cells

C
Ratio of constricting cells

n=10

0.16

0.08

0
0   1
Normalized time τ'

D
Tube diameter

n=10

***
***   **

No constriction
Mech.-resp.
Scheduled

D'
Curvedness

n=10

0.4

***
***   ***

No constriction
Mech.-resp.
Scheduled

A
PDMS
Sample
Gel
Uniaxial strain

B
Lateral compression
Longitudinal compression
External force

C
Untreated (Ctrl)
Lateral (Latr)
Longitudinal (Long)
5 μm
E-cadherin

D
Frequency
70
n=500
Ctrl
Latr
Long
π/2
Angle [rad]
0
1
Aspect ratio
3
0
60
Frequency

D'
n=500
Aspect ratio
3
1
***
*
Ctrl   Ltrl   Long

D''
n=500
Angle [rad]
π/2
***
***
Ctrl   Ltrl   Long

E
Ctrl   Latr   Long
ROCK1
pMRLC
tMRLC
GAPDH

E'
n=9
pMRLC/tMRLC
n.s.
*
1
Ctrl   Latr   Long

E''
n=9
ROCK1/tMRLC
n.s.
n.s.
1
Ctrl   Latr   Long

F
Spontaneous maintenance of tube radial size
Circumferential
Tissue   Cell
1. Mechano sensing
2. Response
Constriction
~ 60 min
3. Cell intercalation