Genetic screen suggests an alternative mechanism for azide-mediated inhibition of SecA

Running title: Mechanism of azide inhibition of SecA

Rachael Chandler[†a] and Mohammed Jamshad[†,a,b], Jack Yule[a], Ashley Robinson[a], Farhana Alam[a],

Karl A. Dunne[a], Naomi Nabi[a,c], Ian Henderson[a] and Damon Huber[a,*]

[a]Institute for Microbiology and Infection, University of Birmingham, Birmingham, United

Kingdom

[b]Current address: Department of Cancer Sciences, University of Birmingham, Birmingham, UK

[c]Current address: Department of Biological Sciences, University of Huddersfield, Huddersfield,

UK

*Address correspondence to: Damon Huber, d.huber@bham.ac.uk

[†]R. C. and M. J. contributed equally to this work.

Keywords: Sec, SecA, azide, protein translocation, TraDIS, phospholipid binding, metal binding

Abstract

Sodium azide prevents bacterial growth by inhibiting the activity of SecA, which is required

for translocation of proteins across the cytoplasmic membrane. Azide inhibits ATP turnover *in*

vitro, but its mechanism of action *in vivo* is unclear. To investigate how azide inhibits SecA in

cells, we used transposon directed insertion-site sequencing (TraDIS) to screen a library of

transposon insertion mutants for mutations that affect the susceptibility of *E. coli* to azide.

Insertions disrupting components of the Sec machinery generally increased susceptibility to

azide, but insertions truncating the C-terminal tail (CTT) of SecA decreased susceptibility of *E.*

coli to azide. Treatment of cells with azide caused increased aggregation of the CTT, suggesting

that azide disrupts its structure. Analysis of the metal-ion content of the CTT indicated that SecA

binds to iron and the azide disrupts the interaction of the CTT with iron. Azide also disrupted

binding of SecA to membrane phospholipids, as did alanine substitutions in the metal-

coordinating amino acids. Furthermore, treating purified phospholipid-bound SecA with azide in

the absence of added nucleotide disrupted binding of SecA to phospholipids. Our results suggest

that azide does not inhibit SecA by inhibiting the rate of ATP turnover *in vivo*. Rather, azide

inhibits SecA by causing it to "backtrack" from the ADP-bound to the ATP-bound conformation,

which disrupts the interaction of SecA with the cytoplasmic membrane.

Significance statement

SecA is a bacterial ATPase that is required for the translocation of a subset of secreted proteins

across the cytoplasmic membrane. Sodium azide is a well-known inhibitor of SecA, but its

mechanism of action *in vivo* is poorly understood. To investigate this mechanism, we examined

the effect of azide on the growth of a library of > 1 million transposon insertion mutations. Our

results suggest that azide causes SecA to backtrack in its ATPase cycle, which disrupts binding

of SecA to the membrane and to its metal cofactor, which is iron. Our results provide insight into

the molecular mechanism by which SecA drives protein translocation and how this essential

biological process can be disrupted.

Introduction

In bacteria, protein substrates of the Sec machinery are transported across, or inserted into, the cytoplasmic membrane through a channel composed of integral membrane proteins SecY, -E and -G (Sec61α, -β and –γ in eukaryotes). In bacteria, translocation of most soluble periplasmic proteins and outer membrane proteins also requires the assistance of a motor ATPase, SecA, which drives translocation through repeated rounds of ATP binding and hydrolysis (1). In *Escherichia coli,* the catalytic core of SecA contains four domains: nucleotide binding domain-1 (NBD-1; amino acids 1-220 & 378-411), nucleotide binding domain-2 (NBD-2; 412-620), the polypeptide crosslinking domain (PPXD; 221-377) and the α-helical C-terminal domain (CTD; 378-830). SecA binds to substrate proteins in a groove formed between the PPXD and the two NBDs (2). ATP binding and hydrolysis cause conformational changes in the PPXD and CTD that drive protein translocation (3).

In addition to the catalytic core, most SecA proteins contain a relatively long C-terminal tail (CTT). In *E. coli*, the CTT contains a structurally flexible linker domain (FLD; amino acids 831-876 and a small metal-binding domain (MBD; 877-901) (4), which co-purifies with Zn^{2+} (5). The CTT is dispensable for translocation (6-8) although cells producing a truncated version of SecA that lacks the CTT display modest growth and translocation defects (6, 7). The only known function of the MBD (or the CTT generally) is in mediating the interaction of SecA with its binding partner SecB (9, 10), a molecular chaperone that is required for the translocation of a subset of Sec substrates (11). However, the MBD is present in many species that lack a SecB homologue (4, 12), suggesting that it may have an additional function.

Several auxiliary components are also required for efficient Sec-dependent translocation *in vivo*. For example, SecYEG forms a supercomplex with SecD, SecF, YajC and YidC (13, 14),

which has been proposed to assist in the insertion and assembly of integral membrane protein

complexes (15). Mutations disrupting components of this complex cause broad translocation

defects *in vivo* (16-18). In addition, a ribonucleoprotein complex known as the signal recognition

particle (SRP) is required for the efficient translocation of a subset of proteins, consisting mostly

of integral membrane proteins (19-21). Finally, many soluble periplasmic and outer membrane

proteins require the assistance of SecB (22).

Sodium azide inhibits the growth of bacteria by inhibiting SecA (23, 24). Mutations that

confer azide resistance map to the *secA* gene (23-25), and azide causes a nearly complete block

in SecA-mediated protein translocation within minutes of addition to growing cells (24). Azide

inhibits ATP turnover by the F-ATPase by inhibiting exchange of ADP for ATP, and it is

thought that azide inhibits SecA by a similar mechanism (26). In support of this notion, most of

the mutations that confer azide resistance result in amino acid substitutions in one of the two

nucleotide binding domains of SecA (23), and many cause an increased rate of nucleotide

exchange *in vitro* (27). However, the concentration of azide required to partially inhibit the rate

of ATP turnover by SecA *in vitro* (10-20 mM) (24, 28) is an order of magnitude greater than that

required completely block translocation *in vivo* (1-2 mM), suggesting that the mechanism by

which azide inhibits SecA is not fully understood.

In this study, we investigated the mechanism of action of azide *in vivo*. Previous genetics

studies have relied on either (i) selections for azide resistant mutants or (ii) the characterisation

of unrelated mutations in the *secA* gene on azide sensitivity (23, 24, 29, 30). These studies have

generally ignored weak effects and effects not attributable to mutations in the *secA* gene. In order

to identify these classes of mutation, we screened a very high-density library of *E. coli*

transposon insertion mutants (>1 million independent insertions) for mutants that became

enriched or depleted during exponential growth in the presence of azide (31). Consistent with the

idea that azide inhibits Sec-dependent protein translocation, insertions in genes encoding many

Sec components were depleted during growth in subinhibitory concentrations of azide. In

contrast, insertions truncating the CTT of SecA caused *E. coli* to grow faster than the parent

strain during exponential growth in the presence of azide. Further investigation indicated that: (i)

the CTT of SecA binds to iron; (ii) the CTT is required for high-affinity binding to phospholipids

in vivo; and (iii) azide disrupts binding of SecA to both iron and phospholipids at physiological

concentrations. Our results suggest that azide inhibits SecA by causing it to backtrack in the

ATPase cycle, which disrupts its interaction with the membrane.

Results

Identification of genes that affect the susceptibility of *E. coli* to sodium azide using

TraDIS. We used transposon-directed insertion-site sequencing (TraDIS) to identify mutations

that increased or decreased the susceptibility of *E. coli* BW25113 to azide (32). To this end, we

grew a library of >1 million independent transposon insertion mutants in the presence of

subinhibitory concentrations of azide. Mutations that increase susceptibility to azide became

depleted during growth in azide, while mutations that decrease susceptibility to azide became

enriched. We then determined the relative number of progeny produced by each insertion mutant

by sequencing the transposon junction using Illumina sequencing. Initial experiments indicated

that 0.5 mM sodium azide partially inhibited growth while concentrations of 0.25 mM or less

had a minimal effect on growth (**supplemental figure S1A**). Consistent with the known activity

of azide on protein translocation (24), growth of the TraDIS library in the presence of 0.25 mM

or 0.5 mM azide resulted in depletion of mutants containing insertions affecting the Sec

machinery (*e.g. yajC*, *secG*, *secF*, *secM*, *yidC*) and cell envelope stress responses (*e.g. cpxR*,

dsbA) from the library (**figure 1A; supplemental data S1**). However, cells containing insertions

in the *secA* and *secB* genes became enriched during growth in azide (**figure 1A**), suggesting that

these insertions decreased the susceptibility of *E. coli* to azide. All of the insertions in *secA*

resulted in truncation of the CTT of SecA, consistent with the essentiality of the catalytic core

for viability. The majority of the insertions enriched during growth in the presence of azide

resulted in truncation of SecA between amino acids 822 and 829 at the junction of the CTD and

the CTT (**figure 1B**), suggesting that the CTT contributed to azide susceptibility. All of the

transposons in *secB* were inserted in the same orientation and maintained expression of the

downstream *gpsA* gene (**supplemental figure S1B**) (31). Because increased production of GpsA

can suppress some Sec defects (33), we did not investigate the *secB* insertion mutants further. In

addition, insertions in genes required for the response to oxidative stress (*e.g. trxC*, *ahpC*, *gor*)

were depleted from the library during growth in the presence of azide (**figure 1A; supplemental**

data S1), suggesting that long-term exposure to azide causes oxidative stress.

Deletion mutations of genes identified in TraDIS have a similar effect on azide

sensitivity. To verify the results of our screen, we examined the growth rates of Δ*secG*, Δ*yajC*,

Δ*dsbA*, Δ*cpxR*, Δ*gor*, Δ*ahpC* and Δ*trxC* deletion mutants relative to the parent in the presence of

azide. With the exception of the Δ*gor* mutant, the presence of 0.25 or 0.5 mM sodium azide had

a greater effect on the growth rate of the mutants than it did on the parent strain (**table 1**).

Because mutations in *yajC* have not previously been shown to cause defects in protein

translocation, we wished to confirm that the increased azide sensitivity of the Δ*yajC* mutant was

not caused by polar effects on the expression of the downstream *secD* and *secF* genes (34).

IPTG-inducible production of YajC from a plasmid restored growth of the Δ*yajC* mutant to that

of the parent in the presence of 0.25 mM sodium azide (**figure 1C**). These results confirmed that

it was the lack of YajC in this mutant (and not SecD or SecF) that caused the increased

sensitivity of the Δ*yajC* mutant to azide.

We also investigated the effect of azide on the growth of an *E. coli* MC4100 mutant

containing an IS*1* insertion at codon ~827 in *secA* (*secA827*::IS*1*) (6). The *secA827*::IS*1* mutant

grew identically to the parent in the absence of azide, but growth of the mutant was significantly

less susceptible to the presence of 0.5 mM azide (**table 1**), confirming that truncation of the CTT

decreases susceptibility of *E. coli* to azide.

Azide disrupts binding of the MBD to iron *in vivo*. Structural studies have suggested that

the FLD inhibits the interaction of SecA with substrate proteins by binding in the substrate

binding groove and that interaction of the MBD with SecB overcomes this inhibition (35, 36).

Our results raised the possibility that azide disrupts the structure of the MBD, resulting in auto-

inhibition of SecA, which cannot be overcome by the interaction of SecA with SecB. To

investigate the effect of azide on the CTT, we purified a protein fusion between the CTT and the

small ubiquitin-like modifier (SUMO) *Saccharomyces cerevisiae* (SUMO-CTT) from cells

treated with 2 mM sodium azide for 10 minutes (*i.e.* conditions routinely used to inhibit SecA *in*

vivo). SUMO-CTT purified from azide-treated cells was more prone to aggregation compared to

protein purified from untreated cells (**figure 2A**), suggesting that azide disrupted the structure of

the CTT. Because the FLD is structurally flexible, we reasoned that the increased propensity of

the CTT to aggregate was likely due to the effect of azide on the structure of the MBD. Because

disrupting the structure of the MBD should result in release of the bound metal ion, we

determined the metal ion content of SUMO-CTT purified from azide-treated cells using mass

spectrometry (ICP-MS) (**figure 2B**). SUMO-CTT copurified with significant amounts of two

biologically relevant transition metals: iron and zinc. The samples also contained trace amounts

of copper, which was likely a contaminant since copper is not a physiological ligand for

cytoplasmic proteins (37). Azide treatment did not significantly affect the amount of zinc that

copurified with SUMO-CTT. However, it did cause a strong reduction in the amount of iron that

copurified with SUMO-CTT (**figure 2B**). These data suggested that the physiological ligand of

the MBD was iron.

SecA copurifies with iron. Because previous studies have suggested that the MBD binds

zinc (5, 38), we wished to confirm that full-length SecA binds to iron. Mis-metallation of iron-

binding proteins with zinc frequently occurs for several reasons (39, 40): first, the form of iron

typically bound by cytoplasmic proteins, Fe^{2+}, can be oxidised by dissolved oxygen in

purification buffers, while zinc is stable under aerobic conditions; in addition, many iron-

metalloproteins have a higher intrinsic affinity for zinc than for iron (37); finally, purification

buffers are often contaminated with low (but non-negligible) concentrations of Zn^{2+}. To

minimize the effect of these issues, we purified SecA as rapidly as possible using a protein that

was covalently modified at its C-terminus with biotin (SecA-biotin; (41)). Production of SecA-

biotin in strain DRH839 was controlled by an IPTG-inducible promoter, and SecA-biotin was the

only version of SecA produced in these cells. Incubation of cells lysates with streptavidin-

coupled beads resulted in the purification of only two proteins: SecA-biotin and streptavidin

(**figure 2C**). Analysis of the zinc and iron content of the eluted protein using optical emission

spectroscopy (ICP-OES) indicated that iron was the dominant metal species present at two

different induction levels (10 μM and 1 mM IPTG) (**figure 2D**). These data confirmed that the

physiological ligand of the MBD is iron.

Effect of azide on the oxidation state of the MBD. The MBD coordinates the bound iron

via three cysteines. Because our genetic analysis suggested that azide causes oxidative stress, we

reasoned that that azide might oxidise of the metal-coordinating cysteines, resulting in

misfolding of the MBD and release of the bound metal. To investigate this possibility, we

examined the effect of azide on binding of the MBD to metal *in vitro*. Because oxidation of iron

by oxygen could affect the affinity of SecA for the bound metal, we used full-length SecA that

had been charged with Zn^{2+} and measured the rate of release of Zn^{2+} using 4-(2-

pyridylazo)resorcinol (PAR). Binding of PAR to Zn^{2+} results in a strong increase in absorbance

at 492 nm. The presence of a strong oxidant, hydrogen peroxide, moderately increased the rate of

Zn^{2+} release (**figure 3**). However, the presence of 20 mM azide did not detectably affect the rate

of Zn^{2+} release by SecA, suggesting that azide does not directly oxidise the metal-coordinating

cysteines.

Azide disrupts binding of SecA to membrane phospholipids. Because the CTT has

previously been implicated in binding of SecA to anionic phospholipids (10), we wished to

investigate whether azide also disrupted binding of SecA to membrane phospholipids. To this

end, we used a hexahistidine-tagged protein fusion between SUMO and SecA (His-SUMO-

SecA). This fusion protein was functional *in vivo* since production of the protein from a plasmid

could complement the growth defect caused by depletion of SecA (**supplemental figure S2**).

His-SUMO-SecA copurified strongly with membrane phospholipids even after several high-salt

washes (**figure 4A**), and the copurifying lipids appeared to be enriched for phosphatidylglycerol

(PG), consistent with the affinity of SecA for acidic phospholipids (42, 43). In contrast, His-

SUMO-SecA containing alanine substitutions in two of the metal-coordinating cysteines (C885

and C887; His-SUMO-SecA$^{CC/AA}$) did not copurify with phospholipids (**supplemental figure**

S3), indicating that the MBD is required for high affinity binding to phospholipids. Because our

results indicated that azide disrupts the structure of the MBD, we investigated the effect of azide

on binding of His-SUMO-SecA to phospholipids. Treating cells producing His-SUMO-SecA

with azide prior to lysis strongly disrupted binding of SecA to phospholipids (**figure 4A**).

To investigate whether azide could disrupt binding of SecA to membranes at similar

concentrations *in vitro*, we examined the effect of azide on phospholipid-bound His-SUMO-

SecA, which was immobilised on a Ni-NTA column. Washing His-SUMO-SecA with buffer

containing 2 mM sodium azide resulted in a rapid loss of the bound phospholipid compared to

washing with buffer without azide (**figure 4B**). These results indicated that azide can disrupt

binding of SecA to the membrane at physiologically relevant concentrations. Furthermore,

because ATP was not added to the wash buffers, the defect in membrane binding was not caused

by the inhibition of nucleotide exchange.

<u>Discussion</u>

Our results indicate that azide does not disrupt SecA-mediated translocation by inhibiting the

rate of ATP turnover *in vivo*. Rather, they suggest that azide causes SecA to backtrack from the

ADP to the ATP-bound conformation, which disrupts the structure of the MBD and binding of

SecA to membrane phospholipids. Normally, SecA cycles through three nucleotide-dependent

conformations (**figure 5**): ATP-bound (**yellow**), ADP-bound (**red**) and nucleotide free (**green**).

Azide inhibits the F-ATPase by binding to the residues that coordinate the γ-phosphate in the

ADP-bound form of the protein, stabilising a conformation that resembles the ATP-bound form,

and it has been suggested that azide acts on other ATPases by a similar mechanism (26). The

ATP-bound form of SecA has a lower affinity for phospholipids than the ADP-bound form (43),

and backtracking to this conformation would result in release of SecA from the membrane.

Because binding of SecA to phospholipids is required for protein translocation (42-46),

disrupting the interaction of SecA with the membrane would also inhibit protein translocation. If

this model is correct, mutations that promote forward progress through the ATPase cycle would

confer resistance to azide, which could explain why (i) the vast majority of mutations that confer

azide resistance alter residues in NBD-1 and NBD-2 (23) and (ii) so many of these alterations

affect the affinity of SecA for nucleotide (27).

The decreased susceptibility of the *secA827* mutant to azide suggests that CTT auto-inhibits

the activity of SecA. This result is consistent with previous studies indicating that disruption of

the CTT inhibits both binding of SecA to membrane vesicles and translocation of proOmpA *in*

vitro (47). Structural studies have suggested that the CTT blocks the interaction of SecA with

substrate proteins *in vitro* and that binding of the MBD to SecB relieves this auto-inhibition (35).

Disrupting the structure of the MBD would simultaneously release SecA from the membrane and

inhibit its interaction with substrate protein. However, because the FLD is required for auto-

inhibition (35) but not for translocation (6-8), cells producing a SecA protein lacking the CTT

would be less susceptible to azide. It is unclear how azide disrupts the structure of the MBD.

However, it is unlikely that azide competes with SecA for binding to iron since *secA827*::IS*1*

mutants are only moderately less susceptible to azide than the parent. We speculate that because

coordination of the metal ligand by the MBD is strained (4), binding of the MBD to the

cytoplasmic membrane stabilises the structure of the MBD. As a result, disrupting the interaction

of SecA with the membrane destabilises the structure of the MBD, resulting in release of the

bound metal.

Our results strongly suggest that the physiological ligand of the MBD is iron. Previous

studies demonstrating that SecA binds to zinc have relied on extensive purification of the protein

under aerobic conditions (5, 38). Under these conditions, reaction of the bound iron with oxygen

likely destabilises its interaction with the MBD. In addition, because binding of SecA to the

membrane could stabilise binding of the MBD to iron, extensive purification of SecA away from

phospholipids could lead to loss of the native ligand. Purification buffers are typically

contaminated with non-negligible concentrations of zinc and other aerobically stable heavy

metals. Indeed, copurification of SUMO-CTT with copper confirms that this sort of exchange

takes place during purification. Although copper is equally as effective as zinc at promoting the

interaction of the MBD with SecB in *in vitro* binding assays (5), it is not a physiological ligand

for cytoplasmic proteins (37).

The increased sensitivity of *yajC* mutants to azide provides the first evidence linking YajC to

Sec-dependent protein translocation *in vivo*. Previous studies have attributed defects in

translocation caused by mutations in *yajC* to polar effects on expression of *secD* and *secF* (34),

which are transcribed in the same poly-cistronic message. However, the ability of a plasmid-born

copy of the *yajC* gene to complement a chromosomal deletion of *yajC* indicates that increased

azide sensitivity of the Δ*yajC* mutant is caused by a defect in YajC. This phenotype provides a

new tool for investigating the function of YajC *in vivo*. These results demonstrate the power of

TraDIS for identifying hidden or subtle phenotypes.

<u>Methods</u>

Chemicals and media. All chemicals were purchased from Fisher or Sigma-Aldrich unless

indicated. Cells were grown using LB (48). Where noted, sodium azide and

isopropylthiogalactoside (IPTG) were added at the indicated concentration. Where required,

ampicillin (200 μg/ml) or kanamycin (30 μg/ml) were added to the growth media.

Strains and plasmids. A description of all the strains and plasmids used in this study can be

found in **supplemental table S1**. Strains and plasmids were constructed using common genetic

methods (48, 49). Single gene deletion mutants from the Keio collection (50) were obtained from

the *E. coli* genetic stock centre (CGSC; Yale University, New Haven, Connecticut). Strains

DRH745 and DRH839 were constructed by integrating the IPTG-inducible copy of *secA* on

plasmid pDH692 or pDH771 onto the chromosome of MC4100 at the phage λ attachment site

using λInCh (51) and transducing the Δ*secA*::Kan mutation into this strain using P1. Strain

MG1115 was a kind gift from M. Grabowicz and T. Silhavy. Plasmid pDH543 was constructed

by PCR amplifying a fragment of the *secA* gene encoding the C-terminal 70 amino acids and

cloning into plasmid pCA597 cut with *BsaI* and *BamHI* (52). Plasmid pTrc-YajC was

constructed by cloning a PCR-amplified DNA fragment corresponding the *yajC* coding sequence

into strain pTrc99a (Promega).

TraDIS. TraDIS was carried out similar to Langridge*, et al.* (32). 50 ml of LB broth

containing 0, 0.25 or 0.5 mM NaN_3 were inoculated with 10 μl of a library of ~1 million *E. coli*

BW25113 mini-Tn*5* insertion mutants, and the cultures were grown to OD_{600} 1.0. Genomic DNA

was extracted using a Qiagen QIAamp DNA blood mini kit and then processed using a two-step

PCR method (53), which results in Illumina-compatible products. The PCR products were

purified using the Agencourt AMPure XP system by Beckman Coulter. The products were

sequenced using an Illumina MiSeq sequencer and the reads were mapped to the *E. coli*

reference genome NC_007779.1 (*E. coli* K-12 substr. W3110). The number of insertions in the

coding sequences (CDS) for each gene was then determined. To reduce the number of false

positives due to sequencing assignment errors, genes that exist in multiple copies on the

chromosome (e.g. insertion elements and rRNA operons) were eliminated from the analysis. In

addition, genes containing 15 or fewer total insertions across all three conditions were eliminated

in the data presented in **figure 1A**. BED files of the aligned sequences, the sequence file used for

alignments and the features file used for assessing the location of the insertions are available at

figshare.com (doi: 10.6084/m9.figshare.5280733).

Determination of metal content of SecA-biotin and Strep-SUMO-CTT. To determine the

effect of azide on binding of Strep-SUMO-CTT to iron, 100 ml cultures of BL21(DE3)

containing plasmid pDH543 was grown to OD_{600} 1.0 in LB, and production of Strep-SUMO-

CTE was induced using 1 mM IPTG. After 1 hour, cultures were split, and half the culture was

treated with 2 mM NaN_3 for 10 minutes. Cells were rapidly cooled, harvested by centrifugation

and lysed using B-PER cell lysis reagent from Pierce (Rockford, Illinois). Cell lysates were

incubated for 15 minutes with 50 μl of a 50% Streptactin-Sepharose slurry that had been pre-

equilibrated with wash buffer (10 mM HEPES (potassium salt) pH 7.5, 100 mM potassium

acetate) and washed extensively with wash buffer. The samples were washed a final time using

10 mM HEPES (potassium salt) pH 7.5 to remove excess salt, and the total protein was eluted

off of the column using using 10 mM HEPES (potassium salt) pH 7.5 buffer containing 7M

guanidinium hydrochloride. The propensity to aggregate was determined by diluting 50 μl of the

guanidinium-denatured protein into 950 μl of a 20 mM HEPES buffer and measuring light

scattering at 500 nm. The metal content of the samples was determined using inductively

coupled plasma-mass spectrometry (ICP-MS) (School of Geography, Earth and Environmental

Sciences, University of Birmingham, UK).

To determine the metal content of full-length SecA-biotin, a 100 ml culture of strain

DRH839 was grown in the indicated concentration of IPTG to OD_{600} ~ 1 and lysed using cell

disruption. Cell lysates were incubated with 100 μl Streptactin-sepharose for 15 minutes on ice,

and the sepharose beads were washed four times with 30 ml binding buffer (50 mM HEPES, pH

7.5, 100 mM K·acetate, 10 mM Mg·acetate, 0.1% nonidet P40). Metal was eluted from the beads

by incubating with metal elution buffer (10 mM HEPES, pH 7.5, 50 mM EDTA) at 55°C for 30

minutes, and the zinc and iron content were determined using inductively coupled plasmid-

optical emission spectroscopy (ICP-OES). Protein was eluted from the beads by boiling in SDS

sample buffer, and the protein content was determined using a Bradford assay. The eluted protein

was resolved on a BioRad 15% TGX gel, and the protein bands were visualised using the

BioRad Strain-Free stain.

In vitro zinc-release assay. Full-length SecA was purified similar to Huber, *et al.* (54) except

that the protein was not subjected to an on-column denaturation step. SecA was charged with

Zn^{2+} by incubating 40 μM of the purified protein with an equimolar concentration of $ZnCl_2$.

Excess metal was removed from the protein using a 5 ml HiTrap Desalting Column (GE

Healthcare). To measure the rate of Zn^{2+} release, 4 μM SecA was incubated in buffer containing

20 mM HEPES (potassium salt), pH 7.5, 100 mM potassium acetate, 20 mM magnesium acetate

and 100 μM PAR in the presence of the indicated concentration of H_2O_2 or NaN_3. The increase

in PAR absorbance was measured every 60 s for 90 minutes at 492 nm using a BMG Labtech

FLUOStar plate reader.

Phospholipid binding assay. Strain DRH625 or MJ118 was grown to late log phase in LB,

and production of His-SUMO-SecA was induced using 1 mM IPTG. Where indicated, the

culture was divided after the addition of IPTG, and half of the culture was treated with 2 mM

sodium azide for 10 minutes. Cells were harvested by centrifugation, resuspended in lysis buffer

(20 mM HEPES [potassium salt] pH7.5, 500 mM NaCl, 1 mM tris(2-carboxyethyl)phosphine

[TCEP]) and lysed by cell disruption. Clarified cell lysates were passed over a 1 ml Ni-HiTrap

column (GE Biosciences), and the bound protein was washed with 50 volumes of lysis buffer

containing 20 mM Imidazole. His-SUMO-SecA was eluted from the column using lysis buffer

containing 500 mM imidazole and dialysed against wash buffer lacking imidazole. The

concentration of the protein was adjusted to 1 mg/ml to a final volume of 2 ml, and the

phospholipids were extracted using 2 ml methanol and 1 ml chloroform. The chloroform phase

was removed, and phospholipids were concentrated by air drying and resuspending in

chloroform to a final volume of 100 μl. The indicated volume of sample was spotted onto TLC

silica gel 60 T$_{254}$ plates (Merck Millipore) and resolved using a mixture of 60 chloroform: 25

methanol: 4 water. For the *in vitro* phospholipid release assay, 0.2 mg of His-SUMO-SecA

purified from untreated cells was bound to 100 μl Ni-NTA agarose beads (Life Technologies).

The beads were incubated with 500 μl of lysis buffer or buffer containing 2 mM sodium azide for

10 minutes and washed three times with 500 μl lysis buffer. The protein was eluted from the

beads with lysis buffer containing 500 mM imidazole, and the phosopholipid content of the

supernatant was analysed using the method described above.

<u>Acknowledgements</u>

We thank Dr. S. Baker and Dr. M. Thompson for technical assistance. We also thank Dr. J.

Bryant, Dr. T. Knowles, E. Goodall, Prof. J. Cole and members of the Huber, Henderson, Lund

and Grainger labs for helpful advice and discussions. This work was funded by the

Biotechnology and Biological Sciences Research Council grant BB/L019434/1 to D. H., M. J.

and R. C..

374 **<u>References</u>**

1. Cranford Smith T & Huber D (2018) The way is the goal: how SecA transports proteins

across the cytoplasmic membrane in bacteria. *FEMS Microbiol Lett*.

2. Zimmer J & Rapoport TA (2009) Conformational flexibility and peptide interaction of

the translocation ATPase SecA. *J Mol Biol* 394(4):606-612.

3. Collinson I, Corey RA, & Allen WJ (2015) Channel crossing: how are proteins shipped

across the bacterial plasma membrane? *Philos Trans R Soc Lond B Biol Sci* 370(1679).

4. Dempsey BR*, et al.* (2004) Solution NMR structure and X-ray absorption analysis of the

C-terminal zinc-binding domain of the SecA ATPase. *Biochemistry* 43(29):9361-9371.

5. Fekkes P, de Wit JG, Boorsma A, Friesen RH, & Driessen AJ (1999) Zinc stabilizes the

SecB binding site of SecA. *Biochemistry* 38(16):5111-5116.

6. Grabowicz M, Yeh J, & Silhavy TJ (2013) Dominant negative lptE mutation that

supports a role for LptE as a plug in the LptD barrel. *J Bacteriol* 195(6):1327-1334.

7. Or E, Boyd D, Gon S, Beckwith J, & Rapoport T (2005) The bacterial ATPase SecA

functions as a monomer in protein translocation. *J Biol Chem* 280(10):9097-9105.

8. Or E, Navon A, & Rapoport T (2002) Dissociation of the dimeric SecA ATPase during

protein translocation across the bacterial membrane. *Embo J* 21(17):4470-4479.

9. Fekkes P, van der Does C, & Driessen AJ (1997) The molecular chaperone SecB is

released from the carboxy-terminus of SecA during initiation of precursor protein

translocation. *Embo J* 16(20):6105-6113.

10.   Breukink E*, et al.* (1995) The C terminus of SecA is involved in both lipid binding and

SecB binding. *J Biol Chem* 270(14):7902-7907.

11.   Randall LL & Hardy SJ (2002) SecB, one small chaperone in the complex milieu of the

cell. *Cell Mol Life Sci* 59(10):1617-1623.

12.   van der Sluis EO & Driessen AJ (2006) Stepwise evolution of the Sec machinery in

Proteobacteria. *Trends Microbiol* 14(3):105-108.

13.   Schulze RJ*, et al.* (2014) Membrane protein insertion and proton-motive-force-dependent

secretion through the bacterial holo-translocon SecYEG-SecDF-YajC-YidC. *Proc Natl*

Acad Sci U S A 111(13):4844-4849.

14.   Duong F & Wickner W (1997) Distinct catalytic roles of the SecYE, SecG and

SecDFyajC subunits of preprotein translocase holoenzyme. *Embo J* 16(10):2756-2768.

15.   Komar J*, et al.* (2016) Membrane protein insertion and assembly by the bacterial holo-

translocon SecYEG-SecDF-YajC-YidC. *Biochem J* 473(19):3341-3354.

16.   Gardel C, Johnson K, Jacq A, & Beckwith J (1990) The secD locus of E.coli codes for

two membrane proteins required for protein export. *Embo J* 9(10):3209-3216.

17.   Pogliano JA & Beckwith J (1994) SecD and SecF facilitate protein export in Escherichia

coli. *Embo J* 13(3):554-561.

18.   Samuelson JC*, et al.* (2000) YidC mediates membrane protein insertion in bacteria.

Nature 406(6796):637-641.

19.   Huber D*, et al.* (2005) Use of thioredoxin as a reporter to identify a subset of Escherichia

coli signal sequences that promote signal recognition particle-dependent translocation. *J*

Bacteriol 187(9):2983-2991.

20.   Ulbrandt ND, Newitt JA, & Bernstein HD (1997) The E. coli signal recognition particle

417   is required for the insertion of a subset of inner membrane proteins. *Cell* 88(2):187-196.

418   21.   Saraogi I & Shan SO (2014) Co-translational protein targeting to the bacterial membrane.

419   *Biochim Biophys Acta* 1843(8):1433-1441.

420   22.   Kumamoto CA & Beckwith J (1985) Evidence for specificity at an early step in protein

421   export in Escherichia coli. *J Bacteriol* 163(1):267-274.

422   23.   Huie JL & Silhavy TJ (1995) Suppression of signal sequence defects and azide resistance

423   in Escherichia coli commonly result from the same mutations in secA. *J Bacteriol*

424   177(12):3518-3526.

425   24.   Oliver DB, Cabelli RJ, Dolan KM, & Jarosik GP (1990) Azide-resistant mutants of

426   Escherichia coli alter the SecA protein, an azide-sensitive component of the protein

427   export machinery. *Proc Natl Acad Sci U S A* 87(21):8227-8231.

428   25.   Fortin Y, Phoenix P, & Drapeau GR (1990) Mutations conferring resistance to azide in

429   Escherichia coli occur primarily in the secA gene. *J Bacteriol* 172(11):6607-6610.

430   26.   Bowler MW, Montgomery MG, Leslie AG, & Walker JE (2006) How azide inhibits ATP

431   hydrolysis by the F-ATPases. *Proc Natl Acad Sci U S A* 103(23):8646-8649.

432   27.   Schmidt M, Ding H, Ramamurthy V, Mukerji I, & Oliver D (2000) Nucleotide binding

433   activity of SecA homodimer is conformationally regulated by temperature and altered by

434   prlD and azi mutations. *J Biol Chem* 275(20):15440-15448.

435   28.   van der Wolk JP, de Wit JG, & Driessen AJ (1997) The catalytic cycle of the escherichia

436   coli SecA ATPase comprises two distinct preprotein translocation events. *EMBO J*

437   16(24):7297-7304.

29.   Das S, *et al.* (2012) The variable subdomain of Escherichia coli SecA functions to

regulate SecA ATPase activity and ADP release. *J Bacteriol* 194(9):2205-2213.

30.   Matsumoto G, Nakatogawa H, Mori H, & Ito K (2000) Genetic dissection of SecA:

suppressor mutations against the secY205 translocase defect. *Genes Cells* 5(12):991-999.

31.   Goodall ECA, *et al.* (2018) The Essential Genome of Escherichia coli K-12. *MBio* 9(1).

32.   Langridge GC, *et al.* (2009) Simultaneous assay of every Salmonella Typhi gene using

one million transposon mutants. *Genome research* 19(12):2308-2316.

33.   Shimizu H, Nishiyama K, & Tokuda H (1997) Expression of gpsA encoding biosynthetic

sn-glycerol 3-phosphate dehydrogenase suppresses both the LB- phenotype of a secB null

mutant and the cold-sensitive phenotype of a secG null mutant. *Mol Microbiol*

26(5):1013-1021.

34.   Pogliano KJ & Beckwith J (1994) Genetic and molecular characterization of the

Escherichia coli secD operon and its products. *J Bacteriol* 176(3):804-814.

35.   Gelis I, *et al.* (2007) Structural basis for signal-sequence recognition by the translocase

motor SecA as determined by NMR. *Cell* 131(4):756-769.

36.   Hunt JF, *et al.* (2002) Nucleotide control of interdomain interactions in the

conformational reaction cycle of SecA. *Science* 297(5589):2018-2026.

37.   Waldron KJ & Robinson NJ (2009) How do bacterial cells ensure that metalloproteins

get the correct metal? *Nat Rev Microbiol* 7(1):25-35.

38.   Woodbury RL, Hardy SJ, & Randall LL (2002) Complex behavior in solution of

homodimeric SecA. *Protein Sci* 11(4):875-882.

39.   Gu M & Imlay JA (2013) Superoxide poisons mononuclear iron enzymes by causing

mismetallation. *Mol Microbiol* 89(1):123-134.

40. Imlay JA (2014) The mismetallation of enzymes during oxidative stress. *J Biol Chem*

289(41):28121-28128.

41. Huber D*, et al.* (2017) SecA Cotranslationally Interacts with Nascent Substrate Proteins

In Vivo. *J Bacteriol* 199(2).

42. Lill R, Dowhan W, & Wickner W (1990) The ATPase activity of SecA is regulated by

acidic phospholipids, SecY, and the leader and mature domains of precursor proteins.

Cell 60(2):271-280.

43. Breukink E, Demel RA, de Korte-Kool G, & de Kruijff B (1992) SecA insertion into

phospholipids is stimulated by negatively charged lipids and inhibited by ATP: a

monolayer study. *Biochemistry* 31(4):1119-1124.

44. Bauer BW, Shemesh T, Chen Y, & Rapoport TA (2014) A "push and slide" mechanism

allows sequence-insensitive translocation of secretory proteins by the SecA ATPase. *Cell*

157(6):1416-1429.

45. Ulbrandt ND, London E, & Oliver DB (1992) Deep penetration of a portion of

Escherichia coli SecA protein into model membranes is promoted by anionic

phospholipids and by partial unfolding. *J Biol Chem* 267(21):15184-15192.

46. Kusters R, Dowhan W, & de Kruijff B (1991) Negatively charged phospholipids restore

prePhoE translocation across phosphatidylglycerol-depleted Escherichia coli inner

membranes. *J Biol Chem* 266(14):8659-8662.

47. den Blaauwen T*, et al.* (1997) Inhibition of preprotein translocation and reversion of the

membrane inserted state of SecA by a carboxyl terminus binding mAb. *Biochemistry*

36(30):9159-9168.

48.   Miller JH (1992) *A Short Course in Bacterial Genetics* (Cold Spring Harbor Press, Cold

484   Spring Harbor, NY).

485   49.   Sambrook J & Russell DW (2001) *Molecular Cloning: A Laboratory Manual, 3rd ed.*

486   (Cold Spring Harbor Press, Cold Spring Harbor, NY).

487   50.   Baba T*, et al.* (2006) Construction of Escherichia coli K-12 in-frame, single-gene

488   knockout mutants: the Keio collection. *Mol Syst Biol* 2:2006 0008.

489   51.   Boyd D, Weiss DS, Chen JC, & Beckwith J (2000) Towards single-copy gene expression

490   systems making gene cloning physiologically relevant: lambda InCh, a simple

491   Escherichia coli plasmid-chromosome shuttle system. *J Bacteriol* 182(3):842-847.

492   52.   Andreasson C, Fiaux J, Rampelt H, Mayer MP, & Bukau B (2008) Hsp110 is a

493   nucleotide-activated exchange factor for Hsp70. *J Biol Chem* 283(14):8877-8884.

494   53.   Christen B*, et al.* (2011) The essential genome of a bacterium. *Mol Syst Biol* 7:528.

495   54.   Huber D*, et al.* (2011) SecA interacts with ribosomes in order to facilitate

496   posttranslational translocation in bacteria. *Mol Cell* 41(3):343-353.

499 **Table 1. Relative growth rates of single gene deletion mutants from the Keio collection.**

Strain	Growth rates*		
	LB†	0.25 mM NaN$_3$†	0.5 mM NaN$_3$†
BW25113	1.00 ± 0.00	1.00 ± 0.09	1.00 ± 0.09
$\Delta yajC$	0.88 ± 0.05	0.59 ± 0.08	0.54 ± 0.11
$\Delta secG$	1.02 ± 0.05	0.53 ± 0.11	0.75 ± 0.1
$\Delta dsbA$	1.01 ± 0.08	0.95 ± 0.09	0.63 ± 0.15
$\Delta cpxR$	1.13 ± 0.13	0.93 ± 0.02	0.65 ± 0.15
Δgor	0.96 ± 0.01	0.97 ± 0.08	1.02 ± 0.04
$\Delta ahpC$	0.95 ± 0.00	0.80 ± 0.06	0.64 ± 0.06
$\Delta trxC$	0.92 ± 0.00	0.89 ± 0.02	0.78 ± 0.04
MC4100	1.00 ± 0.02	ND‡	1.00 ± 0.03
$secA827$::IS1	1.03 ± 0.01	ND‡	1.38 ± 0.18

500 *Growth rates relative to parent strain grown under the same conditions.

501 †Strains were grown in LB alone or in LB containing the indicated concentration of NaN$_3$.

502 ‡not determined

Figure 1. Effect of azide on growth of transposon insertions mutants in TraDIS. (A) Plot of the degree of depletion or enrichment in transposon insertions in all of the non-essential genes in *E. coli* after growth of the TraDIS library to $OD_{600} = 0.9$ in the presence of 0.5 mM sodium azide (see **supplemental data S1B**). The $\log_2$ of the fraction of the number insertions in a gene after growth in the presence of 0.5 mM azide over the number of insertions after growth in LB was plotted according to the degree of enrichment. Data points representing insertions in the *secA*, *secB*, *secF*, *secG*, *secM*, *yidC*, *yajC*, *cpxR*, *dsbA*, *ahpC*, *gor* and *trxC* genes are indicated. The growth of selected single gene deletion mutants in the presence of azide (bolded) was compared with that of the parent (see **Table 1**). (B) Number of mutants containing insertions at the indicated location in the *secA* gene after growth of the TraDIS library in the absence (black) or presence of 0.25 (blue) or 0.5 mM (red) NaN_3. Most of these insertions truncate the *secA* gene between codons 822 and 829 at the junction between the HSD and the CTT. (C) Strains

517 BW25113 (WT) or JW0397 (Δ*yecA*) containing plasmid pTrc99a or pTrc-YajC were grown in

518 the presence of 0.5 mM azide and the indicated concentration of sodium azide in LB at 37°C.

519 Growth of the strains was monitored by measuring the increase in optical density at 600 nm.

Figure 2. Interaction of SecA with iron *in vivo*. (A & B) Cells producing SUMO-CTT were

incubated in the absence (untreated) or presence (+N₃) of 2 mM NaN₃ for 10 minutes. SUMO-

CTT was purified from the cell lysates using streptactin beads and washed extensively with

buffer. (A) The bound protein was eluted from the streptactin beads using 7M guanidinium and

was then diluted into buffer lacking guanidinium. Aggregation of the protein was measured using

light scattering at 500 nm. Confidence intervals are the s.e.m. (B) The iron content of SUMO-

CTT was determined using mass spectrometry (ICP-MS) and normalised to the protein

concentration in the eluted protein. Confidence intervals are one s.d. (C & D) Strain DRH839

(p_{trc}-*secA-biotin*) was grown to $OD_{600} = 1$ in the presence of either 10 μM or 1 mM IPTG, as

indicated. SecA-biotin was purified from the cell lysates using Streptavidin-coupled beads. (C)

The eluted protein was resolved by SDS-PAGE on a 15% gel and visualised using BioRad Stain-

Free dye. (D) The iron and zinc content of the samples was determined using ICP-OES and

normalised to the concentration of protein in the eluted sample. Black, Zn content; Grey, Fe

content. Confidence intervals are one s.d.

Figure 3. Effect of hydrogen peroxide and sodium azide on metal binding by SecA. 4 μM

zinc-bound SecA and 100 μM PAR were incubated in the presence of 2 mM (filled triangles), 10

mM (open squares) or 50 mM (filled squares) hydrogen peroxide, 20 mM sodium azide (open

circles) or in the absence of hydrogen peroxide and sodium azide (closed circles). Binding of

PAR to free Zn^{2+} was measured by the increase in absorbance at 492 nm over a period of 60

minutes. Confidence intervals are one s.d.

Figure 4. Effect of azide on binding of SecA to phospholipids *in vitro* and *in vivo*. (A) A

culture of *E. coli* producing His-SUMO-SecA was grown to late exponential phase. Half of the

culture was treated with 2 mM sodium azide for 10 minutes prior to lysis (lanes 3 & 4), and the

other half was left untreated (lanes 1 & 2). His-SUMO-SecA was purified from the cell lysates

using Ni-affinity purification. Phospholipids from 2 mg of the purified protein were extracted

into 100 μl chloroform, and 5μl of the extracted phospholipids (lanes 1 & 3) and the wash buffer

(lanes 2 & 4) were resolved using thin-layer chromatography (TLC). The positions of

phosphatidylglycerol (PG) and phosphatidylethanolamine (PE) are indicated. (B) His-SUMO-

SecA was purified from untreated cells as in (A) and bound to a Ni-NTA beads. The beads were

then washed with buffer containing (lanes 3 & 4) or lacking (lanes 1 & 2) 2 mM sodium azide.

The lipids from 0.2 mg of protein were extracted from the eluted protein into 100 μl chloroform,

and 20 μl of the extracted lipid (lanes 1 & 3) and the wash buffer (lanes 2 & 4) were resolved

using TLC. The positions of PG, PE and a non-specific band from the wash buffer (*) are

indicated.

Figure 5. Diagrammatic representation of the mechanism of inhibition of SecA by azide.

SecA cycles through three nucleotide bound states: an ATP-bound state (yellow), an ADP-bound

state (red) and a nucleotide-free state (green). Binding of SecA to ATP results in translocation of

the substrate polypeptide through SecYEG (i) followed by hydrolysis ATP to ADP. Exchange of

ADP for ATP requires release of ADP (iii) and is the rate-limiting step the ATPase cycle of

SecA. Our results suggest that binding of azide to the ADP-bound form of SecA causes SecA to

backtrack to the ATP-bound conformation (yellow), which disrupts binding of SecA to the

membrane and to iron (Fe; iv).

1   **Table S1. Strains and plasmids used in this study.**

Strain	Description	Reference
BW25113	*Δ(araD-araB)567 ΔlacZ4787*(::rrnB-3) *rph-1 Δ(rhaD-rhaB)568 hsdR514*	(50)
MC4100	*araD139 Δ(argF-lac)U169 rpsL150 relA1 deoC1 rbsR fthD5301 fruA25*	Lab stock
BL21(DE3)	*E. coli B F- ompT gal dcm hsdSB(rB- mB-) λDE3*	Lab stock
JW0073	BW25113 Δ*leuA*::Kan^R	(50)
JW0397	BW25113 Δ*yajC*::Kan^R	(50)
JW0598	BW25113 Δ*ahpC*::Kan^R	(50)
JW2566	BW25113 Δ*trxC*::Kan^R	(50)
JW3142	BW25113 Δ*secG*::Kan^R	(50)
JW3467	BW25113 Δ*gor*::Kan^R	(50)
JW3832	BW25113 Δ*dsbA*::Kan^R	(50)
JW3883	BW25113 Δ*cpxR*::Kan^R	(50)
MG1115	MC4100 Ara$^+$ *secA827*::IS*1..leuA*::Tn*10*	(50)
DRH543	BL21(DE3) + pDH543	This study
DRH625	BL21(DE3) + pDH625	(54)
DRH745	MC4100 Δ*secA*::Kan^R Δ(λ*att-lom*)::Amp^R p_{lacUV5}-*secA*	This study
DRH839	MC4100 Δ*secA*::Kan^R Δ(λ*att-lom*)::Amp^R p_{lacUV5}-*secA-biotin*	(41)

DRH1075	MC4100 *secA827*::IS*1..leuA*::Tn*10*	This study
MJ118	BL21(DE3) + pMJ118	This study
Plasmid	**Description**	**Source**
pTrc99a	Multi-copy plasmid with ampicillin resistance marker containing IPTG-inducible p_{trc} promoter	Promega
pTrc-YajC	pTrc99a + *yajC*	This study
pCA528	pET24a plasmid producing His$_6$-tagged *S. cerevisiae* SUMO from a T7-inducible promoter	(52)
pDH543	Multi-copy plasmid with kanamycin resistance marker expressing a fusion protein between Strep-SUMO and the C-terminal 70 amino acids of SecA (SUMO-CTT) under control of a T7-inducible promoter	This study
pDH625	pCA528 containing *secA*	(54)
pMJ118	pCA528 containing *secA*$^{CC/AA}$	This study

Figure S1. TraDIS analysis of the effect of azide on the growth of *E .coli* BW25113. (A)

Strain BW25113 was grown in LB in the presence of the indicated concentration of sodium

azide. Growth of the culture was monitored by optical density at 600 nm. (B) Map of unique

transposon insertions in the *yibN-grxC-secB-gpsA* operon. Depicted are the location of all

transposons in the TraDIS transposon library (black), transposons with the internally encoded

promoter transcribing rightwards (red) and transposons with the internally encoded promoter

transcribing leftward (blue). The *gpsA* gene contains no insertions suggesting it is essential under

these growth conditions. In addition, all of the insertions in the *secB* gene preserve expression of

the downstream *gpsA* gene.

DRH745
($\Delta secA\ att_\lambda$::p$_{lacUV5}$-secA)

pDH625	pCA528	pDH625	pCA528
(His-SUMO-SecA)	(His-SUMO)	(His-SUMO-SecA)	(His-SUMO)

+IPTG   -IPTG

Figure S2. Production of His-SUMO-SecA can complement the growth defect caused by depletion of SecA. Strain DRH745 (p$_{lacUV5}$-*secA*) containing either pDH625 or pCA528 was restreaked onto LB plates containing 1 mM IPTG or lacking IPTG and incubated overnight at 37°C.

lane: 1   2   3   4   5   6

Figure S4. Effect of alanine substitution that disrupt the structure of the MBD on

phospholipid binding. Cultures of *E. coli* producing His-SUMO-SecA (lanes 1-4) or His-

SUMO-SecA$^{CC/AA}$ (lanes 5& 6) were grown to late exponential phase. In the case of His-SUMO-

SecA, half of the culture was treated with 2 mM sodium azide for 10 minutes prior to lysis (lanes

3 & 4), and the other half was left untreated (lanes 1 & 2). His-SUMO-SecA and SUMO-

SecA$^{CC/AA}$ were purified from the cell lysates using Ni-affinity purification. Phospholipids from

2 mg of the purified protein were extracted into 100 μl chloroform, and 5μl of the extracted

phospholipids (lanes 1, 3 and 6) and the wash buffer (lanes 2, 4 and 5) were resolved using thin-

layer chromatography (TLC). The positions of phosphatidylglycerol (PG),

phosphatidylethanolamine (PE) and cardiolipin (CL) are indicated.

Supplemental Data S1. Analysis of transposon insertion mutants in TraDIS library after

outgrowth in LB or LB containing NaN₃. (A) Number of insertions in each gene in the *E. coli*

genome after growth of the TraDIS library in LB, LB containing 0.25 mM NaN_3 or LB

containing 0.5 mM NaN_3. Genomic DNA from one experiment was prepared and sequenced

twice to increase the number of sequencing reads and to examine the reproducibility of the

results. For each run the number of insertions in each gene was normalised to the total number of

sequencing reads. (B) The degree of enrichment or depletion of insertion mutations in the library

after growth in the presence of azide was calculated by determining the log (base 2) of the ratio

of the number of insertions in a gene after outgrowth in the azide-treated conditions to the

number of insertions after outgrowth in LB. The BED files used to derive the data presented in

parts A and B is available at figshare.com (doi: 10.6084/m9.figshare.5280733).

43
44