Population genomics of *Wolbachia* and mtDNA in *Drosophila simulans* from California

Sarah Signor*

*Department of Molecular and Computational Biology, University of Southern California, Los Angeles,

California USA

Key words: *Drosophila simulans*, population genetics, *Wolbachia*

Communicating Author:          Sarah Signor

11                                   Section of Molecular and Computational Biology

12                                   University of Southern California

13                                   Los Angeles, California USA

14                                   ssignor@usc.edu

15                                   Phone   213-740-3065

ABSTRACT: *Wolbachia pipientis* is an intracellular endosymbiont infecting many arthropods and filarial

nematodes. Little is known about the short-term evolution of *Wolbachia* or its interaction with its host.

*Wolbachia* is maternally inherited, resulting in co-inheritance of mitochondrial organelles such as

mtDNA.  Here I explore the short-term evolution of *Wolbachia*, and the relationship between *Wolbachia*

and mtDNA, using a large inbred panel of *Drosophila simulans* infected with the *Wolbachia* strain *w*Ri. I

find reduced diversity relative to expectation in both *Wolbachia* and mtDNA, but only mtDNA shows

evidence of a recent selective sweep or population bottleneck. I estimate *Wolbachia* and mtDNA titre in

each genotype, and I find considerable variation in both phenotypes, despite low genetic diversity in

*Wolbachia* and mtDNA. A phylogeny of *Wolbachia* and of mtDNA show that both trees are largely

unresolved, suggesting a recent origin of the infection derived from a single origin. Using *Wolbachia* and

mtDNA titre as a phenotype, I perform an association analysis with the nuclear genome and find several

regions implicated in the phenotype, including one which contains four CAAX-box protein processing

genes. CAAX-box protein processing can be an important part of host-pathogen interactions in other

systems, suggesting interesting directions for future research.

INTRODUCTION:

Heritable symbiotic associations such as that between *Drosophila* and *Wolbachia pipientis* have

widespread impact on host ecology and evolution. These types of heritable endosymbiotic relationships

are recognized as key drivers of evolution, but the intraspecific variation that effects their short-term

evolution is not well explored. *Wolbachia* are $\alpha$-proteobacterial endosymbionts found in up to 40% of all

arthropod species[1-3]. *Wolbachia* are maternally transmitted and spread through manipulating the

reproductive strategies of their host, using mechanisms such as feminization, male-killing, or cytoplasmic

incompatibility. The most common of these is cytoplasmic incompatibility, where mating between males

and females of the same species results in embryonic mortality if they have different *Wolbachia* infection

status[4-8]. *Wolbachia* may also confer certain protections upon their host, such as increased resistance to

certain viruses, or increased survival when exposed to certain environmental stressors[9-15]. *Wolbachia* is

one of the most abundant obligate intracellular parasites, given that 85% of animal species are insects.

This has profound meaning for evolutionary processes such as sexual selection and speciation[16,17].

*Wolbachia* strain *w*Ri is known to have spread recently in the sister species to the model organism

*Drosophila melanogaster*, *D. simulans*[4,5,8,18]. It was at ~95-100% frequency in Southern California

populations at the time its original sampling in the 1980's[19,20]. It likely invaded California less than 25

years before it was first detected in 1984[21]. It is now thought to have been horizontally transmitted to *D.*

*simulans* from *D. ananassae*, though the same strain is also found in *D. suzukkii*[22]. The maternal

transmission of *Wolbachia* means that as the microorganism spreads all maternally inherited organelles

spread along with it. Most notably mtDNA will be forced through a bottleneck, lowering the diversity of

mtDNA in infected populations[18,23,24]. This will cause mtDNA and *Wolbachia* to be more closely

associated than nuclear genes, and this coupling has been demonstrated previously in *D. simulans*[18,25,26]. In

fact, *D. simulans* is known to have three major mitochondrial haplotypes (*si*I, *si*II, and *si*III) and two

subtypes (*si*IIA and *si*IIB) that harbor very little variation and that appear to be nonrandomly associated

with *Wolbachia* strains[27-29]. These mitochondrial haplotypes are largely allopatric, except for the presence

of both *si*II and *si*III in Madagascar and La Reunion[30].

In *D. melanogaster* variation in *Wolbachia* has been well investigated using genomic data, though

this has not been done in the genomic era in *D. simulans*[31]. They found long lived associations between

mitochondrial and *Wolbachia* haplotypes and strong geographic structuring among cytotypes[31-33]. This

study also observed that *Wolbachia* titre varied among fly populations as the result of intraspecific

nuclear genetic variation[31]. However, the assumption that it was due to intraspecific nuclear background

was based on the presence of a constant environment and no polymorphisms were identified that could be

affecting this phenotype. Very little is known about how *Wolbachia* interact with their hosts, though

recent work has uncovered evidence that deubiquitylating enzymes produced by *Wolbachia* and secreted

into the host cytoplasm mediate cytoplasmic incompatibility[34]. *Wolbachia* DNA is also frequently

inserted into the host genome, though this has not occurred with *w*Ri in *D. simulans*[21]. Genes involved in

the formation of germline stem cells such as *benign gonial cell neoplasm* and *bag-of-marbles* are

considered candidates for interacting with *Wolbachia*, and have been found to have unusual population

genetic patterns in *D. melanogaster* [35,36]. *bag-of-marbles* has been suggested to interact with *Wolbachia*

due to fertility rescue in hypomorphs, but the interaction of this gene with *Wolbachia* in natural

populations is not clear[22,35-37]. Notably, *Wolbachia* localizes in tissues differently depending upon the

strain and species so the interactions between the host and *Wolbachia* are likely to also be different[38,39].

*Wolbachia* infections must be maintained in host populations through transovarial transmission,

wherein *Wolbachia* is present in the germline at sufficient copy number to ensure transmission but not to

cause host pathology[40]. *Wolbachia* titre has been shown to have important phenotypic effects on the

host[11,41-49]. However, control of *Wolbachia* replication is not well understood, nor is the dependence of

this control on host background versus bacterial genotype[11,50-52]. Differences in *Wolbachia* titre when it is

transinfected between species suggests a role of host background in controlling copy number, population

genomics in *D. melanogaster* suggest an effect of host background, and there does seem to be host-

specific patterns of tissue colonization[53-55]. However, multiple *Wolbachia* genotypes can also behave

differently in the same genetic background suggesting contributions from the bacterial genome[51,56]. It is

also possible to select for greater *Wolbachia* densities, though the heritability of this is unclear[57,58].

Here I investigate the dynamics of *Wolbachia* and mtDNA in a large panel of *D. simulans* from a

single Californian population.  I determine infection status of *Wolbachia* in the panel of *D. simulans*

genotypes. I look for signatures of selection in both genomes using summary statistics Tajima's $D$ and $\pi$

and find that while *Wolbachia* patterns of variation are not unusual given its demographic history the

reduction in mtDNA diversity is suggestive of a recent bottleneck due either to selection or changes in

population size.  I also measure linkage disequilibrium between mtDNA and *Wolbachia* as a proxy for co-

inheritance. Using whole genome sequences, I investigate the phylogeny of both *Wolbachia* and mtDNA

and find that in this population they are essentially unresolved. I investigate variation in the copy number

of both *Wolbachia* and mtDNA in this population using relative estimates derived from illumina

sequencing coverage compared to nuclear coverage. I find considerable copy number variation in this

population, and an association analysis using this as a phenotype implicates several genomic regions

potentially involved in mediating this phenotype. This includes a region containing multiple genes

involved CAAX-box protein prenylation, a process that is important for mediating the relationship

between host and pathogen in other systems[59-61].

<u>METHODS</u>

***Drosophila* strains**

Strains are as described in[62]. Briefly, the 167 *D. simulans* lines were collected in the Zuma organic orchard in

Zuma beach, California in February of 2012 from a single pile of fermenting strawberries.  Single mated females

were collected and inbred by 15 generations of full sib mating of their progeny.  *Drosophila* were raised at a

constant temperature of 20º C with 12-hour light/dark cycles. They were raised on a standard glucose/yeast media,

and each library was constructed from adult females of similar age (less than one week).

**Data sources and processing**

The sequencing reads were downloaded from the NCBI Short Read Archive from project SRP075682.

Libraries were assembled using BWA mem (v. 0.7.5), and processed with samtools (v. 0.1.19) using

default parameters[63,64]. The *Wolbachia* reference is the *w*Ri strain previously identified in Southern

California (Accession number NC_012416)[21]. The mtDNA reference is from *D. simulans w$^{501}$*, which is

haplogroup *siII* as expected for *D. simulans* from California (Accession number KC244284)[65]. PCR

duplicates were removed using Picard MarkDuplicates (v. 2.9.4) and GATK (v. 3.7) was used for indel

realignment and SNP calling using default parameters (http://picard.sourceforge.net)[66]. SNPs were called

jointly for all genotypes using Haplotypecaller[66]. Individual consensus fasta sequences were produced

using SelectVariants to create individual vcf files and FastaAlternateReferenceMaker.  Vcf files were

filtered for indels and non-biallelic SNPs using VCFtools (v. 0.1.13)[67]. The files were also filtered for

SNPs with more than 10% missing data. The *Wolbachia* genome was filtered for regions of unusual

coverage or SNP density, for example two regions of the *Wolbachia* genome harbored ~40 SNPs within

two kb, far above background levels of variation (Supp. Fig. 1). These two regions coincided with regions

of unusually high coverage suggesting they are repeated elements.

**Prediction of *Wolbachia* infection status**

*Wolbachia* infection status was determined by calculating the mean depth of coverage of the assembly

and the breadth of coverage of the consensus sequence using bedtools[68]. Depth of coverage refers to the

average read depth across the *Wolbachia* genome, while breadth of coverage refers to the number of bases

covered by at least two reads. Depth of coverage at each nucleotide was estimated using the genomecov

function, while breadth was estimated using the coverage function. Predictions of *Wolbachia* infection

status using illumina data have previously been shown to have 98.8% concordance with PCR based

predication of infection status[32].

**Nucleotide diversity**

Levels of polymorphism for mtDNA and *Wolbachia* were estimated as $\pi$ in 10 kb windows using

VCFtools (v0.1.14)[69]. To investigate whether the frequency spectrum conformed to the standard neutral

model of molecular evolution I also calculated Tajima's *D* in 10 kb windows using VCFtools. To assess

the significance of deviations in Tajima's $D$ and $\pi$ 10,000 simulations were performed using msms

conditioned on the number of variable sites and with no recombination[70].

**Linkage disequilibrium**

Linkage between *Wolbachia* and mtDNA SNPs could potentially be a predictor of co-inheritance of

mtDNA and *Wolbachia*. Linkage was estimated using VCFtools (v0.1.14) using inter-chrom-geno-r2 to

estimate $r^2$ between each SNP in the two genomes[67].

**Estimation of mtDNA and *Wolbachia* copy number**

In insects, the phenotypic effect of *Wolbachia* will vary depending upon copy number in the host cells[9,32].

Given that there are two copies of autosomal DNA in a cell, I infer mtDNA and *Wolbachia* copy number

based on the ratio between mtDNA and autosomal DNA. This is intended to provide a relative estimate of

copy number rather than an absolute measure. Relative copy number estimated in this way obscures intra-

individual variation and variation between tissues, though the authors note that all flies used in

constructing the libraries were females of approximately the same age. *Wolbachia* contains several

regions which were excluded due to unusually high coverage across all samples (more than 3x the mean

coverage). Average coverage of autosomal DNA was calculated from randomly chosen and equivalently

sized nuclear regions for each mtDNA (Scf_3L:8000000..8014945) and *Wolbachia*

(Scf_2L:11000000..11445873). The average coverage of each nuclear region, respectively, was then used

to normalize estimates of copy number for each genotype. Previously the results of measuring *Wolbachia*

copy number in the same samples using both qPCR and estimates from illumina read depth had a

Pearson's correlation coefficient of .79, thus this is a robust approach to measuring *Wolbachia* titre[31].

**Phylogenomic analysis**

To understand the relationship between *Wolbachia* infection and mtDNA I reconstructed the genealogical

history of each within the sample population. Multiple alignments were generated for both mtDNA and

*Wolbachia* by concatenating fasta consensus sequence files for each genotype. All indels and non-biallelic

SNPs were excluded from the dataset prior to generating the consensus fasta for each genotype. RAxML

version 8.10.2 was used to reconstruct phylogenies[71]. Maximum likelihood tree searches were conducted

using a general time reversible (GTR) model of nucleotide substitution with CAT rate heterogeneity and

all model parameters estimated by RAxM[72]. Trees were inferred using the rapid bootstrap algorithm and

simultaneous estimation of trees and bootstrapping, with automatic estimation of the necessary number of

bootstrap replicates.

**Association Analysis**

The association analysis focused on a relationship between nuclear polymorphisms and *Wolbachia* and

mtDNA copy number. To reduce the need for correction due to multiple testing and focus on regions that

may have been affected by selection due to the recent invasion of *Wolbachia* I used a subset of the

nuclear genome identified previously as exhibiting haplotype structure suggestive of recent selection[62,73].

These regions are unusually long haplotype blocks, thus many of the SNPs within each block are not

independent, reducing the need for correction due to multiple testing. Heterozygous bases were coded as

missing, and all loci with more than 10% missing data were excluded from the analysis, as well as SNPs

with a minor allele frequency of less than 2%, meaning they were present in the population in at least 3

copies. mtDNA and *Wolbachia* copy number were used for a multivariate analysis of association using

plink.multivariate[74]. To investigate the possibility that *Wolbachia* copy number is affected by

polymorphisms in mtDNA, and vice versa, a single trait analysis was performed using plink v. 1.07[75].

<u>RESULTS</u>

**Sequencing Data**

The autosomal data included in this analysis was reported in[62]. There was very little variation in both

*Wolbachia* and mtDNA in this population. This included 78 SNPs and indels in the *Wolbachia* genome

and 90 in mtDNA. Reduced diversity has been reported previously in *D. simulans* mtDNA[24,25]. The

authors note that previous work has established that there is no unusual relatedness in the nuclear genome

of this population[62].

**Infection status**

In a previous study lines were scored as infected if they had a breadth of coverage greater than 90% and a

mean depth greater than one[32]. However, that dataset had a clearly bimodal distribution between infected

and uninfected lines, where uninfected lines had breadth of coverage less than 10% while infected lines

had a breadth of coverage of greater than 90%. As such that this demarcation was a natural interpretation

of the data[32]. In *D. simulans*, all lines had ~99% breadth of coverage aside from a single line with both a

lower overall depth of coverage and 80% breadth (Fig. 1). For this reason, all lines were scored as

infected. 100% infection is not unusually high for *D. simulans*.

**Nucleotide Diversity**

Estimates of $\pi$ in *Wolbachia* ranged from $5.98 \times 10^{-7}$ to $1 \times 10^{-3}$, with an average of $1.42 \times 10^{-5}$, within the

range of estimates from *Wolbachia* in *D. melanogaster* from another study ($7.9 \times 10^{-6} - 2.8 \times 10^{-5}$)[31]. The

mean of $\pi$ in simulated populations of *Wolbachia* is $1.9 \times 10^{-3}$ suggesting that variation is somewhat

reduced in *w*Ri. $\pi$ in mtDNA is $1 \times 10^{-4}$ which again is similar to estimates from *D. melanogaster* ($4.34 \times$

$10^{-4} - 1.51 \times 10^{-3}$)[31].

Overall Tajima's *D* was estimated to be -2.4 for *D. simulans* mtDNA (Fig 2). This is similar to

estimates in *D. melanogaster*[32]. Significance of this estimate was assessed using 10,000 simulations in

msms conditioned on the number of segregating sites and no recombination, and it is significant at *p*

< .05. Tajima's *D* in *Wolbachia* is not significantly different from expectations under neutrality based on

10,000 simulations. Thus, while a selective sweep or population bottleneck seems to have strongly

effected mtDNA in *D. simulans*, the same is not true of the *Wolbachia* population (Fig 2). This is very

different from *D. melanogaster* where *Wolbachia* and mtDNA had similar patterns of nucleotide

diversity[32].

This is also much more negative than previously reported for mtDNA in *D. simulans*[25]. It is very

different from the general patterns of Tajima's *D* in the nuclear genome, where average Tajima's *D* is 1

and the majority of the genome has a positive Tajima's *D*. Simulations in previous work suggest that the

pervasively positive values in the nuclear genome may be due to a population contraction, which again

indicates that the population dynamics affecting *D*. *simulans* nuclear and mtDNA genomes are very

different[25,62].

**Linkage disequilibrium**

There was no significant linkage disequilibrium between the genomes of *Wolbachia* and *D*. *simulans*

mtDNA. Average LD between *Wolbachia* and mtDNA SNPs was $2.06 \times 10^{-3}$. This may be because the

infection of *D*. *simulans* was too recent for variation to accumulate along particular lineages, and also

suggests that *D*. *simulans* was infected by a single invasion.

**Estimation of mtDNA and *Wolbachia* copy number**

There was considerable heterogeneity in both *Wolbachia* and mtDNA copy number (Fig. 1). Mean

(standard deviation) copy number of *Wolbachia* is 5.56 (2.45). This is similar to one estimate in *D*.

*melanogaster*, where mean copy number is 5.57 (3.95) though the standard deviation is lower in *D*.

*simulans* [32]. The reported mean was lower in other populations of *D*. *melanogaster*, though still within the

same range $(2 - 4.5)$[31]. Similarly mean mtDNA copy number is 33.85 (15.5) in *D*. *simulans* and 32.9

(44.5) in one estimate for *D*. *melanogaster*[32]. This is again not an absolute measure, but relative to nuclear

genomic coverage. The lower standard deviation could be due to more precise staging of the age of *D*.

*simulans*, less background variation effecting copy number (the *D*. *melanogaster* sample was from

multiple populations), or other unknown mechanisms. There was a positive relationship between mtDNA

and *Wolbachia* copy number (Fig. 1) $(p < 2.4 \times 10^{-7})$. While the functional reasons for or consequences of

this are unclear, because they are correlated they will be used in a multivariate analysis of association

rather than as separate analyses.

**Phylogenomic analysis**

To understand the relationship between *Wolbachia* infection status and mtDNA sequence variation I

reconstructed the phylogenetic history of the complete *Wolbachia* and mtDNA genome using the entire

set of 167 strains (Fig 3-4). What I found is consistent with the recent spread of *Wolbachia* in *D*.

*simulans*, as both phylogenies are essentially unresolved. This is not unexpected for mtDNA given

previous work in the species which found little within-haplotype variation among the three major mtDNA

haplotypes in *D. simulans*[25,28]. Furthermore, of the 167 sequences 88 are identical to at least one other

sequence in the sample. While the *Wolbachia* phylogenetic tree gives the impression of having more

resolution than mtDNA, this is likely due to the larger genome, as the branches have similarly low

support. Of the 167 strains included in the tree 18 are identical to one or more *Wolbachia* genomes. Both

trees are essentially star phylogenies with the majority of bootstrap support values being less than 30.

Bootstrap support of greater than 70, for two branches in the mtDNA tree and five in the *Wolbachia* tree,

is shown (Fig 3-4). If uninfected individuals had been included in the dataset perhaps it would be possible

to test for congruence between the two phylogenies, however the essentially unresolved trees make it

clear that both *Wolbachia* and mtDNA swept the population recently.

**Association Analysis**

Association analysis was performed using plink.multivariate by regressing the line means for mtDNA and

*Wolbachia* copy number on each SNP contained within the previously identified in a scan for selection[62].

This scan for selection focused on identifying haplotype blocks in LD. This considerably reduces the

number of SNPs tested for association, in addition to the fact that the SNPs are in haplotype blocks and

are therefore not independent tests[62,73]. This reduces the need for correction due to multiple testing. I used

a *p*-value cut-off of $p < 9 \times 10^{-6}$ and identified 16 SNPs associated with *Wolbachia* and mtDNA copy

number. Of these 16 SNPs 13 are located in the same region on chromosome 2R (Scf_2R: 13550916-

13569038). Given the concentration of significant SNPs in a single region, this is also the region I will

focus on the most in the following discussion. The region containing 13 SNPs contains nine genes, four of

which are involved in CAAX-box protein processing, *ste24a-c* and a recent duplicate of *ste24c* CG30461.

CAAX-box protein processing is a part of a series of posttranslational protein modifications collectively

called protein prenylation which are required for fully functional proteins to be targeted to cell

membranes or organelles. It has been shown that pathogenic bacteria can exploit the host cell's

prenylation machinery, though it is unclear if this occurs in *Wolbachia*[59].

The other five genes are *AsnRs-m*, which is largely unannotated but is thought to a mitochondrial

aminoacyl-tRNA synthetase[76]. *NIPP1Dm* is involved in axon guidance and negative regulation of protein

phosphorylation[77,78]. *CG6805* is generally unannotated but is inferred to be involved in

dephosporylation[76]. *Cbp53E* regulates neural development[79]. Lastly, *Ehbp1* is a developmental gene

implicated in regulation of the Notch pathway and membrane organization[80].

Of the other three SNPs identified in this association analysis two are located at Scf_2R:5814103

and Scf_2R:5811043, while the third is located at Scf_3L:2055556. Scf_2R:5811043 and

Scf_2R:5814103 are located in *Su(var)2-10* and *Phax*, respectively. These are neighboring genes, though

there is a third gene within 10 kb, *Mys45A*. *Su(var)2-10* is involved in development and chromosome

organization, but it has also been implicated in the regulation of the innate immune response and defense

against gram-negative bacteria[81]. *Su(var)2-10* is of particular interest given that *Wolbachia* are gram-

negative bacteria, however the potential role of *Su(var)2-10* in immune response is not clear. *Phax* is not

well annotated but is inferred to be involved in snRNA export from the nucleus[79]. *Mys45A* is potentially

involved in actin cytoskeleton organization[79]. In *D. melanogaster Wolbachia* uses host actin for maternal

transmission, though this has not been verified in *D. simulans*[82]. The last SNP, at Scf_3L:2055556, is in

*Connectin*, a cell adhesion protein also involved in axon guidance[83].

The identification of these SNPs in association with mtDNA and *Wolbachia* copy number does

not imply a functional relationship. Nonetheless, I chose to investigate whether any of these substitutions

had an effect on the coding sequence of any of genes in the region. Of the three SNPs found outside the

region containing the CAAX-box proteins all were either in introns or regulatory regions. Of the 13 SNPs

identified between Scf_2R: 13550916-13569038 eight are in introns or untranslated regions, including

one in the long intron of *Cb53E*, three in the introns or noncoding transcript of *CG6805*, and two in the

introns of *Ephb*. Of the remaining five SNPs four are in coding regions but silent, causing no change in

the amino acid sequence of the protein. This includes silent mutations in the exons of *ste24c* and two

silent mutations in the exons of *Epbh*. One SNP located in an exon of *ste24a*, at 13558515, is an amino

acid substitution from a Leucine to a Valine. This is not an uncommon amino acid substitution[84,85], though

it can be associated with phenotypes[86,87]. Mutations in introns and untranslated regions could also be

having an effect on gene expression or processing, as could other linked SNPs in the region that were not

included in the analysis.

*Association between Wolbachia and mtDNA*

Association analysis was performed using plink by regressing the line means for mtDNA copy number

onto the *Wolbachia* genome and vice versa[75]. There was no association between *Wolbachia* SNPs and

mtDNA copy number, but the opposite was not true. One SNP in the *D. simulans* mtDNA affected

*Wolbachia* copy number at $p < 3.18 \times 10^{-6}$. It is located in the *D. simulans* homolog of *D. melanogaster*

*srRNA* which has been implicated in pole cell formation[88]. *Wolbachia* is incorporated into the pole cells,

the precursor to the germline, in order to be transmitted[89].

<u>DISCUSSION</u>

Using high through-put sequencing of a large panel of *D. simulans* I have reconstructed the complete

genome sequences of mtDNA and *Wolbachia*. I use these genome sequences to investigate the recent

history of *Wolbachia* and mtDNA in this population, as well as to estimate titre of both *Wolbachia* and

mtDNA. The history of *Wolbachia* in this population is reflected in the essentially star-like phylogeny of

both mtDNA and *Wolbachia*, indicating recent spread and co-inheritance. Lack of variation at mtDNA

and *Wolbachia* suggests a single spread of *w*Ri in this population as well as strict vertical transmission in

the maternal cytoplasm. Variation in *Wolbachia* is within the range expected under a neutral model,

however that was not the case for mtDNA which suggests either a selection sweep or a population

bottleneck. Previous studies found similar population genetic patterns at *Wolbachia* and mtDNA in *D.*

*melanogaster*, and thus could not distinguish whether selection on *Wolbachia* was driving similar patterns

in mtDNA or vice versa[31]. The much stronger pattern of negative Tajima's *D* in the mtDNA suggests that

in *D. simulans* selection is in fact mitochondrial. There was no linkage disequilibrium between *Wolbachia*

and mtDNA variants, however this is most likely due to fixation of a single mitochondrial haplotype

without considerable subsequent mutation.

Currently little is known about how *Wolbachia* interacts with its host[37-39,82,90]. Understanding these

interactions, including regulation of *Wolbachia* titre, will be key to understanding the evolution of

*Wolbachia* and its hosts. By normalizing *Wolbachia* and mtDNA copy number using coverage of the

nuclear genome I am able to obtain estimates of its abundance. Much as in previous work, mtDNA copy

number was higher than *Wolbachia* copy number, though both varied across strains[32]. As all of my data

was produced from adult females, at the same time, using the same techniques, there is no danger that this

is due to differences in methodology among samples[32]. Estimates of copy number were very similar to

previous work in *D. melanogaster*, performed with qPCR, and there has been shown to be a high

correlation between qPCR and illumina estimates of copy number[31,32]. These are not absolute measures,

rather they are relative to one another and to nuclear copy number, and they provide robust estimates of

*Wolbachia* titre within the population. As the *Wolbachia* phylogenetic tree is essentially unresolved in

this population but there is considerable variation in *Wolbachia* titre, it is clear that some host factors

must be affecting variation in *Wolbachia* titre.

The history of mtDNA and the nuclear genome is quite divergent in this population. The nuclear

genome has an average Tajima's $D$ of 1 and 5 polymorphisms for every 100 bp[62]. Simulations suggest

that this is due to a combination of population contraction and selection, most likely from standing

variation, though many types of sweeps can produce similar signatures[62]. In contrast the mtDNA genome

contains an abundance of low frequency variation, and in fact many of the mtDNA genomes sampled in

this population are identical. This is consistent with the recent spread, single origin, and maternal

transmission, of *w*Ri in *D. simulans*. This is consistent with previous work which found low levels of

mtDNA variation in *D. simulans* within a haplotype[24,91]. This is also consistent with work on *Wolbachia*

which documented the spread of *w*Ri in *D. simulans* in the 1980's[4,5,8,18-20].

While it has been proposed elsewhere, the author is not aware of another association analysis of

*Wolbachia* and mtDNA copy number[32]. *Wolbachia* copy number is known to be affected by host

background, but the genes or mechanisms involved are not known[54,55,57]. The fact that four of the nine

genes found in the primary region detected in the association analysis are involved in CAAX-box protein

processing is of particular interest, given the history of this type of gene and intracellular pathogens.

CAAX-box protein processing is a part of a series of posttranslational protein modifications collectively

called protein prenylation which are required for fully functional proteins to be targeted to cell

membranes or organelles. Prenylated proteins include Ras, Rac, and Rho. However, it has been shown

that pathogenic bacteria can exploit the host cell's prenylation machinery[59]. For example, *Salmonella-*

*induced filament A* is a protein from *Salmonella typhimurium*, a gram-negative facultative intracellular

bacterium. *Salmonella-induced filament A* has a CAAX motif required for prenylation to occur, it was

shown to be processed by host prenylation machinery, and it is necessary for survival of the

bacterium[60,92,93]. *Legionella pneumophila* Ankyrin B protein exploits the host prenylation machinery in

order to anchor Ankyrin B protein to the membrane of the pathogenic vacuole[61]. Proliferation of

*Legionella pneumophila* requires Ankyrin B, as does the manifestation of Legionnaires disease. Ankyrin

repeat domains are most commonly found in eukaryotes and viruses, though they are rarely found in

bacteria and Archaea[94]. In bacteria they are found in a few obligate or facultative intracellular

Proteobacteria[59]. *Wolbachia* has an unusually high number of Ankyrin repeat domains with rapid

evolution[94]. Ankyrin proteins play a major role in host-pathogen interactions and the evolution of

infections[95,96]. There is no way to know from the current analysis if the Ankyrin repeat genes are

exploiting the host prenylation system but it is an intriguing area for future investigation. The results of

this association analysis suggest that some interaction between the pathogen and its host is targeting the

protein prenylation machinery.

There was also an association between a polymorphism in *srRNA,* which has been implicated in

pole cell formation[88], and *Wolbachia* copy number. Concentration of *Wolbachia* in the posterior of the

embryo, where pole cells are forming, is correlated with degree of cytoplasmic incompatibility[97]. *D.*

*simulans* has been shown to have nearly complete cytoplasmic incompatibility, though it is possible there

are mutations sorting at low frequency that affect this or that mitigate negative phenotypic consequences

of high *Wolbachia* titre. It has also been demonstrated that *gurken* is important for *Wolbachia* titre in the

germline in *D. melanogaster*, and it is involved in pole cell formation beginning at an earlier stage than

*sr*RNA suggesting there could be an interaction between the two factors[88,90]. *D. simulans w*Ri has a

different distribution in the cytoplasm from other strains of *Wolbachia*, as it tends to evenly distribute

throughout the embryo while other strains are either concentrated at the posterior, or at the anterior of the

embryo away from the pole cells[97]. Future work in related species may show that these different

distributions also mitigate different interactions between host and symbiont, including being effected by

different genes and processes within the host.

382
383

**Figure 1. *Wolbachia* infection status and relationship to mtDNA copy number A.** Relationship between depth and breadth of sequencing coverage for *Wolbachia* assemblies in the *D. simulans* panel. Depth of coverage is shown in $\log_{10}$ unites and is calculated as the number of reads present at each nucleotide in the reference averaged over every site. Breadth of coverage is the proportion of covered nucleotides in the consensus sequence relative to the reference. **B.** Relationship between relative mtDNA copy number and *Wolbachia* copy number. Both were normalized relative to nuclear coverage. Although separate regions were used to normalize mtDNA and *Wolbachia*, as they are different sizes, average values were very similar within genotypes. The relationship between mtDNA and *Wolbachia* copy number is positive ($p < 2.4 \times 10^{-7}$).

**Figure 2. *Wolbachia* and mtDNA Tajima's *D*.** 10,000 simulations were performed for *Wolbachia* and *D. simulans* each conditioned upon the number of polymorphisms. The actual values in *D. simulans* mtDNA are outside the 95% confidence interval of the simulations, while *Wolbachia* is not. There is considerable variation in Tajima's *D* across the *Wolbachia* genome while mtDNA is much smaller and invariant in its values of Tajima's *D*.

**Figure 3: Maximum likelihood genealogy of the *D. simulans Wolbachia* pathogen**. All strains were infected with *Wolbachia* and are included in this geneaology. The underlying data consist of an ungapped multiple alignment of 168 sequences of the entire *Wolbachia* genome. The unrooted tree was midpoint rooted for visualization and branches with > 70% RAxML bootstrap support values are shown in bold. Scale bars for branch lengths are in term of mutations per site. The majority of branches are essentially unsupported by bootstrapping.

5.0E-5

**Figure 4: Maximum likelihood genealogy of the *D. simulans* mtDNA genome**. The underlying data consist of an ungapped multiple alignment of 168 sequences of the entire mtDNA genome. The unrooted tree was midpoint rooted for visualization and branches with > 70% RAxML bootstrap support values are shown in bold. Scale bars for branch lengths are in term of mutations per site. The tree is largely unresolved, suggesting recent spread of this mtDNA haplotype through the population.

395
396
397   1.      Zug, R. & Hammerstein, P. Still a host of hosts for *Wolbachia*: analysis of recent data
398          suggests that 40% of terrestrial arthropod species are infected. *PLoS ONE* **7,** e38544 (2012).
399   2.      Hilgenboecker, K., Hammerstein, P., Schlattmann, P., Telschow, A. & Werren, J. H. How

400      many species are infected with *Wolbachia*?--A statistical analysis of current data. *FEMS*
401      *Microbiol. Lett.* **281,** 215–220 (2008).

402 3.   Jeyaprakash, A. & Hoy, M. A. Long PCR improves *Wolbachia* DNA amplification: wsp
403      sequences found in 76% of sixty-three arthropod species. *Insect Mol. Biol.* **9,** 393–405
404      (2000).

405 4.   Turelli, M. & Hoffmann, A. A. Cytoplasmic incompatibility in *Drosophila simulans*:
406      dynamics and parameter estimates from natural populations. *Genetics* **140,** 1319–1338
407      (1995).

408 5.   Turelli, M., Hoffmann, A. A. & McKechnie, S. W. Dynamics of cytoplasmic
409      incompatibility and mtDNA variation in natural *Drosophila simulans* populations. *Genetics*
410      **132,** 713–723 (1992).

411 6.   Charlat, S., Nirgianaki, A., Bourtzis, K. & Merçot, H. Evolution of *Wolbachia*-induced
412      cytoplasmic incompatibility in *Drosophila simulans* and *D. sechellia*. *Evolution* **56,** 1735–
413      1742 (2002).

414 7.   Rousset, F., Vautrin, D. & Solignac, M. Molecular identification of *Wolbachia,* the agent of
415      cytoplasmic incompatibility in *Drosophila simulans*, and variability in relation with host
416      mitochondrial types. *Proc. R. Soc. B* **247,** 163–168 (1992).

417 8.   Hoffmann, A. A., Turelli, M. & Harshman, L. G. Factors affecting the distribution of
418      cytoplasmic incompatibility in *Drosophila simulans*. *Genetics* **126,** 933–948 (1990).

419 9.   Berticat, C., Rousset, F., Raymond, M., Berthomieu, A. & Weill, M. High *Wolbachia*
420      density in insecticide-resistant mosquitoes. *Proc. R. Soc. B* **269,** 1413–1416 (2002).

421 10.  Wong, Z. S., Hedges, L. M., Brownlie, J. C. & Johnson, K. N. *Wolbachia*-mediated
422      antibacterial protection and immune gene regulation in *Drosophila*. *PLoS ONE* **6,** e25430–9
423      (2011).

424 11.  Osborne, S. E., Leong, Y. S., O'Neill, S. L. & Johnson, K. N. Variation in antiviral
425      protection mediated by different *Wolbachia* strains in *Drosophila simulans*. *PLoS Pathog* **5,**
426      e1000656–9 (2009).

427 12.  Fytrou, A., Schofield, P. G., Kraaijeveld, A. R. & Hubbard, S. F. *Wolbachia* infection
428      suppresses both host defence and parasitoid counter-defence. *Proc. R. Soc. B* **273,** 791–796
429      (2006).

430 13.  Brownlie, J. C. *et al*. Evidence for metabolic provisioning by a common invertebrate
431      endosymbiont, *Wolbachia pipientis*, during periods of nutritional stress. *PLoS Pathog* **5,**
432      e1000368 (2009).

433 14.  Teixeira, L., Ferreira, Á. & Ashburner, M. The bacterial symbiont *Wolbachia* induces
434      resistance to RNA viral infections in *Drosophila melanogaster*. *PLoS Biol.* **6,** e1000002
435      (2008).

436 15.  Hedges, L. M., Brownlie, J. C. & O'Neill, S. L. *Wolbachia* and virus protection in insects.
437      **322,** 702 *Science* (2008).

438 16.  Jiggins, F. M., Hurst, G. & Majerus, M. Sex-ratio-distorting *Wolbachia* causes sex-role
439      reversal in its butterfly host B. *Proc. R. Soc. B* **267,** 69–73.

440 17.  Koukou, K. *et al*. Influence of antibiotic treatment and *Wolbachia* curing on sexual isolation
441      among *Drosophila melanogaster* cage populations. *Evolution* **60,** 87–11 (2006).

442 18.  Turelli, M. & Hoffmann, A. A. Rapid spread of an inherited incompatibility factor in
443      California *Drosophila*. *Nature* **353,** 440–442 (1991).

444 19.  Hoffmann, A. A. & Turelli, M. Unidirectional incompatibility in *Drosophila simulans*:
445      inheritance, geographic variation and fitness effects. *Genetics* **119,** 435–444 (1988).

446 20.  Hoffmann, A. A., Turelli, M. & Simmons, G. M. Unidirectional incompatibility between
447      populations of *Drosophila simulans*. *Evolution* **40,** 692–701 (1986).

448 21.  Klasson, L. *et al*. The mosaic genome structure of the *Wolbachia w*Ri strain infecting
449      *Drosophila simulans*. *Proc. Nat. Acad. Sci. USA* **106,** 5725–5730 (2009).

450 22.  Choi, J. Y. & Aquadro, C. F. The coevolutionary period of *Wolbachia pipientis* infecting

*Drosophila ananassae* and its impact on the evolution of the host germline stem cell
regulating genes. *Mol. Biol. Evol.* **31,** 2457–2471 (2014).
23. Ballard, J. W. O. Comparative genomics of mitochondrial DNA in *Drosophila simulans*. *J*
*Mol Evol* **51,** 64–75 (2000).
24. Ballard, J. W., Hatzidakis, J., Karr, T. L. & Kreitman, M. Reduced variation in *Drosophila*
*simulans* mitochondrial DNA. *Genetics* **144,** 1519–1528 (1996).
25. Ballard, J. W. O. Comparative genomics of mitochondrial DNA in *Drosophila simulans*. *J*
*Mol Evol* **51,** 64–75 (2000).
26. Solignac, M., Vautrin, D. & Rousset, F. Widespread occurence of the proteobacteria
*Wolbachia* and partial cytoplasmic incompatibility in *Drosophila melanogaster*. C. R. Acad.
Sci. **317,** 461-479 (1994).
27. Solignac, M. & Monnerot, M. Race formation, speciation, and introgression within
*Drosophila simulans*, *D. mauritiana*, and *D. sechellia* inferred from mitochondrial DNA
analysis. *Evolution* **40,** 531–539 (1986).
28. Baba-Aïssa, F., Solignac, M., Dennebouy, N. & David, J. R. Mitochondrial DNA variability
in *Drosophila simulans*: quasi absence of polymorphism within each of the three
cytoplasmic races. *Heredity* **61,** 419–426 (1988).
29. Montchamp-Moreau, C., Ferveur, J. F. & Jacques, M. Geographic distribution and
inheritance of three cytoplasmic incompatibility types in *Drosophila simulans*. *Genetics*
**129,** 399–407 (1991).
30. James, A. C. & Ballard, J. W. Expression of cytoplasmic incompatibility in *Drosophila*
*simulans* and its impact on infection frequencies and distribution of *Wolbachia pipientis*.
*Evolution* **54,** 1661–1672 (2000).
31. Early, A. M. & Clark, A. G. Monophyly of *Wolbachia pipientis* genomes within *Drosophila*
*melanogaster*: geographic structuring, titre variation and host effects across five
populations. *Mol. Ecol.* **22,** 5765–5778 (2013).
32. Richardson, M. F. *et al.* Population genomics of the *Wolbachia* endosymbiont in *Drosophila*
*melanogaster*. *PLoS Genet.* **8,** e1003129 (2012).
33. Nunes, M. D. S., Nolte, V. & Schl tterer, C. Nonrandom *Wolbachia* infection status of
*Drosophila melanogaster* strains with different mtDNA haplotypes. *Mol. Biol. Evol.* **25,**
2493–2498 (2008).
34. Beckmann, J. F., Ronau, J. A. & Hochstrasser, M. A *Wolbachia* deubiquitylating enzyme
induces cytoplasmic incompatibility. *Nat Microbiol* **2,** 17007 (2017).
35. Civetta, A. Rapid Evolution and gene-specific patterns of selection for three genes of
spermatogenesis in *Drosophila*. *Mol. Biol. Evol.* **23,** 655–662 (2005).
36. Bauer DuMont, V. L., Flores, H. A., Wright, M. H. & Aquadro, C. F. Recurrent positive
selection at *bgcn*, a key determinant of germ line differentiation, does not appear to be
driven by simple coevolution with its partner protein *bam*. *Mol. Biol. Evol.* **24,** 182–191
(2006).
37. Flores, H. A., Bubnell, J. E., Aquadro, C. F. & Barbash, D. A. The *Drosophila bag of*
*marbles* gene interacts genetically with *Wolbachia* and shows female-specific effects of
divergence. *PLoS Genet.* **11,** e1005453 (2015).
38. Serbus, L. R. & Sullivan, W. A Cellular basis for *Wolbachia* recruitment to the host
germline. *PLoS Pathog* **3,** e190 (2007).
39. Serbus, L. R., Casper-Lindley, C., Landmann, F. & Sullivan, W. The genetics and cell
biology of *Wolbachia*-host interactions. *Annu. Rev. Genet.* **42,** 683–707 (2008).
40. McGraw, E. A., Merritt, D. J., Droller, J. N. & O'Neill, S. L. *Wolbachia* density and
virulence attenuation after transfer into a novel host. *Proc. Nat. Acad. Sci. USA* **99,** 2918–
2923 (2002).
41. Bordenstein, S. R., Marshall, M. L., Fry, A. J., Kim, U. & Wernegreen, J. J. The tripartite
associations between Bacteriophage, *Wolbachia*, and Arthropods. *PLoS Pathog* **2,** e43–10

(2006).

42.   Martinez, J. *et al.* Should symbionts be nice or selfish? Antiviral effects of *Wolbachia* are
costly but reproductive parasitism is not. *PLoS Pathog* **11,** e1005021 (2015).

43.   Clark, M. E., Veneti, Z., Bourtzis, K. & Karr, T. L. *Wolbachia* distribution and cytoplasmic
incompatibility during sperm development: the cyst as the basic cellular unit of CI
expression. *Mech. Dev.* **120,** 185–198 (2003).

44.   Chrostek, E., Marialva, M. S. P., Yamada, R., O'Neill, S. L. & Teixeira, L. High anti-viral
protection without immune upregulation after interspecies *Wolbachia* transfer. *PLoS ONE* **9,**
e99025 (2014).

45.   Chrostek, E. & Teixeira, L. Mutualism breakdown by amplification of *Wolbachia* genes.
*PLoS Biol.* **13,** e1002065 (2015).

46.   Chrostek, E. *et al.* *Wolbachia* variants induce differential protection to viruses in *Drosophila*
*melanogaster*: A phenotypic and phylogenomic analysis. *PLoS Genet.* **9,** e1003896 (2013).

47.   Hoffmann, A. A. *et al.* Successful establishment of *Wolbachia* in *Aedes* populations to
suppress dengue transmission. *Nature* **476,** 454–457 (2011).

48.   McMeniman, C. J. *et al.* Host adaptation of a *Wolbachia* strain after long-term serial
passage in mosquito cell lines. *App. Environ. Microbiol.* **74,** 6963–6969 (2008).

49.   Osborne, S. E., Iturbe-Ormaetxe, I., Brownlie, J. C., O'Neill, S. L. & Johnson, K. N.
Antiviral protection and the importance of *Wolbachia* density and tissue tropism in
*Drosophila simulans*. *App. Environ. Microbiol.* **78,** 6922–6929 (2012).

50.   Kondo, N., Shimada, M. & Fukatsu, T. Infection density of *Wolbachia* endosymbiont
affected by co-infection and host genotype. *Biol. Lett.* **1,** 488–491 (2005).

51.   Reynolds, K. T., Thomson, L. J. & Hoffmann, A. A. The effects of host age, host nuclear
background and temperature on phenotypic effects of the virulent *Wolbachia* strain popcorn
in *Drosophila melanogaster*. *Genetics* **164,** 1027–1034 (2003).

52.   Olsen, K., Reynolds, K. T. & Hoffmann, A. A. A field cage test of the effects of the
endosymbiont *Wolbachia* on *Drosophila melanogaster*. *Heredity* (2001).

53.   Boyle, L., O'Neill, S. L., Robertson, H. M. & Karr, T. L. Interspecific and intraspecific
horizontal transfer of *Wolbachia* in *Drosophila*. *Science* **260,** 1796–1799 (1993).

54.   Poinsot, D., Bourtzis, K., Markakis, G. & Savakis, C. *Wolbachia* transfer from *Drosophila*
*melanogaster* into *D. simulans*: host effect and cytoplasmic incompatibility relationships.
*Genetics* (1998).

55.   McGraw, E. A., Merritt, D. J., Droller, J. N. & O'Neill, S. L. *Wolbachia*-mediated sperm
modification is dependent on the host genotype in *Drosophila*. *Proc. R. Soc. B* **268,** 2565–
2570 (2001).

56.   Poinsot, D., Montchamp-Moreau, C. & Mercot, H. *Wolbachia* segregation rate in
*Drosophila simulans* naturally bi-infected cytoplasmic lineages. *Heredity* **85 ( Pt 2),** 191–
198 (2000).

57.   Boyle, L., O'Neill, S. L., Robertson, H. M. & Karr, T. L. Interspecific and intraspecific
horizontal transfer of *Wolbachia* in *Drosophila*. *Science* **260,** 1796–1799 (1993).

58.   Perrot-Minnot & Werren. *Wolbachia* infection and incompatibility dynamics in
experimental selection lines. *J. Evol. Biol.* **12,** 272–282 (1999).

59.   Amaya, M., Baranova, A. & van Hoek, M. L. Protein prenylation: A new mode of host-
pathogen interaction. *Biochemical and Biophysical Research Communications* **416,** 1–6
(2011).

60.   Reinicke, A. T. *et al.* A *Salmonella typhimurium* effector protein SifA is modified by host
cell prenylation and s-acylation machinery. *J. Biol. Chem.* **280,** 14620–14627 (2005).

61.   Price, C. T. D., Al-Quadan, T., Santic, M., Jones, S. C. & Abu Kwaik, Y. Exploitation of
conserved eukaryotic host cell farnesylation machinery by an F-box effector of *Legionella*
*pneumophila*. *J Exp Med* **207,** 1713–1726 (2010).

62.   Signor, S. A., New, F. & Nuzhdin, S. *in review*. An abundance of high frequency variance

| 553 | | uncovered in a large panel of *Drosophila simulans*. |
|-----|-----|----------------------------------------------------------|
| 554 | 63. | Li, H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM |
| 555 | | *arXiv*. 1–3 (2015). |
| 556 | 64. | Li, H. *et al*. The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25,** 2078– |
| 557 | | 2079 (2009). |
| 558 | 65. | Meiklejohn, C. D. *et al*. An Incompatibility between a mitochondrial tRNA and its nuclear- |
| 559 | | encoded tRNA synthetase compromises development and fitness in *Drosophila*. *PLoS* |
| 560 | | *Genet*. **9,** e1003238 (2013). |
| 561 | 66. | McKenna, A. *et al*. The Genome Analysis Toolkit: a MapReduce framework for analyzing |
| 562 | | next-generation DNA sequencing data. *Genome Res*. **20,** 1297–1303 (2010). |
| 563 | 67. | Danecek, P. *et al*. The variant call format and VCFtools. *Bioinformatics* **27,** 2156–2158 |
| 564 | | (2011). |
| 565 | 68. | Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic |
| 566 | | features. *Bioinformatics* **26,** 841–842 (2010). |
| 567 | 69. | Tajima, F. Statistical method for testing the neutral mutation hypothesis by DNA |
| 568 | | polymorphism. *Genetics* **123,** 585–595 (1989). |
| 569 | 70. | Ewing, G. & Hermisson, J. MSMS: a coalescent simulation program including |
| 570 | | recombination, demographic structure and selection at a single locus. *Bioinformatics* **26,** |
| 571 | | 2064–2065 (2010). |
| 572 | 71. | Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large |
| 573 | | phylogenies. *Bioinformatics* **30,** 1312–1313 (2014). |
| 574 | 72. | Izquierdo-Carrasco, F., Smith, S. A. & Stamatakis, A. Algorithms, data structures, and |
| 575 | | numerics for likelihood-based phylogenetic inference of huge trees. *BMC Bioinformatics* |
| 576 | | **12,** 470 (2011). |
| 577 | 73. | Garud, N. R., Messer, P. W., Buzbas, E. O. & Petrov, D. A. Recent selective sweeps in |
| 578 | | North American *Drosophila melanogaster* show signatures of soft sweeps. *PLoS Genet*. **11,** |
| 579 | | e1005004 (2015). |
| 580 | 74. | Ferreira, M. A. R. & Purcell, S. M. A multivariate test of association. *Bioinformatics* **25,** |
| 581 | | 132–133 (2008). |
| 582 | 75. | Purcell, S. *et al*. PLINK: A tool set for whole-genome association and population-based |
| 583 | | linkage analyses. *Am*. *J*. *Hum*. *Genet*. **81,** 559–575 (2007). |
| 584 | 76. | FlyBase Curators, Swiss-Prot Project Members InterPro Project Members. Gene Ontology |
| 585 | | annotation in FlyBase through association of InterPro records with GO terms. (2004). |
| 586 | 77. | Babu, K., Bahri, S., Alphey, L. & Chia, W. Bifocal and PP1 interaction regulates targeting |
| 587 | | of the R-cell growth cone in *Drosophila*. *Dev*. *biol*. **288,** 372–386 (2005). |
| 588 | 78. | Bennett, D., Szoor, B., Gross, S., Vereshchagina, N. & Alphey, L. Ectopic expression of |
| 589 | | inhibitors of Protein phosphatase type 1 (PP1) can be used to analyze roles of PP1 in |
| 590 | | *Drosophila* development. *Genetics* **164,** 235–245 (2003). |
| 591 | 79. | Gaudet, P., Livstone, M. & Thomas, P. Gene Ontology annotation inferences using |
| 592 | | phylogenetic trees. GO Reference Genome Project. (2010). |
| 593 | 80. | Giagtzoglou, N. *et al*. *dEHBP1* controls exocytosis and recycling of *Delta* during |
| 594 | | asymmetric divisions. *J*. *Cell Biol*. **196,** 65–83 (2012). |
| 595 | 81. | Cronin, S. J. F. *et al*. Genome-wide RNAi screen identifies genes involved in intestinal |
| 596 | | pathogenic bacterial infection. *Science* **325,** 340–343 (2009). |
| 597 | 82. | Newton, I. L. G., Savytskyy, O. & Sheehan, K. B. *Wolbachia* utilize host actin for efficient |
| 598 | | maternal transmission in *Drosophila melanogaster*. *PLoS Pathog* **11,** e1004798 (2015). |
| 599 | 83. | Chiba, A. Early development of the *Drosophila* neuromuscular junction: a model for |
| 600 | | studying neuronal networks in development. *Int*. *Rev*. *Neurobiol*. (1999). |
| 601 | 84. | Yampolsky, L. Y. & Stoltzfus, A. The exchangeability of amino acids in proteins. *Genetics* |
| 602 | | **170,** 1459–1472 (2005). |
| 603 | 85. | Creixell, P., Schoof, E. M., Tan, C. S. H. & Linding, R. Mutational properties of amino acid |

604    residues: implications for evolvability of phosphorylatable residues. *Philos. Trans. R. Soc. B*
605    **367,** 2584–2593 (2012).
606    86.    Leone, A. *et al*. Evidence for nm23 RNA overexpression, DNA amplification and mutation
607           in aggressive childhood neuroblastomas. *Oncogene* **8,** 855–865 (1993).
608    87.    Ishiko, A. *et al*. A novel leucine to valine mutation in residue 7 of the helix initiation motif
609           of Keratin10 leads to bullous congenital ichthyosiform erythroderma. *J. Inv. Dermatol.* **116,**
610           991–992 (2001).
611    88.    Mahowald, A. P. Assembly of the *Drosophila* germ plasm. *Int. Rev. Cytol.* **203,** 187–213
612           (2001).
613    89.    Kose, H. & Karr, T. L. Organization of *Wolbachia pipientis* in the *Drosophila* fertilized egg
614           and embryo revealed by an anti-*Wolbachia* monoclonal antibody. *Mech. Dev.* **51,** 275–288
615           (1995).
616    90.    Serbus, L. R. *et al*. A feedback loop between *Wolbachia* and the *Drosophila gurken* mRNP
617           complex influences *Wolbachia* titer. *J. Cell Sci.* **124,** 4299–4308 (2012).
618    91.    Ballard, J. W. O. Comparative genomics of mitochondrial DNA in *Drosophila simulans*. *J.*
619           *Mol. Evol.* **51,** 64–75 (2000).
620    92.    Vinh, D. B. N., Ko, D. C., Rachubinski, R. A., Aitchison, J. D. & Miller, S. I. Expression of
621           the *Salmonella spp*. virulence factor SifA in yeast alters Rho1 activity on peroxisomes. *Mol.*
622           *Biol. Cell* **21,** 3567–3577 (2010).
623    93.    Brumell, J. H., Goosney, D. L. & Finlay, B. B. SifA, a type III secreted effector of
624           *Salmonella typhimurium*, directs *Salmonella*-induced filament (Sif) formation along
625           microtubules. *Traffic* **3,** 407–415 (2002).
626    94.    Siozios, S. *et al*. The diversity and evolution of *Wolbachia* ankyrin repeat domain genes.
627           *PLoS ONE* **8,** e55390 (2013).
628    95.    Bork, P. Hundreds of ankyrin-like repeats in functionally diverse proteins: mobile modules
629           that cross phyla horizontally? *Proteins* **17,** 363–374 (1993).
630    96.    Habyarimana, F. *et al*. Role for the Ankyrin eukaryotic-like genes of *Legionella*
631           *pneumophila* in parasitism of protozoan hosts and human macrophages. *Environ. Microbiol.*
632           **10,** 1460–1474 (2008).
633    97.    Veneti, Z., Clark, M. E., Karr, T. L., Savakis, C. & Bourtzis, K. Heads or Tails: Host-
634           Parasite Interactions in the *Drosophila-Wolbachia* System. *App. Env. Microbiol.* **70,** 5366–
635           5372 (2004).
636

637    **Acknowledgements**
638    I would like to thank J. Butler for helpful commentary on this manuscript, and S. Nuzhdin for helpful
639    discussion and direction.
640
641    **Author Contributions**
642    S. S. performed all of the work described herein.
643
644    **Competing financial interest**
645    The author has no competing financial interests.