1 **Spring flowering habit in field pennycress (*Thlaspi arvense*) has arisen multiple**
2 **independent times**
3

Kevin M. Dorn[1], Evan B. Johnson[1], Erin Daniels[1], Donald L. Wyse[2], M. David
Marks[1]
[1]Department of Plant and Microbial Biology, University of Minnesota,
1445 Gortner Avenue, 250 Biological Sciences Center, Saint Paul, Minnesota 55108
[2]Department of Agronomy and Plant Genetics, University of Minnesota,
411 Borlaug Hall, 1991 Upper Buford Circle, Saint Paul, Minnesota 55108
10

Email addresses:
KMD: dorn@umn.edu
EBJ: john6551@umn.edu
ED: danie711@umn.edu
DLW: wysex001@umn.edu
MDM: marks004@umn.edu
17

Correspondence to:
M David Marks
University of Minnesota
Department of Plant Biology
250 Biological Sciences Center
1445 Gortner Avenue
St Paul, Minnesota 55108
Marks004@umn.edu
26

Word Count, Summary: 200
Word Count, Introduction: 1050
Word Count, Materials and Methods: 1159
Word Count, Results: 1630
Word Count, Discussion: 910
Word Count, Acknowledgements: 98
Total Word Count: 5065
34

35 2 Figures (both in colour)
36 6 Supporting Figures
37 3 Supporting Tables

Summary

- Pennycress (*Thlaspi arvense* L.) is currently being developed as a new cold-season oilseed crop. Like many Brassicaceae, pennycress can exhibit either a winter or spring annual phenotype. In Arabidopsis, mutations in negative regulators of flowering, including FLOWERING LOCUS C (FLC) and FRIGIDA can cause the transition to a spring annual habit. The genetics underlying the difference between spring and winter annual pennycress are currently unknown.

- Using whole genome sequencing across wild spring annual pennycress accessions, co-segregation analyses, and comparative genomics approaches, we identify new alleles of TaFLC and explore their geographic distribution.

- We report that loss of function mutations in TaFLC confer the spring annual phenotype in pennycress. We have also identified four natural alleles of TaFLC that confer a spring annual growth habit. The two spring annual FLC alleles present in European accessions were only identified in accessions collected in Montana, USA.

- In pennycress, the spring annual habit has arisen several independent times. Accessions harboring the two European alleles were introduced to North America, likely after the species became a widespread on the continent. These findings provide new information on the natural history of the introduction and spread of the spring annual phenotype in pennycress.

Key Words: *Thlaspi arvense*, pennycress, Flowering Locus C, spring annual, winter annual, whole genome sequencing

Introduction

Thlaspi arvense L. (field pennycress, pennycress herein) is currently a target for domestication as a new, cold-hardy winter oilseed crop that can fit within the corn/soybean rotation in the Midwestern United States (Dorn *et al.*, 2013; Sedbrook, 2014; Dorn *et al.*, 2015). As a winter annual crop, pennycress can be planted in the late summer to early autumn, either into standing corn or immediately after corn harvest. The pennycress crop establishes a robust vegetative cover prior to winter, providing important ecosystem services such as limiting soil erosion and nutrient runoff. In the spring, pennycress will flower and set seed, yielding upwards of 1,600 pounds per acre of oil-rich seed in time for planting a crop of soybean (Phippen & Phippen, 2012; Sedbrook, 2014). Pennycress seed is high in oils that can be readily converted to biodiesel or jet fuel, with the remaining high-protein seed meal having potential as animal feed or feedstock for industrial uses (Moser *et al.*, 2009; Selling *et al.*, 2011; Evangelista *et al.*, 2012; Fan *et al.*, 2013; Hojilla-Evangelista *et al.*, 2013).

Pennycress is a member of the Thlaspideae, a tribe of the Brassicaceae (Beilstein *et al.*, 2008). The Brassicaceae family is divided into three lineages. Pennycress resides in lineage II, along with the Brassica genus, including the economically important oilseed crops *Brassica rapa* and *Brassica napus*, but not the model species *Arabidopsis thaliana* nor *Capsella rubella*, which are members of lineage I (Franzke *et al.*, 2011). Pennycress is native to Eurasia and is considered naturalized to most temperate to subarctic regions throughout the northern hemisphere, including all of the United States except for Hawaii and Alabama, all provinces of Canada, as well as being considered naturalized in the southern hemisphere, including Australia, New Zealand, and Argentina (Warwick, 2002). Pennycress exhibits numerous traits that make it an attractive winter rotation crop (Sedbrook, 2014). Of particular importance in the upper Midwest is the existence of pennycress lines that complete their life cycle rapidly enough to fit within the corn/soybean rotation. Central to this rapid spring development of pennycress are its underlying growth habits. In wild populations, there are both winter and spring annual

pennycress (Best & McIntyre, 1972; McIntyre & Best, 1978), similar to many other

Brassicaceae species such as *Arabidopsis thaliana* (Stinchcombe *et al.*, 2004; Shindo *et*

al., 2005), *Brassica rapa* (Wu *et al.*, 2012), *Camelina sativa* (Crowley, 1999), and

Brassica napus (Tadege *et al.*, 2001).

Throughout the decades of research on basic developmental questions in Arabidopsis,

and the expanding translational research in other Brassica crops, the underlying

molecular mechanisms controlling flowering time have been identified in many of these

species (Simpson & Dean, 2002; Amasino, 2005; Jung & Muller, 2009; Kim *et al.*, 2009).

In wild accessions of Arabidopsis, only a handful of mutations are responsible for

variation of flowering time and vernalization requirement (Burn *et al.*, 1993; Clarke &

Dean, 1994; Johanson *et al.*, 2000). Most notably, allelic variation in two key negative

regulators, FLOWERING LOCUS C (FLC) and/or FRIGIDA (FRI), underlies the key

difference between spring and winter annual plants (Johanson *et al.*, 2000; Michaels &

Amasino, 2001; Michaels *et al.*, 2003; Michaels *et al.*, 2004; Shindo *et al.*, 2005; Jiang *et*

al., 2009). FLC encodes a MADS box transcription factor (Michaels & Amasino, 1999),

and inhibits flowering prior to vernalization by repressing the expression of

FLOWERING LOCUS T (FT) (Searle *et al.*, 2006). In Arabidopsis, *flc* null mutations

eliminate the vernalization requirement and impart the spring annual, rapid-flowering

phenotype (Michaels & Amasino, 1999). Allelic variation within FRI also impacts

flowering time and the vernalization requirement. Similar to FLC, Arabidopsis accessions

harboring loss of function mutations in FRI flower rapidly without vernalization

(Johanson *et al.*, 2000; Shindo *et al.*, 2005).

The expression of FLC is positively regulated by FRI (Michaels & Amasino, 1999), thus,

mutations in either of these vernalization-responsive negative regulators can lead to a loss

of vernalization requirement and a spring annual growth habit (Michaels & Amasino,

1999; Michaels & Amasino, 2001). The vernalization signal provided by the cold of

winter removes the repression on the transition to flowering through the epigenetic

silencing of FLC. Specifically, vernalization increases histone 3 K27 trimethylation

(H3K27me3) at FLC chromatin, reducing transcriptional activity (Sung *et al.*, 2006;

Finnegan & Dennis, 2007; Greb *et al.*, 2007; Coustham *et al.*, 2012; Yang *et al.*, 2014).

The vernalization-induced silencing of FLC releases the repression on the transition to

flowering, which permits the transition to reproductive growth (Searle *et al.*, 2006).

Additionally, the roles of the COOLAIR long noncoding antisense transcripts transcribed

from the AtFLC locus have been revealed as a key regulatory component in the cold-

induced regulation of FLC (Swiezewski *et al.*, 2009; Csorba *et al.*, 2014; Marquardt *et al.*,

2014).

While this extensive understanding of the molecular genetic pathways controlling

flowering time and vernalization in Arabidopsis has informed similar studies in Brassica

relatives, little is known about the underlying mechanisms controlling flowering time

variation in pennycress. Different accessions of pennycress have been reported to either

act as early flowering or late flowering, with late flowering accessions growing as

rosettes for as long as 150 days prior to flowering (Best & McIntyre, 1972). It was later

found that vernalization increased the rate of flowering in the late flowering accessions

(Best & Mc Intyre, 1976). Analyses of F2 progeny between the late and early flowering

accessions determined the early flowering phenotype (spring annuals lacking the

vernalization requirement) was determined by a single gene, with the late flowering allele

being completely dominant (McIntyre & Best, 1978).

In this report we address the questions concerning the molecular basis for the spring

flowering habit in pennycress, whether or not the spring habit has arisen only once or

multiple times in nature, and we begin to analyze the geographic distribution of

pennycress flowering time variants. These studies were aided by genomic resources that

have been developed for pennycress, including a transcriptome and draft genome (Dorn

et al., 2013; Dorn *et al.*, 2015) These resources capture the pennycress gene space and

greatly aided in the identification of alterations that lead to spring flowering. We show

that key mutations in the FLC gene are responsible for spring flowering in many natural

pennycress accessions, including one allele of FLC that confers the spring annual growth

habit that spread from Europe to North America.

Materials and Methods

1.) *Thlaspi arvense* accessions and F2 population

The spring annual *Thlaspi arvense* line MN108SA was derived from a wild Minnesota

population containing both winter and spring annual plants. Five generations of single

seed descent was performed on a spring annual plant from this collection and sequenced.

A single MN111 plant was also carried through two generations of single seed descent

and sequenced.

Additional pennycress accessions described here with the 'PI' or 'Ames' prefix were

obtained from USDA-GRIN. All pennycress accessions with the 'MN' prefix were

collected by Dr. Donald Wyse (University of Minnesota). The spring annual line

Spring32 was obtained from Dr. Win Phippen at Western Illinois University under a

Materials Transfer Agreement.

Plants were germinated on moist Berger BM2 germination mix (Berger, Inc.,

www.berger.ca), stratified at 4° C for 7 days, and grown in climate-controlled growth

chambers at the University of Minnesota (21° C, 16 hour/8 hour day/night cycles at 100

micromoles/m^2/s PAR). The MN111 plant sequenced in this analysis was vernalized at

six weeks post-germination at 4° C for 30 days in the dark. After vernalization, this plant

was returned to the growth chamber conditions described above. The spring annual

accessions were not vernalized as they flowered immediately.

2.) DNA isolation and Illumina genomic DNA sequencing

DNA was isolated from a single MN108SA and a single MN111 plant using the Omega

Mag-Bind Plant DNA kit (Omega Bio-Tek, www.omegabiotek.com) according to the

manufacturer's recommended protocol. These DNA samples were submitted to the

University of Minnesota Genomics Center for sequencing on the Illumina HiSeq 2000

platform (Illumina Inc, www.illumina.com). Sequencing libraries were prepared using

the Illumina TruSeq Nano DNA Sample Prep kit with an average library insert size of

460 base pairs. The Illumina Universal Adaptor and an Indexed Adaptor (MN111 –

Illumina Indexed Adaptor #12 – barcode = CTTGTA, MN108SA – Illumina Indexed

Adaptor #11 – barcode = GGCTAC) were used to create the sequencing libraries. Each

library was sequenced on a full lane of Illumina HiSeq 2000 (100 base pair, paired-end).

DNA from the six selected European pennycress accessions was isolated using the

Qiagen Plant DNeasy Mini kit (www.qiagen.com) following the manufacturer's

recommended protocol. Illumina sequencing libraries were created for each European

accession using the Illumina TruSeq Nano kit with the 350 bp insert protocol. These six

libraries were sequenced across 1.5 lanes on the Illumina HiSeq 2500 platform (125 base

pair, paired-end) with version 4 chemistry.

All raw sequencing datasets have been deposited in the NCBI Short Read Archive under

BioProject PRJNA237017 / SRA Accession SRP036068 (Table S1).

3.) Sequencing data quality control, read mapping

Illumina sequencing read quality and contamination were examined using FASTQC

(http://www.bioinformatics.babraham.ac.uk/projects/fastqc/). FASTQ files were filtered

and trimmed to remove low quality reads and sequencing adaptors using BBDuk (

https://sourceforge.net/projects/bbmap/) using the following parameters: ftl=10

minlen=50 qtrim=rl trimq=10 ktrim=r k=25 mink=11 hdist=1

ref=/bbmap/resources/adapters.fa tpe tbo. Sequencing reads from accessions MN111 and

MN108SA had the additional BBDuk flag of ftr=95. The resulting high quality reads

from each accession were mapped to the v1 pennycress genome using Bowtie 2 and

visualized in CLC Genomics Workbench.

4.) Analysis of FLC mutations in MN111 x MN108SA F2 population, global spring

varieties, and EMS mutant

DNA was isolated from F2 progeny grown in the conditions described above using the

Omega Bio-Tek Plant MagBind 96 kit according to the manufacturers recommended

protocol. DNA oligos were designed to amplify the 5' end of the pennycress FLC gene

for the *flc-A* allele, approximately 100 base pairs upstream of the transcriptional start site

(TaFLC_1_Forw: 5' – CCGAGGAAGAAAAAGTAGATAGAGACA – 3',

TaFLC_1_Rev: 5' – GAAGCTTAAAGGGGGAAAAAGGAA – 3', Table S2). This

amplicon was also able to identify the *flc-D* allele in MN133 and PI633414 and the *flc-α*

allele from the EMS-mutagenized population as well. Polymerase Chain Reaction (PCR)

was used to amplify this fragment, producing an approximately 450 bp amplicon. New

England Biolabs Q5 High-Fidelity PCR Kit with 2x Master Mix was used, with the

following thermal cycler conditions: 1.) 98°C for 30 seconds, 2.) 98°C for 10 seconds, 3.)

57°C for 20 seconds, 4.) 72°C for 20 seconds, 5.) Go to step #2 34 times , 6.) 72°C for 2

minutes, 7.) 4°C hold. Reactions were visualized to confirm amplification of a single

band using gel electrophoresis. PCR products were submitted to Beckman Coulter

Genomics for PCR product purification and single pass Sanger sequencing. Amplicons

were sequenced in both directions using the forward and reverse primers listed above.

Sanger sequencing reads were analyzed in CLC Genomics Workbench and aligned

against the pennycress MN106 reference genome at the FLC locus to identify sequence

variants.

5.) Identification of deletion alleles using PCR

A diagnostic PCR test was designed to test for the *flc-B* allele. A primer set

(TaFLC_2_Forw – 5' – ATAGTGTGCATCAACTGGTC – 3', and TaFLC_2_Rev – 5'–

CGAACCATAGTTCAGAGCTT – 3', Table S2) were designed to amplify a single

amplicon overlapping the 456 bp deletion (Fig. S1). In the absence of the *flc-B* allele, a

2,088 bp fragment is produced, whereas a 1,632 bp fragment is produced in plants

containing the *flc-B* allele (Fig. 2B). New England Biolabs Q5 High-Fidelity PCR Kit

with 2x Master Mix was used, with the following thermal cycler conditions: 1.) 98°C for

30 seconds, 2.) 98°C for 5 seconds, 3.) 63°C for 15 seconds, 4.) 72°C for 60 seconds, 5.)

Go to step #2 34 times, 6.) 72°C for 2 minutes, 7.) 4°C hold.

A diagnostic PCR test was developed to test for the presence or absence of the *flc-C*

allele (Fig. 2C, Fig. S2). A primer set (TaFLC_4_Forw: 5'–

GCGACGGTGAATATGGAGTTGG – 3', and TaFLC_3_Rev – 5'–

GCTAATTTTTCAGCAAATCTCCCG – 3', Table S2) was designed to amplify a 6,598

bp amplicon overlapping the 4,724 bp deletion. In plants containing the *flc-C* allele, this

region produces a 1,874 bp amplicon, confirming the presence of this deletion. New

England Biolabs Q5 High-Fidelity PCR Kit with 2x Master Mix was used, with the

following thermal cycler conditions: 1.) 98°C for 30 seconds, 2.) 98°C for 5 seconds, 3.)

67°C for 15 seconds, 4.) 72°C for 40 seconds, 5.) Go to step #2 34 times, 6.) 72°C for 2

minutes, 7.) 4°C hold. This amplicon was purified using the Qiagen QiaQuick PCR

Purification Kit (www.qiagen.com) from accessions MN135, Ames22461, and PI650285

and subjected to Sanger sequencing to confirm the exact deletion size and coordinates.

Identical 4,724 bp deletions were identified in these three accessions.

A secondary amplicon was used to confirm the absence of this deletion, with the forward

primer (TaFLC_1_Forw: 5' – CCGAGGAAGAAAAAGTAGATAGAGACA – 3', Table

S2) lying in the middle of the deleted region (Fig. S2). Successful amplification using this

secondary primer set (TaFLC_1_Forw and TaFLC_3_Rev – 5' –

GCTAATTTTTCAGCAAATCTCCCG – 3', Table S2) produces a 1,397 bp fragment

and confirms the absence of this deleted region. New England Biolabs Q5 High-Fidelity

PCR Kit with 2x Master Mix was used, with the following thermal cycler conditions: 1.)

98°C for 30 seconds, 2.) 98°C for 5 seconds, 3.) 64°C for 15 seconds, 4.) 72°C for 40

seconds, 5.) Go to step #2 34 times, 6.) 72°C for 2 minutes, 7.) 4°C hold.

Results

Flowering time phenotypes of winter and spring annual pennycress

While identifying germplasm to be used for the pennycress breeding program, wild isolates of pennycress have been collected from various locations across North America (Table S3). A spring flowering pennycress variant, named MN108SA, was isolated from a mixed population of both spring and winter annual individuals collected from Roseau, Minnesota. Single seed descent for 5 generations showed that the trait was stable and that vernalization was not required for floral induction. MN108SA was crossed to the vernalization-requiring accession MN111. A comparison of MN108SA and MN111 seedlings is shown in Fig. 1A. MN108SA shows internode elongation soon after the formation of the first true leaves (Fig. 1A insert), whereas MN111 forms a rosette. As previously reported by McIntyre and Best, 1978, the F1 derived from the cross between MN111 and MN108SA exhibited the dominant winter flowering habit, segregating in the F2 population at 38:12 winter annual:spring annual, suggesting that a single dominant gene is responsible for maintaining the winter annual phenotype.

Whole genome sequencing of the MN111 and MN108SA parent plants was used to identify key genomic differences underlying the difference in habit between winter annual and spring annual plants. DNA from each parent was sequenced on a full lane of the Illumina HiSeq 2000 platform (100 base pair, paired end). Over 334M sequencing reads were generated per parent (Table S1). A total of 29B high quality (post quality control) were generated per parent, representing >50X coverage of the predicted genome size of 539Mb [1]. Trimmed and filtered reads were mapped to the winter annual MN106 draft genome sequence [2].

Fig. 1: Segregation analysis confirms *flc-A* allele confers spring annual phenotype

(a) Co-grown MN108SA and MN111 pennycress plants. Insert shows immediate

internode elongation in the MN108SA spring annual plant. An F2 population was derived

from a cross between MN108SA x MN111 and the segregation ratio of spring to winter

annual phenotypes was determined. Whole genome sequencing of the parent plants from

each background was also conducted. (b) Gene model of pennycress FLOWERING

LOCUS C (FLC). Gray arrow below the first exon indicates the position of the amplicon

used to genotype the SNP shown in panel C. (c) Nucleotide and predicted peptide

sequence of the 5' end of the pennycress FLC gene in winter annual MN106, winter

annual MN111, and the spring annual accession MN108SA. The *flc-A* mutation in

MN108SA is highlighted in yellow.

Whole genome sequencing identifies the pennycress *flc-A* allele

Read mappings at the pennycress FRI and FLC loci for MN106, MN111, and MN108SA were examined to identify potential mutations underlying the winter to spring annual transition. Comparison of the pennycress FRI homolog (Ta1.0_26225 on scaffold 1344 of the v1.0 pennycress genome) revealed a single SNP between these three lines, however, this SNP was shared by both the MN111 and MN108SA individual, likely indicating a non-causative effect on the spring annual growth habit (Fig. S3). This SNP was found to cause an amino acid change (Threonine to Serine) at position 553 of the TaFRI predicted peptide (position 2,085 of the gene model - c.2,085A>T).

Within the MN106 draft genome, a single copy of FLC with a high degree of similarity to that of Arabidopsis was identified (Fig. 1B, Fig. S4, Fig. S5). We identified a single guanine base insertion in the 7th position of the FLC coding sequence (*flc-A*) of the spring annual MN108SA accession that was absent in both winter annual accessions. This frameshift mutation caused a predicted TGA stop codon in the 12th codon position (Fig. 1C).

To obtain genetic evidence that this *flc-A allele* was the causal variant responsible for the spring flowering habit in MN108SA, an F2 population from the cross between MN111 and MN108SA was examined. A single F1 individual was self pollinated, and 50 F2 progeny were planted for further analysis. Of these F2 individuals, 38 exhibited a winter annual phenotype and 12 exhibited a spring annual phenotype (Fig. 1A).

PCR primers were developed that amplify a 394 bp amplicon that covers the location of the *flc-A* insertion. Amplicons from all 12 spring annual F2 progeny were sequenced and confirmed homozygous for this *flc-A* allele. Amplicons from a random selection of 14 winter annual F2 progeny were also sequenced. Of the winter annual F2 progeny, 5 were homozygous for the wild type FLC gene while 9 were heterozygous. A chi square

analysis ($\chi2$=0.026667, p-value = 0.8703) indicates that the data fits a 1:2:1 segregation

pattern that is predicted for a single recessive mutation in this F2 population.

Identifying a novel EMS-induced allele of FLC in pennycress

As a component of ongoing work to domesticate pennycress, an EMS-mutagenized

MN106 population has been generated (Sedbrook, 2014). This winter flowering

population has been screened to identify early flowering variants. In the course of this

screen, a new spring flowering time mutant was identified (Fig. S6). This line did not

require vernalization and flowered immediately upon germination. Whole genome

sequencing of this mutant revealed a novel mutation in the first exon of FLC (*c.52C>T*),

resulting in an immediate stop codon and truncated predicted protein (Fig. S6), which

was designated *flc-α*. The above segregation data along with the finding of two new

independent spring flowering mutants with FLC mutations confirms the causative role of

FLC mutations in the spring flowering habit in pennycress.

Fig. 2: Identification of *flc-B* and *flc-C* alleles of FLC

(a) Expanded gene model of pennycress FLC region, including the upstream TaUFC gene model. Locations of the *flc-B* and *flc-C* alleles are shown below the gene model. (b) Agarose gel of the diagnostic polymerase chain reaction (PCR) results testing for the *flc-B* allele across the shown pennycress accessions. The wild type MN106 allele generates a 2,088 bp amplicon, where as the *flc-B* allele produces a 1,632 bp amplicon (Fig S1). (c) Agarose gel of the diagnostic polymerase chain reaction (PCR) results testing for the *flc-C* allele across the shown pennycress accessions. Amplification in the top row confirms the presence of the deletion, whereas amplification in the second row confirms absence of the deletion. The sequences of primers used are shown in Table S2, and primer locations and amplicon strategy is shown in Fig. S2. Results for accession PI650286 (Groitzsch, Saxony, Germany) are not shown, but was confirmed to have the *flc-C* allele. (d)

Nucleotide and predicted peptide sequence of FLC in MN106 and PI633414 (Wachstedt,

Germany), with the *flc-D* allele highlighted in yellow. (e) Global distribution of the four

spring annual alleles of FLC in pennycress accessions tested in this study (listed in Table

S3).

Global distribution of three additional spring annual alleles of FLC

With the genetic confirmation that the *flc-A* allele was indeed the causal variant

responsible for the spring annual growth habit in MN108SA, we next examined the

distribution of this allele in spring annual accessions from around the world. We screened

a total of 35 spring annual accessions, which were obtained either from the University of

Minnesota collections or GRIN. Of these, 23 were confirmed as pure spring annual

accessions in both field and growth chamber experiments, with no winter annual

segregation nor contaminating winter annual seed (Table S3). Surprisingly, we only

identified the *flc-A* allele in three other accessions, all from North America (MN131 from

Montana, USA and Ames31489 and Ames31491 from Canada).

As there were a limited number of representative spring annual accessions in our

collection containing the *flc-A* allele, we utilized a PCR-based approach to determine

whether there were additional mutations in FLC underlying the spring annual growth

habit. As we had already amplified and sequenced the first exon of FLC in all of our

spring annual accessions, we moved next to the second exon. Interestingly, we were

unable to amplify the second exon in several spring annual accessions. Thus, we

expanded our PCR experiments into the flanking introns around exon 2, which lead to the

discovery of a second allele of FLC in spring annual lines. This mutation was

characterized by a 456 bp deletion that encompasses the entire second exon of FLC,

which was named *flc-B*. This allele was identified in 9 spring annual accessions

distributed across the United States and Canada, along with the Spring32 accession from

the Western Illinois University breeding program (Fig. 2B and 2E, Table S3) (Sedbrook,

2014).

Having not yet identified a candidate spring annual FLC allele in any accession from

Europe, we again employed whole genome sequencing of the 5 publicly available spring

annual pennycress accessions (collected in Germany, Poland, and France). Using this

approach, we were able to quickly identify a third allele, characterized by a 4,724 bp

deletion (*flc-C*), which removes the first intron of FLC along with the upstream gene

(gene model Ta000918), a homolog of the Arabidopsis gene UPSTREAM OF FLC (UFC,

AT5G10150). In Arabidopsis, UFC and FLC are co-regulated, however, UFC is not

believed to influence the vernalization response (Finnegan *et al.*, 2004; Andersson *et al.*,

2008). A diagnostic amplicon experiment was designed to screen the remaining

unknown spring annual accessions for the *flc-C* allele. This allele was identified in four

accessions collected in Howard Springs, Montana, USA, along with three accessions

from Germany, as well as accessions from Poland and France (Fig. 2C, 2E, Supplemental

Table 3).

A fourth allele, *flc-D*, was identified through whole genome sequencing of the spring

annual accession PI633414 (Germany). This allele is characterized by a single base pair

change at the 100^{th} position of the CDS and caused a precocious TAG stop codon in the

predicted protein sequence (Fig. 2D). As the position of this mutation was covered by the

394 bp amplicon used to identify the *flc-A* allele, we re-examined the Sanger reads from

all of the accessions sequenced in our search for accessions containing the *flc-A* allele.

We did not identify any additional accessions containing the *flc-D* allele.

Discussion

Use of NGS to quickly identify mutations in FLC

At the onset of this experiment, the predicted costs associated with PCR-based cloning and sequencing of all candidate genes was assessed. This analysis indicated that it is now less expensive to use the whole genome sequencing (WGS) approach to first identify a candidate locus in the parent accessions, and then proceed to test F2 progeny via PCR and Sanger sequencing. Here we report the discovery of four natural alleles of FLC that confer the spring annual phenotype in pennycress. The added value with the WGS approach here is that we now also have genome-wide markers for the two parents of the F2 mapping population (MN111 and MN108SA), which is being used to develop a linkage map via Genotype by Sequencing (Elshire *et al.*, 2011; Poland & Rife, 2012). Additionally, as the network controlling flowering time is known to be extremely complex, the gene variants in each of these parents that may have a minor effect on vernalization and flowering provide a wealth of untapped information for later investigation.

Global distribution of FLC alleles and the introduction to North America

Pennycress is native to Europe and Asia. Thought to have been introduced to North America as early as the establishment of Fort Detroit (Michigan, USA) in 1701, pennycress became widely distributed throughout Canada and the United States by the early 1900s (Best & McIntyre, 1975). Despite the limited number (35) of pure spring annual accessions tested in this study, we can begin examining the origins of each of the spring annual alleles of FLC identified here.

As the species originated in Eurasia and only the *flc-B and flc-D* alleles were identified in the seven spring annual European accessions tested here, we hypothesize that these alleles originated prior to introduction to North America (Fig. 2E). As the accessions collected near Howard Spring, Montana, USA (MN129, MN133, MN134, MN135) contained one of these 'European' alleles of FLC (*flc-C*), this suggests that this allele

may have been introduced into North America a limited number of times, potentially

from plants near Saxony or Thuringia, Germany (Fig. 2E). However, it is possible that

the *flc-D* allele is present in populations in North America, but was not identified in the

accessions represented in this experiment.

Similarly, as we did not identify European accessions containing the *flc-A* or *flc-B* alleles,

we predict that these alleles may have originated after the introduction of pennycress to

North America and this allele is unique to North American accessions. It is possible that

there are in fact European pennycress populations that contain the *flc-A* allele, but with

the limited European germplasm availability, were not found in this study. In this

scenario, it is feasible that a mixed seed lot containing *flc-A* and *flc-B* alleles were

introduced to Montana in a single event, and plants with the *flc-A* and *flc-B* alleles spread

significantly more compared to those with the *flc-D* allele.

The accessions collected near Howard Spring, Montana (MN129, MN131, MN133,

MN134, MN135) are an interesting case study of these hypotheses, especially

considering the findings of the *flc-A* allele in MN131, the *flc-B* allele in MN129, MN133,

MN134, and MN135, and a lack of identification of European accessions with the *flc-A*

allele. The above scenarios could be investigated further in these accessions collected

from Montana using genome-wide markers from either genotype-by-sequencing or WGS.

As we have already generated WGS data for five of the European accessions in this study,

sequencing of these accessions from Montana and a comparison of shared

polymorphisms may help in determining the relatedness between the North American and

European lines, and may point to the exact origins of these alleles of FLC in pennycress.

Again, with the limited number of pure spring annual accessions available, future studies

should also focus on the collection of a more comprehensive set of globally distributed

lines, including both spring and winter annuals.

Towards an understanding of flowering time control in pennycress

Our previous efforts to develop genomic resources for the domestication of pennycress identified likely homologs controlling flowering time and the vernalization response via DNA and RNA sequence homology (Dorn *et al.*, 2013; Dorn *et al.*, 2015). Here we present the first sequence-supported genetic information on the underlying mechanisms controlling flowering time in pennycress. While the four natural alleles of FLC reported here are sufficient to confer the spring annual phenotype, there is likely a host of interacting genes and epigenetic factors also essential for the rapid flowering seen in the spring annual accessions investigated here. Of particular interest is FRIGIDA and members of the 'FRIGIDA Complex' (FRI-C), including FRIGIDA LIKE 1 (FRL1), FRIGIDA-ESSENTIAL 1 (FES1), SUPPRESSOR OF FRIGIDA4 (SUF4), and FLC EXPRESSOR (FLX) (Choi *et al.*, 2011). The FRIGIDA Complex acts as a transcriptional activation complex on FLC expression through a diverse range of functions of each complex member (Choi *et al.*, 2011). While no loss of function FRIGIDA alleles were identified in the accessions examined here, it remains possible that an FLC-independent path to the spring annual phenotype exists in light of the limited availability of globally distributed germplasm. The interaction between the vernalization response, photoperiodic, and autonomous pathways are also of great interest, as allelic variation and unique combinations of alleles from each pathway can contribute to quantitative variation in flowering time and lead towards the goal of developing an elite winter annual pennycress variety that flowers rapidly in the spring to fit within the short growing season of northern climates.

Acknowledgements

The authors would like to thank Ryan Emenecker, Nicole Folstad, Matthew Ott, and

Ratan Chopra for their assistance with experiments and for feedback on the manuscript.

This material is based upon work supported by the National Science Foundation Graduate

Research Fellowship under Grant No. 00006595 to KMD. Any opinions, findings, and

conclusions or recommendations expressed in this material are those of the authors and

do not necessarily reflect the views of the National Science Foundation. This project is

funded by the USDA National Institute of Food Agriculture - Institute of Bioenergy,

Climate and Environment, competitive grant no. 2014-67009-22305.

Author Contributions

KMD and MDM conceived and planned the study. EBJ, ED, and MDM constructed the

F2 mapping population. EBJ and ED identified the FLC-α allele. KMD designed and

performed PCR experiments, whole genome sequencing and analyses, and comparative

analyses. DLW collected all 'MN' designated pennycress accessions used in this study.

KMD and MDM wrote the manuscript with feedback from all the co-authors.

References

Amasino RM. 2005. Vernalization and flowering time. *Curr Opin Biotechnol* **16**(2): 154-158.

Andersson CR, Helliwell CA, Bagnall DJ, Hughes TP, Finnegan EJ, Peacock WJ, Dennis ES. 2008. The FLX gene of Arabidopsis is required for FRI-dependent activation of FLC expression. *Plant and cell physiology* **49**(2): 191-200.

Beilstein MA, Al-Shehbaz IA, Mathews S, Kellogg EA. 2008. Brassicaceae phylogeny inferred from phytochrome A and ndhF sequence data: tribes and trichomes revisited. *Am J Bot* **95**(10): 1307-1327.

Best K, Mc Intyre G. 1976. Studies on the flowering of Thlaspi arvense L. III. The influence of vernalization under natural and controlled conditions. *Botanical Gazette*: 121-127.

Best K, McIntyre G. 1972. Studies on the Flowering Time of Thlaspi arvense L. I. The Influence of some Environmental and Genetic Factors. *Botanical Gazette* **133**(4): 454 - 459.

Best K, McIntyre G. 1975. The Biology of Canadian Weeds 9. Thlaspi arvense L. . *Canadian Journal of Plant Science* **55**: 279-292.

Burn J, Smyth D, Peacock W, Dennis E. 1993. Genes conferring late flowering inArabidopsis thaliana. *Genetica* **90**(2-3): 147-155.

Choi K, Kim J, Hwang HJ, Kim S, Park C, Kim SY, Lee I. 2011. The FRIGIDA complex activates transcription of FLC, a strong flowering repressor in Arabidopsis, by recruiting chromatin modification factors. *Plant Cell* **23**(1): 289-303.

Clarke JH, Dean C. 1994. Mapping FRI, a locus controlling flowering time and vernalization response in Arabidopsis thaliana. *Mol Gen Genet* **242**(1): 81-89.

Coustham V, Li P, Strange A, Lister C, Song J, Dean C. 2012. Quantitative modulation of polycomb silencing underlies natural variation in vernalization. *Science* **337**(6094): 584-587.

Crowley JG. 1999. *Evaluation of Camelina sativa as an alternative oilseed crop*: Teagasc.

Csorba T, Questa JI, Sun Q, Dean C. 2014. Antisense COOLAIR mediates the coordinated switching of chromatin states at FLC during vernalization. *Proceedings of the National Academy of Sciences* **111**(45): 16160-16165.

Dorn KM, Fankhauser JD, Wyse DL, Marks MD. 2013. De novo assembly of the pennycress (Thlaspi arvense) transcriptome provides tools for the development of a winter cover crop and biodiesel feedstock. *Plant J* **75**(6): 1028-1038.

Dorn KM, Fankhauser JD, Wyse DL, Marks MD. 2015. A draft genome of field pennycress (Thlaspi arvense) provides tools for the domestication of a new winter biofuel crop. *DNA Res* **22**(2): 121-131.

Elshire RJ, Glaubitz JC, Sun Q, Poland JA, Kawamoto K, Buckler ES, Mitchell SE. 2011. A robust, simple genotyping-by-sequencing (GBS) approach for high diversity species. *PLoS One* **6**(5): e19379.

Evangelista RL, Isbell TA, Cermak SC. 2012. Extraction of pennycress (Thlaspi
arvense L.) seed oil by full pressing. *Industrial Crops and Products* **37**(1): 76-81.

Fan J, Shonnard DR, Kalnes TN, Johnsen PB, Rao S. 2013. A life cycle assessment of
pennycress (Thlaspi arvense L.) -derived jet fuel and diesel. *Biomass and*
Bioenergy **55**: 87-100.

Finnegan EJ, Dennis ES. 2007. Vernalization-induced trimethylation of histone H3
lysine 27 at FLC is not maintained in mitotically quiescent cells. *Current Biology*
17(22): 1978-1983.

Finnegan EJ, Sheldon CC, Jardinaud F, Peacock WJ, Dennis ES. 2004. A cluster of
Arabidopsis genes with a coordinate response to an environmental stimulus.
Current Biology **14**(10): 911-916.

Franzke A, Lysak MA, Al-Shehbaz IA, Koch MA, Mummenhoff K. 2011. Cabbage
family affairs: the evolutionary history of Brassicaceae. *Trends Plant Sci* **16**(2):
108-116.

Greb T, Mylne JS, Crevillen P, Geraldo N, An H, Gendall AR, Dean C. 2007. The
PHD finger protein VRN5 functions in the epigenetic silencing of Arabidopsis
FLC. *Current Biology* **17**(1): 73-78.

Hojilla-Evangelista MP, Evangelista RL, Isbell TA, Selling GW. 2013. Effects of
cold-pressing and seed cooking on functional properties of protein in pennycress
(Thlaspi arvense L.) seed and press cakes. *Industrial Crops and Products* **45**: 223-
229.

Jiang D, Gu X, He Y. 2009. Establishment of the winter-annual growth habit via
FRIGIDA-mediated histone methylation at FLOWERING LOCUS C in
Arabidopsis. *Plant Cell* **21**(6): 1733-1746.

Johanson U, West J, Lister C, Michaels S, Amasino R, Dean C. 2000. Molecular
analysis of FRIGIDA, a major determinant of natural variation in Arabidopsis
flowering time. *Science* **290**(5490): 344-347.

Jung C, Muller AE. 2009. Flowering time control and applications in plant breeding.
Trends Plant Sci **14**(10): 563-573.

Kim DH, Doyle MR, Sung S, Amasino RM. 2009. Vernalization: winter and the timing
of flowering in plants. *Annu Rev Cell Dev Biol* **25**: 277-299.

Marquardt S, Raitskin O, Wu Z, Liu F, Sun Q, Dean C. 2014. Functional
consequences of splicing of the antisense transcript COOLAIR on FLC
transcription. *Mol Cell* **54**(1): 156-165.

McIntyre G, Best K. 1978 Studies on the Flowering of Thlaspi arvense IV. Genetic and
Ecological Differences between Early and Late Flowering Strains *Botanical*
Gazette **139**(2): 190-195.

Michaels SD, Amasino RM. 1999. FLOWERING LOCUS C encodes a novel MADS
domain protein that acts as a repressor of flowering. *Plant Cell* **11**(5): 949-956.

Michaels SD, Amasino RM. 2001. Loss of FLOWERING LOCUS C activity eliminates
the late-flowering phenotype of FRIGIDA and autonomous pathway mutations
but not responsiveness to vernalization. *Plant Cell* **13**(4): 935-941.

Michaels SD, Bezerra IC, Amasino RM. 2004. FRIGIDA-related genes are required for
the winter-annual habit in Arabidopsis. *Proc Natl Acad Sci U S A* **101**(9): 3281-
3285.
Michaels SD, He Y, Scortecci KC, Amasino RM. 2003. Attenuation of FLOWERING
LOCUS C activity as a mechanism for the evolution of summer-annual flowering
behavior in Arabidopsis. *Proc Natl Acad Sci U S A* **100**(17): 10102-10107.
Moser BR, Knothe G, Vaughn SF, Isbell TA. 2009. Production and Evaluation of
Biodiesel from Field Pennycress (Thlaspi arvense L.) Oil. *Energy & Fuels* **23**(8):
4149-4155.
Phippen WB, Phippen ME. 2012. Soybean Seed Yield and Quality as a Response to
Field Pennycress Residue. *Crop Science* **52**(6): 2767.
Poland JA, Rife TW. 2012. Genotyping-by-Sequencing for Plant Breeding and Genetics.
Plant Genome **5**(3): 92-102.
Searle I, He Y, Turck F, Vincent C, Fornara F, Kröber S, Amasino RA, Coupland G.
2006. The transcription factor FLC confers a flowering response to vernalization
by repressing meristem competence and systemic signaling in Arabidopsis. *Genes*
& Development **20**(7): 898-912.
Sedbrook JP, WB; Marks, MD 2014. New approaches to facilitate rapid domestication
of a wild plant to an oilseed crop: example pennycress (Thlaspi arvense L.). *Plant*
Sci **227**: 122-132.
Selling G, Evangelista R, Isbell T, Evangelista MH. 2011. Extraction of proteins from
pennycress. *Abstracts of Papers of the American Chemical Society* **241**.
Shindo C, Aranzana MJ, Lister C, Baxter C, Nicholls C, Nordborg M, Dean C. 2005.
Role of FRIGIDA and FLOWERING LOCUS C in determining variation in
flowering time of Arabidopsis. *Plant Physiol* **138**(2): 1163-1173.
Simpson GG, Dean C. 2002. Arabidopsis, the Rosetta stone of flowering time? *Science*
296(5566): 285-289.
Stinchcombe JR, Weinig C, Ungerer M, Olsen KM, Mays C, Halldorsdottir SS,
Purugganan MD, Schmitt J. 2004. A latitudinal cline in flowering time in
Arabidopsis thaliana modulated by the flowering time gene FRIGIDA. *Proc Natl*
Acad Sci U S A **101**(13): 4712-4717.
Sung S, He Y, Eshoo TW, Tamada Y, Johnson L, Nakahigashi K, Goto K, Jacobsen
SE, Amasino RM. 2006. Epigenetic maintenance of the vernalized state in
Arabidopsis thaliana requires LIKE HETEROCHROMATIN PROTEIN 1. *Nat*
Genet **38**(6): 706-710.
Swiezewski S, Liu F, Magusin A, Dean C. 2009. Cold-induced silencing by long
antisense transcripts of an Arabidopsis Polycomb target. *Nature* **462**(7274): 799.
Tadege M, Sheldon CC, Helliwell CA, Stoutjesdijk P, Dennis ES, Peacock WJ. 2001.
Control of flowering time by FLC orthologues in Brassica napus. *The Plant*
Journal **28**(5): 545-553.
Warwick SI, Francis, A., Susko, D.J. . 2002 The biology of Canadian weeds. 9. Thlaspi
arvense L. (updated) *Canadian Journal of Plant Science* **82**: 803-823.

Wu J, Wei K, Cheng F, Li S, Wang Q, Zhao J, Bonnema G, Wang X. 2012. A
naturally occurring InDel variation in BraA. FLC. b (BrFLC2) associated with
flowering time variation in Brassica rapa. *BMC Plant Biol* **12**(1): 151.
Yang H, Howard M, Dean C. 2014. Antagonistic roles for H3K36me3 and H3K27me3
in the cold-induced epigenetic switch at Arabidopsis FLC. *Current Biology*
24(15): 1793-1797.
679

Supporting Information Brief Legends

Fig. S1 - PCR strategy for identifying *flc-B* allele

Fig. S2 - PCR strategy for identifying *flc-C* allele

Fig. S3 - Peptide alignment of Brassicaceae FRIGIDA predicted peptides and pennycress

FRI gene model

Fig. S4 - Peptide alignment of Brassicaceae FLOWERING LOCUS C predicted peptides

Fig. S5 - Genomic DNA alignment of pennycress FLOWER LOCUS C wild type and *flc-*

a allele

Fig. S6 - Identification and characterization of pennycress flc-α

Table S1 - Summary of whole genome sequencing reads of pennycress accessions used

in this study

Table S2 - Primer sequences used in PCR experiments used in this study

Table S3 - Summary of pennycress accessions and the corresponding alleles of FLC as

identified in this study

(a)

MN108
(Spring Annual)   x   MN111
(Winter Annual)

Whole
Genome
Sequencing

F1
(Winter Annual)

Self pollinated

Whole
Genome
Sequencing

<u>50 F2 individuals</u>
38 Winter Annuals
12 Spring Annuals

(c)

MN106 (WA) ATGGGGAGAAAAAAACTAGAAATCAAGCGAATTGAG
            M G R K K L E I K R I E

MN111 (WA) ATGGGGAGAAAAAAACTAGAAATCAAGCGAATTGAG
            M G R K K L E I K R I E

MN108 (SA) ATGGGGGAGAAAAAAACTAGAAATCAAGCGAATTGA
            M G E K K T R N Q A N *

flc-A

(b)

TaFLC

1,000   2,000   3,000   4,000   5,000