Patterns of thaumarchaeal gene expression in culture and diverse marine environments

Authors:
Paul Carini[1,3], Christopher L. Dupont[2], Alyson E. Santoro[1,4]

[1]Horn Point Laboratory, University of Maryland Center for Environmental Science, Cambridge, MD 21613
[2]J. Craig Venter Institute, San Diego, CA 92037
[3]Present Address: Department of Soil, Water and Environmental Science, University of Arizona, Tucson, AZ 85721
[4]Present address: Department of Ecology, Evolution, and Marine Biology, University of California, Santa Barbara, CA 93106

Keywords: Thaumarchaea, nitrification, metatranscriptome, NirK, nitrogen cycling

Corresponding Author: Alyson Santoro (alyson.santoro@lifesci.ucsb.edu)

Abstract

Thaumarchaea are ubiquitous in marine habitats where they participate in carbon and nitrogen cycling. Although metatranscriptomes suggest thaumarchaea are active microbes in marine waters, we understand little about how thaumarchaeal gene expression patterns relate to substrate utilization and activity. Here, we report the global transcriptional response of the marine ammonia-oxidizing thaumarchaeon 'Candidatus Nitrosopelagicus brevis' str. CN25 to ammonia limitation using RNA-Seq. We further describe the genome and transcriptome of Ca. N. brevis str. U25, a new strain capable of urea utilization. Ammonia limitation in CN25 resulted in reduced expression of transcripts coding for ammonia oxidation proteins, and increased expression of a gene coding an Hsp20-like chaperone. Despite significantly different transcript abundances across treatments, two ammonia monooxygenase subunits (*amoAB*), a nitrite reductase (*nirK*), and both ammonium transporter genes were always among the most abundant transcripts, regardless of growth state. Ca. N. brevis str. U25 cells expressed a urea transporter 139-fold more than the urease catalytic subunit *ureC*. Gene co-expression networks derived from culture transcriptomes and ten thaumarchaea-enriched metatranscriptomes

revealed a high degree of correlated gene expression across disparate environmental conditions and identified a module of genes, including *amoABC* and *nirK*, that we hypothesize to represent the core ammonia oxidation machinery.

Originality-Significance Statement:
Discovering gene function in fastidious or uncultivated lineages remains one of the biggest challenges in environmental microbiology. Here, we use an approach that combines controlled laboratory experiments with *in situ* transcript abundance data from the environment to identify genes that share similar transcription patterns in marine ammonia-oxidizing thaumarchaea. These findings demonstrate how transcriptomes from microbial cultures can be used together with complex environmental samples to identify suites of co-expressed genes that are otherwise enigmatic and provide new insights into the mechanism of ammonia oxidation. Our results add to the growing body of literature showing that relatively small changes in transcript abundance are linked to large changes in growth in organisms with reduced genomes, suggesting they have limited capacity for metabolic regulation or that they rely on mechanisms other than transcriptional regulation to deal with a fluctuating environment.

Introduction:
Ammonia-oxidizing thaumarchaea are ubiquitous and abundant in the oceans, accounting for >30% of all cells below the thermocline (Karner *et al.*, 2001; Schattenhofer *et al.*, 2009) and are integral organisms in oxygen minimum zones (Francis *et al.*, 2005; Coolen *et al.*, 2007; Lam *et al.*, 2009; Pitcher *et al.*, 2011; Stewart *et al.*, 2012; Beman *et al.*, 2012). In many marine environments, thaumarchaeal transcripts are among the most abundant that can be mapped to available prokaryotic genomes (Hollibaugh *et al.*, 2011; Baker *et al.*, 2012; Stewart *et al.*, 2012; Gifford *et al.*, 2013). In these environments, the most frequently detected thaumarchaeal transcripts encode for proteins involved in ammonia oxidation and acquisition, including ammonia monooxygenase subunits (*amoABC*), ammonium transporters (*amtB*), a putative Cu-containing nitrite reductase (*nirK*), and structural cellular components (for example, S-layer proteins (Nakagawa and Stahl, 2013)). In addition to dissolved ammonia, some ammonia-oxidizing archaea utilize ammonia derived from urease-

catalyzed urea hydrolysis as a chemolithoautotrophic growth substrate (Qin
et al., 2014; Bayer *et al.*, 2015) and urease genes and transcripts believed
to be of thaumarchaeal origin have been detected in marine environments
(Shi *et al.*, 2010; Alonso-Sáez *et al.*, 2012; Pedneault *et al.*, 2014; Tolar,
Ross, *et al.*, 2016). Despite the abundance of thaumarchaeal transcripts in
natural assemblages, we still have a poor understanding of how the relative
abundance of thaumarchaeal transcript markers such as *amoA*, *nirK* and
ureC relate to nutrient and energy availability across environmental
gradients.
86
Thaumarchaeal ammonia oxidation is initiated by the oxidation of ammonia
to hydroxylamine (NH_2OH) by the ammonia monooxygenase enzyme
complex (Amo) (Vajrala *et al.*, 2013), but the enzyme(s) catalyzing the
oxidation of NH_2OH to nitrite (NO_2^-) have not been confirmed (Walker *et al.*,
2010). Orthologs of the bacterial hydroxylamine oxidoreductase (Hao) or c-
type cytochrome synthesis and assembly machinery, thought to be required
for NH_2OH oxidation and electron transfer in ammonia-oxidizing bacteria
(AOB) (Arp *et al.*, 2007), are absent from all sequenced thaumarchaeal
genomes (Stahl and la Torre, 2012; Spang *et al.*, 2012; Kerou *et al.*, 2016).
Instead, unidentified Cu-containing metalloenzymes or F_{420}-dependent
monooxygenases are speculated to be involved in NH_2OH oxidation and
electron transfer to archaeal terminal oxidases (Walker *et al.*, 2010; Kerou
et al., 2016) and may involve nitric oxide (NO) as either a direct
intermediate or an electron shuttle (Stieglmeier *et al.*, 2014; Martens-
Habbena *et al.*, 2015; Kozlowski *et al.*, 2016). While the precursor to NO
has not yet been elucidated, all free-living thaumarchaea with complete
genomes encode a Cu-containing multicopper oxidase with homology to
Cu-dependent nitrite reductases (NirK) that may be responsible for the
reduction of NO_2^- to NO (Kerou *et al.*, 2016).
106
'Candidatus Nitrosopelagicus brevis' str. CN25 is a cultured representative
of ubiquitous and abundant pelagic thaumarchaeal populations in the
shallow oligotrophic ocean (Santoro and Casciotti, 2011; Santoro *et al.*,
2015). Here, we describe the genome and transcriptome during urea-based
growth of a *Ca*. N. brevis strain that can utilize ammonia cleaved from urea
as a sole chemolithoautotrophic growth substrate. Additionally, we use *Ca*.
N. brevis str. CN25 to investigate the transcriptional response to ammonia
limitation in laboratory culture. These transcriptomes are further analyzed in

the context of several marine metatranscriptomes and used to identify
conserved gene co-expression networks.
117
Results and Discussion
'*Candidatus* Nitrosopelagicus brevis' strain U25 genome and
transcriptome: A urea-utilizing thaumarchaeon was obtained from an
ammonia-oxidizing enrichment culture (Santoro and Casciotti, 2011) via
subculturing with urea as a sole nitrogen and energy source (see
'Experimental Procedures') and used for the shotgun metagenome
sequencing and experiments described here. After assembly and contig
binning based on nucleotide frequencies and coverage, we obtained a
three contig genome of this urea-utilizing thaumarchaeon (Supplementary
Figure 1a). This genome is nearly identical to the *Ca*. N. brevis CN25
genome with regards to *i)* gene content; *ii)* genome organization
(Supplementary Figure 1b); and *iii)* genome wide average nucleotide
identity (99.99%; Supplementary Figure 2). We found 19 additional genes
at four distinct genomic loci in this strain, relative to CN25 (Supplementary
Table 1). The largest of these insertions (15 contiguous genes) includes 11
genes coding urea utilization machinery, including *ureABCDEFG*, which
codes for urease and its chaperones, two urea sodium:solute symporter
family (SSSF) transporters, a transcriptional regulator and several
hypothetical proteins (Fig. 1a). We designate this urea-utilizing
thaumarchaeon '*Candidatus* Nitrosopelagicus brevis' strain U25.
138
We sequenced a transcriptome from *Ca*. N. brevis str. U25 growing
exponentially with urea as the growth substrate. Only one transcript from
the chromosomal insertion containing the urea transport and metabolism
genes (Fig. 1a) was among the top 50 transcripts detected: A7X95_00990,
coding for a putative urea SSSF transporter (ranked 13.7 ± 0.33; mean ±
SE, $n = 3$). Surprisingly, transcripts coding for catalytic urease components,
or the second putative urea SSSF transporter (A7X95_00985) located
immediately adjacent to A7X95_00990, were not nearly as abundant as
A7X95_00990. For example, the mean expression levels of *ureC*, coding
for the fused catalytic αβ-subunit of urease, and *ureA* (γ urease subunit)
were 139 and 784-fold less abundant than that of A7X95_00990,
respectively (ranked 390 ± 19.7 and 950 ± 50.8, respectively; mean ± SD,
n=3). A7X95_00985, coding for the second urea SSSF transporter, was
also expressed at a low level, comparable to *ureA*, ranked 940 ± 92.0.

During growth on urea, transcripts for genes coding for an ammonium transporter (AMT1), ammonia monooxygenase subunits (*amoAB*) and nitrite reductase (*nirK*) were within the top ten most abundant transcripts detected in strain U25.

Although urea utilization genes have been detected in wild thaumarchaeal populations, we have a poor understanding of how the abundances of urease transcripts relate to growth and activity. To contextualize the expression patterns observed in U25, we compared the relative rank of the transcript abundances for only those genes coding for urea uptake and catabolism (Fig. 1a) under laboratory growth conditions to the relative rank of the transcript abundances of the same genes from several deeply sequenced marine metatranscriptomes. The SSSF urea transporter (A7X95_00990) was the most abundant urea-related gene transcript in 38% of the environmental datasets ($n = 8$) we investigated (Fig. 1b). In contrast to culture conditions, where the SSSF urea transporter was the most abundant urea-related gene transcript, *ureC* was the most abundant transcript in 25% of the environmental datasets (Fig. 1b). This shows that variability in the relative transcriptional activity of urea transport and catabolism genes is not unusual. Our finding that *ureC* was not highly expressed in exponentially growing cells also helps to explain previous field observations of low *ureC* expression, and suggests the abundance of *ureC* transcripts may be a poor molecular biomarker of active urea-based nitrification. For example, *ureC* expression and urea-based nitrification were found to be only weakly correlated across several environments (Tolar, Wallsgrove, *et al.*, 2016), in contrast to high correlation between *amoA* expression and ammonia oxidation rates (J. M. Smith *et al.*, 2014). Similarly, in Arctic samples collected across seasons, *ureC* genes were detected, yet *ureC* transcripts were only sporadically detected and at low abundances (Pedneault *et al.*, 2014).

Transcriptional response to ammonia limitation in '*Ca.* N. brevis' strain CN25*:* To understand adaptive mechanisms during ammonia starvation, we explored the transcriptional response of *Ca.* N. brevis CN25 to ammonia limitation. A total of 51 gene transcripts were differentially abundant when comparing the exponential growth phase of CN25 to ammonia-limited stationary phase (generalized linear model likelihood ratio test FDR ≤ 0.01 and ≥ 2-fold change in abundance; Supplementary Table

2). The gene transcripts that were significantly less abundant in stationary
phase included *amoA, amoB, nirK*, both *amtB*-like ammonium transporters
(AMT1=T478_1378; AMT2=T478_1350), several additional Cu-containing
metalloenzymes, and two ferredoxin-like 4Fe-4S binding domain proteins
(Fd1=T478_1472 and Fd2=T478_1259) (Fig. 2, Supplementary Table 2).
The downregulation of *amoA, amoB* and *nirK* in ammonia-limiting
conditions was recently shown for the ammonia-oxidizing thaumarchaeon
Nitrosopumilus maritimus using DNA microarrays (Qin *et al.*, 2017),
suggesting that one adaptation of ammonia-oxidizing archaea to ammonia
limitation is to reduce the relative expression of energy generation
machinery.
202
Only two genes, T478_1481, coding an Hsp20/α-crystallin domain small
heat shock protein, and T478_1394, annotated as a hypothetical protein,
were significantly more abundant (~10-fold) in ammonia-limited stationary
phase (Fig. 2, Supplementary Table 2). Hsp20 is a molecular chaperone
that enhances thermotolerance and binds to unfolded proteins to prevent
aggregation (Li *et al.*, 2011). The higher proportion of Hsp20 transcripts
and concomitant decrease in proportion of transcripts coding enzymes
integral to energy production suggests that one adaptation *Ca.* N. brevis
employs in ammonia-limited stationary phase may be to protect existing
proteins from degradation. Nutrient stress has been shown to induce the
expression of molecular chaperone proteins in ammonia-oxidizing archaea,
ammonia-oxidizing bacteria, and oligotrophic marine heterotrophs. For
example, in *N. maritimus*, two copies of Hsp20 were differentially
expressed during copper stress and recovery, but not during ammonia
starvation (Qin *et al.*, 2017). Similar to our findings regarding Hsp20,
Nitrosomonas europaea expressed peptides for the molecular chaperone
GroEL in both energy starved and energy replete conditions, but at
significantly greater levels under energy starvation (Pellitteri-Hahn *et al.*,
2011). The authors speculated that energy stress may induce chaperone
expression as part of a generalized stress response, and that these
chaperones are involved in protein protection (Pellitteri-Hahn *et al.*, 2011).
Similarly, the marine chemoorganoheterotroph 'Candidatus Pelagibacter
ubique' induced GroEL protein expression under N-starvation (D. P. Smith
et al., 2013), GroES under iron starvation (D. P. Smith *et al.*, 2010), and the
heat shock protein IpbA in nutrient-limited stationary phase (D. P. Smith *et*
al., 2016). These finding suggest that one role of molecular chaperones

during nutrient stress may be to protect key enzymes from proteolytic
turnover when cells scavenge peptides to support nutrient-limited
sustenance.
232
In contrast to the finding that only two genes were more abundant in
ammonia-limited stationary phase for *Ca*. N. brevis, over 200 genes were
upregulated in ammonia-limited stationary phase for *N. maritimus* (Qin *et*
al., 2017). One explanation for this observation is that *Ca*. N. brevis has a
reduced capacity to sense and respond to environmental change as a
result of its small genome (Santoro *et al.*, 2015). Alternatively, our
conservative statistical thresholding may have resulted in a reduced
number of genes defined as differentially expressed. While it is important to
note that our experimental design cannot distinguish between responses to
energy limitation versus anabolic nitrogen limitation, transcriptional
responses to ammonium limitation that involved the upregulation of only a
few genes has been described for both ammonia-oxidizing bacteria and
oligotrophic marine bacteria. For example, in the bacterium *N. europaea*,
only 0.42% of the genome was upregulated under N-starvation (Wei *et al.*,
2006). Similarly, '*Ca*. P. ubique,' exhibited a weak transcriptional response
to N-starvation (D. P. Smith *et al.*, 2013). The similar lack of transcriptional
responses to nitrogen starvation by *Ca*. N. brevis and *Ca*. P. ubique are
consistent with observations that organisms with small and potentially
streamlined genomes have a limited capacity to respond rapidly to
environmental change (Giovannoni *et al.*, 2014; Cottrell and Kirchman,
2016; Giovannoni, 2017; Satinsky *et al.*, 2017).
254
Despite significantly different transcript abundances of essential ammonia
oxidation and transport genes across growth conditions, the rank order of
these transcripts within a given treatment were similar, irrespective of
growth condition. In particular, the most abundant gene transcripts in
exponential phase were generally still the most abundant transcripts in
ammonium-limited stationary phase (Fig. 2b). For example, transcripts for
32 genes (64%) were in the top 50 most abundant transcripts in both
exponential and stationary phase (Fig. 2b). Interestingly, the abundances of
18 of these transcripts were also significantly different across treatments
(Fig. 2b), illustrating that although transcripts can be differentially abundant
across paired treatments, the changes in their relative cellular abundance
may be subtler. Several of these consistently abundant but differentially

expressed transcripts are common molecular markers predicted to be essential for ammonia oxidation and transport, including the ammonium transporters AMT1 and AMT2, ammonia monooxygenase subunits (*amoAB*) and nitrite reductase (*nirK*). This suggests that the proportional changes we observed in highly-expressed genes may be the result of abundance changes in other genes that make up a smaller proportion of the transcriptome, such as Hsp20, and that even though we observe significant differences across treatments, the underlying transcript abundances might be similar.

The ammonia monooxygenase C subunit (AmoC) has been implicated in stress response (Berube and Stahl, 2012) and recovery from ammonia starvation in *N. europaea* (Berube *et al.*, 2007). The participation of AmoC in ammonium starvation appears to be conserved in thaumarchaeal ammonia oxidizers, where *amoC* transcript levels remained high in ammonia-limited stationary phase (Qin *et al.*, 2017). Consistent with these findings, *amoC* was abundant in both exponential and N-limited stationary phase in CN25 (the 13th and 7th most abundant transcript, respectively), and we did not observe a significant difference in the abundance of *amoC* across growth phases (Fig. 2).

Correlated gene expression across disparate environments: Controlled laboratory experiments such as those described above help us to understand gene regulation by isolating one experimental variable at a time. However, gene expression patterns observed in natural thaumarchaeal populations are the result of cells responding to complex and dynamic environmental conditions that can be cryptic and difficult to mimic in the laboratory. To identify clusters of co-expressed genes across disparate environmental conditions, and relate them to our laboratory findings, we constructed and analyzed a gene expression correlation network constructed from transcriptomes of exponentially growing CN25 and U25 cultures and marine metatranscriptomes. Although 64 metatranscriptomes were mapped to the *Ca.* N. brevis genomes, only ten met our strict criteria for inclusion in the network analysis presented here (see 'Experimental Procedures', below). The transcription of 1,407 of the 1,464 non-redundant genes in the two *Ca.* N. brevis genomes was significantly positively correlated with at least one other gene (Pearson's r ≥ 0.80, *q* ≤ 0.025; Fig. 3, Supplementary Table 3). Network modularity is a

measure of the group connectivity within a network, where connections
contained within a module are denser than connections between modules.
Modularity values range from -0.5 to 1, where 1 describes a highly modular
system. The modularity of this positive correlation network was 0.71,
indicating a high degree of modularity. Genes with positively correlated
expression organized into 38 groups (modules), ranging in membership
size from 2 to 236 genes (mean module size = 38.0 genes).
312
Genes encoding putative components of the core ammonia oxidation and
transport machinery are significantly co-expressed across distinct
environmental and laboratory conditions. A single 15-gene module (module
11 in Fig. 3) contained: *amoABCX*, AMT1, *nirK*, two PEFG-CTERM domain
proteins, Fd1 and Fd2, an Fe-S cluster assembly protein, a membrane
bound cupredoxin-containing protein, and three hypothetical proteins.
Further investigation of these hypothetical proteins suggests that
(T478_0057) is a putative archaeal cell division protein related to the
endosomal sorting complexes involved in membrane trafficking (ESCRT)-III
(Lindås *et al.*, 2008; Spang *et al.*, 2015). Our finding that *amoA, amoB* and
nirK transcripts are abundant and co-expressed with other genes coding for
membrane-bound Cu-containing metalloproteins (T478_1362 and
T478_0895) implies the products of these genes may participate in
ammonia oxidation. Previous speculation implicated membrane-bound
multicopper oxidases (Walker *et al.*, 2010; Stahl and la Torre, 2012;
Kozlowski *et al.*, 2016) or novel F_{420}-dependent monooxygenases (Kerou *et*
al., 2016) in ammonia oxidation chemolithotrophy (specifically NH_2OH
oxidation) based on Cu redox chemistry or ortholog conservation across
thaumarchaeal genomes. However, the genes put forth in those studies
were not present in module 11 (referred to as the AMO module, herein),
suggesting they may not be involved in core energy metabolism (Fig. 3,
Supplementary Table 3). For example, a multicopper oxidase present in the
genomes of both *N. brevis* (T478_0261) and *N. maritimus* (Nmar_1663)
previously implicated to be involved in the oxidation of hydroxylamine to
nitrite (Walker *et al.*, 2010; Qin *et al.*, 2017) was not present in the AMO
module (Supplementary Table 3).
339
On average, the AMO module is expressed at a higher level than other
modules, and was more abundant in conditions where ammonia oxidation
rates and thaumarchaeal abundances would be predicted to be high (Fig.

4). For example, consistent with previous reports of higher ammonia
oxidation rates within hydrothermal plumes (Lam *et al.*, 2004), we show
that the AMO module is expressed highly within the Guyamas Deep
hydrothermal plume, relative to background samples (Fig. 4). Moreover,
similar to reports of increased thaumarchaeal gene expression in the
mesopelagic (Church *et al.*, 2010), the AMO module is less abundant in the
surface waters of Landsort Deep (0 and 5 m), relative to deeper waters (90
and 200 m) (Fig. 4).
351
A new proposed model for thaumarchaeal chemolithotrophy via
ammonia oxidation: A previous model of thaumarchaeal ammonia
oxidation proposed NO is derived from the reduction of NO_2^- by NirK, and
that this NO is subsequently used to oxidize NH_2OH (Kozlowski *et al.*,
2016). However, this model does not agree with tracer experiments using
^{18}O labeled water, which show that only one O atom from water is
incorporated into NO_2^- produced by thaumarchaea (Santoro *et al.*, 2011;
Buchwald *et al.*, 2012). If NO_2^- was reduced to NO and used to produce
additional NO_2^-, the resulting NO_2^- would retain an average of more than
one O atom from water. Based on the co-expression data presented here,
we propose two alternative models of thaumarchaeal ammonia oxidation
that are consistent with previous isotope tracer data regarding the source of
O atoms in NO_2^-. In both models, NirK and two membrane-anchored
cupredoxins (T478_1362 and T478_0895) act in concert to oxidize NH_2OH
to NO_2^- in two steps: a three-electron oxidation of NH_2OH to NO, followed
by a one-electron oxidation of NO to NO_2^- (Supplementary Figure 3). The
involvement of T478_0895 orthologs in electron transfer with NirK, is
consistent with previous predictions based on gene expression data for *N.*
maritimus (Qin *et al.*, 2017). Both proposed pathways are also consistent
with the observed co-expression and high abundance of these transcripts
across distinct environmental conditions (Figs 2-4; (Hollibaugh *et al.*, 2011;
Baker *et al.*, 2012; Stewart *et al.*, 2012; Gifford *et al.*, 2013)). The predicted
localization of ammonia oxidation in the pseudoperiplasm (Walker *et al.*,
2010) is also a key aspect of the proposed models, as protein domain
analysis with InterPro (Jones *et al.*, 2014) indicates all proteins in these
models, and pertinent cupredoxin domains, are likely localized in the
pseudoperiplasmic space. While we cannot determine which reaction is
conducted by which Cu metalloenzyme, both scenarios are more

parsimonious than existing models and are plausible based on existing
bioinorganic chemistry literature.
382
Similar to the *N. maritimus* transcriptome and environmental
metatranscriptomes (Hollibaugh *et al.,* 2011; Williams *et al.,* 2012; Qin *et*
al., 2017), ferredoxins were among the most highly expressed genes in *N.*
brevis (Fig. 2). However, we were surprised to find these ferredoxins (Fd1
and Fd2) and an Fe-S cluster assembly protein co-expressed with
ammonia oxidation genes in the AMO module. The co-expression of
ferredoxins in the AMO module suggests a central role for Fe-S cluster-
containing proteins in the electron transport chain of thaumarchaea. Both
Fd1 and Fd2 lack discernable signal sequences or PEFG domains,
suggesting that they are localized in the cytoplasm. Ferredoxin-containing
DNA binding transcriptional regulators have been implicated as NO sensors
(Kiley and Beinert, 2003). However, there are no predicted DNA-binding
domains in Fd1 or Fd2. Sequence structure threading of Fd1 and Fd2 with
phyre2 (Kelley *et al.,* 2015) returned best structural matches to NADH
dehydrogenase (ubiquinone) iron-sulfur protein 8 (threading confidence
score = 99.8% for both Fd1 and Fd2; coverage of Fd1 was 99% and 71%
for Fd2). Thus, we speculate that Fd1 and Fd2 participate in supplying
electrons to the ubiquinone pool, and thus may be involved in supplying the
electrons necessary to initiate ammonia oxidation via Amo.
402
Microbial gene expression is inherently modular. For example, functionally
related genes are often co-expressed in operons or regulons as a response
to external environmental signals (Freyre-Gonzales *et al.,* 2010). However,
the organization and patterns of population-level gene expression in
dynamic environments are poorly understood. Although we show a high
degree of modularity and correlated expression of a set of genes likely
involved in ammonia oxidation, the gene membership of the remaining
modules did not reveal a clear pattern of how the genes contained within a
given module are functionally related (Supplementary Table 3). For
example, genes coding for the modified 3-hydroxypropioinate/ 4-
hydroxybutryrate carbon fixation pathway proteins are dispersed across
five modules (module numbers 3, 4, 18, 21, and 36; Fig. 3 and
Supplementary Table 3). Similarly, vitamin B_{12}-production genes are spread
across eight modules (module numbers 1, 3, 4, 10, 12, 18, 21, and 26; Fig.
3 and Supplementary Table 3). The structure of the transcription co-

expression network illustrated in Fig. 3 is consistent with the interpretation that thaumarchaeal population-level regulatory organization is structured in a decentralized manner. One explanation for this organization may be that decentralized expression networks may help to buffer gene expression changes or maintain genetic diversity in dynamic environments through a suite of feedbacks without centralized regulatory mechanisms (Hartwell *et al.,* 1999).

Other reasons the gene membership in the co-expression modules may not reveal clear functional relationships may stem from the environments sampled or the analysis techniques used. For example, our methods likely underestimate the true modularity of *Ca*. N. brevis gene expression. Some of the larger expression modules may comprise distinct modules that we could not resolve because we did not sample an environment with physicochemical parameters necessary to resolve subtle gene expression patterns. Second, although our goal was to be conservative in our network construction, we may be missing important network structural components because of the thresholding parameters or our analysis techniques. Further resolution of such gene expression patterns would require deeply sequenced metatranscriptomes from additional distinct environments and transcriptome analysis of additional thaumarchaea grown under diverse culture conditions. Future research into understanding why certain genes or pathways are co-expressed with one another, and how transcript abundances manifest into functional potential in each environment or culture setting would be necessary to fully disentangle the gene expression we observe here.

Conclusions:
Here we show the rank of most thaumarchaeal transcripts reported as being abundant in the environment (*amoABC*, both *amtB* genes and *nirK*, for example) are relatively invariant across growth phases and environmental conditions. That is, within a given treatment, abundant genes are consistently proportionally abundant, irrespective of growth condition. However, consistent with other studies of thaumarchaeal transcription, the proportions of some of these genes were indeed significantly differentially abundant across paired treatments, indicating ammonia availability did affect the proportional abundances of ammonia oxidation and transport transcripts. One explanation for this observation is

456 that although these transcripts are not 'constitutive' in a classic sense (that
457 is, they are differentially abundant across paired experiments), they are
458 instead 'affluent,' in that they make up a large part of the total transcript
459 pool, irrespective of growth condition.
460
Discovering gene function in fastidious or uncultivated lineages remains
one of the biggest challenges in environmental microbiology. Narrowing the
scope of targets for detailed biochemical investigation is difficult because
manipulative experiments are limited in their ability to identify networks of
co-regulated genes by the number of environmental parameters we can
recreate in a laboratory. The approach used here leverages *in situ*
transcript abundance data - in which the environmental conditions are
incompletely characterized - to identify genes that share similar
transcription patterns. In addition to our putative models of ammonia
oxidation in thaumarchaea, this approach shows that 4Fe-4S cluster-
containing proteins likely have an important role in ammonia oxidation,
indicating a role for iron in archaeal nitrification, which has been previously
under appreciated. Detailed biochemical characterization of NirK, other
cupredoxin-containing proteins, Fd1 and Fd2 is the next step in
understanding their specific role in core thaumarchaeal energy metabolism.
476

Methods:
Organism sources: Both 'Candidatus Nitrosopelagicus brevis' str. CN25
and '*Ca.* N. brevis' str. U25 were obtained from an ammonia-oxidizing
enrichment culture previously known as CN25 (Santoro and Casciotti,
2011) grown on Oligotrophic North Pacific Medium. Oligotrophic North
Pacific Medium (ONP) consists of natural seawater, a
chemolithoautotrophic nitrogen source (NH_4Cl or urea), ampicillin (10.8
μM), streptomycin (68.6 μM), potassium phosphate (29.4 μM), and a
chelated trace metal mix consisting of disodium ethylenediaminetetraacetic
acid (14 μM), $FeCl_2$ (7.25 μM), $ZnCl_2$ (0.5 μM), $MnCl_2$ (0.5 μM), H_3BO_3 (1
μM), $CoCl_2$-$6H_2O$ (0.8 μM), $CuCl_2$-$2H_2O$ (0.1 μM), $NiCl_2$-H_2O (0.1 μM),
Na_2MoO_4-$2H_2O$ (0.15 μM). Preliminary metagenomic sequencing of the
original CN25 enrichment indicated the presence of urease genes in a
minority of the archaeal population. Sequential transfers of the initial
enrichment were made into ONP (Santoro and Casciotti, 2011) amended
with 50-100 μM urea, instead of NH_4Cl, over a period of ~48 months. In
parallel, separate transfers of the CN25 enrichment culture were

propagated using NH_4Cl as a nitrogen and energy source, but without the
use of streptomycin, which could potentially serve as a source of urea
(Klein and Pramer, 1961). The enrichment culture resulting from the
propagation with NH_4Cl did not contain amplifiable *ureC* genes by PCR,
using the archaeal-specific ureC primers Thaum_UreC F and Thaum_UreC
R (Yakimov *et al.*, 2011) before the sequencing and genome analysis
described elsewhere (Santoro *et al.*, 2015).
501
General cultivation conditions: All thaumarchaeal enrichments were
propagated in in ONP medium in a base of aged natural seawater
(collected from 10 m depth at 15°S, 173°W on 23 October 2011; 0.2 μm
pore size filtered at sea) amended with 50 or 100 μM NH_4Cl or 100 μM
urea as the chemolithoautotrophic substrate. All cultures were propagated
in 250 mL polycarbonate flasks at 22°C in the dark and monitored for NO_2^-
production using the Griess reagent colorimetric method (Strickland and
Parsons, 1972). Cell counts were obtained with a Millipore Guava
EasyCyte 5HT flow cytometer as described previously (Tripp, 2008).
511
Cell harvesting for and genome sequencing of 'Ca. *N. brevis' str. U25:* The
Ca. N. brevis U25 enrichment culture that was grown exclusively with urea
as the sole chemolithoautotrophic growth substrate for >50 generations,
was harvested by filtration on to 25 mm diameter, 0.22 μm pore-size Supor-
200 filters and frozen at -80°C. DNA was extracted using a DNeasy blood &
Tissue DNA extraction kit (Qiagen, Valencia, CA, USA), following the
manufacturer's instructions. The DNA was treated with RNAse and
examined using a Bioanlayzer 2100 (Agilent) with 500 ng serving as the
input for library construction (NEBNext paired-end DNA Library Prep kit,
New England Biolabs). The sample was sequenced on an Illumina MiSEQ
(v2 chemistry, paired 250 bp reads). Reads were quality trimmed and
served as the inputs to assembly with metaSPAdes (v 0.5, 70mer) (Nurk *et*
al., 2017). The K-mer usage and phylogenetic annotation of the assembled
contigs were then used to visually identify a putative thaumarchaeal bin
(Supplementary Figure 1a) (Laczny *et al.*, 2015). The 3 contig genome was
annotated using the JGI IMG pipeline (img.jgi.doe.gov) and the PGAP
pipeline at NCBI (Zhao *et al.*, 2011).
529
Experimental design, cell harvesting and RNA extraction for culture
transcriptomes: For experiments investigating the effects of ammonia

limitation, ONP medium was amended with 50 µM NH_4Cl as the
chemolithoautotrophic growth substrate. In this experiment, six replicates
were prepared, three of which were harvested in late exponential phase
(Exponential phase) and three of which were harvested in late NH_4Cl-
limited stationary phase (Stationary phase) (Supplementary Fig. 4). We
deliberately harvested the exponential phase cultures in late exponential
phase to ensure maximal cell biomass for transcriptome analysis. The
length of starvation in stationary-phase was based on the exponential-
phase doubling time of seven days, approximately the population doubling
time during exponential growth. NH_4Cl-limitation in this phase is supported
by a linear dose response in maximal cell density to NH_4Cl additions
(Supplementary Fig. 5).
544
Transcriptomes that were included in the network analysis were obtained
from distinct, mid-exponential phase *Ca*. N. brevis strains CN25 and U25,
that were growing on ONP medium amended with either 100 µM NH_4Cl
(str. CN25; n=3) or 100 µM urea (str. U25; n=3) as growth substrates,
respectively (Supplementary Fig. 6). Cells were harvested by filtration on to
25 mm diameter, 0.22 µm pore-size Supor-200 filters and frozen at -80°C.
For RNA extraction, cells were disrupted as described in (Santoro *et al.*,
2010). RNA was extracted using TRIzol LS reagent (Ambion-Life
Technologies) per the manufacturer's instructions and stored in nuclease-
free water at -80°C. Urea consumption by str. U25 was determined
colorimetrically using the diacetyl monoxime method (Price and Harrison,
1987) (Supplementary Fig. 7).
557
Transcriptome sequencing and mapping for culture experiments:
Transcriptome samples were prepared for sequencing using the TotalScript
RNA-Seq kit (Epicentre-Illumina), which biases against rRNA, using the
manufacturer recommended protocol. Libraries were trial sequenced on an
Illumina MiSEQ to determine uniformity between barcodes and then fully
sequenced in one 300 cycle NextSEQ run which generated 246.6 million
paired-end 150 bp reads. Raw Illumina reads in fastq format were
interleaved to match paired ends. Sequencing primers and barcode
indexes were identified by BLAST against the NCBI vector database and
trimmed along with regions with Q scores < 30. Reads mapping to
ribosomal RNAs were identified and removed using ribopicker (Schmieder
et al., 2011). Reads were then mapped to *Ca*. N. brevis genomes at 90%

nucleotide identity with CLC Genomics Workbench (command:
clc_ref_assemble –s 0.9). Raw read counts per open reading frame (ORF)
were compiled.
573
Analysis of differentially abundant gene transcripts: Differential gene
abundance analysis was performed using a generalized linear model
likelihood ratio test in the edgeR software package (v 3.8.5) (Robinson and
Smyth, 2008). We defined significant differential abundance as those
genes with a false discovery rate (FDR) ≤ 0.01 and greater than 2-fold
abundance change across treatments.
580
Rank Analyses: Raw read counts per ORF were scaled to expression units
of reads per base per million reads mapped (RPKM=$(10^6 * C)/(NL/10^3)$)
where C is the number of transcript reads mapped to an ORF; N is total
reads mapped to all ORFs in the genome; and L is the ORF length in base
pairs (Mortazavi *et al.*, 2008). RPKM values were subsequently ranked,
with a rank of 1 depicting the most abundant transcript within a given
treatment. Rank ties within a treatment were averaged.
588
Metatranscriptome mapping to genomes of Ca. N. brevis strains: Sequence
reads from 68 metatranscriptomes were mapped to the *Ca.* N. brevis
genomes at 50% nucleotide identity using CLC Genomics Workbench
(command: clc_ref_assemble –s 0.5) (Supplementary Table 4). The
number of metatranscriptome reads that mapped to the *Ca.* N. brevis
genomes were variable and ranged from 10 reads to 236,954 reads and
mapped to 0.5-89% of the unique genes in the *Ca.* N. brevis genomes
(Supplementary Table 4). Raw read counts per ORF were then compiled
(ORF n=1445 for str. CN25 and n=1461 for U25).
598
Network construction: Only those metatranscriptomes that mapped to
≥45% of the ORFs in the *Ca.* N. brevis genomes, along with the
transcriptomes from *Ca.* N. brevis strains CN25 and U25 growing in
exponential phase initiated with 100 µM NH_4Cl or urea, respectively, were
included for network analysis. Of the 68 metatranscriptomes mapped to the
Ca. N. brevis genomes, only ten passed this filtering step and were used
for network analysis (Supplementary Table 4). Read counts were scaled to
RPKM expression units. RPKM scores calculated for individual culture
transcriptome replicates were averaged (n=3) to avoid pseudo-replication

effects in the network. The resulting RPKM expression values were rank-
normalized to Van der Waerden (VdW) scores using the formula ($s = \Phi^{-1}(r /$
$(n+1)))$, where s is the VdW score for a gene, r is the rank for that
observation, n is the sample size and Φ is the Φ^{th} quantile from the
standard normal distribution using *tRank* in the multic R package (Lunde *et*
al.). Pearson correlation coefficients and P value estimates were calculated
for all gene:gene pairs across the VdW-normalized metatranscriptomes
and culture experiments (n=10 and 2, respectively) with the *rcorr* command
in the Hmisc R package (Harrell and Dupont). To correct for multiple
hypothesis testing, q values were computed from P value estimates using
the qvalue R package (Storey and Tibshirani, 2003). Correlations with a q
value ≤ 0.025 were used for network analysis. All correlations at this
threshold were strongly correlated (Pearson's $r \geq 0.8$).
621
622 *Network Statistics:* Network modularity and module membership were
623 calculated in Gephi (0.8.2 beta) with the following settings: resolution 1.0,
624 randomized and unweighted (Blondel *et al.*, 2008). The resulting network
625 was visualized using the Fruchterman-Reingold algorithm in Gephi.
626
Data Availability/Sources:
Transcriptomes from *Ca.* N. brevis str. CN25 and U25 can be found in the
NCBI BioSample archive under accession numbers SAMN6290440-
6290457. The U25 genome has been deposited at DDBJ/ENA/GenBank
under the accession LXWN00000000. The version described in this paper
is version LXWN01000000. The metatranscriptomic data from Landsort
Deep in the Baltic is available from the Sequence Read Archive under
numbers SAMN04943349-SAMN04943415.
Other metatranscriptomes analyzed in the network are publically available
through iMicrobe (https://imicrobe.us) or NCBI's Short Read Archive
through the following accession numbers: CAM_PROJ_Sapelo2008,
CAM_PROJ_AmazonRiverPlume, CAM_PROJ_PacificOcean,
CAM_P_0000545, SRA023632.1.
640
641 **Acknowledgements:**
642

643 This research was supported by NSF awards OCE-1260006, OCE-
644 1437310, and DBI-1318455 to AES. CLD was supported by NSF OCE-
645 1259994. We thank Matt Rawls for urea measurements, Mike Stukel for

References:

Alonso-Sáez, L., Waller, A.S., Mende, D.R., Bakker, K., Farnelid, H., Yager, P.L., et al. (2012) Role for urea in nitrification by polar marine Archaea. *Proc. Natl. Acad. Sci. U.S.A.* **109**: 17989–17994.

Arp, D.J., Chain, P.S.G., and Klotz, M.G. (2007) The impact of genome analyses on our understanding of ammonia-oxidizing bacteria. *Annu. Rev. Microbiol.* **61**: 503–528.

Baker, B.J., Lesniewski, R.A., and Dick, G.J. (2012) Genome-enabled transcriptomics reveals archaeal populations that drive nitrification in a deep-sea hydrothermal plume. *The ISME Journal* **6**: 2269–2279.

Bayer, B., Vojvoda, J., Offre, P., Alves, R.J.E., Elisabeth, N.H., Garcia, J.A., et al. (2015) Physiological and genomic characterization of two novel marine thaumarchaeal strains indicates niche differentiation. *The ISME Journal* **10**: 1051–1063.

Beman, J.M., Popp, B.N., and Alford, S.E. (2012) Quantification of ammonia oxidation rates and ammonia-oxidizing archaea and bacteria at high resolution in the Gulf of California and eastern tropical North Pacific Ocean. *Limnol. Oceangr.* **57**: 711–726.

Berube, P.M. and Stahl, D.A. (2012) The divergent AmoC3 subunit of ammonia monooxygenase functions as part of a stress response system in *Nitrosomonas europaea*. *Journal of Bacteriology* **194**: 3448–3456.

Berube, P.M., Samudrala, R., and Stahl, D.A. (2007) Transcription of all amoC copies is associated with recovery of *Nitrosomonas europaea* from ammonia starvation. *Journal of Bacteriology* **189**: 3935–3944.

Blondel, V.D., Guillaume, J.-L., Lambiotte, R., and Lefebvre, E. (2008) Fast unfolding of communities in large networks. *Journal of Statistical Mechanics: Theory and Experiment* **10**: 10008–.

Buchwald, C., Santoro, A.E., McIlvin, M.R., and Casciotti, K.L. (2012) Oxygen isotopic composition of nitrate and nitrite produced by nitrifying cocultures and natural marine assemblages. *Limnol. Oceangr.* **57**: 1361–1375.

Church, M.J., Wai, B., Karl, D.M., and DeLong, E.F. (2010) Abundances of

crenarchaeal *amoA* genes and transcripts in the Pacific Ocean.
Environmental Microbiology **12**: 679–688.
Coolen, M.J.L., Abbas, B., van Bleijswijk, J., Hopmans, E.C., Kuypers,
M.M.M., Wakeham, S.G., and Sinninghe Damsté, J.S. (2007) Putative
ammonia-oxidizing Crenarchaeota in suboxic waters of the Black Sea: a
basin-wide ecological study using 16S ribosomal and functional genes
and membrane lipids. *Environmental Microbiology* **9**: 1001–1016.
Cottrell, M.T. and Kirchman, D.L. (2016) Transcriptional control in marine
copiotrophic and oligotrophic bacteria with streamlined genomes.
Applied and Environmental Microbiology **82**: 6010–6018.
Francis, C.A., Roberts, K.J., Beman, J.M., Santoro, A.E., and Oakley, B.B.
(2005) Ubiquity and diversity of ammonia-oxidizing archaea in water
columns and sediments of the ocean. *Proc. Natl. Acad. Sci. U.S.A.* **102**:
14683–14688.
Freyre-Gonzales, J.A., Educ, L.T.-Q.N., 2010 (2010) Analyzing regulatory
networks in bacteria. *Nature Education* **3**: 24–27.
Gifford, S.M., Sharma, S., Booth, M., and Moran, M.A. (2013) Expression
patterns reveal niche diversification in a marine microbial assemblage.
The ISME Journal **7**: 281–298.
Giovannoni, S.J. (2017) SAR11 Bacteria: The most abundant plankton in
the oceans. *Annu. Rev. Marine. Sci.* **9**: 231–255.
Giovannoni, S.J., Cameron Thrash, J., and Temperton, B. (2014)
Implications of streamlining theory for microbial ecology. *The ISME
Journal* **8**: 1553–1565.
Harrell, F.E., Jr and Dupont, C. Hmisc: Harrell Miscellaneous.
Hartwell, L.H., Hopfield, J.J., Leibler, S., and Murray, A.W. (1999) From
molecular to modular cell biology. *Nature* **402**: C47–C52.
Hollibaugh, J.T., Gifford, S., Sharma, S., Bano, N., and Moran, M.A. (2011)
Metatranscriptomic analysis of ammonia-oxidizing organisms in an
estuarine bacterioplankton assemblage. *The ISME Journal* **5**: 866–878.
Jones, P., Binns, D., Chang, H.-Y., Fraser, M., Li, W., McAnulla, C., et al.
(2014) InterProScan 5: genome-scale protein function classification.
Bioinformatics **30**: 1236–1240.
Karner, M.B., Delong, E.F., and Karl, D.M. (2001) Archaeal dominance in
the mesopelagic zone of the Pacific Ocean. *Nature* **409**: 507–510.
Kelley, L.A., Mezulis, S., Yates, C.M., Wass, M.N., and Sternberg, M.J.E.
(2015) The Phyre2 web portal for protein modeling, prediction and
analysis. *Nat Protoc* **10**: 845–858.

Kerou, M., Offre, P., Valledor, L., Abby, S.S., Melcher, M., Nagler, M., et al.
(2016) Proteomics and comparative genomics of *Nitrososphaera*
viennensis reveal the core genome and adaptations of archaeal
ammonia oxidizers. *Proc. Natl. Acad. Sci. U.S.A.* 201601212–19.
Kiley, P.J. and Beinert, H. (2003) The role of Fe–S proteins in sensing and
regulation in bacteria. *Current Opinion in Microbiology* **6**: 181–185.
Klein, D. and Pramer, D. (1961) Bacterial dissimilation of streptomycin.
Journal of Bacteriology **82**: 505–510.
Kozlowski, J.A., Stieglmeier, M., Schleper, C., Klotz, M.G., and Stein, L.Y.
(2016) Pathways and key intermediates required for obligate aerobic
ammonia-dependent chemolithotrophy in bacteria and
Thaumarchaeota. *The ISME Journal* **10**: 1836–1845.
Laczny, C.C., Sternal, T., Plugaru, V., Gawron, P., Atashpendar, A.,
Margossian, H., et al. (2015) VizBin - an application for reference-
independent visualization and human-augmented binning of
metagenomic data. *Microbiome* **3**: 1–7.
Lam, P., Cowen, J.P., and Jones, R.D. (2004) Autotrophic ammonia
oxidation in a deep-sea hydrothermal plume. *FEMS Microbiology
Ecology* **47**: 191–206.
Lam, P., Lavik, G., Jensen, M.M., van de Vossenberg, J., Schmid, M.,
Woebken, D., et al. (2009) Revising the nitrogen cycle in the Peruvian
oxygen minimum zone. *Proc. Natl. Acad. Sci. U.S.A.* **106**: 4752–4757.
Li, D.-C., Yang, F., Lu, B., Chen, D.-F., and Yang, W.-J. (2011)
Thermotolerance and molecular chaperone function of the small heat
shock protein HSP20 from hyperthermophilic archaeon, *Sulfolobus
solfataricus P2. Cell Stress and Chaperones* **17**: 103–108.
Lindås, A.-C., Karlsson, E.A., Lindgren, M.T., Ettema, T.J.G., and
Bernander, R. (2008) A unique cell division machinery in the Archaea.
Proc. Natl. Acad. Sci. U.S.A. **105**: 18942–18946.
Lunde, E., de Andrade, M., Atkinson, B., and Votruba, P. multic:
Quantitative linkage analysis tools using the variance
components approach.
Martens-Habbena, W., Qin, W., Horak, R.E.A., Urakawa, H., Schauer, A.J.,
Moffett, J.W., et al. (2015) The production of nitric oxide by marine
ammonia-oxidizing archaea and inhibition of archaeal ammonia
oxidation by a nitric oxide scavenger. *Environmental Microbiology* **17**:
2261–2274.
Mortazavi, A., Williams, B.A., McCue, K., Schaeffer, L., and Wold, B.

(2008) Mapping and quantifying mammalian transcriptomes by RNA-Seq. *Nat Meth* **5**: 621–628.

Nakagawa, T. and Stahl, D.A. (2013) Transcriptional response of the archaeal ammonia oxidizer *Nitrosopumilus maritimus* to low and environmentally relevant ammonia concentrations. *Applied and Environmental Microbiology* **79**: 6911–6916.

Nurk, S., Meleshko, D., Korobeynikov, A., and Pevzner, P.A. (2017) metaSPAdes: a new versatile metagenomic assembler. *Genome Research* **27**: 824–834.

Pedneault, E., Galand, P.E., Potvin, M., Tremblay, J.-É., and Lovejoy, C. (2014) Archaeal *amoA* and *ureC* genes and their transcriptional activity in the Arctic Ocean. *Sci. Rep.* **4**: 1–11.

Pellitteri-Hahn, M.C., Halligan, B.D., Scalf, M., Smith, L., and Hickey, W.J. (2011) Quantitative proteomic analysis of the chemolithoautotrophic bacterium *Nitrosomonas europaea*: Comparison of growing- and energy-starved cells. *Journal of Proteomics* **74**: 411–419.

Pitcher, A., Villanueva, L., Hopmans, E.C., Schouten, S., Reichart, G.-J., and Sinninghe Damsté, J.S. (2011) Niche segregation of ammonia-oxidizing archaea and anammox bacteria in the Arabian Sea oxygen minimum zone. *The ISME Journal* **5**: 1896–1904.

Price, N.M. and Harrison, P.J. (1987) Comparison of methods for the analysis of dissolved urea in seawater. *Marine Biology* **94**: 307–317.

Qin, W., Amin, S.A., Lundeen, R.A., Heal, K.R., Martens-Habbena, W., Turkarslan, S., et al. (2017) Stress response of a marine ammonia-oxidizing archaeon informs physiological status of environmental populations. *The ISME Journal* 1–12.

Qin, W., Amin, S.A., Martens-Habbena, W., Walker, C.B., Urakawa, H., Devol, A.H., et al. (2014) Marine ammonia-oxidizing archaeal isolates display obligate mixotrophy and wide ecotypic variation. *Proc. Natl. Acad. Sci. U.S.A.* **111**: 12504–12509.

Robinson, M.D. and Smyth, G.K. (2008) Small-sample estimation of negative binomial dispersion, with applications to SAGE data. *Biostatistics* **9**: 321–332.

Santoro, A.E. and Casciotti, K.L. (2011) Enrichment and characterization of ammonia-oxidizing archaea from the open ocean: phylogeny, physiology and stable isotope fractionation. *The ISME Journal* **5**: 1796–1808.

Santoro, A.E., Buchwald, C., McIlvin, M.R., and Casciotti, K.L. (2011)

798  Isotopic signature of N_2O produced by marine ammonia-oxidizing
799  archaea. *Science* **333**: 1282–1285.

800 Santoro, A.E., Casciotti, K.L., and Francis, C.A. (2010) Activity, abundance
801  and diversity of nitrifying archaea and bacteria in the central California
802  Current. *Environmental Microbiology* **12**: 1989–2006.

803 Santoro, A.E., Dupont, C.L., Richter, R.A., Craig, M.T., Carini, P., McIlvin,
804  M.R., et al. (2015) Genomic and proteomic characterization of
805  "*Candidatus* Nitrosopelagicus brevis": an ammonia-oxidizing archaeon
806  from the open ocean. *Proc. Natl. Acad. Sci. U.S.A.* **112**: 1173–1178.

807 Satinsky, B.M., Smith, C.B., Sharma, S., Ward, N.D., Krusche, A.V.,
808  Richey, J.E., et al. (2017) Patterns of Bacterial and Archaeal gene
809  expression through the lower Amazon River. *Front. Mar. Sci.* **4**: 1937–
810  15.

811 Schattenhofer, M., Fuchs, B.M., Amann, R., Zubkov, M.V., Tarran, G.A.,
812  and Pernthaler, J. (2009) Latitudinal distribution of prokaryotic
813  picoplankton populations in the Atlantic Ocean. *Environmental*
814  *Microbiology* **11**: 2078–2093.

815 Schmieder, R., Lim, Y.W., and Edwards, R. (2011) Identification and
816  removal of ribosomal RNA sequences from metatranscriptomes.
817  *Bioinformatics* **28**: 433–435.

818 Shi, Y., Tyson, G.W., Eppley, J.M., and DeLong, E.F. (2010) Integrated
819  metatranscriptomic and metagenomic analyses of stratified microbial
820  assemblages in the open ocean. *The ISME Journal* **5**: 999–1013.

821 Smith, D.P., Kitner, J.B., Norbeck, A.D., Clauss, T.R., Lipton, M.S.,
822  Schwalbach, M.S., et al. (2010) Transcriptional and translational
823  regulatory responses to iron limitation in the globally distributed marine
824  bacterium *'Candidatus* Pelagibacter ubique'. *PLoS ONE* **5**: e10487–10.

825 Smith, D.P., Nicora, C.D., Carini, P., Lipton, M.S., Norbeck, A.D., Smith,
826  R.D., and Giovannoni, S.J. (2016) Proteome remodeling in response to
827  sulfur limitation in "*Candidatus* Pelagibacter ubique." *mSystems* **1**:
828  e00068–16–15.

829 Smith, D.P., Thrash, J.C., Nicora, C.D., Lipton, M.S., Burnum-Johnson,
830  K.E., Carini, P., et al. (2013) Proteomic and transcriptomic analyses of
831  "*Candidatus* Pelagibacter ubique" describe the first PII-independent
832  response to nitrogen limitation in a free-living Alphaproteobacterium.
833  *mBio* **4**: e00133–12.

834 Smith, J.M., Casciotti, K.L., Chavez, F.P., and Francis, C.A. (2014)
835  Differential contributions of archaeal ammonia oxidizer ecotypes to

nitrification in coastal surface waters. *The ISME Journal* **8**: 1704–1714.

Spang, A., Poehlein, A., Offre, P., Zumbrägel, S., Haider, S., Rychlik, N., et
al. (2012) The genome of the ammonia-oxidizing '*Candidatus*
Nitrososphaera gargensis': insights into metabolic versatility and
environmental adaptations. *Environmental Microbiology* **14**: 3122–3145.

Spang, A., Saw, J.H., Jørgensen, S.L., Zaremba-Niedzwiedzka, K., Martijn,
J., Lind, A.E., et al. (2015) Complex archaea that bridge the gap
between prokaryotes and eukaryotes. *Nature* **521**: 173–179.

Stahl, D.A. and la Torre, de, J.R. (2012) Physiology and diversity of
ammonia-oxidizing archaea. *Annu. Rev. Microbiol.* **66**: 83–101.

Stewart, F.J., Ulloa, O., and DeLong, E.F. (2012) Microbial
metatranscriptomics in a permanent marine oxygen minimum zone.
Environmental Microbiology **14**: 23–40.

Stieglmeier, M., Mooshammer, M., Kitzler, B., Wanek, W., Zechmeister-
Boltenstern, S., Richter, A., and Schleper, C. (2014) Aerobic nitrous
oxide production through N-nitrosating hybrid formation in ammonia-
oxidizing archaea. *The ISME Journal* **8**: 1135–1146.

Storey, J.D. and Tibshirani, R. (2003) Statistical significance for
genomewide studies. *Proc. Natl. Acad. Sci. U.S.A.* **100**: 9440–9445.

Strickland, J.D.H. and Parsons, T.R. (1972) A Practical Handbook of
Seawater Analysis. 2.ed.

Tolar, B.B., Ross, M.J., Wallsgrove, N.J., Liu, Q., Aluwihare, L.I., Popp,
B.N., and Hollibaugh, J.T. (2016) Contribution of ammonia oxidation to
chemoautotrophy in Antarctic coastal waters. *The ISME Journal* **10**:
2605–2619.

Tolar, B.B., Wallsgrove, N.J., Popp, B.N., and Hollibaugh, J.T. (2016)
Oxidation of urea-derived nitrogen by thaumarchaeota-dominated
marine nitrifying communities. *Environmental Microbiology*.

Tripp, H.J. (2008) Counting marine microbes with Guava Easy-Cyte 96 well
plate reading flow cytometer. *Protocol Exchange*.

Vajrala, N., Martens-Habbena, W., Sayavedra-Soto, L.A., Schauer, A.,
Bottomley, P.J., Stahl, D.A., and Arp, D.J. (2013) Hydroxylamine as an
intermediate in ammonia oxidation by globally abundant marine
archaea. *Proc. Natl. Acad. Sci. U.S.A.* **110**: 1006–1011.

Walker, C.B., la Torre, de, J.R., Klotz, M.G., Urakawa, H., Pinel, N., Arp,
D.J., et al. (2010) *Nitrosopumilus maritimus* genome reveals unique
mechanisms for nitrification and autotrophy in globally distributed
marine crenarchaea. *Proc. Natl. Acad. Sci. U.S.A.* **107**: 8818–8823.

Wei, X., Yan, T., Hommes, N.G., Liu, X., Wu, L., McAlvin, C., et al. (2006)
Transcript profiles of *Nitrosomonas europaea* during growth and upon
deprivation of ammonia and carbonate. *FEMS Microbiology Letters* **257**:
76–83.
Williams, T.J., Long, E., Evans, F., DeMaere, M.Z., Lauro, F.M., Raftery,
M.J., et al. (2012) A metaproteomic assessment of winter and summer
bacterioplankton from Antarctic Peninsula coastal surface waters. *The*
ISME Journal **6**: 1883–1900.
Yakimov, M.M., La Cono, V., Smedile, F., DeLuca, T.H., rez, S.J.A.,
Ciordia, S., et al. (2011) Contribution of crenarchaeal autotrophic
ammonia oxidizers to the dark primary production in Tyrrhenian deep
waters (Central Mediterranean Sea). *The ISME Journal* **5**: 945–961.
Zhao, Y., Wu, J., Yang, J., Sun, S., Xiao, J., and Yu, J. (2011) PGAP: pan-
genomes analysis pipeline. *Bioinformatics* **28**: 416–418.
888
889

Figures:

Figure 1: Urea transport and catalysis genes are frequently the most highly expressed *Ca*. N. brevis U25-specific genes. (a) Chromosomal orientation of the *Ca*. N. brevis str. U25 indel conferring urea transport and catalysis capability. (b) Heatmap illustrating the relative rank expression level of the five most abundant genes within the urea indel region for culture experiments and environmental metatranscriptomes. Rank was calculated within a given treatment from RPKM normalized expression values, where a rank of 1 is the most abundant transcript of the genes contained in the indel. The mean expression (n=3 replicates) value was used for ranking the culture treatment. ND=Not detected. Note this is not the rank of the transcript within the entire metatranscriptome. Some metatranscriptomes were excluded because they did not have sufficient coverage of the urea transport and catabolism machinery. Numbers in parentheses after gene names refer to A7X95 locus tags.

906
Figure 2: Highly expressed *Ca*. N. brevis str. CN25 transcripts in
exponential phase are also highly expressed in stationary phase, despite
significant differences in abundance. (a) Points are the mean rank (n=3) of
RPKM normalized expression values for all genes in exponential and
stationary growth phases. Red points are transcripts that were significantly
differentially abundant across treatments (Supplementary Table 2). The
abundances of grey points were not significantly different across
treatments. Dashed line is 1:1 line indicating no change in rank. (b) Subset
of panel (a), illustrating the rank of transcripts that are in the top 50 most
abundant transcripts in both exponential and stationary phase. Points in (b)
are the mean rank and error bars represent ± SE (n=3). Gene transcript
abundances in (b) that were significantly different across treatments are
labeled and colored red. The 'T478_' prefix is omitted from labels of genes
annotated as 'hypothetical'. PEFG corresponds to T478_0596.

Module 11 (AMO module) membership details

A: *amoA* (0302)	I: ESCRT-III (0057)
B: *amoB* (0298)	J: PEFG-CTERM (0270)
C: *amoC* (0300)	K: *nirK* (1026)
D: Membrane-bound cupredoxin (0895)	L: *amoX* (0301)
E: 0487	M: AMT1 (1378)
F: Fd1 (1472)	N: Cupredoxin PEFG-CTERM (1362)
G: Fd2 (1259)	O: Fe-S cluster assembly protien (1056)
H: 1317	

Figure 3: Thaumarchaeal gene expression is highly modular and ammonia oxidation genes are co-expressed. Network diagram of strong and significant (Pearson's r ≥ 0.8, *q* value ≤ 0.025) positive correlations across ten environmental metatranscriptomes and the *Ca.* N. brevis U25

transcriptome depicted in Fig. 1 and a *Ca.* N. brevis CN25 transcriptome
from a culture initiated with 100 μM NH_4Cl (12 conditions total, see
methods). Individual nodes are genes. Nodes are sized by the mean
normalized rank abundance (VdW scores), whereby larger nodes are more
abundant transcripts on average. Nodes are colored by module
membership. Circled numbers are the module identity (Supplementary
Table 3). The module 11 (the AMO module) genes are identified by letters
A-O; their annotations are provided below network. Numbers in
parentheses refer to T478 locus tags. Only modules with five or more
nodes are shown for clarity; see Supplementary Table 3 for all module
membership details.
937

938
Fig. 4: Natural variability in the mean rank of expression modules across
environments and culture conditions. Points are the average rank of all
genes contained within a given module at each site; error bars are ± SE.
On average, the AMO module and modules 36, 4 and 21 rank the highest
(that is, they are the most abundant) across all sites. Module 36 is not
displayed because it is only comprised of two genes, which prevents
statistical analysis, and these genes are absent in the U25 genome.