Title

Toward Capturing the Exposome: Exposure Biomarker Variability and Co-Exposure Patterns in

the Shared Environment

Running title:

Exposome variability in the household setting

Ming Kei Chung,[1] Kurunthachalam Kannan,[2] Germaine M. Buck Louis,[3] and Chirag J. Patel[1]

[1]Department of Biomedical Informatics

Harvard Medical School

Harvard University

10 Shattuck Street

Boston, MA 02115

[2]Division of Environmental Health Sciences

Wadsworth Center, New York State Department of Health, and

Department of Environmental Health Sciences

The University at Albany

Albany, New York 12201

Kurunthachalam.kannan@health.ny.gov

[3]Office of the Director

Division of Intramural Population Health Research

Eunice Kennedy Shriver National Institute for Child Health and Human Development

The National Institutes of Health

6710b Rockledge Drive

Bethesda, MD 20892

1-301-496-6155

louisg@mail.nih.gov

<u>Corresponding author:</u>

Chirag J. Patel

Department of Biomedical Informatics

Harvard Medical School

Harvard University

10 Shattuck Street, Room 302

Boston, MA 02115

1-617-432-1195

chirag_patel@hms.harvard.edu

Acknowledgement

The authors declare they have no actual or potential competing financial interests. This study is
supported by the NIH grants (ES025052 and ES023504), and the Intramural Research Program
of the *Eunice Kennedy Shriver* National Institute of Child Health and Human Development
(contracts # N01-HD-3-3355; N01-HD-3-3356; N01-HD-3-3358; HHSN27500002;
HHSN27500003; HHSN27500006)

Abstract

BACKGROUND: Along with time, variation in the exposome is dependent on the location and

sex of study participants. One specific factor that may influence exposure co-variations is a

shared household environment.

OBJECTIVES: To examine the influence of shared household and partner's sex in relation to the

variation in 128 endocrine disrupting chemical (EDC) exposures among couples.

METHODS: In a cohort comprising 501 couples trying for pregnancy, we measured 128 (13

chemical classes) persistent and non-persistent EDCs and estimated 1) sex-specific differences;

2) variance explained by shared household; and 3) Spearman's rank correlation coefficients (r_s)

for females, males, and couples' exposures.

RESULTS: Sex was correlated with 8 EDCs including polyfluoroalkyl substances (PFASs) ($p <$

0.05). Shared household explained 43% and 41% of the total variance for PFASs and blood

metals, respectively, but less than 20% for the remaining 11 EDC classes. Co-exposure patterns

of the exposome were similar between females and males, with within-class r_s higher for

persistent and lower for non-persistent chemicals. Median r_ss of polybrominated compounds

and urine metalloids were 0.45 and 0.09, respectively, for females (0.41 and 0.08 for males),

whereas lower r_ss for these 2 classes were found for couples (0.21 and 0.04).

CONCLUSIONS: Overall, sex did not significantly affect EDC levels in couples. Individual, rather

than shared environment, could be a major factor influencing the co-variation of 128 markers

of the exposome. Correlations between exposures are lower in couples than in individual

partners and have important analytical and sampling implications for epidemiological study.

82   Keywords: exposome, co-exposures, EDCs, combined exposure, POPs, metals, phthalates,

83   persistent pollutants, mixtures, biomarkers, exposure assessment

Variation of environmental exposure levels in the population is a complex phenomenon and is

influenced by factors shared by individuals — such as those within a household — and non-

shared factors specific to individuals, such as their sex. One could apply the exposome

paradigm (Wild 2005), a model to capture the totality of exposures from conception onwards,

to comprehensively characterize the individual-level and shared differences of exposure.

However, the exposome is a dynamic entity with variations across time (temporal) and place

(spatial) (Wild et al. 2013) underscoring the importance of considering variability when

assessing human health.

Instead of trying to "capture all" lifetime exposures, investigators can focus on critical and

sensitive time windows in human development such as pregnancy, infancy, childhood and

adolescence to reduce temporal complexity (Stingone et al. 2017). Furthermore, household-

level ascertainments of exposure (i.e., sampling individuals from households) has been posited

to be sufficient surrogates for all individuals in the household (Potera 2014). These assumptions

may help characterize the exposome and study its time-dependent relationship with health

outcomes. For example, the Human Early-Life Exposome (HELIX) project seeks to define the

pregnancy and early-life exposomes and health (Vrijheid et al. 2014), while the EXPOsOMICS

project has its conceptual framework of a life-course approach to a broader range of exposures

(Vineis et al. 2016).

Another challenge includes the difficulty of interpreting exposure-disease associations due to

the dense correlations among all exposures (Ioannidis 2016). The dense correlation pattern

makes it hard to identify the directionality of the potential causality (Ioannidis et al. 2009).

Second, correlations between exposures vary (e.g. absolute median correlation from almost 0

to above 0.5) and, thus, there is no universal scale to assess the biological significance (Patel

and Ioannidis 2014). In addition, exposome-wide, or equivalently, environment-wide

association studies (EWASs), assess all the associations between exposures and an outcome to

identify potential etiologic signals (Manrai et al. 2017). The data-driven approach assumes little

to no collinearity between environmental predictors, but it is almost impossible to select any

single uncorrelated exposures out from the dense exposome. One strategy for addressing these

analytical issues is to characterize the correlations in diverse cohorts to provide reference levels

to gauge biological significance of associations.

We investigated whether cohabiting couples trying for pregnancy would have similar

concentrations of endocrine disrupting chemicals (EDCs) given their shared households, and

whether concentrations varied across partners in light of their individual exposures arising from

other environments such as lifestyle, recreation or occupation. This avenue of study is

important given that EDCs have been found to affect human fecundity and fertility (Hauser

2006; te Velde et al. 2010), though much of the available evidence relies on research conducted

in either men or women but not couples. We utilized the Longitudinal Investigation of Fertility

and the Environment (LIFE) Study to empirically assess couples' shared and individual variations

in a mixture of EDCs. We selected the LIFE Study because it has 13 classes of EDCs quantified in

both partners of the couple in keeping with the couple based nature of human reproduction

(Buck Louis et al. 2014, 2016; Patel et al. 2016). To meet our overarching aim, we characterize

the dense correlation structure of couples' EDC concentrations and then estimate shared and

individual variability. Lastly, we discuss the implications of the findings in designing future

exposome-related research.

Methods

Study Design and Cohort

Briefly, the LIFE Study enrolled 501 couples planning to discontinue contraception to become

pregnant from 16 counties in Michigan and Texas, 2005–2009. Couples were followed daily

until pregnant or up to 12 months of trying to become pregnant (infertile). Study participants

were screened for eligibility based on a set of criteria and the complete details have been

previously published (Buck Louis et al. 2011).

Data Collection and Toxicologic Analysis

Following enrollment and completion of the baseline interview, couples provided

preconception blood (20 mL) and urine (120 mL) samples for the quantification of both man-

made and natural EDCs (e.g. phytoestrogens). We also included serum cotinine, a metabolite of

nicotine, and have a total of 128 chemicals from 13 different classes (Table 1). Persistent EDCs

included 4 classes of serum persistent organic compounds: 36 congeners of polychlorinated

biphenyls (PCBs), 9 organochlorine pesticides (OCPs), 11 polybrominated chemicals

[polybrominated diphenyl ethers (PBDEs) and 1 polybrominated biphenyl (PBB)], and 7

polyfluoroalkyl substances (PFASs), and were quantified by a single laboratory using published

standard operating procedures (Kuklenyik et al. 2005; Sjödin et al. 2004).

Non-persistent ECDs included 5 classes of urinary non-persistent organic compounds: 6

phytoestrogens, 14 phthalate metabolites, 6 phenols [bisphenol A (BPA) and benzophenones

(BPs)], 12 antimicrobial chemicals [parabens, triclosan (TCS) and triclocarban (TCC)], 2

paracetamol & derivatives, and were quantified by another laboratory using published standard

operating procedures (Asimakopoulos et al. 2014; Guo et al. 2011; Mumford et al. 2015; Smarr

et al. 2016). Other 3 classes of EDCs included 3 blood metals, 17 urinary metals, and 4 urinary

metalloids (Bloom et al. 2015a).

Serum lipids were obtained by measuring total cholesterol, free cholesterol, triglycerides, and

phospholipids by an enzymatic method (Akins et al. 1989) and we calculated the total serum

lipids as described by Phillips et al. (1989). Creatinine was measured by a Roche/Hitachi Model

912 clinical analyzer (Dalla, TX, USA) using the Creatinine Plus Assay (Roche Diagnostics). Serum

cotinine, a metabolite of nicotine, was quantified by isotope dilution tandem mass

spectrometry (Bernert et al. 1997).

Statistical Analysis

Our overall analytical scheme is shown in Figure 1. First, we adjusted each of the indicators of

exposures for potential confounders in addition to total lipids (for lipophilic chemicals, $n = 56$)

and creatinine (for urinary chemicals, $n = 61$) to reduce estimate variability and susceptibility to

bias (Heavner et al. 2006; Schisterman et al. 2005). Specifically, we adjusted PCBs, OCPs, and

polybrominated chemicals for total lipids and all other urinary chemicals for creatinine.

Chemicals were log-transformed (x + 1) and continuous covariates were rescaled to have mean

zero and unit variance (Figure 1A). After extracting the residuals, we calculated the Spearman's

rank correlation (r_s) matrices for EDCs in females, males, and couples.

Adjusting procedures such as Bonferroni correction assume independence between

independent variables. One way to resolve this analytical issue is to account for the correlation.

To study the effect of correlations between exposures on the family-wise error rate, we

calculated the effective number of variables (M_{eff}) (Nyholt 2004) for estimating the Bonferroni

adjusted p values with formula 1, where $Var(\lambda_{obs})$ is the eigenvalue variance of the correlation

matrix and M is the original number of variables.

$$M_{eff} = 1 + (M - 1)\left(1 - \frac{Var(\lambda_{obs})}{M}\right) \dots \text{formula 1}$$

We estimated the sex-specific difference with a paired t-test (by household) after extracting the

residuals from a linear model adjusted for age (Figure 1B). We used a similar approach to

estimate the percentage of variance explained by the shared environment. However, sex and

age variables were excluded in the adjustment step to isolate their effects (Figure 1C).

Afterward, we extracted and regressed the residuals against the household variable to obtain

the adjusted coefficient of determination (R^2).

We computed concordance as the Pearson correlation coefficient (r) between the chemical

relatedness r_s in this study and that in the 2003–2004 National Health and Nutrition

Examination Survey (NHANES) (Patel and Ioannidis 2014) to assess generalizability of the co-

exposure patterns. We estimated the concordance based on a total of 101 matched biomarkers

between the 2 studies. We chose the 2003–2004 NHANES because 1) many of the chemicals

measured in LIFE were also measured in NHANES and 2) by the same laboratory; and 3) the

time period (2003–2004) is close to the beginning of recruitment for the LIFE Study.

We used all the instrument derived concentrations for the analyses (Richardson and Ciampi

2003; Schisterman et al. 2006). For missing values, we substituted them by multiple imputation,

assuming a missing-at-random scenario (Louis et al. 2013). We conducted imputation based on

the information from available demographic, previous history of clinical symptoms and all other

chemical variables and created a total of 10 imputed data sets for males and females

separately.

We visualized the correlations between exposures as exposome globe using the R package

Circlize (Champely 2017). EDCs were sorted from lipophilic to hydrophilic to aid visual

interpretation of the patterns. We combined the final estimates from imputations using Rubin's

method (Schafer 1999) and calculate the p values of correlations by permutation tests. To

adjust for multiple testing, we used the false discovery rate (FDR) q values unless otherwise

specified. We executed all analyses using the computing environment R (v 3.3.1) (R Core Team

2016). For reproducibility purpose, all analytic code is publicly available on GitHub via a MIT

license (github.com/jakemkc/exposome_variability).

Results

Important sociodemographic differences were observed between partners (Table 2). Overall,

female partners were younger, had a lower body mass index (BMI) (< 25), consumed fewer

alcoholic drinks, were less likely to report a hypertension or high cholesterol, and had lower

serum cotinine and lipids and creatinine than male partners ($p < 0.01$).

Nine (7%) chemicals were correlated with partner's sex ($p < 0.05$), i.e., cotinine, blood lead,

mercury and cadmium; and 5 PFASs: perfluorodecanoate (PFDeA), perfluorononanoate (PFNA),

perfluorooctane sulfonamide (PFOSA), perfluorooctane sulfonate (PFOS), and

perfluorooctanoate (PFOA). Of note, findings were robust to the FDR with the exception of

blood cadmium.

Figure 2 shows the boxplot summary of the variance explained by the shared environment. We

estimated that two classes of chemicals, PFASs and blood metals, had higher levels of explained

variance (median 0.43 and 0.41 respectively) than the others. For the rest of the 11 classes,

median explained variances were ranged from 0.04 (phthalates) to 0.21 (cotinine). A few

persistent organic compounds, namely PCB congener 28, PBDE congener 47, and β-

hexachlorocyclohexane (β-HCH), had an explained variance over 50%.

The exposome globe (Patel and Manrai 2015) displays the r_ss for females (right-half), for males

(left-half), and for couples (across the left-right of the globe) (Figure 3). To assist interpretation,

we only presented the r_ss outside the range of −0.25 to 0.25 as lines connecting different parts

on the track, and they represent less than 10% of all the r_ss. For females, we observed two

larger positively correlated "clusters" across EDC classes: A) a dense cluster with serum

persistent organic compounds such as PCBs and OCPs (upper right of Figure 3); B) another

loosely packed cluster with urinary EDCs such as phytoestrogens, phthalates, phenols, and

antimicrobial compounds (lower right of Figure 3). Correlations between serum and urinary

EDCs were mostly small and distributed between −0.25 and 0.25. For males, there were similar

co-exposure patterns to females

While we found similar correlations in the population of males and females separately, we

found that correlations in couple living in the same household were, in fact, less densely packed

and with values attenuated toward the null (Figure 3; see Figure S1).

Summary of the within-class correlations as absolute magnitude is shown in Figure 4. For

females (Figure 4A), we found that polybrominated compounds had the highest median

correlation (r_s = 0.45) while urine metalloids had the lowest (r_s = 0.09). Across different

chemical classes, persistent organic compounds such as polybrominated compounds, PCBs and

OCPs had higher median correlations (0.45, 0.38, and 0.34 respectively). For the rest of the

classes, the median correlations were all below 0.25. Males (Figure 4B) had similar within-class

correlation distributions as were found in females. The class with highest and lowest median

correlations were polybrominated compounds (r_s = 0.41) and paracetamols (r_s = 0.02)

respectively.

In contrast, we found a strong diminishing effects to the within-class correlations, both in terms

of the median and interquartile range (IQR), in couples (Figure 4C). Comparing the couples with

females and males, all chemical classes had the median correlations below 0.25 and urine

metals were the class with greatest percentage drop (83%). We also observed a substantial

reduction in the IQRs of the chemical classes. For example, the IQR of polybrominated

compounds was 0.21, which corresponds to an over 35% drop relatively to females and males.

We found that urine metals had the largest drop in IQR (over 77%). Looking more closely to the

data, correlations between the same chemicals in couples (i.e. the diagonal of the correlation

matrix of couples) were generally higher than that among the same class members. Because of

the low number of exposure indicators in blood metals (3 chemicals), paracetamols (2

chemicals), and cotinine (1 chemical), the diminishing effect to within-class correlations was

countered and thus the drop in medians and IQRs of these classes were not as prominent as

others.

P values obtained from Bonferroni correction with M_{eff} were highly similar between females

and males (Table 3). Both groups had similar M_{eff} values with a difference smaller than 1

variable, suggesting that the overall degree of correlations was also similar. After adjusting for

the correlation, we found the total number of variables decreased from 128 to 112 for females

and 113 for males.

We estimated the concordances between correlations in the LIFE study (as females, males and

couples) and the 2003-2004 NHANES in Table 4. Overall, correlations *r* varied greatly between

chemical groups, from −0.78 to 0.98. Certain chemical groups had small sample size ($n \leq 6$) and

could cause low and inverse correlations. However, correlations *r* were more consistent and

comparable when we discarded group information and considered all chemicals as a whole

(females: 0.88; males: 0.84; couples: 0.67); therefore, we conclude the exposure correlation

patterns captured in LIFE are comparable to that in the U.S. population.

Discussion

Findings

Understanding the co-exposure patterns is an important step toward investigating the joint

health effects of chemical mixtures and for statistical design of exposome-related

investigations. We describe our high level findings here. First, exposure levels of one individual

in the household were not correlated with another individual in the same household. Second,

the percentages of significant r_s ($q < 0.05$) in males and females were 25.3 and 23.1

respectively, in contrast to only 9.5% between couples in the same household (Figure 4).

Chemical correlations in a household setting were concordant to those in 03–04 NHANES,

indicating reproducible co-exposure correlations with respect to the patterns sought in a

generalized and non-institutionalized U.S. population.

Although couples in our study potentially shared a large degree of dietary and indoor

environmental factors, their exposures were only modestly correlated (low r_s). We believe that

there are two additional factors affecting the familial co-exposure patterns in our investigation.

1) The first one is concerned with how long the couples have been living together. In the U.S.,

the median age of first marriage is over 24 since 2000 (U.S. Census Bureau 2016), and newly

married couples could show a greater discordance in chemical co-exposure relationships due to

a different pre-marriage exposures. 2) The second factor is potential physiological dampening

of exposure variability related to the half-life of the target chemicals (Makey et al. 2014;

Rappaport and Kupper 2008). Polybrominated compounds, PCBs, and OCPs are persistent

chemicals with high lipophilicity and longer half-lives (on the order of years). Their serum

concentrations are integrated over a period of time and not completely associated with recent

exposure (Aylward et al. 2014). Since most of the couples recruited in LIFE Study were living

together, we claim this phenomenon can explain the drop in r_s of persistent chemicals relative

to the short-lived urinary chemicals (such as phthalate metabolites, whose half-lives are on the

order of days) in couples (Figure 4).

Wu et al. (2015) conducted a household-based study and measured serum PBDE levels in

children and parents and reported the r_s between child and parent were in the range of 0.66 to

0.74 (median = 0.68, n = 68) for a number of PBDE compounds. The pairs shared a substantial

portion of genes, diet, and living environment and they found that for the latter 2 components,

measured as floor-wipe PBDEs, canned meat, tuna and whitefish, were predictive of serum

PBDEs. Furthermore, they found higher r_s of PBDEs between older couples (age ≥ 55, range =

0.45 to 0.78, median = 0.72). These findings are consistent with our claim.

Persistent organic chemicals are known to cause adverse health effects and are prioritized by

the U.S. Centers for Disease Control and Prevention for health monitoring (Li et al. 2006). Many

of them had wide applications in electrical/electronic equipment, agricultural chemicals, and

furniture (Dodson et al. 2012; Whitehead et al. 2011). Although other emerging EDCs such as

BPA and phthalates have short half-lives, they have extensive modern applications in cosmetics

and consumer products (Bloom et al. 2015b; Buck Louis et al. 2014; Louis et al. 2014; Smarr et

al. 2017). The ubiquity of these chemicals in different microenvironments such as schools and

offices suggests that household environment alone is not a major contributor to body burden,

consistent with our results (Figure 2). PFASs and blood metals had higher variance explained by

the shared environment than other chemical classes and we believe that this could be related

to the lifestyle of the subjects in the LIFE cohort (e.g. PFAS exposures via food items and

personal care products).

Strength and Limitations

Our investigation includes 128 chemical biomarkers with diverse physicochemical properties

that span from persistent lipophilic to non-persistent hydrophilic chemicals and is one of the

first attempts to systematically characterize their correlations in a household setting. However,

we do not know how long the couples have been living together at the baseline of the study to

quantitatively assess how this factor affects co-exposure patterns. Also, we only collected

biological samples at the baseline; thus, it is not possible to study how exposure levels and co-

exposure patterns change longitudinally with time, which could be an important piece of

information for assessing fecundity outcome and chemical exposures.

Analytical and Sampling Implications for Exposome-Wide Investigations

Our findings have implications for high-throughput association tests between correlated

exposures and health outcomes and phenotypes. One of the typical approaches adjusting for

multiplicity in EWASs is controlling the family-wise error rate (e.g. Bonferroni correction).

However, the tests are assumed to be independent. Nyholt (2004) provided one solution to

address correlation in Bonferroni correction by calculating the effective number of variables

(M_{eff}, formula 1), which relies on identifying the number of variables, M, and estimating the

eigenvalue variance, $Var(\lambda_{obs})$ of the correlation matrix. For example, conducting an EWAS of a

response with respect to the chemical class PCBs containing 36 congeners, one can calculate

the correlation matrix (M = 36) and estimate the associated $Var(\lambda_{obs})$ to obtain M_{eff} (Table 3). A

new significance level for this set of comparisons will be α/Meff, which is less stringent than

Bonferroni correction because Meff wil be smaller than M if correlations exist between

congeners. Benjamini and Yekutieli (2001) and Fan et al. (2012) documented a procedure that

considers the correlation structure for better controlling the FDR.

In the future, exposome-related investigations, will include many more variables than ever

before with the emergence of highly sensitive high-resolution mass spectrometry techniques

(Jones 2016) and availability of large data sets from reproducibility initiatives (Manrai et al.

2017). Attempts to assess health outcome associated with a number of exposures may increase

R^2 and lead to "overfitting" (Hawkins 2004). Additionally, dense correlations among exposures

indicate that multicollinearity may also influence the reliability of the association size in

multiple regression or even potential confounding. We claim that these analytic challenges

could be ameliorated through understanding co-exposure patterns. For example, statistical

approaches for evaluating the effects of mixture exposures such as principal component

analysis generally involve dimensionality reduction that relies on estimating the correlation

structure to reduce the number of exposures being considered prior to analysis but are difficult

to interpret (Taylor et al. 2016). Other regression approaches, such as the Elastic Net (Zou and

Hastie 2005) can consider highly co-linear exposure variables while giving coefficients that are

similar to that delivered from a typical regression model; however, inferential estimates (e.g. p

values and confidence intervals) are still in development (Taylor and Tibshirani 2015).

Capturing population-level exposure variability — and the demographic variables that are

associated with the variation, such as sex, location, and time — is a grand ambition in the

exposome concept (Manrai et al. 2017). Given people spend over 90% of their time indoor and

more than 12 hours a day at home (bls.gov/tus), household samples (e.g. such as house dust)

might be a reasonable surrogate to represent home exposures to family members who share

the same living environment. While sampling the household is a tempting approach because of

the simplicity and cost-savings over personal measurement, it may fail to catch a significant

fraction of exposure variability in the population as we found that shared environment could

explain a small percentage of biomarker variance (Figure 2). For epidemiological investigation

that sample study participants who are nested in a unit, for example, schools or homes,

conducting preliminary measurements to assess the influence of shared environment is one of

the ways to justify unit-based measurement.

Finally, correlations of chemical mixtures can also be an important tool used in exposure

science and cumulative risk assessment as groups of correlated chemicals are often released

from a single source (e.g. power plant and vehicular exhaust (Ravindra et al. 2008). Thus,

studying the co-exposure patterns is the first step to identify the sources prior to more in-depth

source apportionment methods such as positive matrix factorization and chemical mass

balance receptor models (Rizzo and Scheff 2007). In cumulative risk assessment, dose addition

is one of the common approaches to estimate the risks from mixture exposures by assuming a

shared toxicity mechanism between chemicals (e.g. binding to the same receptor) (Chen et al.

2001). Knowing the co-exposure correlations could be a first step toward identification of

exposures with similar physicochemical properties to guide follow-up investigations.

Conclusions

While we observed similar co-exposure patterns between females and males, the correlations

were much lower in couples. Our analyses empirically demonstrate that shared environment

explains less than 20% of the biomarker variance in 11 out of 13 EDC classes. The influence of

shared environment to EDC levels is likely conditional on 1) the duration of residence of the

subjects and 2) the lipophilicity and persistency of the chemicals. These factors should be

considered when using surrogate measurement to assess the exposures of family members.

References

Akins JR, Waldrep K, Bernert JT. 1989. The estimation of total serum lipids by a completely

enzymatic "summation" method. Clin. Chim. Acta 184: 219–226.

Asimakopoulos AG, Wang L, Thomaidis NS, Kannan K. 2014. A multi-class bioanalytical

methodology for the determination of bisphenol A diglycidyl ethers, p-hydroxybenzoic acid

esters, benzophenone-type ultraviolet filters, triclosan, and triclocarban in human urine by

liquid chromatography-tandem mass spectrometry. J. Chromatogr. A 1324: 141–148.

Aylward LL, Hays SM, Smolders R, Koch HM, Cocker J, Jones K, et al. 2014. Sources of variability

in biomarker concentrations. J. Toxicol. Environ. Health B Crit. Rev. 17: 45–61.

Benjamini Y, Yekutieli D. 2001. The Control of the False Discovery Rate in Multiple Testing under

Dependency. Ann. Stat. 29: 1165–1188.

Bernert JT Jr, Turner WE, Pirkle JL, Sosnoff CS, Akins JR, Waldrep MK, et al. 1997. Development

and validation of sensitive method for determination of serum cotinine in smokers and

nonsmokers by liquid chromatography/atmospheric pressure ionization tandem mass

spectrometry. Clin. Chem. 43: 2281–2291.

Bloom MS, Buck Louis GM, Sundaram R, Maisog JM, Steuerwald AJ, Parsons PJ. 2015a. Birth

outcomes and background exposures to select elements, the Longitudinal Investigation of

Fertility and the Environment (LIFE). Environ. Res. 138: 118–129.

Bloom MS, Whitcomb BW, Chen Z, Ye A, Kannan K, Buck Louis GM. 2015b. Associations

between urinary phthalate concentrations and semen quality parameters in a general

population. Hum. Reprod. 30: 2645–2657.

Buck Louis GM, Barr DB, Kannan K, Chen Z, Kim S, Sundaram R. 2016. Paternal exposures to

environmental chemicals and time-to-pregnancy: overview of results from the LIFE study.

Andrology 4: 639–647.

Buck Louis GM, Schisterman EF, Sweeney AM, Wilcosky TC, Gore-Langton RE, Lynch CD, et al.

2011. Designing prospective cohort studies for assessing reproductive and developmental

toxicity during sensitive windows of human reproduction and development--the LIFE

Study. Paediatr. Perinat. Epidemiol. 25: 413–424.

Buck Louis GM, Sundaram R, Sweeney AM, Schisterman EF, Maisog J, Kannan K. 2014. Urinary

bisphenol A, phthalates, and couple fecundity: the Longitudinal Investigation of Fertility

and the Environment (LIFE) Study. Fertil. Steril. 101: 1359–1366.

Champely S. 2017. pwr: Basic Functions for Power Analysis. Available: https://CRAN.R-

project.org/package=pwr.

Chen JJ, Chen YJ, Rice G, Teuschler LK, Hamernik K, Protzel A, et al. 2001. Using dose addition to

estimate cumulative risks from exposures to multiple chemicals. Regul. Toxicol. Pharmacol.

34: 35–41.

Dodson RE, Perovich LJ, Covaci A, Van den Eede N, Ionas AC, Dirtu AC, et al. 2012. After the

PBDE phase-out: a broad suite of flame retardants in repeat house dust samples from

California. Environ. Sci. Technol. 46: 13056–13066.

Fan J, Han X, Gu W. 2012. Estimating False Discovery Proportion Under Arbitrary Covariance

Dependence. J. Am. Stat. Assoc. 107: 1019–1035.

Guo Y, Alomirah H, Cho H-S, Minh TB, Mohd MA, Nakata H, et al. 2011. Occurrence of phthalate

metabolites in human urine from several Asian countries. Environ. Sci. Technol. 45: 3138–

3144.

Hauser R. 2006. The environment and male fertility: recent research on emerging chemicals and

semen quality. Semin. Reprod. Med. 24: 156–167.

Hawkins DM. 2004. The problem of overfitting. J. Chem. Inf. Comput. Sci. 44: 1–12.

Heavner DL, Morgan WT, Sears SB, Richardson JD, Byrd GD, Ogden MW. 2006. Effect of

473  creatinine and specific gravity normalization techniques on xenobiotic biomarkers in

474  smokers' spot and 24-h urines. J. Pharm. Biomed. Anal. 40: 928–942.

475 Ioannidis JPA. 2016. Exposure-wide epidemiology: revisiting Bradford Hill. Stat. Med. 35: 1749–

476  1762.

477 Ioannidis JPA, Loy EY, Poulton R, Chia KS. 2009. Researching genetic versus nongenetic

478  determinants of disease: a comparison and proposed unification. Sci. Transl. Med. 1: 7ps8–

479  7ps8.

480 Jones DP. 2016. Sequencing the exposome: A call to action. Toxicol Rep 3: 29–45.

481 Kuklenyik Z, Needham LL, Calafat AM. 2005. Measurement of 18 perfluorinated organic acids

482  and amides in human serum using on-line solid-phase extraction. Anal. Chem. 77: 6085–

483  6091.

484 Li QQ, Loganath A, Chong YS, Tan J, Obbard JP. 2006. Persistent organic pollutants and adverse

485  health effects in humans. J. Toxicol. Environ. Health A 69: 1987–2005.

486 Louis GMB, Kannan K, Sapra KJ, Maisog J, Sundaram R. 2014. Urinary concentrations of

487  benzophenone-type ultraviolet radiation filters and couples' fecundity. Am. J. Epidemiol.

488  180: 1168–1175.

489 Louis GMB, Sundaram R, Schisterman EF, Sweeney AM, Lynch CD, Gore-Langton RE, et al. 2013.

490  Persistent environmental pollutants and couple fecundity: the LIFE study. Environ. Health

491  Perspect. 121: 231.

492 Makey CM, McClean MD, Sjödin A, Weinberg J, Carignan CC, Webster TF. 2014. Temporal

493  variability of polybrominated diphenyl ether (PBDE) serum concentrations over one year.

494  Environ. Sci. Technol. 48: 14642–14649.

495 Manrai AK, Cui Y, Bushel PR, Hall M, Karakitsios S, Mattingly CJ, et al. 2017. Informatics and

496  Data Analytics to Support Exposome-Based Discovery for Public Health. Annu. Rev. Public

Health 38: 279–294.

Mumford SL, Kim S, Chen Z, Boyd Barr D, Buck Louis GM. 2015. Urinary Phytoestrogens Are
Associated with Subtle Indicators of Semen Quality among Male Partners of Couples
Desiring Pregnancy. J. Nutr. 145: 2535–2541.

Nyholt DR. 2004. A simple correction for multiple testing for single-nucleotide polymorphisms
in linkage disequilibrium with each other. Am. J. Hum. Genet. 74: 765–769.

Patel CJ, Ioannidis JPA. 2014. Placing epidemiological results in the context of multiplicity and
typical correlations of exposures. J. Epidemiol. Community Health 68: 1096–1100.

Patel CJ, Manrai AK. 2015. Development of exposome correlation globes to map out
environment-wide associations. Pac. Symp. Biocomput. 231–242.

Patel CJ, Sundaram R, Buck Louis GM. 2016. A data-driven search for semen-related phenotypes
in conception delay. Andrology.

Phillips DL, Pirkle JL, Burse VW, Bernert JT Jr, Henderson LO, Needham LL. 1989. Chlorinated
hydrocarbon levels in human serum: effects of fasting and feeding. Arch. Environ. Contam.
Toxicol. 18: 495–500.

Potera C. 2014. The HELIX Project: tracking the exposome in real time. Environ. Health Perspect.
122: A169.

Rappaport SM, Kupper LL. 2008. *Quantitative exposure assessment*. S. Rappaport.

Ravindra K, Sokhi R, Van Grieken R. 2008. Atmospheric polycyclic aromatic hydrocarbons:
Source attribution, emission factors and regulation. Atmos. Environ. 42: 2895–2921.

R Core Team. 2016. R: A Language and Environment for Statistical Computing.

Richardson DB, Ciampi A. 2003. Effects of exposure measurement error when an exposure
variable is constrained by a lower limit. Am. J. Epidemiol. 157: 355–363.

Rizzo MJ, Scheff PA. 2007. Utilizing the Chemical Mass Balance and Positive Matrix Factorization
models to determine influential species and examine possible rotations in receptor
modeling results. Atmos. Environ. 41: 6986–6998.

Schafer JL. 1999. Multiple imputation: a primer. Stat. Methods Med. Res. 8: 3–15.

Schisterman EF, Vexler A, Whitcomb BW, Liu A. 2006. The limitations due to exposure detection
limits for regression models. Am. J. Epidemiol. 163: 374–383.

Schisterman EF, Whitcomb BW, Louis GMB, Louis TA. 2005. Lipid adjustment in the analysis of
environmental contaminants and human health risks. Environ. Health Perspect. 113: 853–
857.

Sjödin A, Jones RS, Lapeza CR, Focant J-F, McGahee EE 3rd, Patterson DG Jr. 2004.
Semiautomated high-throughput extraction and cleanup method for the measurement of
polybrominated diphenyl ethers, polybrominated biphenyls, and polychlorinated biphenyls
in human serum. Anal. Chem. 76: 1921–1927.

Smarr MM, Grantz KL, Sundaram R, Maisog JM, Honda M, Kannan K, et al. 2016. Urinary
paracetamol and time-to-pregnancy. Hum. Reprod. 31: 2119–2127.

Smarr MM, Sundaram R, Honda M, Kannan K, Louis GMB. 2017. Urinary Concentrations of
Parabens and Other Antimicrobial Chemicals and Their Association with Couples'
Fecundity. Environ. Health Perspect. 125: 730–736.

Stingone JA, Buck Louis GM, Nakayama SF, Vermeulen RCH, Kwok RK, Cui Y, et al. 2017. Toward
Greater Implementation of the Exposome Research Paradigm within Environmental
Epidemiology. Annu. Rev. Public Health 38: 315–327.

Taylor J, Tibshirani RJ. 2015. Statistical learning and selective inference. Proc. Natl. Acad. Sci. U.
S. A. 112: 7629–7634.

Taylor KW, Joubert BR, Braun JM, Dilworth C, Gennings C, Hauser R, et al. 2016. Statistical

Approaches for Assessing Health Effects of Environmental Chemical Mixtures in

Epidemiology: Lessons from an Innovative Workshop. Environ. Health Perspect. 124:

A227–A229.

te Velde E, Burdorf A, Nieschlag E, Eijkemans R, Kremer JAM, Roeleveld N, et al. 2010. Is human

fecundity declining in Western countries? Hum. Reprod. 25: 1348–1353.

U.S. Census Bureau. 2016. Current Population Survey: Annual Social and Economic Supplement.

Vineis P, Chadeau-Hyam M, Gmuender H, Gulliver J, Herceg Z, Kleinjans J, et al. 2016. The

exposome in practice: Design of the EXPOsOMICS project. Int. J. Hyg. Environ. Health;

doi:10.1016/j.ijheh.2016.08.001.

Vrijheid M, Slama R, Robinson O, Chatzi L, Coen M, van den Hazel P, et al. 2014. The human

early-life exposome (HELIX): project rationale and design. Environ. Health Perspect. 122:

535–544.

Whitehead T, Metayer C, Buffler P, Rappaport SM. 2011. Estimating exposures to indoor

contaminants using residential dust. J. Expo. Sci. Environ. Epidemiol. 21: 549–564.

Wild CP. 2005. Complementing the Genome with an "Exposome": The Outstanding Challenge of

Environmental Exposure Measurement in Molecular Epidemiology. Cancer Epidemiol.

Biomarkers Prev. 14: 1847–1850.

Wild CP, Scalbert A, Herceg Z. 2013. Measuring the exposome: a powerful basis for evaluating

environmental exposures and cancer risk. Environ. Mol. Mutagen. 54: 480–499.

Wu XM, Bennett DH, Moran RE, Sjödin A, Jones RS, Tancredi DJ, et al. 2015. Polybrominated

diphenyl ether serum concentrations in a Californian population of children, their parents,

and older adults: an exposure assessment study. Environ. Health 14: 23.

Zou H, Hastie T. 2005. Regularization and variable selection via the elastic net. J. R. Stat. Soc.

Series B Stat. Methodol. 67: 301–320.

Tables

569 Table 1. List of chemical classes and measured chemicals in the current study.

EDC Class	No.	Individual Chemicals	Medium	LOQ
Polychlorinated biphenyls (PCBs)	36	Congeners: 28, 44, 49, 52, 66, 74, 87, 99, 101, 105, 110, 114, 118, 128, 138, 146, 149, 151, 153, 156, 157, 167, 170, 172, 177, 178, 180, 183, 187, 189, 194, 195, 196, 201, 206, 209	Serum	1-3 pg/g, wet weight
Organochlorine pesticides (OCPs)	9	Hexachlorobenzene (HCB), β-hexachlorocyclohexane (β-HCH), γ-hexachlorocyclohexane (γ-HCH), oxychlordane, *trans*-nonachlor, *p,p*'-DDT, *o,p*'-DDT, *p,p*'-DDE, and mirex	Serum	1-3 pg/g, wet weight
Polybrominated chemicals[a]	11	Brominated biphenyl (BB 153); brominated diphenyl ethers (BDEs) congeners: 17, 28, 47, 66, 85, 99, 100, 153, 154, 183	Serum	5 pg/g, wet weight
Polyfluoroalkyl substances (PFASs)	7	2-(*N*-ethyl-perfluorooctane sulfonamido) acetate (Et-PFOSA-AcOH), 2-(*N*-methyl-perfluorooctane sulfonamido) acetate (Me-PFOSA-AcOH), perfluorodecanoate (PFDeA), perfluorononanoate (PFNA), perfluorooctane sulfonamide (PFOSA), perfluorooctane sulfonate (PFOS), and perfluorooctanoate (PFOA)	Serum	0.04-0.1 ng/mL, wet weight
Phytoestrogens	6	Genistein, daidzein, *O*-desmethylangolensin, equol, enterodiol, and enterolactone	Urine	0.2-0.6 ng/mL
Phthalate metabolites	14	Mono (3-carboxypropyl) phthalate (mCPP), monomethyl phthalate (mMP), monoethyl phthalate (mEP), mono (2-isobutyl phthalate) (miBP), mono-n-butyl phthalate (mBP), mono (2-ethyl-5-carboxyphentyl) phthalate (mECPP), mono-[(2-carboxymethyl) hexyl] phthalate (mCMHP), mono (2-ethyl-5-oxohexyl) phthalate (mEOHP), mono (2-ethyl-5-hydroxyhexyl) phthalate (mEHHP), monocyclohexyl phthalate (mCHP), monobenzyl phthalate (mBzP), mono (2-ethylhexyl) phthalate (mEHP), mono-isononyl phthalate (mNP), and monooctyl phthalate (mOP)	Urine	0.2-2 ng/mL
Phenols[b]	6	Total bisphenol A (BPA); benzophenones (BPs): 4-hydroxybenzophenone (4-OH-BP), 2,4-dihydroxybenzophenone (2,4-OH-BP), 2,2′,4,4′-tetrahydroxybenzophenone (2,2′4,4′-OH-BP), 2-hydroxy-4-methoxybenzophenone (2-OH-4-MeO-BP), and 2,2′-dihydroxy-4-methoxybenzophenone (2,2′-OH-4-MeO-BP)	Urine	0.02-0.05 ng/mL
Antimicrobial chemicals[c]	12	Triclosan (TCS) and triclocarban (TCC); parabens: methyl paraben (MP), ethyl paraben (EP), propyl paraben (PP), butyl paraben (BP), benzyl paraben (BzP), heptyl paraben (HP), 4-hydroxy benzoic acid (4-HB), 3,4-dihydroxy benzoic (3,4-DHB), methyl-protocatechuic acid (OH-Me-P), and ethyl-protocatechuic acid (OH-Et-P)	Urine	0.02-0.05 ng/mL
Paracetamol & derivatives	2	Paracetamol and 4-aminophenol	Urine	0.5 ng/mL and 0.25 ng/mL
Blood metals	3	Cadmium (Cd), lead (Pb), and mercury (Hg)		10 ng/L to 10 µg/L
Cotinine[d]	1	Cotinine	Serum	0.01 ng/mL
Urine metals	17	Manganese (Mn), chromium (Cr), beryllium (Be), cobalt (Co), molybdenum (Mo), cadmium (Cd), tin (Sn), caesium (Cs), barium (Ba), nickel (Ni), copper (Cu), zinc (Zn), tungsten (W), platinum (Pt), thallium (Tl), lead (Pb), and uranium (U)	Urine	10 ng/L to 10 µg/L
Urine metalloids	4	Selenium (Se), arsenic (As), antimony (Sb), and tellurium (Te)	Urine	10 ng/L to 10 µg/L

571 Note: LOQ, Limits of quantification.

572   [a]Polybrominated chemicals contain mostly PBDEs with one PBB.
573   [b]Phenols contain mostly benzophenones with one BPA.
574   [c]Antimicrobial chemicals contain mostly parabens with TCS and TCC.
575   [d]Serum cotinine is not an EDC but included for comprehensive investigation.
576
577

578   Table 2. Sociodemographic and lifestyle characteristics of females and males in the LIFE Study.

Characteristics	Females (n = 501)	Males (n = 501)
Age (year) **	29.99 ± 4.14	31.77 ± 4.92
BMI (kg/m^2) **		
< 25	230 (46)	84 (17)
≥ 25 & < 30	136 (27)	206 (41)
≥ 30	135 (27)	311 (62)
Non-Hispanic White	396 (79)	397 (79)
College graduate or higher *	474 (95)	457 (91)
Yearly income $80,000 or over	297 (59)	298 (59)
Regular vigorous exercise in the past 12 months	200 (40)	211 (42)
Smoke at the time of study		
No	445 (89)	440 (88)
Yes (No. of cigarettes on a typical day)		
1 - 3	19 (4)	26 (5)
4 - 6	8 (2)	11 (2)
7 - 10	15 (3)	8 (2)
11-15	7 (1)	2 (0)
16 - 25	3 (1)	10 (2)
> 25	4 (1)	4 (1)
≥ 12 alcoholic drinks in the past 12 months **	374 (75)	428 (85)
No. of alcoholic drinks on a typical occasion **		
0	128 (26)	73 (15)
1	108 (22)	63 (13)
2	169 (34)	150 (30)
3	68 (14)	99 (20)
4	19 (4)	62 (12)
5	9 (2)	54 (11)
History of diabetes	6 (1)	14 (3)
History of high blood pressure **	20 (4)	52 (10)
History of high cholesterol **	41 (8)	78 (16)
Serum cotinine (ng/mL)[a] **	0.62 ± 0.23	1.24 ± 2.17
Serum total lipids (mg/dL)[a] **	2.00 ± 0.03	6.56 ± 0.26
Urinary creatinine (mg/dL)[a] **	4.22 ± 0.86	4.76 ± 0.73

580   Note: BMI, body mass index. Values in mean ± SD or n (%).

581   *p < 0.05; **p < 0.01.

582   [a]log+1 transformed values.

586   Table 3. Number of effective variables for each chemical class after adjusting for the within-class correlation.

| Chemical class | M | Females | | | Males | | |
		M_{eff}	M_{diff}	*P* value	M_{eff}	M_{diff}	*P* value
Serum persistent organic compounds							
Polychlorinated biphenyls (PCBs)	36	28.38	7.62	0.002	28.69	7.31	0.002
Organochlorine pesticides (OCPs)	9	7.94	1.06	0.006	8.17	0.83	0.006
Polybrominated chemicals	11	8.22	2.78	0.006	8.38	2.62	0.006
Polyfluoroalkyl substances (PFASs)	7	6.13	0.87	0.008	6.25	0.75	0.008
Urinary non-persistent organic compounds							
Phytoestrogens	6	5.34	0.66	0.009	5.38	0.62	0.009
Phthalate metabolites	14	12.72	1.28	0.004	12.87	1.13	0.004
Phenols	6	5.48	0.52	0.009	5.51	0.49	0.009
Antimicrobial chemicals	12	11.38	0.62	0.004	11.60	0.40	0.004
Paracetamol & derivatives	2	1.99	0.01	0.025	2.00	0.00	0.025
Others							
Blood metals	3	2.91	0.09	0.017	2.91	0.09	0.017
Serum cotinine	1	1.00	0.00	0.050	1.00	0.00	0.050
Urine metals	17	16.13	0.87	0.003	16.22	0.78	0.003
Urine metalloids	4	3.95	0.05	0.013	3.95	0.05	0.013
Total	128	111.57	16.43	0.0004	112.94	15.06	0.0004

588   Note: M, number of variables in the corresponding chemical classes; M_{eff}, number of effective variables after taking account of
589   within-class correlation; M_{diff}, the different between M and M_{eff}; *P* value, Bonferroni adjusted *p* values as 0.05/M_{eff}.

593 Table 4. Correlations of chemical relatedness between 2003–2004 NHANES and current study by different chemical classes.

Chemical class[a]	Females		Males		Couples	
	Pearson r	n	Pearson r	n	Pearson r	n
Blood_metals	0.07	3	0.13	3	-0.04	6
OCPs	0.69	36	0.51	36	0.25	72
PCBs	0.88	561	0.88	561	0.77	1122
PFASs	0.31	21	0.29	21	0.32	42
Phthalates	0.90	78	0.89	78	0.34	156
Phytoestogens	0.98	15	0.97	15	0.86	30
Polybrominated_cpds	0.76	55	0.74	55	0.77	110
Urine_metalloids	0.41	3	-0.78	3	0.34	6
Urine_metals	0.82	55	0.78	55	-0.01	110
Total	0.84	827	0.84	827	0.67	1654

Note: Pearson r, Pearson correlation coefficients of the chemical relatedness (spearman correlations of chemicals) between National

Health and Nutrition Examination Survey (NHANES) and this study; n, sample size.

[a]Only 9 classes are shown because not all chemicals in 13 chemical classes in this study could be matched with NHANES.

Figure 1. Analytical scheme to investigate the variability and correlations in this study. A) We first extract the residuals from a linear
model after adjusting for the base covariates (total lipids or creatinine) to calculate the effective number of variables (M_{eff}) and
Spearman's rank correlation (r_s). B) Then, we used another linear model with an additional age variable to obtain residuals and
conducted paired t-test to test the difference of biomarkers between females and males living in the same household. C) Afterward,
we further adjusted for sex prior to extracting residuals to calculate the percentage of biomarker variance explained by the shared
environment.
606
607   Figure 2. Summary of the percentage variance explained by the shared environment. Boxplots of the adjusted coefficient of
608   determination (R^2) within different chemical classes are shown. Interquartile range is not shown for Cotinine class because it
609   contains only 1 compound. For each box, median and interquartile range are drawn and the whiskers are extended to the largest
610   values within 1.5*interquartile range. Black dots denote correlations outside of the range covered by the whiskers.
611
Figure 3. Exposome correlation globe showing the relationships between females, males and couples. Right-half represents
biomarkers in females; left-half represents biomarkers in males. Only Spearman's rank correlations greater than 0.25 and smaller
than −0.25 were shown as connections in the globe. Red line denotes positive correlation and dark green line denotes negative one.
Color intensity and line width are proportional to the size of the correlation. Within-class and between-class correlations are shown
outside and inside of the track respectively. Correlations in couples are indicated by the lines linking females and males (i.e. crossing
the vertical-half of the globe).
618
Figure 4. Boxplots of Spearman's rank correlations (r_s) within different chemical classes. A) Females; B) Males; and C) Couples. For
couples, summary statistics were estimated with the full 128 x 128 correlation matrix instead of with the half triangle. Certain
classes contain only 1 pair of correlation (paracetamols in females, paracetamols in males, and cotinine in couples). "All" represents
the grouping by the correlation of all pairs of chemicals available. Horizontal line drawn across the chemical classes is equal to the
95[th] percentile of the null distribution obtained from permuting the concentrations of all chemicals. Definition of whisker and black
dot can be referred to the caption in Figure 2.
625

626
627   Figure 1
628

629
630   Figure 2
631

632
633   Figure 3
634

637
638

639
640   Figure 4
641