Novel approach to quantitative spatial gene expression uncovers genetic stochasticity in the developing

Drosophila eye

Sammi Ali[1,¶], Sarah A. Signor[1,¶,*], Konstantin Kozlov[2], Sergey V. Nuzhdin[1,2]

[1]Molecular and Computational Biology, University of Southern California, Los Angeles, California, 90089 USA

[2]Department of Applied Mathematics, St. Petersburg State Polytechnic University, St. Petersburg, 195251,

Russia

[¶]Co-first author

[*]Corresponding author

Email: ssignor@usc.edu

Abstract

Robustness in development allows for the accumulation of neutral genetically based variation in expression, and here will be termed 'genetic stochasticity'. This largely neutral variation is potentially important for both evolution and complex disease phenotypes. However, it has generally only been investigated as variation exhibited in the response to large genetic perturbations. In addition, work on variation in gene expression has similarly generally been limited to being spatial, or quantitative, but because of technical restrictions not both. Here we bridge these gaps by investigating replicated quantitative spatial gene expression using rigorous statistical models, in different genotypes, sexes, and species (*Drosophila melanogaster* and *D. simulans*). Using this type of quantitative approach with developmental data allows for effective comparison among conditions, including health versus disease. We apply this approach to the morphogenetic furrow, a wave of differentiation that sweeps across the developing eye disc. Within the morphogenetic furrow, we focus on four conserved morphogens, *hairy*, *atonal*, *hedgehog*, and *Delta*. Hybridization chain reaction quantitatively measures spatial gene expression, co-staining for all four genes simultaneously and with minimal effort. We find considerable variation in the spatial expression pattern of these genes in the eye between species, genotypes, and sexes. We also find that there has been evolution of the regulatory relationship between these genes. Lastly, we show that the spatial interrelationships of these genes evolved between species in the morphogenetic furrow. This is essentially the first 'population genetics of development' as we are able to evaluate wild type differences in spatial and quantitative gene expression at the level of genotype, species and sex.

Introduction

Natural genetic variation within populations has long been the purview of evolutionary and population geneticists, while developmental biologists focus on the effect of large mutations in otherwise isogenic backgrounds (Paaby and Gibson, 2016). This dearth of work on developmental variation in wildtype genetic backgrounds is in part because developmental approaches have long been restricted to data that is semi-quantitative (i.e. *in situ* hybridization, antibody staining). Indeed, gene expression studies are generally spatial or quantitative, but not both. In addition, the data is generally not interrogated from a quantitative perspective, including replication and rigorous statistical models. Without quantitative replication and statistical tests one cannot effectively compare developmental processes among conditions, including health versus disease. This is especially important given the potential for complex interactions between conditionally neutral differences in gene expression to result in disease phenotypes. Here we use hybridization chain reaction (HCR) to bridge this gap between developmental and quantitative or population genetics by quantitatively measuring spatial gene expression in multiple genotypes from two sexes of two species (*Drosophila melanogaster* and *D. simulans*) (Choi et al., 2014). Furthermore, we introduce rigorous replication allowing for statistical hypothesis testing. This is essentially the first 'population genetics of development' as we are able to evaluate wild type differences in spatial and quantitative gene expression at the level of genotype, species and sex. Multiplexing of four genes simultaneously also allows more rigorous analysis of gene co-expression, compared to other techniques that require inference across samples that have been individually stained. We use this enormous developmental dataset to focus on the well-known morphogens driving ommatidia specification in *Drosophila* (Atkins et al., 2013; W. Li et al., 1995; Raj et al., 2008; Shah et al., 2016; Tsachaki and Sprecher, 2011).

The *Drosophila* eye is formed from an imaginal disc, which is initially patterned by a wave of differentiation marked by a visible indentation of the tissue, termed the morphogenetic furrow (MF). The MF passes from the posterior to the anterior of the disc over a period of two days (90 minutes per adjacent row), giving each disc an element of both time and space in development (Fig 1) (Roignant and Treisman, 2009). A strength of the eye disc as a model is that within the MF all cells are arrested at G1, and there will be no additional variability introduced due to differences between cells in their stage of cell division (Baker and Yu,

2001; Escudero and M. Freeman, 2007; Firth and Baker, 2005; Firth et al., 2010). Furthermore, while there is

evidence that transcription is bursty, the eye disc is composed of a repeated pattern of cells that will become

ommatidia, akin to natural replication (Bothma et al., 2014; Fukaya et al., 2016; Tantale et al., 2016). By

averaging across it, we reduce the impact of differences in transcriptional bursting across the eye disc.

The furrow is initiated by *hedgehog,* which both represses (short range) and activates (long range) *hairy*

(Fig 1) (Felsenfeld and Kennison, 1995; Strutt and Mlodzik, 1997). *hairy* represses *atonal*, preventing

precocious neural development anterior to the MF (though this role has been recently contested) (Bhattacharya

and Baker, 2012; N. L. Brown et al., 1991; 1995). *hedgehog* activates the expression of *atonal*, driving the MF

anteriorly (Fig 1) (Heberlein et al., 1993; Ma et al., 1993; Greenwood and Struhl, 1999). *atonal* is the proneural

gene in *Drosophila*, establishing the competency to become photoreceptor cells (Jarman et al., 1994). The

relationship between *Delta*/*Notch* and the other members of the pathway is more complex, but it is clear that in

cells posterior to the furrow *Delta*/*Notch* repress *atonal* (Fig 1) (Firth and Baker, 2005; Gavish et al., 2016).

There is also some evidence that *Delta*/*Notch* repress negative regulators of *atonal* at the furrow, such as *hairy*

(Bhattacharya and Baker, 2009; N. L. Brown et al., 1995; A. B. A. M. Freeman, 2001). In addition, there is some

evidence suggesting that *Notch*/*Delta* are involved in the early stages of *atonal* induction, and alternatively that

atonal activates its own transcription (Baker and Yu, 1997; Dominguez and Hafen, 1997; Dominguez et al.,

1998; Y. Li and Baker, 2001; Spencer et al., 1998; Sun et al., 1998). There are many other genes involved in the

specification of the eye disc that will not be mentioned here, in favor of focusing on the genes we have assayed.

We analyze the spatial quantitative expression of *hedgehog*, *hairy*, *atonal*, and *Delta* to understand the evolving

regulatory logic of the gene network and changes in spatial dynamics between sexes and species.

We term differences in gene expression between species, genotypes, and sexes 'genetic stochasticity', as

there are no phenotypic differences between these *Drosophila* eyes other than size and proportion of

photoreceptor subtype. Size is included as a co-factor in the relevant models discussed below, and results only in

a larger area being patterned for ommatidia rather than a difference in pattern. In addition, photoreceptor subtype

is not determined or affected by the genes expressed during the initial patterning phase in the MF (Cook et al.,

2003; Johnston and Desplan, 2014; Johnston et al., 2011; Wernet et al., 2006). We interpret genetic stochasticity

in quantitative spatial patterns of gene expression in light of regulatory relationships among them. We approach

a spatial and quantitative analysis of these gene expression patterns in three ways, first by explicitly creating a

spatial gene expression profile and comparing between genotypes, sexes, and species. Second, we were

interested in examining if the regulatory relationship between these genes had evolved between species or

harbors variation within a species. Lastly, we investigated the possibility that the spatial relationship between

these genes relative to the MF had evolved or harbors variation within populations.

Methods

Fly stocks

109 *D. simulans* were collected from the Zuma organic orchard in Zuma beach, CA in the spring of 2012 (Signor et

110 al., 2017). They were inbred by 15 generations of full sib crosses. *D. melanogaster* were collected in Raleigh,

111 North Carolina and inbred for 20 generations (Mackay et al., 2012).

Staging and dissection of larvae

All flies were reared on a standard medium at 25° C with a 12-h light/12-h dark cycle. 120 hours after hatching,

3^{rd} instar larva were placed in phosphate buffered saline (PBS) and separated by sex. Their guts were carefully

removed posteriorly and their body was inverted anteriorly to expose the brains, eye discs and mouth hooks.

After fixation and labeling (described below), eye discs were isolated and mounted. The authors note that while

there will be variation in the exact row of the eye that is being patterned between images, replicates were not

conducted at the same time nor from the same cross. As such, with up to five replicates per line, any variation in

the exact positioning of the furrow will serve to increase noise within the dataset rather than create false signal.

Hybridization Chain Reaction (HCR)

HCR is unique in that it produces gene expression patterns that are both quantitative and spatial. The DNA

probes were designed and synthesized by Molecular Instruments (Choi et al., 2014) (S1 Table). Four genes were

multiplexed in each preparation as orthogonally-designed hairpins allowed the simultaneous amplification of

their target sequences (Fig 1, S1 Fig). Each target mRNA was detected using five DNA probes to annotate the

position and expression levels for each of the four assayed genes (*hairy, atonal, Delta and hedgehog*). Each

probe contained two-initiator sequences (I1 and I2) that bound to a specific amplifier.

While other approaches such as FISH can be adapted to detect individual transcripts, HCR has a linear signal

that is 20x brighter than FISH, it reduces non-specific background staining, and it can detect 88% of single RNA

molecules in a cell with an appropriately low false discovery rate (Ma and Moses, 1995; Pan and Rubin, 1995).

It is also highly repeatable, with different sets of probes targeted to the same gene showing correlations

of .93-.99 (S. Fraser, pers. comm.).

The protocol for HCR was modified from (Choi et al., 2014) and is described briefly. The full protocol

is available in S1 File. Inverted 3^{rd} instar larva were fixed in 4% paraformaldehyde diluted with PBS

containing .2% Tween 20 (PBST). After fixation, larva were washed with PBST, then increasing concentrations

of methanol (30%, 70% and 100%) at 25° C. Larva were stored in 100% methanol at -20° C. Methanol-fixed

samples were thawed, washed with ethanol, re-permeabalized in 60% xylene, washed with ethanol, then

methanol and rehydrated with PBST. Samples were permeabalized with proteinase K (4 mg/mL), fixed in 4%

formaldehyde then washed with PBST at 25° C. Finally, at 45° C, samples were pre-hybridized for 2 hours

before the addition of all the probes. The probe-hybridized larva were placed in wash buffer (Molecular

Instruments) at 45° C to remove excess probes. Fluorescently labeled hairpins were snap-cooled then added to

the samples at 25° C and placed in the dark to amplify the signal. Afterwards, samples were washed in 5X SSCT

solution, isolated in PBST, then placed in Prolong Gold anti-fade mounting medium (Molecular Probes).

Microscopy

Three dimensional images of mounted, HCR stained 3^{rd} instar larva eye discs were acquired on a Zeiss LSM 780

laser scanning microscope (Carl Zeiss MicroImaging, Inc., Thornwood, NY, USA) with Objective Plan-

Apochromat 63x/1.40 Oil. The gain was adjusted to avoid pixel saturation.

Extraction of gene expression profiles

The first steps in the image analysis is bringing each image to the same orientation and segmenting it. Image

segmentation produces a mask in which pixels are assigned to objects or background. Here the objects are one

or several mRNA molecules. Then the cellular structure of the imaginal disc is approximated using a hexagonal

array. Though the real underlying cell structure of the imaginal disc is technically able to be recognized, this was

unsatisfactory in our data due to imaging noise. Thus, at the second step using the R package hexbin we

constructed a partition of the imaginal disc area into elements that represent pseudo-cells and have a

biologically-relevant hexagonal shape (Brennan et al., 1998). The number of pseudo-cells was selected by visual

inspection of the combined image in which the hexagonal structure was overlaid onto the *atonal* channel to

verify fit. We are primarily interested in expression profiles around the MF, providing us a convenient landmark

to align images from different preparations, thereby assigning coordinates to the pseudo-cells. However,

deformations of the eye disc during growth and preparation sometimes distorts the MF. We used splines to

correct for any bending or deformation of the MF. Next, using the histograms of cumulative pixel intensities of

objects in expression domains and non-expressing areas we estimated the typical intensity of a transcript and

typical background signal, respectively. Consequently, the cumulative intensities greater than the background

are divided by the intensity attributed to single mRNA molecule to yield counts of mRNA molecules. This

normalizes the expression profiles and corrects for differences in microscope gain between images. Finally, the

gene expression profiles are estimated for every pseudo-cell.

Morphological reconstruction and contrast mapping segmentation

To detect gene transcripts within the image stacks we applied a version of the MrComas method that was

modified for processing 3D images (Kozlov et al., 2017). This approach first enhances contrast within the image

and reduces noise. The images were enlarged by a factor of four with the nearest-neighbor algorithm. They were

processed by morphological reconstruction using both opening and closing, where closing (opening) is dilation

(erosion) that removes extraneous dark (bright) spots and connects bright (dark) objects (Vincent, 1993). The

contrast mapping operator assigns each pixel the maximum value between the pixel-by-pixel difference of the

reconstructed images and their pixel-by-pixel product and produces the rough mask for each channel. An image,

I, is mapping from a finite rectangular subset L onto the discrete plane Z^2 into a discrete set 0, 1, …, N - 1 of

gray levels. Let the dilation δ_B and erosion ϵ_B by structural element B be defined as:

$$\delta_B(I) = \bigvee_{\gamma \epsilon_B} I(\gamma) = I \oplus B \qquad \epsilon_B(I) = \bigwedge_{\gamma \epsilon_B} I(\gamma) = I \ominus B \qquad (1)$$

Where ∨ and ∧ denote infimum and supremum respectively. Then formulae:

$$\delta_{I,B}^1(J) = (J \oplus B) \wedge I \qquad \epsilon_{I,B}^1 = (J \ominus B) \vee I \qquad (2)$$

denote geodesic dilation $\delta_{I,B}^1$ and erosion $\epsilon_{I,B}^1$. Binary reconstruction extracts those connected components of the

mask image which are marked on the marker image, and in grayscale it extracts the peaks of the masked image

marked by the marker image. Using the dilated masks image I as the marker $J: J = \delta_B(I)$ defines closing by

reconstruction:

$$\gamma_B(I) = \epsilon^1_{I,B}\epsilon^1_{I,B} \dots \epsilon^1_{I,B}[\delta_B(I)] \tag{3}$$

Opening by reconstructions uses eroded mask I as a marker J:

$$\phi_B(I) = \delta^1_{I,B}\delta^1_{I,B} \dots \delta^1_{I,B}[\epsilon_B(I)] \tag{4}$$

Then the difference between closing and opening by reconstruction has the meaning of the gradient:

$$\nabla_B(I) = \gamma_B(I) - \phi_B(I) \tag{5}$$

To create strong discontinuities at object edges and flatten signal with the objects the contrast mapping operator

takes a maximum between the difference and the pixel-by-pixel produce of the reconstructed images and

produces a rough mask for each channel:

$$R = \max\{\nabla_B(I), \gamma_B(I) \odot \phi_B(I)\} \tag{6}$$

Subsequently, this mask was subjected to distance transform, which substituted each pixel value with the

number of pixels between it and the closest background pixel. This operation creates 'peaks' and 'valleys' of

intensity inside foreground objects. To split erroneously merged objects watershed transform was applied, which

treats the whole image as a surface and intensity of each pixel as its height and determines the watershed lines

along the tops of ridges separating the catchment basins (Meyer, 1994). The quality of segmentation is assessed

visually by inspection of the object boarders overlaid with the original image. Finally, each mask is returned to

its original size and quantitative measures are made of shape and intensity characteristics such as the number of

pixels, as well as their mean and standard deviation in the detected object. MrComas is free and open source

software available at http://sourceforge.net/p/prostack/wiki/mrcomas.

Approximating the MF

We defined the position of the MF as the middle of overlap between *hairy* and *atonal* expression. The shape of

the MF was approximated with a spline using function smooth.spline in R. The degrees of freedom and other

parameters were chosen to make the approximation coincide visually with a MF image.

Inferring counts of transcript number

Segmentation of the image provided a table of coordinates and the shape and intensity characteristics of detected

transcripts. Here, we applied filtering steps to remove false positives and determine the count of mRNA

transcripts. First, we assumed the object we detected as least intense but most frequent corresponds to a single

mRNA molecule. Then, we inferred background intensity for objects outside of well-annotated domains of

expression of the four genes. Assuming that the majority of true objects contain a single molecule, we compare

the distribution of cumulative intensities of particles in expression domains and areas of known non-expression

to obtain the typical intensity of a true single molecule and a false positive, respectively. All detected signals that

were lower than the typical intensity of a false positive were removed from the dataset. The number of removed

objects is typically less then 10%. All other cumulative pixel intensities were divided by the typical intensity of a

true single molecule as normalization coefficient to yield an estimate of the number of mRNA transcripts.

Image registration

We applied an affine coordinate transformation to each eye disc to make the corresponding maxima and the width

of expression patterns of four genes in different eyes coincide as closely as possible. To do so, we shifted the

coordinate system of each eye to its center and also scaled them in the A-P direction. The center of the pattern in

A-P direction is the MF.

220   We mapped the expression patterns to a unified hexagonal structure in order to make comparisons between

pseudo-cells from individual imaginal discs. The unified cell structure was constructed using the R package

hexbin. Each cell in the unified grid represents an 'average' cell from individual eyes. The size of a hexagon in the

unified grid is greater or equal than the cell size in the individual eye. Thus, the number of molecules in each

unified cell in the mapped pattern equals the mean over the cells from native pattern that are covered by this unified

cell. After such coordinate transformation, the MF region is defined as 20 cells on either side of the MF, to focus

the analysis on the area of interest (the MF).

Filtering and quality controls for each eye disc

Some eye imaginal discs were damaged or deformed in the process of dissection or mounting, resulting in regions

of erroneous gene expression, such as disruptions to the MF. The expression profile of each disc was examined by

eye and these regions were individually trimmed out of the final dataset. At the edges of each eye disc the pattern

of the MF was also degraded, so each eye disc was trimmed dorso-ventrally prior to analysis. Outliers were

excluded from the dataset, determined as a single member of the five replicates with more than a 3x difference in

expression values. This resulted in a final dataset of 55 eye discs.

Analysis of individual spatial gene expression patterns

We were primarily interested in variation in gene expression profiles across the eye disc, that is using differences

in expression averaged across rows along the *x*-axis. While the *y*-axis is of interest, variation in the shape, size,

degree of deformation, and occasional damage to the disc made this analysis intractable. We fit curves to each

gene expression profile using the mgcv package in R, using a generalized additive model with integrated

smoothness estimation. Smoothing terms are represented using penalized regression splines. predict.gam was

used to fit the curves to the original range of values and down sample the curves to eight points. MANOVAs

were performed using the "Pillai" test for species x genotype x sex.

Modeling framework to understand variation and evolution of the eye patterning gene network

We wanted to understand if cryptic variation existed within the regulatory logic of *hairy*, *atonal*, *Delta*, and

hedgehog, or if there had been cryptic evolution between species. To understand the regulatory logic between

genes we focused on biologically relevant relationships, such as the regulation of *hairy* by *Delta* and *hedgehog*,

but excluded such relationships as *hairy* and *atonal*. This was due to the low overlap between *hairy* and *atonal*

expression domains, where including cells where only one or another was expressed would artificially create a

relationship between expression levels. Both *hairy* and *atonal* are downstream, directly or indirectly, of *Delta*

and *hedgehog* thus it was these relationships that were modeled. We limited the analysis to cells where all genes

included were expressed in at least ten molecules.

In the previous analysis, we investigated variation in the cryptic spatial quantitative expression pattern of

genes in the MF. Here, we will investigate the possibility that genes in the MF have evolved, or harbor variation,

for how they affect each other in particular cells. For example, is high *atonal* expression associated with high

expression of *hedgehog*, given that *hedgehog* activates *atonal*? We used the following equations to determine the

relationship between the expression of these genes:

$$hairy(i,s) = k^{Dl} \times Dl(i,s) + k^{hh} \times hh(i,s) + \alpha \qquad\qquad (7)$$

$$atonal(i,s) = k^{Dl} \times Dl(i,s) + k^{hh} \times hh(i,s) + \alpha \qquad (8)$$

The coefficient k and constant α were fit using standard methods for multiple regression. Here $hairy(i,s)$

and $atonal(i,s)$ are the measured expression level of each gene in cell i in individual s. $Dl(i,s)$ and $hh(i,s)$ are

vectors containing the corresponding expression levels of *hairy* and *atonal*'s regulators *Delta* and *hedgehog*. To

determine if the regulatory logic is the same between genotypes and species we can then use the regression

coefficients from these models in a MANOVA. We note that we cannot exclude the possibility that other

unmeasured genes are responsible for producing this variation.

Model for understanding overall variation and evolution of MF structure

Lastly, we wanted to understand if there is variation in the relationship between the MF and gene expression, or

if variation existed for the size of the MF overall. For example, the MF was called as the position of overlap

between *atonal* and *hairy* expression, but it is unclear how the overall gene expression pattern of these genes

relates to their overlap. For example, is the position of maximum expression of each always the same relative to

the MF? Two processes occurred to make the MF comparable between samples, they were shifted to occupy the

same position depending upon the position of overlap of *hairy* and *atonal*, and they were scaled to occupy the

same total area. The amount required to scale will depend both on the size of the original disc and the width of

the MF relative to the disc. To account for differences in size we include the number of rows in the original disc

prior to any transformations as a cofactor and perform ANOVA in R.

Results

Individual spatial gene expression patterns

First, to characterize the spatiotemporal dynamics of transcriptional activity along the anterior-posterior axis, we

took the spatial average of signal across the dorsal-ventral axis and compared between genotypes, sexes, and

species (Fig 2A). The authors note that smoothed curves in the figures were created using smooth.spline in

ggplot2, which is not the same method for curve fitting as in the analysis. As such they are meaningful

reflections of the patterns in the data but not depictions of actual analyses. We found abundant spatial

quantitative variation in expression profiles (Fig 2-3). The expression profile of *hairy* around the MF harbors

variation between genotypes and there is an interaction between genotype and sex (Table 1, $p = 2$ x 10^{-3}, $p =$

0.02). There has also been evolution between species for *hairy* (Table 1, $p = 3$ x 10^{-4}). While *atonal* has not

evolved between species, there is variation in expression profile between genotypes, sexes, and there is an

interaction between genotype and sex (Fig 3A-C, Table 1, $p = 4$ x 10^{-4}, $p = .02$, $p = .02$). Surprisingly, given the

conservation of *Delta* in general, *Delta* harbors variation in spatial quantitative expression behind the MF

between genotypes and sexes (Table 1, $p = 2$ x 10^{-3}, $p = 7$ x 10^{-4}) and there are significant interactions between

genotype and sex (Fig 2B-C, Table 1, $p = 2$ x 10^{-3}). There has also been evolution of *Delta* between species, and

evolution of the interaction between species and sex (Table 1, $p = .03$, $p = 3$ x 10^{-4}). *hedgehog* is not different

between species but is significantly different between genotypes, sexes, and there is an interaction between the

two (Table 1, $p = 5$ x 10^{-4}, $p = .05$, $p = 1$ x 10^{-3}). There is also a significant interaction between species and sex

(Table 1, $p = .01$).

Table 1: The results of the full model for each gene.

hairy

Effect	numDF	denDF	F-value	p-value
Species	8	35	5.1	3 x 10^{-4}
Genotype	32	152	2.07	2 x 10^{-3}
Sex	8	35	1.08	0.32
Species x Sex	8	35	1.94	0.08
Genotype x Sex	32	152	1.73	0.02

atonal

Effect	numDF	denDF	F-value	p-value
Species	8	35	1.53	0.18
Genotype	32	152	2.31	4 x 10^{-4}
Sex	8	35	2.6	0.02
Species x Sex	8	35	1.91	0.09
Genotype x Sex	32	152	1.73	0.02

Delta

Effect	numDF	denDF	F-value	p-value
Species	8	35	2.43	0.03
Genotype	32	152	2.07	2 x 10^{-3}
Sex	8	35	4.55	7 x 10^{-4}
Species x Sex	8	35	5.03	3 x 10^{-4}
Genotype x Sex	32	152	2.03	2 x 10^{-3}

hedgehog

Effect	numDF	denDF	F-value	p-value
Species	8	35	0.87	0.55
Genotype	32	152	2.28	5 x 10^{-4}
Sex	8	35	2.23	0.05
Species x Sex	8	35	2.88	0.01
Genotype x Sex	32	152	2.17	1 x 10^{-3}

Significant *p*-values are indicated in bold, with gray shading.

Thus, *hairy* and *Delta* have evolved different spatial quantitative expression patterns between species, while

hairy, *atonal*, *Delta*, and *hedgehog* harbor cryptic variation within species and sexes. Given that there are

regulatory relationships between these genes, it is interesting to see that they do not all harbor variation for the

same factors. This could potentially be due to the influence of other unmeasured regulatory factors, or to

variation in the relationship between these genes and other components in the gene regulatory network.

However, whatever the source of 'buffering' of the network, be it the effect of other genes or threshold effects

on development, the fact that this information is not retained within the steps of the pathway supports our

supposition that this variation does not ultimately have a phenotypic effect.

Variation and evolution of the eye patterning gene network

There has been evolution in the regulatory logic of *hairy* and its upstream regulators *Delta* and *hedgehog*

between species (Fig 4A-C, Table 2, $p = .03$). There is also variation between sexes in the regulatory logic of

hairy and its upstream regulators *Delta* and *hedgehog* (Table 2, $p = .03$). There has been significant evolution of

the regulatory logic of *atonal*, in a significant interaction between species and sex ($p = 1 \times 10^{-3}$). Furthermore,

while there was no significant effect of genotype for *hairy*, there is for *atonal*, indicating that there is variation

segregating in the population affecting the relationship between *atonal*, *hedgehog*, and *Delta* ($p = 1.6 \times 10^{-5}$).

There is also a significant interaction between genotype and sex ($p = 1 \times 10^{-3}$). Thus, the relationship between

hairy and *atonal* and their regulators has evolved between species and sexes in *hairy*, and between genotypes

and sex in *atonal*. We illustrate this difference between species in Figure 3, where a different relationship

between *hairy* and *hedgehog* is visible between *D. melanogaster* and *D. simulans*. In brief, the frequency of cells

with a given log transformed level of expression are plotted against one another for *hairy* and *hedgehog*. *hairy* is

primarily expressed anterior to the MF and *hedgehog* posterior, and they have a different regulatory relationship

in each region with *hedgehog* activating *hairy* long range (anterior) and repressing it short range (posterior). This

is reflected in the frequency of cells expressing both genes for *D. melanogaster*, where anterior to the MF there

is a high frequency of *hairy* expressing cells and a low frequency of co-occurring high *hedgehog* expression.

Posterior to the MF the opposite is true, with high expression of *hedgehog* lacking concordance with any

expression of *hairy*. In *D. simulans*, posterior to the MF, this relationship is the same as in *D. melanogaster*.

However, in anterior to the MF this is not the case. Expression of *hairy* and *hedgehog* both increase as the other

increases, with widespread co-occurrence.

Table 2: The results of the full model for regulatory relationship between *hairy*, *Delta*, **and** *hedgehog*, **and** *atonal*, *Delta*, **and** *hedgehog*.

hairy

Effect	numDF	denDF	F-value	p-value
Species	2	41	4.03	0.25
Genotype	8	84	1.2	0.31
Sex	2	41	3.95	0.027
Species x Sex	2	41	0.78	0.46
Genotype x Sex	8	84	1,11	0.37

atonal

Effect	numDF	denDF	F-value	p-value
Species	2	41	2.64	0.08
Genotype	8	84	5.43	1.6×10^{-5}
Sex	2	41	0.67	0.52
Species x Sex	2	41	8.09	1×10^{-3}
Genotype x Sex	8	84	2.41	0.02

Significant *p*-values are indicated in bold, with gray shading.

Variation and evolution of MF structure

The amount that the eye discs were shifted is not significant for genotype, sex, or species, suggesting that

the relationship of maximum gene expression with the MF does not vary. However, the amount that they were

scaled is, after accounting for original differences in size, between species ($p = 1.38 \times 10^{-6}$). This suggests that

the total relative width of the MF varies between species, but not between genotypes or sexes. This is also

suggestive of evolving interrelationships among genes that could result in broader or narrower areas in which

they enhance or suppress expression of one another.

Discussion

Our results summarize a complicated pattern of variation sorting in the gene network involved in patterning the

MF. For example, the overall shape of the expression of *hedgehog* across the eye disc is different between

genotypes, sexes, and there is an interaction between species and sex and genotype and sex. *hedgehog*

upregulates *hairy*, but *hairy* has differences in expression between species (which *hedgehog* does not),

genotypes, and there is an interaction between genotype and sex. Thus, the differences seen in upstream

regulators, such as *hedgehog*, are not recapitulated in their downstream targets. In another example, *Delta/Notch*

is expected to repress *atonal*, but while *Delta/Notch* is significant for all categories tested *atonal* is only

significant for genotype, sex, and their interaction. It is possible that this variation is being mitigated or

dampened by other regulatory factors not assayed here, or that certain aspects of genetic background are more or

less sensitive to variation. For example, that fixed variation between species dampens variation at *Delta/Notch*

but sorting variation remains sensitive between genotypes, which propagates to *atonal*.

It may be that all of this variation is within levels tolerated by the network, as it has been shown that gene

networks can have thresholds of variation, below which differences in expression are effectively neutral. These

thresholds can also be two sided, creating a sigmoidal curve the center of which is neutral phenotypic space

(Felix and Barkoulas, 2015). Many studies have shown a relative insensitivity to variation in gene dosage, for

example in *Drosophila* early embryos the *bicoid* gradient results in normal development at one to four dosages

of the gene, but markedly abnormal development at six or more (Liu et al., 2013; Lucas et al., 2013; Namba et

al., 1997). It is also possible that the 'genetic stochasticity' documented in these genes is in fact deleterious, and

is being compensated for elsewhere in the network. While most deleterious mutations are purged by selection,

they may rise in frequency due to genetic drift or hitchhiking, among other possible causes (Burch and Chao,

1999; Chun and Fay, 2011; Estes and Lynch, 2003; McKenzie and Clarke, 1988). This type of compensatory

mutation has been documented in microbial and animal systems (K. M. Brown et al., 2010; Burch and Chao,

1999; Charusanti et al., 2010; Estes and Lynch, 2003; Estes et al., 2011; Maisnier-Patin and Andersson, 2004;

McKenzie, 1993; McKenzie et al., 1982; Moore et al., 2000; Stoebel et al., 2009; Szamecz et al., 2014).

Recently cell cycle heterogeneity has been implicated in the appearance of widespread noise in development,

however this is not responsible for the genetic stochasticity observed here as all MF cells are arrested at G1

(Keren et al., 2015; Kumar, 2013).

There have been other semi-quantitative approaches to studying spatial gene expression patterns. In another

study on *orthodenticle*, the authors found that the spatial and temporal pattern of gene expression was conserved

but the amount of gene product was not, though this work was not strictly quantitative given that measurements

were from *in situ* hybridization and reporter constructs and there was no rigorous statistical testing (Goering et

al., 2009). This is in contrast to our results which showed significant differences in the spatial relationship

between gene expression patterns between species. Other semi-quantitative works on the *Drosophila* embryo

using *in situ* hybridization found that the regulatory relationship between genes in the anterior-posterior

blastoderm patterning network were conserved, despite differences between species in their spatio-temporal

pattern (Fowlkes et al., 2011; Wunderlich et al., 2012). Here we find that the regulatory relationship between

atonal and *hairy*, and their regulators *hedgehog* and *Delta*, has evolved between species, sexes, and genotypes.

One of the important messages from this work is that rigorous statistical testing can uncover molecular

variation in spatial and quantitative developmental gene expression. Using the type of replication applied in

quantitative genetics with developmental data we were able to apply rigorous statistical models to micro-

evolutionary variation in development. Despite this variation, observed with repeatable observations of

developmental patterns among natural genotypes, the phenotypes of all flies are normal. This points to a

potential abundance of hidden noise in spatial and quantitative gene expression. The evolutionary approach to

development generally targets large changes that have occurred over broad phylogenetic distances (Ito et al.,

2013; Jeong et al., 2008; Kopp et al., 2000; Reed et al., 2011; Rosenblum et al., 2010; Signor et al., 2016; Yassin

et al., n.d.). Accordingly, the presence of abundant underlying variation is perhaps not a huge surprise. But it

does have large implications, as it demands that we modify developmental models in such a way that such

abundant genetic variation is buffered from perturbing the final phenotype. In the future, application of this type

of replicated, quantitative, spatially resolved data will have unique insights into the penetrance of disease

phenotypes and the origin of developmental defects.

Data Availability Statement

All data associated with the manuscript can be found at:

Author contributions

S. A. performed the staining and imaging of the eye discs, K. K. processed the imaging data, S.S. conceived of

the experiment, analyzed the processed data and wrote the paper, S. N. conceived of the experiment and

coordinated the research.

Acknowledgements

The authors thank S. Restrepo, M. Samsonova, A. Kopp, P. Marjoram, J. Butler, G. Mackerel, R. Hudson, E.

Williams, A. Lana, S. G. Frusina, H. Cagni and R. Hess for help with experimental procedures and manuscript

preparation. This work was supported by grants U01GM103804 and RO1GM102227 to S.V.

Figure 1. **a)** A summary of the eye patterning genes and pathway explored in this paper. The position of the MF

is shown in red, and its direction of movement indicated below. Regulatory relationships are illustrated either as

repression (bar) or activation (arrow). Regulatory relationships which are unclear are shown as gray dotted lines.

b) Example images from the dataset, illustrating gene expression patterns of each gene. The composite image

makes the additional point that we were able to analyze co-expression patterns of all four genes without needing

to stain each gene in different samples and infer gene co-expression patterns. **c)** An illustration of the general

expression pattern of each of the four genes in the study, along the anterior-posterior axis of the eye disc. The

authors note that all curves shown in figures were plotted using smooth.spline in ggplot2 in R, and are meant

only as visual representations of the data rather than formal reflection of the analysis.

Figure 2. **a)** This is an example of a curve being fitted to the gene expression profiles, though note that the curve

corresponds to the average in a given row (*x*-axis). The hexagons are intended to represent cells with varying

amounts of *hairy* expression, from the highest (red) to the lowest (white). **b)** An illustration of variation in *Delta*

expression between species and sexes. Curves shown are fitted to all genotypes within a sex and species with

confidence intervals indicated in gray. **c)** An illustration using the imaginal disc of how *Delta* expression varied

between species and sexes, with lower expression in *D. melanogaster* females and *D. simulans* males **d)**

Evolution of *Delta* illustrated within the context of the gene network, illustrating how changes in *Delta*

expression are not perturbing the gene network and result in phenotypically normal *Drosophila*.

Figure 3. **a)** An illustration of variation in *atonal* expression between genotypes and sexes. Curves shown are

fitted to each genotype and sex with confidence intervals indicated in gray. **b)** An illustration using the imaginal

disc of how *atonal* expression varied between genotypes, with lower expression in females of *D. melanogaster*

R153 and males of *D. simulans* Sz208. *D. simulans* Sz173 has lower expression than females of Sz208 but it is

not sexually dimorphic. **c)** Evolution of *atonal* illustrated within the context of the gene network, illustrating

how changes in *atonal* expression are not perturbing the gene network and result in phenotypically normal

Drosophila.

Figure 4. **a)** An example of variation in regulatory logic between *D. simulans* and *D. melanogaster* for *hairy* and

hedgehog. The heat map illustrates the density of points, and thus reflects the frequency of a given co-expression

profile between *hairy* and *hedgehog*. Gene expression values were log-transformed to better illustrate lower

values and split between anterior to the MF and posterior to the MF. The split between the two regions was to

investigate the possibility that *hedgehog* had a different regulatory relationship with *hairy* depending upon its

relationship to the MF, given that *hedgehog* is thought to activate *hairy* long range and repress *hairy* short range.

b) An illustration of the change in quantitative spatial expression of *hairy* and *hedgehog* between species and

sexes, with the position of the center of the MF marked with a dotted blue line. The red circle emphasizes a large

change in maximum *hedgehog* expression in males of the two species.

Atkins, M., Jiang, Y., Sansores-Garcia, L., Jusiak, B., Halder, G., Mardon, G., 2013. Dynamic rewiring of the
Drosophila retinal determination network switches its function from selector to differentiation. PLoS Genet.
9, e1003731.
Baker, N.E., Yu, S.Y., 2001. The EGF receptor defines domains of cell cycle progression and survival to
regulate cell number in the developing *Drosophila* eye. Cell 104, 699–708.
Baker, N.E., Yu, S.Y., 1997. Proneural function of neurogenic genes in the developing *Drosophila* eye. Curr.
Biol. 7, 122–132.
Bhattacharya, A., Baker, N.E., 2012. The role of the bHLH protein *hairy* in morphogenetic furrow progression
in the developing *Drosophila* eye. PLoS ONE 7, e47503.
Bhattacharya, A., Baker, N.E., 2009. The HLH protein *Extramacrochaetae* is required for R7 cell and cone cell
fates in the *Drosophila* eye. Dev. biol.
Bothma, J.P., Garcia, H.G., Esposito, E., Schlissel, G., Gregor, T., Levine, M., 2014. Dynamic regulation of eve
stripe 2 expression reveals transcriptional bursts in living *Drosophila* embryos. Proc. Nat. Acad. Sci. USA
111, 10598–10603.
Brennan, C.A., Ashburner, M., Moses, K., 1998. Ecdysone pathway is required for furrow progression in the
developing *Drosophila* eye. Development 125, 2653–2664.
Brown, K.M., Costanzo, M.S., Xu, W., Roy, S., Lozovsky, E.R., Hartl, D.L., 2010. Compensatory mutations
restore fitness during the evolution of Dihydrofolate reductase. Mol. Biol. Evol. 27, 2682–2690.
Brown, N.L., Sattler, C.A., Markey, D.R., Carroll, S.B., 1991. *hairy* gene function in the *Drosophila* eye: normal
expression is dispensable but ectopic expression alters cell fates. Development 113, 1245–1256.
Brown, N.L., Sattler, C.A., Paddock, S.W., Carroll, S.B., 1995. *Hairy* and *emc* negatively regulate
morphogenetic furrow progression in the *Drosophila* eye. Cell 80, 879–887.
Burch, C.L., Chao, L., 1999. Evolution by small steps and rugged landscapes in the RNA virus phi6. Genetics
151, 921–927.
Charusanti, P., Conrad, T.M., Knight, E.M., Venkataraman, K., Fong, N.L., Xie, B., Gao, Y., Palsson, B.Ø.,
2010. Genetic basis of growth adaptation of *Escherichia coli* after deletion of *pgi*, a major metabolic gene.
PLoS Genet. 6, e1001186.
Choi, H.M.T., Beck, V.A., Pierce, N.A., 2014. Next-generation *in situ* hybridization chain reaction: higher gain,
lower cost, greater durability. ACS Nano 8, 4284–4294.
Chun, S., Fay, J.C., 2011. Evidence for hitchhiking of deleterious mutations within the human genome. PLoS
Genet. 7, e1002240.
Cook, T., Pichaud, F., Sonneville, R., Papatsenko, D., Desplan, C., 2003. Distinction between color
photoreceptor cell fates is controlled by *Prospero* in *Drosophila*. Dev. Cell 4, 853–864.
Dominguez, M., Hafen, E., 1997. *Hedgehog* directly controls initiation and propagation of retinal differentiation
in the *Drosophila* eye. Genes Dev. 11, 3254–3264.
Dominguez, M., Wasserman, J.D., Freeman, M., 1998. Multiple functions of the EGF receptor in *Drosophila*
eye development. Curr. Biol. 8, 1039–1048.
Escudero, L.M., Freeman, M., 2007. Mechanism of G1 arrest in the *Drosophila* eye imaginal disc. BMC Dev.
Biol. 7, 13.
Estes, S., Lynch, M., 2003. Rapid fitness recovery in mutationally degraded lines of *Caenorhabditis elegans*.
Evolution 57, 1022–1030.
Estes, S., Phillips, P.C., Denver, D.R., 2011. Fitness recovery and compensatory evolution in natural mutant
lines of *C. elegans*. Evolution 65, 2335–2344.
Felix, M.-A., Barkoulas, M., 2015. Pervasive robustness in biological systems. Nat. Rev. Genet. 16, 483–496.
doi:10.1038/nrg3949
Felsenfeld, A.L., Kennison, J.A., 1995. Positional signaling by *hedgehog* in *Drosophila* imaginal disc
development. Development 121, 1–10.
Firth, L.C., Baker, N.E., 2005. Extracellular signals responsible for spatially regulated proliferation in the
differentiating *Drosophila* eye. Dev. Cell 8, 541–551.
Firth, L.C., Bhattacharya, A., Baker, N.E., 2010. Cell cycle arrest by a gradient of *Dpp* signaling during
Drosophila eye development. BMC Dev. Biol. 10, 28.
Fowlkes, C.C., Eckenrode, K.B., Bragdon, M.D., Meyer, M., Wunderlich, Z., Simirenko, L., Luengo Hendriks,
C.L., Keranen, S.V.E., Henriquez, C., Knowles, D.W., Biggin, M.D., Eisen, M.B., DePace, A.H., 2011. A

494   Conserved developmental patterning network produces quantitatively different output in multiple species of
495   *Drosophila*. PLoS Genet. 7, e1002346.
496 Freeman, A.B.A.M., 2001. *Notch* signalling and the initiation of neural development in the *Drosophila* eye.
497   Development 128, 3889-3898.
498 Fukaya, T., Lim, B., Levine, M., 2016. Enhancer control of transcriptional bursting. Cell 166, 358–368.
499 Gavish, A., Shwartz, A., Weizman, A., Schejter, E., Shilo, B.-Z., Barkai, N., 2016. Periodic patterning of the
500   *Drosophila* eye is stabilized by the diffusible activator *Scabrous*. Nat. Comm. 7, 1–10.
501 Goering, L.M., Hunt, P.K., Heighington, C., Busick, C., Pennings, P., Hermisson, J., Kumar, S., Gibson, G.,
502   2009. Association of *orthodenticle* with natural variation for early embryonic patterning in *Drosophila*
503   *melanogaster*. J. Exp. Zool. 312B, 841–854.
504 Heberlein, U., Wolff, T., Rubin, G.M., 1993. The TGF[beta] homolog *dpp* and the segment polarity gene
505   *hedgehog* are required for propagation of a morphogenetic wave in the *Drosophila* retina. Cell 75, 913–926.
506 Ito, Y., Harigai, A., Nakata, M., Hosoya, T., Araya, K., Oba, Y., Ito, A., Ohde, T., Yaginuma, T., Niimi, T.,
507   2013. The role of *doublesex* in the evolution of exaggerated horns in the Japanese rhinoceros beetle. Nat.
508   Neurosci. 14, 561–567.
509 Jarman, A.P., Grell, E.H., Ackerman, L., Jan, L.Y., Jan, Y.N., 1994. *atonal* is the proneural gene for *Drosophila*
510   photoreceptors. Nature 369, 398–400.
511 Jeong, S., Rebeiz, M., Andolfatto, P., Werner, T., True, J., Carroll, S.B., 2008. The evolution of gene regulation
512   underlies a morphological difference between two *Drosophila* sister species. Cell 132, 783–793.
513 Johnston, R.J., Desplan, C., 2014. Interchromosomal communication coordinates intrinsically stochastic
514   expression between alleles. Science 343, 661–665.
515 Johnston, R.J., Jr, Otake, Y., Sood, P., Vogt, N., Behnia, R., Vasiliauskas, D., McDonald, E., Xie, B., Koenig,
516   S., Wolf, R., Cook, T., Gebelein, B., Kussell, E., Nakagoshi, H., Desplan, C., 2011. Interlocked feedforward
517   loops control cell-type-specific rhodopsin expression in the *Drosophila* eye. Cell 145, 956–968.
518 Keren, L., van Dijk, D., Weingarten-Gabbay, S., Davidi, D., Jona, G., Weinberger, A., Milo, R., Segal, E., 2015.
519   Noise in gene expression is coupled to growth rate. Genome Res. 25, 1893–1902.
520 Kopp, A., Duncan, I., Godt, D., Carroll, S.B., 2000. Genetic control and evolution of sexually dimorphic
521   characters in *Drosophila*. Nature 408, 553–559.
522 Kozlov, K., Kosheverova, V., Kamentseva, R., Kharchenko, M., Sokolkova, A., Kornilova, E., Samsonova, M.,
523   2017. Quantitative analysis of the heterogeneous population of endocytic vesicles. J. Bioinform. Comput.
524   Biol. 10, 1750008.
525 Kumar, J.P., 2013. Catching the next wave: Patterning of the *Drosophila* eye by the morphogenetic furrow, in:
526   molecular genetics of axial patterning, growth and disease in the *Drosophila* eye. *in Molecular Genetics of*
527   *Axial Patterning*, Springer New York, New York, NY, pp. 75–97.
528 Li, W., Ohlmeyer, J.T., Lane, M.E., Kalderon, D., 1995. Function of protein kinase A in *hedgehog* signal
529   transduction and *Drosophila* imaginal disc development. Cell 80, 553–562.
530 Li, Y., Baker, N.E., 2001. Proneural enhancement by *Notch* overcomes Suppressor-of-Hairless repressor
531   function in the developing *Drosophila* eye. Curr. Biol. 11, 330–338.
532 Liu, F., Morrison, A.H., Gregor, T., 2013. Dynamic interpretation of maternal inputs by the *Drosophila*
533   segmentation gene network. Proc. Nat. Acad. Sci. USA 110, 6724–6729.
534 Lucas, T., Ferraro, T., Roelens, B., Las Heras Chanes, De, J., Walczak, A.M., Coppey, M., Dostatni, N., 2013.
535   Live imaging of *bicoid*-dependent transcription in *Drosophila* embryos. Curr. Biol. 23, 2135–2139.
536 Ma, C., Moses, K., 1995. *Wingless* and *patched* are negative regulators of the morphogenetic furrow and can
537   affect tissue polarity in the developing *Drosophila* compound eye. Development 121, 2279–2289.
538 Ma, C., Zhou, Y., Beachy, P.A., Moses, K., 1993. The segment polarity gene *hedgehog* is required for
539   progression of the morphogenetic furrow in the developing *Drosophila* eye. Cell 75, 927–938.
540 Mackay, T.F.C., Richards, S., Stone, E.A., Barbadilla, A., Ayroles, J.F., Zhu, D., Casillas, S., Han, Y., Magwire,
541   M.M., Cridland, J.M., Richardson, M.F., Anholt, R.R.H., Barrón, M., Bess, C., Blankenburg, K.P., Carbone,
542   M.A., Castellano, D., Chaboub, L., Duncan, L., Harris, Z., Javaid, M., Jayaseelan, J.C., Jhangiani, S.N.,
543   Jordan, K.W., Lara, F., Lawrence, F., Lee, S.L., Librado, P., Linheiro, R.S., Lyman, R.F., Mackey, A.J.,
544   Munidasa, M., Muzny, D.M., Nazareth, L., Newsham, I., Perales, L., Pu, L.-L., Qu, C., Ràmia, M., Reid,
545   J.G., Rollmann, S.M., Rozas, J., Saada, N., Turlapati, L., Worley, K.C., Wu, Y.-Q., Yamamoto, A., Zhu, Y.,

Bergman, C.M., Thornton, K.R., Mittelman, D., Gibbs, R.A., 2012. The *Drosophila* melanogaster Genetic Reference Panel. Nature 482, 173–178. d

Maisnier-Patin, S., Andersson, D.I., 2004. Adaptation to the deleterious effects of antimicrobial drug resistance mutations by compensatory evolution. R. Microbiol. 155, 360–369.

McKenzie, J.A., 1993. Measuring fitness and intergenic interactions: The evolution of resistance to diazinon in *Lucilia cuprina*. Genetica 90, 227–237.

McKenzie, J.A., Clarke, G.M., 1988. Diazinon resistance, fluctuating asymmetry and fitness in the Australian sheep blowfly, *Lucilia cuprina*. Genetics 120, 213–220.

McKenzie, J.A., Whitten, M.J., Adena, M.A., 1982. The effect of genetic background on the fitness of diazinon resistance genotypes of the Australian sheep blowfly, *Lucilia cuprina*. Heredity 49, 1–9.

Meyer, F., 1994. Topographic distance and watershed lines. Signal processing 38, 113–125.

Moore, F.B., Rozen, D.E., Lenski, R.E., 2000. Pervasive compensatory adaptation in *Escherichia coli*. Proc. R. Soc. B 267, 515–522.

Namba, R., Pazdera, T.M., Cerrone, R.L., Minden, J.S., 1997. *Drosophila* embryonic pattern repair: how embryos respond to *bicoid* dosage alteration. Development 124, 1393–1403.

Paaby, A., Gibson, G., 2016. Cryptic genetic variation in evolutionary developmental genetics. Biology 5, 28–13.

Pan, D., Rubin, G.M., 1995. cAMP-dependent protein kinase and *hedgehog* act antagonistically in regulating decapentaplegic transcription in *Drosophila* imaginal discs. Cell 80, 543–552.

Raj, A., van den Bogaard, P., Rifkin, S.A., van Oudenaarden, A., Tyagi, S., 2008. Imaging individual mRNA molecules using multiple singly labeled probes. Nat. Meth. 5, 877–879.

Reed, R.D., Papa, R., Martin, A., Hines, H.M., Counterman, B.A., Pardo-Diaz, C., Jiggins, C.D., Chamberlain, N.L., Kronforst, M.R., Chen, R., Halder, G., Nijhout, H.F., McMillan, W.O., 2011. *optix* drives the repeated convergent evolution of butterfly wing pattern mimicry. Science 333, 1137–1141.

Roignant, J.-Y., Treisman, J.E., 2009. Pattern formation in the *Drosophila* eye disc. Int. J. Dev. Biol. 53, 795–

Rosenblum, E.B., Römpler, H., Schöneberg, T., Hoekstra, H.E., 2010. Molecular and functional basis of phenotypic convergence in white lizards at White Sands. Proc. Nat. Acad. Sci. USA 107, 2113–2117.

Shah, S., Lubeck, E., Schwarzkopf, M., He, T.F., 2016. Single-molecule RNA detection at depth via hybridization chain reaction and tissue hydrogel embedding and clearing. Development. 143, 2862-2867.

Signor, S.A., Liu, Y., Rebeiz, M., Kopp, A., 2016. Genetic convergence in the evolution of male-specific color patterns in *Drosophila*. Curr. Biol. 26, 2423–2433.

Signor, S.A., New, F.N., Nuzhdin, S., 2017. A Large panel of *Drosophila simulans* reveals an abundance of common variants. Genome Biol. Evol. 10, 189–206.

Spencer, S.A., Powell, P.A., Miller, D.T., Cagan, R.L., 1998. Regulation of EGF receptor signaling establishes pattern across the developing *Drosophila* retina. Development 125, 4777–4790.

Stoebel, D.M., Hokamp, K., Last, M.S., Dorman, C.J., 2009. Compensatory evolution of gene regulation in response to stress by *Escherichia coli* lacking RpoS. PLoS Genet. 5, e1000671.

Greenwood, S., and Struhl, G., 1999. Progression of the morphogenetic furrow in the *Drosophila* eye: the roles of *hedgehog*, *Decapentaplegic* and the Raf pathway. Development. 126, 5795-5808.

Strutt, D.I., Mlodzik, M., 1997. *Hedgehog* is an indirect regulator of morphogenetic furrow progression in the *Drosophila* eye disc. Development 124, 3233–3240.

Sun, Y., Jan, L.Y., Jan, Y.N., 1998. Transcriptional regulation of *atonal* during development of the *Drosophila* peripheral nervous system. Development 125, 3731–3740.

Szamecz, B., Boross, G., Kalapis, D., Kovács, K., Fekete, G., Farkas, Z., Lázár, V., Hrtyan, M., Kemmeren, P., Groot Koerkamp, M.J.A., Rutkai, E., Holstege, F.C.P., Papp, B., Pál, C., 2014. The genomic landscape of compensatory evolution. PLoS Biol. 12, e1001935.

Tantale, K., Kozulic-Pirher, A., Lesne, A., Victor, J.-M., Robert, M.-C.E.C., Capozi, S., Chouaib, R., cker, V.B.A., Mateos-Langerak, J., Darzacq, X., Zimmer, C., Mueller, F., Basyuk, E., Bertrand, E., 2016. A single-molecule view of transcription reveals convoys of RNA polymerases and multi-scale bursting. Nat. Comm. 7, 1–14.

Tsachaki, M., Sprecher, S.G., 2011. Genetic and developmental mechanisms underlying the formation of the *Drosophila* compound eye. Dev. Dyn. 241, 40–56.

Vincent, L., 1993. Morphological grayscale reconstruction in image analysis: Applications and efficient
algorithms. IEEE transactions on image processing 2, 176–201.
Wernet, M.F., Mazzoni, E.O., Çelik, A., Duncan, D.M., Duncan, I., Desplan, C., 2006. Stochastic spineless
expression creates the retinal mosaic for colour vision. Nature 440, 174–180.
Wunderlich, Z., Bragdon, M.D., Eckenrode, K.B., Lydiard-Martin, T., Pearl-Waserman, S., DePace, A.H., 2012.
Dissecting sources of quantitative gene expression pattern divergence between *Drosophila* species. Mol.
Syst. Biol. 8, 1–15.
Yassin, A., Delaney, E.K., Reddieux, A., Bastide, H., Seher, T., Appleton, N.C., Lack, J.B., David, J.R.,
Chenoweth, S.F., Pool, J.E., Kopp, A., 2017. *pdm3* is responsible for recurrent evolution of female-limited
color dimorphism in *Drosophil*. Curr. Biol. 26, 2412-2422.
608
609

Supporting Information

611 **S1 Fig. Typical anterior-to-posterior expression patterns in the MF.** *hairy* (white), *atonal* (red), *Delta*
612 (green), and *hedgehog* (blue), 3rd instar larva eye disc labeled using HCR. Please note that these images contain
613 portions of the antennal disc, which were excluded from any analysis. Scale bars are 30 μm.

614 **S1 File Protocol: Hybridization chain reaction *Drosophila* imaginal discs**

615 **S1 Table The sequence of each of the probes used for hybridization chain reaction.**