REPORT

Quantitative variation and evolution of spatially explicit morphogen expression in *Drosophila*

Sammi Ali[1*], Sarah A. Signor[1*], Konstantin Kozlov[2], Sergey V. Nuzhdin[1,2]

[1] Section of Molecular and Computational Biology, University of Southern California, Los Angeles,

California, 90089 USA

[2] Department of Applied Mathematics, St. Petersburg State Polytechnic University, St. Petersburg,

195251, Russia

*Co-first author

SUMMARY

Developmental gene networks are deployed in complex temporal and spatial patterns. Methods for

studying gene networks and expression typically only capture variation along one axis, i.e. spatial but

not quantitative variation or vice versa. Here we use hybridization chain reaction (HCR) to

quantitatively measure spatial gene expression in multiple genotypes from two sexes of two species

(*Drosophila melanogaster* and *D. simulans*). We do this in the eye imaginal disc, which is initially

patterned by a wave of differentiation marked by a visible indentation of the tissue, termed the

morphogenetic furrow (MF), that passes from the posterior to the anterior of the disc, giving each disc

an element of both time and space in development (Fig 1). The furrow is initiated by *hedgehog,* which

both represses (short range) and activates (long range) *hairy* (Fig 1) [1,2]. *hedgehog* also activates

the expression of *atonal*, driving the furrow anteriorly [3-5]. *Delta/Notch* is under the control of

hedgehog and represses *atonal* [6,7]. We analyze the spatial quantitative expression of *hedgehog*,

hairy, *atonal*, and *Delta* to understand the evolving regulatory logic of the gene network and changes

in spatial dynamics between sexes and species. We find that gene expression varies among

genotypes and sexes, but remains stable between species, with the exception of *Delta* and *hairy*.

Surprisingly, this apparent stability is accomplished through changes in the interrelations between

genes within a network, and is exhibited at both quantitative and spatial levels.

RESULTS AND DISCUSSION

Understanding the relationship between genotype and phenotype requires a mechanistic

understanding of development, as it is the ultimate connection between the two. However, analyzing

this relationship requires understanding how genes are deployed in quantitative replicable way.

Without quantitative replication, there can be no rigorous statistical testing when developmental

processes are compared among conditions, including health versus disease. To date there has been

a surprising deficiency of such quantitative genetic treatments of basic developmental processes. The

goal of this paper is to provide a proof-of-principle for the presence of abundant natural genetic

variation in morphogen deployment during development. We focus on the well-known morphogens

driving ommatidia specification in *Drosophila* [8-12].

We employed recently developed HCR to measure genotype, sex, and species-specific spatial

quantitative gene expression patterns. HCR is unique in that it produces gene expression patterns that

are both quantitative and spatial. While other approaches such as FISH can be adapted to detect

individual transcripts, HCR has a linear signal that is 20x brighter than FISH, it reduces non-specific

background staining, and it can detect 88% of single RNA molecules in a cell with an appropriately low

false discovery rate [13,14]. It is also highly repeatable, with different sets of probes targeted to the

same gene showing correlations of .93-.99 (S. Fraser, pers. comm.). The DNA probes were designed

and synthesized by Molecular Instruments [15] (Table S1). Four genes were multiplexed in each

preparation (*hairy*, *atonal*, *Delta*, and *hedgehog*), as orthogonally-designed hairpins allowed the

simultaneous amplification of their target sequences. Eye imaginal discs were dissected using

standard procedures from three genotypes and both sexes of previously published *D. melanogaster*

and *D. simulans* inbred panels [16,17]. In order to compare gene expression patterns between eye

imaginal discs gene expression was quantified and each image was registered to a common grid such

that the same regions are being compared despite variation in disc size and deformation (Methods

S1).

We interpret the evolution of quantitative spatial patterns of gene expression in light of regulatory

relationships among them. We approach a spatial and quantitative analysis of these gene expression

patterns in three ways, first by explicitly creating a spatial gene expression profile and comparing

between genotypes, sexes, and species. Second, we were interested in examining if the regulatory

relationship between these genes had evolved between species or harbors variation within a species.

Lastly, we investigated the possibility that the spatial relationship between these genes relative to the

MF had evolved or harbors variation within populations.

First, to characterize the spatiotemporal dynamics of transcriptional activity along the anterior-

posterior axis, we took the spatial average of signal across the dorsal-ventral axis and compared

between species, genotypes, and sexes (Fig 2). We fit curves to each gene expression profile using

the mgcv package in R, using a generalized additive model with integrated smoothness estimation

(Fig 2, Supplemental Experimental Procedures). Smoothing terms are represented using penalized

regression splines. The curves were fit to the original range of values and down sampled to eight

points to capture their overall profile.

What we found was that the spatial quantitative gene expression profile of all four genes varied

between genotypes, species and sexes in different ways (Fig 2). The profile of *hairy* expression is

significantly different between genotypes ($F_{40,195}$ = 2.33, p = 6.99 x 10^{-6}) and there is an interaction

between genotype x sex ($F_{40,195}$ = 1.62, p = 0.017) (See Fig 2a for the results of the full models). There

is also a significant difference between *D. simulans* and *D. melanogaster* ($F_{8,43}$ = 4.7, p = 0.0004). The

profile of *atonal* expression is significantly different between genotypes ($F_{40,195}$ = 2.04, p = 7.4 x 10^{-4}),

sexes ($F_{8,35}$ = 2.6, p = .024), and there is an interaction between the two terms ($F_{40,195}$ = 1.73, p

= .0077). The expression profile of *Delta* is significantly different between genotypes $F_{40,195}$ = 2.15, p

= .0003), sexes ($F_{8,35}$ = 4.55, p = .0007), and genotype x sex ($F_{40,195}$ = 1.91, p = .002). The expression

profile of *Delta* is also significantly different between species $F_{8,43}$ = 2.18, p = .048) and species x sex

($F_{8,43}$ = 2.55, p = .024). *hedgehog* is significantly different between genotypes ($F_{40,195}$ = 1.89, p

= .0025), sexes ($F_{8,35}$ = 2.23, p = .049), and genotype x sex ($F_{40,195}$ = 2.01, p = 9.5 x 10^{-4}). Thus, *hairy*

and *Delta* have evolved different spatial quantitative expression patterns between species, while *hairy*,

atonal, *Delta*, and *hedgehog* harbor variation within species and sexes.

Second, it is possible that the genes in the MF have evolved, or harbor variation, for how they

affect each other in a particular cell, termed here the 'regulatory logic'. For example, is high *atonal*

expression associated with high expression of *hedgehog*, given that *hedgehog* activates *atonal*? We

cannot parse changes in regulatory logic versus overall spatial expression pattern, thus we focus on

genes known to regulate one another. We also cannot exclude the role of other unmeasured genes.

We used the following equations to determine the relationship between the expression of these genes:

$$hairy\ (i,s)= k^{Dl} \times Dl(i,s) + k^{hh} \times hh(i,s) + \ \alpha$$

$$atonal(i,s)= k^{Dl} \times Dl(i,s) + k^{hh} \times hh(i,s) + \ \alpha$$

The coefficient k and constant α were fit using standard methods for multiple regression. Here

$hairy(i,s)$ and $atonal(i,s)$ are the measured expression level of each gene in cell i in individual s.

$Dl(i,s)$ and $hh(i,s)$ are vectors containing the corresponding expression levels of *hairy* and *atonal*'s

regulators *Delta* and *hedgehog*. To determine if the regulatory logic is the same between genotypes

and species we can then use the regression coefficients from these models in a MANOVA. The

regulatory logic of *hairy* and its upstream regulators *Delta* and *hedgehog* is different between species

($F_{2,49}$ = 3.52, p = .037) (Fig 3A-C). There is also a difference in the regulatory logic between sexes

($F_{2,41}$ = 3.5, p = .038). We illustrate this difference between species in Figure 3, where a different

relationship between *hairy* and *hedgehog* is visible between *D. melanogaster* and *D. simulans*. The

regulatory logic of *atonal* does not seem strictly to have evolved between species, but there is a

significant interaction between species x sex ($F_{2,49}$ = 4.17, p = .021). Furthermore, while there was no

significant effect of genotype for *hairy*, there is for *atonal*, indicating that there is variation segregating

in the population effecting the relationship between *atonal, hedgehog*, and *Delta* ($F_{10,84}$ = 5.17 p = 7.22

x 10^{-6}). There is also a significant interaction between genotype x sex ($F_{10,84}$ = 3.31, p = .0012). Thus,

the relationship between *hairy* and *atonal* and their regulators has evolved between species and

sexes in *hairy*, and between genotypes and sex in *atonal*. For example, in *D. simulans* this appears as

an approximately linear relationship between *hairy* and *hedgehog* expression that is not present in *D.*

melanogaster (Fig 3A). We cannot, however, rule out a role for other unmeasured genes in the

network.

Significantly different regulatory logic in males and females, and species, is somewhat surprising.

Eye size does seem to vary both within and between species, but the exact nature of this variation is

unclear. By one account female eyes in *D. melanogaster* are smaller than expected based on

allometry, while other work finds that the opposite is the case in both *D. melanogaster* and *D.*

simulans [19,20]. Body size is at least in part controlled by *Sex-lethal,* and sex specific development in

somatic tissues are governed by *doublesex* [21,22]. We have detected *doublesex* expression in the

eye anterior to the morphogenetic furrow, in cells prior to cell cycle arrest at the furrow (Supplemental

Fig 1). It is possible that an interaction between *doublesex* and *hairy*, *atonal*, *Delta*, or *hedgehog* is

mediating the change in regulatory logic between sexes.

Lastly, we wanted to understand if there had been evolution of the relationship between the furrow

and gene expression, or if variation existed for the size of the furrow overall. For example, the furrow

was called as the position of overlap between *atonal* and *hairy* expression, but it is unclear how the

overall gene expression pattern of these genes relates to their overlap. For example, is the position of

maximum expression of each always the same relative to the furrow? Two processes occurred to

make the furrow comparable between samples, they were shifted to occupy the same position

depending upon the position of overlap of *hairy* and *atonal*, and they were scaled to occupy the same

total area. The amount required to scale will depend both on the size of the original disc and the width

of the furrow relative to the disc. To account for differences in size we include the number of rows in

the original disc prior to any transformations as a cofactor and perform ANOVA in R. The amount that

they were shifted is not significant for genotype, sex, or species, suggesting that the relationship of

maximum gene expression with the furrow does not vary. However, the amount that they were scaled

is, after accounting for original differences in size, for both genotype and species ($F_5 = 32.43$, $p = 4.9$ x

10^{-5}, $F_1 = 7.43$, $p = 6.89$ x 10^{-7}). This suggests that the total width of the MF varies between species

and genotypes, but not between sexes. This is also suggestive of evolving interrelationships among

genes that could result in broader or narrower areas in which they enhance or suppress expression of

one another.

It must be noted that all of the *Drosophila* included in this study have phenotypically normal eyes.

This type of developmental variation which does not result in a phenotypic difference is generally

termed cryptic variation [23,24]. During normal development of the eye this variation is buffered by

downstream processes, including cell proliferation and apoptosis [25]. In the past this cryptic variation

has been experimentally exposed through the use of synthesized genetic backgrounds with large

perturbations that prevent buffering of variation [24]. From this work, it has been assumed that

developmental processes vary cryptically among natural genotypes, but it has never been

experimentally observed before. Here, we provide the first empirical support for this hypothesis.

In the past, the developmental approach to understanding gene networks has been to analyze

large effect mutations and their qualitative downstream effects. This results in robust data on

perturbed regulatory networks, but obfuscates information on more normal interrelationships among

genes. Similarly, the evolutionary approach to development generally targets large changes that have

occurred over large phylogenetic distances [26]. Here we take an entirely different approach by

focusing on small, quantitative variation between genotypes and closely related species. Using this

approach, we were able to quantify the regulatory relationships among genes and to observe how

these relationships are altered between sexes and how they evolve between sister taxa. Reconciling

the speed of this cryptic developmental evolution with the patterns of molecular evolution in the

underlying genes will yield new insights on the rules governing microevolution of gene regulatory

networks.

Supplemental Information

AUTHOR CONTRIBUTIONS: S. A. performed the staining and imaging of the eye discs, K. K.

processed the imaging data, S.S. conceived of the experiment, analyzed the processed data and

wrote the paper, S. N. conceived of the experiment, coordinated the research, and co-wrote the paper.

ACKNOWLEDGEMENTS: The authors thank S. Restrepo, M. Samsonova, P. Marjoram, J. Butler for

help with experimental procedures and manuscript preparation. This work was supported by grants

U01GM103804 and RO1GM102227 to S.V.

Figure 1. **a)** A summary of the eye patterning genes and pathway explored in this paper. The position

of the morphogenetic furrow is shown in pink, and its direction of travel indicated below. Regulatory

relationships are illustrated either as repression (bar) or activation (arrow). Shown at right is a

composite image of all four genes (*hairy* (white), *atonal* (red), *Delta* (green), and *hedgehog* (blue)).

Please note that this is a very simplified illustration, intended as a summary between the genes that

are the subject of this paper rather than an exhaustive illustration of the eye patterning gene network.

b) Example gene expression patterns of each gene. *hedgehog* and *Delta* were recolored in Photoshop

(Adobe, v. 21.1) without altering any aspects of expression level or contrast.

d)

hairy

Effect	numDF	denDF	F-value	p-value
Species	8	43	4.7	**0.0004**
Sex	8	43	1.15	0.32
Species x Sex	8	43	1.42	0.21
Genotype	40	195	2.33	**6.99 x 10^{-6}**
Sex	8	35	1.08	0.4
Genotype x Sex	40	195	1.62	**0.017**

atonal

Effect	numDF	denDF	F-value	p-value
Species	8	43	1.23	0.31
Sex	8	43	1.66	0.14
Species x Sex	8	43	1.7	0,12
Genotype	40	195	2.04	**7.4 x 10^{-4}**
Sex	8	35	2.6	**0.024**
Genotype x Sex	40	195	1.73	**0.0077**

Delta

Effect	numDF	denDF	F-value	p-value
Species	8	43	2.18	**0.048**
Sex	8	43	2.55	**0.023**
Species x Sex	8	43	2.52	**0.024**
Genotype	40	195	2.15	**0.0003**
Sex	8	35	4.55	**0.0007**
Genotype x Sex	40	195	1.91	**0.002**

hedgehog

Effect	numDF	denDF	F-value	p-value
Species	8	43	0.68	0.7
Sex	8	43	0.85	0.57
Species x Sex	8	43	1.57	0.16
Genotype	40	195	1.89	**0.0025**
Sex	8	35	2.23	**0.049**
Genotype x Sex	40	195	2.01	**9.5 x 10^{-4}**

Figure 2. **a)** An example of the data-processing used to produce gene expression profiles. *hairy*

expression is shown on the left for an imaginal disc, while on the right is the same disc after it has

been segmented and registered. The bottom is an example of a curve being fitted to the gene

expression profiles, though note that the curve corresponds to the average in a given row (*x-*

axis). **b)** An illustration of variation in hairy expression between species and sexes. Curves

shown are fitted to all genotypes within a sex and species with confidence intervals indicated in

gray. Only the effect of species was large enough to be significant, though it is clear that perhaps

with a larger sample size an interaction effect would also have been significant. **c)** An illustration

of variation in *atonal* expression between genotypes and sexes. Curves shown are fitted to all

samples within a genotype and sex with confidence intervals indicated in gray. The effect of

genotype, sex, and genotype x sex were significant. All *D. melanogaster* genotypes are shown in

shades of purple, *D. simulans* in blue. **d)** The results of the full model for each gene, significant *p*-

values are indicated and bold, with gray shading.

hairy

Effect	numDF	denDF	F-value	*p*-value
Species	2	49	3.52	**0.037**
Sex	2	49	3.5	**0.038**
Species x Sex	2	49	0.92	0.4
Genotype	10	84	1.68	0.1
Sex	2	41	3.95	**0.027**
Genotype x Sex	10	84	1.06	0.4

atonal

Effect	numDF	denDF	F-value	*p*-value
Species	2	49	2.2	0.12
Sex	2	49	0.49	0.62
Species x Sex	2	49	4.17	**0.021**
Genotype	10	84	5.17	**7.22×10^{-6}**
Sex	2	41	0.67	0.52
Genotype x Sex	10	84	3.31	**0.0012**

Figure 3. **a)** An example of the differences in regulatory logic between *D. simulans* and *D.*

melanogaster for *hairy* and *hedgehog*. The heat map illustrates the density of points, and thus reflects

the frequency of a given co-expression profile between *hairy* and *hedgehog*. Gene expression values

were log-transformed to better illustrate lower values and split between anterior to the MF and

posterior to the MF. The split between the two regions was to investigate the possibility that *hedgehog*

had a different regulatory relationship with *hairy* depending upon its relationship to the furrow, given

that *hedgehog* is thought to activate *hairy* long range and repress *hairy* short range. Indeed, anterior

to the MF *D. simulans* has an almost linear relationship between *hairy* and *hedgehog*, while in *D.*

melanogaster almost any value of *hedgehog* can be associated with high *hairy* expression. **b)** An

illustration of the change in the relationship between the expression of *hedgehog* and *hairy* between

D. simulans and *D. melanogaster* anterior to the MF. The dotted, darker line is meant to indicate less

of a regulatory relationship than the thick red line. c) The results of the full model investigating

changes in the regulatory relationship between *hairy*, *Delta*, and *hedgehog*, and *atonal*, *Delta*, and

hedgehog.

1.   Strutt DI, Mlodzik M. (1997). *hedgehog* is an indirect regulator of morphogenetic furrow
progression in the *Drosophila* eye disc. Development *124*, 3233–3240.
2.   Felsenfeld AL, Kennison JA. (1995). Positional signaling by hedgehog in *Drosophila* imaginal
disc development. Development *121*, 1–10.
3.   Heberlein U, Wolff T, Rubin GM. (1993). The TGFβ homolog *dpp* and the segment polarity
gene hedgehog are required for propagation of a morphogenetic wave in the *Drosophila* retina
Cell *75*, 913–926.
4.   Ma C, Zhou U, Beachy PA, Moses K. (2003). The segment polarity gene hedgehog is
required for progression of the morphogenetic furrow in the developing *Drosophila* eye. Cell
75, 927-936.
5.   Struhl SGAG. (1999). Progression of the morphogenetic furrow in the *Drosophila* eye: the
roles of *Hedgehog*, *Decapentaplegic* and the Raf pathway. Development *126*, 5795-5808.
6.   Gavish A, Shwartz A, Weizman A, Schejter E, Shilo B-Z, Barkai N. (2016). Periodic patterning
of the *Drosophila* eye is stabilized by the diffusible activator *Scabrous*. Nat Comm *7*, 1–10.
7.   Firth LC, Baker NE. (2005). Extracellular signals responsible for spatially regulated
proliferation in the differentiating *Drosophila* eye. Dev Cell *8*, 541–551.
8.   Atkins M, Jiang Y, Sansores-Garcia L, Jusiak B, Halder G, Mardon G. (2013). Dynamic
rewiring of the *Drosophila* retinal determination network switches its function from selector to
differentiation. PLoS Genet *9*, e1003731.
9.   Raj A, van den Bogaard P, Rifkin SA, van Oudenaarden A, Tyagi S. (2008). Imaging
individual mRNA molecules using multiple singly labeled probes. Nat Meth *5*, 877–879.
10.   Shah S, Lubeck E, Schwarzkopf M, He TF. (2016). Single-molecule RNA detection at depth
via hybridization chain reaction and tissue hydrogel embedding and clearing. Development
143, 2862-2867.
11.   Tsachaki M, Sprecher SG. (2011). Genetic and developmental mechanisms underlying the
formation of the *Drosophila* compound eye. Dev Dyn *241*:40–56.
12.   Li W, Ohlmeyer JT, Lane ME, Kalderon D. (1995). Function of protein kinase A in *hedgehog*
signal transduction and *Drosophila* imaginal disc development. Cell *80*, 553–562.
13.   Pan D, Rubin GM. (1995). cAMP-dependent protein kinase and hedgehog act antagonistically
in regulating *decapentaplegic* transcription in *Drosophila* imaginal discs. Cell *80*, 543–552.
14.   Ma C, Moses K. (1995). *Wingless* and *patched* are negative regulators of the morphogenetic
furrow and can affect tissue polarity in the developing *Drosophila* compound eye.
Development *121*, 2279–2289.
15.   Choi HMT, Beck VA, Pierce NA. (2014). Next-generation in situ hybridization chain reaction:
higher gain, lower cost, greater durability. ACS Nano *8*, 4284–4294.
16.   Signor SA, New F, Nuzhdin S. *in review*. An abundance of high frequency variance uncovered
in a large panel of Drosophila simulans. Genome Biol. Evol.
17.   Mackay TFC, Richards S, Stone EA, Barbadilla A, Ayroles JF, Zhu D, et al. (2012). The
Drosophila melanogaster Genetic Reference Panel. Nature *482*, 173–178.
18.   Heberlein U, Mosest K. (1995). Mechanisms of *Drosophila* retinal morphogenesis: the virtues
of being progressive. Cell *81*, 987–990.
19.   Posnien N, Hopfen C, Hilbrant M, Ramos-Womack M, Murat S, Schönauer A, et al. (2012).
Evolution of eye morphology and rhodopsin expression in the *Drosophila melanogaster*
species subgroup. PLoS ONE *7*, e37346.
20.   Cowley DE, Atchley WR. (1988). Quantitative genetics of *Drosophila melanogaster*. II.
Heritabilities and genetic correlations between sexes for head and thorax traits. Genetics *119*,
421–433.

21. Mathews KW, Cavegn M, Zwicky M. (2017). Sexual dimorphism of body size is controlled by
dosage of the X-chromosomal gene *Myc* and by the sex-determining gene *tra* in *Drosophila.*
Genetics *205*, 1215-1228.
22. Rideout EJ, Dornan AJ, Neville MC, Eadie S, Goodwin SF. (2010). Control of sexual
differentiation and behavior by the *doublesex* gene in *Drosophila melanogaster*. Nat Neurosci
13, 458–466.
23. Paaby AB, Rockman MV. (2014). Cryptic genetic variation: evolution's hidden substrate. Nat
Rev Genet *15*, 247–258.
24. Gibson G, Dworkin I. (2004). Uncovering cryptic genetic variation. Nat Rev Genet *5*, 681–690.
25. Baker NE, Yu SY. (2001). The EGF receptor defines domains of cell cycle progression and
survival to regulate cell number in the developing *Drosophila* eye. Cell *104*, 699–708.
26. Signor SA, Li Y, Rebeiz M, Kopp A. (2016). Genetic convergence in the evolution of male-
specific color patterns in *Drosophila*. Curr Biol *26*, 2423-2433.
290