Novel approach to quantitative spatial gene expression uncovers cryptic evolution in the developing

Drosophila eye

Sammi Ali[1*], Sarah A. Signor[1*+], Konstantin Kozlov[2], Sergey V. Nuzhdin[1,2]

[1]Molecular and Computational Biology, University of Southern California, Los Angeles, California,

90089 USA

[2]Department of Applied Mathematics, St. Petersburg State Polytechnic University, St. Petersburg,

195251, Russia

*Co-first author

+Corresponding author: 1050 Childs Way, Ray R. Irani Hall,

13   Molecular and Computational Biology

14   University of Southern California

15   Los Angeles, 90089

16   ssignor@usc.edu

17   Phone   213-740-3065

ABSTRACT

Robustness in development allows for the accumulation of cryptic variation, and this largely neutral variation is potentially important for both evolution and complex disease phenotypes. However, it has generally only been investigated as variation in the response to large genetic perturbations. Here we use newly developed methods to quantify spatial gene expression patterns during development of the *Drosophila* eye disc, and uncover cryptic variation in wildtype developmental systems. We focus on four conserved morphogens, *hairy*, *atonal*, *hedgehog*, and *Delta*, that are involved in specifying ommatidia in the developing eye. We find abundant cryptic variation within and between species, genotypes, and sexes, as well as cryptic variation in the regulatory logic between *atonal* and *hairy* and their regulators, *Delta* and *hedgehog*. This work paves the way for a synthesis between population and quantitative genetic approaches with that of developmental biology.

Introduction

Natural genetic variation within populations has long been the purview of evolutionary and population geneticists, while developmental biologists focus on the effect of large mutations in otherwise isogenic backgrounds (Paaby and Gibson 2016). This dearth of work on developmental variation in wildtype genetic backgrounds is in part because developmental approaches have long been restricted to data that is at best semi-quantitative (i.e. *in situ* hybridization, antibody staining). Indeed, gene expression studies are generally spatial or quantitative, but not both. Without quantitative replication, there can be no rigorous statistical testing when developmental processes are compared among conditions, including health versus disease. This is especially important given the potential for cryptic variation to result in disease phenotypes, for example the complex disease phenotypes seen in humans after their recent, rapid change in lifestyle (Gibson and Reed 2008; Gibson 2009; Felix 2012; Ward and Kellis 2012; Hu et al. 2016). Here we use hybridization chain reaction (HCR) to bridge this gap between developmental and quantitative or population genetics by quantitatively measuring spatial gene expression in multiple genotypes from two sexes of two species (*Drosophila melanogaster* and *D. simulans*). This is essentially the first 'population genetics of development' as we are able to evaluate wild type differences in spatial and quantitative gene expression at the level of genotype, species and sex. This includes the ability to multiplex four genes, as more commonly co-expression is inferred across samples. We use this enormous developmental dataset to focus on the well-known morphogens driving ommatidia specification in *Drosophila* (Li et al. 1995; Raj et al. 2008; Tsachaki and Sprecher 2011; Atkins et al. 2013; Shah et al. 2016).

The *Drosophila* eye is formed from an imaginal disc, which is initially patterned by a wave of differentiation marked by a visible indentation of the tissue, termed the morphogenetic furrow (MF). The MF passes from the posterior to the anterior of the disc over a period of two days (90 minutes per adjacent row), giving each disc an element of both time and space in development (Fig 1) (Roignant and Treisman 2009). The furrow is initiated by *hedgehog,* which both represses (short range) and activates (long range) *hairy* (Fig 1) (Felsenfeld and Kennison 1995; Strutt and Mlodzik 1997). *hairy* represses *atonal*, preventing precocious neural development anterior to the MF (though this role has

been recently contested) (Brown et al. 1991; Brown, Sattler, Paddock, and Carroll 1995a;

Bhattacharya and Baker 2012). *hedgehog* activates the expression of *atonal*, driving the MF anteriorly

(Fig 1) (Heberlein et al. 1993; Ma et al. 1993; Struhl 1999). *atonal* is the proneural gene in *Drosophila*,

establishing the competency to become photoreceptor cells (Jarman et al. 1994). The relationship

between *Delta*/*Notch* and the other members of the pathway is not entirely clear, although in cells

posterior to the furrow *Delta/Notch* repress *atonal* (Fig 1) (Firth and Baker 2005; Gavish et al. 2016).

There is also some evidence that *Delta*/*Notch* repress negative regulators of *atonal* at the furrow, such

as *hairy* (Brown, Sattler, Paddock, and Carroll 1995b; Freeman 2001; Bhattacharya and Baker 2009).

In addition, there is some evidence suggesting that *Notch/Delta* are involved in the early stages of

atonal induction, and alternatively that *atona*l activates its own transcription (Baker and Yu 1997;

Dominguez and Hafen 1997; Dominguez et al. 1998; others 1998; Sun et al. 1998; Li and Baker

2001). There are many other genes involved in the specification of the eye disc that will not be

mentioned here, in favor of focusing on the genes we have assayed. We analyze the spatial

quantitative expression of *hedgehog*, *hairy*, *atonal*, and *Delta* to understand the evolving regulatory

logic of the gene network and changes in spatial dynamics between sexes and species.

Variation in gene expression within a species that has no phenotypic effect is termed cryptic

variation, and is thought to be a potential source of adaptive mutations when exposed to selection by

changing environmental factors (Dworkin et al. 2003; Gibson and Dworkin 2004; Gibson and Reed

2008; Duveau and Felix 2012; Kienle and Sommer 2013; Paaby and Rockman 2014; Lavagnino and

Fanara 2015; Taylor and Ehrenreich 2015a). Cryptic variation accumulates due to the robustness of

developmental systems to mutational perturbation (Duveau and Felix 2012; Felix 2012; Felix and

Barkoulas 2012; Kienle and Sommer 2013; Paaby and Rockman 2014; Paaby and Gibson 2016).

Although the final phenotype is the same, the presence of this cryptic variation effects the response to

mutational or environmental perturbation, and thus the evolutionary potential of the phenotype. In the

past cryptic variation has been experimentally exposed through the use of synthesized genetic

backgrounds with large perturbations that prevent buffering of variation (Braendle et al. 2010;

Woodruff et al. 2010; Duveau and Felix 2012; Chandler et al. 2014; Taylor and Ehrenreich 2015b).

While this work provides valuable insights into the importance of cryptic variation, cryptic variation in

developmental processes in natural genotypes has not been experimentally observed. Here, we will

investigate cryptic genetic variation with genotypes and sexes of *D. melanogaster* and *D. simulans*

using the four genes from the eye patterning network described above.

When cryptic genetic variation has fixed between species it has been called developmental

systems drift, compensatory evolution, and cryptic evolution, here we will refer to it as cryptic evolution

(True and Haag 2001; Landry et al. 2005; Felix 2007). These fixed differences do not necessarily

result from the fixation of cryptic variation, it can also be due to selection or compensatory mutation (or

both) (McGregor et al. 2001; Landry et al. 2005; Goncalves et al. 2012; Martinez et al. 2014; Szamecz

et al. 2014; Thompson et al. 2015; Fear et al. 2016). Most often it is investigated in the form of

homologous structures that are patterned by different genes, or homologous regulatory regions that

have evolved in the composition and placement of binding sites. The most well-known case of

homologous structures that are patterned by different genes comes from nematodes, where

homologous cells form phenotypically identical adult vulva in different species but are regulated by

different molecular mechanisms (Braendle et al. 2010; Duveau and Felix 2012; Felix and Barkoulas

2012; Barkoulas et al. 2013). In the latter case of homologous regulatory regions with conserved

expression patterns the most well-known case is the *even-skipped stripe 2* enhancer, where the

output is conserved between multiple species of *Drosophila* but the composition and placement of

binding sites has diverged (Ludwig et al. 1998; Ludwig et al. 2000). Here we will investigate cryptic

evolution on a more micro-evolutionary scale than previously (though it is still a comparison among

species) as rather than different genes or enhancer composition evolving between species we will

examine spatial quantitative expression level and evolution of the quantitative relationship between

genes in the eye patterning network.

We interpret cryptic variation and evolution in quantitative spatial patterns of gene expression in

light of regulatory relationships among them. We approach a spatial and quantitative analysis of these

gene expression patterns in three ways, first by explicitly creating a spatial gene expression profile

and comparing between genotypes, sexes, and species. Second, we were interested in examining if

the regulatory relationship between these genes had evolved between species or harbors variation

within a species. Lastly, we investigated the possibility that the spatial relationship between these

genes relative to the MF had evolved or harbors variation within populations.

Methods

Fly stocks

D. simulans were collected from the Zuma organic orchard in Zuma beach, CA in the spring of 2012

(Signor et al. *in press*). They were inbred by 15 generations of full sib crosses. *D. melanogaster* were

collected in Raleigh, North Carolina and inbred for 20 generations (Mackay et al. 2012).

Image acquisition

Staging and dissection of larvae

All flies were reared on a standard medium at 25° C with a 12-h light/12-h dark cycle. 120 hours after

hatching, 3^{rd} instar larva were placed in phosphate buffered saline (PBS) and separated by sex. Their

guts were carefully removed posteriorly and their body was inverted anteriorly to expose the brains,

eye discs and mouth hooks. After fixation and labeling (described below), eye discs were isolated and

mounted.

Hybridization Chain Reaction (HCR)

HCR is unique in that it produces gene expression patterns that are both quantitative and spatial

(Supplementary Materials). The DNA probes were designed and synthesized by Molecular

Instruments (Choi et al. 2014) (Table S1). Four genes were multiplexed in each preparation as

orthogonally-designed hairpins allowed the simultaneous amplification of their target sequences (Fig

1, S1). Each target mRNA was detected using five DNA probes to annotate the position and

expression levels for each of the four assayed genes (*hairy, atonal, Delta and hedgehog*).

Each probe contained two-initiator sequences (I1 and I2) that bound to a specific amplifier.

While other approaches such as FISH can be adapted to detect individual transcripts, HCR has a

linear signal that is 20x brighter than FISH, it reduces non-specific background staining, and it can

detect 88% of single RNA molecules in a cell with an appropriately low false discovery rate (Ma and

Moses 1995; Pan and Rubin 1995). It is also highly repeatable, with different sets of probes targeted

to the same gene showing correlations of .93-.99 (S. Fraser, pers. comm.).

The protocol for HCR was modified from (Choi et al. 2014) and is described briefly. The full

protocol is available in File S2. Inverted 3[rd] instar larva were fixed in 4% paraformaldehyde diluted with

PBS containing .2% Tween 20 (PBST). After fixation, larva were washed with PBST, then increasing

concentrations of methanol (30%, 70% and 100%) at 25° C. Larva were stored in 100% methanol at -

20° C. Methanol-fixed samples were thawed, washed with ethanol, re-permeabalized in 60% xylene,

washed with ethanol, then methanol and rehydrated with PBST. Samples were permeabalized with

proteinase K (4 mg/mL), fixed in 4% formaldehyde then washed with PBST at 25° C. Finally, at 45° C,

samples were pre-hybridized for 2 hours before the addition of all the probes. The probe-hybridized

larva were placed in wash buffer (Molecular Instruments) at 45° C to remove excess probes.

Fluorescently labeled hairpins were snap-cooled then added to the samples at 25° C and placed in the

dark to amplify the signal. Afterwards, samples were washed in 5X SSCT solution, isolated in PBST,

then placed in Prolong Gold anti-fade mounting medium (Molecular Probes).

Microscopy

Three dimensional images of mounted, HCR stained 3[rd] instar larva eye discs were acquired on a

Zeiss LSM 780 laser scanning microscope (Carl Zeiss MicroImaging, Inc., Thornwood, NY, USA) with

Objective Plan-Apochromat 63x/1.40 Oil. The gain was adjusted to avoid pixel saturation.

Extraction of gene expression profiles

Overview

The first steps in the image analysis is bringing each image to the same orientation and segmenting it.

Image segmentation produces a mask in which pixels are assigned to objects or background. Here

the objects are one or several mRNA molecules. Then the cellular structure of the imaginal disc is

approximated using a hexagonal array. Though the real underlying cell structure of the imaginal disc is

technically able to be recognized, this was unsatisfactory in our data due to imaging noise. Thus, at

the second step using the R package hexbin we constructed a partition of the imaginal disc area into

elements that represent pseudo-cells and have a biologically-relevant hexagonal shape (Brennan et

al. 1998). The number of pseudo-cells was selected by visual inspection of the combined image in

which the hexagonal structure was overlaid onto the *atonal* channel to verify fit. We are primarily

interested in expression profiles around the MF, providing us a convenient landmark to align images

from different preparations, thereby assigning coordinates to the pseudo-cells. However, deformations

of the eye disc during growth and preparation sometimes distorts the MF. We used splines to correct

for any bending or deformation of the MF. Next, using the histograms of cumulative pixel intensities of

objects in expression domains and non-expressing areas we estimated the typical intensity of a

transcript and typical background signal, respectively. Consequently, the cumulative intensities greater

than the background are divided by the intensity attributed to single mRNA molecule to yield counts of

mRNA molecules. This normalizes the expression profiles and corrects for differences in microscope

gain between images. Finally, the gene expression profiles are estimated for every pseudo-cell.

Morphological reconstruction and contrast mapping segmentation

To detect gene transcripts within the image stacks we applied a version of the MrComas method that

was modified for processing 3D images (Kozlov et al. 2017). This approach first enhances contrast

within the image and reduces noise. The images were enlarged by a factor of four with the nearest-

neighbor algorithm. They were processed by morphological reconstruction using both opening and

closing, where closing (opening) is dilation (erosion) that removes extraneous dark (bright) spots and

connects bright (dark) objects (Vincent 1993). The contrast mapping operator assigns each pixel the

maximum value between the pixel-by-pixel difference of the reconstructed images and their pixel-by-

pixel product and produces the rough mask for each channel. An image, *I*, is mapping from a finite

rectangular subset *L* onto the discrete plane Z^2 into a discrete set 0, 1, …, N - 1 of gray levels. Let the

dilation δ_B and erosion ϵ_B by structural element *B* be defined as:

$$\delta_B(I) = \vee_{\gamma \epsilon_B} I(\gamma) = I \oplus B \qquad \epsilon_B(I) = \wedge_{\gamma \epsilon_B} I(\gamma) = I \ominus B$$

Where ∨ and ∧ denote infimum and supremum respectively. Then formulae:

$$\delta^1_{I,B}(J) = (J \oplus B) \wedge I \qquad \epsilon^1_{I,B} = (J \ominus B) \vee I$$

denote geodesic dilation $\delta^1_{I,B}$ and erosion $\epsilon^1_{I,B}$. Binary reconstruction extracts those connected

components of the mask image which are marked on the marker image, and in grayscale it extracts

the peaks of the masked image marked by the marker image. Using the dilated masks image I as the

marker $J: J = \delta_B(I)$ defines closing by reconstruction:

$$\gamma_B(I) = \epsilon^1_{I,B}\epsilon^1_{I,B} \dots \epsilon^1_{I,B}[\delta_B(I)]$$

201 Opening by reconstructions uses eroded mask I as a marker J:

$$\phi_B(I) = \delta^1_{I,B}\delta^1_{I,B} \dots \delta^1_{I,B}[\epsilon_B(I)]$$

203 Then the difference between closing and opening by reconstruction has the meaning of the gradient:

$$\nabla_B(I) = \gamma_B(I) - \phi_B(I)$$

205 To create strong discontinuities at object edges and flatten signal with the objects the contrast

206 mapping operator takes a maximum between the difference and the pixel-by-pixel produce of the

207 reconstructed images and produces a rough mask for each channel:

$$R = \max\{\nabla_B(I), \gamma_B(I) \odot \phi_B(I)\}$$

Subsequently, this mask was subjected to distance transform, which substituted each pixel value with

the number of pixels between it and the closest background pixel. This operation creates 'peaks' and

'valleys' of intensity inside foreground objects. To split erroneously merged objects watershed

transform was applied, which treats the whole image as a surface and intensity of each pixel as its

height and determines the watershed lines along the tops of ridges separating the catchment basins

(Meyer 1994). The quality of segmentation is assessed visually by inspection of the object boarders

overlaid with the original image. Finally, each mask is returned to its original size and quantitative

measures are made of shape and intensity characteristics such as the number of pixels, as well as

their mean and standard deviation in the detected object. MrComas is free and open source software

available at http://sourceforge.net/p/prostack/wiki/mrcomas.

Approximating the MF

We defined the position of the MF as the middle of overlap between *hairy* and *atonal* expression. The

shape of the MF was approximated with a spline using function smooth.spline in R. The degrees of

freedom and other parameters were chosen to make the approximation coincide visually with a MF

image.

Inferring counts of transcript number

Segmentation of the image provided a table of coordinates and the shape and intensity characteristics

of detected transcripts. Here, we applied filtering steps to remove false positives and determine the

count of mRNA transcripts. First, we assumed the object we detected as least intense but most

frequent corresponds to a single mRNA molecule. Then, we inferred background intensity for objects

outside of well-annotated domains of expression of the four genes. Assuming that the majority of true

objects contain a single molecule, we compare the distribution of cumulative intensities of particles in

expression domains and areas of known non-expression to obtain the typical intensity of a true single

molecule and a false positive, respectively. All detected signals that were lower than the typical

intensity of a false positive were removed from the dataset. The number of removed objects is

typically less then 10%. All other cumulative pixel intensities were divided by the typical intensity of a

true single molecule as normalization coefficient to yield an estimate of the number of mRNA

transcripts.

Image registration

We applied an affine coordinate transformation to each eye disc to make the corresponding maxima

and the width of expression patterns of four genes in different eyes coincide as closely as possible. To

do so, we shifted the coordinate system of each eye to its center and also scaled them in the A-P

direction. The center of the pattern in A-P direction is the MF.

We mapped the expression patterns to a unified hexagonal structure in order to make

comparisons between pseudo-cells from individual imaginal discs. The unified cell structure was

constructed using the R package hexbin. Each cell in the unified grid represents an 'average' cell from

individual eyes. The size of a hexagon in the unified grid is greater or equal than the cell size in the

individual eye. Thus, the number of molecules in each unified cell in the mapped pattern equals the

mean over the cells from native pattern that are covered by this unified cell. After such coordinate

transformation, the MF region is defined as 20 cells on either side of the MF, to focus the analysis on

the area of interest (the MF).

Filtering and quality controls for each eye disc

Some eye imaginal discs were damaged or deformed in the process of dissection or mounting, resulting

in regions of erroneous gene expression, such as disruptions to the MF. The expression profile of each

disc was examined by eye and these regions were individually trimmed out of the final dataset. At the

edges of each eye disc the pattern of the MF was also degraded, so each eye disc was trimmed dorso-

ventrally prior to analysis. Outliers were excluded from the dataset, determined as a single member of

the five replicates with more than a 3x difference in expression values. This resulted in a final dataset

of 55 eye discs.

Modeling sources of expression variation and evolution

Analysis of individual spatial gene expression patterns

We were primarily interested in variation in gene expression profiles across the eye disc, that is using

differences in expression averaged across rows along the *x*-axis. While the *y*-axis is of interest,

variation in the shape, size, degree of deformation, and occasional damage to the disc made this

analysis intractable. We fit curves to each gene expression profile using the mgcv package in R, using

a generalized additive model with integrated smoothness estimation. Smoothing terms are

represented using penalized regression splines. predict.gam was used to fit the curves to the original

range of values and down sample the curves to eight points. MANOVAs were performed using the

"Pillai" test for genotype x sex and species x sex.

Modeling framework to understand variation and evolution of the eye patterning gene network

We wanted to understand if cryptic variation existed within the regulatory logic of *hairy*, *atonal*, *Delta*,

and *hedgehog*, or if there had been cryptic evolution between species. To understand the regulatory

logic between genes we focused on biologically relevant relationships, such as the regulation of *hairy*

by *Delta* and *hedgehog,* but excluded such relationships as *hairy* and *atonal*. This was due to the low

overlap between *hairy* and *atonal* expression domains, where including cells where only one or

another was expressed would artificially create a relationship between expression levels. Both *hairy*

and *atonal* are downstream, directly or indirectly, of *Delta* and *hedgehog* thus it was these

relationships that were modeled. We limited the analysis to cells where all genes included were

expressed in at least ten molecules.

In the previous analysis, we investigated variation in the cryptic spatial quantitative expression

pattern of genes in the MF. Here, we will investigate the possibility that genes in the MF have evolved,

or harbor variation, for how they affect each other in particular cells. For example, is high *atonal*

expression associated with high expression of *hedgehog*, given that *hedgehog* activates *atonal*? We

used the following equations to determine the relationship between the expression of these genes:

$$hairy(i,s) = k^{Dl} \times Dl(i,s) + k^{hh} \times hh(i,s) + \alpha$$

$$atonal(i,s) = k^{Dl} \times Dl(i,s) + k^{hh} \times hh(i,s) + \alpha$$

The coefficient k and constant α were fit using standard methods for multiple regression. Here

$hairy(i,s)$ and $atonal(i,s)$ are the measured expression level of each gene in cell i in individual s.

$Dl(i,s)$ and $hh(i,s)$ are vectors containing the corresponding expression levels of *hairy* and *atonal*'s

regulators *Delta* and *hedgehog*. To determine if the regulatory logic is the same between genotypes

and species we can then use the regression coefficients from these models in a MANOVA. We note

that we cannot exclude the possibility that other unmeasured genes are responsible for producing this

variation.

Model for understanding overall variation and evolution of MF structure

Lastly, we wanted to understand if there is cryptic variation and/or evolution for the relationship

between the MF and gene expression, or if cryptic variation and/or evolution existed for the size of the

MF overall. For example, the MF was called as the position of overlap between *atonal* and *hairy*

expression, but it is unclear how the overall gene expression pattern of these genes relates to their

overlap. For example, is the position of maximum expression of each always the same relative to the

MF? Two processes occurred to make the MF comparable between samples, they were shifted to

occupy the same position depending upon the position of overlap of *hairy* and *atonal*, and they were

scaled to occupy the same total area. The amount required to scale will depend both on the size of the

original disc and the width of the MF relative to the disc. To account for differences in size we include

the number of rows in the original disc prior to any transformations as a cofactor and perform ANOVA

in R.

RESULTS

Individual spatial gene expression patterns

First, to characterize the spatiotemporal dynamics of transcriptional activity along the anterior-

posterior axis, we took the spatial average of signal across the dorsal-ventral axis and compared

between genotypes, sexes, and species (Fig 2A). We found abundant cryptic spatial quantitative

variation in expression profiles (Fig 2-3). The expression profile of *hairy* around the MF harbors cryptic

variation between genotypes and there is an interaction between genotype and sex (Table 1, p = 2 x

10^{-3}, p = 0.02). There has also been cryptic evolution between species for *hairy* (Table 1, p = 3 x 10^{-4}).

While *atonal* has not evolved between species, there is cryptic variation in expression profile between

genotypes, sexes, and there is an interaction between genotype and sex (Fig 3A-C, Table 1, p = 4 x

10^{-4}, p = .02, p = .02). Surprisingly, given the conservation of *Delta* in general, *Delta* harbors variation

in spatial quantitative expression behind the MF between genotypes and sexes (Table 1, p = 2 x 10^{-3},

p = 7 x 10^{-4}) and there are significant interactions between genotype and sex (Fig 2B-C, Table 1, p = 2

x 10^{-3}). There has also been cryptic evolution of *Delta* between species, and evolution of the

interaction between species and sex (Table 1, p = .03, p = 3 x 10^{-4}). *hedgehog* is not different between

species but is significantly different between genotypes, sexes, and there is an interaction between

the two (Table 1, p = 5 x 10^{-4}, p = .05, p = 1 x 10^{-3}). There is also a significant interaction between

species and sex (Table 1, p = .01).

Thus, *hairy* and *Delta* have cryptically evolved different spatial quantitative expression patterns

between species, while *hairy*, *atonal*, *Delta*, and *hedgehog* harbor cryptic variation within species and

sexes. Given that there are regulatory relationships between these genes, it is interesting to see that

they do not all harbor variation for the same factors. This could potentially be due to the influence of

other unmeasured regulatory factors, or to variation in the relationship between these genes and other

components in the gene regulatory network.

Variation and evolution of the eye patterning gene network

There has been cryptic evolution in the regulatory logic of *hairy* and its upstream regulators *Delta*

and *hedgehog* between species (Fig 4A-C, Table 2, *p* = .03). There is also cryptic variation between

sexes in the regulatory logic of *hairy* and its upstream regulators *Delta* and *hedgehog* (Table 2, *p*

= .03). There has been significant evolution of the regulatory logic of *atonal*, in a significant interaction

between species and sex (*p* = 1 x 10^{-3}). Furthermore, while there was no significant effect of genotype

for *hairy*, there is for *atonal*, indicating that there is cryptic variation segregating in the population

effecting the relationship between *atonal, hedgehog*, and *Delta* (*p* = 1.6 x 10^{-5}). There is also a

significant interaction between genotype and sex (*p* = 1 x 10^{-3}). Thus, the relationship between *hairy*

and *atonal* and their regulators has cryptically evolved between species and sexes in *hairy*, and

between genotypes and sex in *atonal* (cryptic variation). We illustrate this difference between species

in Figure 3, where a different relationship between *hairy* and *hedgehog* is visible between *D.*

melanogaster and *D. simulans*. In brief, the frequency of cells with a given log transformed level of

expression are plotted against one another for *hairy* and *hedgehog*. *hairy* is primarily expressed

anterior to the MF and *hedgehog* posterior, and they have a different regulatory relationship in each

region with *hedgehog* activating *hairy* long range (anterior) and repressing it short range (posterior).

This is reflected in the frequency of cells expressing both genes for *D. melanogaster*, where anterior to

the MF there is a high frequency of *hairy* expressing cells and a low frequency of co-occurring high

hedgehog expression. Posterior to the MF the opposite is true, with high expression of *hedgehog*

lacking concordance with any expression of *hairy*. In *D. simulans*, posterior to the MF, this relationship

is the same as in *D. melanogaster*. However, in anterior to the MF this is not the case. Expression of

hairy and *hedgehog* both increase as the other increases, with widespread co-occurrence.

Variation and evolution of MF structure

The amount that they were shifted is not significant for genotype, sex, or species, suggesting that

the relationship of maximum gene expression with the MF does not vary. However, the amount that

they were scaled is, after accounting for original differences in size, between species (*p* = 1.38 x 10^{-6}).

This suggests that the total relative width of the MF varies between species, but not between

genotypes or sexes. This is also suggestive of evolving interrelationships among genes that could

result in broader or narrower areas in which they enhance or suppress expression of one another.

Discussion

Our results summarize a complicated pattern of variation sorting in the gene network involved in

patterning the MF. For example, the overall shape of the expression of *hedgehog* across the eye disc

is different between genotypes, sexes, and there is an interaction between species and sex and

genotype and sex. *hedgehog* upregulates *hairy*, but *hairy* has differences in expression between

species (which *hedgehog* does not), genotypes, and there is an interaction between genotype and

sex. Thus, there is no propagation of the pattern seen in *hedgehog* through the network. In another

example, *Delta/Notch* is expected to repress *atonal*, but while *Delta/Notch* is significant for all

categories tested *atonal* is only significant for genotype, sex, and their interaction. It is possible that

this variation is being mitigated or dampened by other regulatory factors not assayed here, or that

certain aspects of genetic background are more or less sensitive to variation. For example, that fixed

variation between species dampens variation at *Delta/Notch* but sorting variation remains sensitive

between genotypes, which propagates to *atonal*.

It may be that all of this variation is within levels tolerated by the network, as it has been shown

that gene networks can have thresholds of variation, below which differences in expression are

effectively neutral. These thresholds can also be two sided, creating a sigmoidal curve the center of

which is neutral phenotypic space (Felix and Barkoulas 2015). Many studies have shown a relative

insensitivity to variation in gene dosage, for example in *Drosophila* early embryos the *bicoid* gradient

results in normal development at one to four dosages of the gene, but markedly abnormal

development at six or more (Namba et al. 1997; Liu et al. 2013; Lucas et al. 2013). It is also possible

that the cryptic evolution documented in these genes is in fact deleterious, and is being compensated

for elsewhere in the network. While most deleterious mutations are purged by selection, they may rise

in frequency due to genetic drift or hitchhiking, among other possible causes (McKenzie and Clarke

1988; Burch and Chao 1999; Estes and Lynch 2003; Chun and Fay 2011). This type of compensatory

mutation has been documented in microbial and animal systems (McKenzie et al. 1982; McKenzie

1993; Burch and Chao 1999; Moore et al. 2000; Estes and Lynch 2003; Maisnier-Patin and Andersson

2004; Stoebel et al. 2009; Brown et al. 2010; Charusanti et al. 2010; Estes et al. 2011; Szamecz et al.

2014). Recently cell cycle heterogeneity has been implicated in the appearance of widespread noise

in development, though that is less of problem here given that cell division is synchronized posterior

the MF and cells are at GI arrest within the MF (Kumar 2013; Keren et al. 2015).

Other than differences in the allometric relationship of eyes between males and females there has

been no documentation of sexual dimorphism in *Drosophila* eyes, which could potentially contribute to

the differences between sexes seen for some of the genes in the MF. However, we note that we

controlled for size of the imaginal disc in this study and the difference between males and females

seems to largely be driven by differences in maximum expression. Furthermore, the size of the eye

disc was not consistently significantly different between sexes. By one account female eyes in *D.*

melanogaster are smaller than expected based on allometry, while other work finds that the opposite

is the case in both *D. melanogaster* and *D. simulans* (Cowley and Atchley 1988; Posnien et al. 2012).

Body size is at least in part controlled by *Sex-lethal,* and sex specific development in somatic tissues

are governed by *doublesex* (Rideout et al. 2010; Mathews et al. 2017). We have detected *doublesex*

expression in the eye anterior to the MF, in cells prior to cell cycle arrest at the MF (Fig S2). It is

possible that an interaction between *doublesex* and *hairy*, *atonal*, *Delta*, or *hedgehog* is mediating the

change in cryptic evolution in regulatory logic between sexes. It is also possible that some of this

variation is not cryptic, and in fact has an undocumented effect on sex-specific development of the

eyes.

There have been other semi-quantitative approaches to studying spatial gene expression

patterns. In another study on *orthodenticle*, the authors found that the spatial and temporal pattern of

gene expression was conserved but the amount of gene product was not, though this work was not

strictly quantitative given that measurements were from *in situ* hybridization and reporter constructs

(Palsson et al. 2014). This is in contrast to our results which showed significant differences in the

spatial relationship between gene expression patterns between species. Other semi-quantitative

works on the *Drosophila* embryo using *in situ* hybridization found that the regulatory relationship between genes in the anterior-posterior blastoderm patterning network were conserved, despite differences between species in their spatio-temporal pattern (Fowlkes et al. 2011; Wunderlich et al. 2012). Here we find that the regulatory relationship between *atonal* and *hairy*, and their regulators *hedgehog* and *Delta*, has evolved between species, sexes, and genotypes.

In the past, the developmental approach to understanding gene networks has been to analyze large effect mutations and their qualitative downstream effects. This results in robust data on perturbed regulatory networks, but obfuscates information on more normal interrelationships among genes. Similarly, the evolutionary approach to development generally targets large changes that have occurred over broad phylogenetic distances (Kopp et al. 2000; Jeong et al. 2008; Rosenblum et al. 2010; Reed et al. 2011; Ito et al. 2013; Signor, Li, et al.; Yassin et al.). Here we take an entirely different approach by focusing on small, quantitative variation between genotypes and closely related species (Johnson 2006; Nunes et al. 2013). Using this approach, we were able to quantify the regulatory relationships among genes and to observe how these relationships are altered between genotypes, sexes and species. Reconciling the speed of this cryptic developmental evolution with the patterns of molecular evolution in the underlying genes will yield new insights on the rules governing microevolution of gene regulatory networks.

AUTHOR CONTRIBUTIONS: S. A. performed the staining and imaging of the eye discs, K. K. processed the imaging data, S.S. conceived of the experiment, analyzed the processed data and wrote the paper, S. N. conceived of the experiment, coordinated the research, and co-wrote the paper.

ACKNOWLEDGEMENTS: The authors thank S. Restrepo, M. Samsonova, P. Marjoram, J. Butler, G. Mackerel, R. Hudson, and E. Williams for help with experimental procedures and manuscript preparation. This work was supported by grants U01GM103804 and RO1GM102227 to S.V.

Figure 1. **a)** A summary of the eye patterning genes and pathway explored in this paper. The position

of the MF is shown in red, and its direction of movement indicated below. Regulatory relationships are

illustrated either as repression (bar) or activation (arrow). Regulatory relationships which are unclear

are shown as gray dotted lines. **b)** Example images from the dataset, illustrating gene expression

patterns of each gene. The composite image makes the additional point that we were able to analyze

co-expression patterns of all four genes without needing to stain each gene in different samples and

infer gene co-expression patterns. **c)** An illustration of the general expression pattern of each of the

four genes in the study, along the anterior-posterior axis of the eye disc.

Figure 2. **a)** This is an example of a curve being fitted to the gene expression profiles, though note that

the curve corresponds to the average in a given row (*x*-axis). The hexagons are intended to represent

cells with varying amounts of *hairy* expression, from the highest (red) to the lowest (white). **b)** An

illustration of variation in *Delta* expression between species and sexes. Curves shown are fitted to all

genotypes within a sex and species with confidence intervals indicated in gray. **c)** An illustration using

the imaginal disc of how *Delta* expression varied between species and sexes, with lower expression in

D. melanogaster females and *D. simulans* males **d)** Cryptic evolution of *Delta* illustrated within the

context of the gene network, illustrating how changes in *Delta* expression are not perturbing the gene

network and result in phenotypically normal *Drosophila*.

Figure 3. **a)** An illustration of cryptic variation in *atonal* expression between genotypes and sexes.

Curves shown are fitted to each genotype and sex with confidence intervals indicated in gray. **c)** An

illustration using the imaginal disc of how *atonal* expression varied between genotypes, with lower

expression in females of *D. melanogaster* R153 and males of *D. simulans* Sz208. *D. simulans* Sz173

has lower expression than females of Sz208 but it is not sexually dimorphic. **d)** Cryptic evolution of

atonal illustrated within the context of the gene network, illustrating how changes in *atonal* expression

are not perturbing the gene network and result in phenotypically normal *Drosophila*.

Figure 4. **a)** An example of cryptic variation in regulatory logic between *D. simulans* and *D. melanogaster* for *hairy* and *hedgehog*. The heat map illustrates the density of points, and thus reflects the frequency of a given co-expression profile between *hairy* and *hedgehog*. Gene expression values were log-transformed to better illustrate lower values and split between anterior to the MF and posterior to the MF. The split between the two regions was to investigate the possibility that *hedgehog* had a different regulatory relationship with *hairy* depending upon its relationship to the MF, given that *hedgehog* is thought to activate *hairy* long range and repress *hairy* short range. **b)** An illustration of the change in quantitative spatial expression of *hairy* and *hedgehog* between species and sexes, with the position of the center of the MF marked with a dotted blue line. The red circle emphasizes a large change in maximum *hedgehog* expression in males of the two species.

hairy

Effect	numDF	denDF	F-value	p-value
Species	8	35	5.1	3×10^{-4}
Genotype	32	152	2.07	2×10^{-3}
Sex	8	35	1.08	0.32
Species x Sex	8	35	1.94	0.08
Genotype x Sex	32	152	1.73	0.02

atonal

Effect	numDF	denDF	F-value	p-value
Species	8	35	1.53	0.18
Genotype	32	152	2.31	4×10^{-4}
Sex	8	35	2.6	0.02
Species x Sex	8	35	1.91	0.09
Genotype x Sex	32	152	1.73	0.02

Delta

Effect	numDF	denDF	F-value	p-value
Species	8	35	2.43	0.03
Genotype	32	152	2.07	2×10^{-3}
Sex	8	35	4.55	7×10^{-4}
Species x Sex	8	35	5.03	3×10^{-4}
Genotype x Sex	32	152	2.03	2×10^{-3}

hedgehog

Effect	numDF	denDF	F-value	p-value
Species	8	35	0.87	0.55
Genotype	32	152	2.28	5×10^{-4}
Sex	8	35	2.23	0.05
Species x Sex	8	35	2.88	0.01
Genotype x Sex	32	152	2.17	1×10^{-3}

Table 1: The results of the full model for each gene, significant *p*-values are indicated in bold, with gray shading.

hairy

Effect	numDF	denDF	F-value	p-value
Species	2	41	4.03	0.25
Genotype	8	84	1.2	0.31
Sex	2	41	3.95	0.027
Species x Sex	2	41	0.78	0.46
Genotype x Sex	8	84	1,11	0.37

atonal

Effect	numDF	denDF	F-value	p-value
Species	2	41	2.64	0.08
Genotype	8	84	5.43	1.6×10^{-5}
Sex	2	41	0.67	0.52
Species x Sex	2	41	8.09	1×10^{-3}
Genotype x Sex	8	84	2.41	0.02

Table 2: The results of the full model investigating changes in the regulatory relationship between *hairy*, *Delta*, and *hedgehog*, and *atonal*, *Delta*, and *hedgehog*. Significant *p*-values are indicated in bold, with gray shading.

Atkins M, Jiang Y, Sansores-Garcia L, Jusiak B, Halder G, Mardon G. 2013. Dynamic Rewiring of the
Drosophila Retinal Determination Network Switches Its Function from Selector to
Differentiation.Desplan C, editor. PLoS Genet. 9:e1003731–17.

Baker NE, Yu SY. 1997. Proneural function of neurogenic genes in the developing Drosophila eye.
Curr. Biol. 7:122–132.

Barkoulas M, van Zon JS, Milloz J, van Oudenaarden A, Felix M-A. 2013. Robustness and Epistasis in
the C. elegans Vulval Signaling Network Revealed by Pathway Dosage Modulation. Dev.
Cell 24:64–75.

Bhattacharya A, Baker NE. 2009. The HLH protein Extramacrochaetae is required for R7 cell and
cone cell fates in the Drosophila eye. Dev. biol. 327:288–300.

Bhattacharya A, Baker NE. 2012. The Role of the bHLH Protein Hairy in Morphogenetic Furrow
Progression in the Developing Drosophila Eye.Hassan BA, editor. PLoS ONE 7:e47503–e47505.

Braendle C, Baer CF, Felix M-A. 2010. Bias and evolution of the mutationally accessible phenotypic
space in a developmental system.Barsh GS, editor. PLoS Genet. 6:e1000877–13.

Brennan CA, Ashburner M, Moses K. 1998. Ecdysone pathway is required for furrow progression in
the developing Drosophila eye. Development 125:2653–2664.

Brown KM, Costanzo MS, Xu W, Roy S, Lozovsky ER, Hartl DL. 2010. Compensatory Mutations
Restore Fitness during the Evolution of Dihydrofolate Reductase. Mol. Biol. Evol. 27:2682–2690.

Brown NL, Sattler CA, Markey DR, Carroll SB. 1991. hairy gene function in the Drosophila eye: normal
expression is dispensable but ectopic expression alters cell fates. Development 113:1245–1256.

Brown NL, Sattler CA, Paddock SW, Carroll SB. 1995a. Hairy and emc negatively regulate
morphogenetic furrow progression in the Drosophila eye. Cell 80:879–887.

Brown NL, Sattler CA, Paddock SW, Carroll SB. 1995b. Hairy and emc negatively regulate
morphogenetic furrow progression in the Drosophila eye. Cell 80:879–887.

Burch CL, Chao L. 1999. Evolution by small steps and rugged landscapes in the RNA virus phi6.
Genetics 151:921–927.

Chandler CH, Chari S, Tack D, Dworkin I. 2014. Causes and consequences of genetic background
effects illuminated by integrative genomic analysis. Genetics 196:1321–1336.

Charusanti P, Conrad TM, Knight EM, Venkataraman K, Fong NL, Xie B, Gao Y, Palsson BØ. 2010.
Genetic Basis of Growth Adaptation of Escherichia coli after Deletion of pgi, a Major Metabolic
Gene.Casadesús J, editor. PLoS Genet. 6:e1001186–13.

Choi HMT, Beck VA, Pierce NA. 2014. Next-generation in situ hybridization chain reaction: higher
gain, lower cost, greater durability. ACS Nano 8:4284–4294.

Chun S, Fay JC. 2011. Evidence for hitchhiking of deleterious mutations within the human
genome.Pritchard JK, editor. PLoS Genet. 7:e1002240.

Cowley DE, Atchley WR. 1988. Quantitative Genetics of Drosophila Melanogaster. II. Heritabilities and
Genetic Correlations between Sexes for Head and Thorax Traits. Genetics 119:421–433.

Dominguez M, Hafen E. 1997. Hedgehog directly controls initiation and propagation of retinal
differentiation in the Drosophila eye. Genes Dev. 11:3254–3264.

Dominguez M, Wasserman JD, Freeman M. 1998. Multiple functions of the EGF receptor in
Drosophila eye development. Curr. Biol. 8:1039–1048.

Duveau F, Felix M-A. 2012. Role of Pleiotropy in the Evolution of a Cryptic Developmental Variation in
Caenorhabditis elegans.Noor MAF, editor. PLoS Biol. 10:e1001230–19.

Dworkin I, Palsson A, Birdsall K, Gibson G. 2003. Evidence that Egfr Contributes to Cryptic Genetic
Variation for Photoreceptor Determination in Natural Populations of Drosophila melanogaster.
Curr. Biol. 13:1888–1893.

Estes S, Lynch M. 2003. Rapid fitness recovery in mutationally degraded lines of Caenorhabditis
elegans. Evolution 57:1022–1030.

Estes S, Phillips PC, Denver DR. 2011. Fitness recovery and compensatory evolution in natural
mutant lines of C. elegans. Evolution 65:2335–2344.

Fear JM, León-Novelo LG, Morse AM, Gerken AR, Van Lehmann K, Tower J, Nuzhdin SV, McIntyre
LM. 2016. Buffering of Genetic Regulatory Networks in Drosophila melanogaster. Genetics
203:1177–1190.

Felix M-A, Barkoulas M. 2012. Robustness and flexibility in nematode vulva development. Trends
Genet. 28:185–195.

Felix M-A, Barkoulas M. 2015. Pervasive robustness in biological systems. Nat. Rev. Genet. 16:483–
496.

Felix M-A. 2007. Cryptic Quantitative Evolution of the Vulva Intercellular Signaling Network in
Caenorhabditis. Curr. Biol. 17:103–114.

Felix M-A. 2012. Evolution in developmental phenotype space. Current Opinion in Genetics &
Development 22:593–599.

Felsenfeld AL, Kennison JA. 1995. Positional signaling by hedgehog in Drosophila imaginal disc
development. Development 121:1–10.

Firth LC, Baker NE. 2005. Extracellular Signals Responsible for Spatially Regulated Proliferation in the
Differentiating Drosophila Eye. Dev. Cell 8:541–551.

Fowlkes CC, Eckenrode KB, Bragdon MD, Meyer M, Wunderlich Z, Simirenko L, Luengo Hendriks CL,
Keranen SVE, Henriquez C, Knowles DW, et al. 2011. A Conserved Developmental Patterning
Network Produces Quantitatively Different Output in Multiple Species of Drosophila.Kopp A,
editor. PLoS Genet. 7:e1002346.

Freeman ABAM. 2001. Notch signalling and the initiation of neural development in the Drosophila
eye. :1–10.

Gavish A, Shwartz A, Weizman A, Schejter E, Shilo B-Z, Barkai N. 2016. Periodic patterning of the
Drosophila eye is stabilized by the diffusible activator Scabrous. Nat. Comm. 7:1–10.

Gibson G, Dworkin I. 2004. Uncovering cryptic genetic variation. Nat. Rev. Genet. 5:681–690.

Gibson G, Reed LK. 2008. Cryptic genetic variation. Curr. Biol. 18:R989–R990.

Gibson G. 2009. Decanalization and the origin of complex disease. Nat. Rev. Genet. 10:134–140.

Goncalves A, Leigh-Brown S, Thybert D, Stefflova K, Turro E, Flicek P, Brazma A, Odom DT, Marioni
JC. 2012. Extensive compensatory cis-trans regulation in the evolution of mouse gene expression.
Genome Res. 22:2376–2384.

Heberlein U, Wolff T, Rubin GM. 1993. The TGF[beta] homolog dpp and the segment polarity gene
hedgehog are required for propagation of a morphogenetic wave in the Drosophila retina. 75:913–
926.

Hu JX, Thomas CE, Brunak S. 2016. Network biology concepts in complex disease comorbidities.
Nat. Rev. Genet. 17:615–629.

Ito Y, Harigai A, Nakata M, Hosoya T, Araya K, Oba Y, Ito A, Ohde T, Yaginuma T, Niimi T. 2013. The
role of doublesex in the evolution of exaggerated horns in the Japanese rhinoceros beetle. Nat.
Neurosci. 14:561–567.

Jarman AP, Grell EH, Ackerman L, Jan LY, Jan YN. 1994. atonal is the proneural gene for Drosophila
photoreceptors. Nature 369:398–400.

Jeong S, Rebeiz M, Andolfatto P, Werner T, True J, Carroll SB. 2008. The evolution of gene
regulation underlies a morphological difference between two *Drosophila* sister species. Cell
132:783–793.

Johnson NA. 2006. The Micro-evolution of development. Genetica 129:1–5.

Keren L, van Dijk D, Weingarten-Gabbay S, Davidi D, Jona G, Weinberger A, Milo R, Segal E. 2015.
Noise in gene expression is coupled to growth rate. Genome Res. 25:1893–1902.

Kienle S, Sommer RJ. 2013. Cryptic variation in vulva development by cis-regulatory evolution of a
HAIRY-binding site. Nat. Comm. 4:1714.

Kopp A, Duncan I, Godt D, Carroll SB. 2000. Genetic control and evolution of sexually dimorphic
characters in *Drosophila*. Nature 408:553–559.

Kozlov K, Kosheverova V, Kamentseva R, Kharchenko M, Sokolkova A, Kornilova E, Samsonova M.
2017. Quantitative analysis of the heterogeneous population of endocytic vesicles. J. Bioinform.
Comput. Biol. 10:1750008.

Kumar JP. 2013. Catching the Next Wave: Patterning of the Drosophila Eye by the Morphogenetic
Furrow. In: Molecular Genetics of Axial Patterning, Growth and Disease in the Drosophila Eye.
New York, NY: Springer New York. pp. 75–97.

Landry CR, Wittkopp PJ, Taubes CH, Ranz JM, Clark AG, Hartl DL. 2005. Compensatory *cis-trans*
evolution and the dysregulation of gene expression in interspecific hybrids of *Drosophila*. Genetics
171:1813–1822.

Lavagnino NJ, Fanara JJ. 2015. Changes Across Development Influence Visible and Cryptic Natural
Variation of Drosophila melanogaster Olfactory Response. Evolutionary biology 43:96–108.

Li W, Ohlmeyer JT, Lane ME, Kalderon D. 1995. Function of protein kinase A in hedgehog signal
transduction and Drosophila imaginal disc development. Cell 80:553–562.

Li Y, Baker NE. 2001. Proneural enhancement by Notch overcomes Suppressor-of-Hairless repressor
function in the developing Drosophila eye. Curr. Biol. 11:330–338.

Liu F, Morrison AH, Gregor T. 2013. Dynamic interpretation of maternal inputs by the Drosophila
segmentation gene network. Proc. Nat. Acad. Sci. USA 110:6724–6729.

Lucas T, Ferraro T, Roelens B, Las Heras Chanes De J, Walczak AM, Coppey M, Dostatni N. 2013.
Live Imaging of Bicoid-Dependent Transcription in Drosophila Embryos. Curr. Biol. 23:2135–2139.

Ludwig MZ, Bergman C, Patel NH, Kreitman M. 2000. Evidence for stabilizing selection in a eukaryotic
enhancer element. Nature 403:564–567.

Ludwig MZ, Patel NH, Kreitman M. 1998. Functional analysis of eve stripe 2 enhancer evolution in
Drosophila: rules governing conservation and change. Development 125:949–958.

Ma C, Moses K. 1995. Wingless and patched are negative regulators of the morphogenetic furrow and
can affect tissue polarity in the developing Drosophila compound eye. Development 121:2279–
2289.

Ma C, Zhou Y, Beachy PA, Moses K. 1993. The segment polarity gene hedgehog is required for
progression of the morphogenetic furrow in the developing Drosophila eye. Cell 75:927–938.

Mackay TFC, Richards S, Stone EA, Barbadilla A, Ayroles JF, Zhu D, Casillas S, Han Y, Magwire
MM, Cridland JM, et al. 2012. The Drosophila melanogaster Genetic Reference Panel. Nature
482:173–178.

Maisnier-Patin S, Andersson DI. 2004. Adaptation to the deleterious effects of antimicrobial drug
resistance mutations by compensatory evolution. Research in Microbiology 155:360–369.

Martinez C, Rest JS, Kim A-R, Ludwig M, Kreitman M, White K, Reinitz J. 2014. Ancestral
Resurrection of the Drosophila S2E Enhancer Reveals Accessible Evolutionary Paths through
Compensatory Change. Mol. Biol. Evol. 31:903–916.

Mathews KW, Cavegn M, Zwicky M. 2017. Sexual dimorphism of body size is controlled by dosage of
the X-chromosomal gene Myc and by the sex-determining gene tra in Drosophila. Genetics.

McGregor AP, Shaw PJ, Hancock JM, Bopp D, Hediger M, Wratten NS, Dover GA. 2001. Rapid
restructuring of bicoid-dependent hunchback promoters within and between Dipteran species:
implications for molecular coevolution. Evolution \& Development 3:397–407.

McKenzie JA, Clarke GM. 1988. Diazinon resistance, fluctuating asymmetry and fitness in the
Australian sheep blowfly, lucilia cuprina. Genetics 120:213–220.

McKenzie JA, Whitten MJ, Adena MA. 1982. The effect of genetic background on the fitness of
diazinon resistance genotypes of the Australian sheep blowfly, Lucilia cuprina. Heredity 49:1–9.

McKenzie JA. 1993. Measuring fitness and intergenic interactions: The evolution of resistance to
diazinon inLucilia cuprina. Genetica 90:227–237.

Meyer F. 1994. Topographic distance and watershed lines. Signal processing 38:113–125.

Moore FB, Rozen DE, Lenski RE. 2000. Pervasive compensatory adaptation in Escherichia coli. Proc.
R. Soc. B 267:515–522.

Namba R, Pazdera TM, Cerrone RL, Minden JS. 1997. Drosophila embryonic pattern repair: how
embryos respond to bicoid dosage alteration. Development 124:1393–1403.

Nunes MDS, Arif S, Schlötterer C, McGregor AP. 2013. A perspective on micro-evo-devo: progress
and potential. Genetics 195:625–634.

others SSA. 1998. Regulation of EGF receptor signaling establishes pattern across the developing
Drosophila retina. :1–14.

Paaby A, Gibson G. 2016. Cryptic Genetic Variation in Evolutionary Developmental Genetics. Biology
5:28–13.

Paaby AB, Rockman MV. 2014. Cryptic genetic variation: evolution's hidden substrate. Nat. Rev.
Genet. 15:247–258.

Palsson A, Wesolowska N, Reynisdóttir S, Ludwig MZ, Kreitman M. 2014. Naturally Occurring
Deletions of Hunchback Binding Sites in the Even-Skipped Stripe 3+7 Enhancer.Moses AM,
editor. PLoS ONE 9:e91924–15.

Pan D, Rubin GM. 1995. cAMP-dependent protein kinase and hedgehog act antagonistically in
regulating decapentaplegic transcription in Drosophila imaginal discs. Cell 80:543–552.

Posnien N, Hopfen C, Hilbrant M, Ramos-Womack M, Murat S, Schönauer A, Herbert SL, Nunes
MDS, Arif S, Breuker CJ, et al. 2012. Evolution of Eye Morphology and Rhodopsin Expression in
the Drosophila melanogaster Species Subgroup.Flatt T, editor. PLoS ONE 7:e37346–11.

Raj A, van den Bogaard P, Rifkin SA, van Oudenaarden A, Tyagi S. 2008. Imaging individual mRNA
molecules using multiple singly labeled probes. Nat. Meth. 5:877–879.

Reed RD, Papa R, Martin A, Hines HM, Counterman BA, Pardo-Diaz C, Jiggins CD, Chamberlain NL,
Kronforst MR, Chen R, et al. 2011. optix Drives the Repeated Convergent Evolution of Butterfly
Wing Pattern Mimicry. Science 333:1137–1141.

Rideout EJ, Dornan AJ, Neville MC, Eadie S, Goodwin SF. 2010. Control of sexual differentiation and
behavior by the *doublesex* gene in *Drosophila* melanogaster. Nat. Neurosci. 13:458–466.

Roignant J-Y, Treisman JE. 2009. Pattern formation in the Drosophila eye disc. Int. J. Dev. Biol.
53:795–804.

Rosenblum EB, Römpler H, Schöneberg T, Hoekstra HE. 2010. Molecular and functional basis of
phenotypic convergence in white lizards at White Sands. Proc. Nat. Acad. Sci. USA 107:2113–
2117.

Shah S, Lubeck E, Schwarzkopf M, He TF. 2016. Single-molecule RNA detection at depth via
hybridization chain reaction and tissue hydrogel embedding and clearing. Genetics.

Signor SA, Li Y, Rebeiz M, Kopp A. Genetic convergence in the evolution of male-specific color
patterns in Drosophila

. Curr. Biol.

Signor SA, New F, Nuzhdin S. An abundance of high frequency variance uncovered in a large panel
of Drosophila simulans. Heredity.

Stoebel DM, Hokamp K, Last MS, Dorman CJ. 2009. Compensatory Evolution of Gene Regulation in
Response to Stress by Escherichia coli Lacking RpoS.Guttman DS, editor. PLoS Genet.
5:e1000671–e1000679.

Struhl SGAG. 1999. Progression of the morphogenetic furrow in the Drosophila eye: the roles of
Hedgehog, Decapentaplegic and the Raf pathway. :1–14.

Strutt DI, Mlodzik M. 1997. Hedgehog is an indirect regulator of morphogenetic furrow progression in
the Drosophila eye disc. Development 124:3233–3240.

Sun Y, Jan LY, Jan YN. 1998. Transcriptional regulation of atonal during development of the
Drosophila peripheral nervous system. Development 125:3731–3740.

Szamecz B, Boross G, Kalapis D, Kovács K, Fekete G, Farkas Z, Lázár V, Hrtyan M, Kemmeren P,
Groot Koerkamp MJA, et al. 2014. The Genomic Landscape of Compensatory Evolution.Barton
NH, editor. PLoS Biol. 12:e1001935–16.

Taylor MB, Ehrenreich IM. 2015a. Transcriptional Derepression Uncovers Cryptic Higher-Order
Genetic Interactions.Wittkopp P, editor. PLoS Genet. 11:e1005606–e1005614.

Taylor MB, Ehrenreich IM. 2015b. Transcriptional Derepression Uncovers Cryptic Higher-Order
Genetic Interactions.Wittkopp P, editor. PLoS Genet. 11:e1005606–e1005614.

Thompson A, Zakon HH, Kirkpatrick M. 2015. Compensatory Drift and the Evolutionary Dynamics of
Dosage-Sensitive Duplicate Genes. Genetics:1–29.

True JR, Haag ES. 2001. Developmental system drift and flexibility in evolutionary trajectories. 3:109–
119.

Tsachaki M, Sprecher SG. 2011. Genetic and developmental mechanisms underlying the formation of
the Drosophila compound eye.Singh A, Irvine KD, editors. Dev. Dyn. 241:40–56.

Vincent L. 1993. Morphological grayscale reconstruction in image analysis: Applications and efficient
algorithms. IEEE transactions on image processing 2:176–201.

Ward LD, Kellis M. 2012. Interpreting noncoding genetic variation in complex traits and human
disease. Nat. Neurosci. 30:1095–1106.

Woodruff GC, Eke O, Baird SE, Felix M-A, Haag ES. 2010. Insights into species divergence and the
evolution of hermaphroditism from fertile interspecies hybrids of *Caenorhabditis* nematodes.
Genetics 186:997–1012.

Wunderlich Z, Bragdon MD, Eckenrode KB, Lydiard-Martin T, Pearl-Waserman S, DePace AH. 2012.
Dissecting sources of quantitative gene expression pattern divergence between Drosophila
species. Mol. Syst. Biol. 8:1–15.

Yassin A, Delaney EK, Reddieux A, Bastide H, Seher T, Appleton NC, Lack JB, David JR, Chenoweth
SF, Pool JE, et al. *pdm3* is responsible for recurrent evolution of female-limited color dimorphism
in *Drosophil*. Curr. Biol.

710